Alteration of mean amplitude of low frequency fluctuations measured by fMRI at 11.7 T in the aging brain of mouse lemur primate

Clément M. Garin [1,2], Marc Dhenain[1,2,*]

[1] Centre National de la Recherche Scientifique (CNRS), Université Paris-Sud, Université Paris-Saclay UMR 9199, Neurodegenerative Diseases Laboratory, 18 Route du Panorama, F-92265 Fontenay-aux-Roses, France

[2] Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), Direction de la Recherche Fondamentale (DRF), Institut François Jacob, MIRCen, 18 Route du Panorama, F-92265 Fontenay-aux-Roses, France

Abbreviated title: Alteration of mALFF in aging primates

Corresponding authors:

Marc Dhenain

MIRCen, UMR CEA-CNRS 9199

18 Route du Panorama

92 265 Fontenay-aux-Roses CEDEX

France

Tel: +33 1 46 54 81 92; Fax: +33 1 46 54 84 51

email: Marc.Dhenain@cnrs.fr

Abstract

Non-human primates are a critical species for the identification of key biological mechanisms in normal and pathological aging. One of these primates, the mouse lemur, has been widely studied as a model of cerebral aging or Alzheimer's disease. The amplitude of low-frequency fluctuations of blood oxygenation level-dependent (BOLD) can be measured with functional MRI. Within specific frequency bands (*e.g.* the 0.01–0.1 Hz), these amplitudes were proposed to indirectly reflect neuronal activity as well as glucose metabolism. Here, we first created whole brain maps of the mean amplitude of low frequency fluctuations (mALFF) in middle-aged mouse lemurs. Then, we extracted mALFF in old lemurs to identify age-related changes. A high level of mALFF was detected in the temporal cortex (Brodmann area 20), somatosensory areas (Brodmann area 5), insula (Brodmann area 13-6) and the parietal cortex (Brodmann area 7) of healthy middle-aged mouse lemurs. Aging was associated with alteration of mALFF in somatosensory areas (area 5) and the parietal cortex (area 7).

Graphical abstract

Highlights

- We characterized mean amplitude of fluctuation at low frequencies (mALFF) in mouse lemurs.
- mALFF was the highest in regions involved in visuo-somatosensory-motor function (Brodmann areas 5, 7, 20) and in more integrative functions (area 13-16).
- mALFF was affected by aging in regions involved in visuo-somatosensory-motor function (parietal areas 5 and 7).
- mALFF is a useful marker to investigate age-related cerebral dysfunction in animals.

Keywords

Amplitude of low frequency fluctuations, fMRI, mouse lemur, resting-state functional MRI, BOLD, ALFF, network, functional, mALFF.

Significance Statement

- The amplitude of low-frequency fluctuations (ALFF) is expected to reflect neuronal activity. It has been proposed as an MRI-based method to evaluate brain function.

- ALFF has been used to investigate different cerebral pathologies in animal models but the regional differences of ALFF signal and the impact of cerebral aging on ALFF has never been characterized.

- Here, we highlight for the first time regional difference of ALFF. High signal was detected in regions involved in visuo-somatosensory-motor function as well as in more integrative functions. ALFF was reduced in regions involved in visuo-somatosensory-motor function during aging. ALFF is thus a useful marker to investigate age-related cerebral dysfunction.

1. Introduction

Human life expectancy has dramatically increased during the last century. This comes with an increased risk for cerebral alterations leading to neurodegenerative diseases or mild cognitive/motor impairments that impair daily living. Monitoring functional impairments associated with cerebral ageing with translational imaging methods is critical to understand mechanisms leading to brain alterations and develop new treatments.

Measures of cerebral glucose metabolism using the radiolabeled glucose analog [18]F-fluorodeoxyglucose (FDG) detected by Positron Emission Tomography (PET) imaging is a largely used marker to investigate brain function in elderlies (Edison et al., 2007; Kalpouzos et al., 2009) and in mouse model of Alzheimer's disease (Franke et al., 2020). The use of PET is however restricted as this requires using radioactive compounds and is limited for animal studies by its low resolution when compared to functional magnetic resonance imaging (fMRI). In consequences, magnetic resonance imaging (MRI) is an interesting alternative to PET.

Low-frequency oscillations (LFO) of blood-oxygen level dependent (BOLD) signal reflects the total power of BOLD signal within the frequency range between 0.01 and 0.1 Hz. The amplitude of low-frequency fluctuations (ALFF) is expected to reflect neuronal activity (Zou et al., 2008) and was associated with markers of glucose metabolism (Aiello et al., 2015). Thus, it has been proposed as an MRI-based method to evaluate brain function (Biswal et al., 1995; Zou et al., 2008) and could be a promising radioactive-free alternative to FDG-PET. Studies in humans have shown that ALFF is negatively correlated with age in several brain regions such as the supplementary motor area, pre-supplementary motor area, anterior cingulate cortex, bilateral dorsal lateral prefrontal cortex, right inferior parietal lobule, and posterior cingulate cortex (Hu et al., 2014). However, healthy aging effects on ALFF indexes remain to our knowledge, unexplored in non-human mammalians.

The mouse lemur (*Microcebus murinus*) is a primate attracting increased attention in neuroscience research. This small animal (typical length 12cm, 60-120g weight) has a decade-long lifespan (Pifferi et al., 2018) and is a model for studying cerebral ageing (Sawiak et al., 2014) and Alzheimer's disease (Kraska et al., 2011). This animal was used to establish the impact of prediabetes in the brain (Djelti et al., 2016) as well as to evaluate interventions modulating cerebral ageing process (Pifferi et al., 2018). The aim of the current study was thus to characterize ALFF in this primate in normal and

ageing conditions. We described regional differences of ALFF signal and showed aged-related changes in specific brain regions.

2. Materials and methods

2.1. Animals and breeding

This study was carried out in accordance with the recommendations of the European Communities Council directive (2010/63/EU). The protocol was approved by the local ethics committees CEtEA-CEA DSV IdF (authorizations 201506051736524 VI (APAFIS#778)).

Thirty-three mouse lemurs (21 males and 12 females) were initially included in this study. They were born in the laboratory-breeding colony of the CNRS/MNHN in Brunoy, France (UMR 7179 CNRS/MNHN) and bred in our laboratory (Molecular Imaging Research Center, CEA, Fontenay-aux-Roses). Four animals that presented brain lesions or MR images with artefacts were excluded from the analysis. Fourteen animals ranged from 1.3 to 3.8 years old (mean±SD: 2.1±0.8 years) were grouped together to form the *"middle-aged lemur cohort"* (Table 1). Fifteen animals ranged from 8.0 to 10.8 years old (mean±SD: 8.8±1.1 years) were grouped together to form the *"old lemur cohort"* (Table 1). Housing conditions were cages containing one or two lemurs with jumping and hiding enrichment, temperatures 24–26°C, relative humidity 55% and seasonal lighting (summer: 14 hours of light/10 hours of dark; winter: 10 hours of light/14 hours of dark). Food consisted of fresh apples and a homemade mixture of bananas, cereals, eggs and milk. Animals had free access to tap water. None of the animals had previously been involved in pharmacological trials or invasive studies.

Subject	Sex	Age (years)	Animal rejected
283EA	M	1.3	NO
283CCA	M	1.3	NO
285AAA	M	1.4	NO
365A	M	1.4	NO
285AB	M	1.4	NO
263BCE	M	1.4	NO
314CA	M	1.5	NO
285D	M	2.3	NO

283CA	M	2.4	NO
276BC	M	2.4	NO
285E	M	2.4	NO
300BA	M	2.5	NO
289BB	F	2.9	NO
208CBF	F	3.1	NO
288BC	F	3.2	cerebral lesion
310C	F	3.8	cerebral lesion
967HACA	M	8.0	NO
184CB	F	8.0	NO
965MBIA	M	8.0	NO
965MBFA	M	8.0	NO
965MBFC	F	8.0	NO
965MBGA	M	8.0	NO
967HACB	F	8.0	NO
965MBFB	M	8.1	NO
169BAB	F	8.1	NO
965FDBB	M	8.7	NO
147BCBB	M	8.7	NO
147BCBA	M	8.8	artifact
943GKBC	F	8.8	NO
153FBA	M	10.3	NO
216B	F	10.4	NO
965MBG	F	10.8	artifact
119BBB	F	10.8	NO

Table 1. Cohort of mouse lemurs involved in the study.

2.2. Animal preparation and MRI acquisition

All animals were scanned once, under isoflurane anesthesia at 1.25-1.5% in air, with respiratory rate monitored to confirm animal stability until the end of the experiment. Body temperature was maintained by an air heating system at 32°C, inducing a natural torpor in mouse lemurs (Aujard, 2001). This has the advantage of allowing a low anesthesia level without reawakening. The MRI system was an 11.7 T Bruker BioSpec (Bruker, Ettlingen, Germany) running ParaVision 6.0.1 with a volume coil for radiofrequency transmission and a quadrature surface coil for reception

(Bruker, Ettlingen, Germany). Anatomical images were acquired using a T2-weighted multi-slice multi-echo (MSME) sequence: TR = 5000 ms, TE = 17.5 ms, FOV = 32 × 32 mm, 75 slices of 0.2 mm thickness, 6 echoes, 5 ms IET, resolution = 200 µm isotropic, acquisition duration 10 min. Resting state time series data were acquired using a gradient-echo echo planar imaging (EPI) sequence: TR = 1000 ms, TE = 10.0 ms, flip angle = 90°, repetitions = 450, FOV = 30 × 20 mm, 23 slices of 0.9 mm thickness and 0.1 mm gap, resolution = 312.5 × 208.33 × 1000 µm, acquisition duration 7m30s.

2.3. MRI pre-processing

Scanner data were exported as DICOM files then converted into NIfTI-1 format. Then spatial preprocessing was performed using the python module sammba-mri (SmAll MaMmals BrAin MRI; (Celestine et al., 2020), http://sammba-mri.github.io) which, using nipype for pipelining (Gorgolewski et al., 2011), leverages AFNI (Cox, 1996) for most steps and RATS (Oguz et al., 2014) for brain extraction. Anatomical images were mutually registered to create a study template, which was further registered to a high-resolution anatomical mouse lemur template of the functional atlas (Garin et al., 2019). Resting state images were corrected for slice timing (interleaved), motion, and B0 distortion (per-slice registration to respective anatomical). Then all the images were brought into the same space of the mouse lemur template by successive application of the individual anatomical to study template and study template to mouse lemur atlas transforms.

Functional images were further pre-treated using AFNI afni_proc.py (Cox, 1996). fMRI images were corrected for slice timing, motion, detrend, smoothed (0.9 mm) and bandpass filtered (0.01 to 0.1 Hz). TRs with excessive motion of 0.07 mm or where too many voxels were flagged as outliers by 3dToutcount (AFNI), were censored. The first five volumes were excluded from analysis to ensure steady-state magnetization.

2.4. mALFF calculation and extraction

LFO measures were performed using the fast Fourier transform index: amplitude of low-frequency fluctuation (ALFF) (Zuo et al., 2010). As ALFF is sensitive to the scale of raw signal and the unit of BOLD signal is arbitrary, the original ALFF value is not adapted for comparisons between animals. In addition, ALFF can be susceptible to signal fluctuations caused by physiological noise unrelated to brain activity (Zou et al., 2008). A standardization procedure has been proposed by dividing the signal of each voxel by the global mean ALFF in each animal (Jia et al., 2020). The newly calculated

index is called mean ALFF (mALFF). mALFF indexes were calculated for each voxel of the pre-processed EPI images in the low-frequencies range 0.01 to 0.1 Hz using the function "3dLombScargle" and "3dAmpToRSFC" from AFNI (Cox, 1996). The mALFF signal of each voxel was extracted within the different regions based on the anatomical atlas (Nadkarni et al., 2019) using NiftiLabelsMasker from Nilearn (Abraham et al., 2014).

2.5. Statistical analysis

Voxel wise analysis was performed using 3dttest++ from AFNI (Cox, 1996). This analysis identifies statistical differences between groups by using an uncorrected pvalue and a cluster size. Analysis space was reduced by thresholding the average mALFF map to the 20% highest voxels. Cluster size (856 voxels) was estimated on these remaining voxels using AFNI (-Clusterize) with p<0.05 (uncorrected). Then, using "map_threshold" from nilearn (Abraham et al., 2014), we extracted on the Ttest map any cluster of statistical voxels superior to 856 associated to p<0.05.

3. Results

3.1. mALFF in middle-aged mouse lemurs

mALFF maps were recorded from 14 middle-aged mouse lemurs. Individual mALFF maps of each animal were averaged to produce 3D maps of the group (Fig. 1). Automatic extractions of the mALFF signal were then performed in various cerebral regions by using a reference anatomical atlas ((Nadkarni et al., 2019); Fig. 1A). The highest cortical signal was observed in the temporal cortex (Brodmann area (BA) 20 *i.e.* secondary visual area expected to be involved in visual processing and recognition memory), parietal cortex (BA 5 *i.e.* a secondary somatosensory areas; BA 7 *i.e.* an integrative area involved in visuo-motor coordination). High signal was also detected in the primary somatosensory cortex (BA 1-3) and the primary motor cortex (BA 4). More integrative areas as the insula (BA 13-16) and in the cingulate cortex (BA 23 and 24) also displayed high signal. In subcortical areas, the basal forebrain exhibited the most elevated signal when compared to the rest of the brain. Conversely, cortical regions such as BA 30 (agranular retrolimbic area involved in vision) and the adjacent BA 27 (area presubicularis involved in vision) or area 25 (antero-ventral part of the cingulate cortex) displayed the lowest levels of mALFF. Subcortical regions such as substantia nigra, subthalamic nucleus or hypophysis also displayed the lowest levels of mALFF.

Figure 1. Whole brain mALFF average-map in middle-aged mouse lemurs.

(A) mALFF in different brain regions of middle-aged adult mouse lemurs (n=14) ranked based on their group median value. Elevated mALFF is observed within regions encompassing cortical Brodmann areas 20, 5, 13-16, 7, 1-3, 4, 23, 24 and basal forebrain. 3D surface average-map of the mALFF showing regional signal variations (B). ant: anterior part of the brain, post: posterior part of the brain.

3.2. Age-related changes of mALFF

BOLD signal was recorded from 15 old mouse lemurs (mean age±SD: 8.8±1.1

years) and as for middle-aged lemurs, mALFF analysis was performed. Average mALFF maps are displayed in Fig. 2A for middle-aged mouse lemurs and in Fig. 2B for old mouse lemurs. Comparison of average mALFF map between middle-aged and aged lemurs indicates lower mALFF in the parietal cortex of aged animals and a trend toward higher signal in their hypothalamic regions. Voxelwise analysis between groups revealed a significant loss of BOLD signal amplitude in parietal regions involving (BA 5 and BA 7) of aged animals (Fig 2C, D).

Figure 2. mALFF contrast in middle-aged and old mouse lemurs.

Average mALFF in middle-aged (A) and old mouse lemurs (B). Significant statistical decrease of the mALFF signal was detected using voxelwise analysis in the Brodmann areas 5 and 7 (C). Anatomical view of the mouse lemur atlas corresponding to the displayed sections (A-C) are shown in D (Nadkarni et al., 2019). BA: Brodmann area. Scale bars in A-C: mALFF contrast, in D: z-score.

4. Discussion

4.1. Cerebral distribution of mALFF index in mouse lemurs

This study evaluated mALFF index in mouse lemur primates at high field (11.7T). The evaluation of mALFF in a middle-aged cohort provided 3D maps of the normal distribution of the mALFF indexes. Highest levels of mALFF were detected in cortical structures involved in secondary visual (area 20 in temporal region), somatosensory (area 5 in parietal region), and in motor processing and integrative regions involved in

visuo-motor coordination (area 7 in parietal region). High signal was also detected in the primary somatosensory cortex (BA 1-3) and the primary motor cortex (BA 4). More integrative areas as the insula (BA 13-16) and in the cingulate cortex (BA 23 and 24) also displayed high signal.

4.2. Age-related changes of mALFF index

As a second part of the study, we assessed age-related changes of mALFF. In humans, healthy ageing mainly affects ALFF in brain regions involved in motor function such as the supplementary motor area (BA 5) or the pre-supplementary motor area. More integrative regions such as the anterior cingulate cortex, the dorso-lateral prefrontal cortex, the posterior cingulate cortex, and the inferior parietal lobule are also impaired (Hu et al., 2014). In mouse lemurs, the only regions significantly affected by healthy ageing were those involved in visuo-somatosensory-motor function (parietal regions as BA 5 *i.e.* a secondary somatosensory areas and BA 7 *i.e.* an integrative area involved in visuo-motor coordination). These two regions are involved in somatosensory processing, in movement such as grasping (Gardner et al., 2007) or object location (Caminiti et al., 2010). These alterations could participate to the described visuo-somatosensory-motor alterations reported in aged lemurs (Le Brazidec et al., 2017).

Atrophy detected in mouse lemurs impacts the whole brain in the latest stages of ageing but the most prominent atrophied areas include the insular (areas 13-16), frontal (area 6), parietal (areas 5, 7), occipital (areas 17, 18), inferior temporal (areas 21, 28) and cingulate cortices (areas 23, 24, 25) (Nadkarni et al., 2019; Sawiak et al., 2014). In consequence, an ageing effect can be detected with mALFF and anatomical measures in both areas 5 and 7 but areas such as 17, 18, 13-16 are impacted only by atrophy and not by mALFF changes. A better understanding of the significance of mALFF changes is now needed to further interpret the significance of mALFF changes.

In previous studies, ALFF or fractional ALFF were used to evaluate the effect of simian immunodeficiency virus (Zhao et al., 2019), spinal injury (Rao et al., 2014) or anesthesia (Wu et al., 2016) in macaques. In mice, Huntington-related pathological effects were detected using ALFF (Chang et al., 2018). In rats, various effects of stress could also be evaluated using ALFF (Li et al., 2018; Yan et al., 2017). By showing age-related changes, the current study increases the range of ALFF changes detected in mammals in pathological situations.

4.3. Limitations

Anesthesia is known to interact with brain function. For example, Isoflurane changes functional connections between brain regions in marmosets (Garin et al., 2022). Thus, one of the possible limitations of the study is anesthesia that could modulate mALFF signal and lead to whole brain mALFF average-map that does not reflect mALFF in non-anesthetized animals. Future studies will thus have to be conducted to investigate the impact of anesthesia on mALFF.

Comparison between middle-aged and old animals revealed age-related differences of mALFF. As both groups were anesthetized, this difference is expected to reflect the ageing effect. We can however not exclude that anesthesia may have decreased our ability to detect mALFF changes in some regions affected by ageing.

5. Conclusion

In conclusion, this study provides the proof of concept that mALFF can be measured in the whole brain of mouse lemurs and can detect aged-related changes. It proposes mALFF as a tool for the exploration of the cerebral function in mammals as well as an interesting candidate for the longitudinal follow up of age-related cerebral dysfunction animal models.

6. Acknowledgements

We thank the France-Alzheimer Association, Plan Alzheimer Foundation and the French Public Investment Bank's "ROMANE" program for funding this study. The 11.7T MRI scanner was funded by a grant from NeurATRIS: A Translational Research Infrastructure for Biotherapies in Neurosciences ("Investissements d'Avenir", ANR-11-INBS-0011). C.G. was financed by the French Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation.

7. Competing interests

The authors do not have financial and non-financial competing interests in relation to the work described.

8. Author contributions

C.M.G. and M.D. contributed to the study conception and design. C.M.G. designed the BOLD sequences, C.M.G. designed registrations strategies and pipelines. C.M.G. and M.D. wrote the manuscript.

9. References

Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., Gramfort, A., Thirion, B., Varoquaux, G., 2014. Machine learning for neuroimaging with scikit-learn. Front Neuroinform 8, 14.

Aiello, M., Salvatore, E., Cachia, A., Pappata, S., Cavaliere, C., Prinster, A., Nicolai, E., Salvatore, M., Baron, J.C., Quarantelli, M., 2015. Relationship between simultaneously acquired resting-state regional cerebral glucose metabolism and functional MRI: a PET/MR hybrid scanner study. Neuroimage 113, 111-121.

Aujard, F., 2001. Effect of ambient temperature on the body temperature rhythm of male gray mouse lemurs (Microcebus murinus). Int J Primatol 22, 43-56.

Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magn Reson Med 34, 537-541.

Caminiti, R., Chafee, M.V., Battaglia-Mayer, A., Averbeck, B.B., Crowe, D.A., Georgopoulos, A.P., 2010. Understanding the parietal lobe syndrome from a neurophysiological and evolutionary perspective. Eur J Neurosci 31, 2320-2340.

Celestine, M., Nadkarni, N.A., Garin, C.M., Bougacha, S., Dhenain, M., 2020. Sammba-MRI: A library for processing SmAll-MaMmal BrAin MRI data in Python. Front Neuroinform 14, 24.

Chang, W.T., Puspitasari, F., Garcia-Miralles, M., Yeow, L.Y., Tay, H.C., Koh, K.B., Tan, L.J., Pouladi, M.A., Chuang, K.H., 2018. Connectomic imaging reveals Huntington-related pathological and pharmaceutical effects in a mouse model. NMR Biomed 31, e4007.

Cox, R.W., 1996. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. Comput Biomed Res 29, 162-173.

Djelti, F., Dhenain, M., Terrien, J., Picq, J.L., Hardy, I., Champeval, D., Perret, M., Schenker, E., Epelbaum, J., Aujard, F., 2016. Impaired fasting blood glucose is associated to cognitive impairment and cerebral atrophy in middle-aged non-human primates. Aging (Albany NY) 9, 173-186.

Edison, P., Archer, H.A., Hinz, R., Hammers, A., Pavese, N., Tai, Y.F., Hotton, G., Cutler, D., Fox, N., Kennedy, A., Rossor, M., Brooks, D.J., 2007. Amyloid, hypometabolism, and cognition in Alzheimer disease: an [11C]PIB and [18F]FDG PET study. Neurology 68, 501-508.

Franke, T.N., Irwin, C., Bayer, T.A., Brenner, W., Beindorff, N., Bouter, C., Bouter,

Y., 2020. In vivo Imaging With (18)F-FDG- and (18)F-Florbetaben-PET/MRI detects pathological changes in the brain of the commonly used 5XFAD mouse model of Alzheimer's Disease. Front Med (Lausanne) 7, 529.

Gardner, E.P., Babu, K.S., Ghosh, S., Sherwood, A., Chen, J., 2007. Neurophysiology of prehension. III. Representation of object features in posterior parietal cortex of the macaque monkey. J Neurophysiol 98, 3708-3730.

Garin, C.M., Hori, Y., Everling, S., Whitlow, C.T., Calabro, F.J., Luna, B., Froesel, M., Gacoin, M., Ben Hamed, S., Dhenain, M., Constantinidis, C., 2022. An evolutionary gap in primate default mode network organization. Cell Rep 39, 110669.

Garin, C.M., Nadkarni, N.A., Landeau, B., Chételat, G., Picq, J.-L., Bougacha, S., Dhenain, M., 2021. Resting state functional atlas and cerebral networks in mouse lemur primates at 11.7 Tesla. NeuroImage 226, 117589.

Gorgolewski, K., Burns, C.D., Madison, C., Clark, D., Halchenko, Y.O., Waskom, M.L., Ghosh, S.S., 2011. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. Front Neuroinform 5, 13.

Hu, S., Chao, H.H., Zhang, S., Ide, J.S., Li, C.S., 2014. Changes in cerebral morphometry and amplitude of low-frequency fluctuations of BOLD signals during healthy aging: correlation with inhibitory control. Brain Struct Funct 219, 983-994.

Jia, X.Z., Sun, J.W., Ji, G.J., Liao, W., Lv, Y.T., Wang, J., Wang, Z., Zhang, H., Liu, D.Q., Zang, Y.F., 2020. Percent amplitude of fluctuation: A simple measure for resting-state fMRI signal at single voxel level. PLoS One 15, e0227021.

Kalpouzos, G., Chételat, G., Baron, J.C., Landeau, B., Mevel, K., Godeau, C., Barré, L., Constans, J.M., Viader, F., Eustache, F., Desgranges, B., 2009. Voxel-based mapping of brain gray matter volume and glucose metabolism profiles in normal aging. Neurobiol Aging 30, 112-124.

Kraska, A., Dorieux, O., Picq, J.L., Petit, F., Bourrin, E., Chenu, E., Volk, A., Perret, M., Hantraye, P., Mestre-Frances, N., Aujard, F., Dhenain, M., 2011. Age-associated cerebral atrophy in mouse lemur primates. Neurobiol Aging 32, 894-906.

Le Brazidec, M., Herrel, A., Thomas, P., Grégoire, B.-A., Aujard, F., Pouydebat, E., 2017. How aging affects grasping behavior and pull strength in captive gray mouse lemurs (Microcebus murinus). Int J Primatol 38, 1120-1129.

Li, J., Yang, R., Xia, K., Wang, T., Nie, B., Gao, K., Chen, J., Zhao, H., Li, Y., Wang, W., 2018. Effects of stress on behavior and resting-state fMRI in rats and evaluation of Telmisartan therapy in a stress-induced depression model. BMC Psychiatry 18, 337.

Nadkarni, N.A., Bougacha, S., Garin, C., Dhenain, M., Picq, J.L., 2019. A 3D population-based brain atlas of the mouse lemur primate with examples of applications in aging studies and comparative anatomy. Neuroimage 185, 85-95.

Oguz, I., Zhang, H., Rumple, A., Sonka, M., 2014. RATS: Rapid Automatic Tissue Segmentation in rodent brain MRI. J Neurosci Methods 221, 175-182.

Pifferi, F., Terrien, J., Marchal, J., Dal-Pan, A., Djelti, F., Hardy, I., Chahory, S., Cordonnier, N., Desquilbet, L., Hurion, M., Zahariev, A., Chery, I., Zizzari, P., Perret, M., Epelbaum, J., Blanc, S., Picq, J.L., Dhenain, M., Aujard, F., 2018. Caloric restriction increases lifespan but affects brain integrity in grey mouse lemur primates. Commun Biol 1, 30.

Rao, J.S., Ma, M., Zhao, C., Zhang, A.F., Yang, Z.Y., Liu, Z., Li, X.G., 2014. Fractional amplitude of low-frequency fluctuation changes in monkeys with spinal cord injury: a resting-state fMRI study. Magn Reson Imaging 32, 482-486.

Sawiak, S.J., Picq, J.L., Dhenain, M., 2014. Voxel-based morphometry analyses of in vivo MRI in the aging mouse lemur primate. Front Aging Neurosci 6, 82.

Wu, T.L., Mishra, A., Wang, F., Yang, P.F., Gore, J.C., Chen, L.M., 2016. Effects of isoflurane anesthesia on resting-state fMRI signals and functional connectivity within primary somatosensory cortex of monkeys. Brain Behav 6, e00591.

Yan, C.G., Rincon-Cortes, M., Raineki, C., Sarro, E., Colcombe, S., Guilfoyle, D.N., Yang, Z., Gerum, S., Biswal, B.B., Milham, M.P., Sullivan, R.M., Castellanos, F.X., 2017. Aberrant development of intrinsic brain activity in a rat model of caregiver maltreatment of offspring. Transl Psychiatry 7, e1005.

Zhao, J., Chen, F., Ren, M., Li, L., Li, A., Jing, B., Li, H., 2019. Low-frequency fluctuation characteristics in rhesus macaques with SIV infection: a resting-state fMRI study. J Neurovirol 25, 141-149.

Zou, Q.H., Zhu, C.Z., Yang, Y., Zuo, X.N., Long, X.Y., Cao, Q.J., Wang, Y.F., Zang, Y.F., 2008. An improved approach to detection of amplitude of low-frequency fluctuation (ALFF) for resting-state fMRI: fractional ALFF. J Neurosci Methods 172, 137-141.

Zuo, X.N., Di Martino, A., Kelly, C., Shehzad, Z.E., Gee, D.G., Klein, D.F., Castellanos, F.X., Biswal, B.B., Milham, M.P., 2010. The oscillating brain: complex and reliable. Neuroimage 49, 1432-1445.