1

Actomyosin remodeling regulates biomineral formation, growth and morphology during eukaryote skeletogenesis

Eman Hijaze[1], Tsvia Gildor[1], Ronald Seidel[2], Majed Layous[1], Mark Winter[3], Luca Bertinetti[2], Yael Politi[2] and Smadar Ben-Tabou de-Leon[1,*]

[1]Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Haifa 31905, Israel.

[2]B CUBE Center for Molecular Bioengineering, Technische Universität Dresden, 01309 Dresden, Germany.

[3]Department of Electrical Engineering, Computer Science and Mathematics, Technische Universiteit Delft, Delft 2628CD, Netherlands

*Correspondence: Smadar Ben-Tabou de-Leon (sben-tab@univ.haifa.ac.il)

Abstract

Biomineralization had apparently evolved independently in different phyla, using distinct minerals, organic scaffolds and gene regulatory networks (GRNs). However, diverse eukaryotes from unicellular organisms, through echinoderms to vertebrates, use the actomyosin network during biomineralization. Specifically, the actomyosin remodeling protein, Rho-associated coiled-coil kinase (ROCK) regulates cell differentiation and gene expression in vertebrates' biomineralizing cells, yet, little is known on ROCK's role in invertebrates' biomineralization. Here we reveal that ROCK controls the formation, growth and morphology of the calcite spicules in the sea urchin larva. ROCK expression is elevated in the sea urchin skeletogenic cells downstream of the Vascular Endothelial Growth Factor (VEGF) signling. ROCK inhibition impairs the organization of F-actin around the spicules, disrupts skeletogenic gene expression and leads to skeletal loss. ROCK inhibition after spicule formation reduces spicule elongation rate and induces ectopic spicule branching. Reduced skeletal growth and enhanced branching are also observed under direct perturbations of the actomyosin network. Similar skeletogenic phenotypes are observed when ROCK is inhibited in a skeletogenic cell culture, indicating that these phenotypes are due to ROCK activity specifically in the skeletogenic cells. We propose that the actomyosin network was employed independently, downstream of distinct GRNs, to regulate biomineral growth and morphology across Eukaryotes.

Key words: Biomineralization, Actomyosin remodeling, Sea urchin, ROCK, Skeletogenesis, Gene regulation, Tubulogenesis

Introduction

Biomineralization is the process in which organisms from the five kingdoms of life, use minerals to produce shells, skeletons and teeth that protect and support them (1-4). Recent studies suggest that biomineralization evolved by the phylum specific co-option of ancestral gene regulatory networks (GRNs) that drove the construction of distinct organic scaffolds (4-9) and by the evolution of specialized sets of biomineralization proteins (3, 7, 10-13). This explanation is in line with the dissimilar GRNs and biomineralization proteins that drive biomineralization in different phyla (5, 12, 14). Yet, there are common cellular processes required for mineralization in highly diverse phyla (4), suggesting that distinct upstream GRNs might have recruited common cellular and molecular mechanisms to drive biomineralization.

The actomyosin network was shown to play a role in biomineralization in various Eukaryote models, from unicellular organisms to vertebrates' bones and teeth (15-22). There is a tight association between actin filaments and the biomineralization compartment in diatoms (19) and a highly dynamic actin organization that forms the biomineralization compartment in foraminifera (17). Perturbations of actin polymerization result with severely deformed shells and inhibition of shell shedding in coccolithophores (15, 16) and diatoms (18). In vertebrates, Rho GTPases and Rho-associated coiled-coil kinase (ROCK), regulate chondrocytes, osteoblasts and odontoblasts differentiation and affect genes expression in these biomineralizing cells (20-22). However, the roles of actomyosin remodeling in controlling mineral deposition and shaping biomineral morphology, are still unclear.

Sea urchin larval skeletogenesis provides an attractive model to study the role of the actomyosin network in biomineralization. Sea urchin larval skeletons are made of two frameworks of interconnected calcite rods, termed "spicules" that are generated by the skeletogenic cells (6, 23). To make the spicules, the skeletogenic cells form a ring with two lateral skeletogenic clusters and fuse through their filopodia forming a pseudopodia cable that links them into a syncytium (24, 25). The mineral is concentrated in the form of amorphous calcium carbonate (ACC) inside intracellular vesicles (26, 27). The vesicles are secreted into the biomineralization compartment generated in the

skeletogenic cell clusters forming two triradiate spicules (Fig. 7A, (26, 28)). The tubular biomineralization compartment, also called the spicule cavity, elongates within the pseudopodia cable by localized mineral deposition at the tip of each rod (29).

The GRN that controls sea urchin skeletogenesis is known in great detail and is highly similar to the GRN that controls vertebrates' vascularization, suggesting a common evolutionary origin of these two tubulogenesis programs (6, 23, 30). As the spicules elongate, the expression of key regulatory and biomineralization related genes becomes restricted to the skeletogenic cells proximal to the growing tips, possibly to promote mineral deposition at these sites (31-33). Localized gene expression is regulated by signaling cues such as the Vascular Endothelial Growth Factor (VEGF) signaling (31, 32). However, how the skeletogenic GRN drives spicule formation and localized mineral deposition at the tips of the rods is poorly understood.

Previous works suggest that the actomyosin network is essential for sea urchin skeletal growth. Calcium bearing vesicles perform an active diffusion motion in the skeletogenic cells with a diffusion length that inversely correlates with the strength and activity of the actomyosin network (34). Actin filaments are formed around the spicule (34, 35) and F-actin signal is enriched at the tips of the growing skeletal rods in skeletogenic cell culture (36). Genetic and pharmacological perturbations of the GTPase, CDC42, prevent the formation of filopodia in the skeletogenic cells and inhibit spicule formation and elongation (37). Pharmacological perturbations of ROCK prevent spicule formation (38) and genetic perturbations of Rhogap24l/2 result with ectopic spicule splitting (6). Despite these phenotypic characterizations, little is known on the regulation and role of the actomyosin network and remodeling proteins in localized mineral deposition.

Here we study the role of ROCK and the actomyosin network in the sea urchin *Paracentraorus lividus (P. lividus)*. Our findings reveal the critical role of ROCK and the actomyosin network in multiple aspects of sea urchin biomineralization, suggesting a common use of these factors in Eukaryote biomineralization, downstream of distinct GRNs.

Results

ROCK is enriched in the skeletogenic cells depending on VEGF signaling

We sought to study the spatial expression of the ROCK protein and its regulation by

VEGF signaling, a prominent regulator of sea urchin skeletogenesis (6, 30). The

sequence of ROCK and especially, its functional domains are highly conserved between

human and sea urchin (Fig. S1, (39, 40)). We therefore used a commercial antibody that

recognizes human ROCK protein to test ROCK expression in *P. lividus* embryos (Fig.

S2). We first used western blot to detect ROCK expression in control and under VEGFR

inhibition at the time of spicule formation and during skeletal elongation (~22 hours post

fertilization [hpf], 27hpf and 33hpf, Fig. S2). The antibody detected a ~150kDa protein

(Fig. S2A) that is the predicted size for *P. lividus* ROCK protein (153kDa). VEGFR

inhibition marginally increased the overall level of ROCK at 22hpf, but did not affect it at

27hpf and 33hpf (Fig. S2B). Yet, this measurement was done on proteins extracted from

whole embryos, and we wanted to study the spatial expression of ROCK and specifically,

its regulation in the skeletogenic cells.

We studied the spatial distribution of ROCK protein using ROCK antibody and the

skeletal cell marker, 6a9, that marks the skeletogenic syncytium membrane ((41), Fig. 1).

We quantified ROCK signal in the skeletogenic cells compared to the neighboring

ectodermal cells in control embryos and under VEGFR inhibition, at the three time points

mentioned above. At 22hpf, ROCK expression in the skeletogenic cells is mildly

enriched compared to the neighboring ectodermal cells and this enrichment is not

observed under VEGFR inhibition (Fig. 1A-F). At 27hpf, ROCK enrichment in the

skeletogenic cells increases, but similar enrichment is observed under VEGFR inhibition

(Fig. 1G-L). At 33hpf, ROCK enrichment in the skeletogenic cells is most apparent (Fig.

1M-O) and depends on VEGF signaling (Fig. 1P-S). At both 27hpf and 33hpf in control

embryos, ROCK seems localized near the skeletogenic cell membranes, which further

supports its activation in these cells, since ROCK activation leads to its localization to the

cell membranes in other systems (42). Overall, this data demonstrate that ROCK

expression is elevated in the skeletogenic cells and this enrichment strengthens with

skeletal elongation and depends on VEGF signaling.

ROCK activity in the skeletogenic cells controls spicule initiation, growth and

morphology

Next, we studied the role of ROCK in sea urchin skeletogenesis using genetic
perturbations. We downregulated the expression of ROCK by the injection of two
different translation morpholino anti-sense oligonucleotides (MASOs). ROCK MASO-1
matches the start of translation region, ROCK MASO-2 matches the 5' region upstream
the start of translation, and random MASO was used as a control. Embryos injected with
either ROCK MASO-1 or MASO-2 show strongly reduced skeletons (Fig. 2C, D) and
embryos injected with ROCK MASO-2 show an additional phenotype of severely
branched skeleton (Fig. 2B). The additional branching phenotype and the larger
percentage of affected embryos in ROCK MASO-2 are probably due to its higher
efficiency resulting from its more 5' location relative to translation start codon (Fig. 2E).
Thus, the genetic perturbations of ROCK expression indicate that sea urchin ROCK is
important for skeletal elongation and normal branching.

To elucidate the dependence of ROCK phenotypes on ROCK activity level and identify
its effect on different stages of skeletogenesis, we tested the skeletogenic phenotypes of
different concentrations of ROCK inhibitor, Y27632, applied before or after spicule
initiation (Fig. S3A). Y27632 binds to the ATP site of ROCK's kinase domain and
prevents its activity (43). The amino-acids to which Y27632 binds, are conserved in the
sea urchin ROCK protein, supporting its specific inhibition of this protein (Fig. S1C).
Continuous ROCK inhibition, beginning at egg fertilization, using 80µM Y27632, did
not affect skeletogenic cell migration but eliminated skeletal formation at 27hpf (Fig. SB,
C). At two days post fertilization (dpf) the majority of the embryos completely lack
skeleton and the rest show only spicule initiations, yet their overall embryonic
development seems normal (Fig. 2G, S3I, in agreement with (38)). Similar results are
observed when adding the inhibitor in this concentration at 20hpf (Fig. 2I, S3I). Yet, the
addition of 80µM of the inhibitor at 25hpf, after spicule initiation, results in a majority of
embryos with severe ectopic spicule branching, which resembles the branching
phenotype observed in ROCK MASO-2 (Fig. 2H, I). Reducing the concentration of the
added inhibitor leads to milder branching and elongation phenotypes (Fig. S3F, H, K).
However, even at the lowest inhibitor concentration, we detect a reduction of skeletal
growth rate and ectopic branching a few hours after the addition of ROCK inhibitor (Fig.
S4, 30µM). Washing the inhibitor after 25hpf results in partial recovery of

skeletogenesis, with normal or mildly delayed skeletons (Fig. S3E, K). Overall, our genetic and pharmacological perturbations indicate that ROCK activity is necessary for spicule formation, skeletal elongation and normal branching pattern.

To test if ROCK skeletogenic phenotypes are due to its activity specifically in the skeletogenic cells, we inhibited ROCK in a culture of isolated skeletogenic cells (Fig. 2J-N). The addition of ROCK inhibitor to the skeletogenic cell culture at 16hpf, before the spicules form, completely abolished spicule formation (Fig. 2K). The addition of the inhibitor at 48hpf, after spicule formation, resulted with significantly shorter spicules (Fig. 2L, N) and increased branching compared to control spicules (Fig. 2M). The reduction of the spicule length becomes more significant with increasing concentration of the ROCK inhibitor (Fig. 2N). Overall, ROCK skeletogenic phenotypes in the isolated skeletogenic cell culture are similar to its phenotypes in whole embryos, verifying that it's specific activity in the skeletogenic cells is essential for spicule formation, normal elongation and prevention of branching.

ROCK inhibition reduces spicule volume, surface area and total length, but not thickness (SR-µCT)

To quantify the effect of ROCK inhibition on mineral deposition and skeletal shape we used synchrotron radiation micro-computed tomography (SR-µCT) that enables three-dimensional visualization and quantification of spicule morphology at 2 and 3dpf (Figs. 3, S5, see methods for details). The aberrant morphology of ROCK inhibited spicules is visible at both developmental time points (Fig. 3A). The ectopic branching occurs both at the tips where tip-splitting is detected (arrowheads, Fig. 3A), as well as in the back, where small spicule extensions, or mineral 'dripping' are observed (arrow at 48hpf, Fig. 3A). This implies that the regulation of mineral deposition is perturbed both at the tip and in the back of the spicules under ROCK inhibition.

When compared to control spicules, ROCK inhibition leads to a ~2.5-fold reduction of the spicule volume and surface area in both 2 and 3dpf (Fig. 3B-D, see Table S2 for measured values and standard deviations and S3 for statistical analyses). These SR-µCT measurements show that the rate of mineral deposition is significantly reduced under ROCK inhibition.

To understand the specific effect of ROCK inhibition on spicule elongation vs. thickening, we compared the total length and the mean thickness of the spicules between control and ROCK inhibition (Fig. 3E-G). The total length of control spicules was on average about 2.5-times longer than the length under ROCK inhibition in both 2dpf and 3dpf (Fig. 4E, F, tables S2 and S3). In contrast to its effect on the spicule length, ROCK inhibition caused a minor increase of the spicule thickness at 2dpf and at 3dpf the thickness was not affected by the inhibition (Fig. 3E, G, Tables S2, S3). These SR-μCT measurements indicate that ROCK activity plays a crucial role in controlling the rate of mineral deposition at the tips of the rods and in spicule elongation, and is not affecting spicule thickening.

ROCK activity does not control mineral uptake but is required for "tip-dominance"

To test the effect of ROCK inhibition on mineral intake, we used the fluorescent chromophore, calcein that binds to divalent ions including Ca^{+2} and is widely used to track the calcium pathway in biomineralization (6, 28, 33, 34). We measured the number of calcein stained vesicles per area as well as the number of pixels of calcein signal per area in the skeletogenic cells, in control and under ROCK inhibition at 1dpf and 2dpf (Fig. S6A-E). Continuous ROCK inhibition does not change the number of calcium bearing vesicles at 1dpf, but the number of calcein stained pixels significantly increases in this condition, possibly indicting that the vesicle volume is larger (Fig. S6F, G). At 2dpf, the number of calcein stained pixels is similar in control and in continuous ROCK inhibition, suggesting that the overall calcium content does not change in this condition (Fig. S6H, I). The number of calcein stained vesicles is however decreased under ROCK inhibition, possibly indicating that there are fewer vesicles with larger volume. Addition of ROCK inhibitor at 25hpf affects neither the number of calcein stained vesicles nor the number of calcein stained pixels in the skeletogenic cells. Together, these measurements show that ROCK is not required for the uptake of calcium into the cells. Therefore, the skeletal phenotypes occurring under ROCK inhibition are not related to a decrease in calcium uptake or shortage in intracellular calcium, but are due to defects in mineral deposition.

To monitor the effect of ROCK inhibition on mineral deposition rate and distribution, we

applied two pulses of different calcein dyes: blue calcein was added at 29hpf and washed

after one hour, followed by green calcein added at 44hpf and washed after one hour.

Hence, blue calcein marks the initial triradiate region and green calcein marks the edges

of the spicules that were recently deposited (blue and green arrows in Fig. 3H, I). The dye

FM4-64 was used to mark the cell membranes. Under ROCK inhibition, the green

labeled region is much closer to the blue labeled region compared to control, due to the

reduction in skeletal elongation rate, in agreement with our SR-μCT measurements.

However, while in control embryos each body rod has a single tip stained in green

calcein, under ROCK inhibition one of the body rods has two tips and both are stained in

green. This indicates that mineral is being deposited in both edges, and the mechanism

that prevents the growth of multiple tips in each rod, enabling "tip-dominance", (in

analogy to plant stem apical dominance), is disrupted under ROCK inhibition.

231 The activity of the actomyosin network is essential for normal spicule elongation

232 and inhibition of ectopic branching

In other systems, major roles of ROCK are to control F-actin polymerization and MyoII

activation (44-48), hence we wanted to directly test the role of these actomyosin

components in sea urchin skeletogenesis. To directly inhibit F-actin polymerization we

used Latrunculin-A (Lat-A), that prevents the polymerization of actin monomers (49) and

was shown to be effective in sea urchin embryos (50). To directly inhibit actomyosin

contractility we used Blebbistatin (Blebb), that interferes with MyoII ATPase activity,

prevents actin-myosin interactions (51) and was shown to be effective in sea urchin

embryos (52). To prevent interference with the early cell divisions where the actomyosin

network plays a critical role (53-55), we added the inhibitors, individually or together,

before or after spicule formation (20hpf and 25hpf, see methods for details). We tested

the resulting phenotypes at 2dpf (Fig. 4A-G).

Our results indicate that F-actin polymerization and to a lesser extent, actomyosin

contractility, are essential for normal skeletal growth and inhibition of ectopic branching.

The majority of the embryos where MyoII activation was inhibited show mild ectopic

branching mostly at the tips of the rods, and a minor delay in skeletal growth (Fig. 4B, C,

G). The majority of the embryos where F-actin formation was inhibited show significant

reduction of skeletal growth and severe ectopic skeletal branching (Fig. 4E, F and similar

phenotypes to Fig. 4D, statistics in Fig. 4G). In some embryos, additional spicules form

(Fig. 4F, G). The co-inhibition of F-actin polymerization and MyoII activity results in

skeletogenic phenotypes that resemble, but are slightly stronger, than those of the

inhibition of F-actin alone (Fig. 4D, G).

To test whether actin polymerization and myosin activity specifically in the skeletogenic

cells underlies the observed skeletogenic phenotypes, we inhibit both in an isolated

skeletogenic cell culture (Fig. 4H-K). We added the inhibitors after the spicule formed

(>48hpf) and observed the phenotypes at 3dpf. Both inhibitors reduced skeletal growth,

with Lantruculin-A having a significantly stronger effect than Blebbistatin (Fig. 4K), in

agreement with the effect of these inhibitors in whole embryos. However, differently than

in whole embryos, skeletal branching in both Lantruculin-A and Blebbistatin is not

affected and is similar to the observed branching in control cultures. The absence of the

branching phenotype in the skeletogenic cell culture could be due to the increased rigidity

of the substrate that could compensate for the reduce actomyosin activity. Another option

is that the branching phenotype in whole embryos could be due to the reduction of the

activity of the actomyosin network in non-skeletogenic cells. Nevertheless, these results

indicate that normal skeletal growth depends on F-actin polymerization and myosin

contractility in the skeletogenic cells.

ROCK activity is required for F-actin organization around the forming spicule

Next, we tested whether ROCK regulates the actomyosin structure and activity during

skeletal formation. To accomplish this we used phalloidin, a fluorophore that binds F-

actin specifically, an antibody that detects phosphorylated MyoII (MyoIIP, (34)) and the

skeletogenic marker, 6a9 (Fig. 5). In control embryos, F-actin is detected around the tri-

radiate calcite spicules at 27hpf (green arrow in Fig. 5B, in agreement with (34, 35)). The

pseudopodia cable that connects the skeletogenic cells shows lower F-actin signal

compared to the signal around the spicule (blue line in Fig. 5B, C, E blue arrows in Fig.

S7B, C, E). MyoIIP signal is not enriched in the skeletogenic cells nor in the spicule (Fig.

5D, S7D). Under continuous ROCK inhibition, the pseudopodia that connects the

skeletogenic cells still forms (Fig. S7H) and F-actin accumulation is still observed in the

lateral skeletogenic cell clusters; but unlike control embryos, F-actin is not organized and

the spicule cavity is not formed (Fig. 5G). MyoIIP signal seems unchanged by ROCK

inhibition (Fig. 5I, S7I), in agreement with the weak and late phenotype of MyoII

inhibition (Figs. 4). These findings suggest that ROCK activity is essential for F-actin

organization around the spicules, but not for the formation of the pseudopodia cable.

F-actin is enriched at the tips of the elongating spicules independently of ROCK

activity

To assess the role of ROCK in actomyosin organization during skeletal elongation we

compared F-actin and myoIIP signals between control embryos and embryos where

ROCK inhibitor was added at 25hpf (Fig. 5K-T, S7). At the prism stage, F-actin is

enriched at the tips of the spicules in both control and ROCK inhibited embryos (white

arrowheads in Figs. 5L, O). We quantified the F-actin signal at the tips compared to the

back in both treatments (Fig. S7K, L, see methods for details). The phalloidin signal is on

average 3-fold stronger at the tips of the spicules compared to the back, in both control

and ROCK inhibition, indicating that F-actin enrichment at the tips is significant and

independent of ROCK activity. In both control and ROCK inhibited embryos, the F-actin

signal was markedly higher in the regions of the pseudopodia cable where the spicule

cavity had formed, compared to the regions where the spicule cavity had not yet formed

(blue arrows in Fig. 5L-O, Q-T). We did not detect an enhanced signal of MyoIIP around

the spicules or in the skeletogenic cells, in either condition (Fig. 5N, S). At the pluteus

stage, F-actin is enriched at the tips of the spicules in both control and ROCK inhibited

embryos (white arrowheads in Fig. S7N-Q, S-V). Some non-skeletogenic cells that are

enriched with MyoIIP signal are detected at this time (red arrowheads in Fig. S7N-P).

Together these data demonstrate that F-actin coats the spicule cavity and is enriched at

the tip of the rods, independently of ROCK activity.

We tested the effect of ROCK inhibition on F-actin in skeletogenic cell cultures. F-actin

is detected at the tips, in both control cultures and cultures where ROCK was inhibited

after spicule formation (Fig. 5U, V). Yet, under ROCK inhibition, branching is enhanced

and F-actin is enriched at the splitting tips of the spicule rods (Fig, 5V). The enrichment

of F-actin at the splitting tips of the spicule under ROCK inhibition in skeletogenic cell

culture resembles the calcein staining at the two tips in the embryo in this condition (Fig.

3I). These observation further supports the role of ROCK activity in regulating tip-

dominance during sea urchin skeletogenesis.

ROCK activity is essential for normal skeletogenic gene expression

The role of ROCK in regulating gene expression during vertebrates' chondrocytes,

osteoblasts and odontoblasts differentiation (20-22) intrigued us to study the role of

ROCK in regulating skeletogenic gene expression in the sea urchin embryo. ROCK

continuous inhibition significantly downregulates the expression of multiple skeletogenic

genes, including the cytoskeleton remodeling gene, *rhogap24l/2*, the biomineralization

genes, *caral7, scl26a5* and *SM30* at 27hpf and 2dpf (Fig. 6A, (6)). ROCK inhibition did

not affect the expression level of VEGFR and ROCK itself at these times (Fig. 6A).

Washing the inhibitor after 25hpf or adding the inhibitor at 25hpf had a weaker effect, but

still led to downregulation of *caral7*, *SM30*, *angio1* and *notch1* (Fig. 6B). Thus, ROCK

activity is required for the normal expression level of multiple skeletogenic genes during

sea urchin skeletogenesis.

Both continuous and late ROCK inhibition, strongly affect the spatial expression of key

regulatory and biomineralization genes that are normally expressed differentially within

the skeletogenic lineage at the pluteus stage (Figs. 6D-F, S8). These include, the VEGF

receptor (VEGFR), the biomineralization genes *SM50* and *SM30,* skeletogenic

transcription factors, Ets1, Alx1 and Erg1 that are essential for skeletogenic cell

specification and biomineralization (23, 56) and the transcription factors Hex and

MyoD1. In normal plutei these genes are differentially expressed within skeletogenic

lineage, but in both continuous and late ROCK inhibition the expression of these genes

becomes localized to the two lateral skeletogenic cell clusters and their vicinity (Fig. 6D-

F, S8). The lateral skeletogenic cells clusters are the cells where these genes are

expressed in the gastrula stage (24-27hpf) and where spicule formation begins in normal

embryos (6, 32). The expression of VEGFR and SM50 under ROCK inhibition is more

expanded than the other tested genes and is also observed at the anterolateral rods (Fig.

6B, C).The VEGF ligand that is normally expressed in the ectodermal cells near the four

tips of the post-oral and anterolateral rods (32), is expressed at four ectodermal domains

near the lateral skeletogenic cell clusters and near the end of the anterolateral chain,

under ROCK inhibition (Fig. S8G).

Overall, under ROCK inhibition, the expression of key regulatory and biomineralization

genes is downregulated (Fig. 6A) and remains localized to the area near the lateral

skeletogenic cells clusters (Fig. 6D-F, S8). This localization is despite the proper

formation of the dorsal, anterolateral and mid-ventral chains (Fig. 6C). This implies that

ROCK activity provides regulatory cues to the skeletogenic GRN required for the

dynamic progression of gene expression within the skeletogenic lineage.

Discussion

Biomineralization is a complex morphogenetic process, encoded in the species genome

and executed by the GRNs and the cellular machinery. Here we used the sea urchin larval

skeletogenesis to investigate the role of ROCK and the actomyosin network in biomineral

formation and in regulating gene expression, as we summarize in Fig. 7. ROCK

expression is enriched in the skeletogenic cells, partially depending on VEGF signaling

(Fig. 1). ROCK activity in the skeletogenic cells is necessary for spicule formation,

skeletal elongation and prevention of ectopic skeletal branching (Figs. 2-3, S3, 4). Direct

inhibition of F-actin polymerization results in similar but not identical skeletogenic

phenotypes (Fig. 4) and F-actin is organized around the spicule cavity and enriched at the

spicule tips (Figs. 5, S7). ROCK activity feeds back into the skeletogenic GRN and

affects regulatory and biomineralization gene expression (Fig. 6). Below we discuss our

findings and their implications for the biological control and evolution of

biomineralization.

The first step in generating the sea urchin spicules is the construction of the spicule

cavity, a membrane filled with calcium carbonate and coated with F-actin (Fig. 7A).

ROCK activity is necessary for the formation of the spicule cavity and for the

organization of F-actin around it (Figs. 2, 5). A direct perturbation of F-actin

polymerization strongly disrupts spicule growth and shape (Fig. 4). F-actin is enriched in

the regions of the pseudopodia cable that are filled with mineral, supporting its role in

mineral deposition (Figs. 5, S7, Fig. 7A). Overall, our studies indicate that ROCK

activity in the skeletogenic cells is essential for the formation of the spicule cavity and for
the initial F-actin organization around it, which could be essential for mineral deposition
into it (Fig. 7C).

After the spicule cavity has formed, the spicules elongate inside the pseudopodia chord
by mineral deposition at the tips of the spicule rods, which depends on ROCK activity
and on F-actin polymerization (Figs. 2-4, S7, 7D-F). F-actin is detected along the spicules
and is significantly enriched at the tips of the elongating spicules (Fig. 4, S7 (57)). F-actin
enrichment at the tips is independent on ROCK activity (Fig. 5K-T, S7K, F), yet under
ROCK inhibition and under the inhibition of F-actin polymerization, ectopic branching is
observed at the tips and in the back (tip splitting and spicule dripping, Figs. 2J, N, 3A, 4,
5V). The participation of F-actin and ROCK in polarized growth and vesicle exocytosis
has been observed in both animals and plants (57, 58). F-actin is used for short range
myosin trafficking of vesicles toward the secretion site (57, 59). Once the secretory
vesicle fuses with the membrane, it is coated with F-actin that stabilizes the vesicle and
enables MyoII binding and ROCK-regulated contraction, necessary for content secretion
(60, 61). In plant cells, F-actin accumulate at the growing tip of the cell and assists
vesicle exocytosis necessary for polarized tip-growth (58). Our observations suggest that
ROCK, F-actin and MyoII could be regulating the localized exocytosis of mineral-
bearing vesicles during sea urchin skeletogenesis. The delay in spicule growth and the
ectopic branching could be due to impaired and misplaced vesicle trafficking and
deposition under ROCK and F-actin inhibition, but the exact protein regulatory network
is still to be determined.

ROCK activity is necessary for correct spatiotemporal expression of regulatory and
biomineralization genes (Fig. 6). ROCK inhibition downregulates the expression level of
skeletogenic genes (Fig. 6A) and prevents the spatial progression of key skeletogenic
genes toward the edges of the skeletal rods (Fig. 7E). In other systems, ROCK activity
was shown to regulate gene expression through various intracellular pathways (62-64).
Furthermore, ROCK has an important role in mechano-sensing of matrix rigidity and
transducing it into changes in gene expression (65). Relatedly, stiff matrix activates
ROCK in pre-osteoblastic cells, which activates Erk that leads to osteogenic
differentiation and upregulation of osteogenic gene expression (64). Sea urchin ROCK

could be sensing the spicule stiffness and transducing it into gene expression, providing a necessary feedback between spicule elongation and the skeletogenic GRN. Further studies are required to identify the specific molecular mechanisms activated by ROCK to affect gene expression in the skeletogenic cells.

Overall, our findings and the role of actomyosin remodeling in biomineralization, from calcifying and silicifying single cells to vertebrates' bones and teeth (15-22, 64), suggest that this molecular machinery is a part of the common molecular tool-kit of biomineralization. Most likely, the actomyosin network was employed independently, downstream of distinct GRNs across Eukaryotes, to control mineral growth and morphogenesis.

Methods

Animal and embryos

Adult *Paracentrotous lividus* were obtained from the Institute of Oceanographic and Limnological Research (IOLR) in Eilat, Israel. Eggs and sperm were obtained by injection of 0.5M KCl solution into the adult sea urchin species. Embryos were cultured at 18°C in 0.2μ filtered ASW.

Imaging

Embryonic phenotypes and WMISH were imaged by Zeiss Axioimager 2. Fluorescent markers were imaged using Nikon A1-R Confocal microscope. All images were aligned in Adobe photoshop and Illustrator.

Axitinib (AG013736) treatment

Axitinib (AG013736, Selleckchem, Houston, TX, USA) was applied as described in (32)

Western Blot

Embryo cultures treated with 150 nM Axitinib, or DMSO as control, were collected at 22hpf, 27hpf and 33hpf by centrifugation of 10,000g for 2 minutes. We lysed the pellet in 150μL lysis buffer (20 mM Tris-HCl, 150 mM NaCl, 1% Triton X-100, pH 8) with protease inhibitors (protease inhibitor cocktail w/o metal chelator; Sigma, P8340; 1:100) and phosphatase inhibitors (50 mM NaF, 1 mM Na3VO4, 1 mM Na4P2O7, 1mM, βGP)

as described in (66). 40µg protein were loaded on 8% SDS-acrylamide gel, transferred to

PVDF membranes for Western blot analysis. Anti-ROCK antibody (ab45171, Abcam)

was used in a 1:300 dilution followed by anti-rabbit HRP secondary antibody diluted to

1:2500 (Jackson ImmunoResearch 111-035-003). For the loading control, the membrane

was incubated with anti-β-tubulin antibody (1:5000; Sigma) followed by anti-mouse

antibody 1:5000 (Jackson ImmunoResearch 115-035-003). Quantification of ROCK

signal was done using Image studio lite vr. 5.2. ROCK signal was divided by Tubulin

signal and then the ratio between ROCK/Tubulin in control vs. VEGFR inhibition was

calculated. The graph in Fig. S2 was generated based on three biological replicates

(different sets of parents) at 27hpf and 33hpdf and four biological replicates at 22hpf.

Immunostaining procedure

Phalloidin labeling, MyoIIP (p20-MyosinII), ROCK (anti ROCK2+ROCK1 antibody

[EP786Y], AB-ab45171, Abcam), and 6a9 immunostaining were done similarly to (34).

Quantification of Anti-ROCK and Phalloidin signal

A graphical program was written to allow manual quantification of the fluorescent signal

from the spicule regions by identifying "stained" pixels per selected area. The ratio of

stained anti-ROCK to region total area was compared between ectodermal and

skeletogenic cells, and between control and VEGFR inhibited embryos at 22, 27 and

33hpf. The average ratios and ztest for significance difference from 1, are 22hpf control,

~1.3, z=0.0006; VEGFR inhibition ~1, z=0.5, 27hpf control ~1.8, z<10-11; VEGFR

inhibition ~1.6, z<10-8, 33hpf control ~4.6, z<10-23; VEGFR inhibition ~1.5, z<10-5.

For phalloidin analyses, embryos at 33hpf of control and ROCK inhibitor addition at

25hpf were used and for each image, two regions were selected: an area at the tip of the

spicule and an area at the back of the spicule, about ~10 microns apart. The ratio of

stained phalloidin to region total area was compared between tip and back, in control and

ROCK inhibited embryos. Paired t-test was used to compare the differences in phalloidin

signal between the tip and back of the spicules across all groups and between control and

ROCK inhibited embryos.

ROCK MASOs injections

Microinjections were conducted similarly to (32). The eggs were injected with an

injection solution containing 0.12M KCl, 0.5µg/µl Rhodamine Dextran and 900µM of

MASOs. Translation MASOs were designed and synthesized by Gene Tools, Philomath,

OR, according to ROCK sequence (39). The sequence of ROCK MASO-1 (5'-

AGACATATTTGGAGCCGA[CAT]CCTG-3') matches the start of translation region

and ROCK MASO-2 (5'-TCTCTTGCGTTATATTCCACTAAGT-3') matches the 5'

region upstream of the start of translation (39). Control embryos were injected in the

same concentration with Random MASO. Injected embryos were cultured at 18°C,

imaged and scored at 2dpf.

ROCK inhibitor (Y27632) treatment

Y27632 stock (10005583 CAS Regisrty No. 129830-38-2, Cayman chemical), was

prepared by reconstituting the chemical in DMSO. The embryos were treated with

Y27632 at a final concentration between 30-80µM, as mentioned in the results.

Throughout the paper, control embryos were cultured in DMSO at the same volume as

Y27632 solution and no more than 0.1% (v/v).

ROCK inhibition in isolated skeletogenic cell culture

Skeletogenic cell culture was performed as described in (36) with minor changes.

Isolated skeletogenic cells were cultured in CultureSlides (4 chambers polystyrene vessel

tissue culture treated glass slide REF 354104) with ASW (Red Sea Fish LTD) containing

Gentamicin and Penicillin-Streptomycin (GPS) at 18°C. Media containing 4% (v/v) horse

serum (Sigma-Aldrich H1270) in ASW+GPS was added to the culture. ROCK inhibitor,

Y27632, was added to the cell culture at 16hpf (before spicule initiation) or 48hpf (after

spicule initiation), and images were taken at 72hpf. Each treatment was conducted in

three biological repeats except from the >48hpf addition of 120µM which was done in

two biological repeats.

Quantification of skeletal length and statistical analysis

Skeletal length measurement was done as described in (32). The measurements were

repeated for three biological repeats for control, ROCK inhibition with 30µM and 80µM,

Latrunculin-A 2nM, Blebbistatin 2µM, and 2 biological repeats for 120µM. In ROCK

inhibition experiments, a total of 275 skeletons were measured for control, 105 skeletons

for 30μM, 122 skeletons for 80μM and 80 for 120μM. In Latrunculin-A and Blebbistatin

experiments, a total of 116 skeletons were measured for control, 149 for Latrunculin-A

and 107 for Blebbistatin. The data were analyzed in Excel and the statistical analysis was

performed using Kruskal-Wallis non-parametric test in Excel as described in (67) and

SPSS statistics 27.

Spicule preparation for micro-computed tomography (SR-μCT) measurement

Control embryos and embryos where 40μM Y27632 was added at 25hpf, were collected

by centrifugation of 50mL Falcon tubes at 2000rpm, at 48h and 72hpf. Every tube was

washed in 10 mL cold distilled water (DW), transferred into epi-tube and washed three

times in 1mL cold DW (incubation at 4°C between every wash, centrifuge at 2000rpm in

room temperature for 2min). Then the skeletons were washed three times in 3% NaOCl

followed by three washes in DW, one wash in 70% EtOH and one wash in Acetone.

Skeletal spicules were dried at room temperature and stored at -20 °C.

Synchrotron and lab-based SR-μCT

Dry and loose skeletal spicules prepared as described above, were used for the SR-μCT

analysis. We analyzed 14 sets of spicules from three (control) and four (ROCK inhibited)

pairs of parents, divided into four groups: 2dpf, control, n=44; 2dpf ROCK inhibited,

n=93; 3dpf control, n=51; 3dpf ROCK inhibited, n=93, see Table S1). Spicules were

glued to sharpened toothpick tips, either in bulk or individually when picked by hand by

means of a brush bristle. Each toothpick loaded with spicules, was mounted on a 3ml vial

twist-off lid and for transportation securely stored in the vial (Fig. S5A, B). For each set

2-3 μCT-samples were prepared, which allowed the acquisition of statistically relevant

sample sizes (40+ spicules/group, Table S1). Tomographic data sets were acquired using

the following scanning parameters; for the lab based μCT (RX Solutions, Chavanod,

France) 90kV, 143-160μA, 1440 images, average frames 10, frame rates 0.4-1, voxel

sizes: 0.45-59μm; and for the synchrotron radiation μCT (Anatomix, Synchrotron

SOLEIL, Gif-sur-Yvette, France) 2048 images, angle between projections 0.09°, voxel

size: 0.65μm. Tomographic data reconstruction was performed with commercial (RX

Solutions, Chavanod, France) and PyHST2 software (A. Mirone et al., Nucl. Instrum. Meth. B 324 (2014) 41-48, doi:10.1016/j.nimb.2013.09.030) at the MPICI and at Anatomix, SOLEIL, respectively.

Three-dimensional (3D) data analysis

Data visualization, pre-processing including cropping and image type conversion to 16bit (of the Sr- µCT data), and user-augmented segmentation of intact spicules were performed in Amira using off-the-shelf modules (Fig. S5C-E, (68)). Volume and area measurements were performed on the segmented spicules, *i.e.* the label-fields using Amira's material.statistics module (Volume3D and Area3D). Average spicule thickness and total length measurements were performed with python code, utilizing available python libraries (*e.g.*, skimage, skan, scipy). The code was applied to 3D-tif files created from Amira's label-fields, which contained the segmented spicules. For length measurements, the spicules were skeletonized and the chord lengths of the branches forming a spicule-skeleton were summed up to obtain the total spicule length. For thickness measurements, we performed Euclidean distance transforms for each spicule using scipy's ndimage.morphology.distance_transform_edt package, and calculated the average spicule thickness based on the distance measurements obtained at each voxel from spicule-skeleton to the nearest spicule surface. Quality control of our measurements was performed with open source software, Fiji and available plugins (e.g. BoneJ, FiberJ, DiameterJ).

Calcein staining

A 2mg/ml stock solution of blue calcein (M1255, Sigma) and a 2mg/ml stock solution of green calcein (C0875, Sigma) were prepared by dissolving the chemicals in distilled water. Working solution of 250μg/ml and 150μg/ml, respectively, was prepared by diluting the stock solution in artificial sea water. Blue calcein was added to the embryo culture at 29hpf for one hour and then washed. At 44hpf the green calcein was added for one hour and washed. For FM4-64 staining A 100µg/ml stock solution of FM4-64 (T13320, Life technologies, OR, USA) was prepared by dissolving the chemical in distilled water. Working solution of 2.5 µg/ml was prepared by diluting the stock solution

543 in artificial sea water. Embryos were immersed in working solution about 10 minutes

544 before visualization.

Latrunculin-A and Blebbistatin treatments

1mM of Lat-A stock (L12370, Thermo Fisher Scientific) was prepared by reconstituting

the chemical in DMSO. The embryos were treated with Lat-A at final concentration of 2

nM. 1mM of Blebbistatin stock (13013-5, Enco) was prepared by reconstituting the

chemical in DMSO. The embryos were treated with Blebbistatin at final concentration of

2 µM. For the double inhibition experiments, embryos were treated with Blebb at final

concentration of 1.5 µM, and with Lat-A at final concentration of 2 nM.

WMISH procedure

553 WMISH was performed as described in (32). List of the primers used to make the

554 WMISH probes is available in (32).

cDNA preparation for QPCR experiments

For ROCK inhibition experiments total RNA was extracted from >1000 sea urchin

embryos in each condition using RNeasy Mini Kit (50) from QIAGEN (#74104)

according to the kits' protocol. DNase treatment on column was done using RNase-Free

DNase Set-Qiagen (50) (#79254). RNA was reverse-transcribed using the High Capacity

cDNA RT kit, AB-4368814 (Applied Biosystems) following the manufacturer's

instructions.

Quantitative polymerase chain reaction (QPCR)

563 QPCR was carried out as described in (32). Complete list of primer sequences used for

564 the experiments in provided there.

Acknowledgments

566 We are grateful to SOLEIL for provision of synchrotron radiation facilities and we would

567 like to thank Dr. Mario Scheel, Dr. Timm Weitkamp and Dr. Jonathan Perrin for

568 assistance in using beamline ANATOMIX for nano and µCT. We thank the Bioimaging

569 Unit, Faculty of Natural Sciences, University of Haifa for assistance with the use of

confocal. We thank Charles Ettensohn for the generous gift of the 6a9 antibody produced in his lab. We thank David Ben-Ezra and Michael Kantorovitz for their help with sea urchin handling. We thank Haguy Wolfenson for illuminating discussions of the results and conceptual framework. We thank Nir Sapir and Tovah Nehrer for their advice and help with the statistical analysis of spicule length. We thank Alvaro Israel and Omri Nahor for their help with sea urchin maintenance. We thank Yulia Kagan for her help with the western blot.

Funding: Israel Science Foundation grant number 211/20 (S.B.D.), and Zuckerman fellowship (M.R.W.), TUD internal funding (YP). ANATOMIX is an Equipment of Excellence (EQUIPEX) funded by the Investments for the Future program of the French National Research Agency (ANR), project NanoimagesX, grant no. ANR-11-EQPX-0031. Access to ANATOMIX beamline is granted for proposal number proposal 20200811.

Figures

Figure 1 ROCK enrichment in the skeletogenic cells increases with time and depends on VEGF signaling. A-R, ROCK immunostaining at different times points in control and under VEGFR inhibition. In each condition and time point, the left image shows ROCK immunostaining and the skeletogenic cell marker, 6a9 (A, D, G, J, M and P). The middle image shows ROCK immunostaining alone in the whole embryo (B, E, H, K, N and Q). The right image shows the enlargement of the white rectangle region of the middle image (C, F, I, L, O, R). Scale bar in whole embryo is 50µm and in enlargements is 10µm. E – ectoderm, SC – skeletogenic cells. S, quantification of the ratio between ROCK signal/area in the skeletogenic cells compared to the ectodermal cells (see methods for details). Each box plot shows the average marked in x, the median, the first and the third quartiles (edges of boxes) and all experimental measurement (dots). Experiments were performed in three independent biological replicates and in each condition at least 33 embryos were measured. Statistical significance was measured using paired 2-tailed t-test where, * indicates P<0.05, ** indicates P<0.005, and *** indicates P<0.0005.

Figure 2 ROCK activity is essential for spicule formation, normal elongation and branching in whole embryos and in skeletogenic cultures. A-E, genetic perturbation of ROCK translation using two different MASOs observed at 2dpf. A, control embryo injected with Random MASO. B, embryo injected with ROCK MO-2 shows ectopic spicule branching. C, D, embryos injected with ROCK MO-1 or MO-2 show spicule initiations. E, summary of MASO injection phenotypes based on 4-6 independent biological replicates. F-I, pharmacological perturbations of ROCK activity using 80µM of the inhibitor Y27632 observed at 2dpf. F, representative control embryo with normal skeletal rods, B, body; AL, anterolateral; PO, post-oral and MV, midventral. G, complete skeletal loss in embryo treated continuously with ROCK inhibitor. J, reduced skeletal growth and enhanced ectopic branching in embryo where ROCK inhibitor was added at 25hpf. I, summary of perturbation phenotypes based on 3-8 biological replicates for each treatment. See additional phenotypes and summary in Fig. S3 and Table S1. J-M, representative spicules from skeletogenic cell cultures in control and under 30µM Y27632 at 72hpf. J, linear spicule in control culture, K, Y27632 addition at 16hpf, before spicule initiation, completely blocks spiculogenesis. L, Y27632 addition after spicule initiation, at 48hpf, reduces spicule elongation, M, and enhances branching, N, Quantification of spicule length in control and ROCK inhibition (>48hpf) at 72hpf. *P < 0.05, **P < 0.001, Kruskal-Wallis non-parametric test. Results are based on three biological repeats for each treatment, except from 120µm that was done in two biological repeats. Scale bars are 50µm. In J, L, M, the numbers at the bottom indicate the number of spicules that show this phenotypes (left) over all observed spicules (right).

Figure 3 SR-µCT studies of ROCK inhibited spicuels show reduction in skeletal volume, surface area and total length, but not thickness and 2-pulse calcein shows loss of tip-dominance. A, Exemplary 3D-renderings of control spicules (top, blue) and spicules where 40µM of ROCK inhibitor was added at 25hpf (bottom, orange), dissected at 48hpf and 72hpf. Arrowheads point to tip splitting and arrow at 48hpf point to spicule dripping at the back. B, Spicule volume vs. area for control and ROCK-inhibited spicules at 48 and 72 hpf. Each data point represents a single spicule. C-D, Frequency distributions for volume and surface area of control and ROCK-inhibited spicules at 48 hpf and 72 hpf (left and right violin plots, respectively). E, Spicules' total branch length and average thickness for control and ROCK-inhibited spicules at 48 and 72 hpf. F-G, Frequency distributions for spicule lengths and thickness of control and ROCK-inhibited spicules dissected at 48hpf and 72hpf. H, I, Calcein two-pulses experiment. Embryos were exposed to blue calcein from 29-30hpf, to green calcein from 44-45hpf and stained with FM4-64 membrane marker (red) prior to image acquisition at 48hpf. H, Control embryo, G, Embryo where 30µM of Y27632 was added at 25hpf. The experiments were done in three biological replicates and the numbers at the bottom indicate the number of embryos that show this phenotype out of all embryos scored.

Figure 4 Actin polymerization and myosin activation affect skeletal growth and branching. A-F, representative embryos showing the effect of actomyosin perturbations at 2df. A, control embryo B, embryo treated with skeletogenic cell culture were, C, embryo treated with 2µM Blebbistatin >20hpf, D, embryo treated with 2nM Latrunculin-A and 1.5µM Blebbistatin after 25hpf, E, embryo treated with 2nM Latrunculin-A after 20hpf, F, embryo treated with 2nM Latrunculin-A after 25hpf, arrow pointing to the additional spicule rod. G, Statistics of Latrunculin-A and Blebbistatin phenotypes, color code of phenotype is indicated in the representative images. Error bars indicate standard deviation. All treatments were conducted in at least three biological replicates and exact number of replicates and scored embryos are provided in table S4. H-J, representative spicules recorded at 72hpf from H, control skeletogenic cell culture, I, skeletogenic cell culture were 2nM Latrunculin-A was added at 48hpf and J, skeletogenic cell culture were 2µM Blebbistatin was added at 48hpf. K, Quantification of spicule length in the different treatments at 72hpf **P < 0.001, Kruskal-Wallis non-parametric test. Results are based on three biological repeats for each treatment. Scale bars are 50µm.

Figure 5 ROCK inhibition effect on F-actin organization and MyoII activity. A-T, Representative images showing normal embryos and embryos treated with 80µM ROCK inhibitor at different times. Right panels show enlargements of the rectangle sections marked in the left most panels. Phalloidin (green) marks F-actin, MyoIIP (red) marks phosphorylated MyoII and 6a9 (blue) marks the skeletogenic cells. A-E, Representative images at 27hpf showing control embryo and F-J, embryos treated with ROCK inhibitor from fertilization. Green arrows point to regions that show enriched F-actin signal. K-T, representative images at 33hpf, showing normal embryo K-O, and P-T, embryos where 30µM ROCK inhibitor was added at 25hpf. White arrowheads point to the enriched F-actin signal at the tips. Blue arrows point to the region of the pseudopodia cable that is not filled with the spicule cavity. The experiments were done in 3 biological replicates, the numbers at the bottom left of (A, F, K, P) indicate the number of embryos that show this phenotype out of all embryos scored. Scale bar in A, F, K, P is 50µm, and in B, G, L, Q is 10µm. U, V, representative spicules out of three biological replicates from control skeletogenic cell culture (n=60), U, and skeletogenic cell cultures that was treated with 30µM ROCK inhibitor was added at 48hpf (n=52), recorded at 72hpf. Left panel is phase image, middle panel is

phalloidin staining and right panel shows the overlay. Green arrows point to the enhanced F-actin
signal at the tips. Scale bar is 20μm.

Figure 6 ROCK activity is essential for normal gene expression in the skeletogenic cells. A,
B, the effect of 80μM ROCK inhibition on gene expression. A, continuous ROCK inhibition at
27hpf and 50 hpf (n=4). B, addition of ROCK inhibitor at 25hpf and the wash of ROCK inhibitor
at 25hpf, measured at 50hpf (n=3). Asterisks indicate p<0.001, one tailed z-test. Error bars show
standard deviation. C-F, Representative images of control embryo (top panels), embryos where
ROCK inhibitor was added at 25hpf (middle panels), and embryos that were exposed to
continuous ROCK inhibition (bottom panels), at the pluteus stage (~48hpf). C, skeletogenic cell
marker, 6a9. MV, midventral; AL, anterolateral, and PO, Post-oral rods, D, dorsal skeletogenic
chain. D-F, spatial expression of skeletogenic genes. Gene names are indicated at the top of each
panel. Numbers at the bottom of each image indicate the number of embryos that show this
phenotype (left) out of all embryos scored (right), conducted in at least three independent
biological replicates.

Figure 7 The role of ROCK in sea urchin skeletogenesis, summary. A, at the gastrula stage (~27hpf in *P. lividus*), the triradiate spicule is coated with F-actin. B, ROCK activity is required for spicule formation and F-actin organization. C, model of the regulatory interactions between the skeletogenic GRN and cytoskeleton remodeling during spicule initiation (~24-27hpf in *P. lividus*). D, during skeletal elongation F-actin is detected around the spicule cavity and is enriched at the tips of the rods. The expression of SM50 and some other skeletogenic genes, is localized to the tips of the rods. E, under the addition of ROCK inhibitor, skeletal growth rate is reduced and ectopic spicule branching is observed. The expression of skeletogenic genes is localized to the vicinity of the growing rods. F, the regulatory interactions between the skeletogenic GRN and cytoskeleton remodeling during skeletal elongation.

References

1.   Weiner S, Addadi I. Crystallization Pathways in Biomineralization. Annu Rev Mater Res. 2011;41:21-40.

2.   Murdock DJ, Donoghue PC. Evolutionary origins of animal skeletal biomineralization. Cells Tissues Organs. 2011;194(2-4):98-102.

3.   Knoll AH. Biomineralization and Evolutionary History. Reviews in Mineralogy and Geochemistry. 2003;54:329-56.

4.   Gilbert P, Bergmann KD, Boekelheide N, Tambutte S, Mass T, Marin F, et al. Biomineralization: Integrating mechanism and evolutionary history. Sci Adv. 2022;8(10):eabl9653.

5.   Ben-Tabou de-Leon S. The Evolution of Biomineralization through the Co-Option of Organic Scaffold Forming Networks. Cells. 2022;11(4).

6.   Morgulis M, Gildor T, Roopin M, Sher N, Malik A, Lalzar M, et al. Possible cooption of a VEGF-driven tubulogenesis program for biomineralization in echinoderms. Proc Natl Acad Sci U S A. 2019;116(25):12353-62.

7.   Murdock DJE. The 'biomineralization toolkit' and the origin of animal skeletons. Biol Rev Camb Philos Soc. 2020;95(5):1372-92.

8.   Hojo H, McMahon AP, Ohba S. An Emerging Regulatory Landscape for Skeletal Development. Trends Genet. 2016;32(12):774-87.

9. Fraser GJ, Hulsey CD, Bloomquist RF, Uyesugi K, Manley NR, Streelman JT. An ancient
gene network is co-opted for teeth on old and new jaws. PLoS Biol. 2009;7(2):e31.
10. Evans JS. The Biomineralization Proteome: Protein Complexity for a Complex Bioceramic
Assembly Process. Proteomics. 2019;19(16):e1900036.
11. Khor JM, Ettensohn CA. Functional divergence of paralogous transcription factors
supported the evolution of biomineralization in echinoderms. eLife. 2017;6.
12. Kawasaki K, Buchanan AV, Weiss KM. Biomineralization in humans: making the hard
choices in life. Annu Rev Genet. 2009;43:119-42.
13. Drake JL, Mass T, Haramaty L, Zelzion E, Bhattacharya D, Falkowski PG. Proteomic
analysis of skeletal organic matrix from the stony coral Stylophora pistillata. Proc Natl Acad Sci U
S A. 2013;110(10):3788-93.
14. Tucker A, Sharpe P. The cutting-edge of mammalian development; how the embryo
makes teeth. Nat Rev Genet. 2004;5(7):499-508.
15. Langer G, Jan de Nooijer L, Oetjen K. On the role of cytoskeleton in coccolith
morphogenesis: the effect of cytoskeleton inhibitors. Journal of Phycology. 2010;46:1252–6.
16. Durak GM, Brownlee C, Wheeler GL. The role of the cytoskeleton in biomineralisation in
haptophyte algae. Scientific reports. 2017;7(1):15409.
17. Tyszka J, Bickmeyer U, Raitzsch M, Bijma J, Kaczmarek K, Mewes A, et al. Form and
function of F-actin during biomineralization revealed from live experiments on foraminifera.
Proc Natl Acad Sci U S A. 2019;116(10):4111-6.
18. Tesson B, Hildebrand M. Dynamics of silica cell wall morphogenesis in the diatom
Cyclotella cryptica: substructure formation and the role of microfilaments. J Struct Biol.
2010;169(1):62-74.
19. Tesson B, Hildebrand M. Extensive and intimate association of the cytoskeleton with
forming silica in diatoms: control over patterning on the meso- and micro-scale. PLoS One.
2010;5(12):e14300.
20. Strzelecka-Kiliszek A, Mebarek S, Roszkowska M, Buchet R, Magne D, Pikula S. Functions
of Rho family of small GTPases and Rho-associated coiled-coil kinases in bone cells during
differentiation and mineralization. Biochim Biophys Acta Gen Subj. 2017;1861(5 Pt A):1009-23.
21. Qin L, Liu W, Cao H, Xiao G. Molecular mechanosensors in osteocytes. Bone Res.
2020;8:23.
22. Huang X, Chen X, Chen H, Xu D, Lin C, Peng B. Rho/Rho-associated protein kinase
signaling pathway-mediated downregulation of runt-related transcription factor 2 expression
promotes the differentiation of dental pulp stem cells into odontoblasts. Exp Ther Med.
2018;15(5):4457-64.
23. Oliveri P, Tu Q, Davidson EH. Global regulatory logic for specification of an embryonic
cell lineage. Proc Natl Acad Sci U S A. 2008;105(16):5955-62.
24. Ettensohn CA, Dey D. KirrelL, a member of the Ig-domain superfamily of adhesion
proteins, is essential for fusion of primary mesenchyme cells in the sea urchin embryo. Dev Biol.
2017;421(2):258-70.
25. Gildor T, Winter MR, Layous M, Hijaze E, Ben-Tabou de-Leon S. The biological regulation
of sea urchin larval skeletogenesis - From genes to biomineralized tissue. J Struct Biol.
2021:107797.
26. Kahil K, Varsano N, Sorrentino A, Pereiro E, Rez P, Weiner S, et al. Cellular pathways of
calcium transport and concentration toward mineral formation in sea urchin larvae. Proc Natl
Acad Sci U S A. 2020;117(49):30957-65.

27.   Kahil K, Kaplan-Ashiri I, Wolf SG, Rechav K, Weiner S, Addadi L. Elemental compositions
of sea urchin larval cell vesicles evaluated by cryo-STEM-EDS and cryo-SEM-EDS. Acta Biomater.
2023;155:482-90.
28.   Vidavsky N, Addadi S, Mahamid J, Shimoni E, Ben-Ezra D, Shpigel M, et al. Initial stages
of calcium uptake and mineral deposition in sea urchin embryos. Proc Natl Acad Sci U S A.
2014;111(1):39-44.
29.   Wilt FH, Killian CE, Hamilton P, Croker L. The dynamics of secretion during sea urchin
embryonic skeleton formation. Exp Cell Res. 2008;314(8):1744-52.
30.   Adomako-Ankomah A, Ettensohn CA. Growth factor-mediated mesodermal cell
guidance and skeletogenesis during sea urchin gastrulation. Development. 2013;140(20):4214-
25.
31.   Sun Z, Ettensohn CA. Signal-dependent regulation of the sea urchin skeletogenic gene
regulatory network. Gene Expr Patterns. 2014;16(2):93-103.
32.   Tarsis K, Gildor T, Morgulis M, Ben-Tabou de-Leon S. Distinct regulatory states control
the elongation of individual skeletal rods in the sea urchin embryo. Dev Dyn. 2022.
33.   Morgulis M, Winter MR, Shternhell L, Gildor T, Ben-Tabou de-Leon S. VEGF signaling
activates the matrix metalloproteinases, MmpL7 and MmpL5 at the sites of active skeletal
growth and MmpL7 regulates skeletal elongation. Dev Biol. 2021;473:80-9.
34.   Winter MR, Morgulis M, Gildor T, Cohen AR, Ben-Tabou de-Leon S. Calcium-vesicles
perform active diffusion in the sea urchin embryo during larval biomineralization. PLoS Comput
Biol. 2021;17(2):e1008780.
35.   Stepicheva NA, Dumas M, Kobi P, Donaldson JG, Song JL. The small GTPase Arf6
regulates sea urchin morphogenesis. Differentiation. 2017;95:31-43.
36.   Moreno B, DiCorato A, Park A, Kellen Mobilia, Knapp R, Bleher R, et al. Culture of and
experiments with sea urchin embryo primary mesenchyme cells. Methods in Cell Biology.
2019;150:293-330.
37.   Sepulveda-Ramirez SP, Toledo-Jacobo L, Henson JH, Shuster CB. Cdc42 controls primary
mesenchyme cell morphogenesis in the sea urchin embryo. Dev Biol. 2018;437(2):140-51.
38.   Croce J, Duloquin L, Lhomond G, McClay DR, Gache C. Frizzled5/8 is required in
secondary mesenchyme cells to initiate archenteron invagination during sea urchin
development. Development. 2006;133(3):547-57.
39.   Marletaz F, Couloux A, Poulain J, Labadie K, Da Silva C, Mangenot S, et al. Analysis of the
P. lividus sea urchin genome highlights contrasting trends of genomic and regulatory evolution
in deuterostomes. Cell Genom. 2023;3(4):100295.
40.   Arshinoff BI, Cary GA, Karimi K, Foley S, Agalakov S, Delgado F, et al. Echinobase:
leveraging an extant model organism database to build a knowledgebase supporting research on
the genomics and biology of echinoderms. Nucleic Acids Res. 2022;50(D1):D970-D9.
41.   Ettensohn CA, McClay DR. Cell lineage conversion in the sea urchin embryo. Dev Biol.
1988;125(2):396-409.
42.   Chen W, Mao K, Liu Z, Dinh-Xuan AT. The role of the RhoA/Rho kinase pathway in
angiogenesis and its potential value in prostate cancer (Review). Oncol Lett. 2014;8(5):1907-11.
43.   Jacobs M, Hayakawa K, Swenson L, Bellon S, Fleming M, Taslimi P, et al. The structure of
dimeric ROCK I reveals the mechanism for ligand selectivity. J Biol Chem. 2006;281(1):260-8.
44.   Liu J, Wada Y, Katsura M, Tozawa H, Erwin N, Kapron CM, et al. Rho-Associated Coiled-
Coil Kinase (ROCK) in Molecular Regulation of Angiogenesis. Theranostics. 2018;8(21):6053-69.
45.   Julian L, Olson MF. Rho-associated coiled-coil containing kinases (ROCK): structure,
regulation, and functions. Small GTPases. 2014;5:e29846.

46.   Lecuit T, Lenne PF, Munro E. Force generation, transmission, and integration during cell
and tissue morphogenesis. Annu Rev Cell Dev Biol. 2011;27:157-84.
47.   Amin E, Dubey BN, Zhang SC, Gremer L, Dvorsky R, Moll JM, et al. Rho-kinase:
regulation, (dys)function, and inhibition. Biol Chem. 2013;394(11):1399-410.
48.   Pavlov D, Muhlrad A, Cooper J, Wear M, Reisler E. Actin filament severing by cofilin. J
Mol Biol. 2007;365(5):1350-8.
49.   Yarmola EG, Somasundaram T, Boring TA, Spector I, Bubb MR. Actin-latrunculin A
structure and function. Differential modulation of actin-binding protein function by latrunculin
A. J Biol Chem. 2000;275(36):28120-7.
50.   Schatten G, Schatten H, Spector I, Cline C, Paweletz N, Simerly C, et al. Latrunculin
inhibits the microfilament-mediated processes during fertilization, cleavage and early
development in sea urchins and mice. Exp Cell Res. 1986;166(1):191-208.
51.   Kovacs M, Toth J, Hetenyi C, Malnasi-Csizmadia A, Sellers JR. Mechanism of blebbistatin
inhibition of myosin II. J Biol Chem. 2004;279(34):35557-63.
52.   Moorhouse KS, Gudejko HF, McDougall A, Burgess DR. Influence of cell polarity on early
development of the sea urchin embryo. Dev Dyn. 2015;244(12):1469-84.
53.   Mabuchi I. Cleavage furrow: timing of emergence of contractile ring actin filaments and
establishment of the contractile ring by filament bundling in sea urchin eggs. J Cell Sci. 1994;107
(Pt 7):1853-62.
54.   Schroeder TE. Fourth cleavage of sea urchin blastomeres: microtubule patterns and
myosin localization in equal and unequal cell divisions. Dev Biol. 1987;124(1):9-22.
55.   Wong GK, Allen PG, Begg DA. Dynamics of filamentous actin organization in the sea
urchin egg cortex during early cleavage divisions: implications for the mechanism of cytokinesis.
Cell motility and the cytoskeleton. 1997;36(1):30-42.
56.   Khor JM, Ettensohn CA. Transcription Factors of the Alx Family: Evolutionarily Conserved
Regulators of Deuterostome Skeletogenesis. Frontiers in genetics. 2020;11:569314.
57.   Miklavc P, Frick M. Actin and Myosin in Non-Neuronal Exocytosis. Cells. 2020;9(6).
58.   Bibeau JP, Kingsley JL, Furt F, Tuzel E, Vidali L. F-Actin Mediated Focusing of Vesicles at
the Cell Tip Is Essential for Polarized Growth. Plant Physiol. 2018;176(1):352-63.
59.   Bond LM, Brandstaetter H, Sellers JR, Kendrick-Jones J, Buss F. Myosin motor proteins
are involved in the final stages of the secretory pathways. Biochem Soc Trans. 2011;39(5):1115-
9.
60.   Segal D, Zaritsky A, Schejter ED, Shilo BZ. Feedback inhibition of actin on Rho mediates
content release from large secretory vesicles. J Cell Biol. 2018;217(5):1815-26.
61.   Rousso T, Schejter ED, Shilo BZ. Orchestrated content release from Drosophila glue-
protein vesicles by a contractile actomyosin network. Nat Cell Biol. 2016;18(2):181-90.
62.   Olson EN, Nordheim A. Linking actin dynamics and gene transcription to drive cellular
motile functions. Nat Rev Mol Cell Biol. 2010;11(5):353-65.
63.   Knipe RS, Tager AM, Liao JK. The Rho kinases: critical mediators of multiple profibrotic
processes and rational targets for new therapies for pulmonary fibrosis. Pharmacol Rev.
2015;67(1):103-17.
64.   Khatiwala CB, Kim PD, Peyton SR, Putnam AJ. ECM compliance regulates osteogenesis by
influencing MAPK signaling downstream of RhoA and ROCK. J Bone Miner Res. 2009;24(5):886-
98.
65.   Zhong J, Yang Y, Liao L, Zhang C. Matrix stiffness-regulated cellular functions under
different dimensionalities. Biomater Sci. 2020;8(10):2734-55.
66.   Luo YJ, Su YH. Opposing nodal and BMP signals regulate left-right asymmetry in the sea
urchin larva. PLoS Biol. 2012;10(10):e1001402.

67.   Siegel S, & Castellan, N. J., Jr. . Nonparametric statistics for the behavioral sciences 2ed:
Mcgraw-Hill Book Company; 1988.
68.   Stalling D, Westerhoff M, Hege H-C. Amira: A highly interactive system for visual data
analysis. The visualization handbook. 2005;38:749-67.