TrAGEDy: Trajectory Alignment of Gene Expression Dynamics

Ross F. Laidlaw[1], Emma M. Briggs[1,2], Keith R. Matthews[2], Richard McCulloch[1], Thomas D. Otto[1*]

[1]Wellcome Centre for Integrative Parasitology, School of Infection & Immunity, University of Glasgow, Glasgow, UK

[2]Institute for Immunology and Infection Research, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

*Correspondence to thomasdan.otto@glasgow.ac.uk

Abstract

To compare biological processes between conditions, alignment of single cell transcriptomic trajectories can be performed. Current tools place constraints on what data can be aligned. We present Trajectory Alignment of Gene Expression Dynamics (TrAGEDy) which overcomes these constraints, allowing the alignment of asymmetric biological processes. Across simulated and real datasets, TrAGEDy returns correct alignment based on underlying simulated process, where current tools fail. Across different biological contexts, TrAGEDy can capture biologically relevant genes which other differential expression methods fail to detect. TrAGEDy provides a new tool for in-depth analysis of many emerging single cell transcriptomic datasets.

Background

First described in 2014 (1), Trajectory Inference (TI) methods order cells based on the gradual change in transcript levels underlying biological process/es captured by single cell

RNA-sequencing (scRNA-seq) (2). Cells are assigned a "pseudotime" value based on their

position in the inferred trajectory, allowing differential gene expression tests over

pseudotime. Various methods have been developed to identify differentially expressed (DE)

genes in a biological process of interest, in some cases allowing targeted comparisons.

Notably, TradeSeq (3) uses negative binomial general additive models (GAM) and Wald

tests to assess whether a gene is differentially expressed over pseudotime within a trajectory

or between lineages of a trajectory. Alternatively, Monocle's BEAM (4)identifies genes

associated with a particular branch of a trajectory whereas pseudotimeDE (5) uses

permutation to account for uncertainty in pseudotime and a zero-inflated negative binominal

GAM to account for expression value dropout. By applying these techniques, dynamic gene

expression patterns associated with a variety of interesting biological events have been

investigated, such as regulators of myogenesis (1), effector gradients in CD4+ T cells (6),

and life cycle transitions of the pathogen *Trypanosoma brucei* (7).

These methods can be extended to identify DE genes between two conditions where the

common biological process under analysis varies, such as before and after mutation or in an

overlapping response to different stimuli. One approach is to integrate the separate datasets

together and do a cluster-based comparison between conditions, treating cells as being at a

discrete stage in development captured in each cluster. This method misses the nuance that

development may be continuous, with some cells falling between stage boundaries. Another

approach is to integrate the datasets together, perform TI to capture the shared, possibly

branching, trajectory and differential expression tests across this pseudotime axis for each

condition, using tools like TradeSeq's conditionTest. However, the integration may force

similarity between the trajectories, thus obscuring DE genes.

Alternatively, trajectory alignment can align cells from independently generated trajectories

to find a common pseudotime axis that retains the original ordering of cells without dataset

integration. Analysing aligned trajectories can reveal DE genes across the captured process,

as well as differences between conditions. The discrete point at which the genes are DE

between conditions can also be found. The first method described to perform such trajectory

alignment was CellAlign (8), which uses dynamic time warping (DTW) to align two

trajectories together. Interpolated points are created across the trajectory and gene

expression values are assigned to the interpolated points based on the cells that are in

pseudotime vicinity to the points. Similarity between the conditions is assessed by computing

the distance or correlation between the scaled gene expression values for the interpolated

points on opposing conditions and, from this, the minimum cost path through the interpolated

points is calculated using DTW.

The use of DTW imposes some limitations on the type of alignment that can be analysed.

Trajectories must start and end at the same point, and each part of a trajectory must be

matched to at least one point in the other trajectory. The practical implication of such

limitations is that cell types which may not be represented on the opposing trajectory, are

nevertheless matched to a point on the other trajectory, making interpreting the subsequent

alignment difficult. Here, we build on the work of cellAlign and develop a tool we called

Trajectory Alignment of Gene Expression Dynamics (TrAGEDy), where we make post-hoc

changes to the alignment, allowing us to overcome the limitations of DTW and better reflect

differences that may occur across the alignment, allowing us to implement our own

differential gene expression test, in order to better identify differences between the two

conditions. We test TrAGEDy with a variety of simulated scRNA-seq datasets with different

underlying alignments, as well datasets of *Trypanosoma brucei* lifecycle and T cell

development under different genetic conditions, revealing gene expression changes

undetected by current methods.

Results

TrAGEDy overview

A typical workflow for TrAGEDy is as follows (Fig.1). Prior to trajectory alignment with TrAGEDy, a predefined list of genes is selected for TI (Trajectory Inference) and dissimilarity is calculated between the datasets. Top variable genes or cluster markers can be selected, for example. The two datasets are then projected into a reduced dimension space using the preselected features and TI performed on the two datasets independently (Fig.1A). There is no limitation on what TI method can be used, provided pseudotime values are generated for each cell.

TrAGEDy then uses the cellAlign method of creating interpolated points to sample gene expression at different points in the trajectory, allowing inherently noisy scRNA-seq data to be smoothed across the trajectory. The closer a cell is to the interpolated point, the more it contributes to its gene expression. By changing a parameter which controls a gaussian pseudotime window, the user can alter the contribution level more distal cells have to interpolated point gene expression (Fig.1B).

Dissimilarity between the gene expression of the interpolated points is next assessed. While interpolation can smooth out noise, possible batch effects between datasets mean calculating dissimilarity with methods such as Euclidean distance is problematic without scaling the data first. We used Spearman correlation, as it is less sensitive to outliers than Pearson and, unlike Euclidean distance, does not require scaling of gene expression values.

Figure 1. Graphical overview of the TrAGEDy process.

Trajectory Inference is carried out on two datasets which share a common process but have

a difference in condition (A). TrAGEDy samples gene expression across interpolated points

of the trajectory, with cells with closer pseudotime values to the interpolated points

contributing more to their gene expression (B). TrAGEDy then aligns the pseudotime of the

interpolated points then the cells, finally performing a sliding window comparison between

cells at similar points in aligned pseudotime, to extract differentially expressed genes (C).

TrAGEDy next finds the optimal path through the dissimilarity matrix of the interpolated

points, which constitutes the shared process between the two trajectories (Fig.1B). DTW,

with alterations, is used to find the optimal path. To overcome the constraint in DTW that the

two processes must start and end at the same point, we first cut the dissimilarity matrix so

the path starts and ends at points which are most transcriptionally similar to one another.

Another constraint of DTW is that all points must be matched to at least one other point; thus

(post-DTW) we pruned any matches that have high transcriptional dissimilarity, enabling

processes which may have diverged in the middle of their respective trajectories to be

compared. Finally, we used the modified alignment to adjust the pseudotime values of the

interpolated points, generating new values as the basis to then adjust the pseudotime value

of each cell (Fig.1C).

To extract DE genes, we performed a sliding window comparison, clustering cells at similar

points in aligned pseudotime together then testing for significance with a Mann-Whitney U

test, extracting genes that are DE at different points in the shared process (Fig.1C).

Clustering with a sliding window allows comparisons over a continuous process. Users can

define how much overlap there is between the windows of comparison.

Aligning simulated scRNA-seq datasets

We tested TrAGEDy's ability to correctly identify the alignment of two trajectories on

datasets where the alignment points were known *a priori*. Using Dyngen (9) we simulated

scRNA-seq datasets with inherent trajectories encoded in the expression values, resulting in

three different test cases: two positive controls and one negative control.

The first positive test case used two datasets, the first simulating a full developmental

progression involved sequential expression of many genes (wild type; WT) and the second

resulting from knockout (KO) of one of these genes (Fig.2A). Thus, the KO trajectory is

truncated at the point of gene KO. TrAGEDy correctly aligned these datasets: the WT and

the KO datasets initially align with one another, but the KO finishes its full process before the

WT. For comparison, CellAlign was also performed using these datasets, with Euclidean

distance selected as the metric to assess similarity, as this was the metric used in the source

publication. Like TrAGEDy, CellAligns distance matrix revealed the initial alignment of the

WT and KO, before the KO trajectory ends while the WT continues. However, CellAlign

failed to position the correct end point of the KO trajectory relative to the WT due to running

DTW on the whole dissimilarity matrix.

TrAGEDy cellAlign

KO Process

A

KO C Process

B

A process

C

Figure 2. Alignment of simulated single cell RNA sequencing datasets with TrAGEDy

and cellAlign.

Three simulated datasets were generated with Dyngen in which there was initial alignment

(A), alignment at the beginning and end only (B) and no alignment (C), between the two

simulated processes. TrAGEDy (left column) and cellAlign (right column) were used to align

the datasets with Euclidean distance being used to measure transcriptional dissimilarity in

cellAlign, while TrAGEDy used Spearman correlation. The transcriptional dissimilarity score

between individual interpolated points of the two processes is shown in the corresponding

key, with blue being low dissimilarity and red being high. If the interpolated points are part of

the optimal path, their index colour is grey.

To assess if TrAGEDy can align trajectories that have shared start and end points, but

deviate in the middle, we simulated two datasets with Dyngen that have a diverge-converge

backbone. By knocking out one of the divergent branches in one dataset (KO C) and the

other divergent branch in the other dataset (KO D), we simulated linear trajectories which do

not share a common process in their middle points (Fig.2B). Applying TrAGEDy to this

scenario accurately captured the initial and terminal alignment of the two trajectories, while

leaving the middle sections unaligned. CellAlign failed to align KO C and KO D correctly, as

CellAlign does not include features to prune matches.

For the negative control, two datasets with distinct gene regulatory networks and

transcription kinetics were generated (A and one) (Fig.2C). As there is no shared process

between the conditions, the dissimilarity scores are expected to be high, and no alignment

should be found. The dissimilarity score calculated by TrAGEDy between process A and

process one showed that most of the indexes have a dissimilarity score close to 1, and thus

do not share a common process. As TrAGEDy finds the optimal path by looking at the

context of the whole dissimilarity matrix, even if the dissimilarity scores are high, it will still

find a path, as seen in the negative control. TrAGEDy will give a warning to the user if the

median of the cut dissimilarity matrix scores is over 0.6 and thus it is up to the user to

interpret the resulting alignment in the context of the overall dissimilarity score of the path.

CellAlign also creates a path through the dissimilarity matrix, which has a higher overall

dissimilarity range than the other two datasets, with the Euclidean distance ranging from ~7-

14, while the other two datasets ranged from ~4-12. If, instead, Pearson correlation is

selected for CellAlign, it becomes clearer that the two datasets share little transcriptional

similarity (supp. Fig.4C).

181   WT vs Bcl11b KO T cell development analysis

Simulated data represents a 'best case' scenario for benchmarking tools, as they often lack

the complexity of real datasets. We thus decided to apply TrAGEDy to real scRNA-seq

datasets where the underlying process has been characterised. We first applied TrAGEDy to

a scRNA-seq dataset of *in vitro* T cell development in WT and Bcl11b KO condition mice

over two timepoints (10). The data contain cells undergoing the early stages of T cell

development in the thymus from thymus-seeding precursors (TSP) to the double negative

three stage (DN3), where the cells undergo beta selection (11). The authors find that when

Bcl11b is removed, the cells deviate from the T cell commitment pathway and develop

distinct transcriptomic signatures, including expression of Id2 and Rora. However, some

Bcl11b KO cells still express markers of T cell pathway commitment such as the gene for the

pre-T cell Antigen receptor alpha (Ptcra) (12) and Rag1, one of the central genes for TCR

gene (13) (supp. Fig. 6). We thus aimed to compare the WT and Bcl11b KO trajectories as

development proceeds and diverges between the two conditions to profile the differences in

transcriptome.

Figure 3. Analysis of WT and Bcl11b KO trajectories of T cell development with

TrAGEDy

TrAGEDy alignment of the WT and Bcl11b KO trajectories of T cell development with

transcriptional dissimilarity calculated with Spearman correlation (A). Alignment of the

interpolated points of the WT and Bcl11b KO trajectories, showing connections between the

two processes as calculated by TrAGEDy. The interpolated points are displayed across

pseudotime before (B) and after (C) the pseudotime values have been modified by

TrAGEDy. (D) UpSet plot showing the top 10 intersections of differentially expressed genes

captured by TrAGEDy across 7 windows of comparison and Seurat over 7 clusters and

TradeSeq over 7 knot comparisons. (E) Heatmap showing DE genes with the highest log 2

fold-change (Log2FC) captured (but not uniquely captured) between the WT and Bcl11b KO

trajectories at 7 different windows of comparison. Each index is coloured by the Log2FC (red

= higher expression in KO, blue = higher expression in WT) and an * symbol indicates the

difference is significant (bonferroni adjusted $p < 0.05$, Mann-Whitney U test, abs(Log2FC) >

0.5, minimum percentage expressed > 0.1). Full list of heatmap genes can be found in

supplementary table 7. (F) TradeSeq plotSmoother plots showing the expression changes

that occur across the WT (blue) and Bcl11b KO (red) trajectories within the cells of aligned

pseudotime, as calculated by TrAGEDy, which share a common process. The 6 genes

plotted were only identified as being significantly DE by TrAGEDy.

We first tested whether TrAGEDy could overcome batch effects between the two replicates

to still return a biologically accurate alignment. The cells were sequenced over two runs, but

each run contained a variety of biological and technical replicates, as well as cell from

different sampling days. For simplicity, the WT cells sequenced in run 1 will be referred to as

WT1, the Bcl11b KO cells sequenced in run 1 will be referred to as Bcl11b KO1 and those

sequenced in run 2 as WT2 and Bcl11b KO2, resulting in four individual datasets. PHATE

(14) embeddings were generated independently and used as the basis for Slingshot TI to

calculate pseudotime values for each of the cells. When projected into the PHATE space,

some of the WT1 cells were separated from the main body of the trajectory, which had

downstream effects on the alignment, and so these cells were removed as they did not

appear to constitute part of the biological process under examination (supp. Fig. 7).

TrAGEDy was used to first align the two datasets for each condition together, resulting in

two separate alignments: a WT alignment of WT1 and WT2, and a Bcl11b KO alignment of

Bcl11b KO1 and Bcl11b KO2 (supp. Fig. 8). The WT1 dataset only contains cells from day

10 of sampling while WT2 only contains cells from day 13, meaning the alignment may be

skewed with the WT1 process starting before the WT2 process, but finishing before the WT2

process; a pattern captured by TrAGEDy (supp. Fig. 8). TrAGEDy was then performed on

the aligned WT and Bcl11b KO trajectories. The resulting final alignment shows the two

conditions sharing an initial common process, but the Bcl11b KO developmental progression

finished before that of the WT (Fig. 3A, B, C).

	TrajDE	tradeSeq	Seurat
-----------------------0.25 Log2FC threshold-----------------------			
Total genes	1064	980	196
Unique genes	554	479	12
-----------------------0.5 Log2FC threshold-----------------------			
Total genes	300	484	40
Unique genes	164	348	5

Runtime (minutes)	2.7	450	0.13

Table 1 Summary of DE gene numbers for the WT vs Bcl11b KO T cell comparison.

For the WT vs Bcl11b KO T cell development dataset comparison, the number of DE genes

captured by the three different methods in total, and DE genes that they captured uniquely

by each method, using two different Log2FC cutoffs (0.25 (top) and 0.5 (bottom)) for

comparisons. The time taken to run the necessary functions to get differentially expressed

genes (in minutes) is shown on the final row.

By aligning trajectories rather than integrating the datasets, we expected to preserve more

variability between them, which may be reflected in the DE genes captured. We thus

compared TrAGEDy against other methods which can extract DE genes between conditions.

We compared the differential gene expression tests by TrAGEDy (TrajDE) with established

methods of extracting DE genes: we compared TrajDE to a standard Seurat pipeline of

comparing DE genes between conditions, and TradeSeq's method of comparing DE genes

across trajectories from different conditions. Analysis was carried out using each method,

and genes were classified as DE using two different thresholds, 0.25 and 0.5 log fold change

(log2FC). A brief overview of these results can be seen in Table 1, which indicates that

TrajDE was able to detect the most DE genes, including 554 not detected by TradeSeq or

Seurat, at the lower threshold of 0.25 log2FC. Below we discuss biological implications of

the genes classified as DE with the more stringent threshold of 0.5 log2FC. With this

threshold, TrajDE returned 300 genes as being DE across the 7 windows of comparison,

while Seurat only returned 40 across 7 clusters, and TradeSeq returned 484 genes across 7

comparison knots. TrajDE uniquely captured 164 genes that neither Seurat nor TradeSeq

captured, whereas Seurat found 5 unique DE genes and TradeSeq uniquely captured 348

DE genes (table 1). The number of DE genes that were captured per window/cluster and the

intersection of the different lists can be viewed in Fig.3C. Of the 164 DE genes captured only

by TrAGEDy, 121 were expressed higher in the WT condition and 45 were expressed higher

in the Bcl11b KO condition. Genes with higher expression in the WT include those

associated with T cell signaling (Lat (12)), cell cycle associated genes (Top2a, Dut, Cenpa

(15), Btg1 (16)), and those with functions for protecting against reactive oxygen species

(ROS) (Txn1, Prdx1, Gsr) (17). The genes only TrAGEDy identified as being significantly

higher expressed in Bcl11b KO cells include factors involved in immune cell activation (Jak1,

Nfkb1 (18), Jund (12)), cytotoxic elements (Srgn, (19)), and transcription factors (Runx3 (20),

Sox4 (21)).

Both TrAGEDy and TradeSeq captured many of the same biological relevant genes. Genes

TrAGEDy captured as being significantly higher expressed in the WT, which TradeSeq also

captured as being DE, included the pre-TCR alpha chain gene (Ptcra), CD3 chain genes

(Cd3e, Cd3g and Cd3d) (12), TCR recombination associate gene Dntt (22), immune

synapse associated genes (Cd48 (23), Cd7 (24,25)). Genes identified as significantly higher

expressed in the Bcl11b KO T cells by both methods included: NK cell marker (Cd244) (26),

cytotoxic molecule Gzma and IgE receptor Fcer1g (27). While it took TradeSeq around 7

hours to return its DE results (time taken to execute the fitGAM and conditionTest functions),

it only took TrAGEDy around 3 minutes to perform its process (Table 1). TrAGEDy

additionally allows the relative timing of expression changes between conditions across the

trajectory to be identified, by investigation of expression at each window (Fig. 3E). Plotting

some of the DE genes that only TrAGEDy captured show they are concurrent with the

patterns of expression TrAGEDy suggests, with Notch1, Rag1 and Hivep3 being higher

expressed in WT T cells, while Jak1, Gzmb and Nfkb1 are higher expressed in Bcl11b KO

cells (Fig. 3F).

WT vs ZC3H20 KO *Trypanosoma brucei* alignment

In its mammalian host, WT *Trypanosoma brucei* parasites transition from replicative slender

forms to non-replicative stumpy forms, through a quorum sensing mechanism (28–30)

Mutant *T. brucei* where an RNA binding protein essential for slender to stumpy

differentiation, ZC3H20, has been knocked out (ZC3H20 KO) are unable to undergo this

transition (31,32) and fail to express stumpy associated genes. Briggs and colleagues (7)

used 10x Chromium to sequence two biological replicates of WT *T. brucei* parasites

undergoing this transition *in vitro*, as well as ZC3H20 KO failing to differentiate in the same

conditions. The authors defined four main clusters Long Slender (LS) A and B, and Short

Stumpy (SS) A and B to show the progression of the transition, with LS A being the start and

SS B being the end. Thus, the final alignment is expected to show that the WT and the

ZC3H20 KO datasets share an initial common process, but that the KO process is truncated

or branched relative to the WT, as suggested by previous analysis using data integration and

Slingshot TI.

Figure 4. Analysis of WT and ZC3H20 KO *Trypanosoma brucei* development with

TrAGEDy.

TrAGEDy alignment of the WT and ZC3H20 KO trajectories of *Trypanosoma brucei*

development with transcriptional dissimilarity calculated with Spearman correlation (A).

Alignment of the interpolated points of the WT and ZC3H20 KO trajectories, showing

connections between the two processes as calculated by TrAGEDy. The interpolated points

are displayed across pseudotime before (B) and after (C) the pseudotime values have been

modified by TrAGEDy. (D) UpSet plot showing the top 10 intersections of DE genes as

captured by TrAGEDy across four windows of comparison and Seurat over the four long

slender (LS) clusters and TradeSeq over 4 knot comparisons. (E) Heatmap showing the log

2 fold-change (log2FC) of genes found only by TrAGEDy to be DE between the WT and

ZC3H20 KO trajectories in only 1 of the 4 different windows of comparison. Each index is

coloured by the Log2FC (red = higher expression in KO, blue = higher expression in WT)

and a * symbol indicates the difference is significant (bonferroni adjusted $p < 0.05$ Mann-

Whitney U test, abs(Log2FC) > 0.5, minimum percentage expressed > 0.1). Genes names

were taken from TriTrypDB. Full list of heatmap genes can be found in Supplementary table

7. (F) TradeSeq plotSmoother plots showing the expression changes that occur across the

WT (blue) and ZC3H20 KO (red) trajectories within the cells of aligned pseudotime, as

calculated by TrAGEDy, which share a common process. The 6 genes plotted were only

identified as being significantly DE by TrAGEDy.

Applying TrAGEDy to WT01 and WT02 returned an alignment which started at the same

point, and ended just before the final endpoint, returning a nearly 1:1 alignment, as predicted

of biological replicates (supp. Fig.5). We next used the aligned WT values to perform

TrAGEDy alignment between the combined WT replicates and the ZC3H20 KO dataset. The

resulting alignment showed that the two process have an initial point of alignment with the

332 ZC3H20 KO process finishing before the WT process does (Fig.4A). Using the DTW

333 calculated path, we could then adjust the pseudotime values of the trajectories to get a

334 common pseudotime axis (Fig.4B, C).

	TrajDE	tradeSeq	Seurat
----------------------0.25 Log2FC threshold----------------------			
Total genes	604	253	179
Unique genes	344	36	10
----------------------0.5 Log2FC threshold----------------------			
Total genes	264	78	14
Unique genes	208	25	0
--			
Runtime (minutes)	0.85	96	0.024

Table 2 Summary of DE gene numbers for the WT vs ZC3H20 KO *Trypanosoma brucei*

comparison.

For the WT vs ZC3H20 KO *Trypanosoma brucei* development dataset comparison, the

number of DE genes captured by the three different methods in total, and DE genes that

they captured uniquely by each method, using two different Log2FC cutoffs (0.25 (top) and

0.5 (bottom)) for comparisons. The time taken to run the necessary functions to get

differentially expressed genes (in minutes) is shown on the final row.

In this case, comparing DE results between methods revealed TrAGEDy was able to detect

more genes compared to TradeSeq and Seurat, at both a lower and high log2FC threshold,

and detect a greater number of DE genes missed by the other methods. Using a 0.5 log2FC

threshold, TrajDE extracted 264 DE genes from 4 windows of comparison, Seurat extracted

14 across the 4 annotated slender clusters, and TradeSeq captured 78 genes across the 4

knot comparisons (Table 2). Of the genes only captured by TrajDE, most had significantly

higher expression in the WT (n=202) compared with the ZC3H20 KO (n=7) (Fig 4D). The

202 genes upregulated in the WT condition uniquely identified by TrAGEDy includes genes

involved in metabolism and catabolism of amino acids (glutamate dehydrogenase (33), L-

threonine 3-dehydrogenase, and 2-amino-3-ketobutyrate coenzyme A ligase (AKCT) (34)

and that act in the cytoskeleton and motor activity (actin-B and cytoskeleton-associated

proteins, kinesin-C and dynein (35)) and post-transcriptional regulation of gene expression

(Splicing factor TSR1 (36), RBP42 (37), UBP1 & UBP2 (38)). TrAGEDy was the only method

able to detect the known stumpy marker, Protein Associated with Differentiation (PAD)2 (39),

as being DE and upregulated in the WT trajectory, as expected. The 7 genes TrAGEDy

identified as being significantly higher expressed in the ZC3H20 KO cells include GAPDH,

expression site-associated gene 5 (ESAG5), and ZC3H38. By assessing DE over sequential

windows, genes with transient transcript level changes between conditions across

pseudotime could be identified (Fig. 4E).

To visualise changes in expression between conditions, we used TradeSeq to fit a GAM to

each gene in the WT and ZC3H20 KO datasets using the new TrAGEDy calculated

pseudotime to plot the smoothed expression. While the GAM was fitted across the whole

TrAGEDY aligned pseudotime, visualisations were made for only cells TrAGEDy defined as

being aligned between the two datasets and showed that the GAMs follow a similar pattern

as identified by TrAGEDy, with AKCT, L-threonine dehydrogenase, glutamate

dehydrogenase & RBP48 being higher expressed in the WT cells and ESAG5 and ZC3H38

being higher expressed in the ZC3H20 KO cells (Fig.4F).

Discussion

In this paper we present TrAGEDy, a tool for aligning and comparing single cell trajectories

between conditions. TrAGEDy allows the alignment of processes which may start and end at

different points by identifying the point at which dissimilarity increases from the minimum

dissimilarity point. TrAGEDy assesses alignments to remove matches where the dissimilarity

score is unaligned in portions of the trajectories, allowing TrAGEDy to additionally capture

alignments where the processes diverge and subsequently converge. TrajDE then identifies

differences in gene expression over pseudotime between the conditions by taking a sliding

window soft clustering approach, allowing it to identify transient changes in gene expression

over the shared process.

Across multiple simulated datasets, TrAGEDy can find the underlying alignment between

two processes, where cellAlign fails, due to its ability to prune matches and identify optimal

start and end points in the dissimilarity matrix (Figure 2A). Across biological replicates and

different conditions of real datasets, TrAGEDy overcomes batch effects to deliver accurate

alignments. We additionally find that applying TrAGEDy to real scRNA-seq datasets

identifies biologically relevant genes where TradeSeq and Seurat DE fail, with these

comparisons being performed in a fraction of the time when compared to TradeSeq (Table 1

& 2).

A key consideration when applying TrAGEDy is the density of cells across pseudotime.

However, the window TrAGEDy uses to calculate expression at the interpolated points can

be adjusted to account for regions with few cells. Notably, if cells are present which are

separated from the main bulk of the process, like in the WT1 T cell dataset (supp. Fig.7), this

leads to an uneven dissimilarity score between neighboring interpolated points, thus biasing

the end alignment. These outlier cells should thus be removed before TrAGEDy is carried

out, to ensure a smooth and accurate alignment.

As of 2019, there were more than 70 TI methods, all of which have varying performance,

functionality and constraints (2). Additionally, the chosen method of dimensionality reduction

(PCA, UMAP, PHATE, etc) used to generate the embeddings that the trajectory is calculated

from also affects the pseudotime results. We chose PHATE for the reduced dimension

embeddings as its purpose is to identify transitions in scRNA-seq data (14). User discretion

is advised when choosing the TI and dimensionality reduction method. Given this variability

that can occur before TrAGEDy is applied, a core assumption of TrAGEDy is that the

pseudotime must be an accurate reflection of the underlying biological process. In all

simulated datasets discussed here, TrAGEDy was able to correctly align trajectories, and

with greater accuracy than cellAlign. Additionally, alignment of real data sets resulted in

expected outcomes, including the complete alignment of biological replicates and partial

alignments between conditions.

To compare DE results of TrAGEDy we also performed differential gene expression tests

using two widely used tools: Seurat and TradeSeq. Although, Seurat does not take

pseudotime into account when it performs differential expression tests, it is possible to

compare conditions which is not implemented in most tools designed to specifically analyse

trajectories, including pseudotimeDE (5) and Monocle (4). Low detection of DE genes by

Seurat is thus likely explained by the fact it does not take pseudotime into account when

carrying out the differential expression tests. TradeSeq, however, is designed for

pseudotime-based differential gene expression tests and is able to compare conditions. We

thus compared TrAGEDy results with these two tools, using both a low and higher stringency

logFC threshold.

When analysing the T cell development dataset, TradeSeq captured more DE genes at a

log2FC threshold of 0.5 than TrAGEDy, but at a log2FC of 0.25 TrAGEDy captured more DE

genes. While getting a higher number of DE genes is an important benchmark, it is also

important to assess whether these genes fit into the context of the existing literature

surrounding the datasets. TrAGEDy uniquely captured many cell-cycle associated genes as

significantly upregulated in the WT compared with Bcl11b KO. After beta-selection, before

they begin to rearrange their TCR alpha chain, T cells undergo a proliferative burst and then

become quiescent (40). In accordance with this, cell cycle genes are mostly DE in the final

two windows (windows 6 and 7) and the anti-proliferation factor Btg1 (16) (which was only

detected by TrAGEDy) is significantly higher expressed in the WT in only the final window

(window 7) (supp. Table 1). This indicates that TrAGEDy has managed to capture the

transient changes in proliferative state as WT T cells prepare to rearrange their TCR alpha

chain. It also indicates that Bcl11b KO cells may have reduced ability to carry out this

proliferative burst. TrAGEDy also captured the transcription factor Hivep3 as significantly

more highly expressed in the WT compared with Bcl11b KO cells. This gene has been

shown to bind regions of the DNA associated with V(D)J recombination of antigen receptors

in T and B cells (41). Interestingly, Dntt is also significantly higher expressed in the WT, and

so Bcl11b KO cells may have reduced capacity to rearrange their TCR; affecting their

signaling ability with a knock-on effect on the proliferative burst.

While the effect of ROS on T cell development has not been fully elucidated, increased ROS

generation leads to reduced levels of mature single positive T cells in the thymus and

treatment of T cells with factors which limit ROS show increased survival (42). This suggests

that reduced ROS positively influences T cell development, and with genes associated with

antioxidation being significantly higher expressed in WT cells, Bcl11b KO cells may be more

susceptible to ROS, limiting their maturation ability.

In the Bcl11b KO T cells, TrAGEDy identified Gzmb and Srgn as being significantly more

highly expressed compared with WT T cells. Gzmb is a key enzyme expressed in cytotoxic

immune cells such as Natural killer (NK) cells while Srgn plays a role in storage of Gzmb in T

cells (19). Expression of such genes (and Cd244 which both TradeSeq and TrAGEDy

detected) is in keeping with Zhou and colleagues' (10) observation that the Bcl11b KO T

cells appear to develop a NK cell-like phenotype.

While TradeSeq returns more unique DE genes than the other methods tested here for the T

cell dataset, it does so at the cost of runtime, which stretched into several hours, compared

with the minutes taken for TrAGEDy to run its process. As TrAGEDy not only uniquely

captures biological relevant DE genes but captures many that TradeSeq does, at the fraction

of the time, TrAGEDy is well poised to quickly and efficiently uncover the biological

differences that occur across a process between conditions.

When applied to *T. brucei* development*,* TrAGEDy captured more DE genes compared to

Seurat and TradeSeq at both low and high logFC cut-offs, suggesting greater analytical

power. Many of the 208 genes identified only by TrAGEDy have known associations with the

biological process under investigation (discussed below), indicating TrAGDEy can uncover

DE genes with biological relevance as opposed to increasing the false-discovery rate.

For example, several DE genes reflect the parasite's need to alter its metabolism when

moving from a high to low glucose environment. Glutamate hydrogenase, found to be

significantly more highly expressed in the WT, is important in proline metabolism, a key

feature of survival in the tsetse fly, as proline is a major energy source for the parasite (33).

Stumpy form *T. brucei,* a specialised transmission stage, preadapts for survival in the tsetse

fly midgut where they differentiate to procyclic forms, and so upregulation of glutamate

hydrogenase is likely to aid rapid adjustment to nutrient availability when ingested during a

bloodmeal (43). Other observations include WT upregulation of AKCT and L-threonine 3-

dehydrogenase, both of which are involved in the catabolic pathway converting L-threonine

into glycine (34). L-threonine is readily metabolised by procyclic forms and preferably

metabolised over other amino acids in glucose-depleted environments (44), where it is used

in lipid biosynthesis (45). Further reinforcing this metabolic change is TrAGEDy identifying

RBP42 as being significantly more highly expressed in the WT. This protein has been

implicated as a key regulator of metabolism associated genes (46), and it is significantly DE

in the final window along with the previously mentioned genes. Thus, TrAGEDy captured the

WT parasite pre-adaptation to the tsetse fly environment through the upregulation of genes

that will help the parasite respond to oxidative stress and alter metabolism, and failure of the

ZC3H20 KO parasites to do the same. Taken together, TrAGEDy captures DE genes not

captured by other differential expression test methods, revealing greater insight into

functions that differ between the WT and ZC3H20 KO *T. brucei* developmental programmes.

TI represents an approximation of how a developmental trajectory might look and may not

represent the actual ordering of the cells through the process. Adding extra temporal

information from lineage tracing will help add more biological validity to single cell data (47).

TrAGEDy could thus take lineage trace inferred time, rather than pseudotime, as it's input to

allow the analysis of such datasets, allowing it to remain relevant as more sophisticated

single cell sequencing techniques are introduced. Furthermore, the advent of perturb-seq

allows researchers to more easily create and analyse the effect of genetic perturbations on

processes (48), creating multiple and diverse opportunities for TrAGEDy to be applied.

Both TrAGEDy and cellAlign are restricted to aligning linear processes due to DTW only

being able to operate with linear processes. To overcome this limitation, branching

processes could be treated as multiple root-to-leaf linear processes and aligned and

analysed in a pairwise fashion but how accurate this process would be is uncertain as the

two paths are not independent of one another. Thus, future work needs to be carried out to

identify a method permitting the alignment of complex trajectories in order to better expand

the datasets which trajectory alignment can be applied to.

Conclusion

Our paper demonstrates the ability of TrAGEDy to overcome batch effects between datasets

and reveal the underlying alignment between two datasets, identifying shared and dissimilar

points across the independent trajectories. For some cases, TrAGEDy can identify more DE

genes than other methods, and in all cases, TrAGEDy can identify biologically relevant DE

genes between conditions and at specific points in the trajectory. With the advent of perturb-

seq methods, the need for TrAGEDy becomes more apparent and with lineage tracing it will

make the time axis more biological accurate. Thus, this tool increases the capacity to

analyse developmental progression with scRNA-seq to reveal novel detail.

520   Methods

521   The following is an overview of the steps that TrAGEDy takes to analyse a dataset. A

522   workflow diagram of the TrAGEDy process and its core steps can be seen in supplementary

523   figure 1 and implementation of the method can be seen, in full, in the code contained on our

524   GitHub.

526   Create interpolated points

Performing alignment directly on every cell in a dataset would not only be computationally

and time expensive, but also prone to noise. As such, the cellAlign method was used to

smooth gene expression over the trajectory by creating a user-defined number of

interpolated points that sample the gene expression patterns of surrounding cells at specific

points in the process and perform alignment on them.

In the cellAlign method, each interpolated point is given a set window of pseudotime around

it. Cells within that window contribute highly to the gene expression of the interpolated point,

while ones further away contribute less. In cellAlign, the size of the window is the same for

every interpolated point. For TrAGEDy, the user specifies a window size, but this value is

then weighted by the density of cells around each interpolated point. Interpolated points with

many cells around it will have a smaller window, while those with few surrounding cells will

have a larger window.

Scoring dissimilarity

For every interpolated point in the two trajectories, we calculate the transcriptome dissimilarity between it and all the interpolated points on the other trajectory and store it in a matrix. This can be done using any metric, e.g. Pearson correlation, Spearman correlation, Euclidean distance, given the right processing of the data (e.g. scaling prior of gene expression prior to Euclidean distance). For our experiments, Spearman correlation was used to assess dissimilarity. Hence, all the dissimilarity scales start from 0 (i.e. least dissimilarity), and we change the correlation scores to be 1 minus the correlation score.

Identifying the optimal path of the two trajectories

To find the optimal alignment between the interpolated points of the two trajectories, Dynamic Time Warping (DTW) was utilised. DTW requires that every point on the two processes is matched with at least one other point on the opposing process. It also requires that the first and last points of each process must be matched to one another. To tackle this constraint, TrAGEDy scans the first and last row and column of the dissimilarity matrix and identifies the indexes with the lowest score from the beginning and end. These become the initial start and ends points of the process. To make sure no valid matches are missed from the downstream analysis, TrAGEDy takes the following steps. First, TrAGEDy finds out how much the absolute dissimilarity changes across the row/column of the beginning and end changes as we move across them (supp. Fig.2A). The user then defines whether to find the median or mean of these points and this value is used as a cutting threshold. To give a value to each index to compare against the cutting threshold, TrAGEDy finds the absolute difference between the scores of each index in the chosen row/column and the current start/end point (supp. Fig.2B). If there is an index value which occurs before the current start point or after the current end point whose dissimilarity score is less than the threshold, it becomes the new start/end point. This repeats until all the points that come before/after it, have a dissimilarity score higher than the threshold (supp. Fig.2B). The matches that occur

before the new start point and after the new end point are removed and DTW is performed

on this cut dissimilarity matrix, assigning indexes to either be matched (part of the shared

process) or unmatched.

To deal with the scenario where two processes might differ somewhere in the middle of their

processes, TrAGEDy first finds two thresholds for the matched and unmatched indexes. The

user chooses to use either the mean or median of the dissimilarity score of all the

unmatched indexes and all the matched indexes. If a matched index's dissimilarity score is

closer to that of the unmatched threshold than the matched threshold then it is cut, otherwise

it remains matched (supp. Fig.2C).

Align pseudotime of interpolated points and cells

The aim of aligning the pseudotime of the interpolated points is to give matched interpolated

points similar pseudotime values. As we have two processes, we have two vectors:

$y_i^1\{i \in 1:m\}$ and, $y_j^2\{j \in 1:n\}$ where m and n are the number of interpolated points on

process 1 and 2 respectively, with y_i^1 being the pseudotime of the ith interpolated point in

process 1 and y_j^2 being the pseudotime of the jth interpolated point in process 2.

Interpolated points may only match one another (supp. Fig.2A) or be a part of a multi match,

where one interpolated point on one of the trajectories is connected to 2 or more interpolated

points on the other trajectory (supp. Fig.3B). For adjusting pseudotime of the former

scenario, given that y_i^1 and y_j^2 are matched and that $y_j^2 < y_i^1: y_{i:m}^1 = y_{i:m}^1 + (y_j^2 - y_i^1)$ (supp.

Fig.3A). For a multi match, first TrAGEDy aligns the pseudotime of the next match not

involved in the current multi match (supp. Fig.3B, equation 1), using the steps outlined for

the individual match, then it does the same for the first match in the multi match (supp.

Fig.3B, equation 2). The interpolated points that are multi matched to one interpolated point

on the other process, have their pseudotime values scaled between the aligned pseudotime

value of the first match in the multi match and the aligned pseudotime value of the match

that occurs after the multi match, adjusted by the difference between pseudotime of the next

match and the pseudotime of the first match in the multi match, normalized by the number of

interpolated points being scaled (supp. Fig.3B, equation 3). This leaves us with an aligned

pseudotime axis, where the matched interpolated points have similar pseudotime values

(supp. Fig.3C). We used the cellAlign method of mapping gene expression values from

individual cells onto interpolated points to map pseudotime values from interpolated points

onto the individual cells.

Differential expression analysis with TrAGEDy

TrAGEDy identifies DE genes across pseudotime, between the two conditions by taking a

sliding window soft clustering approach.

First the cells were assigned to the closest interpolated point in terms of pseudotime via one

round k-means clustering, with the interpolated points acting as the cluster centroid. The

interpolated points that were matched together were grouped, with the cells that were

assigned to these interpolated points acting as a cluster which the window slides over when

making comparisons. The user defines how many windows of comparison will be made

across the pseudotime, and how much overlap there will be, in terms of matched

interpolated points, between each successive window.

The user chooses a method of assessing significance using either a t-test or a Mann-

Whitney U test. Log2 fold change (Log2FC) of the genes between conditions is calculated

using the same formula used by Seurat. To adjust the p-values to account for multiple

testing, Bonferroni correction is used across all the genes in the dataset.

Simulating trajectories with Dyngen

All simulated datasets were simulated with Dyngen (9). The first positive control experiment

trajectories were simulated with a linear backbone with parameters drawn from the same

transcription factor, feature and kinetic network distributions. One of the datasets was not

modified beyond the standard pipeline, while the other had a full knockout in the B2 gene

module, causing a stunted process in comparison to the other dataset. The second positive

control experiment trajectories were simulated with a bifurcating-converging backbone with

parameters drawn from the same transcription factor, feature and kinetic network

distributions. One of the datasets had a full knockout in the C1 gene module while the other

dataset had a full knockout in the D1 gene module. The negative control experiment

trajectories were simulated with a linear backbone with parameters drawn from different

transcription factor, feature and kinetic network distributions. The specific parameters and

program calls used to simulate the individual datasets can be found on the GitHub link in the

"Availability of data and materials" section.

Applying TrAGEDy and cellAlign to simulated trajectories

All simulated datasets were processed using the following. In Seurat, the simulated counts

were normalized prior to scaling across all simulated genes. Principal component analysis

(PCA) was performed on the scaled data and using the first 15 PCA dimensions clustering

was performed at a resolution of 0.5 for all the datasets as well as UMAP. Slingshot (49) was

performed to get pseudotime values for each of the cells, using the UMAP space as the

basis for the trajectory. To get the feature space, the top 100 DE genes in terms of Log2FC

for each cluster in each dataset were collected for both datasets being compared. The prior

analysis steps were kept the same for both cellAlign and TrAGEDy analysis, as well as the

window size and the number of interpolated points. For cellAlign Euclidean distance and

Pearson correlation was used to calculate dissimilarity, while for TrAGEDy Spearman

correlation was used. Specific parameters such as number of dimensions used for each

dataset can be found on the GitHub linked in the "Availability of data and materials" section.

Pre-processing of WT vs ZC3H20 KO *Trypanosoma brucei* dataset

The processed RDS (R dataset) file containing the final cluster labels was acquired from the authors (https://zenodo.org/record/5163554#.Yyb113bMKUl). The 'LS A.1' and 'LS A.2' clusters were merged into the cluster 'LS A' and the 'LS B.1' and 'LS B.2' were merged into the cluster 'LS B'.

TrAGEDy analysis of WT vs ZC3H20 KO *Trypanosoma brucei* dataset

The integrated object was split into the individual datasets (WT01, WT02 and ZC3H20_KO). The top 300 DE genes across the clusters were found for each dataset and used as the basis for constructing the PHATE space and assessing similarity with TrAGEDy. The number of genes in the feature space amounted to 455 genes. PHATE embeddings were constructed from the normalised gene expression count matrix. Pseudotime was calculated using Slingshot for each of the three datasets, with the 'LS A' cluster used as the starting point. The WT01 and WT02 datasets were then aligned with TrAGEDy, with 50 interpolated points used to align the processes. This TrAGEDy aligned WT dataset was then aligned against the ZC3H20 KO dataset. Differential expression was carried out over 4 windows, with a gene identified as being DE if the absolute Log2FC was more than 0.5, bonferroni corrected p-value < 0.05 and the gene expressed in at least 10% of cells in one of the conditions.

Seurat analysis of WT vs ZC3H20 KO *Trypanosoma brucei* dataset

The 'SS A' and 'SS B' clusters were removed from the integrated dataset and the FindMarkers function was performed for all the four slender clusters between the WT and ZC3H20 KO conditions. A gene identified as being DE if the absolute Log2FC was more than 0.5, bonferroni corrected p-value < 0.05 and the gene expressed in at least 10% of cells in one of the conditions.

TradeSeq analysis of WT vs ZC3H20 KO *Trypanosoma brucei* dataset

The 'SS A' and 'SS B' clusters were removed from the integrated dataset. Using the authors PHATE space, Slingshot was applied to build trajectories. The optimal number of knots for fitting the general additive model to the dataset was assessed through the evaluateK function, with the optimal number determined to be 9. fitGAM was run on the dataset followed by the conditionTest function to identify DE genes between the WT and Bcl11b KO conditions. 4 knot comparisons were made, and a gene were determined to be DE if the absolute Log2FC was greater than 0.5, bonferroni p-value < 0.05 and the gene was expressed in at least 10% of the cells that fell within the current knot comparison, in one of the conditions.

For all cases, genes were analysed in TriTrypDB to get their gene short names and functions (50).

WT vs Bcl11b KO *in vitro* T cell development dataset

The RDS (R dataset) file containing the scRNA-seq information was acquired from the authors (10).

Preprocessing of WT vs Bcl11b KO *in vitro* T cell development dataset prior to TrAGEDy analysis

The dataset was split into the WT and Bcl11b KO conditions and then further split into the different sequencing runs. For each of the four scRNA-seq objects derived from the previous, the data was analysed using the Seurat V4 pipeline. When the data was scaled, the effects of the cell cycle were regressed out by passing the cell cycle associated genes Seurat provides into the var.to.regress parameter. For each dataset, PCA was carried out on the scaled-regressed expression matrix, reducing the dimensions down to 50. An elbow plot

was used to determine the number of principal components (PC) used to create a UMAP

space embedding. Clustering was performed using the same number of PCs for UMAP

embedding calculation. Clustree (51) was used to help choose a resolution for clustering that

leads to stable clusters. Clusters were annotated as follows. Clusters that did not express

Cd3 genes but expressed Cd34 were defined as 'Cd34_TSP', clusters which did not express

Cd34 or Cd3 were characterized as 'early_T', clusters which had low expression of Cd3 but

no Cd34 expression were characterised as 'Cd3_lo_T', those with high Cd3 expression were

'Cd3_T'. Clusters which had high expression of Cd3 and expressed Ptcra and Rag1 were

defined as 'rearrange_TCR' and those that expressed the TCR signal transduction molecule

Zap70 and some cells that express the gene Trac, were defined as 'signaling_T'. Clusters

that express Rora were defined as 'Rora_T', clusters that expressed high levels of interferon

associated genes were defined as 'Interferon_response' and cells which clustered out with

the main body of the UMAP were classed as 'Outlier'. These three clusters were removed

before downstream analysis with TI, TrAGEDy, Seurat DE or TradeSeq.

TrAGEDy analysis of WT vs Bcl11b KO *in vitro* T cell development

dataset.

In order to build a gene feature space to construct PHATE embeddings and assess

dissimilarity with TrAGEDy, the top 250 DE genes (in terms of Log2FC) for each annotated

cluster for each of the four datasets were extracted. To reduce the effect of the cell cycle,

genes supplied by Seurat as being important in the cell cycle and genes included in the MGI

GO term 'cell cycle' were removed from the top 250 DE genes extracted, leaving 952 genes.

For all four of the individual datasets, 10 dimensional phate embeddings were generated

from the normalised gene expression count matrix for the features selected previously. For

the WT1 dataset, cells with a pseudotime higher than 0.1 (11 cells) were removed. TI was

then carried out using Slingshot on the PHATE embeddings with the Cd34_TSP cluster

chosen as the starting point. TrAGEDy was carried out on each of the conditions, using 100

interpolated points with a window size of the maximum pseudotime value, from either of the

two datasets being analysed, divided by 90. TrAGEDy aligned pseudotime values were then

created, resulting in a TrAGEDy aligned WT and a TrAGEDy aligned Bcl11b KO dataset.

The WT and Bcl11b KO TrAGEDy aligned datasets were then analysed using TrAGEDy.

TrAGEDy was carried out using 100 interpolated points with a window size of the maximum

pseudotime value, from either of the two datasets being analysed, divided by 90. TrAGEDy

differential expression test was then carried out across 7 windows of comparison. A gene

was returned as being differentially expressed if the Log2FC < 0.5, Bonferroni adjusted p-

value < 0.05 and the gene was expressed in at least 10% of cells in the window of

comparison, in one of the conditions. The final TrAGEDy aligned WT and Bcl11b KO

datasets were then analysed independently by TradeSeq to create the smooth gene

expression plots across the aligned pseudotime.

Seurat analysis of WT vs Bcl11b KO *in vitro* T cell development dataset

Using Seurat V4 integration (52–54), the WT condition dataset was integrated based on their

sequencing run with the same carried out for the Bcl11b KO. The top 2000 integration

anchor genes were identified by Seurat and used as the basis of the integrated expression

space. When the data was scaled, the effects of the cell cycle were regressed out by

passing the cell cycle associated genes Seurat provides into the var.to.regress parameter.

The resulting two integrated datasets were then integrated together to create a final

integrated Bcl11b KO/WT dataset. The top 3000 integration anchor genes were identified by

Seurat and used as the basis of the integrated expression space.

The integrated gene expression values were then scaled, and the cell cycle regressed out as

done previously. PCA was carried out on the scaled-regressed expression matrix, reducing

the dimensions down to 50. An elbow plot was used to determine the number of principal

components (PC) used to create a UMAP space embedding. Clustering was performed

using the same number of PCs for UMAP embedding calculation. Clustree was used to help

choose a resolution for clustering that leads to stable clusters and Clusters were annotated

as described previously. The FindMarkers function was used to identify DE genes between

the conditions for each cluster. Genes were determined to be DE if the absolute Log2FC

was greater than 0.5, bonferroni p-value < 0.05 and the gene was expressed in at least 10%

of the cells in one of the conditions for the cluster being compared.

764 TradeSeq analysis of WT vs Bcl11b KO *in vitro* T cell development

765 dataset

Using the Seurat integrated, cell cycle regressed dataset; PHATE was carried out, using the

scaled and cell cycle regressed gene expression matrix. Slingshot was then applied to the

dataset, using the Cd34_TSP cluster as the starting point of the trajectory. The optimal

number of knots for fitting the general additive model to the dataset was assessed through

the evaluateK function, with the optimal number determined to be 8. fitGAM was run on the

dataset followed by the conditionTest function to identify DE genes between the WT and

Bcl11b KO conditions. 7 different knot comparisons were made, and a gene were

determined to be DE if the Log2FC was greater than 0.5, bonferroni p-value < 0.05 and the

gene was expressed in at least 10% of the cells that fell within the current knot comparison,

in one of the conditions.

777 Runtime experiments

778   Packages were run single threaded on a virtual machine with 70 gigabytes of allotted RAM

779   and 3 GHz clock speed. Runtime experiment code detailing what sections were included for

780   each package when calculating runtime can be found on our GitHub.

782 Package versions details

783 All analysis and experiments were carried out on R version 4.1.2 except for the Dyngen

784 dataset simulation and the TradeSeq experiments which were carried out on R version 4.1.0.

785 Seurat version 4.2.0, Dyngen version 1.0.5, PhateR version 1.0.7, Slingshot version 2.2.1,

786 Single Cell Experiment version 1.16.0, TradeSeq version 1.6.0 were used.

787 References

1.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics
and regulators of cell fate decisions are revealed by pseudotemporal ordering of
single cells. Nat Biotechnol [Internet]. 2014;32(4):381–6. Available from:
https://doi.org/10.1038/nbt.2859

2.   Saelens W, Cannoodt R, Todorov H, Saeys Y. A comparison of single-cell trajectory
inference methods. Nat Biotechnol [Internet]. 2019;37(5):547–54. Available from:
https://doi.org/10.1038/s41587-019-0071-9

3.   van den Berge K, Roux de Bézieux H, Street K, Saelens W, Cannoodt R, Saeys Y, et
al. Trajectory-based differential expression analysis for single-cell sequencing data.
Nat Commun [Internet]. 2020;11(1):1201. Available from:
https://doi.org/10.1038/s41467-020-14766-3

4.   Qiu X, Hill A, Packer J, Lin D, Ma YA, Trapnell C. Single-cell mRNA quantification and
differential analysis with Census. Nat Methods [Internet]. 2017;14(3):309–15.
Available from: https://doi.org/10.1038/nmeth.4150

5.   Song D, Li JJ. PseudotimeDE: inference of differential gene expression along cell
pseudotime with well-calibrated p-values from single-cell RNA sequencing data.
Genome Biol [Internet]. 2021;22(1):124. Available from:
https://doi.org/10.1186/s13059-021-02341-y

6.   Cano-Gamez E, Soskic B, Roumeliotis TI, So E, Smyth DJ, Baldrighi M, et al. Single-
cell transcriptomics identifies an effectorness gradient shaping the response of CD4+
T cells to cytokines. Nat Commun [Internet]. 2020;11(1):1801. Available from:
https://doi.org/10.1038/s41467-020-15543-y

7.   Briggs EM, Rojas F, McCulloch R, Matthews KR, Otto TD. Single-cell transcriptomic
analysis of bloodstream Trypanosoma brucei reconstructs cell cycle progression and
developmental quorum sensing. Nat Commun [Internet]. 2021;12(1):5268. Available
from: https://doi.org/10.1038/s41467-021-25607-2

8.   Alpert A, Moore LS, Dubovik T, Shen-Orr SS. Alignment of single-cell trajectories to
compare cellular expression dynamics. Nat Methods [Internet]. 2018;15(4):267–70.
Available from: https://doi.org/10.1038/nmeth.4628

9.   Cannoodt R, Saelens W, Deconinck L, Saeys Y. Spearheading future omics analyses
using dyngen, a multi-modal simulator of single cells. Nat Commun [Internet].
2021;12(1):3942. Available from: https://doi.org/10.1038/s41467-021-24152-2

10. Zhou W, Gao F, Romero-Wolf M, Jo S, Rothenberg E v. Single-cell deletion analyses
show control of pro–T cell developmental speed and pathways by Tcf7, Spi1, Gata3,
Bcl11a, Erg, and Bcl11b. Sci Immunol [Internet]. 2022 Dec 21;7(71):eabm1920.
Available from: https://doi.org/10.1126/sciimmunol.abm1920

11.   Seo W, Taniuchi I. Transcriptional regulation of early T-cell development in the
825   thymus. Eur J Immunol [Internet]. 2016 Mar 1;46(3):531–8. Available from:
826   https://doi.org/10.1002/eji.201545821
827   12.   Hwang JR, Byeon Y, Kim D, Park SG. Recent insights of T cell receptor-mediated
828   signaling pathways for T cell activation and development. Exp Mol Med [Internet].
829   2020;52(5):750–61. Available from: https://doi.org/10.1038/s12276-020-0435-8
830   13.   Mombaerts P, Iacomini J, Johnson RS, Herrup K, Tonegawa S, Papaioannou VE.
831   RAG-1-deficient mice have no mature B and T lymphocytes. Cell [Internet]. 1992 Mar
832   6;68(5):869–77. Available from: https://doi.org/10.1016/0092-8674(92)90030-G
833   14.   Moon KR, van Dijk D, Wang Z, Gigante S, Burkhardt DB, Chen WS, et al. Visualizing
834   structure and transitions in high-dimensional biological data. Nat Biotechnol [Internet].
835   2019;37(12):1482–92. Available from: https://doi.org/10.1038/s41587-019-0336-3
836   15.   Giotti B, Chen SH, Barnett MW, Regan T, Ly T, Wiemann S, et al. Assembly of a parts
837   list of the human mitotic cell cycle machinery. J Mol Cell Biol [Internet]. 2019 Aug
838   19;11(8):703–18. Available from: https://doi.org/10.1093/jmcb/mjy063
839   16.   Yuniati L, Scheijen B, van der Meer LT, van Leeuwen FN. Tumor suppressors BTG1
840   and BTG2: Beyond growth control. J Cell Physiol [Internet]. 2019 May 1;234(5):5379–
841   89. Available from: https://doi.org/10.1002/jcp.27407
842   17.   Rashida Gnanaprakasam JN, Wu R, Wang R. Metabolic Reprogramming in
843   Modulating T Cell Reactive Oxygen Species Generation and Antioxidant Capacity.
844   Front Immunol [Internet]. 2018;9. Available from:
845   https://www.frontiersin.org/articles/10.3389/fimmu.2018.01075
846   18.   Liau NPD, Laktyushin A, Lucet IS, Murphy JM, Yao S, Whitlock E, et al. The
847   molecular basis of JAK/STAT inhibition by SOCS1. Nat Commun [Internet].
848   2018;9(1):1558. Available from: https://doi.org/10.1038/s41467-018-04013-1
849   19.   Grujic M, Braga T, Lukinius A, Eloranta ML, Knight SD, Pejler G, et al. Serglycin-
850   deficient Cytotoxic T Lymphocytes Display Defective Secretory Granule Maturation
851   and Granzyme B Storage*. Journal of Biological Chemistry [Internet].
852   2005;280(39):33411–8. Available from:
853   https://www.sciencedirect.com/science/article/pii/S0021925820790708
854   20.   Korinfskaya S, Parameswaran S, Weirauch MT, Barski A. Runx Transcription Factors
855   in T Cells—What Is Beyond Thymic Development? Front Immunol [Internet]. 2021;12.
856   Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2021.701924
857   21.   Schilham MW, Moerer P, Cumano A, Clevers HC. Sox-4 facilitates thymocyte
858   differentiation. Eur J Immunol [Internet]. 1997 May 1;27(5):1292–5. Available from:
859   https://doi.org/10.1002/eji.1830270534
860   22.   B RD. V(D)J Recombination: Mechanism, Errors, and Fidelity. Microbiol Spectr
861   [Internet]. 2014 Nov 21;2(6):2.6.18. Available from:
862   https://doi.org/10.1128/microbiolspec.MDNA3-0041-2014
863   23.   Moran M, Miceli MC. Engagement of GPI-Linked CD48 Contributes to TCR Signals
864   and Cytoskeletal Reorganization: A Role for Lipid Rafts in T Cell Activation. Immunity
865   [Internet]. 1998;9(6):787–96. Available from:
866   https://www.sciencedirect.com/science/article/pii/S1074761300806445
867   24.   Lobach DF, Hensley LL, Ho W, Haynes BF. Human T cell antigen expression during
868   the early stages of fetal thymic maturation. J Immunol. 1985 Sep;135(3):1752–9.
869   25.   Aandahl EM, Sandberg JK, Beckerman KP, Taskén K, Moretto WJ, Nixon DF. CD7 Is
870   a Differentiation Marker That Identifies Multiple CD8 T Cell Effector Subsets1. The

871   Journal of Immunology [Internet]. 2003 Mar 1;170(5):2349–55. Available from:
872   https://doi.org/10.4049/jimmunol.170.5.2349

873 26.   Agresta L, Hoebe KHN, Janssen EM. The Emerging Role of CD244 Signaling in
874   Immune Cells of the Tumor Microenvironment. Front Immunol [Internet]. 2018;9.
875   Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2018.02809

876 27.   Dong K, Chen W, Pan X, Wang H, Sun Y, Qian C, et al. FCER1G positively relates to
877   macrophage infiltration in clear cell renal cell carcinoma and contributes to
878   unfavorable prognosis by regulating tumor immunity. BMC Cancer [Internet].
879   2022;22(1):140. Available from: https://doi.org/10.1186/s12885-022-09251-7

880 28.   Reuner B, Vassella E, Yutzy B, Boshart M. Cell density triggers slender to stumpy
881   differentiation of Trypanosoma brucei bloodstream forms in culture. Mol Biochem
882   Parasitol [Internet]. 1997;90(1):269–80. Available from:
883   https://www.sciencedirect.com/science/article/pii/S0166685197001606

884 29.   Rojas F, Silvester E, Young J, Milne R, Tettey M, Houston DR, et al. Oligopeptide
885   Signaling through TbGPR89 Drives Trypanosome Quorum Sensing. Cell [Internet].
886   2019;176(1):306-317.e16. Available from:
887   https://www.sciencedirect.com/science/article/pii/S0092867418313977

888 30.   Matthews KR. Trypanosome Signaling—Quorum Sensing. Annu Rev Microbiol
889   [Internet]. 2021 Oct 8;75(1):495–514. Available from: https://doi.org/10.1146/annurev-
890   micro-020321-115246

891 31.   Liu B, Kamanyi Marucha K, Clayton C. The zinc finger proteins ZC3H20 and ZC3H21
892   stabilise mRNAs encoding membrane proteins and mitochondrial proteins in insect-
893   form Trypanosoma brucei. Mol Microbiol [Internet]. 2020 Feb 1;113(2):430–51.
894   Available from: https://doi.org/10.1111/mmi.14429

895 32.   Cayla M, McDonald L, MacGregor P, Matthews K. An atypical DYRK kinase connects
896   quorum-sensing with posttranscriptional gene regulation in Trypanosoma brucei.
897   Engstler M, Soldati-Favre D, Acosta Serrano Á, editors. Elife [Internet].
898   2020;9:e51620. Available from: https://doi.org/10.7554/eLife.51620

899 33.   Mantilla BS, Marchese L, Casas-Sánchez A, Dyer NA, Ejeh N, Biran M, et al. Proline
900   Metabolism is Essential for Trypanosoma brucei brucei Survival in the Tsetse Vector.
901   PLoS Pathog [Internet]. 2017 Jan 23;13(1):e1006158-. Available from:
902   https://doi.org/10.1371/journal.ppat.1006158

903 34.   Marcus JP, Dekker EE. Threonine formation via the coupled activity of 2-amino-3-
904   ketobutyrate coenzyme A lyase and threonine dehydrogenase. J Bacteriol. 1993
905   Oct;175(20):6505–11.

906 35.   Fletcher DA, Mullins RD. Cell mechanics and the cytoskeleton. Nature [Internet].
907   2010;463(7280):485–92. Available from: https://doi.org/10.1038/nature08908

908 36.   Gupta SK, Chikne V, Eliaz D, Tkacz ID, Naboishchikov I, Carmi S, et al. Two splicing
909   factors carrying serine-arginine motifs, TSR1 and TSR1IP, regulate splicing, mRNA
910   stability, and rRNA processing in Trypanosoma brucei. RNA Biol [Internet]. 2014 Jun
911   1;11(6):715–31. Available from: https://doi.org/10.4161/rna.29143

912 37.   Tyler Weisbarth R, Das A, Castellano P, Fisher MA, Wu H, Bellofatto V. The
913   Trypanosoma cruzi RNA-binding protein RBP42 is expressed in the cytoplasm
914   throughout the life cycle of the parasite. Parasitol Res [Internet]. 2018;117(4):1095–
915   104. Available from: https://doi.org/10.1007/s00436-018-5787-9

916 38.   Claudia H, Corinna B, Stefanie B, Louise E, Van-Duc L, Mhairi S, et al. Small
917   Trypanosome RNA-Binding Proteins TbUBP1 and TbUBP2 Influence Expression of F-

Box Protein mRNAs in Bloodstream Trypanosomes. Eukaryot Cell [Internet]. 2007
Nov 1;6(11):1964–78. Available from: https://doi.org/10.1128/EC.00279-07

39.   Dean S, Marchetti R, Kirk K, Matthews KR. A surface transporter family conveys the
trypanosome differentiation signal. Nature [Internet]. 2009;459(7244):213–7. Available
from: https://doi.org/10.1038/nature07997

40.   Li Y, Li K, Zhu L, Li B, Zong D, Cai P, et al. Development of double-positive
thymocytes at single-cell resolution. Genome Med [Internet]. 2021;13(1):49. Available
from: https://doi.org/10.1186/s13073-021-00861-7

41.   Allen CE, Mak C ho, Wu LC. The κB transcriptional enhancer motif and signal
sequences of V(D)J recombination are targets for the zinc finger protein
HIVEP3/KRC: a site selection amplification binding study. BMC Immunol [Internet].
2002;3(1):10. Available from: https://doi.org/10.1186/1471-2172-3-10

42.   Peng HY, Lucavs J, Ballard D, Das JK, Kumar A, Wang L, et al. Metabolic
Reprogramming and Reactive Oxygen Species in T Cell Immunity. Front Immunol
[Internet]. 2021;12. Available from:
https://www.frontiersin.org/articles/10.3389/fimmu.2021.652687

43.   Geigy R, Huber M, Weinman D, Wyatt GR. Demonstration of trehalose in the vector of
African trypanosomiasis: the tsetse fly. Acta Trop. 1959;16:255–62.

44.   Lamour N, Rivière L, Coustou V, Coombs GH, Barrett MP, Bringaud F. Proline
Metabolism in Procyclic Trypanosoma brucei Is Down-regulated in the Presence of
Glucose*. Journal of Biological Chemistry [Internet]. 2005;280(12):11902–10.
Available from: https://www.sciencedirect.com/science/article/pii/S0021925820808990

45.   Millerioux Y, Ebikeme C, Biran M, Morand P, Bouyssou G, Vincent IM, et al. The
threonine degradation pathway of the Trypanosoma brucei procyclic form: the main
carbon source for lipid biosynthesis is under metabolic control. Mol Microbiol
[Internet]. 2013 Oct 1;90(1):114–29. Available from:
https://doi.org/10.1111/mmi.12351

46.   Das A, Morales R, Banday M, Garcia S, Hao L, Cross GAM, et al. The essential
polysome-associated RNA-binding protein RBP42 targets mRNAs involved in
Trypanosoma brucei energy metabolism. RNA. 2012 Nov;18(11):1968–83.

47.   Wagner DE, Klein AM. Lineage tracing meets single-cell omics: opportunities and
challenges. Nat Rev Genet [Internet]. 2020;21(7):410–27. Available from:
https://doi.org/10.1038/s41576-020-0223-2

48.   Schraivogel D, Gschwind AR, Milbank JH, Leonce DR, Jakob P, Mathur L, et al.
Targeted Perturb-seq enables genome-scale genetic screens in single cells. Nat
Methods [Internet]. 2020;17(6):629–35. Available from:
https://doi.org/10.1038/s41592-020-0837-5

49.   Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell lineage
and pseudotime inference for single-cell transcriptomics. BMC Genomics [Internet].
2018;19(1):477. Available from: https://doi.org/10.1186/s12864-018-4772-0

50.   Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bażant W, et al.
VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center.
Nucleic Acids Res [Internet]. 2022 Jan 7;50(D1):D898–911. Available from:
https://doi.org/10.1093/nar/gkab929

51.   Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. Gigascience [Internet]. 2018 Jul 1;7(7):giy083. Available from:
https://doi.org/10.1093/gigascience/giy083

52.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nat
Biotechnol [Internet]. 2018;36(5):411–20. Available from:
https://doi.org/10.1038/nbt.4096
53.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated
analysis of multimodal single-cell data. Cell [Internet]. 2021;184(13):3573-3587.e29.
Available from: https://www.sciencedirect.com/science/article/pii/S0092867421005833
54.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
Comprehensive Integration of Single-Cell Data. Cell [Internet]. 2019;177(7):1888-
1902.e21. Available from:
https://www.sciencedirect.com/science/article/pii/S0092867419305598
976

Acknowledgements

This research was funded by the Medical Research Council through their Precision Medicine

DTP (MRC grant number: MR/N013166/1). This research was funded in whole, or in part, by

the Wellcome Trust [Grant numbers 218648/Z/19/Z to EMB, 104111/Z/14/ZR to TDO/RM,

206815/Z/17/Z & 224501/Z/21/Z to RM, and 103740/Z14/Z to KRM] and the BBSRC Project

grants (BB/R017166/1 & BB/W001101/1 to RM).

Availability of data and materials

The functions to run TrAGEDy are available on our GitHub

(https://github.com/No2Ross/TrAGEDy) along with the code required to repeat our analysis.

Author Information

Authors and Affiliations

Wellcome Centre for Integrative Parasitology, Institute of Infection, Immunity and

Inflammation, University of Glasgow, Glasgow, UK

Ross F. Laidlaw, Emma M. Briggs, Richard McCulloch & Thomas D. Otto

Contributions

993   R.F.L. and T.D.O. conceived the project and R.F.L. implemtented the TrAGEDy package.

994   R.F.L., T.D.O., E.M.B. and R.M. designed experiments. R.F.L. performed the experiments

995   and collected data. R.F.L., T.D.O., E.M.B. and R.M. analyzed the data. R.F.L. wrote the

996   manuscript with editing from R.M., E.M.B., K.R.M. and T.D.O. All authors read, edited and

997   approved the manuscript.

998   Corresponding authors

999   Correspondence to Ross F. Laidlaw or Thomas D. Otto.

1000

1001   Ethics declarations

1002   Ethics approval and consent to participate

1003   Not applicable.

1004   Competing interests

1005   The authors declared that they have no competing interests.

1006   Supplementary figures and tables

1007

Process data

- Construct feature space
- Perform Trajectory Inference

Create interpolated points

- Get pseudotime values (cellAlign)
- Adjust sampling windows
- Get gene expression values (cellAlign)

Find alignment path

- Identify start and end points
- Run DTW
- Cut matches with high dissimilarity

Pseudotime adjustment

- Adjust pseudotime of interpolated points
- Adjust pseudotime of cells

Perform TrajDE

- Soft cluster cells
- Perform statistical tests
- Get summary statistics

Visualise

- Get window significance heatmap
- Do Upset plot over windows

1008

1009

1010 **Supplementary figure 1. Workflow of TrAGEDy process.**

1011 Workflow detailing the different parts of the TrAGEDy process. Sections which were derived

1012 from cellAlign are noted in the process boxes.

1013

1014

1015

Supplementary figure 2. Graphical explanation of the TrAGEDy process of modifying

the penalty matrix.

TrAGEDy finds the minimum dissimilarity score index from the beginning and end of the

matrix from either the row or column. The absolute change in dissimilarity score across the

row/column is captured and the median or mean threshold of change is found (A). To identify

whether a new index will become the start/end point we find the absolute difference between

each index and the current start/end point and identify a new start/end point if it lies below

the threshold (B). In order to cut matches from the DTW aligned path, a mean/median

threshold for the unmatched and matched indexes are calculated and if an index in the DTW

path is closer to the unmatched threshold than the matched, it is cut from the path (C).

1026

Supplementary figure 3. Graphical explanation of the alignment of interpolated point

pseudotime values, from the DTW calculated path.

There are two types of movement that can occur when aligning pseudotime values, an

individual match (A) or a multi-match (B). For the individual match, the pseudotime value of

the interpolated point with the highest value (and all the subsequent interpolated points) are

brought down by the difference between the lower pseudotime value and the higher

pseudotime value. For multi-match points, the match that occurs after it is adjusted then the

first match in the multi-match. All the interpolated points that are matched to the same

interpolated point are then scaled between pseudotime of the initial match in the multi-match

and the pseudotime of the next match modified by the difference between the two,

normalised by the number of interpolated points to be scaled. This then gives us aligned

pseudotime values for all the interpolated points (C).

Supplementary figure 4. cellAlign alignments of simulated datasets using Pearson

correlation dissimilarity score.

The alignments of the three simulated datasets experiments as calculated by cellAlign. WT

vs KO started shared simulation (A), KO C vs KO D diverge to converge simulation (B),

negative control A vs One simulation (C). Pearson correlation was used to assess

transcriptional dissimilarity between the conditions.

1047

1048

1049

1050

Supplementary figure 5. TrAGEDy alignment of biological replicates of WT slender to

stumpy *Trypanosoma brucei* transition.

TrAGEDy alignment of the WT01 and WT02 slender to stumpy *Trypanosoma brucei*

development trajectories. Dissimilarity in gene expression of interpolated points was

calculated using Spearman correlation with blue meaning low dissimilarity and red meaning

high dissimilarity.

1057

1058

1059 **Supplementary figure 6. Expression of markers associated with DN3 stage of T cell**

1060 **development in Bcl11b KO and WT T cells.**

1061 UMAP showing the normalised gene expression levels of markers of double negative 3

1062 (DN3) stage of T cell development: Ptcra (pre-TCR alpha chain) and Rag1 (recombination

1063 activating gene 1) in T cells under WT (left) and Bcl11b KO (right) conditions.

1064

Supplementary Figure 7. Impact of outlier cells on TrAGEDy alignments.

Plot showing the PHATE embeddings of the WT1 T cell dataset with each cell coloured by

its cell type annotation. Circle identifies cells which are separate from the main body of the

trajectory (A). TrAGEDy alignment of the WT1 and WT2 T cell datasets when the cells

circled in A are kept in the trajectory. Dissimilarity in gene expression of interpolated points

was calculated using Spearman correlation with blue meaning low dissimilarity and red

meaning high dissimilarity.

A

B

Supplementary figure 8. TrAGEDy alignment of WT and Bcl11b KO T cell sequencing

run trajectories.

TrAGEDy alignment of the WT1 and WT2 T cell development trajectories (A) and the Bcl11b

KO1 and Bcl11b KO2 T cell development trajectories (B). Dissimilarity in gene expression of

interpolated points was calculated using Spearman correlation with blue meaning low

dissimilarity and red meaning high dissimilarity.

1082

1083 Supplementary table information:

1084 Sup table 1 – the DE genes and associated statistics for TrAGEDy applied to T cell dataset

1085 for 0.25 and 0.5 Log2FC

1086

1087 Sup table 2 - the DE genes and associated statistics for TradeSeq applied to T cell dataset

1088 for 0.25 and 0.5 Log2FC

1089

1090 Sup table 3 - the DE genes and associated statistics for Seurat applied to T cell dataset for

1091 0.25 and 0.5 Log2FC

1092

1093 Sup table 4 – the DE genes and associated statistics for TrAGEDy applied to *T. brucei*

1094 dataset for 0.25 and 0.5 Log2FC

1095

1096 Sup table 5 - the DE genes and associated statistics for TradeSeq applied to *T. brucei*

1097 dataset for 0.25 and 0.5 Log2FC

1098

1099 Sup table 6 - the DE genes and associated statistics for Seurat applied to *T. brucei* dataset

1100 for 0.25 and 0.5 Log2FC

1101

1102 Sup table 7 – full list of genes in heatmap for T cell and *T. brucei* analysis