***Legionella pneumophila* inhibits type I interferon signaling to avoid cell-intrinsic host cell**

defense.

C.N.S. Allen*[1,] D. A. Banks*[1], M. Shuster[1], S. N. Vogel[2], T.J. O'Connor[3] and V. Briken[1]

[1] University of Maryland, Department of Cell Biology and Molecular Genetics

[2]Department of Microbiology and Immunology, University of Maryland School of Medicine,

Baltimore, Maryland, USA

[3] Johns Hopkins University School of Medicine, Department of Biological Chemistry

* authors contributed equally

Corresponding author: vbriken@umd.edu

Abstract:

The host type I interferon (IFN) response protects against a *Legionella pneumophilia* infection.
Other pathogens inhibit type I IFN-mediated cell signaling; however, the interaction between this
signaling pathway and *L. pneumophila* has not been described. This study demonstrates that *L.
pneumophila* inhibits the IFN-β signaling pathway in an IFN-β dose-dependent manner.
However, *L. pneumophila* does not inhibit IFN-γ-mediated cell signaling. The addition of IFN-β to
L. pneumophila-infected macrophages limited bacterial growth in a nitric oxide (NO)-dependent
pathway since inhibiting NO production via a chemical inhibitor increased bacterial growth.
Finally, this study demonstrates that the type IV secretion system of *L. pneumophila* is required
to inhibit IFN-β-mediated cell signaling. In conclusion, this study describes a novel host cell
signaling pathway inhibited by *L. pneumophila,* which improves the survival of bacteria in
infected macrophages.

Introduction:

Legionella species are ubiquitous in the environment and have adapted to survive within freshwater amoebas (Rowbotham, 1980; Boamah et al., 2017). The bacteria rely on their type 4 secretion system (T4SS), named the defective in organelle trafficking and intracellular multiplication (Dot/Icm), to inject proteins into the host cell to manipulate host response pathways and increase their chances of survival (Berger and Isberg, 1993; Brand et al., 1994; Segal and Shuman, 1998; Vogel et al., 1998). Legionella can infect a wide variety of amoeba species and, thus, through evolution, carry multiple versions of bacterial effectors to successfully modify the response of various hosts (Park et al., 2020). Humans are accidental hosts that can be exposed to Legionella via the aerosolization of contaminated water supplies, which often results in respiratory infections (McDade et al., 1977).

Type I interferons (IFNs), such as IFN-β, bind to the heterodimeric interferon-α/β receptor (IFNAR) and induce cell signaling, leading to the upregulation of hundreds of IFN-stimulated genes (ISGs) (Stetson and Medzhitov, 2006; Lazear et al., 2019). Type I IFNs are critical for the host defense against viral pathogens (Stetson and Medzhitov, 2006; Lazear et al., 2019). However, during bacterial infections, the activity of type I IFNs is more complex, where in some cases, they mediate host resistance, and in other cases, they increase the susceptibility (Boxx and Cheng, 2016; Peignier and Parker, 2021). L. pneumophila has been shown to induce IFN-β production by host cells via various cytosolic surveillance pathways, including the nucleic acid surveillance pathway (Liu and Shin, 2019). Type I IFNs then signal through IFNARs expressed on the cell surface of infected and bystander cells, contributing to L. pneumophila resistance observed in mice (Lippmann et al., 2011). Multiple studies have demonstrated that cell-intrinsic host defense mechanisms, including the production of nitric oxide (NO) (Plumlee et al., 2009) and itaconic acid (IA) (Naujoks et al., 2016), are triggered by exogenous IFN-β and subsequent IFNAR signaling during ex vivo L. pneumophila infections (Schiavoni et al., 2004;

Opitz et al., 2006; Plumlee et al., 2009; Majoros et al., 2016). Thus, the IFN-β response plays a

role in determining the outcome of *L. pneumophila*.

Here, we demonstrate that *L. pneumophila* has a capacity to inhibit IFN-β-mediated cell

signaling. Additionally, we show that the ability to reduce IFN-β signaling depends on the T4SS

of *L. pneumophila* and thus is likely to be mediated by secreted effectors. The inhibition of the

IFN-β response pathway improves the intracellular survival of *L. pneumophila* by limiting the

cellular production of NO.

Materials and Methods:

Cell culture and mice

$Ifnb^{-/-}$ mice were originally obtained from Dr. E. Fish (University of Toronto) and have been previously described (Deonarain et al., 2000). Bone marrow-derived macrophages (BMDMs) were prepared from bone marrow cells flushed from the tibias and femurs of these mice. The BMDMs were cultured in Gibco Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 1% penicillin/streptomycin, and 20% L929 conditioned medium for six days prior to infection. L929 conditioned medium was used for all the BMDM infection experiments. RAW 264.7 and RAW-Lucia ISG-KO-IRF3 cells (InvivoGen, rawl-koirf3) were grown in DMEM supplemented with 10% heat-inactivated FBS. RAW-Lucia ISG-KO-IRF3 cells were plated in FBS-free DMEM 16 hours before the infection since the absence of FBS reduces the background activation of the integrated ISRE-Luciferase gene cassette.

Ethics statement

All animals were handled in accordance with the NIH guidelines for the housing and care of laboratory animals. Studies were approved by the Institutional Animal Care and Use Committee at the University of Maryland.

Bacteria

L. pneumophila Philadelphia-1 strain Lp02 were cultured at 37°C in liquid N-(2-87Acetamido)-2-aminoethanesulfonic acid (ACES)-buffered yeast extract (AYE) medium or on solid charcoal ACES-buffered yeast extract (CYET) containing 0.4 mg/ml L-cysteine (Sigma), 0.135 mg/ml ferric nitrate (Sigma), and 0.1 mg/ml thymidine (Sigma). The *L. pneumophila* Lp02 *dotA- deficient* mutant was previously described (Berger and Isberg, 1993).

Cellular infections

L. pneumophila were grown in AYE medium to the post-exponential phase (A600 = 3.7–4.1). Raw 264.7 cells, RAW-Lucia ISG-KO-IRF3 cells, or BMDMs were infected with *L.*

pneumophila for two hours at the indicated multiplicity of infections (MOI). The cells were

washed three times with phosphate-buffered saline (PBS) and incubated in DMEM media

supplemented with or without IFN for the given time.

IFN-β enzyme-linked immunosorbent assay (ELISA)

The levels of secreted IFN-β were measured by ELISA (R&D Systems). RAW 264.7

cells were infected with *L. pneumophila* at an MOI of 0.05 for four hours. The supernatant of *L.*

pneumophila infected and uninfected cells were collected 24-, 48-, 72-, and 96-hours post-

infection (h.p.i). The ELISA was performed according to the manufacturer's instructions, and the

IFN-β signal was measured by a plate reader (BioTek ELx800).

Luciferase Reporter Assay

RAW-Lucia ISG-KO-IRF3 cells were infected with *L. pneumophila* at an MOI of 10 for 20

hours. IFN-β (R&D Systems) or IFN-γ (S&D Systems), at the indicated concentrations, was

added to the post-infection media at 0 h.p.i. The supernatant from the infected and uninfected

cells was collected, and the amount of type I IFN signaling was indirectly measured using

QUANTI-Luc Gold, following the manufacturer's protocol.

Bacterial Survival

Macrophages were infected with *L. pneumophila* at the indicated MOI. Cells were

collected at the provided time points and lysed in sterile water. The bacteria were enumerated

based on the colony forming units (CFUs) from plating cellular lysates on the CYE medium agar

plates.

Antibody neutralization

Antibody neutralization studies were performed using *InVivo*Plus anti-mouse IFNAR1

and *InVivo*MAB mouse IgG1 isotype control antibodies (Bio X Cell). Antibodies were added to

the infected cells at 0 h.p.i. at a final concentration of 20 µg/ml.

NO assay

The production of NO in response to IFN-β was quantified using a Griess assay kit

(Promega) following the manufacturer's protocol. Briefly, the supernatants from the infected

cells with or without IFN-β were collected at the indicated time points. The production of nitrite, a

derivative of NO, was measured at 548 nm using a microplate reader (BioTek). The nitrite

concentration was determined using a sodium nitrite standard curve.

Statistical Analysis

All statistical analyses were performed using GraphPad 9.0. For all the experiments, the

data are shown as the mean ± standard deviation (*SD*) of three biological replicates unless

otherwise noted. Significance is indicated by a *p*-value less than or equal to 0.05.

Results:

To test if *L. pneumophila* infections activated IFN-β production, RAW 264.7 cells were infected with *L. pneumophila* at an MOI of 0.05 for four hours, rinsed, and the amount of secreted IFN-β was measured at 24-hour intervals via an ELISA, where each data point is equivalent to a single round of infection. After 24 hours, 163.5 ± 55.2 pg/ml of IFN-β was detected and stayed constant through 48 h.p.i,, which was a ten-fold increase compared with the uninfected cells. At 72 h.p.i., the level of IFN-β increased to 342.4 ± 103.9 pg/ml and continued to rise to 460.9 ± 109.5 pg/ml by 96 h.p.i. (Figure 1A). These data indicate that *L. pneumophila* induces IFN-β production in RAW 264.7 macrophages and that IFN-β accumulates over time after the infection.

Since it has been previously reported that *Mycobacterium tuberculosis* and *Shigella sonnei* inhibits IFNAR signaling in infected cells (Banks et al., 2019; Alphonse et al., 2022) we aimed to determine if *L. pneumophila* inhibited type I interferon signaling. RAW-Lucia ISG-KO-IRF3 cells were used for these studies because they are deficient in *irf3*[-/-] which means that infection with *L. pneumophilia* will not induce production of endogenous IFN-β. Consequently, the only IFN-β detected by the IFN-β-inducible reporter gene is the IFN-β which we are adding exogenously to the cells. Cells were infected with *L. pneumophila* in the presence of increasing amounts of IFN-β (200–2000 pg/ml) for 20 hours to measure the effect *L. pneumophila* has on IFN-β-mediated cell signaling. The secretion of luciferase into the supernatant is representative of the activity of the IFN-β responsive promotor. The secreted luciferase was measured via a chemiluminescence assay. Comparing the response of infected cells to the uninfected cells, cells challenged with *L. pneumophila* showed lower levels of IFN-β-associated gene expression at all the doses of IFN-β tested (Figure 1B). The reduction in the transcriptional response was most significant at the lower doses of IFN-β, with a 75% decrease relative to the uninfected cells. At the highest IFN-β dose (2000 pg/ml), the transcriptional response between the infected and uninfected cells was reduced by 20% (Figure 1C). However, the IFN-β signaling pathway

reduction was reduced in a dose-dependent manner, suggesting that as the IFN-β concentration

increases, *L. pneumophila*'s ability to impede the signaling pathway is overwhelmed.

The IFN-β and IFN-γ signal transduction pathways share some of their intracellular

signaling components and thus we asked the question if *L. pneumophila* can also inhibit IFN-γ

signaling. The RAW-Lucia ISG-KO-IRF3 reporter construct is also induced by IFN-γ signaling

and could thus serve as a model to study the potential of *L. pneumophila* to inhibit IFN-γ

signaling. An increasing amount of IFN-γ (200–2000 pg/ml) was added to *L. pneumophila*-

infected or uninfected cells, and the luciferase production was analyzed (Figures 1C and D).

Overall, there was no reduction in the IFN-γ response in the infected cells; however, at IFN-γ

doses of 1000 and 2000 pg/ml, there was a significant increase in transcription response in the

L. pneumophila-infected cells compared with the uninfected cells (Figures 1D and E). These

results indicate that *L. pneumophila* partially inhibits IFN-β signaling while not affecting IFN-γ

signaling.

Since IFN-β has been shown to contribute to the clearance of *L. pneumophila* in mice

(Lippmann et al., 2011), *ifnb*[-/-]-derived BMDMs were infected with *L. pneumophila* and treated

with IFN-β to determine the effects of IFN-β on bacterial survival within macrophages. The CFUs

were determined from *L. pneumophila*-infected *ifnb*[-/-]-derived BMDMs that were untreated or

treated with 1000 pg/ml IFN-β. The untreated *ifnb*[-/-]-derived BMDMs cells showed a delayed

reduction compared with the IFN-β-treated cells in the percentage of CFUs starting at 48 h.p.i.

and continued until 96 h.p.i. (Figure 2A). The reduction in the percentage of CFUs was

significantly different between the IFN-β treated and untreated groups, indicating that IFN-β

signaling decreases *L. pneumophila* survival and enhances *L. pneumophila* clearance in *ifnb*[-/-]-

derived BMDMs. The decrease in CFUs in the untreated *ifnb*[-/-]-derived BMDMs could indicate

that another innate cellular immune response independent of IFN-β, such as TNFα (Liu et al.,

2008), contributes to the clearance of *L. pneumophila* in these cells.

To determine if the protective role of IFN-β is due to IFN-receptor signaling, *L.*

pneumophila-infected RAW-Lucia ISG-KO-IRF3 cells were treated with IFNAR1-neutralizing

antibodies with or without IFN-β. The RAW-Lucia ISG-KO-IRF3 cells were infected with *L.*

pneumophila at an MOI of 1, and then 20 µg/ml of IFNAR1-neutralizing or isotype control

antibodies were added. The IFN-β treated groups were then treated with 1000 pg/ml IFN-β. In

contrast to the BMDMs, RAW-Lucia ISG-KO-IRF3 cells have been shown to support the

intracellular growth of *L. pneumophila* (Gilchrist et al., 2020). The addition of IFN-β to the RAW-

Lucia ISG-KO-IRF3 cells did not affect the growth for the first two time points (24 and 48 h.p.i.),

but at 72 and 96 h.p.i. there was a significant reduction in the CFUs in the isotype IFN-β-treated

cells compared with the untreated isotype cells (Figure 2B). However, blocking the IFNAR

receptor by the IFNAR1-neutralizing antibodies inhibited the IFN-β-mediated reduction in the

CFUs (Figure 2B). These results indicate that IFN-β plays a role in the clarence of *L.*

pneumophila in macrophages and that this clarence results from signaling through the IFNAR1

receptor.

Nitric oxide (NO) has been shown to have cell-intrinsic anti-bacterial activity and limits

the growth of *L. pneumophila* (Plumlee et al., 2009). NO production has been shown to be

induced via the IFN-β signaling pathway in RAW 264.7 cells in response to lipopolysaccharide

through the induction of inducible nitric-oxide synthase (iNOS) (Jacobs and Ignarro, 2001).

Therefore, the NO production was analyzed using a Griess assay, which detects nitrite as a

proxy for the induced production of NO. *L. pneumophila*-infected RAW-Lucia ISG-KO-IRF3 cells

showed a slight increase in nitrite levels in the IFN-β-treated cells (1000 pg/ml) compared with

untreated cells at 24 h.p.i. (Figure 3A). After 48 hours, the IFN-β treatment increased the nitrite

level to about 18 µM; however, the nitrite level in the untreated cells was about 5 µM. The IFN-β

treatment led to nitrite production of about 25 µM at 96 hours, whereas the level in the untreated

cells remained low and was around 8 µM (Figure 3A). These results indicate that IFN-β

signaling induced NO production in the RAW-Lucia ISG-KO-IRF3 cells and that this production

is reduced in cells infected with *L. pneumophila* through the *L. pneumophila*-associated

inhibition of the IFNAR signaling pathway.

To determine if the reduced NO production influenced the clearance of *L. pneumophila,*

Raw-Lucia ISG-KO-IRF3 cells were infected with *L. pneumophila* and then were treated with or

without the iNOS inhibitor L-NMMA and with or without IFN-β. First, it was determined that L-

NMMA had a dose-dependent effect on the nitrite production of RAW-Lucia ISG-KO-IRF3 cells.

At the highest dose of L-NMMA, the nitrite production was approximately 12 μM, about 15% of

the nitrite produced by the untreated cells (Figure 3B). Next, RAW-Lucia ISG-KO-IRF3 cells

were infected with *L. pneumophila* and treated with or without IFN-β and with the solvent control

(DMSO) or 12 μM of L-NMMA to observe the effect of NO production on the survival of *L.*

pneumophila (Figure 3C). For all the experimental groups, a similar bacteria growth was

observed up to 48 h.p.i. At 72 h.p.i., there was a ten-fold reduction in the CFUs observed in the

DMSO and IFN-β-treated cells compared with the DMSO-treated cells. In contrast, the L-

NMMA-treated cells had a similar growth regardless of whether IFN-β was added (Figure 3C).

At the final measured timepoint of 96 h.p.i., the CFU difference between the DMSO and IFN-β-

treated cells was reduced about 100-fold compared with the DMSO-treated cells. However,

there was no significant fold reduction in the CFUs between the L-NMMA DMSO and L-NMMA

IFN-β-treated cells (Figure 3C). These results suggest that IFNAR signaling stimulates cellular

NO production, which leads to the clearance of *L. pneumophila*, and through the inhibition of the

IFN-β signaling pathway, *L. pneumophila* gains some resistance to the cellular anti-bacterial

response.

L. pneumophila T4SS Dot/Icm is a critical virulence factor that deploys effector proteins

to the host cell to manipulate host cellular processes (Berger and Isberg, 1993; Brand et al.,

1994; Segal and Shuman, 1998; Vogel et al., 1998). One mechanism by which *L. pneumophila*

might inhibit IFN-β signaling is through the activity of a secreted effector. To test this hypothesis,

an *L. pneumophila dotA-* deficient mutant was used to explore the importance of the Dot/Icm

system and, thus, the effectors in inhibiting the IFN-β signaling pathway. RAW-Lucia ISG-KO-

IRF3 cells were infected with wild-type *L. pneumophila* or a *dotA*-deficient mutant at an MOI of

10 and were treated with 600, 1000, or 2000 pg/ml of IFN-β for 20 hours (Figure 4A). The IFN-

β-mediated signaling was reduced in the cells infected with the wild-type bacteria by

approximately 60% for all the tested IFN-β concentrations (Figure 4A). In contrast, the *dotA*-

deficient mutant did not reduce the IFN-β-mediated signaling as much as the wild-type *L.*

pneumophila (Figure 4A). In addition to the wild-type and *dotA*- deficient mutant, RAW-Lucia

ISG-KO-IRF3 cells were infected with heat-inactivated wild-type and *dotA*- deficient mutant

bacteria to explore if the IFN-β inhibition requires live bacteria or if the inhibition is a result of

pre-existing/heat resistant bacterial lipids. The heat inactivation was achieved by incubating the

bacteria at 70°C for 30 minutes. The two heat-inactivated bacterial strains did not inhibit the IFN-

β signaling compared with the uninfected Raw-Lucia ISG-KO-IRF3 cells (Figure 4A). Overall,

these results suggest that the inhibition of the IFN-β pathway results from an effector secreted

through the Dot/Icm T4SS.

Next, the effect of impairing the T4SS on the capacity of infected cells to produce NO

upon IFN-β stimulation was analyzed. Uninfected RAW-Lucia ISG-KO-IRF3 cells treated with

100 ng/ml LPS and 1000 pg/ml IFN-β were used as a positive control. The wild-type *L.*

pneumophila-infected cells had significantly less NO production than the dot*A*- deficient mutant

and the positive control upon treatment with 1000 pg/ml IFN-β (Figure 4B). The level of NO

produced was not significantly different between the *dotA- deficient* mutant *L. pneumophila*-

infected cells and the positive control. Overall, these results demonstrate that the Dot/Icm T4SS

is necessary for the bacteria to inhibit IFN-β-mediated cell signaling and suggest that one or

more *L. pneumophila* effectors secreted into the host cell are responsible for inhibiting the

IFNAR signaling pathway.

Discussion

Type I IFNs have mainly been shown to play a protective role during viral infections

(Lazear et al., 2019); however, the role of type I IFNs is more complex in bacterial infections

since IFN-β has been shown to produce protective and detrimental effects depending on the

bacterial genus (Boxx and Cheng, 2016; Peignier and Parker, 2021). Previous studies have

identified other bacteria that inhibit IFN-β signaling through various means. For instance, *M.*

tuberculosis has been shown to inhibit IFN-β signaling by limiting the activation of IFNAR-

associated tyrosine kinase (Banks et al., 2019). Conversely, *Shigella* has been shown to inhibit

IFN signaling through the effector OspC, which binds to calmodulin, inhibiting calmodulin kinase

II and Stat1 phosphorylation (Alphonse et al., 2022). While previous studies have focused on

IFN-β and other bacteria, our work is the first to demonstrate that IFN-β plays a protective role

against an *L. pneumophila* infection and contributes to the clarence of *L. pneumophila.*

However, our results also demonstrate that *L. pneumophila* employs mechanisms to inhibit IFN-

β signaling to enhance their survival.

Our study suggests that *L. pneumophila* employs an effector(s) to inhibit the IFN-β

response since a T4SS-deficient mutant fails to inhibit IFNAR signaling. Future studies should

focus on defining the effector(s) and the mechanisms by which *L. pneumophila* disrupts the IFN-

β signaling pathway. *L. pneumophila* has adapted to survive in amoebas not human

macrophages. Additionally, amoebas do not have an equivalent receptor to IFNAR, or a class of

receptors associated with tyrosine kinases (Goldberg et al., 2006). The evolutionary advantage

for *L. pneumophila* to inhibit IFNAR-signaling when amoebas lack equivalent receptors is

currently unknown. However, amoebas, such as *Dictyostelium sp.* and *Acanthamoeba sp.,*

express a homolog to the mammalian STAT proteins (Kawata et al., 1997; Kicinska et al.,

2014), and STAT1 and STAT2 are secondary signaling components activated via

phosphorylation by receptor-associated kinases during IFNAR signaling (Lazear et al., 2019). In

addition, the amoeba STAT1 is closely related to the human and mouse STAT2 (Kicinska et al.,

2014), which is the STAT isoform involved in type I, but not type II, IFN signaling (Villarino et al.,

2020). Consequently, *L. pneumophila* could have evolved mechanisms to suppress STAT1-

mediated transcriptional regulation which allows the growth of *L. pneumophila* in amoebas, and

these mechanisms might have a similar effect on mammalian STAT2. Due to the homology of

mammalian STAT2 with the amoeba STAT1 (Kicinska et al., 2014), we hypothesize that IFNAR

signaling could be inhibited at the STAT2 level; however, future studies are needed to elucidate

the role of STAT2 in *L. pneumophila*-mediated inhibition of IFN-β signaling.

Acknowledgements

C.N.S.A., D.A.B., M.S. and V.B. were supported by grants R21AI142396 and R21 AI151973.

S.A.V. by grant R01 AI123371. M.S. was supported by "The Stefanie & Richard Vogel Graduate

Student Award" and T32 AI089621.

Fig. 1

Figure 1: *Legionella pneumophila* inhibits type I but not type II IFN signaling. (A) RAW

264.7 cells were infected with *L. pneumophila* (Lp), and the amount of secreted IFN-β was

detected by an ELISA. (B) Infected or uninfected cells were challenged with increasing

quantities of IFN-β, and the reporter ISG-response was measured after 20 hours post-infection

(h.p.i.). (C) The fold change of the raw data in B compared with the uninfected cells. (D) Infected

or uninfected cells were challenged with increasing quantities of IFN-γ, and the reporter ISG-

response was measured after 20 hours post-infection (h.p.i.). (E) The fold change of the raw

data in D compared with the uninfected cells. Individual points indicate the averages of

independent experiments. Statistical analyses were done using multiple unpaired *t*-tests with

Welch correction for each time point (A) or each dose concentration of IFN-β or IFN-γ (B, D).

The fold change is relative to the uninfected cells for the same treatment dose of IFN-β or IFN-

γ. Statistical analyses were done using multiple unpaired *t*-tests with Welch correction, and each

fold change was compared to the fold change of the uninfected cells for each treatment dose of

IFN-β or IFN-γ (C, E). Significance is indicated as follows: *$p \leq 0.05$, and **$p \leq 0.01$.

A

B

Figure 2: IFN-β-treatment increases cell-intrinsic resistance to *L. pneumophila*. (A) Bone

marrow-derived macrophages (BMDMs) from ifnb$^{-/-}$ mice were infected with *L. pneumophila* (Lp)

and treated with 1000 ng/ml IFN-β throughout the infection. The number of colony-forming units

(CFUs) was counted at the listed time points and normalized to the CFUs of the untreated

control cells at 0 hours post-infection (h.p.i.). (B) RAW-ISG-KO-IRF3 cells were infected with *L.*

pneumophila (Lp) and treated with 1000 ng/ml IFN-β throughout the infection. Isotype control

antibodies or IFNAR1-blocking antibodies and IFN-β were added to cells simultaneously. The

points are the mean ± standard deviation (*SD*) of two biological replicates. (A) An asterisk

indicates multiple unpaired *t*-tests with Welch correction relative to the untreated cells and (B)

comparing isotype ± IFN-β to the untreated isotype cells or α-IFNAR1 ± IFN-β to the untreated

α-IFNAR1 at each time point, where *$p \leq 0.05$ and **$p \leq 0.01$.

Figure 3: IFN-β-treatment increases cell-intrinsic resistance to *L. pneumophila* via increased NO production. (A) RAW-ISG-KO-IRF3 cells were infected with *L. pneumophila* (Lp) at an MOI of 10. The amount of produced nitrite was measured using a Griess assay and shown as the averages of three individual experiments. (B) The dose-response analysis of the iNOS2 inhibitor L-NMMA on RAW-ISG-KO-IRF3 cells is the averages of four independent experiments. (C) RAW-ISG-KO-IRF3 cells were infected with *L. pneumophila* (Lp) at MOI of 1 and treated with 1000 ng/ml IFN-β and solvent control (DMSO) or L-NMMA. The averages of three

333   independent experiments are shown. Statistical analyses were done using an unpaired *t*-test

334   with Welch correction per row with individual variances (A), a repeat-measure (RM) one-way

335   ANOVA (B), or an unpaired *t*-test with Welch correction comparing DMSO ± IFN-β or L-NMMA ±

336   IFN-β (C). Significance is indicated as follows: *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, and ****$p \leq$

337   0.0001.

Figure 4: The inhibition of IFN-β signaling by *L. pneumophila* is dependent on the type IV

secretion system. (A) RAW-ISG-KO-IRF3 cells were infected with wild-type *L. pneumophila*

(WT), the *L. pneumophila dotA*- deficient mutant (*dotA⁻*), the heat-inactivated wild-type *L.*

pneumophila (HI WT), or the heat-inactivated *dotA*- deficient mutant (HI *dotA⁻*) at an MOI of 10.

All experimental groups were treated with 1000 ng/ml IFN-β. The fold differences between the

wild-type, *dotA*- deficient mutant, heat-inactivated wild-type *L. pneumophila,* and heat-

inactivated *dotA*- deficient mutant were determined. Data points are shown as the averages of

three independent experiments. (B) RAW-ISG-KO-IRF3 cells were infected with the wild-type *L.*

pneumophila (WT) or the *L. pneumophila dotA*- deficient mutant (*dotA*⁻) at an MOI of 10 and

treated with or without 1000 ng/ml IFN-β. The production of NO was measured via a Griess

assay, and the data points are shown as the averages of three independent experiments. A two-

way ANOVA with Geisser-Greenhouse correction and Tukey's multiple comparisons test was

performed on all data. Significance is indicated as follows: *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$,

and ****$p \leq 0.0001$.

References

357
358

Alphonse, N., Wanford, J. J., Voak, A. A., Gay, J., Venkhaya, S., Burroughs, O., et al. (2022). A
family of conserved bacterial virulence factors dampens interferon responses by blocking
calcium signaling. *Cell* 185, 2354-2369.e17. doi: 10.1016/j.cell.2022.04.028.

Banks, D. A., Ahlbrand, S. E., Hughitt, V. K., Shah, S., Mayer-Barber, K. D., Vogel, S. N., et al.
(2019). Mycobacterium tuberculosis Inhibits Autocrine Type I IFN Signaling to Increase
Intracellular Survival. *Journal of immunology (Baltimore, Md. : 1950)*, ji1801303. doi:
10.4049/jimmunol.1801303.

Berger, K. H., and Isberg, R. R. (1993). Two distinct defects in intracellular growth
complemented by a single genetic locus in Legionella pneumophila. *Mol Microbiol* 7, 7–19.
doi: 10.1111/j.1365-2958.1993.tb01092.x.

Boamah, D. K., Zhou, G., Ensminger, A. W., and O'Connor, T. J. (2017). From Many Hosts,
One Accidental Pathogen: The Diverse Protozoan Hosts of Legionella. *Front Cell Infect Mi* 7,
477. doi: 10.3389/fcimb.2017.00477.

Boxx, G. M., and Cheng, G. (2016). The Roles of Type I Interferon in Bacterial Infection. *Cell
Host and Microbe* 19, 760–769. doi: 10.1016/j.chom.2016.05.016.

Brand, B. C., Sadosky, A. B., and Shuman, H. A. (1994). The Legionella pneumophila icm
locus: a set of genes required for intracellular multiplication in human macrophages. *Mol
Microbiol* 14, 797–808. doi: 10.1111/j.1365-2958.1994.tb01316.x.

Deonarain, R., Alcami, A., Alexiou, M., Dallman, M. J., Gewert, D. R., and Porter, A. C. G.
(2000). Impaired Antiviral Response and Alpha/Beta Interferon Induction in Mice Lacking
Beta Interferon. *Journal of Virology* 74, 3404–3409. doi: 10.1128/jvi.74.7.3404-3409.2000.

Gilchrist, A. K., Smolensky, D., Ngwaga, T., Chauhan, D., Cox, S., Perumal, R., et al. (2020).
High-polyphenol extracts from Sorghum bicolor attenuate replication of Legionella
pneumophila within RAW 264.7 macrophages. *Fems Microbiol Lett* 367, fnaa053. doi:
10.1093/femsle/fnaa053.

Goldberg, J. M., Manning, G., Liu, A., Fey, P., Pilcher, K. E., Xu, Y., et al. (2006). The
Dictyostelium Kinome—Analysis of the Protein Kinases from a Simple Model Organism. *Plos
Genet* 2, e38. doi: 10.1371/journal.pgen.0020038.

Jacobs, A. T., and Ignarro, L. J. (2001). Lipopolysaccharide-induced Expression of Interferon-β
Mediates the Timing of Inducible Nitric-oxide Synthase Induction in RAW 264.7
Macrophages*. *J Biol Chem* 276, 47950–47957. doi: 10.1074/jbc.m106639200.

Kawata, T., Shevchenko, A., Fukuzawa, M., Jermyn, K. A., Totty, N. F., Zhukovskaya, N. V., et
al. (1997). SH2 Signaling in a Lower Eukaryote: A STAT Protein That Regulates Stalk Cell
Differentiation in Dictyostelium. *Cell* 89, 909–916. doi: 10.1016/s0092-8674(00)80276-7.

Kicinska, A., Leluk, J., and Jarmuszkiewicz, W. (2014). Acanthamoeba castellanii STAT Protein.
Plos One 9, e111345. doi: 10.1371/journal.pone.0111345.

Lazear, H. M., Schoggins, J. W., and Diamond, M. S. (2019). Shared and Distinct Functions of
Type I and Type III Interferons. *Immunity* 50, 907–923. doi: 10.1016/j.immuni.2019.03.025.

Lippmann, J., Müller, H. C., Naujoks, J., Tabeling, C., Shin, S., Witzenrath, M., et al. (2011).
Dissection of a type I interferon pathway in controlling bacterial intracellular infection in mice.
Cell Microbiol 13, 1668–1682. doi: 10.1111/j.1462-5822.2011.01646.x.

Liu, M., Conover, G. M., and Isberg, R. R. (2008). Legionella pneumophila EnhC is required for
efficient replication in tumour necrosis factor α-stimulated macrophages. *Cell Microbiol* 10,
1906–1923. doi: 10.1111/j.1462-5822.2008.01180.x.

Liu, X., and Shin, S. (2019). Viewing Legionella pneumophila pathogenesis through an
immunological lens. *J Mol Biol* 431, 4321–4344. doi: 10.1016/j.jmb.2019.07.028.

Majoros, A., Platanitis, E., Szappanos, D., Cheon, H., Vogl, C., Shukla, P., et al. (2016).
Response to interferons and antibacterial innate immunity in the absence of tyrosine-
phosphorylated STAT1. *Embo Rep* 17, 367–382. doi: 10.15252/embr.201540726.

McDade, J. E., Shepard, C. C., Fraser, D. W., Tsai, T. R., Redus, M. A., and Dowdle, W. R.
(1977). Legionnaires' Disease — Isolation of a Bacterium and Demonstration of Its Role in
Other Respiratory Disease. *New Engl J Medicine* 297, 1197–1203. doi:
10.1056/nejm197712012972202.

Naujoks, J., Tabeling, C., Dill, B. D., Hoffmann, C., Brown, A. S., Kunze, M., et al. (2016). IFNs
Modify the Proteome of Legionella-Containing Vacuoles and Restrict Infection Via IRG1-
Derived Itaconic Acid. *Plos Pathog* 12, e1005408. doi: 10.1371/journal.ppat.1005408.

Opitz, B., Vinzing, M., Laak, V. van, Schmeck, B., Heine, G., Günther, S., et al. (2006).
Legionella pneumophila Induces IFNβ in Lung Epithelial Cells via IPS-1 and IRF3, Which
Also Control Bacterial Replication. *J Biol Chem* 281, 36173–36179. doi:
10.1074/jbc.m604638200.

Park, J. M., Ghosh, S., and O'Connor, T. J. (2020). Combinatorial selection in amoebal hosts
drives the evolution of the human pathogen Legionella pneumophila. *Nat Microbiol* 5, 599–
609. doi: 10.1038/s41564-019-0663-7.

Peignier, A., and Parker, D. (2021). Impact of Type I Interferons on Susceptibility to Bacterial
Pathogens. *Trends in microbiology*, 1–13. doi: 10.1016/j.tim.2021.01.007.

Plumlee, C. R., Lee, C., Beg, A. A., Decker, T., Shuman, H. A., and Schindler, C. (2009).
Interferons Direct an Effective Innate Response to Legionella pneumophila Infection*. *J Biol
Chem* 284, 30058–30066. doi: 10.1074/jbc.m109.018283.

Rowbotham, T. J. (1980). Preliminary report on the pathogenicity of Legionella pneumophila for
freshwater and soil amoebae. *J Clin Pathol* 33, 1179. doi: 10.1136/jcp.33.12.1179.

Schiavoni, G., Mauri, C., Carlei, D., Belardelli, F., Pastoris, M. C., and Proietti, E. (2004). Type I
IFN Protects Permissive Macrophages from Legionella pneumophila Infection through an
IFN-γ-Independent Pathway. *J Immunol* 173, 1266–1275. doi: 10.4049/jimmunol.173.2.1266.

Segal, G., and Shuman, H. A. (1998). Intracellular multiplication and human macrophage killing
by Legionella pneumophila are inhibited by conjugal components of IncQ plasmid RSF1010.
Mol Microbiol 30, 197–208. doi: 10.1046/j.1365-2958.1998.01054.x.

Stetson, D. B., and Medzhitov, R. (2006). Type I interferons in host defense. *Immunity* 25, 373–
381. doi: 10.1016/j.immuni.2006.08.007.

Villarino, A. V., Gadina, M., O'Shea, J. J., and Kanno, Y. (2020). SnapShot: Jak-STAT Signaling
II. *Cell* 181, 1696-1696.e1. doi: 10.1016/j.cell.2020.04.052.

Vogel, Joseph. P., Andrews, H. L., Wong, S. K., and Isberg, R. R. (1998). Conjugative Transfer
by the Virulence System of Legionella pneumophila. *Science* 279, 873–876. doi:
10.1126/science.279.5352.873.