1   **Transcriptomic and functional analysis of *ANGPTL4* overexpression in pancreatic**

2   **cancer nominates targets that reverse chemoresistance**

Authors: Emily R. Gordon[1], Carter A. Wright[1,2], Mikayla James[1], Sara J. Cooper[1*]

Affiliations:

[1]HudsonAlpha Institute for Biotechnology, 601 Genome Way, Huntsville, AL 35806

[2]The University of Alabama in Huntsville, 301 Sparkman Drive, Huntsville, AL 35899

*Corresponding Author:

Sara J. Cooper

HudsonAlpha Institute for Biotechnology, 601 Genome Way

Huntsville, AL 35806, USA

Email: sjcooper@hudsonalpha.org

phone: 256-327-9491

ORCID: 0000-0002-9627-0309

ABSTRACT

Background

Pancreatic ductal adenocarcinoma (PDAC) is one of the deadliest cancers based on five-

year survival rates. Genes contributing to chemoresistance represent novel therapeutic

targets that can improve treatment response. Increased expression of *ANGPTL4* in

tumors correlates with poor outcomes in pancreatic cancer.

Methods

We used statistical analysis of publicly available gene expression data (TCGA-PAAD) to test whether expression of *ANGPTL4* and its downstream targets, ITGB*4* and *APOL1*, were correlated with patient survival. We measured the impact of *ANGPTL4* overexpression in a common pancreatic cancer cell line, MIA PaCa-2 cells, using CRISPRa for overexpression and DsiRNA for knockdown. We characterized global gene expression changes associated with high levels of *ANGPTL4* and response to gemcitabine treatment using RNA-sequencing. Gemcitabine dose response curves were calculated on modified cell lines by measuring cell viability with CellTiter-Glo (Promega). Impacts on cell migration were measured using a time course scratch assay.

Results

We show that *ANGPTL4* overexpression leads to *in vitro* resistance to gemcitabine and reduced survival times in patients. Overexpression of *ANGPTL4* induces transcriptional signatures of tumor invasion and metastasis, proliferation and differentiation, and inhibition of apoptosis. Analyses revealed an overlapping signature of genes associated with both *ANGPTL4* activation and gemcitabine response. Increased expression of the genes in this signature in patient PDAC tissues was significantly associated with shorter patient survival. We identified 42 genes that were both co-regulated with *ANGPTL4* and were responsive to gemcitabine treatment. *ITGB4* and *APOL1* were among these genes. Knockdown of either of these genes in cell lines overexpressing *ANGPTL4* reversed the

observed gemcitabine resistance and inhibited cellular migration associated with

epithelial to mesenchymal transition (EMT) and *ANGPTL4* overexpression.

Conclusions

These data suggest that *ANGPTL4* promotes EMT and regulates the genes *APOL1* and

ITGB4. Importantly, we show that inhibition of both targets reverses chemoresistance and

decreases migratory potential. Our findings have revealed a novel pathway regulating

tumor response to treatment and suggest relevant therapeutic targets in pancreatic

cancer.

KEYWORDS

ANGPTL4, APOL1, ITGB4, gemcitabine, chemoresistance, pancreatic cancer, PDAC,

EMT, transcriptomic analysis, TCGA

BACKGROUND

Pancreatic ductal adenocarcinoma (PDAC) is among the deadliest cancers with a five-

year survival of only 11% (SEER) **[1]**. The high mortality of pancreatic cancer patients is

explained by the lack of early diagnostic markers leading to late-stage diagnosis of the

disease and the high rate of treatment resistance **[2]**. Standard of care includes resection

for about 20% of patients with localized disease who are eligible and treatment with

cytotoxic chemotherapies such as gemcitabine or combination agents like FOLFIRINOX

(5-FU, oxaliplatin, folinic acid, and irinotecan). This combination, when tolerated,

improves progression-free survival by 11 months **[3]**. Despite the aggressive treatment

of this disease with combinations of cytotoxic drugs, the vast majority of patients do not

have an enduring response to the treatment. Through improvements in precision cancer

treatments based on genetic mutations, implementation of combination therapies like

FOLFIRINOX, and advances in immunotherapy over the last decade the five-year survival

has increased from 6% to 11%, nevertheless, pancreatic cancer remains extremely

deadly. These factors combined with the increased incidence of risk factors like diabetes

and obesity. It is predicted that pancreatic cancer will be the second leading cause of

cancer deaths by 2030 **[4]**.

Doubling the five-year survival rate is remarkable progress, but novel treatment strategies

are still needed. One effective strategy is to identify cellular markers or pathways that are

enriched in pancreatic cancer cells. For example, over 90% of PDAC tumors have a

mutation in the KRAS gene, however, no KRAS targeting treatments are available and

existing targeted therapies like PARP inhibitors **[5]** and EGFR inhibitors **[6]** are limited to

a subset of patients. It is clear that targeted therapies are impactful for the eligible patients

with significant improvement in progression-free survival among patients receiving a

targeted treatment **[7]**. Unfortunately, patients who do not respond to standard

chemotherapeutics and are not eligible for existing targeted therapies are left with few

options. Thus, there is significant interest in understanding the development of resistance,

and in pursuing targets that might sensitize resistant cells to existing treatments.

Our previous transcriptomic study identified genes whose expression was positively or

negatively correlated with patient survival and also linked to *in vitro* response to

gemcitabine, a common PDAC treatment. We identified ANGPTL4 among those genes

whose expression in patient tumors is associated with poor survival and whose

knockdown in PDAC cell lines could increase sensitivity to gemcitabine **[8]**. ANGPTL4 is

a member of the family of angiopoietin-like proteins that were first described for their role

in angiogenesis **[9, 10]**. ANGPTL4 can be proteolytically cleaved **[11]** and the two

products each have their own functions. The N-terminal domain plays a role in lipid

metabolism and genetic variation in this domain is linked to cardiovascular disease risk.

The C-terminal domain has been implicated in tumorigenesis, the promotion of

proliferation, and wound healing **[11, 12, 13]**.

Overall, the role of ANGPTL4 in cardiovascular disease and lipid metabolism has been

much better described than its roles in cancer. ANGPTL4 has a described role in known

cancer pathways including its ability to regulate CREB, FOS, and STAT3 via ERK

signaling **[14, 15]**. ANGPTL4's ability to alter metabolism and ATP abundance can also

impact drug transport **[15]**. The picture, however, is not perfectly clear since ANGPTL4

expression seems to have different impacts in cancers of different primary sites. For

example, methylation and downregulation of ANGPTL4 are associated with progression

and metastasis in colon cancer **[16]** but overexpression is associated with progression

and poor prognosis in breast cancer **[17]**. Breast cancers of the triple negative subtype

may be different since ANGPTL4 overexpression in that context has been associated with

inhibition of invasion **[18]**. In pancreatic cancer, *ANGPTL4* overexpression has been

associated with tumorigenesis **[19]**, cellular resistance to chemotherapy **[8]**, hypoxia

response, and poor patient outcomes **[20]**. The complicated role for ANGPTL4 motivates

our further exploration of its function in pancreatic cancer.

Here, we describe the role of ANGPTL4's in PDAC by exploring the molecular and cellular

changes associated with altered activity of ANGPTL4, the impact of ANGPTL4 on

chemoresistance, and the potential for inhibiting downstream pathways driven by

ANGPTL4 activation to sensitize tumor cells to treatment. We show that overexpression

of *ANGPTL4* leads to chemoresistance, increased migratory potential, and proliferation.

Our transcriptomic analysis revealed altered gene expression signatures downstream of

ANGPTL4 overexpression that are linked to epithelial to mesenchymal transition (EMT)

and predict patient outcomes. We showed that knockdown of downstream effectors

including APOL1 and ITGB4 reversed resistance to treatment and reduced migratory

potential. The expression of these genes is also associated with patient survival. These

data support the hypothesis that ANGPTL4 and its downstream pathways are potential

therapeutic targets for the reversal of treatment resistance in pancreatic cancer.

METHODS

Cell Culture

HEK293FT cells (ThermoFisher #70007), MIA PaCa-2 cells (ATCC #CRM-CRL-1420),

and Panc-1 (ATCC #CRL-1469) were cultured in D10 media: DMEM (Lonza #12-614Q)

supplemented with 10% FBS, and 0.5% Penicillin-Streptomycin. All cell lines were

maintained at 37 °C and 5% CO_2. Cells were cryopreserved with the addition of 10%

DMSO (EMD #MX1458-6) to D10 media.

Plasmids

LentiSAMv2 (Addgene #92062) and lenti-MS2-p65-HSF1-Hygro (Addgene #89308) were

used to generate stable cell lines for gene activation. pMD2.G (Addgene #12259) and

psPAX2 (Addgene #12260) were used to facilitate viral packaging of sgRNA vector

plasmids.

sgRNA Cloning

sgRNA oligos were designed and cloned into their respective plasmids as described

previously (ANGPTL4: 5'-CACGGCCCTGGGGATGCCAAACTGTGG-3' and NTC: 5'-

ACGGAGGCTAAGCGTCGCAA-3') **[21]**.

DsiRNA

IDT TriFECTa RNAi kit was used per manufacturer's protocol. The DsiRNA sequences

used were as follows: ANGPTL4 (IDT hs.Ri.ANGPTL4.13.1), APOL1 (IDT

hs.Ri.APOL1.13.3), and ITGB4 (IDT hs.Ri.ITGB4.13.3). 100,000 cells were seeded per

well of a 12 well tissue culture treated plate 24 hours prior to transfection. Cells were

transfected using RNAiMax (ThermoFisher #13778-030) following manufacturer's

recommended protocol. As directed in the TriFecta kit (IDT #hs.Ri.HDAC1.13), TYE 563

transfection efficiency control, positive HPRT-S1 control, and negative (DS NC1)

scrambled sequence control were utilized. Functional assays were performed 48 hours

after transfection. Expression was validated with each transfection with IDT PrimeTime

qPCR Assay system according to manufacturer's recommendations on an Agilent

QuantStudio 6 Flex Real-Time PCR system (ANGPTL4: Hs.PT.58.40146104, ACTB:

Hs.PT.39a.22214847, GAPDH: Hs.PT.39a.22214836, and HPRT:

Hs.PT.58v.45621572).

Protein Quantification using AlphaLISA

Briefly, cells were seeded at 25,000 cells in 50uL of media per well of a 96-well tissue-

culture treated plate in triplicate per cell type. The following day, the media from each well

was transferred to a non-treated 96-well plate and 25ul of AlphaLISA Lysis Buffer

(PerkinElmer, #AL001C) was added per well. Plates were then shaken at 250 RPM at

room temperature for 10 minutes. Lysates were immediately used. Samples or standards

were transferred to a 384 well AlphaPlate (PerkinElmer, #600535, lot:8220-21331) and

the AlphaLISA ANGPTL4 manufacturer's kit protocol (PerkinElmer, #AL3017HV,

lot:2905318) was followed for standards, lysate, and supernatant (media) except for the

following: lids were covered in foil to prevent light exposure when possible and during

incubation periods plates were shaken at 150 RPM for the recommended time. The

AlphaLISA plate was read on a BioTek Synergy H5 following previously published

protocols **[22]**. Data were analyzed in R (version 4.1.2) and GraphPad Prism 9.

RNA-sequencing

Cells were seeded at a density of 0.6x10^6 cells per well of a 6-well plate in triplicate. The

following day media was removed and 3ml of media per well with or without 1.5nM of

gemcitabine (Sigma #G6423) was added and incubated for 24 hours. Media was

removed, cells were washed with PBS (Gibco #10010072), and released from the plate

with 2 ml of TrypLE (Gibco #12604-021) per well. After centrifugation, pellets were frozen

at -80C until RNA extraction. For RNA extraction, of cell pellets 350 ul of RL Buffer plus

1% BME from the Norgen Total RNA extraction kit and extraction proceeded per

manufacturer's instructions including use of the DNase kit (Norgen # 37500, 25720). RNA

quality was verified with the Agilent BioAnalyzer RNA Nano 600 kit (cat# 5067-1512) with

the RIN range between 9.2-10. RNA-sequencing libraries were made using Lexogen

QuantSeq 3' mRNA-Seq Library Prep Kit FWD for Illumina kit (cat# 015.24) with 250 ng

of RNA input. They were pooled and sequenced on an Illumina NextSeq 500 instrument

with 75 bp single-end reads. Read counts averaged 4 million reads and 93.65% of bases

exceeding Q30. Lexogen's BlueBee integrated QuantSeq data analyses pipeline was

used for trimming, mapping, and alignment and DESeq2 was used for differential

expression **[23]**. Heatmaps were generated using iDEP.95 **[24]**.

Scratch-wound Assay

Cells were seeded at 100,000 cells per well of a 96-well plate. After 24 hours uniform

scratches were made across the diameter of the wells using a multichannel pipette with

200ul pipette tips and even pressure applied across the wells to cause a wound. The

media was then vacuumed off and 200ul of new media was added. Cells were imaged on

a Lionheart FX Automated Microscope every 8 hours for 48 hours. Cell culture growth

conditions of 37 °C and 5% CO_2 were maintained throughout. Forty images were taken of

each scratch (4 wide by 10 long) with no overlap, autofocus, and auto brightness.

Images were integrated using R and the "tiff" package. Images were further processed

using GIMP 2.10. ImageJ and the MRI_wound_Healing_Tool.ijm macro plugin **[25]** were

used to process the images and calculate the area of the wound the cells have not

covered. Relative wound closure over time using time 0 for each condition as the control

was plotted and a curve line equation was formed by fitting the curve to a non-linear fit

one phase decay least squares fit with Yo=0 as a constraint. This base equation (Y=(Y0

- Plateau)*exp(-K*X) + Plateau) where Y0 is the Y value when X (time) is zero, Plateau is

the Y value at infinite times, and K is the rate constant, was used to determine the value

of X or half the time it takes to close the wound. Data were analyzed using GraphPad

Prism 9.

Drug Resistance Screening

Cells were seeded in 96-well plates at 2000 cells/well. Seeded cells were treated with a

range of gemcitabine concentrations. Cells were treated again 48 hours later. The number

of viable cells surviving drug treatment were measured with CellTiter-Glo (Promega

#G7571) 24 hours after the last drug treatment per the manufacturer's protocol using a

BioTek Synergy H5 plate reader. Sample size ranged from 4-11.

Pathway Analysis

1198 DEG from the ANGPTL4 OE vs KD analysis were imported into the KEGG Mapper-

Color **[26]** and Panther GO Enrichment Analysis **[27]**. The following parameters were

used for KEGG: Search mode: hsa, used uncolored diagrams, included aliases. The

following parameters were used for PANTHER: overrepresentation test, homo sapiens

reference list, GO biological process complete, Fisher's exact test, and calculate FDR.

Top pathways were merged manually and drawn using BioRender.

Overall Survival Analysis (OS)

To conduct survival analysis, clinical and RNA-seq expression data was retrieved from The

Cancer Genome Atlas for 178 PDAC (TCGA-PAAD) patients (https://portal.gdc.cancer.gov/).

Data were normalized using the R package DESeq2 and differentially expressed genes

with an FDR < 0.1 were used to generate Kaplan-Meier survival curves. We classified

tissues based on their average expression of a given gene set (bottom 25%, middle 50%,

and top 25% of gene expression). We compared the patients with the lowest and highest

quartile of average gene expression and performed survival analysis. Survival curves and

analyses were generated using the "ggplot2", "survminer", and "survival" R packages. P

values were generated using a log rank test.

Recurrence Free Survival (RFS)

A Kaplan Meier curve of recurrence-free survival survival plots for ANGPTL4 was created

using GEPIA (http://gepia.cancer-pku.cn/) single gene analysis. The relevant parameters

were as follows: Disease-free Survival (RFS), Group Cutoff: Quartile (75% high, 25%

low), Hazards Ratio: Yes, 95% Confidence Interval: Yes, Axis Units: Months, and

datasets Selection: PAAD.

Correlation Analysis

To compute correlation matrices, RNA-seq expression data was retrieved from The

Cancer Genome Atlas for 178 PDAC (TCGA-PAAD) patients

(https://portal.gdc.cancer.gov/).

Rank-based correlation coefficients were computed for each of the DEG with $\log_2$ fold

change greater than 0.7 and baseMean>10 (MP2_ANGPTL4_KD vs

MP2_ANGPTL4_OE with or without treatment) for 1) all TCGA-PAAD data DEG per gene

vs ANGPTL4 expression and 2) all TCGA-PAAD data DEG gene versus OS time. These

correlation values were used to generate a list of genes that are co-expressed with

ANGPTL4 is overexpressed.

Correlation coefficients and p-values were computed using the "Hmisc" (version 4.6-0)

R package. We classified tissues based on their ANGPTL4 expression of a given gene

set (bottom 25%, middle 50%, and top 25% of gene expression). We computed

correlation matrices using the expression data for patients with the lowest and highest

quartile of ANGPTL4 expression which was used to divide samples into quantiles with

highest and lowest average gene expression; based on expression of these 42 genes.

(40 genes were measured in TCGA: KDM7a and STN1 were omitted).

Statistical Analysis

Statistical analysis was conducted in R (version 3.6.1 and R version 4.1.2 for RNAseq

analysis). The following R packages and software were used for analysis:

survival (version 1.2.1335) **[28]**

survminer (version 0.4.9) **[29]**

ggplot2 (version 3.3.6) **[30]**

DESeq2 (version 1.24.0) **[31]**

pheatmap (version 1.0.12) **[32]**

Hmisc" (version 4.6-0) **[33]**

tiff (version 0.1-11) **[34]**

RESULTS

ANGPTL4 **overexpression increases gemcitabine resistance genes in PDAC**

We previously showed that increased *ANGPTL4* expression in tumors is associated with

poor survival in PDAC patients (n=51) **[8]**. Here, we confirm the association with survival

in an independent cohort of 178 PDAC patients from The Cancer Genome Atlas (TCGA,

PAAD expression dataset). Patients with the highest (top 25%) expression of *ANGPTL4*

in tumor tissue had reduced overall survival (OS) and disease-free survival, (p=0.043 and

p=0.059 (log-rank test) (**Fig. 1A**, **Supplementary Fig. S1A)**.

Our previous findings motivated the current study to understand possible mechanisms by

which ANGPTL4 contributes to patient survival and specifically how it alters survival

outcomes through a possible role in drug resistance. We measured the impact of

ANGPTL4 overexpression and knockdown in the MIA PaCa-2 cell line using CRISPRa

(MP2_ANGPTL4_OE) **[21]** and siRNA knockdown (MP2_ANGPTL4_KD). qPCR showed

CRISPRa successfully increased expression of the ANGPTL4 transcript by 8-fold

compared to a non-targeting control guide. Similarly, siRNA knockdown reduced

expression by over 90% of the control line (**Fig. 1B, Supplementary Table S1**). An ELISA

assay showed that protein levels were affected similarly (**Supplementary Fig. S1B**). We

measured global gene expression changes using RNA-sequencing on the modified cell

lines and controls which revealed 1198 differentially expressed genes (DEG) with a mean

read count greater than 10, absolute $\log_2$Fold Change (log2FC) of at least 0.7, and

adjusted p-value (padj) less than 0.05) when comparing overexpressed (OE) ANGPTL4

cells with the knockdown (KD) ANGPTL4 cells (**Supplementary Fig. S1C,**

Supplementary Table S2). Given the observed association with drug resistance, we

asked whether these DEGs were also associated with gemcitabine resistance by

overlapping the 1198 DEG with 114 known gemcitabine resistance genes

(**Supplementary Table S3**) **[35, 36, 37, 38]**. This revealed 15 gemcitabine resistance

genes that are altered with overexpression of *ANGPTL4*, 10 of which overlap genes that

are part of the EMTome (**Fig. 1C) [39].** Several of these genes are linked to activation of

TGF**β** and its downstream signaling pathways: ERK, PI3K/AKT, and JNK (SH3RF2,

TGFBR3, PPP1R15A, INPP4B, WWTR1, and FGF2), contributors to EMT (SOX9, CUL7,

and TGFBR3), and transcriptional regulation of pluripotent stem cells (STAT3, KLF4, and

FGF2) **[26, 27]**. Additional pathway analysis showed a connection between the

expression of ANGPTL4 and EMT (GO:0001837) in the presence (3.24-fold enrichment,

0.0273 FDR) or absence (3.04-fold enrichment, 0.195 FDR) of gemcitabine treatment

(**Supplementary Table S4-S5**).

Figure 1

Figure 1: Expression of *ANGPTL4* is associated with patient outcomes and modifies expression of resistance-associated genes.

a) A Kaplan Meier curve for overall survival using TCGA-PAAD data. The patients with the top (red) and bottom (blue) 25% average *ANGPTL4* gene expression have significantly different OS. (p=0.043, log-rank test) **b)** Relative expression of ANGPTL4 transcript in MP2_ANGPTL4_KD (black circles), MP2 (pink squares), and MP2_ANGPTL4_OE (turquoise triangles) as measured by qPCR. P-values from unpaired, two-tailed parametric t-tests performed with 95% confidence intervals are displayed. Values are normalized with *ACTB* as the housekeeper. (P-values ≤

0.0001=****) **c)** Heatmap displaying RNA-seq results from MP2_ANGPTL4_KD (black),

MP2_NTC (pink), and MP2_ANGPTL4_OE (turquoise) cell lines. Displayed are 15 genes

associated with both ANGPTL4 overexpression and gemcitabine resistance. Normalized

expression for each of the 15 genes is plotted in the heatmap. Red (*) indicates ITGB4

used for later experiments.

Increased *ANGPTL4* expression alters cellular response to gemcitabine

We assessed whether altering *ANGPTL4* expression impacted response to gemcitabine

to explore the potential impact of *ANGPTl4* overexpression on treatment response in

patients. We measured cell viability in MP2_ANGPTL4_KD and MP2_ANGPTL4_OE cell

lines before and after treatment with gemcitabine. As expected, treatment with 12.5nM

gemcitabine reduces viability compared to untreated cells. Comparing cells with

overexpression of *ANGPTL4* to controls, we observed an increase in cell viability with the

same gemcitabine concentrations. Similarly, there was a significant reduction in viability

with the *ANGPTL4* knockdown compared to cells overexpressing *ANGPTL4*. This

difference in viability is not observed in the untreated cells **(Fig. 2A, Supplementary**

Table S1). These results demonstrate a greater sensitivity to gemcitabine with reduction

of ANGPTL4. To understand the link between ANGPTL4 expression and gemcitabine

response, we measured transcriptional response to gemcitabine in MP2_ANGPTL4_OE

cells and MP2_ANGPTL4_KD cells using RNA-sequencing. We were not surprised to

note that 955 genes were differentially expressed upon treatment with the cytotoxic

gemcitabine alone. We explored the intersection of the 1198 DEGs identified upon

ANGPTL4 overexpression and the 955 DEGs associated with gemcitabine treatment.

This analysis revealed 42 genes associated with gemcitabine response and *ANGPTL4*

overexpression (**Fig. 2B, Supplementary Table S6).** This strategy nominated genes that

might contribute to altered response to gemcitabine in cells overexpressing *ANGPTL4*

and are potentially important for patient outcomes. We tested whether these genes were

predictive of patient overall survival using data from TCGA. Since all 42 transcripts were

positively correlated with ANGPTL4 expression, we calculated the mean gene expression

of these genes and found that patients whose tumors had the highest average expression

had significantly reduced overall survival (p=0.0073, **Figure 2C**). The fraction of patients

surviving 1500 days or more is dramatically increased in the individuals with low

expression of these 42 ANGPTL4 and gemcitabine response genes. This compares

favorably to the average 3-year relative survival for PDAC patients in the United States

with resectable stage I/II disease- 17% **[40]**.

Figure 2

Figure 2: Gemcitabine resistance signature in PDAC cells with high ANGPTL4

a) Cell viability was measured for MP2_ANGPTL4_KD (black), MP2_NTC (pink), and MP2_ANGPTL4_OE (turquoise) cells treated with 0, 5nM, or 12.5nM gemcitabine. Relative viability reflects normalization to cells not treated with gemcitabine. P-value asterisks: ≤0.01=**, ≤0.001=*** & ≤0.0001=****, unpaired, two-tailed t-test **b)** Heatmap of gene expression measured by RNA-seq in MP2_ANGPTL4_KD (black/gray), MP2 (pink/light pink), and MP2_ANGPTL4_OE (turquoise/aqua) cell lines both untreated and treated with 1.5nM gemcitabine. Normalized expression of the 42 genes that resulted

from an intersection of ANGPTL4 expression DEGs and gemcitabine treatment response

DEGs are included in the heatmap. **c)** Kaplan-Meier curve for overall survival using

TCGA-PAAD data where patients were divided into quartiles with highest (red) and lowest

(blue) using mean expression of the 42 genes in the intersection (40 genes were available

in TCGA: *KDM7a* and *STN1* were omitted). P-values were derived from a log-rank test.

***ITGB4* and *APOL1* inhibition sensitizes high *ANGPTL4* cells to gemcitabine**

I*n vitro* overexpression of *ANGPTL4* alters cellular drug response and correlates with

patient survival but the pathway(s) by which this resistance is achieved are unknown. We

reasoned that genes co-regulated with ANGPTL4 may function with ANGPTL4 and

contribute to resistance and found those genes by identifying genes that are co-

expressed with *ANGPTL4* in patient tissues. For each of the 1198 DEGs identified when

ANGPTL4 was overexpressed *in vitro*, we calculated rank-based correlation coefficients

and identified co-expressed genes in patient tissues (TCGA-PAAD). For each gene with

a significant correlation with ANGPTL4 in patient tissues, we determined whether

expression of that gene alone was significantly correlated with patient overall survival

(**Supplementary Table S7**). This allowed further filtering of the list based on the

assumption that true resistance-associated genes would impact patient survival. From

this list, ITGB4 and APOL1 are both positively correlated with *ANGPTL4* expression *in*

vitro and in patients, altered with gemcitabine response *in vitro*, and correlated with

patient survival in tissues. While there are several genes of interest, we focused on ITGB4

and APOL1 in this study, in part based on their drugability (pharos.nih.gov) and availability

of efficient siRNA knockdown reagents. Based on the co-expression with ANGPTL4 and

the link to patient survival, we tested whether expression of these genes in combination

with ANGPTL4 was relevant for patient outcomes. Comparing tissues that were among

the top 25% expression of both *ANGPTL4* and *APOL1* (**Fig. 3A**) or ITGB4 (**Fig. 3B**) to

those that were among the bottom 25% expression of both *ANGPTL4* and *APOL1* or

ITGB4 we saw significant differences in overall survival, with overall survival being worse

for TCGA-PAAD patients with high ANGPTL4 combined with high APOL1 (p = 0.014) or

ITGB4 (p = 0.01) expression. Tumors with high levels (top 25%) of all three genes;

APOL1, *ITGB4*, and *ANGPTL4,* also had reduced survival compared with the bottom

25%, but when APOL1 and ITGB4 are combined with ANGPTL4 expression the reduction in

survival did not become more significant (p = 0.015) (**Supplementary Fig. S1D).**

Given the association of ANGPTL4 with resistance and patient outcome, we tested

whether, like ANGPTL4, loss of *ITGB4* or *APOL1* expression increased sensitivity to

gemcitabine *in vitro*. Using siRNAs, we knocked down *APOL1* and *ITGB4* in the

MP2_ANGPTL4_OE cell line. We showed that treatment with the siRNAs reduced mRNA

levels of the target genes (**Supplementary Figure S2A-B**), and we also found that

knockdown of *ITGB4* or *APOL1* reduced expression of *ANGPTL4* significantly compared

to a non-targeting siRNA control with ANGPTL4 overexpression background; expression

was reduced to 2% in MP2, 5% in ITGB4 knockdown, and 25% in APOL1 knockdown

(Fig. 3C, Supplementary Table S1). Given our previous finding that overexpression of

ANGPTL4 leads to resistance, we hypothesized that knockdown of *APOL1* and *ITGB4*

might reverse resistance given the impact on *ANGPTL4* expression. We tested the impact

of *ITGB4* and *APOL1* knockout on gemcitabine sensitivity in the MP2_ANGPTL4_OE line.

After gemcitabine treatment, cells with knockdown of *ITGB4* or *APOL1* in the

ANGPTL4_OE background (MP2_ANGPTL4_OE_ITGB4_KD or

MP2_ANGPTL4_OE_APOL1_KD) showed reversal of drug resistance when compared

to the ANGPTL4 OE cells **(Fig. 3D).** There was no difference in growth rate among the

lines that explained the differential viability observed with gemcitabine treatment

(Supplementary Fig. S2C).

Figure 3

Figure 3: ITGB4 and APOL1 inhibition sensitizes cells overexpressing ANGPTL4 to

gemcitabine.

a-b) Kaplan-Meier plot for overall survival using TCGA-PAAD data. Patients with the top

(red) and bottom (blue) 25% average ANGPTL4 gene expression and APOL1 or ITGB4

gene expression (p=0.01 and p=0.014, respectively, log-rank test). **c)** ANGPTL4

expression measured by qPCR normalized to MP2_ANGPTL4_OE+DsiRNA_NTC

(turquoise) control: MP2 (pink), MP2_ANGPTL4_OE_ITGB4_KD (lavender), and

MP2_ANGPTL4_OE_APOL1_KD   (purple)   compared   to   mean   of

MP2_ANGPTL4_OE+DsiRNA_NTC (MP2_ANGPTL4_OE cells transduced with non-

targeting control siRNA). Expression is normalized to the housekeeping gene ACTB. P-

values ≤ 0.0001=****; unpaired two-tailed parametric t-test **d)** Relative viability is plotted

by normalizing all data to cells not treated with gemcitabine. MP2_ANGPTL4_KD (black),

MP2_NTC   (pink),   MP2_ANGPTL4_OE+DsiRNA_NTC   (turquoise),

MP2_ANGPTL4_OE_ITGB4_KD (lavender), and MP2_ANGPTL4_OE_APOL1_KD

(purple) cells treated with 0, 5nM, or 12.5nM gemcitabine. Sample size ranges from 4-

11 per condition. P-value asterisks: ≤0.001=*** & ≤0.0001=****, unpaired, two-tailed

parametric t-tests.

ANGPTL4 **overexpression increases cell migration**

To further understand the pathways that might be involved in resistance, we performed

pathway enrichment analysis on the 1198 DEG associated with overexpression of

ANGPTL4 (OE v KD). Those included tumor invasion and metastasis, proliferation,

differentiation, inhibition of apoptosis, and cell membrane formation. Genes participating

in invasion and metastasis included RhoEGF which was increased with *ANGPTL4*

expression. RhoEGF is linked to tumor invasion and metastasis **[41]**. Many genes in the

Ras-ARF6 and Ras-MEK2 signaling pathways are also upregulated (e.g. *FGF, APOL1,*

ITGB4, EGFR, MPK, CREB, cFOS, JUN, and *VEGF*) impacting proliferation and

differentiation. In addition, ANTPGL4 overexpression is associated with the upregulation

of anti-apoptotic pathways including JAK-STAT, Bcl_XS, and SOCS pathways triggered

by upregulation of cytokine, cytokine receptors, *CNTF*, and *IL22RA2*. The formation of

cellular structure was also affected by *ANGPTL4* and increase in Claudin via the decrease

in cell permeability (*GEF-H1, Claudin2*), increased actin assembly (*ARP2/3*), and cell

polarity (*AMPK, DLG1*) (**Fig. 4A, Supplementary Table S8).**

Others have shown that on their own *ANGPTL4, APOL1,* and *ITGB4* are each associated

with cell proliferation and contribute to PDAC progression **[17, 42, 43]**. We measured

migration using a wound healing time course assay to determine if the increased

migration associated with *ANGPTL4* overexpression could be reversed with the KD of

ITGB4 or *APOL1* (**Fig. 4B).** We observed that knockdown of either *APOL1* or *ITGB4* in

lines overexpressing *ANGPTL4* reduced the migratory potential of these cells. This was

apparent 48 hours after the scratch was made and was quantified by calculating the time

to close half of the gap (t1/2 gap) for each of the four cell types is plotted. (**Fig. 4C,**

Supplementary Table S1).

Figure 4

Figure 4: ANGPTL4 overexpression correlation with EMT.

a) Pathway analysis of 1198 DEG revealed common cancer pathways. Red=log$_2$FC >0.7 and Blue=log$_2$FC <0.7. **b)** Wound healing assay results at 0 hrs (after initial wound) and 48 hours MP2_NTC (pink), MP2_ANGPTL4_OE+DsiRNA_NTC (turquoise), MP2_ANGPTL4_OE_ITGB4_KD (lavender), and MP2_ANGPTL4_OE_APOL1_KD (purple). Each set was performed in triplicate or quadruplicate. **c)** Time to close half the gap (t1/2 gap) for each of the four cell types is plotted. t1/2: MP2=57.7hrs, MP2_ANGPTL4_OE= 31.6 hrs., MP2_ANGPTL4_OE_ITGB4_KD= 192 hrs., and MP2_ANGPTL4_OE_APOL1_KD=52.96 hrs. P-values ≤0.01=**, unpaired parametric t-

test.

DISCUSSION

Here we have described how overexpression of *ANGPTL4* in pancreatic cancer

contributes to disease progression and resistance. We have shown that overexpression

of this gene and protein leads to cellular resistance to gemcitabine, one of the most

commonly used chemotherapeutics in PDAC. It follows that if overexpression leads to

chemoresistance, increased expression of this gene would also be associated with poor

patient outcomes, and we confirmed that in independent patient cohorts. The goal of our

study is to further understand the role of this gene in chemoresistance in the hopes that

a mechanistic understanding will facilitate the development of new treatment strategies.

Our transcriptomic analysis revealed that overexpression of *ANGPTL4* broadly impacts

transcriptomic profiles in pancreatic cancer cells. This is perhaps not surprising given a

rather large body of literature describing not only ANGPTL4's role in cancer but also

cardiovascular disease risk and metabolism **[9]**. In fact, ANGPTL4's role in cancer can

be difficult to summarize because it has been shown to have both protective and

promoting effects in different cancer types **[13]**. Our initial findings highlighted the

potential importance of this gene and existing literature was not sufficient to understand

how this gene functions in pancreatic cancer. Using our transcriptomic data, we narrowed

our analysis to a list of *ANGPTL4*-impacted genes that are also responsive to gemcitabine

treatment *in vitro*. We identified 42 genes associated with *ANGPTL4* overexpression and

gemcitabine response and determined that the expression of those genes effectively

predicted patient outcomes.

Further narrowing our focus, we identified *APOL1* and *ITGB4* among the genes that are

associated with *ANGPTL4* overexpression, gemcitabine resistance, and patient survival

and explored them further. We found that knockdown of either *APOL1* or *ITGB4* increased

sensitivity to gemcitabine in cells overexpressing *ANGPTL4*. This finding confirms that

inhibition of *APOL1* or *ITGB4* can reverse resistance associated with *ANGPTL4*

expression in pancreatic cancer cells. One unexpected finding was that knockdown of

these genes also reduced expression of *ANGPTL4* itself. Rather than our initial

hypothesis that expression of *ANGPTL4* was correlated with *APOL1* or *ITGB4* because

they were regulated by *ANGPTL4*, this data supports the hypothesis that *ANGPTL4* is

regulated, directly or indirectly, by *APOL1* and *ITGB4* and these genes could be involved

in a feedback loop that explains our results.

Considering *APOL1* first, APOL1 (apolipoprotein L1) is part of the HDL lipid complex

which plays a key role in lipid metabolism **[44]**. APOL1 can activate NOTCH1 signaling

leading to reduced proliferation and increased apoptosis **[42]**. NOTCH1 also upregulates

ABCC1, a transporter that can also lead to multidrug resistance **[45]**. In other contexts,

NOTCH1 also has been shown to stabilize and activate PPARγ, a transcription factor that

promotes transcription of ANGPTL4 **[46]**. These results suggest a possible mechanism

by which knockdown of *APOL1* may reduce *ANGPTL4* expression and further suggests

that inhibition of *APOL1*, especially in patients with high expression of *ANGPTL4* might

reduce cellular capacity for migration and proliferation. Additional studies would need to

be performed to demonstrate whether the link between ANGPTL4 and APOL1 requires

NOTCH signaling.

ITGB4 knockdown also reduces *ANGPTL4* overexpression. ITGB4 (integrin β4, α6β4) is

an integrin protein that plays a role in cell-extracellular matrix adhesion. It is upregulated

in several cancers including pancreatic cancer and has been linked to poor prognosis,

high tumor grade, lymph node metastasis, and drug response **[47, 48, 49]**. Knockdown

of *ITGB4* increases cisplatin sensitivity in lung cancer models **[47]** suggesting that *ITGB4*

is relevant for resistance to multiple chemotherapeutics. ITGB4 may promote receptor

tyrosine kinase (RTK) activation through ERBB2 **[50]**. RTKs generally promote

proliferation, invasion, and sustained cell migration, and ITGB4 has a synergistic effect in

combination with increased RTKs expression **[50, 51]**. In our study, we observed

increased expression of *CSF3, VEGFA, HBEGF,* and *DDR2*, which all play a role in the

epithelial-to-mesenchymal transition in cells overexpressing *ANGPTL4*. An alternate

hypothesis is that *ITGB4* promotes cell migration and invasion through regulation of the

MEK1-ERK1/2 signaling cascade **[49]**. Reduction of *ITGB4* in the context of

overexpressed *ANGPTL4* could also affect this pathway, explaining the reduced

migratory potential of *ITGB4* KD cells in the cells overexpressing *ANGPTL4*. Additional

transcriptomic and biochemical studies are needed to determine which of these pathways

are most relevant for the chemoresistance phenotypes.

CONCLUSIONS

Our study provides evidence that *ANGPTL4* overexpression in pancreatic cancer impacts several key signaling and metabolic pathways associated with patient survival, in part through controlling cellular resistance to chemotherapy and cellular migration. *ITGB4* and *APOL1* knockdown are both able to reverse drug resistance and migration increases associated with overexpression of *ANGPTL4* but appear to do so through different mechanisms. These findings emphasize the complexity of developing treatments to target resistance and metastasis; a single approach may not be sufficient. Nonetheless, both *APOL1* and *ITGB4* represent potential novel targets and our data support the further exploration of these genes and the pathways in which they function.

ABBREVIATIONS

AKT: AKT Serine/Threonine Kinase 1

AMPK: Protein Kinase AMP-Activated Catalytic Subunit Alpha 1

ANGPTL4: Angiopoietin Like 4

ARP2/3: Actin Related Protein 2/3

APOL1: Apolipoprotein L1

Bcl_XS: BCL2 Apoptosis Regulator

cFOS: Fos Proto-Oncogene, AP-1 Transcription Factor Subunit

CNTF: Ciliary Neurotrophic Factor

CO$_2$: carbon dioxide

CREB: CAMP Responsive Element Binding Protein

CRISPRa: dCas9 activation system

CSF3: colony-stimulating factor 3

CUL7: Cullin 7

DAVID: Database for Annotation, Visualization, and Integrated Discovery

DEG: differentially expressed genes

DDR2: Discoidin Domain Receptor Tyrosine Kinase 2

DLG1: Discs Large MAGUK Scaffold Protein 1

DsiRNA: Dicer-substrate small interfering RNA

EMT: epithelial to mesenchymal transition

EGFR: Epidermal Growth Factor Receptor

ERK: Mitogen-Activated Protein Kinase 1

ERK1/2: Mitogen-Activated Protein Kinase 1/2

FGF: Fibroblast Growth Factor 1

FGF2: Fibroblast Growth Factor Receptor 2

FOS: Fos Proto-Oncogene, AP-1 Transcription Factor Subunit

GEO: Gene Expression Omnibus

GEPIA: Gene Expression Profiling Interactive Analysis

GEF-H1: Rho/Rac Guanine Nucleotide Exchange Factor 2

HBEGF: Heparin Binding EGF-Like Growth Factor

INPP4B: Inositol Polyphosphate-4-Phosphatase Type II B

ITGB4: Integrin Subunit Beta 4

JAK:Janus Kinase 1

JNK: Mitogen-Activated Protein Kinase 8

JUN: Jun Proto-Oncogene, AP-1 Transcription Factor Subunit

KD: knockdown

KEGG: Kyoto Encyclopedia of Genes and Genomes

KDM7a: Lysine Demethylase 7A

KLF4: KLF Transcription Factor 4

KRAS: KRAS Proto-Oncogene, GTPase

IL22RA2: Interleukin 22 Receptor Subunit Alpha 2

MEK1: Mitogen-Activated Protein Kinase

MP2: MIA PaCa-2 cells

MPK: Mitogen-Activated Protein Kinase

n: number

ng: nanograms

nM: nanomolar

NOTCH: Notch Receptor

OE: overexpression

OS: overall survival

*p:*p-value

PAAD: pancreatic ductal adenocarcinoma

padj: adjusted p-value

PARP: Poly(ADP-Ribose) Polymerase

PDAC: pancreatic ductal adenocarcinoma

PI3K: Phosphatidylinositol-4,5-Bisphosphate 3-Kinase Catalytic Subunit Delta

PPP1R15A: Protein Phosphatase 1 Regulatory Subunit 15A

qPCR: quantitative polymerase chain reaction

RFS: recurrence-free survival

SEER: Surveillance, Epidemiology, and End Results

SH3RF2: SH3 Domain Containing Ring Finger 2

siRNA: silencing RNA

SOCS: Suppressor of Cytokine Signaling

SOX9: SRY-Box Transcription Factor 9

STAT: Signal Transducer and Activator of Transcription 3

STAT3: Signal Transducer and Activator of Transcription 3

STN1: STN1 subunit of CTS complex

TCGA: The Cancer Genome Atlas

TGFβ: TGFB Induced Factor Homeobox 1

TGFBR3: Transforming Growth Factor Beta Receptor 3

uL: microliter

v: versus

VEGF: Vascular Endothelial Growth Factor A

VEGFA: Vascular Endothelial Growth Factor A

WWTR1: WW Domain Containing Transcription Regulator 1

DECLARATIONS

Consent for Publication

NA

Data Availability

The RNA-seq data generated in this study are available in Gene Expression Omnibus

(GEO) at GSE207359. Clinical data and RNA-sequencing data for TCGA-PAAD

(PDAC) samples were retrieved on 04/01/2020 using the GDC Data Portal

(https://portal.gdc.cancer.gov/projects/TCGA-PAAD). Our analyses included 178 samples

in this cohort that had matched clinical and RNA sequencing data.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by NIH 1R43CA232844-01A1 and the State of Alabama Cancer

Research Fund.

Authors' contributions

ERG, CAW, and SJC designed the experiments. ERG, CAW, and MJ collected data. ERG

and CAW analyzed the data. ERG and SJC wrote the first draft. All authors contributed

to the writing of the paper and read and approved the final manuscript.

Acknowledgements

This work was funded by NIH 1R43CA232844-01A1 and the State of Alabama Cancer

Research Fund. SJC is supported by UL1TR003096. We would like to thank Anuj Singhal

and his team at CFD Research as well as our HudsonAlpha colleagues, especially those

in the Myers lab for their feedback, support, and discussions of this work. Figure 4A

created with BioRender.com agreement number RD24SQPZ3K.

Ethics Declaration

All methods were carried out in accordance with relevant guidelines and regulations.

REFERENCES

1. U.S. Mortality Data 1969-2019 when Using SEER*Stat: Surveillance,

Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov)

SEER*Stat Database: Mortality - All COD, Aggregated With State, Total U.S. (1969-

2019) [Internet]. 2021. Available from: http://www.cdc.gov/nchs

2. Sturm N, Ettrich TJ, Perkhofer L. The Impact of Biomarkers in Pancreatic Ductal

Adenocarcinoma on Diagnosis, Surveillance and Therapy. *Cancers* 2022;14(1)217.

http://dx.doi.org/10.3390/cancers14010217

3. Conroy T, Desseigne F, Ychou M, Bouché O, Guimbaud R, Bécouarn Y, Adenis A,

Raoul JL, Gourgou-Bourgade S, de la Fouchardière C, Bennouna J, Bachet JB,

Khemissa-Akouz F, Péré-Vergé D, Delbaldo C, Assenat E, Chauffert B, Michel P,

Montoto-Grillot C, Ducreux M; Groupe Tumeurs Digestives of Unicancer;

PRODIGE Intergroup. FOLFIRINOX versus gemcitabine for metastatic pancreatic

cancer. *N Engl J Med*. 2011;364(19):1817-25. doi: 10.1056/NEJMoa1011923.

PMID: 21561347.

4. American Cancer Society. Cancer Facts & Figures 2022. Atlanta; 2022.

5. Javle M, Shacham-Shmueli E, Xiao L, Varadhachary G, Halpern N, Fogelman D,

et al. Olaparib Monotherapy for Previously Treated Pancreatic Cancer With DNA

Damage Repair Genetic Alterations Other Than Germline BRCA Variants: Findings

From 2 Phase 2 Nonrandomized Clinical Trials. *JAMA Oncology* 2021;7:693–9.

6. Moore MJ, Goldstein D, Hamm J, Figer A, Hecht JR, Gallinger S, et al. Erlotinib

plus gemcitabine compared with gemcitabine alone in patients with advanced

pancreatic cancer: A phase III trial of the National Cancer Institute of Canada

Clinical Trials Group. *J Clin Oncol. American Society of Clinical Oncology*

2007;25:1960–6.

7. Pishvaian MJ, Bender RJ, Halverson D, Rahib L, Hendifar AE, Mikhail S, et al.

Molecular profiling of patients with pancreatic cancer: Initial results from the know

your tumor initiative. Clin Cancer Res. *American Association for Cancer Research*

Inc. 2018;24:5018–27.

8. Kirby MK, Ramaker RC, Gertz J, Davis NS, Johnston BE, Oliver PG, et al. RNA

sequencing of pancreatic adenocarcinoma tumors yields novel expression patterns

associated with long-term survival and reveals a role for ANGPTL4. *Mol Oncol.*

2016;10:1169–82.

9. Fernández-Hernando C, Suárez Y. ANGPTL4: A multifunctional protein involved in

metabolism and vascular homeostasis. *Current Opinion in Hematology* 2020. page

206–13. Available from: http://dx.doi.org/10.1097/MOH.0000000000000580

10. La Paglia L, Listì A, Caruso S, Amodeo V, Passiglia F, Bazan V, et al. Potential

Role of ANGPTL4 in the Cross Talk between Metabolism and Cancer through

PPAR Signaling Pathway. *PPAR Research* 2017. http://dx.doi.org/10.1155/2017/8187235

11. Lei X, Shi F, Basu D, Huq A, Routhier S, Day R, et al. Proteolytic processing of angiopoietin-like protein 4 by proprotein convertases modulates its inhibitory effects on lipoprotein lipase activity. *J Biol Chem. American Society for Biochemistry and Molecular Biology Inc.* 2011;286:15747–56.

12. Kim SH, Park YY, Kim SW, Lee JS, Wang D, DuBois RN. ANGPTL4 induction by prostaglandin E 2 under hypoxic conditions promotes colorectal cancer progression. *Cancer Res.* 2011;71:7010–20.

13. Tan MJ, Teo Z, Sng MK, Zhu P, Tan NS. Emerging roles of angiopoietin-like 4 in human cancer. *Molecular Cancer Research* 2012;10(6):677–88. http://dx.doi.org/10.1158/1541-7786.MCR-11-0519

14. Zhu P, Tan MJ, Huang RL, Tan CK, Chong HC, Pal M, et al. Angiopoietin-like 4 Protein Elevates the Prosurvival Intracellular O2-:H2O2 Ratio and Confers Anoikis Resistance to Tumors. Cancer Cell. *Cell Press* 2011;19:401–15.

15. Teo Z, Sng MK, Chan JSK, Lim MMK, Li Y, Li L, et al. Elevation of adenylate energy charge by angiopoietin-like 4 enhances epithelial-mesenchymal transition by inducing 14-3-3γ expression. *Oncogene* 2017;36:6408–19.

16. Zhang K, Zhai Z, Yu S, Tao Y. DNA methylation mediated down-regulation of ANGPTL4 promotes colorectal cancer metastasis by activating the ERK pathway. *J Cancer* 2021;12:5473–85.

17. Zhao J, Liu J, Wu N, Zhang H, Zhang S, Li L, et al. ANGPTL4 overexpression is

associated with progression and poor prognosis in breast cancer. *Oncol Lett*.

2020;20:2499–505.

18. Cai YC, Yang H, Wang KF, Chen TH, Jiang WQ, Shi YX. ANGPTL4 overexpression

inhibits tumor cell adhesion and migration and predicts favorable prognosis of triple-

negative breast cancer. *BMC Cancer* 2020;20. http://dx.doi.org/10.1186/s12885-

020-07343-w

19. Yan HH, Jung KH, Lee JE, Son MK, Fang Z, Park JH, et al. ANGPTL4 accelerates

KRASG12D-Induced acinar to ductal metaplasia and pancreatic carcinogenesis.

Cancer Lett. 2021;519:185–98.

20. Gupta N, Park JE, Tse W, Keem Low J, Kon OL, Mccarthy N, et al. ERO1α

promotes hypoxic tumor progression and is associated with poor prognosis in

pancreatic cancer. *Oncotarget* 2019;10(5):5870-5982.

https://www.oncotarget.com/article/27235/text/

21. Joung, J., Konermann, S., Gootenberg, J. S., Abudayyeh, O. O., Platt, R. J.,

Brigham, M. D., Sanjana, N. E., & Zhang, F. Genome-scale CRISPR-Cas9

knockout and transcriptional activation screening. *Nature Protocols*; 2017; *12*(4),

828–863. https://doi.org/10.1038/nprot.2017.016

22. Larson B, Held P, Banks P. Optimization of Perkin Elmer's AlphaScreen® cAMP

Assay on the Synergy[TM] H4 Hybrid Multi-Mode Microplate Reader [Internet]. 2010.

http://www.biotek.com

23. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol*. 2014;15.

http://dx.doi.org/10.1186/s13059-014-0550-8

24. Ge SX, Son EW, Yao R. iDEP: An integrated web application for differential

expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics* 2018;19.

http://dx.doi.org/10.1186/s12859-018-2486-6

25. Baecker V. ImageJ Macro Tool Sets for Biological Image Analysis. Conference

26. Kanehisa M, Sato Y. KEGG Mapper for inferring cellular functions from protein

sequences. *Protein Sci.* 2020;29:28–35.

27. Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, Thomas PD. PANTHER

version 14: more genomes, a new PANTHER GO-slim and improvements in

enrichment analysis tools. *Nucleic Acids Res.* 2019;47(D1):D419-D426.

28. Therneau T. A Package for Survival Analysis in R. R package version 3.4-0

[Internet]. 2022. https://CRAN.R-project.org/package=survival

29. Kassambara, A., Kosinski, M., & Biecek, F. S.Package "survminer" Type Package

Title Drawing Survival Curves using "ggplot2;" 2022

30. Wickham H. Ggplot2 : elegant graphics for data analysis. Springer; 2009.

31. Love, M. I., Huber, W., & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology* 2014; *15*(12).

https://doi.org/10.1186/s13059-014-0550-8

32. Kolde R. pheatmap: Pretty Heatmaps. R package version 1.0.12 [Internet]. 2019.

Available from: https://CRAN.R-project.org/package=pheatmap

33. Frank M, Harrell E. Hmisc: A package of miscellaneous R functions [Internet]. 2020.

Available from: http://biostat.mc.vanderbilt.edu/Hmisc

34. Urbanek S, Johnson K. tiff: Read and Write TIFF Images. R package version 0.1-

11. 2020; Available from: https://CRAN.R-project.org/package=tiff

35. Jia Y, Xie J. Promising molecular mechanisms responsible for gemcitabine

resistance in cancer. *Genes and Diseases* 2015;2(4):299-306.

http://dx.doi.org/10.1016/j.gendis.2015.07.003

36. Wei X, Zhou X, Zhao Y, He Y, Weng Z, Xu C. A 14-gene gemcitabine resistance

gene signature is significantly associated with the prognosis of pancreatic cancer

patients. *Sci Rep* 2021;11(6087). http://dx.doi.org/10.1038/s41598-021-85680-x

37. Akada M, Crnogorac-Jurcevic T, Lattimore S, Mahon P, Lopes R, Sunamura M, et

al. Intrinsic Chemoresistance to Gemcitabine Is Associated with Decreased

Expression of BNIP3 in Pancreatic Cancer. *Clin Cancer Res.* 2005;11(8):3094-101.

https://aacrjournals.org/clincancerres/article/11/8/3094/188241/Intrinsic-

Chemoresistance-to-Gemcitabine-Is

38. Nakano Y, Tanno S, Koizumi K, Nishikawa T, Nakamura K, Minoguchi M, et al.

Gemcitabine chemoresistance and molecular markers associated with gemcitabine

transport and metabolism in human pancreatic cancer cells. *Br J Cancer*

2007;96:457–63.

39. Vasaikar SV, Deshmukh AP, den Hollander P, Addanki S, Kuburich NA, Kudaravalli

S, Joseph R, Chang JT, Soundararajan R, Mani SA. EMTome: a resource for pan-

cancer analysis of epithelial-mesenchymal transition genes and signatures. *Br J*

Cancer. 2021;124(1):259-269. doi: 10.1038/s41416-020-01178-9. Epub 2020 Dec 10. PMID: 33299129; PMCID: PMC7782839.

40. Huang L, Jansen L, Balavarca Y, Babaei M, van der Geest L, Lemmens V, Van Eycken L, De Schutter H, Johannesen TB, Primic-Žakelj M, Zadnik V, Besselink MG, Schrotz-King P, Brenner H. Stratified survival of resected and overall pancreatic cancer patients in Europe and the USA in the early twenty-first century: a large, international population-based study. BMC Med. 2018 Aug 21;16(1):125. doi: 10.1186/s12916-018-1120-9. PMID: 30126408; PMCID: PMC6102804.

41. Ghanem NZ, Matter ML, Ramos JW. Regulation of Leukaemia Associated Rho GEF (LARG/ARHGEF12). *Small GTPases* 2022;13(1)196-204; http://dx.doi.org/10.1080/21541248.2021.1951590

42. Lin, J., Xu, Z., Xie, J., Deng, X., Jiang, L., Chen, H., Peng, C., Li, H., Zhang, J., & Shen, B. Oncogene APOL1 promotes proliferation and inhibits apoptosis via activating NOTCH1 signaling pathway in pancreatic cancer. *Cell Death and Disease*; 2021; 12(8). https://doi.org/10.1038/s41419-021-03985-1

43. Cruz-Monserrate Z, Qiu S, Evers BM, O'Connor KL. Upregulation and redistribution of integrin α6β4 expression occurs at an early stage in pancreatic adenocarcinoma progression. *Mod Pathol*. 2007;20:656–67.

44. Duchateau PN, Pullinger CR, Orellana RE, Kunitake ST, Naya-Vigne J, O'Connor PM, et al. Apolipoprotein L, a new human high density lipoprotein apolipoprotein expressed by the pancreas. Identification, cloning, characterization, and plasma distribution of apolipoprotein L. *J Biol Chem.* 1997;272:25576–82.

45. Cho S, Lu M, He X, Ee PLR, Bhat U, Schneider E, et al. Notch1 regulates the

expression of the multidrug resistance gene ABCC1/MRP1 in cultured cancer cells.

Proc Natl Acad Sci 2011;108:20778–83.

46. Sangphech N, Keawvilai P, Palaga T. Notch signaling increases PPARγ protein

stability and enhances lipid uptake through AKT in IL-4-stimulated THP-1 and

primary human macrophages. *FEBS Open Bio.* 2020;10:1082–95.

47. Mohanty A, Nam A, Pozhitkov A, Yang L, Srivastava S, Nathan A, et al. A Non-

genetic Mechanism Involving the Integrin β4/Paxillin Axis Contributes to

Chemoresistance in Lung Cancer. *iScience* 2020;23.

http://dx.doi.org/10.1016/j.isci.2020.101496

48. Yoshioka T, Otero J, Chen Y, Kim YM, Koutcher JA, Satagopan J, et al. β4 Integrin

signaling induces expansion of prostate tumor progenitors. *J Clin Invest*.

2013;123:682–99.

49. Masugi Y, Yamazaki K, Emoto K, Effendi K, Tsujikawa H, Kitago M, Itano O,

Kitagawa Y, Sakamoto M. Upregulation of integrin β4 promotes epithelial-

mesenchymal transition and is a novel prognostic marker in pancreatic ductal

adenocarcinoma. *Lab Invest.*; 2015;95(3):308-19. doi: 10.1038/labinvest.2014.166.

PMID: 25599535.

50. Giancotti FG, Ruoslahti E. Integrin signaling. Science. 2017;285:1028–32.

51. Meng X, Liu P, Wu Y, Liu X, Huang Y, Yu B, et al. Integrin beta 4 (ITGB4) and its

tyrosine-1510 phosphorylation promote pancreatic tumorigenesis and regulate the

MEK1-ERK1/2 signaling pathway. *Bosn J Basic Med Sci*. 2020;20:106–16.

52. ImageJ User and Developer Conference 2012 At: Luxembourg. 2012.

ADDITIONAL FILES

Additional file 1: Figure S1.pdf

a) Kaplan-Meier curve of Recurrence/Disease Free Survival (RFS) using GEPIA PPAD

dataset. Log rank p=0.047 **b)** Protein abundance of ANGPTL4 in cell lysate and

supernatant as measured by AlphaLISA assay for normalization by cell input.

MP2_ANGPTL4_OE vs MP2_ANPGTL4_KD p=0.0075 and 0.0018 (**) for lysate and

supernatant. MP2_ANGPTL4_OE vs MP2 p=0.0055 (**) for lysate. **c)** Heatmap of

RNA-seq data for 1198 DEG from MP2_ANGPTL4_OE vs MP2_ANGPTL4_KD with

thresholds: baseMean>10 padj<0.05, and $\log_2$ fold change +/-0.7.

Additional file 2: Figure S2.pdf

a) ITGB4 expression measured by qPCR and normalized by the housekeeping gene

ACTB. All data points are plotted relative to the ANGPTL4_OE line. **b)** *APOL1* expression

measured by qPCR and normalized by the housekeeping gene *ACTB*. All data points are

plotted relative to the ANGPTL4_OE line. **c)** Cell viability over time normalized to control

MP2 cells (lt. pink) at time 0 for MP2_NTC (pink), MP2_ANGPTL4_OE+DsiRNA_NTC

(turquoise),   MP2_ANGPTL4_OE_ITGB4_KD   (lavender),   and

MP2_ANGPTL4_OE_APOL1_KD (purple).

Additional file 3: Table S1.txt

P-values associated with qPCR and viability data

Additional file 4: Table S2.txt

DEG from MP2_ANGPTL4_OE vs MP2_ANGPTL4_KD analysis with or without

gemcitabine treatment

Additional file 5: Table S3.txt

114 Gemcitabine Resistance Genes

Additional file 6: Table S4.txt

AMIGO2 Panther GO biological process complete analysis for DEG retrieved from with

or without gemcitabine analysis

Additional file 7: Table S5.txt

AMIGO2 Panther GO biological process EMT results for with or without gemcitabine

analysis

Additional file 8: Table S6.txt

List of 42 DEG genes for both ANGPTL4 expression and gemcitabine resistance

Additional file 9: Table S7.txt

Correlation analysis of TCGA-PAAD data using DEG gene list from MP2_ANGPTL4_OE

vs MP2_ANGPTL4_KD analysis with or without gemcitabine treatment

Additional file 10: Table S8.txt

Pathway analysis results from DEG from MP2_ANGPTL4_OE vs MP2_ANGPTL4_KD

analysis

FIGURE LEGENDS

Figure 1: Expression of *ANGPTL4* is associated with patient outcomes and

modifies expression of resistance-associated genes.

a) A Kaplan Meier curve for overall survival using TCGA-PAAD data (n=90). Top (red)

and bottom (blue) 25% average *ANGPTL4* gene expression are significantly different.

(p=0.043, log-rank test) **b)** Relative expression of ANGPTL4 transcript in

MP2_ANGPTL4_KD (black circles), MP2 (pink squares), and MP2_ANGPTL4_OE

(turquoise triangles) as measured by qPCR. P-values from unpaired, two-tailed

parametric t-tests performed with 95% confidence intervals are displayed. Values are

normalized with *ACTB* as the housekeeper. (P-values ≤ 0.0001=****) **c)** Heatmap

displaying RNA-seq results from MP2_ANGPTL4_KD (black), MP2_NTC (pink), and

MP2_ANGPTL4_OE (turquoise) cell lines. Displayed are 15 genes associated with both

ANGPTL4 overexpression and gemcitabine resistance. Normalized expression for each

of the 15 genes is plotted in the heatmap. Red (*) indicate ITGB4 used for later

experiments.

Figure 2: Gemcitabine resistance signature in PDAC cells with high ANGPTL4

a) Cell viability was measured for MP2_ANGPTL4_KD (black), MP2_NTC (pink), and

MP2_ANGPTL4_OE (turquoise) cells treated with 0, 5nM, or 12.5nM gemcitabine. A

relative viability reflects normalization to cells not treated with gemcitabine. P-value

asterisks: ≤0.01=**, ≤0.001=*** & ≤0.0001=****, unpaired, two tailed t-test **b)** Heatmap of

gene expression measured by RNA-seq in MP2_ANGPTL4_KD (black/gray), MP2

(pink/light pink), and MP2_ANGPTL4_OE (turquoise/aqua) cell lines both untreated and

treated with 1.5nM gemcitabine. Normalized expression of the 42 genes that resulted

from an intersection of ANGPTL4 expression DEGs and gemcitabine treatment response

DEGs are included in the heatmap. **c)** Kaplan-Meier curve for overall survival using

TCGA-PAAD data (n=90) where patients were divided into quartiles with highest (red)

and lowest (blue) using mean expression of the 42 genes in the intersection (40 genes

were available in TCGA: *KDM7a* and *STN1* were omitted). P-values were derived from a

log-rank test.

Figure 3: ITGB4 and APOL1 inhibition sensitizes cells overexpressing ANGPTL4 to

gemcitabine.

a-b) Kaplan-Meier plot for overall survival using TCGA-PAAD data. Patients (n=90) with

top (red) and bottom (blue) 25% average *ANGPTL4* gene expression and *APOL1* or

ITGB4 gene expression (p=0.01 and p=0.014, respectively, log-rank test). **c)** ANGPTL4

expression measured by qPCR normalized to MP2_ANGPTL4_OE+DsiRNA_NTC

(turquoise) control: MP2 (pink), MP2_ANGPTL4_OE_ITGB4_KD (lavender), and

MP2_ANGPTL4_OE_APOL1_KD (purple) compared to mean of

MP2_ANGPTL4_OE+DsiRNA_NTC (MP2_ANGPTL4_OE cells transduced with non-

targeting control siRNA). Expression is normalized to the housekeeping gene *ACTB*. P-

values ≤ 0.0001=****; unpaired two-tailed parametric t-test **d)** Relative viability is plotted

by normalizing all data to cells not treated with gemcitabine. MP2_ANGPTL4_KD (black),

MP2_NTC (pink), MP2_ANGPTL4_OE+DsiRNA_NTC (turquoise),

MP2_ANGPTL4_OE_ITGB4_KD (lavender), and MP2_ANGPTL4_OE_APOL1_KD

(purple) cells treated with 0, 5nM, or 12.5nM gemcitabine. Sample size ranges from 4-

11 per condition. P-value asterisks: ≤0.001=*** & ≤0.0001=****, unpaired, two-tailed

parametric t-tests.

Figure 4: ANGPTL4 overexpression correlation with EMT.

a) Pathway analysis of 1198 DEG revealed common cancer pathways. Red=$\log_2$FC >0.7

and Blue=$\log_2$FC <0.7. **b)** Wound healing assay results at 0 hrs (after initial wound) and

48 hours MP2_NTC (pink), MP2_ANGPTL4_OE+DsiRNA_NTC (turquoise),

MP2_ANGPTL4_OE_ITGB4_KD (lavender), and MP2_ANGPTL4_OE_APOL1_KD

(purple). Each set was performed in triplicate or quadruplicate. **c)** Time to close half the

half the gap (t1/2 gap) for each of the four cell types is plotted. t1/2: MP2=57.7hrs,

MP2_ANGPTL4_OE= 31.6 hrs., MP2_ANGPTL4_OE_ITGB4_KD= 192 hrs., and

MP2_ANGPTL4_OE_APOL1_KD=52.96 hrs. P-values ≤0.01=**, unpaired parametric t-

test

Figure 4

Figure 3

Figure 1

Figure 2