Functional genomic mechanisms of opioid action and opioid use disorder: a systematic

review of animal models and human studies

Camille Falconnier[1], Alba Caparros-Roissard[1], Charles Decraene[1,2], Pierre-Eric Lutz[1,3]

[1] Centre National de la Recherche Scientifique, Université de Strasbourg, Institut des

Neurosciences Cellulaires et Intégratives UPR3212, 67000 Strasbourg, France

[2] Centre National de la Recherche Scientifique, Université de Strasbourg, Laboratoire de

Neurosciences Cognitives et Adaptatives, 67000 Strasbourg, France

[3] Douglas Mental Health University Institute, Montreal, QC, Canada

Corresponding author:

Pierre-Eric Lutz, INCI UPR 3212, 8 allée du général Rouvillois, 67000 Strasbourg, France

Email: pierreeric.lutz@gmail.com

Word count: 7495

6 Figures, 4 Supplementary Figures, 6 Supplementary Tables, 1 Supplementary Material, 1

Supplementary File

Abstract

In the past two decades, over-prescription of opioids for pain management has driven a steep increase in opioid use disorder (OUD) and death by overdose, exerting a dramatic toll on western countries. OUD is a chronic relapsing disease associated with a lifetime struggle to control drug consumption, suggesting that opioids trigger long-lasting brain adaptations, notably through functional genomic and epigenomic mechanisms. Current understanding of these processes, however, remain scarce, and have not been previously reviewed systematically. To do so, the goal of the present work was to synthesize current knowledge on genome-wide transcriptomic and epigenetic mechanisms of opioid action, in primate and rodent species. Using a prospectively registered methodology, comprehensive literature searches were completed in PubMed, Embase, and Web of Science. Of the 2709 articles identified, 73 met our inclusion criteria and were considered for qualitative analysis. Focusing on the 5 most studied nervous system structures (nucleus accumbens, frontal cortex, whole striatum, dorsal striatum, spinal cord; 44 articles), we also conducted a quantitative analysis of differentially expressed genes, in an effort to identify a putative core transcriptional signature of opioids. Only one gene, Cdkn1a, was consistently identified in eleven studies, and globally, our results unveil surprisingly low consistency across published work, even when considering most recent single-cell approaches. Analysis of putative sources of variability detected significant contributions from species, brain structure, duration of opioid exposure, strain, time-point of analysis, and batch effects, but not type of opioid. To go beyond those limitations, we leveraged threshold-free methods to illustrate how genome-wide comparisons may generate new findings and hypotheses. Finally, we discuss current methodological development in the field, and their implication for future research and, ultimately, better care.

Keywords

Opioid, substance use disorder, genomics, epigenomics, next generation sequencing

Introduction

Opioids have been used for millennia for their analgesic and euphoric properties[1]. While they remain reference pain treatments, their chronic use also associates with tolerance, physical dependence and, in some individuals, the emergence of an opioid use disorder (OUD). OUD is defined as a problematic use leading to significant impairment. This severe and chronic disorder associates with a life expectancy decreased by more than 10 years due to somatic and psychiatric comorbidities. Over the last 2 decades, western countries have faced an increase in death by opioid overdose, due to more frequent prescription for pain management, increasing use of illegal compounds, or misuse of substitution therapies. OUD imposes a major socio-economic burden, with an estimated annual cost in the trillion dollar range in the US[2], and increasing use and harm in Europe[3]. This results, overall, in a worsening public health crisis.

OUD results from interacting biological, psychological and socio-economic factors[4]. At a biological level, it originates from pharmacological effects of opioid drugs, which trigger chemical and molecular brain adaptations, under modulation by genetic vulnerability and epigenetic regulation. In turn, these effects mediate behavioral and cognitive dysfunctions, including the inability to control drug use despite harmful consequences, or persistent and intense desire for the drug, even after years of abstinence.

Since the 1990s, high-throughput approaches have been harnessed to characterize the molecular underpinnings of opioid effects in brain tissue, throughout the full genome[5], across various experimental models and species. In this context, the goal of the present systematic review was to synthesize current knowledge on functional genomic mechanisms of opioid action and OUD, defined as changes in gene expression or epigenetic regulation. To do so, we used a preregistered methodology, and performed an unbiased survey of bibliographic repositories, focusing on high-throughput studies, most notably microarrays and next-generation sequencing. While we acknowledge the contribution of candidate gene studies (already reviewed in the past[6, 7]), this focus on genome-wide analyses reflects the

conviction that understanding heterogeneous phenotypes such as OUD requires analyzing

the full genome.

Based on studies identified through this screening, a qualitative synthesis was first

performed to summarize experimental designs and findings. Second, a focused quantitative

analysis of bulk tissue transcriptomic studies was conducted, with the goal of defining a core

signature of opioids. To our knowledge, while numerous reviews are regularly published in the

opioid field[8–12], such a systematic synthesis across brain structures and technologies had

never been performed. Our results indicate that, although more than 40 genome-wide studies

have been published, available evidence for convergent findings is surprisingly limited. Third,

we reviewed most recent work that leveraged cell-type specific and single-cell approaches, or

that interrogated epigenetic regulatory mechanisms contributing to opioid plasticity. Finally,

we discuss current challenges, as well as avenues and recommendations for future work.

Methods

Full methodology and code are available as Supplementary Material and File.

Results

The identification and selection of eligible studies is presented in Fig.1. For qualitative

synthesis, 73 papers were selected, covering both transcriptomic and epigenomic

approaches. Among these, the 44 articles that performed transcriptomic analyses in the 5

most frequently investigated regions were selected for more detailed quantitative synthesis:

of these, 34 investigated a single brain region, 9 covered 2 regions, and only 1 characterized

5 regions[13]. This resulted in 52 differential expression analyses: 18, 11, 11, 6 and 6 in the

nucleus accumbens (NAc), frontal cortex, whole striatum, dorsal striatum and spinal cord,

respectively.

1. Qualitative analysis (n=73 studies)

Species and sex ratio. Most studies were conducted in rodents (Fig.2A): 30 (41%) in rats, 29 (40%) in mice, and 1 in both species. Only a small number investigated humans (n=11, 15%) and macaques (n=2, 3%), reflecting well-known practical limitations. As consequence, current knowledge on species-specific opioid effects is limited.

Regarding sex, epidemiological surveys indicate that OUD prevalence is higher in males than in females[3, 14] (although smaller differences have been reported with DSM-5 criteria[15]), with a number of sex-specific risk factors (e.g., ethnicity, income). This clinical situation has long been modeled in rodents, as sex differences in behavioral effects of opioids[16] have been described for analgesia[17], tolerance[18], withdrawal symptoms[19], reward[20] or motivation[21]. In comparison, molecular mechanisms underlying these differences are poorly characterized. The majority of OUD studies reviewed here investigated males only (n=51, 70%; Fig.2B-D), with few conducted in females only (n=5, 7%), and 15 (20%) that included both sexes (7 in rodents, 8 in humans). These numbers are consistent with the current landscape of neuroscience research, in which the ratio of articles reporting on males only, as opposed to females only, is around 5 to 1[22]. We nevertheless note an encouraging trend, as the majority of studies published since 2021 (8/12) investigated both sexes. Several factors likely underlie this evolution: policies from funding agencies (such as the NIH; https://grants.nih.gov/grants/guide/notice-files/not-od-15-102.html); increased recognition that females do not display more experimental variability than males[23]; and evidence that pathophysiological mechanisms may differ across sexes, as shown for depression[24], schizophrenia[25] or alcohol use disorder[26].

Specifically in rodents (Fig.2C), 80% of studies investigated males only (n=48). Seven articles investigated both sexes (n=7, 12%), using surprisingly low sample sizes (n=1-4 in male/female, and opioid-treated/control, groups; Fig.2D). Among these, 2 studies treated sex as a covariate[27, 28], and 3 pooled both sexes for the analysis of opioid effects[29–31], therefore providing no direct description of male/female differences. This is concerning, considering that the 2 remaining studies found little overlap, from 5 to 35% (see[32] and[33]), among morphine-induced changes occurring in each sex. Therefore, available evidence

points toward substantial sex-specificity in opioid-induced transcriptomic plasticity, which will have to be confirmed in future work. In human, even less is known (Fig.2E). Among 11 articles, 8 investigated males and females, while 3 investigated males only. However, these 8 studies showed a strong bias towards examination of men (Fig.2F), with only one using similar sample sizes in both sexes[34]. In the latter, sex was treated as a covariate, with no specific description of its impact. Altogether, this significant gap in the literature calls for additional work, with the hope that better understanding of sex-specific pathophysiological routes may unravel distinct and more efficient therapeutic strategies.

Route and duration of opioid administration. Among studies included in qualitative analysis (n=73), the majority used chronic administration (n=59/73, 81%; see Fig.3A), mostly via self-administration (SA, n=22/59, 37%, including 11 human studies) and intraperitoneal injection (ip; n=19, 32%), followed by subcutaneous injection (n=8, 14%). Other procedures were less frequently used: continuous administration (with pellets or mini-pumps; n=5, 8%), intrathecal administration (n=3, 5%), or gavage (n=2, 3%).

There is limited information regarding differences among administration procedures. Interestingly, Lefevre et al[35] recently compared 3 conditions: continuous and systemic morphine infusion using osmotic mini-pumps, with ("Interrupted") or without ("Continuous") the induction, twice daily, of a precipitated withdrawal episode (using naloxone, a non-specific opioid antagonist), compared to Controls (saline mini-pumps). Results indicated 687 and 407 differentially expressed genes (DEG) when comparing Interrupted and Control groups in the NAc and dorsal striatum, respectively, with only 1 DEG between Continuous and Control groups. Therefore, rather than opioid exposure per se, it is the repetition of intermittent withdrawal episodes that may drive the intensity of transcriptomic adaptations affecting the mesolimbic system (whether precipitated, as in this study, or spontaneous, in-between bolus injections). This is consistent with the clinical notion that the alternation of intoxication and withdrawal phases is critical for the development of OUD.

Opioid effects have also been investigated following acute administration (10/73

studies, 14%), corresponding mostly to ip injection (9 articles, with only 1 using subcutaneous

injection). While these studies were interested in the use of opioids as analgesics or

anesthetics[36, 37], or compared acute effects of various psychoactive drugs[38], their results

may be relevant for the understanding of OUD. Indeed, comparisons of inbred mouse strains

found striking variability in the intensity of opioid physical dependence, with the same strains

showing severe or mild physical signs when withdrawal was precipitated after acute or chronic

opioid injections[39, 40]. In other words, there may be shared genetic determinants, within

each strain, for brain adaptations triggered by acute or chronic opioid exposure. A recent

single-cell RNA-sequencing study (scRNA-Seq) provides an interesting illustration of how

such strain-specific genetic traits may translate into at least partially similar transcriptional

plasticity during acute or chronic morphine exposure ([41] and *3. High cellular resolution*

approaches below).

Another important consideration is that most studies (51/73, 70%) used passive

administration, which do not model volitional or compulsive drug-taking. This reflects the

technical difficulty of intravenous SA experiments (IVSA), particularly in the mouse. Few

studies directly compared passive and active exposure. Among these, Tapocik et al[42]

analyzed "yoker" mice and groups of "yoked-morphine" or "yoked-saline" animals which, each

time the yoker self-administered morphine, were passively exposed to morphine or saline IV

injections. In C57BL/6J, much more DEG were identified between the yoker and yoked-saline

groups (n=1176) than between yoked-morphine and yoked-saline (n=244), with their overlap

likely reflecting passive morphine effects (n=103). In DBA/2J mice, which did not develop

morphine self-administration (as shown by the absence of differences in operant responding

among the 3 groups), transcriptional effects were milder, as expected (yoked-morphine *vs*

yoked-saline, n=262 DEG; yoker *vs* yoked-saline, n=107). This suggests that distinct

transcriptional regulations may mediate voluntary opioid consumption and reflect their

pharmacological effects, with both aspects ultimately contributing to OUD.

New paradigms have been developed to facilitate such studies, which notably include

oral SA[43–45]), or devices for opioid vapor SA[46]. In addition, optogenetic tools now enable

light-induced self-stimulation (optogenetic intracranial self-stimulation, oICSS)[47]. The latter

models are based on selective manipulation, using opsins, of the activity of anatomically- or

genetically-defined neuronal populations. While these approaches do not recapitulate the

systemic effects of opioid IVSA, they are nevertheless able to trigger similar behavioral

deficits, including uncontrolled consumption, resistance to punishment, or excessive

motivation[48, 49]. It is therefore possible that they may allow dissecting the neuronal

pathways that are necessary or sufficient for each of these behavioral dimensions, as well as

underlying molecular mechanisms. Another avenue for improvement relates to the mode of

action of genetic tools used for oICSS. Upon light-stimulation, opsins open ion channels that

modulate neuronal excitability. Their high temporal resolution does not necessarily represent

the best model for the longer time-frame of pharmacological action of drugs of abuse. To more

faithfully mimic such kinetics and associated intracellular signaling, chemogenetics or chimeric

opsins (for light-induced metabotropic signaling, with a proof-of-concept available for

MOR[50]) might represent better tools.

Nervous system structures. Among articles eligible for qualitative analysis, 18 regions of the

nervous system were explored (Fig.2B). Unsurprisingly, a large majority focused on the

mesocorticolimbic dopaminergic pathway: the NAc (22 articles[13, 27, 30, 33–35, 37, 51–65]),

frontal cortex (16; [13, 29, 32, 34, 62, 66–76]), whole/unspecified striatal complex (11; [31, 38,

77–85]), dorsal striatum (7; [13, 35, 59–61, 86, 87]) and ventral tegmental area (VTA, 5; [33,

62, 88–90]). Other regions included the spinal cord (7; [77, 91–96]); hippocampus (5; [13, 78,

97–99]); amygdala (3; [13, 100, 101]); locus coeruleus[89, 102], ventral midbrain[42, 103],

hypothalamus[104, 105] (2 each), whole brain with[106] or without[107] cerebellum,

periaqueductal gray matter[108], pituitary gland[105], arcuate nucleus[109], nucleus of the

tractus solitarius[36], brainstem[28], cerebellum[99] or dorsal root ganglia[110] (DRG, 1 each).

Below, we briefly describe the rationale for studying such diverse structures.

The course of OUD involves 3 stages: intoxication, withdrawal/negative affect, and anticipation of next intoxication. These stages reflect gradual adaptation of the brain to drug exposure, and rely on distinct mechanisms. The intoxication stage, mostly driven by acute reward, notably associates with dopamine release, in the striatum and frontal cortex, by neurons located in the VTA. The second stage corresponds to negative reinforcement, whereby drug consumption alleviates physical and negative affective states of withdrawal, involving the locus coeruleus[111], NAc[112], and amygdala[113], among others. Finally, anticipation results from memories of drug-associated cues and contexts (hippocampus), as well as impaired goal-directed behaviors, implicating higher-order structures (e.g. prefrontal and orbital cortices[114]). These relationships are not exclusive, with some structures implicated at multiple stages (e.g., the striatum, also involved in habit formation). Other regions (spinal cord, DRG, brainstem, periaqueductal gray matter, PAG) were investigated in relation to opioid analgesia, tolerance and hyperalgesia. Finally, few articles were interested in how opioids modulate neuroendocrine systems (hypothalamus, arcuate nucleus, pituitary gland).

In addition to these associations, it is also important to consider the distribution of the endogenous opioid system. This system is composed of 3 opioid receptors (mu, delta and kappa; MOR, DOR and KOR), among which the MOR is necessary for opioid-induced analgesia, reward, and physical dependence[115]. Neurons expressing this receptor therefore represent pharmacological entry points for brain adaptations to opioids. Comparison of studies reviewed here with MOR brain distribution (assessed using ligand autoradiography[116–121]; Fig.3C) indicates that: i) MOR is highly expressed among structures most frequently studied, as expected; ii) surprisingly, other areas with similarly high expression have received little to no attention (NTS, locus coeruleus, dorsal raphe nucleus, medial habenula, thalamus), calling for more work.

Techniques. Thirty-seven and 29 studies used microarrays and RNA-seq, respectively, to analyze the transcriptome (Fig.3D). While the first RNA-seq study was published in 2014[37], this technique has since progressively replaced microarrays, reflecting well-recognized

advantages (higher sensitivity and throughput, gene discovery). For epigenomic profiling, 3 studies used DNA methylation arrays[73, 75, 76], 2 bisulfite sequencing[72, 74], and 1 ATACseq (Assay for Transposase-Accessible Chromatin with sequencing)[87].

2. Quantitative analysis of transcriptomics studies

To identify reproducible transcriptomic opioid signatures, we ran quantitative analyses of bulk-tissue results available for the 5 most studied structures (44 articles; Fig.4A). To do so, gene-level results were retrieved when available. This yielded 24 lists of DEG comparing opioid-treated animals and control groups, originating from 17 articles. Among these, 6 articles did not report any p-values, while the remaining 11 used very diverse thresholds on p-values and fold changes (FC; Supplementary Table 1). Only a small minority (3/44, 6.8%[31, 33, 34]) provided p-values and FC for all genes. In addition, raw data were available in the public repository GEO for less than half of the studies that reported DEG (8/17, 47%). Overall, this underscores strong heterogeneity in data analysis and reporting, which limits re-use[122]. To counter such issues, efforts are being made to harmonize practices[123, 124], with guidelines enforced by funding agencies[125]. For the present review, a genome-wide meta-analysis combining effect and sample sizes across studies was not possible. We nevertheless: i) Systematically assessed concordance among available lists of DEG; ii) Conducted a threshold-free comparison of the 2 NAc studies for which genome-wide results were available.

2.1 Nucleus accumbens

Among 18 articles, only 8 provided lists of DEGs: 4 in mice (males only[35, 37, 51, 54]), 2 in rats (1 in males[56], 1 in both sexes[33]), 2 in humans (both sexes[34, 58]). The 8 articles provided a total of 3502 DEG (Fig.4B-E). Among these, the vast majority were identified by 1 study only (n=3017/3502, 86%), while 424, 57 and 4 DEG were identified across 2, 3 or 4 studies, respectively. No single genes were common to 5 or more articles. In addition, the best overlap was found among the 2 DEG lists reported in a single study by Imperio et al[56] (Fig.3E). Overall, this indicates low concordance among NAc studies.

To define factors contributing to this heterogeneity, we intersected DEG lists in each

species separately (Fig.3C-E). This, however, did not improve concordance: in humans, only

54 genes were common to both studies (Jaccard Index, JI=2.4%, Fig.4C, Supplementary

Table 2 for the matrix of pair-wise JI across all studies); in mice, 7 were identified by at least

2 studies (JI=1%, Fig. 4D); in rats, the JI increased to 15.5% (Fig.4E), mostly driven by high

overlap among the 2 aforementioned DEG lists from Imperio et al. In the latter work, the

authors identified large- (LS) and small-suppressors (SS): in LS rats, greater avoidance of a

natural reward (saccharine) that predicted heroin availability was associated, compared to SS

rats, with worse outcomes during later heroin IVSA (increased consumption, higher motivation

and relapse). Gene expression changes occurring in the LS group (compared to control rats

given access to saccharine but not heroin; n=597 genes), and in the SS group (same controls,

n=205) showed a JI of 25.5% (with striking concordance in up- and down-regulation). This

was much higher than overlap between Imperio et al and the other rat study, by Townsend et

al (Imperio-SS/Townsend, JI=1.35%; Imperio-LS/Townsend, 3.43%), suggesting that

differences across laboratories and animal facilities may significantly contribute to poor

concordance. Importantly, while the relatively small number of studies in the NAc precluded

definitive conclusions, a more powerful analysis of sources of variability was possible across

our 5 regions of interest (see section *2.6*).

2.2 Dorsal striatum

Only 2 dorsal striatum studies provided DEG (Fig.3F). One study used chronic morphine

administration interrupted by naloxone injections in male mice[35], as described above, while

the other used acute morphine injection in male rats[86]. Reflecting these differences, a very

small overlap was found between the 2 (0.62%). Therefore, while the dorsal striatum is thought

to gradually gain control over drug-taking in OUD[126], underlying transcriptomic plasticity is

poorly characterized.

2.3 Whole striatum

Six of the 11 whole striatum studies provided DEG[31, 38, 79–82] (Fig.5B): 2 investigated

chronic heroin[82] or morphine[31], while 2 compared acute effects of distinct opioids

(morphine and heroin)[38, 80], and 2 compared acute and chronic morphine[79, 81]. Most

DEG were identified by 1 study only (3308/3736, 88.5%), with only 112 genes common to 3

or more studies. Consistent with NAc results, strongest overlaps were observed for lists of

DEG coming from the same article, with for example 54 (JI=17%, Korostynski 2013[38]) and

16 (21%, Piechota 2010[80]) DEG common to acute morphine and heroin injections,

respectively, and 139 DEG (31%, Korostynski 2007 [81]; Fig.5A) common to acute and chronic

morphine administration. In sharp contrast, comparisons across, rather than within, studies

yielded poor overlaps, even when considering those that used similar drug administration:

among chronic morphine studies (Fig.S1A), the JI varied from 0.1 to 2.6% (Korostynski

2007/Skupio 2017), and, among acute morphine studies, from 2.7% to 7.8% (Fig.S1B), with

only one comparison standing out among acute heroin studies (30 common DEG, JI=19%,

Piechota 2010/Korostinsksy 2013). Importantly, 5 of these 6 studies in the whole striatum were

conducted by the same group, suggesting that significant variability may persist even among

studies conducted in the same facility, but at different periods (see also section 2.6).

2.4 Frontal cortex

Seven studies provided DEG (Fig.5B): 2 in humans[34, 70], 5 in rats[29, 32, 67–69]. Data from

Odegaard et al[29] were discarded, as they corresponded to transgenerational effects in

animals not directly exposed to opioids. For 2 additional studies [67, 69], we could not map

microarray probes to genes. Among the 4 remaining articles, little overlaps were again found

(Kuntz-Melcavage/Mendez, JI=0.5%; Kuntz-Melcavage/Seney, 1.1%; Liu/Mendez, 1.4%;

Liu/Seney, 1.6%), with no improvement within species (Liu/Kuntz-Melcavage in rats, JI=0%;

Seney/Mendez in humans, 4.1%).

2.5 Spinal cord

Among the 6 spinal cord studies, 5 did not provide DEG[91–95], while the 6th one used a

homemade microarray[77], for which correspondence between probe IDs and genes was only

partially provided. As such, no comparison was possible.

2.6 Sources of variability

Considering that pharmacological recruitment and intra-cellular signaling of MOR may trigger

a set of common transcriptional adaptations across the 5 regions of interest, we next pooled

all lists of DEGs (Fig.5D). Again, the vast majority of DEG (70.1%, 4940/7047) were unique to

a single study, while only 21.4% common to 2 studies (1509), 5.9% to 3 (418), and 1.7% to 4

or more (117; Supplementary Table 3). Interestingly, Cdkn1a (Cyclin Dependent Kinase

Inhibitor 1A) was identified by 11 studies. This gene encodes for p21, a protein involved in the

regulation of the cell-cycle and oligodendrocytes[127]. Avey et al[51] found Cdkn1a to be one

of the most significantly upregulated genes in response to acute morphine. This upregulation

occurred in oligodendrocytes, and was blocked by the opioid antagonist naltrexone, indicating

a MOR-dependent mechanism. Of note, cocaine, another major drug of abuse, similarly

recruits p21, as cocaine-induced behavioral responses were modified in p21 knockout

mice[128]. Using this knock-out line, or viral approaches for p21 manipulation, represent

appealing perspectives to further investigate this gene in OUD.

We also performed Gene Ontology (GO) analysis using DEG common to at least 3, 4

or 5 studies (Supplementary Table 4). No results were significant for the small list of 63 DEG

common to 5 or more studies. When considering the 180 DEGs common to at least 4 studies,

only 3 GO terms were enriched: cell periphery, plasma membrane and hormone activity.

Results more directly related to the nervous system emerged for the 598 DEG common to at

least 3 studies, with 46 significant terms partly related to synaptic signaling and cell-cell

communication, which may reflect general opioid-induced electrophysiological and synaptic

adaptations across brain structures.

Because weak DEG overlaps were identified within individual regions (even when

grouping studies based on species, Fig.4C-E, or treatment chronicity, Fig.S1), we sought to

more systematically assess, across the 5 regions, the impact of following factors: species,

brain structure, treatment chronicity, drug, strain, time-point of assessment following last

exposure, and article (Fig.5D). For each factor (e.g., species), pair-wise JI were classified in

2 categories depending on whether they belonged to the same sub-group within that factor (2

studies in humans; "intra"), or not (1 human study compared to another in rats; "inter"). As

expected, grouping list of DEG by species (p=2.5E-3), brain structures (3.4E-8), treatment

chronicity (p=1.4E-3), mouse genetic background (3.9E-5) or the time-point of analysis after

opioid exposure (6.7E-10) all increased concordance, with higher JI. This is consistent with

the notion that, although they remain poorly characterized, these 5 variables contribute to

opioid-induced transcriptional plasticity. In contrast, the type of opioid used had no significant

impact (p=0.059) when considering the 13 and 5 rodent DEG lists related to morphine or

heroin, respectively (human studies were discarded, as the types of opioids consumed are

diverse and poorly characterized). This is surprising, considering that these 2 MOR agonists

exhibit strong pharmacokinetic differences and would be expected to generate distinct

adaptations (heroin being a lipophilic prodrug that rapidly crosses the blood brain barrier to be

metabolized in 6-monoacetylmorphine and morphine, which in turn activate MOR). Time-

course and dose-response experiments may be necessary to detect such differences.

Importantly, concordance increased most when DEG lists came from the same publication

(p=2.5E-5), despite the fact that only 7 comparisons fell under this category. As already

mentioned, this suggests that differences across laboratories may explain a substantial part

of the lack of concordance across studies.

Finally, we refined our analysis by considering the directionality (up/down) of DEG,

focusing on genes identified by at least 5 studies (n=63). Among results from 17 out of 24

studies that reported on directionality (Fig.S2), most genes (37/63, 59%) showed discordant

results (while, for 8 others, directionality was reported by only 1 or no study). Only 18 genes

showed 100% concordance across studies (29%; 17 up-regulated, 1 down), among which 11

were identified by at least 3 studies. The latter included the aforementioned Cdkn1a, as well

as Slc2a1, Fkbp5, Sult1a1, Arrdc3, Ccdc117, Plin4, Wscd1, Arid5b, Pla2g3, and Tsc22d3.

Slc2a1 encodes for the glutamate transporter GLT1, which plays an important role in

extracellular glutamate uptake and the regulation of excitatory transmission. A recent study

showed that during prolonged morphine withdrawal, activation of KOR in the amygdala leads

to increased GLT1 expression and excitatory drive in the NAc, thereby mediating withdrawal-

induced depressive-like behaviors. Another interesting hit is Fkbp5, a chaperone and intra-

cellular negative regulator of glucocorticoid signaling that has been largely investigated in

stress-related disorders[129], including OUD[130]. The 8 remaining genes, although

comparatively less studied in relation to opioids (Fig.S2), represent candidates for future work.

2.7 Threshold-free genome-wide comparison

Next, we compared the 2 NAc studies for which full genome-wide results were available

(Seney et al[34], Townsend et al[33]). While both included males and females, only Townsend

reported sex-specific results. Therefore, we reprocessed Seney's raw data to identify DEG in

each sex separately (*Supplementary Material*). This slightly increased concordance, arguing

for the importance of accounting for sex: while JI were equal to 7.5 and 4% when comparing

initial Seney data (male/females pooled) with Townsend data in females or males, respectively

(Fig.S3A-B), they increased to 9.5% and 4.5% when considering Seney's sex-specific results

(Fig.6A-B).

We then used RRHO2 (Rank–Rank Hypergeometric Overlap[131]) to compare the 2

studies. RRHO2 iteratively performs hypergeometric testing for all combinations of thresholds

applied to each dataset, generating a "threshold-free" analysis. Because this approach takes

directionality into account, we also computed, for comparison, overlaps among either up- or

down-regulated DEG reported by each study (white crosses, Fig.6C-D). Compared to the

latter, much more significant overlaps were identified when RRHO2 considered less stringent

p-value thresholds (best hypergeometric results are shown as black crosses). By definition,

these increased overlaps corresponded to large gene lists which, although commonly

dysregulated in similar direction, individually exhibited milder FC. Interestingly, these overlaps

were detected despite the fact that Seney investigated human post mortem tissue from

individuals with OUD, while Townsend focused on rat tissue collected after 24 hours of

withdrawal from fentanyl IVSA. Also, different patterns of convergence were observed in each

sex: in females, overlaps were highly significant among both up- and down-regulated genes

while, in males, a lower concordance predominantly affected down-regulated genes. This

suggests that the rat dataset may differentially capture sex-specific signatures of the human

disorder. Overall, these analyses illustrate the utility of reporting full genome-wide data which,

in combination with threshold-free approaches, may generate deeper understanding (see

Discussion).

3. High cellular resolution approaches

Studies reviewed above indicate that opioids induce genome-wide transcriptomic changes,

with striking variability across laboratories, experimental designs, and brain regions. This likely

reflects a combination of: i) direct pharmacological opioid effects, which correspond to the

recruitment of MOR intracellular signaling and result in decreased neuronal excitability[132];

ii) indirect effects, as the chemical identity and connectivity pattern of MOR-expressing cells

determine   how   their   inhibition   leads   to   secondary   changes   affecting   other

neurotransmitters[117]. This raises 2 questions: how is each cell type, or even each brain cell,

affected by these changes? What are underlying molecular regulatory principles?

To address the first question, several groups recently used cell-type specific and

single-cell methodologies. Coffey et al[31] focused on microglial cells of the whole striatum,

using translational ribosome affinity purification (TRAP, to capture ribosomes and the

"translating transcriptome"[133, 134]). Chronic morphine effects were analyzed with, or

without, the induction of naloxone-precipitated withdrawal. Results showed that both chronic

morphine and withdrawal triggered numerous changes in gene expression, which mostly

affected GO terms related to cAMP signaling, regulation of synapses, and the cytoskeleton. A

striking negative correlation was observed between the 2 sets of results, providing a correlate,

at the level of the microglial transcriptome, of the pharmacological antagonism between

morphine and naloxone. Considering the inter-strain variability in the sensitivity to both acute

and chronic opioid physical dependence described earlier (see section *1.Qualitative analysis-*

Route and duration of opioid administration and[39, 40]), an interesting perspective would be

to determine whether naloxone similarly antagonizes transcriptomic effects of acute opioid

exposure, and how this is modulated by underlying genetic variation. Of note, surprisingly,

such cell-type specific approaches have not been used to interrogate opioid effects in neuronal

cells, let alone opioid-responsive MOR-expressing neurons which, arguably, may be

particularly responsive. This is feasible, as knock-in lines in which the Cre recombinase is

selectively active in MOR-expressing cells are now available[135–137].

Achieving higher resolution, Avey and colleagues applied single-cell RNA-seq

(scRNA-Seq, using Drop-seq[138]) to analyze transcriptomic changes induced by acute

morphine in more than 23,000 cells of the mouse NAc[51]. As is typical in scRNA-Seq,

sequencing depth yielded data for ~1,600 genes, corresponding to a reduced genomic

representation. Changes were widespread and, interestingly, most frequent in

oligodendrocytes, followed by neurons, astrocytes, and microglia, therefore uncovering a

cellular hierarchy in transcriptional sensitivity to opioids. More recently, Reiner et al[41] used

single-nucleus RNA-Seq (snRNA-seq, 10x Genomics 3') to analyze a larger number of nuclei

(190,030) and genes (>2000) in the rat NAc, also following passive and acute morphine

injection. Compared to the former study, DEG mostly affected neurons and oligodendrocytes,

followed by astrocytes and microglia. Morphine IVSA during 10 days was also investigated,

and compared to acute effects. Among DEG shared across the 2 conditions (n=431), the vast

majority were dysregulated in a similar direction (up/down), suggesting that they may reflect

morphine exposure rather than volitional drug-taking (a "yoked" group chronically exposed to

morphine would help substantiate this distinction). Importantly, a larger number of DEG were

specifically dysregulated in only one condition (n=868), with more DEG associated with IVSA

(reminiscent of the study by Tapocik et al described above), and distinct distributions among

NAc cell-types (e.g., Dopamine receptor D1- or D2-expressing neurons). By providing the first

single-cell description of morphine IVSA, these results reinforce the notion that voluntary drug-

intake may recruit partly distinct molecular mechanisms from those of non-contingent

administration.

We next compared results on acute morphine from these 2 studies. Both used

Seurat[139] and dimensionality reduction (t-SNE[51] or UMAP[41]) to identify clusters, which

were then assigned to cell-types using external data. We first compared marker genes

reported to define those clusters, when available (Avey: 11/29 clusters; Reiner: 11/19

clusters). Low agreement was found (Fig.S4A), with JI ranging from 0 up to 7.1% (for Avey's

activated medium spiny neurons, MSNs, and Reiner's D1R MSN-1), possibly because both

studies used few marker genes (Avey: 39-51; Reiner: 4-12). We then confronted morphine-

induced DEG for each cell-type (Fig.6E): low JI were found, with maximal overlap (6.4%)

obtained when comparing Avey's "myelin-forming and mature oligodendrocytes" (MO_all) with

Reiner's "Oligodendrocyte-3". Surprisingly, while some clusters assigned to similar cell-types

in the 2 studies showed some similarity in DEG (green squares, Fig.6E), strong concordance

was also observed across clusters with different identities (e.g. Reiner's "Astrocyte-2" and

Avey's MO_all; yellow squares). Overlaps among DEG (Fig.6E) did not increase as a function

of cell-type similarity across studies (Fig.S4B), as would have been expected. While a

particular emphasis has been put in sc/snRNA-Seq studies on the number of individual cells

to be sequenced, these results suggest that in order to maximize their utility: i) marker genes

used to assign cellular identities to clusters should be reported comprehensively; and ii) rather

than short lists of DEG, differential expression results should be provided for all genes

analyzed in each cluster. In addition, classical issues related to sample size, power or

statistical models for group comparisons remain pregnant. The Avey and Reiner studies used

similar sample sizes (4 and 5 replicates per opioid or control group, respectively), in the upper

range of bulk tissue studies (Fig.2D), but applied distinct tools to identify DEG (the SCDE and

MAST packages, respectively), which may partly account for discrepancies. With the rapid

evolution of this field and emerging controversies (e.g. regarding dimensionality

reduction[140]), additional efforts will be necessary to reach consensus on designs and

analyses[141, 142].

4. Non-coding and epigenomic mechanisms

In parallel to efforts to deepen cellular resolution, others have sought to decipher underlying non-coding RNA and epigenomic regulations[143]. Although neurons are post-mitotic cells, they exhibit peculiar plasticity features that may contribute to the encoding of life events, including exposure to psychoactive drugs[144]. Several non-coding RNA species have been investigated. The first report used custom-made arrays to examine micro-RNA (miR) following morphine or heroin exposure, in rat primary neuronal cultures or the mouse hippocampus[99]. 7 miR exhibited significant expression changes, among which miR-339-3p was later shown to target the 3'-UTR and down-regulate MOR in response to opioid treatment[145]. A second study used more recent arrays to interrogate all mouse miR in the NAc, following chronic heroin injections[63]. 111 miR exhibited >25% expression changes (no significance reported). The study focused on miR-218, and showed that its over-expression inhibited heroin-induced reinforcement (in conditioned place preference and IVSA), possibly by targeting MeCP2 (a protein that binds methylated DNA), suggesting a potential link between opioids, miR and DNA methylation. Other studies explored long non-coding (lncRNA) or circular RNA (circRNA). One group used 2 arrays interrogating 9,000 lncRNA[93] and 15,000 circRNA[96] to characterize chronic morphine effects in the rat spinal cord: 136 lncRNA (84 up, 52 down; nominal p-value<0.05, |log2FC|>1.5) and 2038 circRNA (896 up, 1142 down; nominal p-value<0.05, FC>2) were affected. Another report used RNA-Sequencing to focus on circRNA in the NAc[64], among which 112 were differentially expressed between mice that had received chronic morphine injections with or without electroacupuncture (51 up, 61 down; nominal p-value<0.05, |log2FC|>1.5). Understanding how these non-coding RNA contribute to OUD will require further work, as numerous mechanisms have been implicated in their physiological effects (including the sequestering of miR by circRNA, or complementarity interactions among lncRNA and mRNA).

Regarding the epigenome, a handful of candidate studies had initially provided evidence for differential DNA methylation at specific loci: the MOR gene in human blood[146–

148], and a few loci in the rat brain[149]. At the genome-wide scale, 6 recent studies (1 in

rats[62], 5 in humans investigated DNA methylation. The rat study used Whole Genome

Bisulfite Sequencing (WGBS) to analyze DNA methylation changes after heroin IVSA, in a

standard or enriched environment, in 3 brain regions (VTA, NAc and medial prefrontal

cortex)[62]. Because of low sequencing depth, DNA methylation was averaged across the

whole genome for selected features (promoters, CG islands). As expected, neither IVSA nor

environmental enrichment had any impact on these metrics, suggesting that the methylome is

not massively reprogrammed. Opioids may still, however, trigger more discrete DNA

methylation changes, consistent with the intuition that the large transcriptomic changes

associated with opioid exposure reviewed above are unlikely to occur in the absence of any

methylomic plasticity.

Two human studies conducted on bulk brain tissue strengthen this hypothesis.

Focusing on Brodmann area 9 (BA9), the first applied EPIC arrays to a cohort of OUD patients

(n=19) and controls (n=11)[75]. Among >800,000 CG sites (where DNA methylation mostly

occurs in the mammalian brain), 11,917 located at transcription start sites (TSS) showed

nominally significant differences (p-value<0.05; unfortunately, genomic localization of these

probes, and full genome-wide results, were not provided). Such studies face the difficult task

of controlling for confounding factors, as variation in genotype, age, sex, tissue cellular

composition or socio-demographic factors have all been associated with DNA methylation

changes. The second study addressed these questions[76]. Adjusting for socio-demographic

characteristics, ancestry and cellular composition, the authors investigated dorsolateral

prefrontal cortex from a large cohort (n=153) of 72 individuals who died of acute opioid

intoxication, 53 psychiatric controls, and 28 healthy controls. Although no individual CG site

survived multiple testing correction, 13 passed a relaxed significance threshold (p<1.0E-05)

for an association with opioid intoxication, one of which was located in Netrin-1, a gene

regulating KOR synthesis. At the genome-wide level, opioid use was also associated with

enriched differential methylation in the KEGG pathway related to dopaminergic signaling.

While aforementioned studies focused on bulk tissue, 3 others used Fluorescence-Activated Nuclei Sorting (FANS) to achieve cell-type specific analysis of the neuronal methylome, in the orbitofrontal cortex[72–74]. The first used 450K arrays and, comparing 37 heroin users who died from heroin overdose with 28 controls, identified 1298 differentially methylated CG sites, which were more frequently annotated to genes expressed by glutamatergic than GABAergic neurons[73]. The 2 last studies focused on a smaller cohort of male subjects only (12 OUD patients, 26 controls), using Reduced Representation oxidative Bisulfite Sequencing (RRoxBS). Compared to arrays, this allowed for the investigation of a broader set of cytosines (3.5 million CG sites at an average 10X coverage) and, importantly, the characterization of 2 types of non-canonical modifications enriched in the brain: DNA methylation in the non-CG context (also called mCH, where H stands for A, C or T)[150, 151] and hydroxymethylation at CG sites (or hmCG, generated by oxidation of methylated CG, mCG, by *TET* enzymes[152]). Differential methylation events were found to be more frequent in the non-CG (n=2352) than in the reference CG context (n=397), with enrichments in GO terms and gene sets (from Genome-Wide Association Studies, GWAS) related to neuronal function and psychiatric diseases[72]. Similarly, hmCG levels were found more frequently modified (n=1740) than mCG, at least in genomic regions covered by RRoxBS, and enriched in GO terms and co-methylation modules relevant to neuronal physiology[74]. Together, these 2 studies suggest that OUD pathophysiology may be strongly dependent on atypical, brain-enriched forms of methylomic plasticity. Of note, these human studies identified OUD-associated differences at the level of individual cytosines. However, evidence from other research fields and animal models[153, 154] argues for a model whereby DNA methylation changes with functional impact (e.g. on transcription factor binding, or interactions with other epigenetic factors) implicate groups of cytosines in close proximity and with differential methylation in similar direction (differentially methylated regions, DMR), rather than individual sites. Future work should therefore seek to combine DMR calling with the statistical models necessary to account for human confounding factors.

Interacting with these DNA modifications, epigenomic plasticity also relies on histone proteins and chromatin structure or accessibility. Surprisingly, a single study characterized histone modifications in the mouse NAc[65]. This pioneering work showed that behavioral effects of chronic morphine (including locomotor sensitization and precipitated withdrawal) were bidirectionally modulated by down- or over-expression of the histone methyltransferase G9a, and accompanied by modulation of H3K9me2 levels across thousands of genomic sites. While these results, and later work on cocaine[152, 155], suggest widespread histone changes, no follow-up work has been published in relation to opioids, to our knowledge. Similarly, only one study has investigated chromatin accessibility. Focusing on a human cohort of 10 heroin users who died from heroin overdose, and 10 healthy individuals[87], genomic regions with "open" chromatin were identified using ATAC-Seq, in the NAc and putamen. The technique was applied to neuronal and non-neuronal cells isolated by FANS. Heroin use was associated with enriched chromatin accessibility in specific gene (CG islands, promoters, 5' untranslated regions) or regulatory (peaks for the H3K4me1 histone mark, binding sites for the EZH2 transcription factor) features, as well as more frequent changes in neurons than in non-neurons (affecting GO terms related to the synapse, among others). At the behavioral level, FYN, a tyrosine kinase prioritized from human data, was then shown in the rat to significantly contribute to heroin IVSA. Overall, this work nicely illustrates how epigenetic analyses may uncover chromatin mechanisms contributing to OUD.

Discussion

Here we systematically assessed functional genomic mechanisms of opioid action and OUD, and uncovered a surprisingly low convergence across studies. As a result, a lot remains to be done to move the field forward. Below, we propose recommendations for study design and data sharing, as well as perspectives to refine behavioral modelling, improve cellular resolution, and advance multiomic data integration.

Recommendations for study design and data sharing. The low sample sizes observed in

the present review (Fig.2D,F) were surprising, considering that this factor is arguably the most

important one influencing statistical power[156–158]. Accordingly, a more rigorous analysis of

power, and larger sample sizes[159], will lead in future work to better reproducibility, and

should be enforced, especially when studying rodent outbred strains, or humans, who display

higher inter-individual variability[160, 161]. In humans, similar to what has been observed for

GWAS[162], functional genomic studies of OUD are expected to evolve in the next decade

towards collaborative efforts involving multiple labs (e.g. the SCORCH consortium[163]), and

the progressive aggregation of ever larger cohorts. Another major recommendation relates to

data sharing. While public repositories have been specifically implemented for collecting and

disseminating omic data (GEO[164], the IHEC data portal[165]) or bioinformatic pipelines

(Zenodo, GitHub), their systematic use is not yet current practice in the OUD field, as

exemplified in the present work (with similar criticisms raised in other research domains[166]).

This hampers re-analysis of published data, comparisons of results across laboratories, and

the use of threshold-free methods. We believe this last point is particularly meaningful. Most

opioid studies report subsets of DEG identified using arbitrary significance cut-offs, which can

mask subtle changes holding relevant biological information (Fig.6[131]). We therefore

strongly advocate for generalized use of threshold-free algorithms for enrichment analysis of

single datasets (GSEA), or comparison across pairs of differential analysis results (RRHO2).

Within this line, newer packages are expected to scale-up the applicability of such tools to

epigenomic data, which frequently involve larger number of observations than gene-level

counts[167].

Refine behavioral modeling. The face validity of SA or oICSS models of OUD has

significantly increased over last 2 decades[168], and we now face the task of more

systematically combining them with most recent molecular approaches. Another important

step will be to decipher the molecular mechanisms that may differentially account for distinct

behavioral dimensions, among impulsive or persistent drug-seeking, resistance to

punishment, or excessive motivation (including those modelled using passive administration,

such as tolerance, sensitization, or withdrawal). While the vast majority of studies reviewed

here used a categorical comparison of animals exposed or not to opioids, another design,

based on a dimensional approach within the exposed group, could help address the inter-

individual variability in susceptibility or resilience to OUD. The latter design has been used by

only one article in the present review[56] (section *2.1*), which contrasts with the amount of data

available for models of other psychiatric disorders, in particular depression (e.g. chronic social

defeat[169–171]. To better understand such inter-individual susceptibility, epigenetic

mechanisms are obvious candidates[172], as illustrated by 2 recent studies on DNA

methylation and oral SA of alcohol in macaques[173], and miRNA in a rat model of cocaine

SA[174]. While similar mechanisms are likely at play in OUD, they have not been addressed

by open-ended approaches yet, pointing toward an important avenue for future work.

Improve cellular resolution. This is another major avenue for future work, considering the

extreme heterogeneity of brain tissue. In addition to cell-type specific (TRAP) and sc/snRNA-

Seq (already applied to OUD, see above), a newer generation of tools now enables similar

molecular profiling while taking neuroanatomical context into consideration, with 2 possible

factors: i) connectivity patterns, probed using antero- or retrograde viruses, in combination

with TRAP (transcriptomics[175]) or FANS (epigenomics[176]); or ii) spatial context, with

methodologies based on sequencing or microscopy to detect RNA "in histological space";

while spatial capture followed by transcriptome sequencing (e.g. Slide-seq[177]) enables

genome-wide analysis with 10µm resolution, single-molecule fluorescence in situ hybridization

(smFISH) allows for better spatial resolution (1µm), but requires probe design and target gene

selection[146,147] (covering a few hundred to a few thousand genes, lower than sc/snRNA-

Seq[180]). Future work using these techniques, by characterizing molecular effects of opioids

in a particular neuronal pathway or cell-type, or with exquisite spatial resolution, should

deepen the understanding of OUD.

Advance multiomic data integration. The epigenomic and transcriptomic mechanisms

reviewed above contribute to changes at protein level that ultimately mediate

neurophysiological adaptations. Proteomic studies were initially not included in our systematic

bibliography, considering the low number of studies available and because, compared to RNA-

or DNA-sequencing methodologies, they only cover a few hundred proteins[181–184]).

Nevertheless, to go further in the understanding of OUD, future work will need to combine

transcriptomic and proteomic analyses with the characterization of upstream epigenetic

regulation. The next challenge, therefore, will be to integrate those various molecular layers

into multiomic models of pathophysiology.

Illustrating these concepts, we recently conducted a human study suggesting that

early-life adversity and depression associate with 2 forms of DNA methylation changes in the

CG and non-CG contexts (specifically focusing on CAC sites, where non-CG methylation is

most abundant)[150]. By combining WGBS with ChIP-Seq analysis of 6 histone marks, the 2

forms of plasticity were found to affect distinct genomic sites and features, and to be defined

by different histones modifications and chromatin states. Similar multiomic investigations of

opioid effects should be conducted and, importantly, will require dedicated bioinformatic

frameworks, which are only emerging in psychiatry[185]. Only 2 studies reviewed here

followed that path: Mendez et al performed transcriptomic and proteomic analyses, while Liu

et al analyzed the transcriptome and methylome (see *4.Non-coding and epigenomic*

mechanisms of opioid plasticity). In both studies, multiomic integration consisted: first, in the

identification of features (single mRNA, DNA methylation site, or protein) passing a

significance threshold, and their overlap; second, in the construction of co-expression or co-

methylation networks, and the prioritization of groups of features, defined as modules, most

significantly associated with OUD, and their biological interpretation (enrichment in functional

GO terms, significantly affected features, GWAS results, etc). While such systems-biology

approaches will undoubtedly contribute to the understanding of OUD, they remain based on

step-by-step aggregation of results generated individually for each layer. In the coming years,

it is expected that advanced multiomic integration approaches (e.g. similarity network fusion,

or multiblock methods[186–188]), may more globally define, throughout the genome, how

multiple regulatory layers interact, and the chain of events linking genetic, developmental, and

environmental factors to abnormal behavior and compulsive drug use[189].

Conclusion

Although numerous functional genomic studies of opioid action and OUD have been

conducted across animal models and patient cohorts, replicable findings appear limited. In the

future, combining refined behavioral models with multiomic integration, at the single-cell scale,

will hopefully pave the way towards the identification of innovative therapeutic targets and,

ultimately, contribute to better care.

Acknowledgements

This research was supported by the Centre National de la Recherche Scientifique (CNRS, France), 'Université de Strasbourg' (Idex Recherche Exploratoire 2022; PEL), French National Research Agency (ANR-19-CE37-0010; PEL), 'Fondation Fyssen' (Subvention recherche 2021), 'Fondation Avenir' (AAP 2022 Recherche médicale Appliquée), 'Fondation pour la Recherche Médicale' (FDT202204015236 ; CF), and 'Institut pour la Recherche en Santé Publique – Institut National du Cancer' (IReSP-INCa doctoral fellowship 2022-166 ; ACR).

Conflict of Interest

The authors declare no conflicts of interest.

Code availability

All code used in the present review is available as Supplementary File 1.

Figure legends

Figure 1. PRISMA diagram. The review protocol was registered prospectively in Prospero (record ID#CRD42022270113), and followed the Preferred Reporting Items for Systematic Review and Meta-Analysis protocol (PRISMA[190]). ***Identification of studies***. Three databases were used for systematic screening: MEDLINE, Embase and Web of Science. The literature search was performed using the following keywords: "microarray" OR "RNA sequencing" OR "bisulfite sequencing" OR "chromatin immunopurification sequencing" OR "single cell RNA sequencing" AND each of the following terms: "opiates", "opioid", "morphine", "fentanyl", "oxycodone", "heroin", "methadone" or "buprenorphine". The initial search provided 2709 potentially eligible studies. ***Screening***. Articles were screened based on their title and abstract, and included if they used a high-throughput, genome-wide methodology to assess modifications of gene expression or epigenetic mechanisms as a function of opioid exposure, in the nervous system of primates or rodents (rat, mouse), and were published in English in peer-reviewed journals. Exclusion criteria included tissue other than the nervous system and candidate genes approaches. Once duplicates were removed, a total of 1229 articles were screened, among which 1067 were excluded because their title or abstract didn't meet eligibility criteria. ***Eligibility***. After a more thorough examination of articles' full-text, an additional 89 articles were excluded. As a result, 73 papers were selected for qualitative synthesis. Among these, 44 articles reporting on transcriptomic analyses in the 5 most frequently investigated regions were selected for a more detailed quantitative synthesis. Of these, 34 articles investigated a single brain region, 9 covered 2 regions, and only 1 characterized 5 regions [13]. Overall, this resulted in 52 differential expression analyses: 18, 11, 11, 6 and 6 in the nucleus accumbens (NAc), frontal cortex, whole striatum, dorsal striatum and spinal cord, respectively.

Figure 2. Characteristics of studies included: Species & Sex. (A) Species repartition in studies identified during systematic search. **(B)** Distribution of sex and year of publication across studies. **(C-D)** Sex distribution specifically in rodent and human studies. **(E-F)** Sample

sizes in rodent and human studies that investigated both sexes and compared opioid-exposed

animals to controls (rodents, panel E), or individuals with a diagnosis of opioid use disorder

(OUD) to healthy controls (humans, panel F).

Figure 3. Characteristics of studies included: Injection parameters, Brain structure &

Techniques. Among the 73 studies considered during qualitative analysis, a large panel of

injection parameters, nervous system structures and techniques were represented. **(A)**

Injection parameters used for opioid administration. NB: "mixed" refers to a single study that

used a combination of chronic oral administration (through bottles) and acute ip injections of

morphine[79]. **(B)** Number of articles investigating each nervous system structure. **(C)** Side-

by-side comparison of the number of articles investigating each structure with the protein

expression profile of the molecular target of opioids, the mu opioid receptor (MOR; as

assessed using [^{3}H]-DAMGO autoradiography, data in fmol/mg of tissue, adapted from[115,

118–120]). **(D)** Distribution of high-throughput methodologies and year of publication across

studies. *Abbreviations:* FCx, Frontal cortex; NAc, Nucleus accumbens; dStr, Dorsal Striatum;

HT, Hypothalamus; Th, Thalamus; mHb, Medial Habenula; Hc, Hippocampus; Amy,

Amygdala; VTA, Ventral Tegmental Area; PAG, periaqueductal gray; DRN, Dorsal Raphe

Nucleus; LC, Locus Coeruleus; NTS, Nucleus Tractus Solitarius.

Figure 4. Data availability for eligible studies and quantitative analysis in the nucleus

accumbens. (A) The graph depicts the number of: i) differential expression analysis results

considered for quantitative analysis (with the corresponding number of articles between

brackets); ii) cases when lists of differentially expressed genes (DEG) were available; iii) cases

when full genome-wide results were available (with both fold changes, FC, and p-values). **(B)**

Number of DEG identified in studies that investigated the nucleus accumbens. **(C-E)** Venn

diagram of DEG identified in each species (human, mice and rat). **(F)** Venn diagram of DEG

in the 2 studies that investigated the dorsal striatum.

Figure 5. Quantitative analyses in the dorsal striatum, whole striatum and frontal cortex.

(A) Number of differentially expressed genes (DEG) identified in studies that investigated the

striatum (see Fig.S1 for separate analyses of studies that used acute or chronic opioid

administration). **(B)** Number of DEG in studies of the frontal cortex. **(C)** Number of genes

identified as differentially expressed (DE) across different studies. **(D)** Across the 5 brain

regions considered during quantitative analysis, we assessed the impact of the following 7

factors on consistency among studies, using pairwise Jaccard indexes: species, brain

structure, treatment chronicity (acute/chronic), opioid drug (morphine/heroin), strain

(C57BL6J/others), time of analysis following last exposure (5 lists of DEG from tissue collected

1 hour after opioid exposure, 5 after 2 hours, 10 after 4 hours, 4 after 8 hours, 1 after 14 hours,

and 3 after 24 hours) and article. For each factor (e.g., species), pair-wise JI were classified

in 2 categories describing whether they belonged to the same sub-group (e.g., 2 studies

conducted in humans; "intra"), or not (e.g., 1 study in humans compared to another in rats;

"inter"), within that factor. For the drug factor, studies in human were not considered as they

cannot be assigned to specific opioids; a single study investigated fentanyl, and was

excluded[33]. For the strain factor, among the 14 mouse studies available, JI significantly

increased among the 12 that used C57BL6J mice, while a similar analysis was not possible

for 129P3/J, DBA/2J, SWR/J or Kunming strains, as only 1 list of DEG was available for each.

One study reporting on the combined analysis of C57BL6J and DBA2J mice was excluded[54].

No similar analysis was possible in rats, as all available studies used Sprague-Dawley (except

one that did not state the strain used[68]). Note that the numbers of JI values considered in

each comparison are indicated at the bottom of each bar.

Figure 6. (A-D) Threshold-free comparison of 2 bulk tissue RNA-sequencing studies

(Seney et al and Townsend et al), and cell-type specific comparison of 2 single-cell

(Avey et al) and single-nucleus (Reiner et al) RNA-sequencing studies. (A-B) Venn

diagram of differentially expressed genes (DEG) identified across the 2 bulk tissue RNA-

Sequencing studies by Seney et al and Townsend et al (DEG defined by arbitrary significance

thresholds in each study: p-value < 0.05 for both) for women/female rats (A) and men/male

rats (B). **(C-D)** Genome-wide, threshold-free comparison of the same 2 studies using Rank-

Rank Hypergeometric Overlap (RRHO2) for women/female (C) rats and men/male rats (D). In

RRHO2, briefly, genes are ranked using p-values (signed according to directionality), from

upregulated (Seney: left, Townsend: bottom) to down-regulated genes (Seney: right, fentan:

top). Overlaps along the 2 distributions are then iteratively assessed using hypergeometric

testing. High p-values in the bottom-left and top-right quadrants indicates concordance,

whereas high p-values in the other 2 quadrants (bottom-right and top-left) indicate discordance

between the 2 datasets. The white crosses indicate the location of the overlap corresponding

to Venn diagrams depicted in panels A-B (split in subgroups of up- or down-regulated genes),

while black crosses correspond to most significant overlaps identified using RRHO2. **(E)**

Heatmap of pairwise Jaccard indexes for DEG identified across each of the 26 and 27 cell-

types defined by Avey et al and Reiner et al, respectively.

References

References

1.   Rosenblum A, Marsch LA, Joseph H, Portenoy RK. Opioids and the treatment of chronic pain:
controversies, current status, and future directions. Exp Clin Psychopharmacol. 2008;16:405–
416.

2.   Florence C, Luo F, Rice K. The economic burden of opioid use disorder and fatal opioid
overdose in the United States, 2017. Drug Alcohol Depend. 2021;218:108350.

3.   European Monitoring Centre for Drugs and Drug Addiction. European drug report 2017 :trends
and developments. LU: Publications Office; 2017.

4.   Heilig M, MacKillop J, Martinez D, Rehm J, Leggio L, Vanderschuren LJMJ. Addiction as a brain
disease revised: why it still matters, and the need for consilience. Neuropsychopharmacology.
2021;46:1715–1723.

5.   Maze I, Shen L, Zhang B, Garcia BA, Shao N, Mitchell A, et al. Analytical tools and current
challenges in the modern era of neuroepigenomics. Nat Neurosci. 2014;17:1476–1490.

6.   Przewlocki R. Opioid abuse and brain gene expression. Eur J Pharmacol. 2004;500:331–349.

7.   McClung CA. The molecular mechanisms of morphine addiction. Rev Neurosci. 2006;17:393–
402.

8.   Koob GF, Volkow ND. Neurobiology of addiction: a neurocircuitry analysis. Lancet Psychiatry.
2016;3:760–773.

9.   Reed B, Kreek MJ. Genetic Vulnerability to Opioid Addiction. Cold Spring Harb Perspect Med.
2021;11:a039735.

10. Bodnar RJ. Endogenous opiates and behavior: 2020. Peptides. 2022;151:170752.

11. Przewlocki R. Opioid abuse and brain gene expression. European Journal of Pharmacology.
2004;500:331–349.

12. McClung CA. The molecular mechanisms of morphine addiction. Reviews in the Neurosciences.
2006;17:393–402.

13. Yamada K, Nagai T, Nabeshima T. Drug dependence, synaptic plasticity, and tissue plasminogen
activator. J Pharmacol Sci. 2005;97:157–161.

14. UNODC, World Drug Report 2022 (United Nations publication, 2022).

15. Kerridge BT, Saha TD, Chou SP, Zhang H, Jung J, Ruan WJ, et al. Gender and nonmedical
prescription opioid use and DSM-5 nonmedical prescription opioid use disorder: Results from
the National Epidemiologic Survey on Alcohol and Related Conditions - III. Drug Alcohol
Depend. 2015;156:47–56.

16. Bravo IM, Luster BR, Flanigan ME, Perez PJ, Cogan ES, Schmidt KT, et al. Divergent behavioral
responses in protracted opioid withdrawal in male and female C57BL/6J mice. Eur J Neurosci.
2020;51:742–754.

17.   Craft RM. Sex differences in opioid analgesia: 'from mouse to man'. Clin J Pain. 2003;19:175–
849   186.

850   18.   Craft RM, Stratmann JA, Bartok RE, Walpole TI, King SJ. Sex differences in development of
851   morphine tolerance and dependence in the rat. Psychopharmacology (Berl). 1999;143:1–7.

852   19.   Cicero TJ, Nock B, Meyer ER. Gender-linked differences in the expression of physical
853   dependence in the rat. Pharmacol Biochem Behav. 2002;72:691–697.

854   20.   Karami M, Zarrindast MR. Morphine sex-dependently induced place conditioning in adult
855   Wistar rats. Eur J Pharmacol. 2008;582:78–87.

856   21.   Cicero TJ, Aylward SC, Meyer ER. Gender differences in the intravenous self-administration of
857   mu opiate agonists. Pharmacol Biochem Behav. 2003;74:541–549.

858   22.   Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. Neurosci Biobehav Rev.
859   2011;35:565–572.

860   23.   Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in neuroscience and
861   biomedical research. Neurosci Biobehav Rev. 2014;40:1–5.

862   24.   Labonté B, Engmann O, Purushothaman I, Menard C, Wang J, Tan C, et al. Sex-specific
863   transcriptional signatures in human depression. Nat Med. 2017;23:1102–1111.

864   25.   Hoffman GE, Ma Y, Montgomery KS, Bendl J, Jaiswal MK, Kozlenkov A, et al. Sex Differences in
865   the Human Brain Transcriptome of Cases With Schizophrenia. Biol Psychiatry. 2022;91:92–101.

866   26.   Hitzemann R, Bergeson SE, Berman AE, Bubier JA, Chesler EJ, Finn DA, et al. Sex Differences in
867   the Brain Transcriptome Related to Alcohol Effects and Alcohol Use Disorder. Biol Psychiatry.
868   2022;91:43–52.

869   27.   Vassoler FM, Oliver DJ, Wyse C, Blau A, Shtutman M, Turner JR, et al. Transgenerational
870   attenuation of opioid self-administration as a consequence of adolescent morphine exposure.
871   Neuropharmacology. 2017;113:271–280.

872   28.   Borrelli KN, Yao EJ, Yen WW, Phadke RA, Ruan QT, Chen MM, et al. Sex Differences in
873   Behavioral and Brainstem Transcriptomic Neuroadaptations following Neonatal Opioid
874   Exposure in Outbred Mice. ENeuro. 2021;8:ENEURO.0143-21.2021.

875   29.   Odegaard KE, Schaal VL, Clark AR, Koul S, Sankarasubramanian J, Xia Z, et al. A Holistic Systems
876   Approach to Characterize the Impact of Pre- and Post-natal Oxycodone Exposure on
877   Neurodevelopment and Behavior. Front Cell Dev Biol. 2020;8:619199.

878   30.   Odegaard KE, Schaal VL, Clark AR, Koul S, Gowen A, Sankarasubramani J, et al. Characterization
879   of the intergenerational impact of in utero and postnatal oxycodone exposure. Transl
880   Psychiatry. 2020;10:329.

881   31.   Coffey KR, Lesiak AJ, Marx RG, Vo EK, Garden GA, Neumaier JF. A cAMP-Related Gene Network
882   in Microglia Is Inversely Regulated by Morphine Tolerance and Withdrawal. Biol Psychiatry
883   Glob Open Sci. 2022;2:180–189.

884   32.   Liu SX, Gades MS, Swain Y, Ramakrishnan A, Harris AC, Tran PV, et al. Repeated morphine
885   exposure activates synaptogenesis and other neuroplasticity-related gene networks in the

886   dorsomedial prefrontal cortex of male and female rats. Drug Alcohol Depend.
887   2021;221:108598.

888   33.   Townsend EA, Kim RK, Robinson HL, Marsh SA, Banks ML, Hamilton PJ. Opioid withdrawal
889   produces sex-specific effects on fentanyl-vs.-food choice and mesolimbic transcription. Biol
890   Psychiatry Glob Open Sci. 2021;1:112–122.

891   34.   Seney ML, Kim S-M, Glausier JR, Hildebrand MA, Xue X, Zong W, et al. Transcriptional
892   Alterations in Dorsolateral Prefrontal Cortex and Nucleus Accumbens Implicate
893   Neuroinflammation and Synaptic Remodeling in Opioid Use Disorder. Biol Psychiatry.
894   2021;90:550–562.

895   35.   Lefevre EM, Pisansky MT, Toddes C, Baruffaldi F, Pravetoni M, Tian L, et al. Interruption of
896   continuous opioid exposure exacerbates drug-evoked adaptations in the mesolimbic dopamine
897   system. Neuropsychopharmacology: Official Publication of the American College of
898   Neuropsychopharmacology. 2020;45:1781–1792.

899   36.   Sugino S, Konno D, Abe J, Imamura-Kawasawa Y, Kido K, Suzuki J, et al. Crucial involvement of
900   catecholamine neurotransmission in postoperative nausea and vomiting: Whole-transcriptome
901   profiling in the rat nucleus of the solitary tract. Genes Brain Behav. 2021;e12759.

902   37.   Ye J, Yang Z, Li C, Cai M, Zhou D, Zhang Q, et al. NF-κB signaling and vesicle transport are
903   correlated with the reactivation of the memory trace of morphine dependence. Diagn Pathol.
904   2014;9:142.

905   38.   Korostynski M, Piechota M, Dzbek J, Mlynarski W, Szklarczyk K, Ziolkowska B, et al. Novel drug-
906   regulated transcriptional networks in brain reveal pharmacological properties of psychotropic
907   drugs. BMC Genomics. 2013;14:606.

908   39.   Klein G, Juni A, Waxman AR, Arout CA, Inturrisi CE, Kest B. A survey of acute and chronic heroin
909   dependence in ten inbred mouse strains: evidence of genetic correlation with morphine
910   dependence. Pharmacol Biochem Behav. 2008;90:447–452.

911   40.   Kest B, Palmese CA, Hopkins E, Adler M, Juni A, Mogil JS. Naloxone-precipitated withdrawal
912   jumping in 11 inbred mouse strains: evidence for common genetic mechanisms in acute and
913   chronic morphine physical dependence. Neuroscience. 2002;115:463–469.

914   41.   Reiner BC, Zhang Y, Stein LM, Perea ED, Arauco-Shapiro G, Ben Nathan J, et al. Single nucleus
915   transcriptomic analysis of rat nucleus accumbens reveals cell type-specific patterns of gene
916   expression associated with volitional morphine intake. Transl Psychiatry. 2022;12:374.

917   42.   Tapocik JD, Luu TV, Mayo CL, Wang B-D, Doyle E, Lee AD, et al. Neuroplasticity, axonal
918   guidance and micro-RNA genes are associated with morphine self-administration behavior.
919   Addiction Biology. 2013;18:480–495.

920   43.   Berger AC, Whistler JL. Morphine-induced mu opioid receptor trafficking enhances reward yet
921   prevents compulsive drug use. EMBO Mol Med. 2011;3:385–397.

922   44.   Slivicki RA, Earnest T, Chang Y-H, Pareta R, Casey E, Li J-N, et al. Oral oxycodone self-
923   administration leads to features of opioid misuse in male and female mice. Addict Biol.
924   2023;28:e13253.

925   45.   Phillips AG, McGovern DJ, Lee S, Ro K, Huynh DT, Elvig SK, et al. Oral prescription opioid-
926   seeking behavior in male and female mice. Addict Biol. 2020;25:e12828.

927   46.   Moussawi K, Ortiz MM, Gantz SC, Tunstall BJ, Marchette RCN, Bonci A, et al. Fentanyl vapor
928   self-administration model in mice to study opioid addiction. Sci Adv. 2020;6:eabc0413.

929   47.   Corre J, van Zessen R, Loureiro M, Patriarchi T, Tian L, Pascoli V, et al. Dopamine neurons
930   projecting to medial shell of the nucleus accumbens drive heroin reinforcement. Elife.
931   2018;7:e39945.

932   48.   Corre J, van Zessen R, Loureiro M, Patriarchi T, Tian L, Pascoli V, et al. Dopamine neurons
933   projecting to medial shell of the nucleus accumbens drive heroin reinforcement. ELife.
934   2018;7:e39945.

935   49.   Pascoli V, Terrier J, Hiver A, Lüscher C. Sufficiency of Mesolimbic Dopamine Neuron Stimulation
936   for the Progression to Addiction. Neuron. 2015;88:1054–1066.

937   50.   Siuda ER, Copits BA, Schmidt MJ, Baird MA, Al-Hasani R, Planer WJ, et al. Spatiotemporal
938   control of opioid signaling and behavior. Neuron. 2015;86:923–935.

939   51.   Avey D, Sankararaman S, Yim AKY, Barve R, Milbrandt J, Mitra RD. Single-Cell RNA-Seq
940   Uncovers a Robust Transcriptional Response to Morphine by Glia. Cell Rep. 2018;24:3619-
941   3629.e4.

942   52.   Liang J, Chen J-H, Chen X-H, Peng Y-H, Zheng X-G. Gene expression of conditioned locomotion
943   and context-specific locomotor sensitization controlled by morphine-associated environment.
944   Behav Brain Res. 2011;216:321–331.

945   53.   Tapocik JD, Luu TV, Mayo CL, Wang B-D, Doyle E, Lee AD, et al. Neuroplasticity, axonal
946   guidance and micro-RNA genes are associated with morphine self-administration behavior.
947   Addict Biol. 2013;18:480–495.

948   54.   Grice DE, Reenilä I, Männistö PT, Brooks AI, Smith GG, Golden GT, et al. Transcriptional
949   profiling of C57 and DBA strains of mice in the absence and presence of morphine. BMC
950   Genomics. 2007;8:76.

951   55.   Ikeda H, Miyatake M, Koshikawa N, Ochiai K, Yamada K, Kiss A, et al. Morphine modulation of
952   thrombospondin levels in astrocytes and its implications for neurite outgrowth and synapse
953   formation. J Biol Chem. 2010;285:38415–38427.

954   56.   Imperio CG, McFalls AJ, Colechio EM, Masser DR, Vrana KE, Grigson PS, et al. Assessment of
955   individual differences in the rat nucleus accumbens transcriptome following taste-heroin
956   extended access. Brain Res Bull. 2016;123:71–80.

957   57.   Sillivan SE, Whittard JD, Jacobs MM, Ren Y, Mazloom AR, Caputi FF, et al. ELK1 transcription
958   factor linked to dysregulated striatal mu opioid receptor signaling network and OPRM1
959   polymorphism in human heroin abusers. Biol Psychiatry. 2013;74:511–519.

960   58.   Albertson DN, Schmidt CJ, Kapatos G, Bannon MJ. Distinctive profiles of gene expression in the
961   human nucleus accumbens associated with cocaine and heroin abuse.
962   Neuropsychopharmacology. 2006;31:2304–2312.

59. Zhang Y, Liang Y, Randesi M, Yuferov V, Zhao C, Kreek MJ. Chronic Oxycodone Self-
administration Altered Reward-related Genes in the Ventral and Dorsal Striatum of C57BL/6J
Mice: An RNA-seq Analysis. Neuroscience. 2018;393:333–349.

60. Zhang Y, Liang Y, Levran O, Randesi M, Yuferov V, Zhao C, et al. Alterations of expression of
inflammation/immune-related genes in the dorsal and ventral striatum of adult C57BL/6J mice
following chronic oxycodone self-administration: a RNA sequencing study.
Psychopharmacology (Berl). 2017;234:2259–2275.

61. Yuferov V, Zhang Y, Liang Y, Zhao C, Randesi M, Kreek MJ. Oxycodone Self-Administration
Induces Alterations in Expression of Integrin, Semaphorin and Ephrin Genes in the Mouse
Striatum. Front Psychiatry. 2018;9:257.

62. Imperio CG, McFalls AJ, Hadad N, Blanco-Berdugo L, Masser DR, Colechio EM, et al. Exposure to
environmental enrichment attenuates addiction-like behavior and alters molecular effects of
heroin self-administration in rats. Neuropharmacology. 2018;139:26–40.

63. Yan B, Hu Z, Yao W, Le Q, Xu B, Liu X, et al. MiR-218 targets MeCP2 and inhibits heroin seeking
behavior. Sci Rep. 2017;7:40413.

64. Zhang H, Wang Q, Wang Q, Liu A, Qin F, Sun Q, et al. Circular RNA expression profiling in the
nucleus accumbens: Effects of electroacupuncture treatment on morphine-induced
conditioned place preference. Addict Biol. 2020;25:e12794.

65. Sun H, Maze I, Dietz DM, Scobie KN, Kennedy PJ, Damez-Werno D, et al. Morphine
epigenomically regulates behavior through alterations in histone H3 lysine 9 dimethylation in
the nucleus accumbens. J Neurosci. 2012;32:17454–17464.

66. Martin JA, Caccamise A, Werner CT, Viswanathan R, Polanco JJ, Stewart AF, et al. A Novel Role
for Oligodendrocyte Precursor Cells (OPCs) and Sox10 in Mediating Cellular and Behavioral
Responses to Heroin. Neuropsychopharmacology. 2018;43:1385–1394.

67. Ammon-Treiber S, Tischmeyer H, Riechert U, Höllt V. Gene expression of transcription factors
in the rat brain after morphine withdrawal. Neurochem Res. 2004;29:1267–1273.

68. Kuntz-Melcavage KL, Brucklacher RM, Grigson PS, Freeman WM, Vrana KE. Gene expression
changes following extinction testing in a heroin behavioral incubation model. BMC Neurosci.
2009;10:95.

69. Ammon S, Mayer P, Riechert U, Tischmeyer H, Höllt V. Microarray analysis of genes expressed
in the frontal cortex of rats chronically treated with morphine and after naloxone precipitated
withdrawal. Brain Res Mol Brain Res. 2003;112:113–125.

70. Mendez EF, Wei H, Hu R, Stertz L, Fries GR, Wu X, et al. Angiogenic gene networks are
dysregulated in opioid use disorder: evidence from multi-omics and imaging of postmortem
human brain. Mol Psychiatry. 2021;26:7803–7812.

71. Jiang C, Wang X, Le Q, Liu P, Liu C, Wang Z, et al. Morphine coordinates SST and PV
interneurons in the prelimbic cortex to disinhibit pyramidal neurons and enhance reward. Mol
Psychiatry. 2021;26:1178–1193.

72. Nagamatsu ST, Rompala G, Hurd YL, Núñez-Rios DL, Montalvo-Ortiz JL, Traumatic Stress Brain
Research Group. CpH methylome analysis in human cortical neurons identifies novel gene
pathways and drug targets for opioid use disorder. Front Psychiatry. 2022;13:1078894.

73. Kozlenkov A, Jaffe AE, Timashpolsky A, Apontes P, Rudchenko S, Barbu M, et al. DNA
Methylation Profiling of Human Prefrontal Cortex Neurons in Heroin Users Shows Significant
Difference between Genomic Contexts of Hyper- and Hypomethylation and a Younger
Epigenetic Age. Genes (Basel). 2017;8:152.

74. Rompala G, Nagamatsu ST, Martínez-Magaña JJ, Nuñez-Ríos DL, Wang J, Girgenti MJ, et al.
Profiling neuronal methylome and hydroxymethylome of opioid use disorder in the human
orbitofrontal cortex. Nat Commun. 2023;14:4544.

75. Liu A, Dai Y, Mendez EF, Hu R, Fries GR, Najera KE, et al. Genome-Wide Correlation of DNA
Methylation and Gene Expression in Postmortem Brain Tissues of Opioid Use Disorder Patients.
The International Journal of Neuropsychopharmacology. 2021;24:879–891.

76. Shu C, Sosnowski DW, Tao R, Deep-Soboslay A, Kleinman JE, Hyde TM, et al. Epigenome-wide
study of brain DNA methylation following acute opioid intoxication. Drug and Alcohol
Dependence. 2021;221:108658.

77. Loguinov AV, Anderson LM, Crosby GJ, Yukhananov RY. Gene expression following acute
morphine administration. Physiol Genomics. 2001;6:169–181.

78. Choi MR, Jin Y-B, Bang SH, Im C-N, Lee Y, Kim H-N, et al. Age-related Effects of Heroin on Gene
Expression in the Hippocampus and Striatum of Cynomolgus Monkeys. Clin Psychopharmacol
Neurosci. 2020;18:93–108.

79. Skupio U, Sikora M, Korostynski M, Wawrzczak-Bargiela A, Piechota M, Ficek J, et al. Behavioral
and transcriptional patterns of protracted opioid self-administration in mice. Addict Biol.
2017;22:1802–1816.

80. Piechota M, Korostynski M, Solecki W, Gieryk A, Slezak M, Bilecki W, et al. The dissection of
transcriptional modules regulated by various drugs of abuse in the mouse striatum. Genome
Biol. 2010;11:R48.

81. Korostynski M, Piechota M, Kaminska D, Solecki W, Przewlocki R. Morphine effects on striatal
transcriptome in mice. Genome Biol. 2007;8:R128.

82. Piechota M, Korostynski M, Sikora M, Golda S, Dzbek J, Przewlocki R. Common transcriptional
effects in the mouse striatum following chronic treatment with heroin and methamphetamine.
Genes Brain Behav. 2012;11:404–414.

83. Korostynski M, Piechota M, Golda S, Przewlocki R. High-throughput gene expression profiling
of opioid-induced alterations in discrete brain areas. Methods Mol Biol. 2015;1230:65–76.

84. Choi MR, Jin Y-B, Kim H-N, Chai YG, Im C-N, Lee S-R, et al. Gene expression in the striatum of
cynomolgus monkeys after chronic administration of cocaine and heroin. Basic Clin Pharmacol
Toxicol. 2021;128:686–698.

85. Cha HJ, Choi M-J, Ahn J-I, Jeon S-H, Kang H, Kim EJ, et al. Comparison of gene expression
profiles in drug-withdrawn rats. Mol Cell Toxicol. 2016;12:197–207.

86. Valderrama-Carvajal A, Irizar H, Gago B, Jiménez-Urbieta H, Fuxe K, Rodríguez-Oroz MC, et al.
Transcriptomic integration of D4R and MOR signaling in the rat caudate putamen. Sci Rep.
2018;8:7337.

87. Egervari G, Akpoyibo D, Rahman T, Fullard JF, Callens JE, Landry JA, et al. Chromatin
accessibility mapping of the striatum identifies tyrosine kinase FYN as a therapeutic target for
heroin use disorder. Nat Commun. 2020;11:4634.

88. Heller EA, Kaska S, Fallon B, Ferguson D, Kennedy PJ, Neve RL, et al. Morphine and cocaine
increase serum- and glucocorticoid-inducible kinase 1 activity in the ventral tegmental area. J
Neurochem. 2015;132:243–253.

89. McClung CA, Nestler EJ, Zachariou V. Regulation of gene expression by chronic morphine and
morphine withdrawal in the locus ceruleus and ventral tegmental area. J Neurosci.
2005;25:6005–6015.

90. Zhang H, Wang Q, Sun Q, Qin F, Nie D, Li Q, et al. Effects of Compound 511 on BDNF-TrkB
Signaling in the Mice Ventral Tegmental Area in Morphine-Induced Conditioned Place
Preference. Cell Mol Neurobiol. 2021;41:961–975.

91. Jokinen V, Sidorova Y, Viisanen H, Suleymanova I, Tiilikainen H, Li Z, et al. Differential Spinal
and Supraspinal Activation of Glia in a Rat Model of Morphine Tolerance. Neuroscience.
2018;375:10–24.

92. Cheng Y-C, Tsai R-Y, Sung Y-T, Chen I-J, Tu T-Y, Mao Y-Y, et al. Melatonin regulation of
transcription in the reversal of morphine tolerance: Microarray analysis of differential gene
expression. Int J Mol Med. 2019;43:791–806.

93. Shao J, Wang J, Huang J, Liu C, Pan Y, Guo Q, et al. Identification of lncRNA expression profiles
and ceRNA analysis in the spinal cord of morphine-tolerant rats. Mol Brain. 2018;11:21.

94. Peng Y, Guo G, Shu B, Liu D, Su P, Zhang X, et al. Spinal CX3CL1/CX3CR1 May Not Directly
Participate in the Development of Morphine Tolerance in Rats. Neurochem Res. 2017;42:3254–
3267.

95. Santiago JM, Rosas O, Torrado AI, González MM, Kalyan-Masih PO, Miranda JD. Molecular,
anatomical, physiological, and behavioral studies of rats treated with buprenorphine after
spinal cord injury. J Neurotrauma. 2009;26:1783–1793.

96. Weng Y, Wu J, Li L, Shao J, Li Z, Deng M, et al. Circular RNA expression profile in the spinal cord
of morphine tolerated rats and screen of putative key circRNAs. Mol Brain. 2019;12:79.

97. Juul SE, Beyer RP, Bammler TK, Farin FM, Gleason CA. Effects of neonatal stress and morphine
on murine hippocampal gene expression. Pediatr Res. 2011;69:285–292.

98. Marie-Claire C, Courtin C, Robert A, Gidrol X, Roques BP, Noble F. Sensitization to the
conditioned rewarding effects of morphine modulates gene expression in rat hippocampus.
Neuropharmacology. 2007;52:430–435.

99. Zheng H, Zeng Y, Zhang X, Chu J, Loh HH, Law P-Y. mu-Opioid receptor agonists differentially
regulate the expression of miR-190 and NeuroD. Mol Pharmacol. 2010;77:102–109.

100. Befort K, Filliol D, Ghate A, Darcq E, Matifas A, Muller J, et al. Mu-opioid receptor activation
1079   induces transcriptional plasticity in the central extended amygdala. Eur J Neurosci.
1080   2008;27:2973–2984.

1081   101. Rodriguez Parkitna JM, Bilecki W, Mierzejewski P, Stefanski R, Ligeza A, Bargiela A, et al. Effects
1082   of morphine on gene expression in the rat amygdala. J Neurochem. 2004;91:38–48.

1083   102. Zachariou V, Liu R, LaPlant Q, Xiao G, Renthal W, Chan GC, et al. Distinct roles of adenylyl
1084   cyclases 1 and 8 in opiate dependence: behavioral, electrophysiological, and molecular studies.
1085   Biol Psychiatry. 2008;63:1013–1021.

1086   103. Saad MH, Rumschlag M, Guerra MH, Savonen CL, Jaster AM, Olson PD, et al. Differentially
1087   expressed gene networks, biomarkers, long noncoding RNAs, and shared responses with
1088   cocaine identified in the midbrains of human opioid abusers. Scientific Reports. 2019;9:1534.

1089   104. Befort K, Filliol D, Darcq E, Ghate A, Matifas A, Lardenois A, et al. Gene expression is altered in
1090   the lateral hypothalamus upon activation of the mu opioid receptor. Ann N Y Acad Sci.
1091   2008;1129:175–184.

1092   105. Anghel A, Jamieson C a. M, Ren X, Young J, Porche R, Ozigbo E, et al. Gene expression profiling
1093   following short-term and long-term morphine exposure in mice uncovers genes involved in
1094   food intake. Neuroscience. 2010;167:554–566.

1095   106. Hayase T, Yamamoto Y, Yamamoto K, Muso E, Shiota K. Microarray profile analysis of toxic
1096   cocaine-induced alterations in the expression of mouse brain gene sequences: a possible
1097   'protective' effect of buprenorphine. J Appl Toxicol. 2004;24:15–20.

1098   107. Hassan HE, Myers AL, Lee IJ, Chen H, Coop A, Eddington ND. Regulation of gene expression in
1099   brain tissues of rats repeatedly treated by the highly abused opioid agonist, oxycodone:
1100   microarray profiling and gene mapping analysis. Drug Metab Dispos. 2010;38:157–167.

1101   108. Gaspari S, Purushothaman I, Cogliani V, Sakloth F, Neve RL, Howland D, et al. Suppression of
1102   RGSz1 function optimizes the actions of opioid analgesics by mechanisms that involve the
1103   Wnt/β-catenin pathway. Proc Natl Acad Sci U S A. 2018;115:E2085–E2094.

1104   109. Toorie AM, Vassoler FM, Qu F, Schonhoff CM, Bradburn S, Murgatroyd CA, et al. A history of
1105   opioid exposure in females increases the risk of metabolic disorders in their future male
1106   offspring. Addict Biol. 2021;26:e12856.

1107   110. Thibault K, Calvino B, Rivals I, Marchand F, Dubacq S, McMahon SB, et al. Molecular
1108   mechanisms underlying the enhanced analgesic effect of oxycodone compared to morphine in
1109   chemotherapy-induced neuropathic pain. PLoS One. 2014;9:e91297.

1110   111. Delfs JM, Zhu Y, Druhan JP, Aston-Jones G. Noradrenaline in the ventral forebrain is critical for
1111   opiate withdrawal-induced aversion. Nature. 2000;403:430–434.

1112   112. Zhu Y, Wienecke CF, Nachtrab G, Chen X. A thalamic input to the nucleus accumbens mediates
1113   opiate dependence. Nature. 2016;530:219–222.

1114   113. Koob GF. The dark side of emotion: the addiction perspective. Eur J Pharmacol. 2015;753:73–
1115   87.

114. Hearing M. Prefrontal-accumbens opioid plasticity: Implications for relapse and dependence.
Pharmacol Res. 2019;139:158–165.

115. Lutz P-E, Kieffer BL. The multiple facets of opioid receptor function: implications for addiction.
Curr Opin Neurobiol. 2013;23:473–479.

116. Lutz P-E, Almeida D, Filliol D, Jollant F, Kieffer BL, Turecki G. Increased functional coupling of
the mu opioid receptor in the anterior insula of depressed individuals.
Neuropsychopharmacology. 2021;46:920–927.

117. Lutz P-E, Kieffer BL. Opioid receptors: distinct roles in mood disorders. Trends Neurosci.
2013;36:195–206.

118. Kitchen I, Slowe SJ, Matthes HW, Kieffer B. Quantitative autoradiographic mapping of mu-,
delta- and kappa-opioid receptors in knockout mice lacking the mu-opioid receptor gene. Brain
Res. 1997;778:73–88.

119. Goody RJ, Oakley SM, Filliol D, Kieffer BL, Kitchen I. Quantitative autoradiographic mapping of
opioid receptors in the brain of delta-opioid receptor gene knockout mice. Brain Res.
2002;945:9–19.

120. Slowe SJ, Simonin F, Kieffer B, Kitchen I. Quantitative autoradiography of mu-,delta- and
kappa1 opioid receptors in kappa-opioid receptor knockout mice. Brain Res. 1999;818:335–
345.

121. Xia Y, Haddad GG. Ontogeny and distribution of opioid receptors in the rat brainstem. Brain
Res. 1991;549:181–193.

122. Miyakawa T. No raw data, no science: another possible source of the reproducibility crisis. Mol
Brain. 2020;13:24.

123. Field D, Sansone S-A, Collis A, Booth T, Dukes P, Gregurick SK, et al. Megascience. 'Omics data
sharing. Science. 2009;326:234–236.

124. FAIR Principles. GO FAIR. https://www.go-fair.org/fair-principles/. Accessed 14 February 2023.

125. NIH Genomic Data Sharing.

126. Murray JE, Belin-Rauscent A, Simon M, Giuliano C, Benoit-Marand M, Everitt BJ, et al.
Basolateral and central amygdala differentially recruit and maintain dorsolateral striatum-
dependent cocaine-seeking habits. Nat Commun. 2015;6:10088.

127. Zezula J, Casaccia-Bonnefil P, Ezhevsky SA, Osterhout DJ, Levine JM, Dowdy SF, et al. p21cip1 is
required for the differentiation of oligodendrocytes independently of cell cycle withdrawal.
EMBO Rep. 2001;2:27–34.

128. Scholpa NE, Briggs SB, Wagner JJ, Cummings BS. Cyclin-Dependent Kinase Inhibitor 1a (p21)
Modulates Response to Cocaine and Motivated Behaviors. J Pharmacol Exp Ther. 2016;357:56–
65.

129. Häusl AS, Brix LM, Hartmann J, Pöhlmann ML, Lopez J-P, Menegaz D, et al. The co-chaperone
Fkbp5 shapes the acute stress response in the paraventricular nucleus of the hypothalamus of
male mice. Mol Psychiatry. 2021;26:3060–3076.

130. Levran O, Peles E, Randesi M, Li Y, Rotrosen J, Ott J, et al. Stress-related genes and heroin
addiction: a role for a functional FKBP5 haplotype. Psychoneuroendocrinology. 2014;45:67–76.

131. Cahill KM, Huo Z, Tseng GC, Logan RW, Seney ML. Improved identification of concordant and
discordant gene expression signatures using an updated rank-rank hypergeometric overlap
approach. Scientific Reports. 2018;8:9588.

132. Al-Hasani R, Bruchas MR. Molecular mechanisms of opioid receptor-dependent signaling and
behavior. Anesthesiology. 2011;115:1363–1381.

133. Nectow AR, Moya MV, Ekstrand MI, Mousa A, McGuire KL, Sferrazza CE, et al. Rapid Molecular
Profiling of Defined Cell Types Using Viral TRAP. Cell Rep. 2017;19:655–667.

134. Heiman M, Schaefer A, Gong S, Peterson JD, Day M, Ramsey KE, et al. A translational profiling
approach for the molecular characterization of CNS cell types. Cell. 2008;135:738–748.

135. Bailly J, Del Rossi N, Runtz L, Li J-J, Park D, Scherrer G, et al. Targeting morphine-responsive
neurons: generation of a knock-in mouse line expressing Cre recombinase from the mu opioid
receptor gene locus. ENeuro. 2020. 7 May 2020. https://doi.org/10.1523/ENEURO.0433-
19.2020.

136. Okunomiya T, Hioki H, Nishimura C, Yawata S, Imayoshi I, Kageyama R, et al. Generation of a
MOR-CreER knock-in mouse line to study cells and neural circuits involved in mu opioid
receptor signaling. Genesis. 2020;58:e23341.

137. Märtin A, Calvigioni D, Tzortzi O, Fuzik J, Wärnberg E, Meletis K. A Spatiomolecular Map of the
Striatum. Cell Rep. 2019;29:4320-4333.e5.

138. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel Genome-
wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell. 2015;161:1202–
1214.

139. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al. Comprehensive
Integration of Single-Cell Data. Cell. 2019;177:1888-1902.e21.

140. Chari T, Pachter L. The Specious Art of Single-Cell Genomics. Genomics; 2021.

141. Gibson G. Perspectives on rigor and reproducibility in single cell genomics. PLoS Genet.
2022;18:e1010210.

142. Schmid KT, Höllbacher B, Cruceanu C, Böttcher A, Lickert H, Binder EB, et al. scPower
accelerates and optimizes the design of multi-sample single cell transcriptomic studies. Nat
Commun. 2021;12:6625.

143. Browne CJ, Godino A, Salery M, Nestler EJ. Epigenetic Mechanisms of Opioid Addiction. Biol
Psychiatry. 2020;87:22–33.

144. Belzeaux R, Lalanne L, Kieffer BL, Lutz PE. Focusing on the Opioid System for Addiction
Biomarker Discovery. Trends Mol Med. 2018;24:206–220.

145. Wu Q, Hwang CK, Zheng H, Wagley Y, Lin H-Y, Kim DK, et al. MicroRNA 339 down-regulates μ-
opioid receptor at the post-transcriptional level in response to opioid treatment. FASEB J.
2013;27:522–535.

146. Ebrahimi G, Asadikaram G, Akbari H, Nematollahi MH, Abolhassani M, Shahabinejad G, et al.
Elevated levels of DNA methylation at the OPRM1 promoter region in men with opioid use
disorder. Am J Drug Alcohol Abuse. 2017:1–7.

147. Nielsen DA, Yuferov V, Hamon S, Jackson C, Ho A, Ott J, et al. Increased OPRM1 DNA
methylation in lymphocytes of methadone-maintained former heroin addicts.
Neuropsychopharmacology. 2009;34:867–873.

148. Doehring A, Oertel BG, Sittl R, Lotsch J. Chronic opioid use is associated with increased DNA
methylation correlating with increased clinical pain. Pain. 2013;154:15–23.

149. Barrow TM, Byun HM, Li X, Smart C, Wang YX, Zhang Y, et al. The effect of morphine upon DNA
methylation in ten regions of the rat brain. Epigenetics. 2017;12:1038–1047.

150. Lutz P-E, Chay M-A, Pacis A, Chen GG, Aouabed Z, Maffioletti E, et al. Non-CG methylation and
multiple histone profiles associate child abuse with immune and small GTPase dysregulation.
Nat Commun. 2021;12:1132.

151. Lutz P-E, Mechawar N, Turecki G. Neuropathology of suicide: recent findings and future
directions. Mol Psychiatry. 2017;22:1395–1412.

152. Feng J, Shao N, Szulwach KE, Vialou V, Huynh J, Zhong C, et al. Role of Tet1 and 5-
hydroxymethylcytosine in cocaine action. Nat Neurosci. 2015;18:536–544.

153. Benelli M, Franceschini GM, Magi A, Romagnoli D, Biagioni C, Migliaccio I, et al. Charting
differentially methylated regions in cancer with Rocker-meth. Commun Biol. 2021;4:1249.

154. Kaluscha S, Domcke S, Wirbelauer C, Stadler MB, Durdu S, Burger L, et al. Evidence that direct
inhibition of transcription factor binding is the prevailing mode of gene and repeat repression
by DNA methylation. Nat Genet. 2022;54:1895–1906.

155. Feng J, Wilkinson M, Liu X, Purushothaman I, Ferguson D, Vialou V, et al. Chronic cocaine-
regulated epigenomic changes in mouse nucleus accumbens. Genome Biol. 2014;15:R65.

156. Ching T, Huang S, Garmire LX. Power analysis and sample size estimation for RNA-Seq
differential expression. RNA. 2014;20:1684–1696.

157. Baccarella A, Williams CR, Parrish JZ, Kim CC. Empirical assessment of the impact of sample
number and read depth on RNA-Seq analysis workflow performance. BMC Bioinformatics.
2018;19:423.

158. Hoffman GE, Jaffe AE, Gandal MJ, Collado-Torres L, Sieberts SK, Devlin B, et al. Comment on:
What genes are differentially expressed in individuals with schizophrenia? A systematic review.
Mol Psychiatry. 2023;28:523–525.

159. Khan AH, Bagley JR, LaPierre N, Gonzalez-Figueroa C, Spencer TC, Choudhury M, et al. Genetic
pathways regulating the longitudinal acquisition of cocaine self-administration in a panel of
inbred and recombinant inbred mice. Cell Rep. 2023;42:112856.

160. Hekman JP, Johnson JL, Kukekova AV. Transcriptome Analysis in Domesticated Species:
Challenges and Strategies. Bioinform Biol Insights. 2015;9:21–31.

161. Williams AG, Thomas S, Wyman SK, Holloway AK. RNA-seq Data: Challenges in and
Recommendations for Experimental Design and Analysis. Curr Protoc Hum Genet.
2014;83:11.13.1-20.

162. Crist RC, Reiner BC, Berrettini WH. A review of opioid addiction genetics. Curr Opin Psychol.
2019;27:31–35.

163. scorch - Home. https://scorch.igs.umaryland.edu/. Accessed 19 May 2023.

164. Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI GEO: archive
for functional genomics data sets--update. Nucleic Acids Res. 2013;41:D991-995.

165. Data Portal | IHEC. https://epigenomesportal.ca/ihec/. Accessed 19 May 2023.

166. Savage CJ, Vickers AJ. Empirical study of data sharing by authors publishing in PLoS journals.
PLoS One. 2009;4:e7078.

167. Piron A, Szymczak F, Alvelos MI, Defrance M, Lenaerts T, Eizirik DL, et al. RedRibbon: A new
rank-rank hypergeometric overlap pipeline to compare gene and transcript expression
signatures. Bioinformatics; 2022.

168. Deroche-Gamonet V, Belin D, Piazza PV. Evidence for addiction-like behavior in the rat.
Science. 2004;305:1014–1017.

169. Domingo-Rodriguez L, Ruiz de Azua I, Dominguez E, Senabre E, Serra I, Kummer S, et al. A
specific prelimbic-nucleus accumbens pathway controls resilience versus vulnerability to food
addiction. Nat Commun. 2020;11:782.

170. Laine MA, Trontti K, Misiewicz Z, Sokolowska E, Kulesskaya N, Heikkinen A, et al. Genetic
Control of Myelin Plasticity after Chronic Psychosocial Stress. ENeuro. 2018;5:ENEURO.0166-
18.2018.

171. Bagot RC, Cates HM, Purushothaman I, Lorsch ZS, Walker DM, Wang J, et al. Circuit-wide
Transcriptional Profiling Reveals Brain Region-Specific Gene Networks Regulating Depression
Susceptibility. Neuron. 2016;90:969–983.

172. Roy B, Wang Q, Dwivedi Y. Long Noncoding RNA-Associated Transcriptomic Changes in
Resiliency or Susceptibility to Depression and Response to Antidepressant Treatment. Int J
Neuropsychopharmacol. 2018;21:461–472.

173. Cervera-Juanes R, Wilhelm LJ, Park B, Grant KA, Ferguson B. Alcohol-dose-dependent DNA
methylation and expression in the nucleus accumbens identifies coordinated regulation of
synaptic genes. Transl Psychiatry. 2017;7:e994.

174. Domingo-Rodriguez L, Cabana-Domínguez J, Fernàndez-Castillo N, Cormand B, Martín-García E,
Maldonado R. Differential expression of miR-1249-3p and miR-34b-5p between vulnerable and
resilient phenotypes of cocaine addiction. Addict Biol. 2022;27:e13201.

175. Ekstrand MI, Nectow AR, Knight ZA, Latcha KN, Pomeranz LE, Friedman JM. Molecular profiling
of neurons based on connectivity. Cell. 2014;157:1230–1242.

176. Zhang Z, Zhou J, Tan P, Pang Y, Rivkin AC, Kirchgessner MA, et al. Epigenomic diversity of
cortical projection neurons in the mouse brain. Nature. 2021;598:167–173.

177. Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al. Slide-seq: A
scalable technology for measuring genome-wide expression at high spatial resolution. Science.
2019;363:1463–1467.

178. Eng C-HL, Lawson M, Zhu Q, Dries R, Koulena N, Takei Y, et al. Transcriptome-scale super-
resolved imaging in tissues by RNA seqFISH. Nature. 2019;568:235–239.

179. Borm LE, Mossi Albiach A, Mannens CCA, Janusauskas J, Özgün C, Fernández-García D, et al.
Scalable in situ single-cell profiling by electrophoretic capture of mRNA using EEL FISH. Nat
Biotechnol. 2022. 22 September 2022. https://doi.org/10.1038/s41587-022-01455-3.

180. Armand EJ, Li J, Xie F, Luo C, Mukamel EA. Single-Cell Sequencing of Brain Cell Transcriptomes
and Epigenomes. Neuron. 2021;109:11–26.

181. Mendez EF, Wei H, Hu R, Stertz L, Fries GR, Wu X, et al. Angiogenic gene networks are
dysregulated in opioid use disorder: evidence from multi-omics and imaging of postmortem
human brain. Molecular Psychiatry. 2021;26:7803–7812.

182. Li KW, Jimenez CR, van der Schors RC, Hornshaw MP, Schoffelmeer ANM, Smit AB. Intermittent
administration of morphine alters protein expression in rat nucleus accumbens. Proteomics.
2006;6:2003–2008.

183. Van den Oever MC, Lubbers BR, Goriounova NA, Li KW, Van der Schors RC, Loos M, et al.
Extracellular matrix plasticity and GABAergic inhibition of prefrontal cortex pyramidal cells
facilitates relapse to heroin seeking. Neuropsychopharmacology. 2010;35:2120–2133.

184. Bu Q, Yang Y, Yan G, Hu Z, Hu C, Duan J, et al. Proteomic analysis of the nucleus accumbens in
rhesus monkeys of morphine dependence and withdrawal intervention. J Proteomics.
2012;75:1330–1342.

185. Mokhtari A, Porte B, Belzeaux R, Etain B, Ibrahim EC, Marie-Claire C, et al. The molecular
pathophysiology of mood disorders: From the analysis of single molecular layers to multi-omic
integration. Prog Neuropsychopharmacol Biol Psychiatry. 2022;116:110520.

186. Li C-X, Wheelock CE, Sköld CM, Wheelock ÅM. Integration of multi-omics datasets enables
molecular classification of COPD. Eur Respir J. 2018;51:1701930.

187. Yang M, Matan-Lithwick S, Wang Y, De Jager PL, Bennett DA, Felsky D. Multi-omic integration
via similarity network fusion to detect molecular subtypes of ageing. Brain Commun.
2023;5:fcad110.

188. Revilla L, Mayorgas A, Corraliza AM, Masamunt MC, Metwaly A, Haller D, et al. Multi-omic
modelling of inflammatory bowel disease with regularized canonical correlation analysis. PLoS
One. 2021;16:e0246367.

189. Hasin Y, Seldin M, Lusis A. Multi-omics approaches to disease. Genome Biol. 2017;18:83.

190. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, et al. The PRISMA
statement for reporting systematic reviews and meta-analyses of studies that evaluate
healthcare interventions: explanation and elaboration. BMJ. 2009;339:b2700–b2700.

1304

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6