Functional genomic mechanisms of opioids: a systematic review of animal and human studies

Camille Falconnier[1], Alba Caparros-Roissard[1], Charles Decraene[1,2], Pierre-Eric Lutz[1]

[1] Centre National de la Recherche Scientifique, Université de Strasbourg, Institut des Neurosciences Cellulaires et Intégratives UPR3212, 67000 Strasbourg, France

[2] Centre National de la Recherche Scientifique, Université de Strasbourg, Laboratoire de Neurosciences Cognitives et Adaptatives, 67000 Strasbourg, France

Corresponding author:

Pierre-Eric Lutz, INCI UPR 3212, 8 allée du général Rouvillois, 67000 Strasbourg, France

Email: pierreeric.lutz@gmail.com

Word count: 6924

6 Figures, 4 Supplementary Figures, 6 Supplementary Tables, 1 Supplementary Material, 1 Supplementary File

Abstract

In the past two decades, over-prescription of opioids for pain management has driven a steep increase in opioid use disorder (OUD) and death by overdose, exerting a dramatic toll on western countries. OUD is a chronic relapsing disease associated with a lifetime struggle to control drug consumption, suggesting that opioids trigger long-lasting brain adaptations, notably through functional genomic and epigenomic mechanisms. Current understanding of these processes, however, remain scarce, and have not been previously reviewed systematically. To do so, the goal of the present work was to synthesize current knowledge on genome-wide transcriptomic and epigenetic mechanisms of opioid action, in primate and rodent species. Using a prospectively registered methodology, comprehensive literature searches were completed in PubMed, Embase, and Web of Science. Of the 2705 articles identified, 69 met our inclusion criteria and were considered for qualitative analysis. Focusing on the 5 most studied nervous system structures (nucleus accumbens, frontal cortex, whole striatum, dorsal striatum, spinal cord; 44 articles), we also conducted a quantitative analysis of differentially expressed genes, in an effort to identify a putative core transcriptional signature of opioids. Only one gene, Cdkn1a, was replicated in six studies and, globally, our results unveil surprisingly low consistency across published work, even when considering most recent single-cell approaches. Analysis of putative sources for this poor reproducibility suggest contributions from species, brain structures, duration of opioid exposure, as well as batch effects. To go beyond those limitations, we leveraged threshold-free methods to illustrate how genome-wide comparisons may generate new findings and hypotheses. Finally, we discuss current methodological development in the field, and their implication for future research and, ultimately, better care.

Keywords

Opioid, substance use disorder, genomics, epigenomics, next generation sequencing

Introduction

Opioids are compounds derived from opium that have been used for millennia for their analgesic and euphoric properties [1]. While they remain reference treatment for severe pain, their chronic use also associates with a plethora of adverse effects, including tolerance, physical dependence and, in some individuals, the emergence of an opioid use disorder (OUD). OUD is defined as a problematic pattern of use that leads to clinically significant impairment. This severe and chronic disorder associates with increased mortality, with life expectancy decreased by more than 10 years due to somatic and psychiatric comorbidities. Over the last 2 decades, western countries have faced an increase in death by opioid overdose, due to more frequent prescription for pain management, increasing use of illegal compounds, or misuse of substitution therapies. From a socio-economic point-of-view, OUD imposes a major burden, with an estimated annual cost in the trillion dollar range in the US [2], and increasing use and harm in Europe [3]. This results, overall, in a worsening public health crisis.

OUD results from a variety of biological, psychological and socio-economic factors that show reciprocal interactions [4]. At biological level, it originates from pharmacological effects of opioid drugs, which trigger neurochemical and molecular adaptations within the brain, under modulation by genetic vulnerability and epigenetic regulatory factors. In turn, these effects are thought responsible for behavioral and cognitive dysfunctions that define the disease, including the inability to control drug use despite harmful consequences, or persistent and intense desire for the drug, even after years of abstinence.

Over the last 2 decades, high-throughput approaches have been harnessed to characterize at unprecedented scale, throughout the full genome [5], the molecular underpinnings of opioid effects in brain tissue, across various experimental models and species. In this context, the goal of the present systematic review was to synthesize and discuss current knowledge on functional genomic mechanisms of opioids, defined as changes in gene expression or epigenetic regulation that occur as a function of opioid exposure. To do so, we used a preregistered methodology, and performed an unbiased survey of bibliographic

repositories, focusing on studies that leveraged high-throughput approaches, most notably

microarrays and next-generation sequencing. This focus on genome-wide analyses is

motivated by the well-known biases associated with candidate gene approaches. It also

reflects the conviction that better understanding of heterogeneous behavioral phenotypes

such as OUD requires analyzing the full genome.

Based on studies identified through this screening, a qualitative synthesis was first

performed to summarize experimental designs and findings. Second, a focused quantitative

analysis of bulk tissue transcriptomic studies was conducted, with the goal of defining a core

signature of opioids. To our knowledge, previous reviews on opioids and OUD [6,7] were

completed more than 15 years ago, and such a systematic synthesis across brain structures

and technologies had never been performed. Our results indicate that, although more than 40

genome-wide studies have been published, available evidence for reproducible findings is

surprisingly limited. Third, we reviewed most recent work that leveraged cell-type specific and

single-cell approaches, or that interrogated epigenetic regulatory mechanisms contributing to

opioid plasticity. Finally, we discuss current challenges in the field, as well as avenues for

future work.

Methods

Full methodology and bioinformatic code are available as Supplementary Material and File.

Results

The search provided 2705 potentially eligible studies (Fig.1). Once duplicates were removed,

a total of 1225 were screened. 1067 articles were excluded because their title or abstract didn't

meet eligibility criteria, with an additional 89 articles excluded after full text evaluation. As a

result, 69 papers were selected for qualitative synthesis, covering both transcriptomic and

epigenomic approaches. Among these, 44 articles that performed transcriptomic analyses in

the 5 most frequently investigated regions were selected for more detailed quantitative

synthesis: of these, 34 articles investigated a single brain region, 9 covered 2 regions, and

only 1 characterized 5 regions [8]. Overall, this resulted in 52 differential expression analyses: 18, 11, 11, 6 and 6 in the nucleus accumbens (NAc), frontal cortex, whole striatum, dorsal striatum and spinal cord, respectively.

1. Qualitative analysis (n=69 studies)

Species and sex ratio. Most studies were conducted in rodents (Fig.2A): 30 (43%) in rats, 28 (41%) in mice, and 1 in both species. Only a small number investigated humans (n=8, 12%) and macaques (n=2, 3%), reflecting well-known practical limitations. As consequence, current knowledge on conserved or species-specific opioids effects is limited.

Regarding sex, epidemiological surveys indicate that OUD prevalence is higher in males than in females [3, 9] (although smaller differences have been reported with DSM-5 criteria [10]), with a number of sex-specific risk factors (e.g., ethnicity, income). This clinical situation has long been modeled in rodents, as sex differences in behavioral effects of opioids [11] have been described for analgesia [12], tolerance [13], withdrawal symptoms [14], reward [15] or motivation [16]. In comparison, molecular mechanisms underlying these differences are poorly characterized. The majority of OUD studies reviewed here investigated males only (n=48, 70%; Fig.2B-D), with few analyses conducted in females (n=5, 7%), and only 14 studies (20%) that included both sexes (7 in rodents, 7 in humans). These numbers are consistent with the current landscape in neuroscience research, in which the ratio of articles reporting on males only, as opposed to females only, is around 5 to 1 [17]. We nevertheless note an encouraging trend, as the large majority of studies (8/10) published in 2021 investigated both sexes. Several factors likely underlie this recent evolution, including the recognition that females do not display more experimental variability than males [18], and evidence that pathophysiological mechanisms may differ across sexes, as shown for disorders such as depression [19], schizophrenia [20] or alcohol use disorder [21].

Specifically in rodents (Fig.2C), more than three quarters of studies investigated males only (n=47, 80%). Seven articles investigated both sexes (n=7, 12%), using surprisingly low sample sizes (n=1-4 in male/female, and opioid-treated/control, groups; Fig.2D). Among

these, 2 studies treated sex as a covariate [22,23], and 3 pooled both sexes for the analysis

of opioid effects [24–26], therefore providing no direct description of male/female differences.

This is concerning, considering that the 2 remaining studies found little overlap, from 5 to 35%

(see [27] and [28]), among morphine-induced changes occurring in each sex. Therefore,

available evidence points toward a substantial degree of sex-specificity in opioid-induced

transcriptomic plasticity, which will have to be confirmed in future work. Even less is known in

human (Fig.2E). Among 8 articles, 7 investigated males and females, while only 1 investigated

males only. However, these 7 studies showed a strong bias towards examination of men

(Fig.2F), with only one that used equal sample sizes in both sexes [29]. In the later, sex was

treated as a covariate, with no direct description of its impact. Altogether, this underscores a

significant gap in the opioid literature, and calls for additional work on sex as a biological

variable. The hope is that better understanding of divergent pathophysiological routes in males

and females may unravel sex-specific therapeutic strategies.

Route and duration of opioid administration. Among studies included in qualitative analysis

(n=69), the majority used chronic administration (n=55, 80%; see Fig.3A), mostly via self-

administration (SA, n=19, 34%, among which 8 corresponded to human studies),) and

intraperitoneal injection (ip; n=18, 33%), followed by subcutaneous injection (n=8, 15%). Other

procedures were less frequently used: continuous administration (with pellets or mini-pumps;

n=5, 9%), intrathecal administration (n=3, 5%), or gavage (n=2, 4%).

There is limited information regarding differences among administration procedures.

Interestingly, Lefevre et al [30] recently compared 3 conditions: continuous and systemic

morphine infusion using osmotic mini-pumps, with ("Interrupted") or without ("Continuous") the

induction, twice daily, of a precipitated withdrawal episode (using naloxone, a non-specific

opioid antagonist), compared to Controls (saline minipumps). Results indicated 687 and 407

differentially expressed genes (DEG) when comparing Interrupted and Control groups in the

NAc and dorsal striatum, respectively, with only 1 DEG between Continuous and Control

groups. Therefore, rather than opioid exposure per se, it is the repetition of intermittent

withdrawal episodes that may be driving the intensity of transcriptomic adaptations affecting

the mesolimbic system (whether precipitated, as in this study, or spontaneous, in-between

bolus injections). This is consistent with the clinical notion that the alternation of intoxication

and withdrawal is critical for the development of OUD.

Opioid effects have also been investigated using acute administration (10 studies,

14%), corresponding mostly to ip injection (9 articles), with only 1 using subcutaneous

injection. While these studies were mainly interested in opioids used as analgesics or

anesthetics [31,32], or sought to compare acute effects of various psychoactive drugs [33],

available data supports the notion that they may be relevant for the understanding of OUD.

Indeed, systematic comparisons across inbred mouse strains have shown that variability in

the intensity of opioid physical dependence is strikingly similar when withdrawal is precipitated

after acute or chronic injections [34,35]. In other words, there are shared genetic determinants

for brain adaptations triggered by acute or chronic exposure. A recent single-cell RNA-

sequencing (scRNA-Seq) study [36] provides further evidence that these genetic determinants

translate into at least partially similar transcriptional plasticity during acute or chronic opioid

exposure (see *3. High cellular resolution approaches*).

Another important consideration is that most studies reviewed here (51/69, 73%) used

passive modes of administration, which do not model the volitional or compulsive drug-taking

aspects of OUD. This reflects the technical difficulty of intravenous SA experiments (IVSA),

particularly in the mouse. Interestingly, new paradigms now facilitate such studies, including

devices for opioid vapor [37] SA, or the use of optogenetics for light-induced self-stimulation

[38]. Few comparisons of passive and active administrations have been conducted. Among

these, Tapocik et al [39] compared "yoker" mice (IVSA group) with groups of "yoked-morphine"

and "yoked-saline" animals which, each time the yoker self-administered morphine, received

passive IV morphine or saline injections. A much higher number of DEG were identified

between the yoker and yoked-saline groups (n=1177) than between yoked-morphine and

yoked-saline mice (n=244), with only limited overlap between the 2. This suggests that

voluntary drug-taking may drive or reflect transcriptional adaptations that are only partially

recapitulated by non-contingent administration. Additional studies will be required to deepen

this understanding.

Nervous system structures. Among articles eligible for qualitative analysis, 18 regions of the

nervous system were explored (Fig.2B). Not surprisingly, a large majority focused on the

mesocorticolimbic dopaminergic pathway: the NAc (21 articles [8,22,26,28–30,32,40–53]),

frontal cortex (13 articles [8,25,27,29,51,54–61]), whole/unspecified striatal complex (11

articles [24,33,62–70]), dorsal striatum (7 articles [8,30,48–50,71,72]) and ventral tegmental

area (VTA, 5 articles [28,51,73–75]). Other regions included the spinal cord (7 articles [62,76–

81]); hippocampus (5 articles [8,63,82–84]); amygdala (3 articles [8,85,86]); locus coeruleus

[74, 87], ventral midbrain [42, 88], hypothalamus [89, 90], whole brain with [91] or without [92]

cerebellum (1 each); periaqueductal gray matter [93], pituitary gland [90], arcuate nucleus [94],

nucleus of the tractus solitarius [31], brainstem [23], cerebellum [84] or dorsal root ganglia [95]

(DRG, 1 each). Below, we briefly describe the rationale for studying such diverse structures.

The course of OUD involves 3 stages: intoxication, withdrawal/negative affect, and

anticipation of next intoxication. These stages reflect gradual adaptation of the brain to drug

exposure, and rely on distinct mechanisms. The intoxication stage, mostly driven by acute

reward, notably associates with dopamine release, in the striatum and frontal cortex, from

neurons located in the VTA. The second stage corresponds to negative reinforcement,

whereby drug consumption alleviates physical and negative affective states of withdrawal,

involving the locus coeruleus [96], NAc [97], and amygdala [98], among others. Finally, the

anticipation stage results from memories of drug-associated cues and contexts

(hippocampus), as well as impaired goal-directed behaviors, implicating higher-order

structures (e.g. prefrontal and orbital cortices [99]). These relationships are not exclusive, with

some structures implicated at multiple stages of the disease (e.g., the striatum, also involved

in habit formation through its dorsal part). Remaining regions (spinal cord, DRG, brainstem,

periaqueductal gray matter, PAG) were investigated in relation to opioid analgesia, tolerance

and hyperalgesia. Finally, few articles were interested in how opioids modulate

neuroendocrine systems (hypothalamus, arcuate nucleus, pituitary gland), with implications

for their interactions with the stress or reward systems.

In addition to these associations, it is also important to consider the distribution of the

endogenous opioid system. This system is composed of 3 opioid receptors (mu, delta and

kappa; MOR, DOR and KOR), among which the MOR is central and necessary for opioid-

induced analgesia, reward, and physical dependence [100]. This receptor, and neurons by

which it is expressed, therefore represent pharmacological entry points for brain adaptations

to opioids. Comparison of studies reviewed here with MOR distribution in the brain (as

assessed using ligand autoradiography [101–106]; Fig.3C) indicates that: i) MOR is highly

expressed among structures most frequently studied, as expected; ii) surprisingly, other areas

with similarly high expression have received little to no attention (NTS, locus coeruleus, dorsal

raphe nucleus, medial habenula, thalamus), calling for more work.

Techniques. Thirty-seven and 29 studies used microarrays and RNA-seq, respectively, to

analyze the transcriptome (Fig.3D). While the first RNA-seq study was published in 2014[12],

this technique has since progressively replaced microarrays, reflecting its well-recognized

advantages (higher sensitivity and throughput, potential for gene discovery). For epigenomic

profiling, 3 studies recently performed bisulfite sequencing [51, 59, 61], and only 1 used

ATACseq (Assay for Transposase-Accessible Chromatin with sequencing) [72]. Considering

rapid improvements in throughput and resolution of sequencing methodologies (see below),

these trends are expected to strengthen.

2. Quantitative analysis of transcriptomics studies

To identify reproducible transcriptomic opioid signatures, we ran quantitative analyses of bulk-

tissue results available for the 5 most studied structures (corresponding, after removal of

epigenomic studies, to 44 articles; Fig.4A). To do so, gene-level results were retrieved from

each study, when available. This yielded only 24 lists of DEG comparing opioid-treated

animals and control groups, originating from 17 articles. Among these, 6 articles did not report

any p-values, while the remaining 11 used very diverse thresholds on p-values and fold

changes (FC; details in Supplementary Table 1). Surprisingly, a small minority (3/44, 6.8%

[24,28,29]) provided full results with p-values and FC for all genes. In addition, raw data were

identified as available in the public repository GEO in less than half of the cases (8/17, 47%).

Overall, this underscores strong heterogeneity in analytical procedures and data reporting

which, as discussed elsewhere [107], prevents re-analysis and limits creation of new

knowledge. To counter this, efforts have been made to harmonize omics practices [108], with

policies enforced by funding agencies [109]. For the present review, a genome-wide meta-

analysis combining, for every gene, effect sizes and sample sizes across studies was not

possible. We nevertheless: i) Systematically assessed concordance among available lists of

DEG; ii) Conducted a threshold-free comparison of the 2 NAc studies for which genome-wide

results were available.

2.1 Nucleus accumbens (18 studies)

Among 18 articles, only 8 provided lists of DEGs: 4 in mice (in males only [30,32,40,43]), 2 in

rats (1 in males only [45], 1 in both sexes [28]), 2 in humans (both sexes [29,47]). The 8 articles

provided a total number of 3502 DEG (Fig.4B-E). Among these, the vast majority were

identified by 1 study only (n=3017/3502, 86%), while 424, 57 and 4 DEG were identified across

2, 3 or 4 studies, respectively. No single genes were common to 5 or more articles. In addition,

the best overlap was found among the 2 DEG lists reported by Imperio et al [45] (Fig.3E).

Overall, this indicates poor reproducibility in the NAc.

To try to define which factors contribute to this heterogeneity, we intersected DEG lists

in each species separately (Fig.3C-E). This, however, did not improve concordance: in

humans, only 54 genes were common to both studies (Jaccard Index, JI=2.4%, Fig.4C; see

Supplementary Table 2 for the full matrix of pair-wise JI across all studies); in mice, 7 genes

were identified by at least 2 studies (JI=1%, Fig. 4D); in rats, the JI increased to 15.5%

(Fig.4E), mostly driven by high overlap among the 2 aforementioned DEG lists from Imperio

et al. In the later work, the authors identified large- (LS) and small-suppressors (SS): in LS rats, greater avoidance of a natural reward (saccharine) that predicted heroin availability was associated, compared to SS rats, with worse outcomes during later heroin IVSA (increased consumption, higher motivation and relapse). Gene expression changes occurring in the LS group (compared to control rats given access to saccharine but not heroin; n=597 genes), and in the SS group (same controls, n=205) showed a JI of 25.5% (with striking concordance in up- and down-regulation). This was much higher than overlaps observed between Imperio et al and the other rat study by Townsend et al, which yielded JI of 1.35% (Imperio-SS/Townsend) and 3.43% (Imperio-LS/Townsend). Overall, this indicates that differences across laboratories, animal facilities or experimental designs lead to poor concordance across studies.

2.2 Dorsal striatum (6 studies)

Only 2 dorsal striatum studies provided lists of DEG (Fig.3F). One study used chronic morphine administration interrupted by naloxone injections in male mice [30], as described above, while the other used acute morphine injection in male rats [71]. Reflecting these differences, a very small overlap was found between the 2 (0.62%). Therefore, while the dorsal striatum is thought to gradually gain control over drug-taking behaviors in OUD [110], underlying transcriptomic plasticity is poorly characterized.

2.3 Whole striatum (11 studies)

Six of the 11 whole striatum studies provided lists of DEG [24,33,64–67] (Fig.5B): 2 investigated chronic heroin [67] or morphine [24] treatment, while 2 compared acute effects of distinct opioids (morphine and heroin, with lists of DEG for each) [33,65], and 2 compared acute and chronic morphine exposure [64,66]. Most DEG were identified by 1 study only (3308/3736, 88.5%), with only 112 genes common to 3 or more studies. Strikingly, consistent with NAc results, strongest overlaps were observed for lists of DEGs coming from the same article, with for example 54 (JI=17%, Korostynski 2013 [33]) and 16 (21%, Piechota 2010 [65])

DEG common to acute morphine and heroin injections, respectively, and 139 DEG (31%, Korostynski 2007 [66]; Fig.5A) common to acute and chronic morphine administration. In sharp contrast, comparisons across, rather than within, studies yielded poor overlaps, even when considering those that used similar drug administration: among chronic morphine studies (Fig.S1A), the JI varied between 0.1% (Coffey 2020 versus Skupio 2017), 1.2% (Korostynski 2007/Skupio 2017) or 2.6% (Korostynski 2007/Skupio 2017), and, among acute morphine studies, from 2.7% to 7.8% (Fig.S1B; Piechota 2010/Skupio 2017, JI=2.7%; Piechota 2010/Korostynski 2007, 3.2%; Piechota 2010/Korostynski 2013, 6.4%; Korostynski 2013/Skupio 2017, 6.6%; Korostynski 2013/Korostynski 2007, 7.3%; Korostynski 2007/Skupio 2017, 7.8%), with only one comparison standing out among acute heroin studies (30 common DEG, JI=19%, Piechota 2010/Korostinsksy 2013). Importantly, because 5 of these 6 studies were conducted by the same group, it is possible to speculate that this variability may reflect batch effects among experiments conducted in the same facilities but at different time points, rather than variations across different opioids or administration parameters (see also section 2.6 below).

2.4 Frontal cortex (11 studies)

Seven studies provided DEG lists (Fig.5B): 2 in humans [29, 58], 5 in rats [25,27,55–57]. Data from Odegaard et al [25] were discarded, as they corresponded to transgenerational effects in animals not directly exposed to opioids (administered to dams during pregnancy, or postnatally). For 2 additional studies [55,57], we could not map microarray probes to genes. Among the 4 remaining articles, little overlaps were again found (Kuntz-Melcavage/Mendez, JI=0.5%; Kuntz-Melcavage/Seney, 1.1%; Liu/Mendez, 1.4%; Liu/Seney, 1.6%), with no improvement when conducting comparisons within each species (Liu/Kuntz-Melcavage in rats, JI=0%; Seney/Mendez in humans, 4.1%).

2.5 Spinal cord (6 studies)

Among the 6 spinal cord studies, 5 did not provide DEG [76–80], while the 6[th] one used a

homemade microarray[62], for which correspondence between probe IDs and genes was only

partially provided. As such, no comparison was possible.

2.6 Sources of variability

Considering that pharmacological recruitment of MOR and resulting intra-cellular signaling

may trigger a set of common transcriptional adaptations across the 5 regions of interest, we

next pooled all lists of DEGs (Fig.5D). Again, the vast majority of DEG (70.1%, 4940/7047)

were unique to a single study, while only 21.4% common to 2 studies (1509), 5.9% to 3 studies

(418), and 1.7% to 4 or more studies (117; full lists in Supplementary Table 3). Interestingly,

Cdkn1a (Cyclin Dependent Kinase Inhibitor 1A) was identified by 11 studies. This gene

encodes for p21, a protein involved in the regulation of the cell-cycle and oligodendrocytes

[111]. Interestingly, Avey et al [40] found Cdkn1a to be one of the most significantly

upregulated genes in response to acute morphine. This upregulation occurred in

oligodendrocytes (confirmed by FISH), and was blocked by the opioid antagonist naltrexone,

indicating a MOR-dependent mechanism. Therefore, Cdnk1a may contribute to opioid effects

across multiple brain sites, possibly through the regulation of the oligodendrocyte lineage. Of

note, cocaine, another major drug of abuse, similarly recruits p21, as cocaine-induced

behavioral responses were modified in p21 knockout mice [112].

We also performed Gene Ontology (GO) analysis using DEG common to at least 3, 4

or 5 studies (Supplementary Table 4). No results were significant for the small list of 63 DEG

common to 5 or more studies. When considering the 180 DEGs common to at least 4 studies,

only 3 unspecific GO terms were enriched: cell periphery, plasma membrane and hormone

activity. More consistent results emerged for the 598 DEG common to at least 3 studies, with

46 significant GO terms mostly related to synaptic signaling and cell-cell communication, likely

reflecting opioid-induced electrophysiological and synaptic plasticity across brain structures.

Because weak DEG overlaps were identified within each individual region (even when

grouping studies based on species, Fig.4C-E, or treatment chronicity, Fig.S1), we sought to

more systematically assess, across the 5 regions, the impact of following factors: species,

brain structure, treatment chronicity, drug, and article (Fig.5D). For each factor (e.g., species),

pair-wise JI were classified in 2 categories depending on whether they belonged to the same

sub-group within that factor (e.g., 2 studies conducted in humans; "intra"), or not (e.g., 1 study

in humans compared to another in rats; "inter"). As expected, grouping studies by species

(p=6.0E-4) or brain structures (2.3E-4) increased concordance, with higher JI. This is

consistent with the notion that, although they remain poorly characterized, these 2 variables

significantly contribute to variability in opioid-induced transcriptional plasticity. Treatment

chronicity also had a significant, albeit milder, impact (p=0.012). In contrast, the type of opioid

used had none (p=0.43), possibly because, despite pharmacokinetic or dynamic differences,

all drugs considered act as agonists at the same molecular target, MOR. Importantly,

concordance increased most when lists of DEG came from the same publication (p=0.011),

despite the fact that only 7 comparisons fell under this category. As already mentioned, this

suggests that differences across laboratories explain a substantial part of the lack of

reproducibility across studies.

Finally, we refined our analysis by considering the directionality (up/down) of DEG,

focusing on genes identified by at least 5 studies. Among the 17/24 studies that reported on

directionality (Fig.S2), most genes showed discordant results (37/63, 59%): for example,

Cables1 was upregulated in Imperio et al and Coffey et al, but downregulated in Seney et al

in both the frontal cortex and NAc. In contrast, only 18/63 (29%) DEGs showed concordant

directions, with 17 up-regulated genes, and 1 down-regulated. For the remaining 8 genes

(13%), directionality was reported by none or only 1 study. Altogether, these results further

highlight low reproducibility.

2.7 Threshold-free genome-wide comparison

Next, we compared the 2 NAc studies (Seney et al [29], Townsend et al [28]) for which full

genome-wide results were available. While both included males and females, only Townsend

et al reported sex-specific results. Therefore, we reprocessed Seney et al raw data to identify

DEG in each sex separately (see *Supplementary Material*). This slightly increased

concordance: while JI were equal to 7.5 and 4% when comparing initial Seney data

(male/females pooled) with Townsend data in females or males, respectively (Fig.S3A-B),

they increased to 9.5% and 4.5% when considering sex-specific Seney results (Fig.6A-B).

These results therefore argue for the importance of accounting for sex when identifying opioid-

induced transcriptional changes.

We then used RRHO2 (Rank–Rank Hypergeometric Overlap [113]) to conduct

genome-wide comparison of the 2 studies. RRHO2 iteratively performs hypergeometric testing

for all possible combinations of thresholds applied to each dataset, generating a so-called

"threshold-free" analysis. Because this approach takes directionality into account, we also

computed, for comparison, selective overlaps among either up- or down-regulated DEG only

(using p-value thresholds from each study, and represented as white crosses in RRHO2

matrices; Fig.6C-D). Compared to those, much more significant overlaps were identified using

RRHO2 when less stringent p-value thresholds were considered within each dataset (best

hypergeometric results are shown as black crosses). These increased overlaps, however,

came at the expense of the identification of large lists of genes which, although commonly

dysregulated in similar direction, individually exhibited milder FC. Of note, these overlaps were

detected despite the fact that Seney investigated human post mortem tissue from individuals

with OUD, while Townsend focused on rat tissue collected after 24 hours of withdrawal from

fentanyl IVSA. In addition, different patterns of convergence were observed in each sex: in

females, overlaps were highly significant among both up- and down-regulated genes while, in

males, a lower concordance predominantly affected down-regulated genes. This suggests that

these rat datasets may differentially capture sex-specific signatures of human OUD. Overall,

these analyses demonstrate the utility of reporting full genome-wide results which, in

combination with threshold-free approaches, could generate deeper comparisons and

meaningful insight, even across species.

3. High cellular resolution approaches

Studies reviewed above indicate that opioids induce genome-wide transcriptomic changes,

with striking variability across laboratories, experimental designs, and brain regions. The later

likely reflects a combination of direct and indirect effects: i) direct pharmacological opioid

effects correspond to the recruitment of MOR and downstream intracellular signaling. In case

of neurons, this classically results in decreased excitability and activity [114]; ii) then,

depending on the chemical identity and connectivity pattern of MOR-expressing cells, their

inhibition leads to changes in the release of multiple neuromodulators and transmitters,

triggering indirect adaptations [102]. Accordingly, transcriptomic changes described above

reflect the sum of these events. This raises 2 important questions: how is each cell type, or

even each brain cell, affected by these changes? And what are underlying molecular

regulatory principles?

To address the first question, several groups recently used cell-type specific and

single-cell methodologies. Coffey et al [115] focused on microglial cells of the whole striatum,

using translational ribosome affinity purification (TRAP, to capture their ribosomes and

"translating transcriptome" [116,117]). Chronic morphine effects were analyzed with, or

without, the induction of naloxone-precipitated withdrawal. Results showed that both chronic

morphine and withdrawal triggered numerous gene expression changes, which mostly

affected GO terms related to cAMP signaling, regulation of synapses, and the cytoskeleton. A

striking anticorrelation was observed between morphine- and withdrawal-induced changes,

providing an interesting transcriptomic correlate, at the level of microglia, of the

pharmacological antagonism between morphine and naloxone. Surprisingly, to our

knowledge, no study has used such approaches to interrogate opioid effects specifically in

neuronal cells, let alone MOR-expressing neurons which, arguably, may exhibit particular

sensitivity to opioids. This is feasible, as several knock-in mouse lines are now available, in

which the Cre enzyme is selectively expressed by MOR-expressing cells [118–120].

Achieving even higher resolution, Avey and colleagues applied single-cell RNA-seq

(scRNA-Seq, using Drop-seq [121]) to analyze transcriptomic changes occurring, in more than

23,000 cells of the mouse NAc, 4 hours after an acute morphine injection (20 mg/kg, ip) [40].

As is typical in scRNA-Seq, the sequencing depth achieved yielded expression data for around

1,600 genes, corresponding to a reduced representation of the genome. Acute morphine

induced gene expression changes that were widespread and, interestingly, most frequent in

oligodendrocytes, followed by neurons, astrocytes, and microglia. This therefore uncovers a

hierarchy in transcriptional sensitivity to opioids among NAc cell-types, and extend on the

previous investigation of microglia[24]. More recently, Reiner et al [36] used single-nucleus

RNA-Seq (snRNA-seq, 10x Genomics 3' gene expression assay) to analyze a larger number

of nuclei (190,030) and genes (>2000) in the rat NAc, also following acute morphine injection

(10 mg/kg, ip). DEG mostly affected neurons and oligodendrocytes, followed by astrocytes

and microglia, reminiscent of the former study. Effects of morphine were also analyzed after

10 days of IVSA and compared to acute ip effects. Among DEG shared across the 2 conditions

(n=431), the vast majority were dysregulated in a similar direction (up/down), suggesting that

they may reflect morphine exposure rather than volitional drug-taking (an additional "yoked"

group, chronically exposed to morphine, would help substantiate this distinction). Importantly,

an even larger number of DEG were specifically dysregulated in only one of the 2 conditions

(n=868, with more DEG associated with IVSA), with distinct distributions among NAc cell types

(e.g., sub-populations of Dopamine receptor D1- or D2-expressing neurons). By providing the

first single-cell description of morphine IVSA, these results reinforce the notion that voluntary

drug-intake may recruit molecular mechanisms that are not recapitulated by non-contingent

administration.

We next directly confronted results from these 2 single-cell studies regarding acute

morphine exposure. Both used Seurat [122] and dimensionality reduction (t-SNE [40] or UMAP

[36]) to identify clusters, which were then assigned to cell-types using external data. We first

compared marker genes reported to define those clusters, when available (Avey: 11/29

clusters; Reiner: 11/19 clusters). Low agreement was found, with JI ranging from 0 up to 7.1%

(for activated medium spiny neurons, MSNs, in Avey et al and D1R MSN-1 in Reiner et al,

Fig.S4A), possibly reflecting the fact that both studies reported low numbers of marker genes

(Avey: 39 to 51 marker genes; Reiner: 4 to 12). We then confronted morphine-induced DEG

for each cell-type (Fig.6E). Low JI values were found, with maximal overlap (6.4%) obtained

when comparing DEG affecting "myelin-forming and mature oligodendrocytes" (MO_all) in

Avey et al to "Oligodendrocyte-3" in Reiner et al. In addition, surprisingly, while a group of cell

clusters assigned to similar cell-types in the 2 studies showed some similarity in DEG (shown

as green squares, Fig.6E), strong concordance was also observed across cell clusters with

different identities (e.g. "Astrocyte-2" in Reiner et al compared to MO_all in Avey et al; yellow

squares). Also, overlaps among DEG (Fig.6E) did not increase when considering more similar

cell-types (Fig.S4B), as would be expected. Therefore, these analyses indicate: i) significant

discrepancies across the 2 studies (Fig.S4B), possibly reflecting methodological factors not

yet standardized in single-cell approaches, and ii) that, reminiscent of issues raised at bulk

tissue level, a more comprehensive reporting of marker genes and differential expression

results should be enforced.

4. Non-coding and epigenomic mechanisms of opioid plasticity

In parallel to efforts to deepen cellular resolution of transcriptomic analyses, others have

sought to decipher underlying non-coding RNA and epigenomic modes of regulation [123].

This is motivated by the recognition that, although neurons are post-mitotic cells, they exhibit

peculiar plasticity features that may contribute to the encoding of life events, including

exposure to drugs of abuse [124].

Several non-coding RNA species have been investigated. The first report used

custom-made arrays to characterize an unknown number of micro-RNA (miR), and reported

bidirectional expression changes affecting 7 of them in rat primary neuronal cultures or the

mouse hippocampus, following morphine or heroin exposure [84]. Among these, the miR-339-

3p was later shown to target the 3'-UTR and down-regulate MOR in response to opioid

treatment [125]. A second study used more recent arrays to interrogate all mouse miR in the

NAc, following chronic heroin injections [52]. 111 miR were found to exhibit >25% expression

changes (although no statistical significance was reported). This study then focused on miR-

218, and showed that its over-expression could inhibit heroin-induced reinforcement (in both

conditioned place preference and IVSA), possibly through the targeting of MeCP2 (a protein that binds methylated DNA), suggesting a potential link between opioids, miR and DNA methylation. Other studies explored long non-coding (lncRNA) or circular RNA (circRNA). One group used 2 arrays interrogating 9,000 lncRNA [78] and 15,000 circRNA [81] to characterize chronic morphine effects in the rat spinal cord: 136 lncRNA (84 up, 52 down; nominal p-value<0.05, |log2FC|>1.5) and 2038 circRNA (896 up, 1142 down; nominal p-value<0.05, FC>2) showed expression changes. In parallel, another report used RNA-Sequencing to focus on circRNA in the NAc [53], and found differences for 112 of them when comparing groups of mice receiving chronic morphine injections with or without electroacupuncture (51 up, 61 down; nominal p-value<0.05, |log2FC|>1.5). These results remain descriptive, and a large variety of mechanisms have been proposed to explain how non-coding RNA may exert biological functions (including the sequestering of miR by circRNA, or complementary interactions among lncRNA and mRNA). Understanding how these mechanisms are recruited by opioids will require further investigation.

Regarding the epigenome, 2 recent studies (1 in the rat [51], 2 in humans [126,127]) investigated DNA methylation, a mark increasingly implicated in psychiatric pathophysiology [128]. The rat study used Whole Genome Bisulfite Sequencing (WGBS) to analyze DNA methylation changes during heroin IVSA, in a standard or enriched environment, in 3 brain regions (VTA, NAc and medial prefrontal cortex) [51]. Because of low sequencing depth, DNA methylation levels were averaged for selected gene features across the whole genome (promoters, CG islands). As expected, neither IVSA nor environmental enrichment had any impact on these global metrics, suggesting that the methylome is not massively reprogrammed. It remains possible, however, that opioids may trigger DNA methylation changes at more localized genomic sites, a hypothesis supported by: i) the intuition that large transcriptomic changes associated with opioid exposure or OUD reviewed above are unlikely to occur in the absence of any chromatin plasticity; ii) candidate gene studies which, although outside the scope of the present review, provide evidence for differential DNA methylation at specific loci (the MOR gene in human blood [129–131], and a few loci in the rat brain [132]);

iii) and, finally, the 2 aforementioned human studies. Focusing on Broadmann area 9, the first used EPIC arrays to assess DNA methylation in a post-mortem cohort of OUD patients (n=19) and healthy controls (n=11) [127]. Among the >800,000 CG sites investigated (where DNA methylation mostly occurs in the mammalian brain), 11,917 sites located at transcription start sites (TSS) showed nominally significant difference (p-value<0.05), with bidirectional changes. While genomic localization of these probes or full genome-wide results were not described, these results nevertheless support the existence of a form of methylomic plasticity in OUD.

Human studies face the difficult task of accounting for confounding factors. Variations in genotype, age, sex, or socio-demographic factors, for example, have all been associated with DNA methylation changes. Also, because epigenetic landscapes determine cellular identity, differences observed in bulk tissue may reflect variations in cellular composition (which may contribute to pathogenesis), rather than transcriptional regulation. The second human study started to address these questions [126]. Using EPIC arrays to analyze bulk tissue, the authors adjusted for socio-demographic characteristics, ancestry and cellular composition. They investigated a large cohort (n=153), focusing on death from acute opioid overdose rather than OUD. Although no CG site survived multiple testing correction, 13 passed a relaxed significance threshold (p<1.0E-05), one of which was located in Netrin-1, a gene implicated in KOR activity. At genome-wide level, an association between opioid use and accelerated epigenetic aging was reported, along with enriched differential methylation in GO and KEGG pathways related to substance use. Overall, these studies support the notion that the brain DNA methylome may be modulated by opioid exposure. Additional studies will be necessary to better understand these effects, as well as potential roles for non-canonical, brain-specific forms of DNA methylation (methylation in non-CG context [133], hydroxymethylation at CG sites [134]).

Beyond covalent modifications of the DNA, epigenomic plasticity also relies on histone modifications and changes in compaction or 3D structure of chromatin. Surprisingly, no data characterizing histone changes in relation to opioid exposure were identified during our systematic screening. Studies available on cocaine [134,135] nevertheless suggest potentially

broad adaptations. Interestingly, a recent study compared chromatin accessibility in a cohort

of 10 heroin abusers and 10 healthy individuals [72], in the NAc and putamen. ATAC-Seq was

used to identify genomic regions where chromatin was "opened" and DNA accessible for

tagmentation (i.e., tagging with sequencing adapters by a mutated, hyperactive transposase).

The technique was applied to neuronal and non-neuronal cells, isolated using NeuN staining

and Fluorescence-Activated Nuclear Sorting (FANS). Heroin use associated with an

enrichment of ATAC-Seq peaks in specific gene features (CG islands, promoters, 5'

untranslated regions) and annotated regulatory regions (the histone mark H3K4me1, or

binding sites for the EZH2 transcription factor), as well as more frequent changes in neurons

than in non-neuronal cells (affecting GO terms related to the synapse, among others). Using

in vitro reporter assays, primary cultures and behavioral rat experiments, the authors then

showed that FYN (a member of the Src tyrosine kinase family that exhibited strong heroin-

induced changes in chromatin accessibility in human), also significantly contributed to heroin

IVSA in the rat. Overall, this work nicely illustrates how epigenetic analyses may uncover

chromatin regulatory mechanisms contributing to OUD.

Discussion

In the present review, we systematically assessed current knowledge on functional genomic

mechanisms of opioids and OUD, and uncovered a surprisingly small overlap in genes for

which transcriptomic changes were consistently identified across studies. As a result, a lot

remains to be done to move the field forward, and we discuss below most promising

perspectives to refine behavioral modelling, improve cellular resolution, and use more diverse

sequencing technologies and advanced data integration.

While the complexity of OUD is difficult to recapitulate in animal models, their face

validity has nevertheless increased over last decades, and it is admitted that addiction-related

phenotypes (e.g., uncontrolled consumption, resistance to punishment, increased motivation)

can be induced in rodents [136]. We now face the task of deciphering the molecular

mechanisms that may account for each of these dimensions, including for example the

susceptibility to develop compulsive drug intake. The later susceptibility aspect has been

addressed by only one article covered in the present review (Imperio et al [45]), which

contrasts with the amount of data available for other psychiatric disorders, in particular

depression. Among other examples [137,138], Bagot et al [139] examined transcriptomic

adaptations occurring in mice categorized as susceptible or resilient to chronic social defeat.

Using network analysis, 2 hub genes (Dkkl1, Neurod2) were identified, whose overexpression

was sufficient to potentiate social stress effects, thus unravelling molecular substrates of

susceptibility. At epigenetic level, regulatory mechanisms implicating long non-coding RNAs

may contribute to mediate such susceptibility [140]. In the addiction field, little is known about

molecular substrates of susceptibility. Two reports applied high-throughput methodologies to

describe such inter-individual differences in relation to DNA methylation and alcohol oral SA

in macaques [141], or to micro-RNA expression in a rat model of cocaine SA [142]. While

similar mechanisms are likely recruited in OUD, they have not yet been addressed by open-

ended approaches, pointing toward an important goal for future work.

Another avenue for improvement is cellular resolution. The brain tissue heterogeneity

has long hampered characterization of the role of each cell-type in complex phenotypes. In

complement to cell-type specific (TRAP) and single-cell sequencing (sc/snRNA-Seq), which

have already been applied to the understanding of OUD (see above), other tools are emerging

for molecular profiling of cell or neuronal sub-populations based on 2 factors that determine

their functional organization and plasticity : i) connectivity pattern, probed using antero- or

retrograde viruses, in combination with TRAP [143] (transcriptomics) or FANS (epigenomics

[144]); or ii) spatial context, with methodologies based either on sequencing or microscopy to

detect mRNA in space. Among the former, spatial capture followed by transcriptome

sequencing (e.g. Slide-seq [145]) enables genome-wide interrogation with 10µm resolution;

among the later, single-molecule fluorescence in situ hybridization (smFISH) allows for better

spatial resolution (1µm), but requires probe design and target gene selection [146,147]. These

spatial transcriptomic technologies have the drawback of interrogating only a fraction of the

genome, with a coverage of around a few hundred to a few thousand genes, lower than current

single-cell sequencing [148]. Future work using these techniques, by characterizing molecular

effects of opioids in a particular neuronal pathway, or with exquisite spatial resolution, shall

deepen the understanding of OUD.

A third aspect relates to gaps in the use of available technologies (ChIP-Seq, Cut&Tag,

ATAC-Seq) and their integration in multi-omic models of pathophysiology. Accordingly,

several epigenetic layers of regulation have been left unaddressed so far in relation to opioids.

This includes chromatin conformation and histone modifications, which have not yet been

investigated at genome-wide scale, in contrast with cocaine [72,149]. Because cocaine

studies uncovered widespread histone post-translational modifications, it is likely that opioids,

notably through hyperdopaminergic activity, may recruit similar plasticity. Along this line, we

also speculate that opioid effects on DNA methylation, histone modifications, and other

epigenetic substrates (lncRNA, chromatin conformation, etc) may show reciprocal molecular

interactions, with possibly locus- and cell-type specific mechanisms that remain unknown. As

illustration, we recently conducted a post mortem human study indicating that early-life

adversity and depression associate with 2 distinct forms of DNA methylation, in the canonical

CG or in the brain-enriched CAC contexts [133]. By combining WGBS with 6 types of ChIP-

Seq analysis, the 2 forms of plasticity were found to affect distinct genomic features and sites,

and defined by divergent histones modifications and chromatin states. Therefore, combined

investigation of multiple epigenetic marks, possibly at the single-cell scale, have the potential

to suggest molecular interactions that can then be dissected and manipulated *in vivo*.

Importantly, such multi-omic assessment requires implementing dedicated bioinformatic

frameworks, which are only emerging in psychiatry [150]. These integrative approaches, by

describing how multiple molecular layers of regulation interact, are expected to better capture

the chain of events linking disease causes (whether genetic, developmental, environmental,

or a combination thereof) to their behavioral manifestation [151].

In conclusion, this is to our knowledge the first systematic review of studies of

functional genomic mechanisms of opioids and OUD. Although numerous molecular

adaptations have been described across animal models and patient cohorts, replicable

findings appear limited. In the future, it is expected that leveraging multi-omic and integrative

analyses will pave the way towards the identification of innovative therapeutic strategies, and

ultimately contribute to better care.

Acknowledgements

This research was supported by the Centre National de la Recherche Scientifique (CNRS,

France), 'Université de Strasbourg', French National Research Agency (ANR-19-CE37-0010 ;

PEL), 'Fondation de France' (N° Engt: 00081244 ; PEL), 'Fondation pour la Recherche sur le

Cerveau' (FRC 2019 ; PEL), American Foundation for Suicide Prevention (AFSP YIG-1-102-

19 ; PEL), 'Fondation pour la Recherche Médicale' (FDT202204015236 ; CF), and 'Institut

pour la Recherche en Santé Publique – Institut National du Cancer' (IReSP-INCa doctoral

fellowship 2022-166 ; ACR).

Conflict of Interest

The authors declare no conflicts of interest.

Code availability

All code used in the present review is available as Supplementary File 1.

Figure legends

Figure 1. PRISMA diagram. The review protocol was registered prospectively in Prospero (record ID#CRD42022270113), and followed the Preferred Reporting Items for Systematic Review and Meta-Analysis protocol (PRISMA[152]). *Identification of studies*. Three databases were used for systematic screening: MEDLINE, Embase and Web of Science. The literature search was performed using the following keywords: "microarray" OR "RNA sequencing" OR "bisulfite sequencing" OR "chromatin immunopurification sequencing" OR "single cell RNAsequencing" AND each of the following terms: "opiates", "opioid", "morphine", "fentanyl", "oxycodone", "heroin", "methadone" or "buprenorphine". The initial search provided 2705 potentially eligible studies. *Screening*. Articles were screened based on their title and abstract, and included if they used a high-throughput, genome-wide methodology to assess modifications of gene expression or epigenetic mechanisms as a function of opioid exposure, in the nervous system of primates or rodents (rat, mouse), and were published in English in peer-reviewed journals. Exclusion criteria included tissue other than the nervous system and candidate genes approaches. Once duplicates were removed, a total of 1225 articles were screened, among which 1057 were excluded because their title or abstract didn't meet eligibility criteria. *Eligibility*. After a more thorough examination of articles' full-text, an additional 89 articles were excluded. As a result, 69 papers were selected for qualitative synthesis. Among these, 44 articles reporting on transcriptomic analyses in the 5 most frequently investigated regions were selected for a more detailed quantitative synthesis. Of these, 37 articles investigated a single brain region, 6 covered 2 regions, and only 1 characterized 5 regions [8]. Overall, this resulted in 52 differential expression analyses: 18, 11, 11, 6 and 6 in the nucleus accumbens (NAc), frontal cortex, whole striatum, dorsal striatum and spinal cord, respectively.

Figure 2. Characteristics of studies included: Species & Sex. (A) Species repartition in studies identified during systematic search. **(B)** Distribution of sex and year of publication across studies. **(C-D)** Sex distribution specifically in rodent and human studies. **(E-F)** Sample

sizes in rodent and human studies comparing opioid-exposed animals to controls (rodent,

panel E), or individuals with a diagnosis of opioid use disorder (OUD) to healthy controls.

Figure 3. Characteristics of studies included: Injection parameters, Brain structure &

Techniques. Among the 69 studies considered during qualitative analysis, a large panel of

injection parameters, nervous system structures and techniques were represented. **(A)**

Injection parameters used for opioid administration. NB: "mixed" refers to a single study that

used a combination of chronic oral administration (through bottles) and acute ip injections of

morphine[64]. **(B)** Number of articles investigating each nervous system structure. **(C)** Side-

by-side comparison of the number of articles investigating each structure with the protein

expression profile of the molecular target of opioids, the mu opioid receptor (MOR; as

assessed using [^{3}H]-DAMGO autoradiography, data in fmol/mg of tissue, adapted from[100,

103–105]). **(D)** Distribution of high-throughput methodologies and year of publication across

studies. *Abbreviations:* FCx, Frontal cortex; NAc, Nucleus accumbens; dStr, Dorsal Striatum;

Hc, Hippocapus; HT, Hypothalamus; Th, Thalamus; Amy, Amygdala; VTA, Ventral Tegmental

Area; MHb, Medial Habenula; LC, Locus Coeruleus; DRN, Dorsal Raphe Nucleus.

Figure 4. Data availability for eligible studies and quantitative analysis in the nucleus

accumbens. (A) The graph depicts the number of: i) differential expression analysis results

considered for quantitative analysis (with the corresponding number of articles between

brackets); ii) cases when lists of differentially expressed genes (DEG) were available; iii) cases

when full genome-wide results were available (with both fold changes, FC, and p-values). **(B)**

Number of DEG identified in studies that investigated the nucleus accumbens. **(C-E)** Venn

diagram of DEG identified in each species (human, mice and rat). **(F)** Venn diagram of DEG

in the 2 studies that investigated the dorsal striatum.

Figure 5. Quantitative analyses in the dorsal striatum, whole striatum and frontal cortex.

(A) Number of differentially expressed genes (DEG) identified in studies that investigated the

striatum (see Fig.S1 for separate analyses of studies that used acute or chronic opioid

administration). **(B)** Number of DEG in studies of the frontal cortex. **(C)** Number of genes

identified as differentially expressed (DE) across different studies. **(D)** Accros the 5 brain

regions considered during quantitative analysis, we assessed the impact of 5 factors (species,

brain structure, treatment chronicity, drug, and article) on consistency among studies, using

pairwise Jaccard indexes. For each factor (e.g., species), pair-wise JI were classified in 2

categories describing whether they belonged to the same sub-group (e.g., 2 studies conducted

in humans; "intra"), or not (e.g., 1 study in humans compared to another in rats; "inter"), within

that factor. NB: for the drug factor, studies in human were not considered as they cannot be

assigned to specific opioids.

Figure 6. (A-D) Threshold-free comparison of 2 bulk tissue RNA-sequencing studies

(Seney et al and Townsend et al), and cell-type specific comparison of 2 single-cell

(Avey et al) and single-nucleus (Reiner et al) RNA-sequencing studies. (A-B) Venn

diagram of differentially expressed genes (DEG) identified across the 2 bulk tissue RNA-

Sequencing studies by Seney et al and Townsend et al (DEG defined by arbitrary significance

thresholds in each study: p-value < 0.05 for both) for women/female rats (A) and men/male

rats (B). **(C-D)** Genome-wide, threshold-free comparison of the same 2 studies using Rank-

Rank Hypergeometric Overlap (RRHO2) for women/female (C) rats and men/male rats (D). In

RRHO2, briefly, genes are ranked using p-values (signed according to directionality), from

upregulated (Seney: left, Townsend: bottom) to down-regulated genes (Seney: right,

Townsend: top). Overlaps along the 2 distributions are then iteratively assessed using

hypergeometric testing. High p-values in the bottom-left and top-right quadrants indicates

concordance, whereas high p-values in the other 2 quadrants (bottom-right and top-left)

indicate discordance between the 2 datasets. The white crosses indicate the location of the

overlap corresponding to Venn diagrams depicted in panels A-B (split in subgroups of up- or

down-regulated genes), while black crosses correspond to most significant overlaps identified

using RRHO2. **(E)** Heatmap of pairwise Jaccard indexes for DEG identified across each of the

26 and 27 cell-types defined by Avey et al and Reiner et al, respectively.

References

749   1.   Rosenblum A, Marsch LA, Joseph H, Portenoy RK. Opioids and the treatment of chronic

750   pain: controversies, current status, and future directions. Exp Clin Psychopharmacol.

751   2008;16:405–416.

752   2.   Florence C, Luo F, Rice K. The economic burden of opioid use disorder and fatal opioid

753   overdose in the United States, 2017. Drug Alcohol Depend. 2021;218:108350.

754   3.   European Monitoring Centre for Drugs and Drug Addiction. European drug report

755   2017 :trends and developments. LU: Publications Office; 2017.

756   4.   Heilig M, MacKillop J, Martinez D, Rehm J, Leggio L, Vanderschuren LJMJ. Addiction

757   as a brain disease revised: why it still matters, and the need for consilience.

758   Neuropsychopharmacology. 2021;46:1715–1723.

759   5.   Maze I, Shen L, Zhang B, Garcia BA, Shao N, Mitchell A, et al. Analytical tools and

760   current challenges in the modern era of neuroepigenomics. Nat Neurosci.

761   2014;17:1476–1490.

762   6.   Przewlocki R. Opioid abuse and brain gene expression. Eur J Pharmacol.

763   2004;500:331–349.

764   7.   McClung CA. The molecular mechanisms of morphine addiction. Rev Neurosci.

765   2006;17:393–402.

766   8.   Yamada K, Nagai T, Nabeshima T. Drug dependence, synaptic plasticity, and tissue

767   plasminogen activator. J Pharmacol Sci. 2005;97:157–161.

768   9.   Substance Abuse and Mental Health Services Administration, Results from the 2013

769   National Survey on Drug Use and Health: Summary of National Findings, NSDUH

770   Series H-48, HHS Publication No. (SMA) 14-4863. Rockville, MD: Substance Abuse

771   and Mental Health Services Administration, 2014.

772   10.   Kerridge BT, Saha TD, Chou SP, Zhang H, Jung J, Ruan WJ, et al. Gender and

773   nonmedical prescription opioid use and DSM-5 nonmedical prescription opioid use

774   disorder: Results from the National Epidemiologic Survey on Alcohol and Related

775   Conditions - III. Drug Alcohol Depend. 2015;156:47–56.

11.   Bravo IM, Luster BR, Flanigan ME, Perez PJ, Cogan ES, Schmidt KT, et al. Divergent

behavioral responses in protracted opioid withdrawal in male and female C57BL/6J

mice. Eur J Neurosci. 2020;51:742–754.

12.   Craft RM. Sex differences in opioid analgesia: 'from mouse to man'. Clin J Pain.

2003;19:175–186.

13.   Craft RM, Stratmann JA, Bartok RE, Walpole TI, King SJ. Sex differences in

development of morphine tolerance and dependence in the rat. Psychopharmacology

(Berl). 1999;143:1–7.

14.   Cicero TJ, Nock B, Meyer ER. Gender-linked differences in the expression of physical

dependence in the rat. Pharmacol Biochem Behav. 2002;72:691–697.

15.   Karami M, Zarrindast MR. Morphine sex-dependently induced place conditioning in

adult Wistar rats. Eur J Pharmacol. 2008;582:78–87.

16.   Cicero TJ, Aylward SC, Meyer ER. Gender differences in the intravenous self-

administration of mu opiate agonists. Pharmacol Biochem Behav. 2003;74:541–549.

17.   Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. Neurosci

Biobehav Rev. 2011;35:565–572.

18.   Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in

neuroscience and biomedical research. Neurosci Biobehav Rev. 2014;40:1–5.

19.   Labonté B, Engmann O, Purushothaman I, Menard C, Wang J, Tan C, et al. Sex-

specific transcriptional signatures in human depression. Nat Med. 2017;23:1102–1111.

20.   Hoffman GE, Ma Y, Montgomery KS, Bendl J, Jaiswal MK, Kozlenkov A, et al. Sex

Differences in the Human Brain Transcriptome of Cases With Schizophrenia. Biol

Psychiatry. 2022;91:92–101.

21.   Hitzemann R, Bergeson SE, Berman AE, Bubier JA, Chesler EJ, Finn DA, et al. Sex

Differences in the Brain Transcriptome Related to Alcohol Effects and Alcohol Use

Disorder. Biol Psychiatry. 2022;91:43–52.

22. Vassoler FM, Oliver DJ, Wyse C, Blau A, Shtutman M, Turner JR, et al.

Transgenerational attenuation of opioid self-administration as a consequence of

adolescent morphine exposure. Neuropharmacology. 2017;113:271–280.

23. Borrelli KN, Yao EJ, Yen WW, Phadke RA, Ruan QT, Chen MM, et al. Sex Differences

in Behavioral and Brainstem Transcriptomic Neuroadaptations following Neonatal

Opioid Exposure in Outbred Mice. ENeuro. 2021;8:ENEURO.0143-21.2021.

24. Coffey KR, Lesiak AJ, Marx RG, Vo EK, Garden GA, Neumaier JF. A cAMP-Related

Gene Network in Microglia Is Inversely Regulated by Morphine Tolerance and

Withdrawal. Biol Psychiatry Glob Open Sci. 2022;2:180–189.

25. Odegaard KE, Schaal VL, Clark AR, Koul S, Sankarasubramanian J, Xia Z, et al. A

Holistic Systems Approach to Characterize the Impact of Pre- and Post-natal

Oxycodone Exposure on Neurodevelopment and Behavior. Front Cell Dev Biol.

2020;8:619199.

26. Odegaard KE, Schaal VL, Clark AR, Koul S, Gowen A, Sankarasubramani J, et al.

Characterization of the intergenerational impact of in utero and postnatal oxycodone

exposure. Transl Psychiatry. 2020;10:329.

27. Liu SX, Gades MS, Swain Y, Ramakrishnan A, Harris AC, Tran PV, et al. Repeated

morphine exposure activates synaptogenesis and other neuroplasticity-related gene

networks in the dorsomedial prefrontal cortex of male and female rats. Drug Alcohol

Depend. 2021;221:108598.

28. Townsend EA, Kim RK, Robinson HL, Marsh SA, Banks ML, Hamilton PJ. Opioid

withdrawal produces sex-specific effects on fentanyl-vs.-food choice and mesolimbic

transcription. Biol Psychiatry Glob Open Sci. 2021;1:112–122.

29. Seney ML, Kim S-M, Glausier JR, Hildebrand MA, Xue X, Zong W, et al. Transcriptional

Alterations in Dorsolateral Prefrontal Cortex and Nucleus Accumbens Implicate

Neuroinflammation and Synaptic Remodeling in Opioid Use Disorder. Biol Psychiatry.

2021;90:550–562.

30.   Lefevre EM, Pisansky MT, Toddes C, Baruffaldi F, Pravetoni M, Tian L, et al.

830   Interruption of continuous opioid exposure exacerbates drug-evoked adaptations in the

831   mesolimbic dopamine system. Neuropsychopharmacology. 2020;45:1781–1792.

832   31.   Sugino S, Konno D, Abe J, Imamura-Kawasawa Y, Kido K, Suzuki J, et al. Crucial

833   involvement of catecholamine neurotransmission in postoperative nausea and vomiting:

834   Whole-transcriptome profiling in the rat nucleus of the solitary tract. Genes Brain Behav.

835   2021:e12759.

836   32.   Ye J, Yang Z, Li C, Cai M, Zhou D, Zhang Q, et al. NF-κB signaling and vesicle transport

837   are correlated with the reactivation of the memory trace of morphine dependence. Diagn

838   Pathol. 2014;9:142.

839   33.   Korostynski M, Piechota M, Dzbek J, Mlynarski W, Szklarczyk K, Ziolkowska B, et al.

840   Novel drug-regulated transcriptional networks in brain reveal pharmacological

841   properties of psychotropic drugs. BMC Genomics. 2013;14:606.

842   34.   Klein G, Juni A, Waxman AR, Arout CA, Inturrisi CE, Kest B. A survey of acute and

843   chronic heroin dependence in ten inbred mouse strains: evidence of genetic correlation

844   with morphine dependence. Pharmacol Biochem Behav. 2008;90:447–452.

845   35.   Kest B, Palmese CA, Hopkins E, Adler M, Juni A, Mogil JS. Naloxone-precipitated

846   withdrawal jumping in 11 inbred mouse strains: evidence for common genetic

847   mechanisms in acute and chronic morphine physical dependence. Neuroscience.

848   2002;115:463–469.

849   36.   Reiner BC, Zhang Y, Stein LM, Perea ED, Arauco-Shapiro G, Ben Nathan J, et al.

850   Single nucleus transcriptomic analysis of rat nucleus accumbens reveals cell type-

851   specific patterns of gene expression associated with volitional morphine intake. Transl

852   Psychiatry. 2022;12:374.

853   37.   Moussawi K, Ortiz MM, Gantz SC, Tunstall BJ, Marchette RCN, Bonci A, et al. Fentanyl

854   vapor self-administration model in mice to study opioid addiction. Sci Adv.

855   2020;6:eabc0413.

38. Corre J, van Zessen R, Loureiro M, Patriarchi T, Tian L, Pascoli V, et al. Dopamine

neurons projecting to medial shell of the nucleus accumbens drive heroin

reinforcement. Elife. 2018;7:e39945.

39. Tapocik JD, Luu TV, Mayo CL, Wang BD, Doyle E, Lee AD, et al. Neuroplasticity, axonal

guidance and micro-RNA genes are associated with morphine self-administration

behavior. Addict Biol. 2012. 15 July 2012. https://doi.org/10.1111/j.1369-

1600.2012.00470.x.

40. Avey D, Sankararaman S, Yim AKY, Barve R, Milbrandt J, Mitra RD. Single-Cell RNA-

Seq Uncovers a Robust Transcriptional Response to Morphine by Glia. Cell Rep.

2018;24:3619-3629.e4.

41. Liang J, Chen J-H, Chen X-H, Peng Y-H, Zheng X-G. Gene expression of conditioned

locomotion and context-specific locomotor sensitization controlled by morphine-

associated environment. Behav Brain Res. 2011;216:321–331.

42. Tapocik JD, Luu TV, Mayo CL, Wang B-D, Doyle E, Lee AD, et al. Neuroplasticity,

axonal guidance and micro-RNA genes are associated with morphine self-

administration behavior. Addict Biol. 2013;18:480–495.

43. Grice DE, Reenilä I, Männistö PT, Brooks AI, Smith GG, Golden GT, et al.

Transcriptional profiling of C57 and DBA strains of mice in the absence and presence

of morphine. BMC Genomics. 2007;8:76.

44. Ikeda H, Miyatake M, Koshikawa N, Ochiai K, Yamada K, Kiss A, et al. Morphine

modulation of thrombospondin levels in astrocytes and its implications for neurite

outgrowth and synapse formation. J Biol Chem. 2010;285:38415–38427.

45. Imperio CG, McFalls AJ, Colechio EM, Masser DR, Vrana KE, Grigson PS, et al.

Assessment of individual differences in the rat nucleus accumbens transcriptome

following taste-heroin extended access. Brain Res Bull. 2016;123:71–80.

46. Sillivan SE, Whittard JD, Jacobs MM, Ren Y, Mazloom AR, Caputi FF, et al. ELK1

transcription factor linked to dysregulated striatal mu opioid receptor signaling network

and OPRM1 polymorphism in human heroin abusers. Biol Psychiatry. 2013;74:511–519.

47. Albertson DN, Schmidt CJ, Kapatos G, Bannon MJ. Distinctive profiles of gene expression in the human nucleus accumbens associated with cocaine and heroin abuse. Neuropsychopharmacology. 2006;31:2304–2312.

48. Zhang Y, Liang Y, Randesi M, Yuferov V, Zhao C, Kreek MJ. Chronic Oxycodone Self-administration Altered Reward-related Genes in the Ventral and Dorsal Striatum of C57BL/6J Mice: An RNA-seq Analysis. Neuroscience. 2018;393:333–349.

49. Zhang Y, Liang Y, Levran O, Randesi M, Yuferov V, Zhao C, et al. Alterations of expression of inflammation/immune-related genes in the dorsal and ventral striatum of adult C57BL/6J mice following chronic oxycodone self-administration: a RNA sequencing study. Psychopharmacology (Berl). 2017;234:2259–2275.

50. Yuferov V, Zhang Y, Liang Y, Zhao C, Randesi M, Kreek MJ. Oxycodone Self-Administration Induces Alterations in Expression of Integrin, Semaphorin and Ephrin Genes in the Mouse Striatum. Front Psychiatry. 2018;9:257.

51. Imperio CG, McFalls AJ, Hadad N, Blanco-Berdugo L, Masser DR, Colechio EM, et al. Exposure to environmental enrichment attenuates addiction-like behavior and alters molecular effects of heroin self-administration in rats. Neuropharmacology. 2018;139:26–40.

52. Yan B, Hu Z, Yao W, Le Q, Xu B, Liu X, et al. MiR-218 targets MeCP2 and inhibits heroin seeking behavior. Sci Rep. 2017;7:40413.

53. Zhang H, Wang Q, Wang Q, Liu A, Qin F, Sun Q, et al. Circular RNA expression profiling in the nucleus accumbens: Effects of electroacupuncture treatment on morphine-induced conditioned place preference. Addict Biol. 2020;25:e12794.

54. Martin JA, Caccamise A, Werner CT, Viswanathan R, Polanco JJ, Stewart AF, et al. A Novel Role for Oligodendrocyte Precursor Cells (OPCs) and Sox10 in Mediating Cellular and Behavioral Responses to Heroin. Neuropsychopharmacology. 2018;43:1385–1394.

55.   Ammon-Treiber S, Tischmeyer H, Riechert U, Höllt V. Gene expression of transcription factors in the rat brain after morphine withdrawal. Neurochem Res. 2004;29:1267–1273.

56.   Kuntz-Melcavage KL, Brucklacher RM, Grigson PS, Freeman WM, Vrana KE. Gene expression changes following extinction testing in a heroin behavioral incubation model. BMC Neurosci. 2009;10:95.

57.   Ammon S, Mayer P, Riechert U, Tischmeyer H, Höllt V. Microarray analysis of genes expressed in the frontal cortex of rats chronically treated with morphine and after naloxone precipitated withdrawal. Brain Res Mol Brain Res. 2003;112:113–125.

58.   Mendez EF, Wei H, Hu R, Stertz L, Fries GR, Wu X, et al. Angiogenic gene networks are dysregulated in opioid use disorder: evidence from multi-omics and imaging of postmortem human brain. Mol Psychiatry. 2021;26:7803–7812.

59.   Liu A, Dai Y, Mendez EF, Hu R, Fries GR, Najera KE, et al. Genome-Wide Correlation of DNA Methylation and Gene Expression in Postmortem Brain Tissues of Opioid Use Disorder Patients. Int J Neuropsychopharmacol. 2021;24:879–891.

60.   Jiang C, Wang X, Le Q, Liu P, Liu C, Wang Z, et al. Morphine coordinates SST and PV interneurons in the prelimbic cortex to disinhibit pyramidal neurons and enhance reward. Mol Psychiatry. 2021;26:1178–1193.

61.   Shu C, Sosnowski DW, Tao R, Deep-Soboslay A, Kleinman JE, Hyde TM, et al. Epigenome-wide study of brain DNA methylation following acute opioid intoxication. Drug Alcohol Depend. 2021;221:108658.

62.   Loguinov AV, Anderson LM, Crosby GJ, Yukhananov RY. Gene expression following acute morphine administration. Physiol Genomics. 2001;6:169–181.

63.   Choi MR, Jin Y-B, Bang SH, Im C-N, Lee Y, Kim H-N, et al. Age-related Effects of Heroin on Gene Expression in the Hippocampus and Striatum of Cynomolgus Monkeys. Clin Psychopharmacol Neurosci. 2020;18:93–108.

64.   Skupio U, Sikora M, Korostynski M, Wawrzczak-Bargiela A, Piechota M, Ficek J, et al.
Behavioral and transcriptional patterns of protracted opioid self-administration in mice.
Addict Biol. 2017;22:1802–1816.

65.   Piechota M, Korostynski M, Solecki W, Gieryk A, Slezak M, Bilecki W, et al. The
dissection of transcriptional modules regulated by various drugs of abuse in the mouse
striatum. Genome Biol. 2010;11:R48.

66.   Korostynski M, Piechota M, Kaminska D, Solecki W, Przewlocki R. Morphine effects on
striatal transcriptome in mice. Genome Biol. 2007;8:R128.

67.   Piechota M, Korostynski M, Sikora M, Golda S, Dzbek J, Przewlocki R. Common
transcriptional effects in the mouse striatum following chronic treatment with heroin and
methamphetamine. Genes Brain Behav. 2012;11:404–414.

68.   Korostynski M, Piechota M, Golda S, Przewlocki R. High-throughput gene expression
profiling of opioid-induced alterations in discrete brain areas. Methods Mol Biol.
2015;1230:65–76.

69.   Choi MR, Jin Y-B, Kim H-N, Chai YG, Im C-N, Lee S-R, et al. Gene expression in the
striatum of cynomolgus monkeys after chronic administration of cocaine and heroin.
Basic Clin Pharmacol Toxicol. 2021;128:686–698.

70.   Cha HJ, Choi M-J, Ahn J-I, Jeon S-H, Kang H, Kim EJ, et al. Comparison of gene
expression profiles in drug-withdrawn rats. Mol Cell Toxicol. 2016;12:197–207.

71.   Valderrama-Carvajal A, Irizar H, Gago B, Jiménez-Urbieta H, Fuxe K, Rodríguez-Oroz
MC, et al. Transcriptomic integration of D4R and MOR signaling in the rat caudate
putamen. Sci Rep. 2018;8:7337.

72.   Egervari G, Akpoyibo D, Rahman T, Fullard JF, Callens JE, Landry JA, et al. Chromatin
accessibility mapping of the striatum identifies tyrosine kinase FYN as a therapeutic
target for heroin use disorder. Nat Commun. 2020;11:4634.

73.   Heller EA, Kaska S, Fallon B, Ferguson D, Kennedy PJ, Neve RL, et al. Morphine and
cocaine increase serum- and glucocorticoid-inducible kinase 1 activity in the ventral
tegmental area. J Neurochem. 2015;132:243–253.

74.   McClung CA, Nestler EJ, Zachariou V. Regulation of gene expression by chronic
966   morphine and morphine withdrawal in the locus ceruleus and ventral tegmental area. J
967   Neurosci. 2005;25:6005–6015.

968   75.   Zhang H, Wang Q, Sun Q, Qin F, Nie D, Li Q, et al. Effects of Compound 511 on BDNF-
969   TrkB Signaling in the Mice Ventral Tegmental Area in Morphine-Induced Conditioned
970   Place Preference. Cell Mol Neurobiol. 2021;41:961–975.

971   76.   Jokinen V, Sidorova Y, Viisanen H, Suleymanova I, Tiilikainen H, Li Z, et al. Differential
972   Spinal and Supraspinal Activation of Glia in a Rat Model of Morphine Tolerance.
973   Neuroscience. 2018;375:10–24.

974   77.   Cheng Y-C, Tsai R-Y, Sung Y-T, Chen I-J, Tu T-Y, Mao Y-Y, et al. Melatonin regulation
975   of transcription in the reversal of morphine tolerance: Microarray analysis of differential
976   gene expression. Int J Mol Med. 2019;43:791–806.

977   78.   Shao J, Wang J, Huang J, Liu C, Pan Y, Guo Q, et al. Identification of lncRNA
978   expression profiles and ceRNA analysis in the spinal cord of morphine-tolerant rats. Mol
979   Brain. 2018;11:21.

980   79.   Peng Y, Guo G, Shu B, Liu D, Su P, Zhang X, et al. Spinal CX3CL1/CX3CR1 May Not
981   Directly Participate in the Development of Morphine Tolerance in Rats. Neurochem
982   Res. 2017;42:3254–3267.

983   80.   Santiago JM, Rosas O, Torrado AI, González MM, Kalyan-Masih PO, Miranda JD.
984   Molecular, anatomical, physiological, and behavioral studies of rats treated with
985   buprenorphine after spinal cord injury. J Neurotrauma. 2009;26:1783–1793.

986   81.   Weng Y, Wu J, Li L, Shao J, Li Z, Deng M, et al. Circular RNA expression profile in the
987   spinal cord of morphine tolerated rats and screen of putative key circRNAs. Mol Brain.
988   2019;12:79.

989   82.   Juul SE, Beyer RP, Bammler TK, Farin FM, Gleason CA. Effects of neonatal stress and
990   morphine on murine hippocampal gene expression. Pediatr Res. 2011;69:285–292.

991   83.   Marie-Claire C, Courtin C, Robert A, Gidrol X, Roques BP, Noble F. Sensitization to the

992   conditioned rewarding effects of morphine modulates gene expression in rat

993   hippocampus. Neuropharmacology. 2007;52:430–435.

994   84.   Zheng H, Zeng Y, Zhang X, Chu J, Loh HH, Law P-Y. mu-Opioid receptor agonists

995   differentially regulate the expression of miR-190 and NeuroD. Mol Pharmacol.

996   2010;77:102–109.

997   85.   Befort K, Filliol D, Ghate A, Darcq E, Matifas A, Muller J, et al. Mu-opioid receptor

998   activation induces transcriptional plasticity in the central extended amygdala. Eur J

999   Neurosci. 2008;27:2973–2984.

86.   Rodriguez Parkitna JM, Bilecki W, Mierzejewski P, Stefanski R, Ligeza A, Bargiela A,

et al. Effects of morphine on gene expression in the rat amygdala. J Neurochem.

2004;91:38–48.

87.   Zachariou V, Liu R, LaPlant Q, Xiao G, Renthal W, Chan GC, et al. Distinct roles of

adenylyl cyclases 1 and 8 in opiate dependence: behavioral, electrophysiological, and

molecular studies. Biol Psychiatry. 2008;63:1013–1021.

88.   Saad MH, Rumschlag M, Guerra MH, Savonen CL, Jaster AM, Olson PD, et al.

Differentially expressed gene networks, biomarkers, long noncoding RNAs, and shared

responses with cocaine identified in the midbrains of human opioid abusers. Sci Rep.

2019;9:1534.

89.   Befort K, Filliol D, Darcq E, Ghate A, Matifas A, Lardenois A, et al. Gene expression is

altered in the lateral hypothalamus upon activation of the mu opioid receptor. Ann N Y

Acad Sci. 2008;1129:175–184.

90.   Anghel A, Jamieson C a. M, Ren X, Young J, Porche R, Ozigbo E, et al. Gene

expression profiling following short-term and long-term morphine exposure in mice

uncovers genes involved in food intake. Neuroscience. 2010;167:554–566.

91.   Hayase T, Yamamoto Y, Yamamoto K, Muso E, Shiota K. Microarray profile analysis of

toxic cocaine-induced alterations in the expression of mouse brain gene sequences: a

possible 'protective' effect of buprenorphine. J Appl Toxicol. 2004;24:15–20.

92. Hassan HE, Myers AL, Lee IJ, Chen H, Coop A, Eddington ND. Regulation of gene
expression in brain tissues of rats repeatedly treated by the highly abused opioid
agonist, oxycodone: microarray profiling and gene mapping analysis. Drug Metab
Dispos. 2010;38:157–167.

93. Gaspari S, Purushothaman I, Cogliani V, Sakloth F, Neve RL, Howland D, et al.
Suppression of RGSz1 function optimizes the actions of opioid analgesics by
mechanisms that involve the Wnt/β-catenin pathway. Proc Natl Acad Sci U S A.
2018;115:E2085–E2094.

94. Toorie AM, Vassoler FM, Qu F, Schonhoff CM, Bradburn S, Murgatroyd CA, et al. A
history of opioid exposure in females increases the risk of metabolic disorders in their
future male offspring. Addict Biol. 2021;26:e12856.

95. Thibault K, Calvino B, Rivals I, Marchand F, Dubacq S, McMahon SB, et al. Molecular
mechanisms underlying the enhanced analgesic effect of oxycodone compared to
morphine in chemotherapy-induced neuropathic pain. PLoS One. 2014;9:e91297.

96. Delfs JM, Zhu Y, Druhan JP, Aston-Jones G. Noradrenaline in the ventral forebrain is
critical for opiate withdrawal-induced aversion. Nature. 2000;403:430–434.

97. Zhu Y, Wienecke CF, Nachtrab G, Chen X. A thalamic input to the nucleus accumbens
mediates opiate dependence. Nature. 2016;530:219–222.

98. Koob GF. The dark side of emotion: the addiction perspective. Eur J Pharmacol.
2015;753:73–87.

99. Hearing M. Prefrontal-accumbens opioid plasticity: Implications for relapse and
dependence. Pharmacol Res. 2019;139:158–165.

100. Lutz P-E, Kieffer BL. The multiple facets of opioid receptor function: implications for
addiction. Curr Opin Neurobiol. 2013;23:473–479.

101. Lutz P-E, Almeida D, Filliol D, Jollant F, Kieffer BL, Turecki G. Increased functional
coupling of the mu opioid receptor in the anterior insula of depressed individuals.
Neuropsychopharmacology. 2021;46:920–927.

102. Lutz P-E, Kieffer BL. Opioid receptors: distinct roles in mood disorders. Trends
1047   Neurosci. 2013;36:195–206.

1048   103. Kitchen I, Slowe SJ, Matthes HW, Kieffer B. Quantitative autoradiographic mapping of
1049   mu-, delta- and kappa-opioid receptors in knockout mice lacking the mu-opioid receptor
1050   gene. Brain Res. 1997;778:73–88.

1051   104. Goody RJ, Oakley SM, Filliol D, Kieffer BL, Kitchen I. Quantitative autoradiographic
1052   mapping of opioid receptors in the brain of delta-opioid receptor gene knockout mice.
1053   Brain Res. 2002;945:9–19.

1054   105. Slowe SJ, Simonin F, Kieffer B, Kitchen I. Quantitative autoradiography of mu-,delta-
1055   and kappa1 opioid receptors in kappa-opioid receptor knockout mice. Brain Res.
1056   1999;818:335–345.

1057   106. Xia Y, Haddad GG. Ontogeny and distribution of opioid receptors in the rat brainstem.
1058   Brain Res. 1991;549:181–193.

1059   107. Miyakawa T. No raw data, no science: another possible source of the reproducibility
1060   crisis. Mol Brain. 2020;13:24.

1061   108. Field D, Sansone S-A, Collis A, Booth T, Dukes P, Gregurick SK, et al. Megascience.
1062   'Omics data sharing. Science. 2009;326:234–236.

1063   109. NIH Genomic Data Sharing.

1064   110. Murray JE, Belin-Rauscent A, Simon M, Giuliano C, Benoit-Marand M, Everitt BJ, et al.
1065   Basolateral and central amygdala differentially recruit and maintain dorsolateral
1066   striatum-dependent cocaine-seeking habits. Nat Commun. 2015;6:10088.

1067   111. Zezula J, Casaccia-Bonnefil P, Ezhevsky SA, Osterhout DJ, Levine JM, Dowdy SF, et
1068   al. p21cip1 is required for the differentiation of oligodendrocytes independently of cell
1069   cycle withdrawal. EMBO Rep. 2001;2:27–34.

1070   112. Scholpa NE, Briggs SB, Wagner JJ, Cummings BS. Cyclin-Dependent Kinase Inhibitor
1071   1a (p21) Modulates Response to Cocaine and Motivated Behaviors. J Pharmacol Exp
1072   Ther. 2016;357:56–65.

113. Cahill KM, Huo Z, Tseng GC, Logan RW, Seney ML. Improved identification of
concordant and discordant gene expression signatures using an updated rank-rank
hypergeometric overlap approach. Sci Rep. 2018;8:9588.

114. Al-Hasani R, Bruchas MR. Molecular mechanisms of opioid receptor-dependent
signaling and behavior. Anesthesiology. 2011;115:1363–1381.

115. Coffey KR, Lesiak AJ, Marx RG, Vo EK, Garden GA, Neumaier JF. A cyclic AMP related
gene network in microglia is inversely regulated by morphine tolerance and withdrawal.
Biological Psychiatry Global Open Science. 2021:S2667174321000884.

116. Nectow AR, Moya MV, Ekstrand MI, Mousa A, McGuire KL, Sferrazza CE, et al. Rapid
Molecular Profiling of Defined Cell Types Using Viral TRAP. Cell Rep. 2017;19:655–
667.

117. Heiman M, Schaefer A, Gong S, Peterson JD, Day M, Ramsey KE, et al. A translational
profiling approach for the molecular characterization of CNS cell types. Cell.
2008;135:738–748.

118. Bailly J, Del Rossi N, Runtz L, Li J-J, Park D, Scherrer G, et al. Targeting morphine-
responsive neurons: generation of a knock-in mouse line expressing Cre recombinase
from the mu opioid receptor gene locus. ENeuro. 2020. 7 May 2020.
https://doi.org/10.1523/ENEURO.0433-19.2020.

119. Okunomiya T, Hioki H, Nishimura C, Yawata S, Imayoshi I, Kageyama R, et al.
Generation of a MOR-CreER knock-in mouse line to study cells and neural circuits
involved in mu opioid receptor signaling. Genesis. 2020;58:e23341.

120. Märtin A, Calvigioni D, Tzortzi O, Fuzik J, Wärnberg E, Meletis K. A Spatiomolecular
Map of the Striatum. Cell Rep. 2019;29:4320-4333.e5.

121. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel
Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell.
2015;161:1202–1214.

122. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177:1888-1902.e21.

123. Browne CJ, Godino A, Salery M, Nestler EJ. Epigenetic Mechanisms of Opioid

Addiction. Biol Psychiatry. 2020;87:22–33.

124. Belzeaux R, Lalanne L, Kieffer BL, Lutz PE. Focusing on the Opioid System for

Addiction Biomarker Discovery. Trends Mol Med. 2018;24:206–220.

125. Wu Q, Hwang CK, Zheng H, Wagley Y, Lin H-Y, Kim DK, et al. MicroRNA 339 down-

regulates μ-opioid receptor at the post-transcriptional level in response to opioid

treatment. FASEB J. 2013;27:522–535.

126. Shu C, Sosnowski DW, Tao R, Deep-Soboslay A, Kleinman JE, Hyde TM, et al.

Epigenome-wide study of brain DNA methylation following acute opioid intoxication.

Drug Alcohol Depend. 2021;221:108658.

127. Liu A, Dai Y, Mendez EF, Hu R, Fries GR, Najera KE, et al. Genome-Wide Correlation

of DNA Methylation and Gene Expression in Postmortem Brain Tissues of Opioid Use

Disorder Patients. Int J Neuropsychopharmacol. 2021;24:879–891.

128. Lutz P-E, Mechawar N, Turecki G. Neuropathology of suicide: recent findings and future

directions. Mol Psychiatry. 2017;22:1395–1412.

129. Ebrahimi G, Asadikaram G, Akbari H, Nematollahi MH, Abolhassani M, Shahabinejad

G, et al. Elevated levels of DNA methylation at the OPRM1 promoter region in men with

opioid use disorder. Am J Drug Alcohol Abuse. 2017:1–7.

130. Nielsen DA, Yuferov V, Hamon S, Jackson C, Ho A, Ott J, et al. Increased OPRM1 DNA

methylation in lymphocytes of methadone-maintained former heroin addicts.

Neuropsychopharmacology. 2009;34:867–873.

131. Doehring A, Oertel BG, Sittl R, Lotsch J. Chronic opioid use is associated with increased

DNA methylation correlating with increased clinical pain. Pain. 2013;154:15–23.

132. Barrow TM, Byun HM, Li X, Smart C, Wang YX, Zhang Y, et al. The effect of morphine

upon DNA methylation in ten regions of the rat brain. Epigenetics. 2017;12:1038–1047.

133. Lutz P-E, Chay M-A, Pacis A, Chen GG, Aouabed Z, Maffioletti E, et al. Non-CG

methylation and multiple histone profiles associate child abuse with immune and small

GTPase dysregulation. Nat Commun. 2021;12:1132.

134. Feng J, Shao N, Szulwach KE, Vialou V, Huynh J, Zhong C, et al. Role of Tet1 and 5-
hydroxymethylcytosine in cocaine action. Nat Neurosci. 2015;18:536–544.

135. Feng J, Wilkinson M, Liu X, Purushothaman I, Ferguson D, Vialou V, et al. Chronic
cocaine-regulated epigenomic changes in mouse nucleus accumbens. Genome Biol.
2014;15:R65.

136. Deroche-Gamonet V, Belin D, Piazza PV. Evidence for addiction-like behavior in the
rat. Science. 2004;305:1014–1017.

137. Domingo-Rodriguez L, Ruiz de Azua I, Dominguez E, Senabre E, Serra I, Kummer S,
et al. A specific prelimbic-nucleus accumbens pathway controls resilience versus
vulnerability to food addiction. Nat Commun. 2020;11:782.

138. Laine MA, Trontti K, Misiewicz Z, Sokolowska E, Kulesskaya N, Heikkinen A, et al.
Genetic Control of Myelin Plasticity after Chronic Psychosocial Stress. ENeuro.
2018;5:ENEURO.0166-18.2018.

139. Bagot RC, Cates HM, Purushothaman I, Lorsch ZS, Walker DM, Wang J, et al. Circuit-
wide Transcriptional Profiling Reveals Brain Region-Specific Gene Networks
Regulating Depression Susceptibility. Neuron. 2016;90:969–983.

140. Roy B, Wang Q, Dwivedi Y. Long Noncoding RNA-Associated Transcriptomic Changes
in Resiliency or Susceptibility to Depression and Response to Antidepressant
Treatment. Int J Neuropsychopharmacol. 2018;21:461–472.

141. Cervera-Juanes R, Wilhelm LJ, Park B, Grant KA, Ferguson B. Alcohol-dose-
dependent DNA methylation and expression in the nucleus accumbens identifies
coordinated regulation of synaptic genes. Transl Psychiatry. 2017;7:e994.

142. Domingo-Rodriguez L, Cabana-Domínguez J, Fernàndez-Castillo N, Cormand B,
Martín-García E, Maldonado R. Differential expression of miR-1249-3p and miR-34b-
5p between vulnerable and resilient phenotypes of cocaine addiction. Addict Biol.
2022;27:e13201.

143. Ekstrand MI, Nectow AR, Knight ZA, Latcha KN, Pomeranz LE, Friedman JM. Molecular
profiling of neurons based on connectivity. Cell. 2014;157:1230–1242.

144. Zhang Z, Zhou J, Tan P, Pang Y, Rivkin AC, Kirchgessner MA, et al. Epigenomic diversity of cortical projection neurons in the mouse brain. Nature. 2021;598:167–173.

145. Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. Science. 2019;363:1463–1467.

146. Eng C-HL, Lawson M, Zhu Q, Dries R, Koulena N, Takei Y, et al. Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. Nature. 2019;568:235–239.

147. Borm LE, Mossi Albiach A, Mannens CCA, Janusauskas J, Özgün C, Fernández-García D, et al. Scalable in situ single-cell profiling by electrophoretic capture of mRNA using EEL FISH. Nat Biotechnol. 2022. 22 September 2022. https://doi.org/10.1038/s41587-022-01455-3.

148. Armand EJ, Li J, Xie F, Luo C, Mukamel EA. Single-Cell Sequencing of Brain Cell Transcriptomes and Epigenomes. Neuron. 2021;109:11–26.

149. Feng J, Wilkinson M, Liu X, Purushothaman I, Ferguson D, Vialou V, et al. Chronic cocaine-regulated epigenomic changes in mouse nucleus accumbens. Genome Biol. 2014;15:R65.

150. Mokhtari A, Porte B, Belzeaux R, Etain B, Ibrahim EC, Marie-Claire C, et al. The molecular pathophysiology of mood disorders: From the analysis of single molecular layers to multi-omic integration. Prog Neuropsychopharmacol Biol Psychiatry. 2022;116:110520.

151. Hasin Y, Seldin M, Lusis A. Multi-omics approaches to disease. Genome Biol. 2017;18:83.

152. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. BMJ. 2009;339:b2700–b2700.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6