The inhibitory mechanism of a small protein reveals its role in antimicrobial peptide sensing

Shan Jiang[1], Lydia C. Steup[1], Charlotte Kippnich[1], Symela Lazaridi[2,3], Gabriele Malengo[1], Thomas Lemmin[2] and Jing Yuan[1]*

Abstract

A large number of small membrane proteins have been discovered in bacteria, but their mechanism of action has remained mostly elusive. Here, we investigate the mechanism of a physiologically important small protein, MgrB, which represses the activity of the sensor kinase PhoQ and is widely distributed among enterobacteria. The PhoQ/PhoP two-component system is a master regulator of the bacterial virulence program and interacts with MgrB to modulate bacterial virulence, fitness, and drug resistance. A combination of crosslinking approaches with functional assays and protein dynamic simulations revealed structural rearrangements due to interactions between MgrB and PhoQ near the membrane/periplasm interface and along the transmembrane helices. These interactions induce the movement of the PhoQ catalytic domain and the repression of its activity. Without MgrB, PhoQ appears to be much less sensitive to antimicrobial peptides, including the commonly used C18G. In the presence of MgrB, C18G promotes MgrB to dissociate from PhoQ, thus activating PhoQ via derepression. Our findings reveal the inhibitory mechanism of the small protein MgrB and uncover its importance in antimicrobial peptide sensing.

Significance Statement

Small proteins have high prevalence, vast diversity, and primarily regulatory functions in biological processes across all domains of life. However, their mechanisms of action remain largely elusive. In this study, we investigate the mechanism of the small protein, MgrB. It interacts with the sensor kinase PhoQ, rearranges its conformation, represses its kinase activity, and regulates bacterial response to environmental changes. In particular for antimicrobial peptides, MgrB is required for bacteria to have a selective response to this host-exclusive stimulus. Our findings underline the importance of a small protein in bacterial fitness and drug resistance and provide a molecular basis for engineering novel peptide-based regulators.

Introduction

Small proteins are defined by their short sequences (usually < 50 aa) and their direct synthesis via ribosomal translation. An increasing number of small proteins have been identified in individual species in all domains of life and complex biological communities, suggesting a high prevalence and vast diversity in biological systems (reviewed in [1-4]). Many of these small proteins (~30%) are located in the cell membrane or have been predicted to be located there (e.g., [5, 6]). Although only a few detailed studies of small membrane proteins have been conducted, results show that small proteins may target larger proteins and the lipid bilayer, regulating protein function, abundance, and changing membrane properties, demonstrating their predominant role as regulators in the lipid bilayer (reviewed in [7]).

The membrane located PhoQ sensor kinase, a master regulator of a bacterial virulence program, detects host-associated stimuli (review in [8]), such as cationic antimicrobial peptides (CAMPs) [9], mild acidic pH [10, 11], increased osmolarity [12], biliary unsaturated long-chain fatty acids [13], and magnesium limitation [14]. Three small membrane proteins have been identified to control PhoQ activity via direct protein-protein interactions. In *E. coli,* MgrB (47aa) and SafA (65aa) modulate PhoQ activity, with MgrB acting as an inhibitor and SafA as an activator [15,

49   16]. Deletion of *mgrB* or *safA* has physiological consequences, resulting in a cell division block
50   under magnesium starvation or lowered acid resistance, respectively [17, 18]. In *Salmonella*
51   *enterica* serovar Typhimurium, a species-specific small protein, UgtS (34 aa), was discovered
52   recently and shown to control the timing of virulence gene expression by preventing premature
53   PhoQ activation when inside macrophages [19].
54
The PhoQ sensor kinase is found as a dimer on the bacterial inner membrane. It consists of a
periplasmic, a transmembrane (TM), a HAMP, a DHp, and a catalytic-ATP binding domain. In the
periplasmic domain, several molecular interactions are essential for the functional state of the
PhoQ molecule. These include the central hydrogen bonding network (T48 D179 K186) in the
PAS-fold [20-23], the inter-monomeric salt bridge (D179-R50') [24], the membrane-facing acidic
cluster ([148]EDDDDAE) and the juxta-membrane region at the periplasm/inner-membrane
interface [9, 23, 25]. Mutations altering these molecular interactions change the conformation
of the periplasmic domain and the sensitivity of PhoQ to magnesium, low pH, and/or CAMPs.
Additionally, different stimuli may result in distinct conformational states of the periplasmic
domain, at least in an *in vitro* setting. For example, the PhoQ periplasmic domain showed
divergent NMR spectra in low pH compared to that in low magnesium conditions [10]. Besides
the periplasmic domain, the PhoQ transmembrane domain was shown to detect environmental
osmotic upshifts, where the transmembrane four-helix bundle adopts a more compact and
kinase-active conformation [12].
69
Among the known PhoQ-regulating small proteins, MgrB is the most widely distributed PhoQ
repressor in enterobacteria. In *Salmonella*, the deletion of *mgrB* was shown to affect its
infection of macrophages and epithelial cells [26]. In *Klebsiella pneumonia*, the disruption of the
mgrB gene is the most prevalent among other chromosomal mutations in colistin-resistant
strains [27]. The expression of *mgrB* is PhoPQ dependent. Thus, the inhibitory function of MgrB
provides negative feedback to the two-component system resulting in a partial adaptation of *E.*
coli to environmental changes [15, 28]. MgrB is a bitopic inner membrane protein with a short
cytosolic N-terminus and a periplasmic C-terminal region (Fig. 1A). Two conserved cysteines
(C28 and C39) in the periplasmic region were shown to be essential for MgrB function and
suggested to form a disulfide bond [15]. This disulfide bond was proposed to allow MgrB to act
as a sensor for the redox state of the bacterial periplasm and create an entry point for redox
sensing by the PhoQ/PhoP system [29]. Besides the two cysteines, eight functionally essential
residues were identified spread across the entire molecule [30]. However, the mechanism of
MgrB inhibiting PhoQ remained largely unclear.
84
In this study, we combined crosslinking approaches with functional assays and protein dynamic
simulations to identify interactions between PhoQ and MgrB, probe MgrB-induced PhoQ
conformational changes, and examine the impact of MgrB on PhoQ sensing environmental
stimuli. We propose that the binding of MgrB (i) changes the conformation of the linker region
between PhoQ periplasmic and transmembrane domains, (ii) rearranges the transmembrane
four-helix bundle, and (iii) initiates the translocation of the catalytic domain. MgrB interferes
with PhoQ signaling, resulting in an overall lowered activity but still allows PhoQ activation
under low pH, osmotic upshift, and magnesium limitation. We further reveal that MgrB
mediates PhoQ sensing antimicrobial peptides, enabling bacteria to have a selective response to
this exclusively host-associated stimulus.
95
96

97   **Results**
98
Structural predictions by Alphafold2
We used Alphafold2 multimer [31-33] to predict the structures of the PhoQ/MgrB complex as
well as the individual component of the complex (Fig. 1B, C, and S1). The overall prediction of
structural elements agrees with previous experimental evidence. For example, the two
conserved periplasmic cysteine residues (C28 and C39) in MgrB were positioned in close
proximity in the models, structurally supporting the presence of a disulfide bond. In the PhoQ
predictions, the PAS fold in the periplasmic domain, the HAMP, the DHp, and the cytosolic
catalytic-ATP binding domains resembled the experimental results of these structural motifs [24,
34]. Notably, the region connecting the periplasmic and TM domains and that connecting the
HAMP and DHp domains showed a lower predicted local distance difference test loss (pLDDT)
score (colored in yellow to green in Fig. S1), indicating these linker regions might have a higher
capability of adopting different conformations. Unlike the asymmetric crystal structure of the
PhoQ sensor domain [24], the Alphafold2 predicted PhoQ dimer was symmetric, which likely
represented only one of many conformations that PhoQ could adopt. In all five PhoQ/MgrB
complex models (Fig. 1B and S1), MgrB was placed in between the two PhoQ protomers, with
the transmembrane helix parallel to the PhoQ TM helices and the C-terminal short helix (A40-
W47) protruding away from PhoQ. The five predicted complex models had only minor
differences regarding the positioning of MgrB. Because the PhoQ dimer was mostly symmetrical,
Alphafold2 randomly placed MgrB on one or the other side of the PhoQ dimer (Fig. S1). When
bound to PhoQ, MgrB showed a major conformational change in the region flanked by the two
conserved cysteines (C28 and C39). This region had reduced helicity and rigidness, appearing as
an extended flexible loop (Fig. 1C).
121
Disulfide crosslinking between MgrB and PhoQ TM helices
Before Alphafold was available, we used the disulfide crosslinking approach to identify the
interacting residues in PhoQ and MgrB. We started with residue W20 in the MgrB TM helix,
which was suggested to have direct contact with PhoQ [30]. Sequence alignment of MgrB
homologs from different bacterial species revealed both W and Y conserved at position 20 (Fig.
S2). The MgrB W20Y variant also showed strong inhibitory activity comparable to the wild type
(Fig. S2). Since both W and Y have an aromatic ring structure and a polar moiety, we predicted
that the nature of the interactions at position 20 likely include hydrophobic interactions and
hydrogen bonding. Scanning through the sequence of the two PhoQ transmembrane helices,
Y32 in TM1 appeared to be the optimal candidate to fulfill both types of interaction with MgrB
W20.
133
To test this prediction with disulfide crosslinking, we constructed a PhoQ TM_Cys library and
generated the MgrB C16A W20C variant, which has only one cysteine residue in the
transmembrane helix. The Cys to Ala mutation at MgrB position 16 did not appear to have a
functional effect (Fig. S2B). However, the Trp to Cys mutation at position 20 significantly
reduced MgrB activity (Fig. S2A). To compensate for the loss of function, we increased the
induction of MgrB C16A W20C (pTrc promoter with 10 μM IPTG), which ensured significant
repression of PhoQ during crosslinking (Fig. S2D). Additionally, a FLAG tag was fused to the N-
terminus of MgrB (Fig. S2E) to facilitate the detection of the PhoQ/MgrB complex in Western
blot. Cu-phenanthroline was added to the cells to catalyze the disulfide formation between
sulfhydryl groups at a close distance. When MgrB C16A W20C was co-expressed with PhoQ
Y32C, we observed a higher molecular weight band at around 55 kDa on the Western blot

corresponding to the PhoQ/MgrB complex (Fig. 1D, S3A). This band was reduced by the addition
of dithiothreitol (DTT) and was absent when the cysteine residue was at position 16 (WT), or no
cysteine was in the MgrB transmembrane helix (C16A). The results are consistent with our
prediction suggesting the close proximity of MgrB W20 with the PhoQ TM1 residue Y32 in the
complex. Additionally, we found that PhoQ L204C crosslinked with MgrB W20C, indicating
position 20 near L204 in PhoQ TM2 as well (Fig. 1D, S3A). Furthermore, we tested PhoQ Cys
variants in the neighboring positions of 32 and 204 (PhoQ 31C, 33C, 203C, and 205C). None of
the variants formed detectable crosslinked complexes with MgrB W20C in the Western blot (Fig.
S2G). Functionally, both PhoQ Y32C and L204C mutants partially restored the MgrB inhibition
(Fig. S2F), supporting the hypothesis that MgrB W20 interacts with residue Y32 in PhoQ TM1
and L204 in PhoQ TM2.
156
Using MgrB residue 20 as a reference point, we next identified interacting amino acids of two
more functionally important residues (Q22 and F24) in the MgrB TM helix [30]. In both cases, we
observed stronger crosslinking bands and more than one crosslinked position in both
transmembrane helices of PhoQ (Fig. 1E, F, S3B, C). For MgrB Q22C, stronger crosslinks were
observed to residues in the PhoQ TM2 helix (colored blue). The strongest crosslinking band was
shown when PhoQ Y197C was co-expressed. The crosslinking efficiency decreased when the
cysteine residue moved to the neighboring positions and further away (196C, 195C, and 198C).
It was then partially recovered when the cysteine (194C) was roughly one helical turn away from
Y197 (Fig. 1E). Functionally, the Y197C mutation in PhoQ restored the MgrB Q22C activity the
best when compared to other PhoQ TM_Cys mutants, supporting the notion that MgrB Q22
primarily interacts with PhoQ Y197.
168
For MgrB F24C (Fig. 1F, S3C), the strongest crosslinking was found with a PhoQ TM1 mutant,
I38C. Functionally, PhoQ I38C showed the highest extent of activity reduction when co-
expressed with MgrB F24C, thus restoring the activity of MgrB F24C the best among the tested
PhoQ/MgrB pairs. The minor crosslinking events observed could be due to the greater distance
between the cysteine residues or an indication of intermediate binding states of MgrB. Notably,
one PhoQ Cys variant, F195C, retained only about 1% of the wild-type activity (Fig. 1E). Mutants
with such a severe functional defect may have altered MgrB binding, thus unsuitable for the
crosslinking approach. For other PhoQ variants in Fig. 1E and F, some showed moderately
reduced activity, possibly due to the reduced protein expression level (i.e., PhoQ W192C in Fig.
S3F). More importantly, the presence of MgrB significantly repressed the activity of these PhoQ
variants, indicating that the change caused by cysteine substitutions does not have a major
impact on MgrB inhibition.
181
Collectively, we identified four-pair residues of MgrB and PhoQ located in close proximity at the
protein interface of the complex: W20-Y32, W20-L204, Q22-Y197, and F24-I38. In the
AlphaFold2 predicted complex models, these crosslinked pairs were positioned less than 14 Å in
the top four ranked models (Table 1). Furthermore, Q22-Y197 and F24-I38 had shorter Cα - Cα
distances than the W20-Y32 and W20-L204 pairs, consistent with our experimental results that
the former two pairs showed higher crosslinking efficiency.
188
Crosslinking in the periplasmic domains of PhoQ and MgrB
To gain insights into PhoQ/MgrB interactions at the periplasmic domains, we attempted to
identify interacting residue pairs using site-specific photo-crosslinking [35]. We introduced an
amber stop codon at each residue in the MgrB periplasmic region (residues 25 to 47). The

photo-crosslinking amino acid, *p*-benzoyl-l-phenylalanine (*p*Bpa), was then incorporated into the
flag-tagged MgrB at the amber stop codon via the orthogonal tRNA[Amber] and the aminoacyl-tRNA
synthetase in *E. coli*. After UV irradiation, we detected bands with a molecular weight of ~55
kDa by anti-FLAG antibody when *p*Bpa was incorporated at positions 26, 30, 32, and 38,
indicating the formation of the UV-dependent crosslinked PhoQ/MgrB complex (Fig. 2A, S4).
198
The crosslinked products from positions 30, 32, and 38 could be purified and subjected to mass
spectrometry (MS) analysis (Fig. S4 and File S1). PhoQ was detected in all samples as the most
prominent protein. Only a small number of other proteins were identified, indicating the protein
purification steps were highly effective. MgrB was challenging to detect due to its small size,
high hydrophobicity, and a low number of potential MS-detectable peptides. However, we
increased the sample injection amount, which allowed for the successful identification of MgrB
in crosslinked complex samples, except when *p*Bpa was incorporated at position 38. This was
most likely due to the interference of the digestion step with the incorporated *p*Bpa. Despite the
successful identification of both PhoQ and MgrB in the crosslinked samples, confident MS
identification of MgrB-PhoQ crosslinked peptides providing clear evidence towards positioning
the two proteins in the complex was not possible. The lack of crosslinked products with other
mgrB_amber constructs (Fig. S4) could be due to the low efficiency of *p*Bpa incorporation (i.e.,
position 29) and/or the obstruction of the bulky unnatural amino acid on the binding of MgrB to
PhoQ. Most of the residues that resulted in crosslinked products were within the MgrB C28-C39
loop region, suggesting its possible direct interaction with PhoQ.
214
In the Alphafold2 complex model, the MgrB C28-C39 loop was close to the PhoQ linker region
connecting the periplasmic and the TM domains. To test whether the MgrB C28-C39 loop
interacts directly with the linker region, we performed disulfide crosslinking with MgrB D31C
and PhoQ_linker_Cys variants. Because of two essential cysteines in the MgrB periplasmic
region, we first tested if introducing a third cysteine would affect MgrB function. We observed
that the introduction of a third cysteine at position 31 did not reduce MgrB inhibition as much
as the single cysteine variant, C39A (Fig. S5), suggesting that the cysteine at position 31 does not
severely interfere with the C28-C39 disulfide bond formation. Therefore, MgrB D31C is suitable
for crosslinking experiments. The crosslinking results showed that MgrB 31C formed disulfide
linkages with cysteines in several positions in the PhoQ linker region (184-188) with a similar
amount of crosslinked complex (Fig. 2B, S3D). No crosslinked products were detectable in the
control sample, where the wild-type MgrB (C28 and C39) was co-expressed with
PhoQ_linker_Cys mutants (Fig. 2B, S3E). Our data thus support that MgrB D31 interacts with the
PhoQ linker region. The comparable amount of crosslinked complex at multiple positions is
probably due to the flexible nature of the MgrB C28-C39 loop.
230
The conformational change of PhoQ linker region upon MgrB binding
After mapping the MgrB binding site on PhoQ, we next sought to detect MgrB-induced PhoQ
conformational change by comparing the PhoQ dimer crosslinking pattern with MgrB to the one
without MgrB. For the PhoQ linker region, we focused on residues 187 and 188, which showed
different conformations in PhoQ monomer and dimer models (Fig. S6A). These two residues
were located at the periplasmic end of the TM2 helix and at a similar lateral position as the TM1
residue 43 (Fig. S6B). By monitoring the crosslinking efficiency between residues at positions 43
and 188, we intended to detect local helical movements upon MgrB binding. In the absence of
MgrB, we observed that 43C and 188C formed an intramolecular disulfide linkage, manifesting
as a faster-migrating band below the PhoQ monomer band (Fig. 3A). DTT treatment of the same

sample retarded the migration to the exact location as the PhoQ monomer, excluding the
likelihood of protein degradation. Nearly no PhoQ dimer band was detected under this
condition, further supporting that the orientation of the 188C side chain was more toward the
TM1 helix of the same protomer. In the presence of MgrB, the crosslinking pattern changed. The
intramolecular crosslinking band reduced, and the intensity of the PhoQ dimer band increased
(Fig. 3A, S6C). In this case, crosslinked PhoQ dimer could be formed in four ways: 43C-43'C, 43C-
188'C, 43'C-188C, and 188C-188'C. Our control samples showed that the crosslinking was
reduced for 43C-43'C and 188C-188'C when MgrB was present. Therefore, the increase in PhoQ
dimer formation was likely due to the inter-monomeric crosslinking between positions 43 and
188, suggesting that the binding of MgrB changes the crosslinking pattern from intra- to inter-
monomeric linkage and likely the relative orientation between PhoQ residue 188 to residue 43
(Fig. 3B). We also observed a similar trend of change in crosslinking for the residue 43 and 187
pair but with reduced intramolecular crosslinking efficiency (Fig. S6C). Taken together, we
hypothesize that MgrB may promote a rotational or translational movement of one TM2
periplasmic terminus towards the TM1 from the other promoter. Because the stoichiometry of
PhoQ and MgrB remains unknown, whether the conformational change detected here occurs in
both PhoQ protomers awaits further investigations.
258
MgrB repression of PhoQ is divergent from high magnesium repression
Besides the small protein MgrB, high magnesium also represses PhoQ kinase activity [14]. The
structural change of PhoQ under magnesium repression was studied in detail by monitoring the
arrangement of α helices along the PhoQ dimer interface via disulfide crosslinking [36]. It was
proposed that under high magnesium conditions, the N-terminal α helix across the TM and the
periplasmic domains undergoes a scissoring-type of motion with the pivot point near the
periplasm/membrane interface [36].
266
Using a similar approach, we monitored the arrangement of α helices along the PhoQ dimer
interface in the presence and absence of MgrB. Our data showed that most TM1 residues had
reduced crosslinking between PhoQ protomers upon MgrB binding (Fig. 3C blue residues, Fig.
S7). This reduction was more prominent for residues close to the cytoplasm, suggesting that
MgrB binding pushes TM1-TM1' apart and the propagation of PhoQ conformational change
from the periplasm to the cytoplasm. In contrast, the presence of MgrB did not show a similar
crosslinking effect on residues along the periplasmic helical interface (Fig. 3C purple residues,
Fig. S7). Most residues in this region showed little change in the crosslinking pattern. Some
showed a significantly decreased (i.e., residues 46 and 60) or a slightly increased dimer fraction
(i.e., residues 62). The overall PhoQ dimer rearrangement upon MgrB binding differed from the
scissoring motion proposed for high magnesium repression, suggesting that MgrB and high
magnesium act differently on PhoQ. We then did functional analyses of PhoQ under high
magnesium, with MgrB, or with both repressive factors. The results showed that PhoQ was
further repressed significantly when both high magnesium and MgrB were present (Fig. 3D),
supporting the hypothesis that magnesium and MgrB may have divergent mechanisms of
repression.
283
Molecular dynamics simulations of PhoQ/MgrB complex models
When comparing the predicted PhoQ dimer with its complex with MgrB, there was only minor
conformational change, which is in stark contrast to our experimental results. The Alphafold2
prediction might have only captured a transient arrangement of the complex but not the final
state. To investigate the effect of MgrB on the conformation of PhoQ in silico, we carried out a

set of four molecular dynamic simulations. The initial atomic models were obtained using the
best-ranked model predicted by the multimeric version of Alphafold2. In total, we modeled four
different complexes: PhoQ dimer (i) alone, and interacting with (ii) the wild-type MgrB, (iii) MgrB
W20Y, and (iv) MgrB W20A. Each complex was inserted into a bacterial mimetic membrane
bilayer, and simulations were carried out up to 1 μs under near-physiological conditions. To
evaluate the impact of the starting conformation on the stability of the PhoQ/MgrB complex, we
carried out two additional 500 ns simulations seeded from the second and third predictions. All
protein complexes remained stable throughout the entire simulations, with minor fluctuation in
the secondary structure. We did not observe any significant differences between the three
PhoQ/MrgB simulations. The MgrB C-terminal short helix (A40-W47) was buried close to the
membrane surface. The C-terminal carboxylate group interacted with the head group of PE
lipids in the simulations. The orientation of this C-terminal helix was primarily restrained by the
disulfide linkage (C28-C39) and remained stable in the simulations. A principal component
analysis of the transmembrane domain clearly showed that the functional MgrB variants (WT
and W20Y) induced a similar conformational change (Fig. 4A), corresponding to an increase in
the distance between PhoQ TM1 helices, coupled with a decrease in distance between TM2
helices. This change is more evident in TM residues near the membrane/cytoplasm interface
(Fig. 4B, S8). The MgrB W20A had only a marginal impact on the conformation of the
transmembrane domain.
308
In all the simulations with MgrB variants, we also observed the translocation of one of the PhoQ
catalytic-ATP binding (CA) domains (Fig. 4C). This is most prominent in the 1 μs simulations with
the wild-type MgrB, where one CA domain moved fully away from the helix harboring the
phosphorylatable histidine and located itself close to the inner membrane. For PhoQ alone, we
did not observe any significant movement of the CA domains throughout the entire simulation.
The functionally impaired MgrB W20A also induced PhoQ CA domain movement, though to a
lesser extent compared to the wild type (Fig. 4C). In this simulation, the CA domain remained
close to its initial position but rotated by about 90°. Our previous work showed that the W20A
mutant had low activity due to reduced affinity to PhoQ [30]. However, since the simulation was
seeded with a complex where MgrB W20A was already interacting with PhoQ, this likely
represents the scenario where overexpression of MgrB W20A compensates for its weak binding
to PhoQ and partially recovers its inhibitory function.
321
322   Overall, the simulation results confirm our experimental observations, showing the PhoQ
323   structural changes at the MgrB binding site. Furthermore, MgrB induces movements of the
324   distantly located CA domain, suggesting that the local conformational change due to MgrB
325   binding results in the repression of PhoQ kinase activity.
326
MgrB mediates PhoQ sensing cationic antimicrobial peptides
Our crosslinking data (Fig. 2B) suggested that MgrB directly interacts with PhoQ E184, a residue
involved in CAMP sensing [9]. We, therefore, hypothesized that MgrB might affect PhoQ
activation by CAMPs. To investigate this as well as the impact of MgrB on PhoQ sensing other
stimuli, we tested PhoQ activation by cationic antimicrobial peptides, low pH, and high
osmolarity in the presence and absence of MgrB (Fig. 5). Under a repressive magnesium
condition (LB medium supplemented with 10 mM Mg^{2+}), only slight or no PhoQ activation was
detected by antimicrobial peptide C18G regardless of MgrB, consistent with previous work
where high magnesium repressed the activation of PhoQ by CAMPs in *Salmonella* [9, 10].
Conversely, with the same magnesium condition (10 mM), a relatively large activation of PhoQ

was detected by low pH or osmolarity upshift in the presence of MgrB. The activation did not
appear to have a qualitative change when MgrB was absent, except that the overall PhoQ
activity increased due to the lack of MgrB repression (Fig. 5A).
340
We next followed PhoQ activities in the presence of antimicrobial peptide LL37 and C18G under
a physiologically relevant magnesium concentration (1 mM Mg^{2+} in LB) that does not repress
CAMPs from inducing PhoQ activity [9]. Up to a nearly 5-fold increase of PhoQ activity by
antimicrobial peptides was detected when the wild-type MgrB was present (Fig. 5B). However,
in the absence of MgrB, there was no significant further increase of PhoQ activity in the
presence of C18G. In the case of LL37, only a marginal increase (1.14-fold) of PhoQ activity was
observed, consistent with a previous report [15], suggesting that LL37 may be slightly different
from C18G in terms of PhoQ activation. To test if PhoQ was already maximally activated in the
ΔmgrB strain with 1 mM magnesium, we treated the cells with additional stimuli. PhoQ activity
was significantly further increased by lowered pH or elevated osmolarity (Fig. S9A), indicating
that PhoQ activity did not reach saturation with 1 mM magnesium alone. To be precise on the
total magnesium concentration in the growth medium, we tested our strains in the defined
minimal A medium [37] supplemented with 1 mM magnesium. Consistent with what we
observed in LB, PhoQ activity was higher in the absence of MgrB but could not be further
increased by C18G. Therefore, we hypothesize that MgrB may mediate PhoQ to sense
antimicrobial peptides under a physiologically relevant magnesium concentration.
357
The lack of PhoQ activation by antimicrobial peptides in the absence of MgrB might also be due
to the protective effect of the overall elevated activity of the PhoQ/PhoP two-component
system, which could increase the outer membrane modification and reduce the entry of
antimicrobial peptides to the periplasm. To obtain more direct evidence, we monitored PhoQ
and MgrB interactions *in vivo* using the acceptor photobleaching Förster resonance energy
transfer (FRET) approach. Functional PhoQ-mNeonGreen and mCherry-MgrB fusion proteins
(Fig. S9C and D) were expressed in *E. coli*. Cells were collected in the mid-log phase, treated with
C18G, and then used for FRET measurements (Fig. 5C). The control cells without C18G
treatment, showed energy transfer from mNeonGreen to mCherry with about 10% efficiency,
comparable to our previous report [12], indicating the formation of the PhoQ/MgrB complex *in*
vivo. With the C18G treatment, the FRET efficiency reduced to about 2%, slightly higher than the
negative controls, suggesting that C18G induces the dissociation of the PhoQ/MgrB complex.
370
To confirm the *in vivo* FRET results, we performed disulfide crosslinking experiments with or
without C18G using the PhoQ I38C MgrB F24C variant pair. We chose this pair because: i) they
formed a good amount of PhoQ/MgrB crosslinked complex readily detectable by Western blot
(Fig. 1F); ii) PhoQ I38C showed wild-type-like activity (Fig. 1F); iii) MgrB F24C repressed PhoQ
I38C to a relatively low level (Fig. 1F); iv) the variant pair responded to C18G showing
significantly increased PhoQ activity (Fig. 5D). Compared to the crosslinking in the absence of
C18G, we observed a lighter band corresponding to the PhoQ/MgrB complex (Fig. 5E). Assuming
that C18G does not adversely affect Cu: phenanthroline catalysis, the reduction in the
crosslinked complex suggests that C18G induces MgrB to move away from PhoQ, consistent
with our FRET results.
381
Taken together, we propose that MgrB is involved in PhoQ sensing CAMPs. Under
physiologically relevant magnesium conditions, PhoQ is active but partially repressed due to the
bound MgrB. This state allows PhoQ to detect and respond sensitively to CAMPs at a sub-

inhibitory concentration. CAMPs bind to PhoQ and induce the release of the bound MgrB,
further activating PhoQ/PhoP-regulated genes and potentially increasing bacterial resistance to
CAMPs (Fig. 6).
388
389 **Discussion**
390
In this study, we report that MgrB directly interacts with the TM helices and the periplasmic/TM
linker region of PhoQ, changing its conformation in a way that divergent from high magnesium
repression. It reduces the overall activity of PhoQ yet still allows its activation by low
magnesium, low pH, and increased osmolarity. We further reveal that PhoQ activation by
CAMPs is MgrB-dependent under physiologically relevant magnesium conditions. CAMPs induce
MgrB dissociating from PhoQ and thus increase PhoQ activity via derepression. A higher PhoQ
activity leads to an increased outer membrane modification and improved resistance to CAMPs.
Since the septum is the initial attacking site of certain CAMPs (i.e., LL37) on the membrane [38],
the possible division block caused by strong PhoQ activation may also protect bacteria against
the host's innate immune response. CAMPs were suggested to bind to an acidic surface in the
PhoQ periplasmic domain, though the exact binding site has not been well defined [9]. The
acidic cluster ([148]EDDDDAE), a major component of the acidic surface, is relatively distant from
the MgrB binding site. However, it remains unclear whether the acidic residue E184 in the linker
region is directly involved in CAMP binding. It is plausible that the binding of CAMPs induces
PhoQ conformational change, which leads to the dissociation of MgrB from PhoQ.
406
We observed that a high level of magnesium (10 mM) repressed PhoQ activation by CAMPs in
the presence or absence of MgrB in *E. coli*, consistent with a previous report on PhoQ in
Salmonella, though MgrB had not yet been discovered at that time [9]. The same report
proposed that CAMPs activate PhoQ by replacing magnesium ions from the acidic cluster in
PhoQ, which is different from our MgrB-mediated CAMP sensing mechanism. Both hypotheses,
however, do not conflict with each other but rather describe two aspects of PhoQ sensing
CAMPs under different magnesium conditions. It is also possible that magnesium may affect
MgrB binding to PhoQ, thus connecting these two mechanisms of CAMP sensing by PhoQ.
Interestingly, the magnesium replacement mechanism can not explain PhoQ activation by
antimicrobial peptides in some cases. For example, a recent study revealed that antimicrobial
peptides with a ß-sheet structure or a neutral charge could also activate PhoQ [39]. These newly
found PhoQ activators do not possess the properties to interact with the PhoQ acidic cluster.
Furthermore, there are PhoQ homologs that do not contain an acidic cluster in their sequences.
Our MgrB-dependent antimicrobial peptide sensing mechanism may provide a suitable
explanation for both cases.
422
By monitoring PhoQ conformational change and activities, we show that MgrB repression
diverges from high magnesium repression. This is consistent with the different binding sites of
magnesium and MgrB on PhoQ. Magnesium was suggested to bind to several locations,
including the acidic cluster near the inner membrane (148-152) and residues G93, W97, H120,
and T156 [14, 23, 40]. Mutations of these residues were shown to abolish magnesium
protection from iron-mediated protein cleavage [40]. The majority of these residues are located
in the peripheral of the periplasmic domain, distinct from the MgrB binding site.
430
Our crosslinking results revealed extensive interactions between MgrB and the PhoQ linker
region, implying its importance in functional regulations. Highly conserved and functionally

important residues are found in or near this region, such as K186, T48, R50, E184, and L185,
mutations of which lead to alterations in PhoQ activity as well as signaling capability [20, 23, 24].
Curiously, the wild-type MgrB maintained its inhibition on PhoQ linker variants, suggesting that
the flexible MgrB loop may have various interaction modes with the PhoQ linker region.
Interestingly, the PhoQ activating protein, SafA, appears to act near this linker region in terms of
the putative SafA binding site, and the activation required inter-monomeric salt bridge (R50'-
D179) [41, 42]. Whether MgrB repression competes with SafA activation of PhoQ and how these
regulations are coordinated await further studies.
441
Sensor kinases in bacterial two-component systems are evolutionarily related and have
structural resemblance. We found that the predicted structures of EnvZ, QseC, and CreC sensor
kinases closely resemble the PhoQ structure. EnvZ was also shown to form a complex with MgrB
experimentally [30]. With the mechanism of MgrB revealed here, it is foreseeable that novel
small protein regulators can be designed for PhoQ, EnvZ, and possibly many other sensor
kinases. This will contribute to the repertoire of new antimicrobial agents targeting sensor
kinases that regulate bacterial virulence [43] and to the treatment of infections caused by drug-
resistant pathogens.
450
451 **Materials and methods**
452
Bacterial strains, plasmids, and growth conditions
The complete lists of strains, plasmids, and primers used in this study are summarized in Table
S1, S2, and S3, respectively. The *E. coli* MG1655 *ΔmgrB* and *ΔphoQΔmgrB* strains were from the
previous study [30]. All strains were grown in the lysogeny broth (LB) medium or minimal A
medium [37] containing 0.2% glycerol, 0.1% casamino acids supplemented with magnesium as
indicated. Cells were grown at 37 °C with shaking unless otherwise indicated. Genes in pTrc99a
vector were induced with isopropyl β-D-1-thiogalactopyranoside (IPTG) at a final concentration
of 10 μM. Genes in pBAD vector were induced with 0.008% arabinose. Genes in pEVOL-pBpF
vector were induced with 0.02% arabinose. Genes in pETduet-1 vector were induced with IPTG
at a final concentration of 0.5 mM. When appropriate, the antibiotics ampicillin, kanamycin, and
chloramphenicol were used at concentrations of 100 μg/ml, 50 μg/ml, and 34 μg/ml,
respectively. A detailed description of cloning and plasmid construction can be found in the SI
Materials and Methods.
466
Disulfide crosslinking *in vivo*
The *in vivo* disulfide-crosslinking experiments were performed as previously described with
modifications [44]. Briefly, cells carrying pBAD33 *phoQ* variant and pTrc99A *mgrB* variant were
grown at 37 °C in LB medium supplemented with 10 mM $MgSO_4$ overnight, then diluted 1:100 to
fresh LB medium supplemented with 1 mM $MgSO_4$. The expression of plasmid-encoded proteins
was induced with arabinose and IPTG. When the OD_{600} value reached 0.6, cells were harvested,
washed, and resuspended in the crosslinking buffer (137 mM NaCl, 2.7 mM KCl, 10 mM
Na_2HPO_4, 1.8 mM KH_2PO_4, 1 mM $MgSO_4$, and 0.1 mM Cu-phenanthroline), followed by 10
minutes incubation at 25 °C. The reaction was stopped by pelleting the cells and then
resuspending in the 1x stop-solution (8 mM sodium phosphate buffer, pH 7.8, 12.5 mM EDTA,
and 12.5 mM N-ethylmaleimide) and incubation at 25 °C for 10 minutes. Cells were collected,
resuspended in the lysis buffer (50 mM Tris-HCl, pH 8.0, 100 mM NaCl, and protease inhibitors),
mixed with lysing matrix B beads (MP Biomedicals), and broken with a homogenizer following
the manufacturer's instructions. The mixture was spun down at 2,300 x g for 10 minutes at 4 °C

to pellet the beads and cell debris. The supernatant was centrifuged at 21,000 x g for one hour
at 4 °C to pellet the membrane fraction, which was then dissolved in the SDS sample buffer at 4
°C overnight. The membrane proteins were separated on an SDS tris-tricine or tris-glycine
polyacrylamide gel via electrophoresis and analyzed with Western blot.
485
Western blot analysis
Proteins were transferred from a polyacrylamide gel to a polyvinylidene fluoride membrane or
nitrocellulose membrane. PhoQ-His and Flag-MgrB were detected with the anti-His and anti-Flag
primary antibody (Sigma), respectively. The IRDye 800CW-conjugated secondary antibody (LI-
COR) was used, and the protein bands were visualized with the Odyssey CLx imaging system (LI-
COR) and quantified with ImageJ. The crosslinking efficiency was calculated by dividing the
amount of PhoQ/MgrB complex with the total amount of PhoQ/MgrB complex and MgrB
monomer.
494
Functional assay with the *gfp* reporter gene
The GFP reporter assay was performed as a previous study [12]. Briefly, the reporter plasmid
pUA66 P_{mgtLA}-*gfp* was transformed to strains indicated. The resulting transformants were grown
overnight at 37 °C in LB medium with 10 mM $MgSO_4$, then diluted 1:100 to fresh LB medium
with magnesium and/or stimuli as indicated. The cultures were grown at 37 °C with vigorous
shaking till the early log phase (OD=0.4-0.5). The fluorescence of cells was monitored with a BD
LSRFortessa SORP flow cytometer (BD Biosciences) equipped with a 100-mW 488-nm laser and a
510/20 bandpass filter. BD FACSDiva software version 8.0 (BD Biosciences) was used to analyze
the acquired data.
504
Unnatural amino acid incorporation in MgrB and site-specific UV crosslinking *in vivo*
The site-specific photo-crosslinking experiments were performed as previous studies with
modifications [35, 45]. *E. coli* BL21 DE3 cells carrying pETduet-1 *phoQ-his_6 flag-mgrB* and pEVOL-
pBpF vectors were grown in LB medium at 37 °C overnight, then diluted 1:100 to fresh LB
medium containing 0.5 mM *p*Bpa. When the OD_{600} value reached around 0.6, cells were induced
with arabinose (0.02%) and IPTG (0.5 mM) for 4h at 30 °C, followed by harvesting, washing two
times, and resuspension in 1 ml cold PBS buffer. The bacterial suspension was spread out onto a
35 mm round dish on ice and irradiated for 5 minutes with 365 nm UV light using an LED UV
hand lamp (Honle UV technology). Cells were then harvested at 4 °C, and the formation of
crosslinked PhoQ/MgrB complex was then detected by the separation of total protein with
polyacrylamide gel electrophoresis and Western blot with anti-Flag antibody. The crosslinked
complex was purified by affinity chromatography, separated in an SDS polyacrylamide gel, and
verified by mass spectrometry (see SI Materials and Methods for details).
518
Molecular Modeling and Molecular Dynamics simulations
The PhoQ/MgrB complex was modeled using Alphafold2 multimer on Colab [PMID: 35637307].
We considered six different complexes: i. PhoQ, ii. PhoQ / MgrB (predictions ranked 1 to 3), iii.
PhoQ / MgrB W20Y variant, and iv. PhoQ / MgrB W20A variant. Each complex was inserted into
a bacterial membrane mimetic composed of a mixture of phosphatidylethanolamine (POPE) and
phosphatidylglycerol (POPG) with a 3:1 ratio. The systems were solvated with a 17 Å water
padding and neutralized with NaCl at a concentration of 150 mM. Each system was assembled
using the CHARMM-GUI webserver [46].
527

The simulation was performed with the CHARMM36m force field, including CMAP corrections
for the protein [47]. The water molecules were described with the TIP3P water parameterization
[48]. The simulations were carried out with OPENMM molecular engine [49] following the
minimization and equilibration protocols provided by CHARMM-GUI. The cutoff for non-bonded
interactions was set to 12 Å with a switching distance at 10 Å. The periodic electrostatic
interactions were computed using particle-mesh Ewald (PME) summation with a grid spacing
smaller than 1 Å. Constant temperature of 310 K was imposed by Langevin dynamics with a
damping coefficient of 1.0 ps. Constant pressure of 1 atm was maintained with Monte Carlo
barostat [50]. The hydrogen mass repartitioning scheme was used to achieve a 4 fs time-step
[51]. Each simulation was carried out up to 1 μs and up to 500 ns for the second and third
ranked predictions of PhoQ / MgrB.
539

540   Acceptor photobleaching FRET analysis
FRET measurements were performed as described before [30]. The expression of PhoQ-
mNeonGreen and mCherry-MgrB was induced in *E. coli* lptD4213 strain with 0.008% arabinose
and 10 μM IPTG, respectively. The cultures were grown to the mid-log phase (OD=0.6) in LB
media. Cells were harvested and washed twice with pre-chilled tethering buffer (20 mM
potassium phosphate pH 7.0, 1 μM methionine, and 10 mM lactic acid). The surface of a glass-
bottom 96-well plate (Greiner) was treated with 0.1% poly-L-lysine for 15 minutes at room
temperature, followed by removing poly-L-Lysine and air-dry for 5 minutes. The wells were
washed twice using a pre-chilled tethering buffer. Subsequently, cells were incubated in the
wells at room temperature for 25 minutes to allow attachment. Unattached cells were removed
by washing with tethering buffer twice, and attached cells were overlaid with tethering buffer
with or without 10 μg/ml C18G and incubated at room temperature for one hour. Acceptor
photobleaching FRET was performed using the same microscope settings previously [30]. The
details of the data acquisition and analysis can be found in the SI Materials and Methods.
554
555

556   Acknowledgments
557   We thank L. Cassidy and A. Throley (the Christian-Albrecht University of Kiel) for performing
558   mass spectrometry analyses, writing and editing the manuscript; H. Koch (the Albert Ludwig
559   University of Freiburg) for pEVOL-pBpF vector and the site-directed photo-crosslinking protocol;
560   M. Stopp for the disulfide crosslinking protocol; S. G. Sierra for technical assistance; members of
561   the Yuan and the Sourjik laboratories for helpful discussions.
562

563   We thank G. Storz (National Institutes of Health), V. Sourjik (Max Planck Institute for Terrestrial
564   Microbiology), R. Lill (Philipps University of Marburg), and S. Yadavalli (Rutgers University) for
565   their suggestions to improve the manuscript.
566

567   This work is supported by the German Research Foundation (DFG) priority program 2002 YU
568   247/3-1 and the Max Planck Society (to J.Y.), the Swiss National Science Foundation (SNSF) grant
569   PCEFP3_194606 (to T.L. and S.L.).
570

571   J.Y. and T.L. designed the experiments; S.J., L.C.S., C.K., S.L., and J.Y. performed the experiments;
572   S.J., G.M., L.C.S., C.K., S.L., T.L., V.S., and J.Y. analyzed data; J.Y., T.L., and S.J. wrote the
573   manuscript.
574

575   References

1. Duval, M. and P. Cossart, *Small bacterial and phagic proteins: an updated view on a rapidly moving field.* Curr Opin Microbiol, 2017. **39**: p. 81-88.

2. Ramamurthi, K.S. and G. Storz, *The small protein floodgates are opening; now the functional analysis begins.* BMC Biol, 2014. **12**: p. 96.

3. Su, M., et al., *Small proteins: untapped area of potential biological importance.* Front Genet, 2013. **4**: p. 286.

4. Weidenbach, K., et al., *Small Proteins in Archaea, a Mainly Unexplored World.* J Bacteriol, 2022. **204**(1): p. e0031321.

5. Sberro, H., et al., *Large-Scale Analyses of Human Microbiomes Reveal Thousands of Small, Novel Genes.* Cell, 2019. **178**(5): p. 1245-1259 e14.

6. Hemm, M.R., J. Weaver, and G. Storz, *Escherichia coli Small Proteome.* EcoSal Plus, 2020. **9**(1).

7. Yadavalli, S.S. and J. Yuan, *Bacterial Small Membrane Proteins: the Swiss Army Knife of Regulators at the Lipid Bilayer.* J Bacteriol, 2022. **204**(1): p. e0034421.

8. Groisman, E.A., A. Duprey, and J. Choi, *How the PhoP/PhoQ System Controls Virulence and Mg(2+) Homeostasis: Lessons in Signal Transduction, Pathogenesis, Physiology, and Evolution.* Microbiol Mol Biol Rev, 2021. **85**(3): p. e0017620.

9. Bader, M.W., et al., *Recognition of antimicrobial peptides by a bacterial sensor kinase.* Cell, 2005. **122**(3): p. 461-72.

10. Prost, L.R., et al., *Activation of the bacterial sensor kinase PhoQ by acidic pH.* Mol Cell, 2007. **26**(2): p. 165-74.

11. Choi, J. and E.A. Groisman, *Acidic pH sensing in the bacterial cytoplasm is required for Salmonella virulence.* Mol Microbiol, 2016. **101**(6): p. 1024-38.

12. Yuan, J., et al., *Osmosensing by the bacterial PhoQ/PhoP two-component system.* Proc Natl Acad Sci U S A, 2017. **114**(50): p. E10792-E10798.

13. Carabajal, M.A., et al., *PhoQ is an unsaturated fatty acid receptor that fine-tunes Salmonella pathogenic traits.* Sci Signal, 2020. **13**(628).

14. Garcia Vescovi, E., F.C. Soncini, and E.A. Groisman, *Mg^{2+} as an extracellular signal: environmental regulation of Salmonella virulence.* Cell, 1996. **84**(1): p. 165-74.

15. Lippa, A.M. and M. Goulian, *Feedback inhibition in the PhoQ/PhoP signaling system by a membrane peptide.* PLoS Genet, 2009. **5**(12): p. e1000788.

16. Eguchi, Y., et al., *B1500, a small membrane protein, connects the two-component systems EvgS/EvgA and PhoQ/PhoP in Escherichia coli.* Proceedings of the National Academy of Sciences of the United States of America, 2007. **104**(47): p. 18712-18717.

17. Yadavalli, S.S., et al., *Antimicrobial peptides trigger a division block in Escherichia coli through stimulation of a signalling system.* Nat Commun, 2016. **7**: p. 12340.

18. Eguchi, Y., et al., *Regulation of acid resistance by connectors of two-component signal transduction systems in Escherichia coli.* J Bacteriol, 2011. **193**(5): p. 1222-8.

19. Salvail, H., J. Choi, and E.A. Groisman, *Differential synthesis of novel small protein times Salmonella virulence program.* PLoS Genet, 2022. **18**(3): p. e1010074.

20. Sanowar, S., A. Martel, and H.L. Moual, *Mutational analysis of the residue at position 48 in the Salmonella enterica Serovar Typhimurium PhoQ sensor kinase.* J Bacteriol, 2003. **185**(6): p. 1935-41.

21. Regelmann, A.G., et al., *Mutational analysis of the Escherichia coli PhoQ sensor kinase: differences with the Salmonella enterica serovar Typhimurium PhoQ protein and in the mechanism of Mg2+ and Ca2+ sensing.* J Bacteriol, 2002. **184**(19): p. 5468-78.

22. Minagawa, S., et al., *Isolation and molecular characterization of the locked-on mutant of Mg2+ sensor PhoQ in Escherichia coli.* Biosci Biotechnol Biochem, 2005. **69**(7): p. 1281-7.

23. Cho, U.S., et al., *Metal bridges between the PhoQ sensor domain and the membrane regulate transmembrane signaling.* J Mol Biol, 2006. **356**(5): p. 1193-206.

24. Cheung, J., et al., *Crystal structure of a functional dimer of the PhoQ sensor domain.* J Biol Chem, 2008. **283**(20): p. 13762-70.

25. Waldburger, C.D. and R.T. Sauer, *Signal detection by the PhoQ sensor-transmitter. Characterization of the sensor domain and a response-impaired mutant that identifies ligand-binding determinants.* J Biol Chem, 1996. **271**(43): p. 26630-6.

26. Venturini, E.S., S. L.; Maaß, S.; Gelhausen, R.; Eggenhofer, F.; Li, L.; Cain, A. K.; Parkhill, J.; Becher, D.; Backofen, R.; Barquist, L.; Sharma, C. M.; Westermann, A. J.; Vogel, J., *A global data-driven census of Salmonella small proteins and their potential functions in bacterial virulence.* microLife, 2020. **1**(1).

27. Shein, A.M.S., et al., *High prevalence of mgrB-mediated colistin resistance among carbapenem-resistant Klebsiella pneumoniae is associated with biofilm formation, and can be overcome by colistin-EDTA combination therapy.* Sci Rep, 2022. **12**(1): p. 12939.

28. Salazar, M.E., A.I. Podgornaia, and M.T. Laub, *The small membrane protein MgrB regulates PhoQ bifunctionality to control PhoP target gene expression dynamics.* Mol Microbiol, 2016. **102**(3): p. 430-445.

29. Lippa, A.M. and M. Goulian, *Perturbation of the oxidizing environment of the periplasm stimulates the PhoQ/PhoP system in Escherichia coli.* J Bacteriol, 2012. **194**(6): p. 1457-63.

30. Yadavalli, S.S., et al., *Functional determinants of a small protein controlling a broadly conserved bacterial sensor kinase.* J Bacteriol, 2020.

31. Mirdita, M., et al., *ColabFold: making protein folding accessible to all.* Nat Methods, 2022. **19**(6): p. 679-682.

32. Evans, R., et al., *Protein complex prediction with AlphaFold-Multimer.* bioRxiv, 2022: p. 2021.10.04.463034.

33. Akdel, M., et al., *A structural biology community assessment of AlphaFold2 applications.* Nat Struct Mol Biol, 2022.

34. Ferris, H.U., et al., *Mechanism of regulation of receptor histidine kinases.* Structure, 2012. **20**(1): p. 56-66.

35. Ryu, Y. and P.G. Schultz, *Efficient incorporation of unnatural amino acids into*
proteins in Escherichia coli. Nat Methods, 2006. **3**(4): p. 263-5.
36. Molnar, K.S., et al., *Cys-scanning disulfide crosslinking and bayesian modeling*
probe the transmembrane signaling mechanism of the histidine kinase, PhoQ.
Structure, 2014. **22**(9): p. 1239-1251.
37. Miller, J.H., *A Short Course in Bacterial Genetics - A Laboratory Manual and*
Handbook for Escherichia coli and Related Bacteria. 1992: Cold Spring Harbor
Laboratory Press.
38. Sochacki, K.A., et al., *Real-time attack on single Escherichia coli cells by the*
human antimicrobial peptide LL-37. Proc Natl Acad Sci U S A, 2011. **108**(16): p.
E77-81.
39. Brink, K.R., et al., *An E. coli display method for characterization of peptide-sensor*
kinase interactions. Nat Chem Biol, 2022.
40. Chamnongpol, S., M. Cromie, and E.A. Groisman, *Mg2+ sensing by the Mg2+*
sensor PhoQ of Salmonella enterica. J Mol Biol, 2003. **325**(4): p. 795-807.
41. Eguchi, Y., et al., *The connector SafA interacts with the multi-sensing domain of*
PhoQ in Escherichia coli. Mol Microbiol, 2012. **85**(2): p. 299-313.
42. Yoshitani, K., et al., *Identification of an internal cavity in the PhoQ sensor domain*
for PhoQ activity and SafA-mediated control. Biosci Biotechnol Biochem, 2019.
83(4): p. 684-694.
43. Cabezudo, I., et al., *Effect-Directed Synthesis of PhoP/PhoQ Inhibitors to Regulate*
Salmonella Virulence. J Agric Food Chem, 2022. **70**(22): p. 6755-6763.
44. Stopp, M., et al., *Transmembrane signaling and cytoplasmic signal conversion by*
dimeric transmembrane helix 2 and a linker domain of the DcuS sensor kinase. J
Biol Chem, 2021. **296**: p. 100148.
45. Kuhn, P., et al., *Ribosome binding induces repositioning of the signal recognition*
particle receptor on the translocon. J Cell Biol, 2015. **211**(1): p. 91-104.
46. Jo, S., et al., *CHARMM-GUI: a web-based graphical user interface for CHARMM.* J
Comput Chem, 2008. **29**(11): p. 1859-65.
47. Best, R.B., et al., *Optimization of the additive CHARMM all-atom protein force*
field targeting improved sampling of the backbone phi, psi and side-chain chi(1)
and chi(2) dihedral angles. J Chem Theory Comput, 2012. **8**(9): p. 3257-3273.
48. Jorgensen, W.L., Chandrasekhar, J., and Madura, J. D., *Comparison of simple*
potential functions for simulating liquid water. J Chem Phys, 1983. **79**(2).
49. Eastman, P., et al., *OpenMM 7: Rapid development of high performance*
algorithms for molecular dynamics. PLoS Comput Biol, 2017. **13**(7): p. e1005659.
50. Chow, K.a.F., D. M. , *Isothermal-isobaric molecular dynamics simulations with*
Monte Carlo volume sampling. Comp. Phys. Comm., 1995. **91**(1-3): p. 283-289.
51. Balusek, C., et al., *Accelerating Membrane Simulations with Hydrogen Mass*
Repartitioning. J Chem Theory Comput, 2019. **15**(8): p. 4673-4686.
703
704

Figures and tables

Fig. 1. Identification of the interacting residues between MgrB and PhoQ in the transmembrane helices. (**A**) The sequence of MgrB from *E. coli*. Residues predicted to be in the transmembrane helix are underlined. (**B**) The tertiary structure of the top ranked PhoQ/MgrB complex predicted by Alphafold2. The two PhoQ protomers are colored in wheat and teal. MgrB is colored in magenta. (**C**) The structural models of MgrB predicted by Alphafold2 are shown in ribbon. The model of MgrB alone (left) is colored in rainbow spectrum with the N-terminus in blue and the C-terminus in red. The positions involved in crosslinking (20, 22 and 24) are in

purple and shown with side chains in ball-and-stick. The predicted structure of MgrB in complex
(right) is colored in magenta. All structural figures were prepared using PyMOL unless otherwise
stated. (**D, E top, F top**) Western blot analysis of *E. coli* membrane extracts after disulfide
crosslinking. The indicated PhoQ and FLAG-tagged MgrB variants were expressed in *E. coli*,
followed by Cu-phenanthroline catalyzed disulfide crosslinking (details in Materials and
Methods). The total protein stain of the PVDF membranes serves as loading controls. Residues
from PhoQ TM1 and TM2 are colored in wheat and teal, respectively. Magenta indicates the
residues in the MgrB TM helix. Data are representative of at least three independent
experiments**. (D bottom, F bottom**) The functional assay of PhoQ Cys-variants with or without
MgrB. An GFP reporter plasmid carrying the *gfp* gene fused with a PhoP regulated promoter
P$_{mgtLA}$ was introduced to the *E. coli* strain. The expression of *phoQ* and *mgrB* variants was
induced and cells were grown to the early log phase. The function of PhoQ variants was then
monitored by measuring the GFP fluorescence of the cell culture. Each data point is shown with
the calculated average and standard deviation from three independent biological replicates.
730

Fig. 2. Site-specific photo crosslinking in the periplasmic domain. (**A**) Western blot analysis of
the site-specific photo crosslinking between PhoQ and MgrB. The *E. coli ΔphoQΔmgrB* strain
harboring pETduet-1 *phoQ-his₆ flag-mgrB* and pEVOL-pBpF vectors were grown in LB containing
the unnatural amino acid, *p*Bpa, which was incorporated into the FLAG tagged MgrB at the
indicated positions via the amber stop codon and the cognate tRNA/aminoacyl-tRNA synthetase
pair. Cells were treated with UV light. The crosslinked MgrB-PhoQ complex was detected with
Western blot using antibodies directed to the FLAG epitopes. Samples without UV irradiation
served as controls. (**B**) Western blot analysis of disulfide crosslinking from MgrB residue 31 to
the PhoQ linker region connecting the periplasmic domain to the TM2 helix. The experimental
procedure is as in Fig. 1D and details can be found in Materials and Methods. Data are
representative of at least three independent experiments.

747
748
Fig. 3. The conformational change of the PhoQ sensor kinase upon MgrB binding. (A) Western
blot analysis of disulfide crosslinking within the PhoQ molecule in the presence or absence of
MgrB. The His$_6$-tagged PhoQ cysteine variants were expressed with or without the wild type
MgrB in *E. coli*. The disulfide crosslinking reaction was initiated with Cu-phenanthroline. The
resulting crosslinked PhoQ species was then detected using Western blot with an anti-His
antibody. The red asterisk indicates PhoQ molecules with an intramolecular disulfide crosslink.
The total protein stain serves as a loading control. Data are representative of three independent
experiments. **(B)** A diagram of MgrB affecting the disulfide crosslinking between the cysteine
residues at position 43 and 188. Brown and cyan circles represent the TM helices in each PhoQ
monomer. The magenta circle represents MgrB. Orange arrows indicate the crosslinking
tendency of Cys188 to Cys43 in the same or different PhoQ monomer. **(C)** MgrB induced PhoQ
conformational change along the PhoQ dimer interface. The His$_6$-tagged PhoQ cysteine variants
were expressed with or without the wild type MgrB in *E. coli*. The experimental procedure is as
described in A and details can be found in Materials and Methods. The position of cysteine is
indicated as residue numbers in the PhoQ N-terminal helix. Numbers colored in blue and purple
are residues in TM and periplasmic domains, respectively. Data are representative of three
independent experiments. **(D)** The repression of PhoQ activity by high magnesium and MgrB.
The activity of PhoQ was monitored with the GFP reporter system using the *E. coli ΔphoQΔmgrB*
strain as described in Fig. 1E. Data are representative of three independent experiments. Each
data point is shown with the calculated average and standard deviation from three biological
replicates.
770
771
772
773

774
Fig. 4. Molecular dynamics simulations of PhoQ/MgrB complex models. (**A**) Projection of the
molecular dynamics simulations into the eigenspace formed by the two first components of a
Principal Component Analysis (PCA) of the transmembrane domain of PhoQ (residues 21 – 38
and 194 – 215). Each point corresponds to a conformation of the TM and is colored in shades of
teal, purple, red and blue, for the simulations of PhoQ alone, and in complex with MgrB, MrgB
W20Y and W20A, respectively. Lighter shades correspond to conformations sampled earlier in
the simulation. (**B**) Distribution in distances between TM1 – TM1' (left) and TM2 – TM2' (right)
at the top, middle, and bottom of PhoQ TM domain. (**C**) The overlay of the final conformations
of PhoQ/MgrB complex models with PhoQ dimer. PhoQ structures are colored as in A. The MgrB
wild-type, W20A, and W20Y variants are colored in pink, light blue, and light red, respectively.

785
786
Fig. 5. The impact of MgrB on PhoQ sensing external stimuli. (**A**) The PhoQ activation in the
presence or absence of MgrB under high magnesium conditions. *E. coli* MG1655 *ΔmgrB* strain
harboring pBAD33 with (left panel) or without (right panel) *mgrB* gene, were induced with
arabinose and grown in the presence of indicated stimuli in LB media supplemented with 10 mM
magnesium. The activity of PhoQ was monitored using the GFP reporter as described in Fig. 1E.
(**B**) The activation of PhoQ in the presence of antimicrobial peptides and MgrB under 1 mM

magnesium conditions. The orange lines indicate the fold change of PhoQ activity in the
presence of antimicrobial peptides. *E. coli* MG1655 *ΔmgrB* strain expressing MgrB variants were
grown in LB supplemented with 1 mM magnesium and antimicrobial peptides, LL37 (5 µg/ml) or
C18G (5 µg/ml) as indicated. The PhoQ activity was followed as in A. (**C**) The FRET measurements
of PhoQ-mNeonGreen and mCherry-MgrB *in vivo*. *E. coli* cells expressing PhoQ-mNeonGreen
and mCherry-MgrB from plasmids were grown to mid-log phase, then treated with or without
the antimicrobial peptide, C18G (10 µg/ml). The energy transfer from mNeonGreen to mCherry
was then measured for each condition. The strains expressing no FRET receptor (mCherry-MgrB)
serve as negative controls. (**D**) The activation of PhoQ variants in the presence of MgrB and
C18G (5 µg/ml). The growth and induction conditions for the PhoQ and MgrB variants were the
same as described before (Fig. 1F bottom). The PhoQ activity was monitored by measuring the
GFP reporter fluorescence. (**E**) Western blot analysis of PhoQ I38C MgrB 16A24C disulfide
crosslinking in the presence or absence of C18G (5 µg/ml). The crosslinking and Western blot
experiments were performed as described before (Fig. 1F top). The total protein stain of the
membrane serves as a loading control. Data are representative of three independent
experiments. Each data point in A-D is shown with the calculated average and standard
deviation from at least three independent experiments.
810

811
812
Fig. 6. The schematic representation of MgrB-mediated PhoQ activation by antimicrobial
peptides. PhoQ is active under physiological magnesium conditions, inducing the expression of
mgrB. The expressed MgrB binds to PhoQ, resulting in a partially repressed state. Antimicrobial
peptides (AMPs) enter bacterial periplasm, bind to PhoQ, and cause MgrB to dissociate from
PhoQ.

Table 1. The Cα - Cα distance of crosslinked residue pairs in the PhoQ/MgrB structural model.

Cα - Cα distance (Å)	W20-Y32	W20-L204	Q22-Y197	F24-I38
model 1	12.10	12.80	8.30	9.10
model 2	12.90	11.90	9.60	10.80
model 3	12.90	10.10	10.00	11.30
model 4	13.60	11.90	10.20	11.40
model 5	14.20	16.40	16.20	11.60

820
821

     Supporting Information
SI Materials and Methods

Cloning and plasmid construction
E. coli phoQ-his$_6$ gene was cloned into pET-Duet-1 vector at NcoI/BamHI restriction sites, and *E.*
coli flag-mgrB gene was cloned into pET-Duet-1 vector at NdeI/ EcoRV restriction sites. The
pEVOL-pBpF vector is a gift from Dr. Hans-Georg Koch (Freiburg, Germany). Mutations in *phoQ*
and *mgrB* were generated using the Q5 site-directed mutagenesis kit (New England BioLabs)
following the manufacturer's instruction. All constructs were verified by DNA sequencing.

Purification of the crosslinked PhoQ-His/FLAG-MgrB complex
A two-liter culture was prepared for the purification of the crosslinked PhoQ-His/FLAG-MgrB
complex. After UV crosslinking, cells were harvested, resuspended in the resuspension buffer
(20 mM Tris-HCl, pH 7.5, 300 mM NaCl, 0.1 mM PMSF), then lysed with LM10 microfluidizer at 4
°C. The lysate was centrifuged at 11,000 for 10 min to remove cell debris. The supernatant was
collected and centrifuged at 13,500 g for 2 hours to pellet the total membrane, which was then
dissolved in the resuspension buffer containing 1% (wt/vol) n-dodecyl-β-D-maltopyranoside
(DDM) with gentle shaking at 4 °C overnight. Insoluble fraction was spun down by centrifugation
at 21,000 g for 45 min. The supernatant was loaded onto a TALON column (Cytiva), which was
then washed three times with buffer containing 50 mM Tris-HCl, pH 8.0, 500 mM NaCl, 10%
glycerol, 10 mM imidazole, 0.1 mM PMSF and 0.03% DDM. The protein was eluted with elution
buffer containing 50 mM Tris-HCl, pH 8.0, 500 mM NaCl, 10% glycerol, 200 mM imidazole, 0.1
mM PMSF and 0.03% DDM, and concentrated to ~200 μL. The elution was then incubated with
anti-FLAG M2 magnetic beads (Sigma) at 4 °C overnight. The magnetic beads were washed with
Tris buffered saline, and then incubated with 3 X FLAG peptide solution (Sigma) at room
temperature for one hour to elute FLAG-tagged proteins. The supernatant was collected,
concentrated to ~50 μl, and analyzed using gel electrophoresis with a 7.5% tris-glycine
polyacrylamide gel.

Protein verification with mass spectrometry
Crosslinked and control bands were excised from SDS-PAGE gels and de-stained and sliced into
approx. 1mm^3 pieces. Proteins in the gel pieces were reduced (10 mM dithiothreitol, 56°C for 30
min), and alkylated (50 mM chloroacetamide, 30 min, RT, in the dark), prior to overnight
enzymatic hydrolysis with chymotrypsin (Promega, WI, USA) (0.1 μg of chymotrypsin per band in
50 mM ammonium bicarbonate buffer pH 7.4, 2 mM CaCl2) at 37°C. Following overnight
digestion, the peptides were collected and dried down via vacuum centrifugation (Concentrator
Plus, Eppendorf). Peptides were resuspended in 10 μl of HPLC loading buffer (3 % ACN + 0.1 %
trifluoroacetic acid) prior to LC-MS/MS analysis.

Chromatographic separation was performed on a Dionex U3000 nanoHPLC system (Thermo,
Germany) equipped with an Acclaim pepmap100 C18 column (2 μm particle size, 75 μm × 500
mm) coupled online to a QExactive Plus mass spectrometer (Thermo, Bremen). The separation
was performed over a 60-minute run: 5 % B for 3 minutes followed by linear gradients from 5 %
to 50 % B over 30 minutes, then 50 % to 90 % over 1 minute, and 10 minutes at 90 % B. Inter-run
equilibration was achieved by 15 minutes at 5 % B. Eluent A: 0.05 % formic acid (FA), eluent B:
80 % ACN + 0.04 % FA. A flow rate of 300 nl/min was used and 1 μl of sample was injected per
run. For crosslink samples a second injection of 6 μl was also performed. Full scan MS
acquisition was performed (300-1500 m/z, resolution 70,000, AGC target 3e6, max injection

time (IT) 100 ms) with subsequent data dependent MS/MS of the top 10 most intense ions via
HCD ion activation at NCE 27 (resolution 17,500, AGC target 1e5, isolation window 1.6 m/z, max
IT 50 ms); dynamic exclusion (20 s duration) and lock mass at 445.12003 m/z were enabled.
873
For protein identifications, the MS data files were processed with the Proteome Discoverer™
software suite (Ver. 2.5.0.400) (Thermo, Germany). Raw files were searched using the
SequestHT algorithmand Target Decoy PSM validator node, against a combined database that
included the proteins of interest (PhoQ-Histagged, the modified MgrB-pBpA incorporation
variants), the UniProt canonical proteins for *Escherichia coli* (strain K12) (accessed: 2022.03.10),
and common laboratory contaminants (cRAP list). Searches were performed with full
chymotrypsin specificity, and a maximum 3 missed cleavages. Precursor mass tolerance 10 ppm.
Fragment mass tolerance 0.02 Da. Fixed modification of carbamidomethyl (Cys), dynamic
modification oxidation (Met). Strict parsimony criteria were applied: Target FDR <1% was
applied, and at least one unique high confidence peptide was required for identification. For the
small protein MgrB, manual assessment of the peptide spectral matches was also performed.
885
Acceptor photobleaching FRET data acquisition and analysis
For each well, we acquired three or more sequences of images on isolated fields of view with
the following protocol: (a) 2 images in the acceptor channel; (b) 140 images in the donor
channel followed by (c) 16 seconds acceptor photobleaching (no image acquisition); (d) 50
images in the donor channel; and then (e) 2 images in the acceptor channel. FRET efficiency was
calculated as the increase of donor fluorescence signal upon acceptor photobleaching divided by
the total donor signal after acceptor photobleaching. In order to correct for the donor
photobleaching present during steps b to d, we performed linear fitting (RStudio) of the donor
fluorescence signal versus time for both pre- and postbleaching curves.
895
896

897   **SI Figures and Tables**
898

899
900
Fig. S1. Structural models predicted by Alphafold2. (A-B) The predicted PhoQ monomer and
dimer are colored according to the pLDDT score to show the regions with lower prediction
confidence. Red indicates highest confidence and blue indicates lowest confidence. The four
predicted PhoQ/MgrB complex models ranked from two to five by Alphafold2. Models are
shown with a PhoQ dimer (colored in wheat and deep teal) and four MgrB molecules overlayed
(**C**) or separated (**D**) in green, yellow, pink and grey ranked from 2 to 5, respectively. All
structural figures were prepared using PyMOL unless otherwise stated.
908
909

910
911
Fig. S2. The function of MgrB variants and the crosslinking with PhoQ from MgrB position
20. The activity analysis of MgrB W20 variants (**A**) and MgrB C16A (**B**) using the reporter plasmid

pUA66 P$_{mgtLA}$-*gfp*. The *E. coli ΔmgrB* strain harboring the reporter plasmid and the pBAD plasmid
with *mgrB* variants were grown to the early log phase in LB media supplemented with 0.008%
arabinose. GFP fluorescence in cells was then measured by flow cytometry. (**C**) Sequence
alignment of MgrB homologs from fourteen bacterial species. The conservation of the amino
acid at position 20 (*E. coli* numbering) is highlighted with a red rectangle. (**D**) The functional
assay of MgrB variants expressed from pTrc99A vector. The pUA66 P$_{mgtLA}$-*gfp*, pBAD33 *phoQ*,
and pTrc99A plasmid with flag tagged *mgrB* variants, were transformed into *E.*
coli ΔphoQΔmgrB cells. The resulting transformants were grown overnight at 37 °C in LB
medium supplemented with 10 mM MgSO4, then diluted 1:100 to fresh LB medium
supplemented with 1 mM MgSO4, 0.008% arabinose, 10 µM IPTG, and antibiotics when
appropriate. The cultures were grown at 37 °C with vigorous shaking till the early log phase. The
GFP fluorescence of cells was then monitored by flow cytometry. (**E**) The functional assay of
MgrB and Flag-MgrB expressed from pTrc99A vector. The pUA66 P$_{mgtLA}$-*gfp* and pTrc99A plasmid
with *mgrB* or *flag-mgrB*, were transformed into *E. coli ΔmgrB* cells. The resulting transformants
were grown overnight at 37 °C in LB medium supplemented with 10 mM MgSO4, then diluted
1:100 to fresh LB medium supplemented with 1 mM MgSO4, 10 µM IPTG, and antibiotics when
appropriate. The cultures were grown at 37 °C with vigorous shaking till the early log phase. The
GFP fluorescence of cells was then monitored by flow cytometry. (**F**) The functional assay of the
indicated PhoQ and MgrB variants using the same protocol as in D. (**G**) Western blot analysis
of *E. coli* membrane extracts after disulfide crosslinking. The indicated PhoQ and FLAG-tagged
MgrB variants were expressed in *E. coli*, followed by Cu-phenanthroline catalyzed disulfide
crosslinking (details in Materials and Methods). The total protein stain of the PVDF membranes
serves as loading controls. Data are representative of at least three independent experiments. In
A, B, D, E and F, each data point is shown with the calculated average and standard deviation
from three independent biological replicates.
939
940

946
947
Fig. S3. The full Western blot results of the figures in the main text. (**A**) is the full blot of Fig.
1D. (**B**) is the full blot of Fig. 1E. (**C**) is the full blot of Fig. 1F. (**D, E**) are the full blots of Fig. 2B.
Data are representative of at least three independent experiments. (**F**) Western blot analysis of
the expression level of PhoQ variants. The indicated PhoQ-His variants were expressed in *E. coli*
by inducing with 0.008% arabinose. Cells were harvested in the mid log phase and the total
membrane was extracted for the western blot analysis. The total protein stain of the PVDF
membranes serves as a loading control. Data are representative of two independent
experiments.
956

957
Fig. S4. Mapping the binding interface of MgrB on PhoQ by using site-specific UV crosslinking
in *vivo*. (**A**) *E. coli* BL21 DE3 cells expressing PhoQ-His$_6$, FLAG-MgrB or *p*Bpa-containing FLAG-
MgrB variants were irradiated with or without UV light. The cells were lysed in SDS sample
buffer. Proteins were separated on 7.5% tris-glycine polyacrylamide gels and transfer to PVDF
membranes. FLAG tagged proteins were detected with an anti-FLAG primary antibody and an
IRDye 800CW-conjugated secondary antibody. Data are representative of three independent
experiments. (**B**) Purification of PhoQ-His/FLAG-MgrB crosslinked complex. Elution from a two-
step affinity purification was separated on 7.5% tris-glycine gels. The protein bands in the gel
were visualized with the ready-blue protein stain. Data are representative of two independent
experiments.
968

969
Fig. S5. The functional assay of PhoQ variants with or without MgrB. *E. coli* MG1655
ΔphoQΔmgrB cells harboring pBAD33 *phoQ* variants, the GFP reporter plasmid pUA66 P*mgtA*-*gfp*,
pTrc99A with or without *mgrB* variants were grown in LB supplemented with 0.008% arabinose
and 10 µM IPTG till the early-log phase. The GFP fluorescence of the cells was measured with
flow cytometry. Each data point is shown with the calculated average and standard deviation
from three biological replicates. The results are the representative of at least three independent
experiments.
977

Fig. S6. The conformational change of PhoQ linker region in the presence of MgrB. (**A**) The structural superimposition of the periplasmic and TM domains of the PhoQ dimer (wheat and deep teal) and monomer (pLDDT) is shown. The periplasmic domain was used for structural alignment in Pymol. (**B**) Zoom in view of the linker region of PhoQ. Residue 43 is shown in ball-and-stick. Residues 187 and 188 are colored in purple in both monomer and dimer structures. (**C**) Western blot analysis of disulfide crosslinking within the PhoQ molecule in the presence or absence of MgrB. The crosslinked PhoQ species was detected with an anti-His antibody. Data are representative of at least three independent experiments.

988
Fig. S7. **PhoQ_Cys variants in the presence or absence of MgrB.** (**A-B**) Quantification of the
disulfide crosslinking efficiency of PhoQ_Cys variants in the presence (blue curve) and absence
(black curve) of MgrB. The PhoQ dimer and monomer bands in Fig. 4C were quantified with
ImageJ. The fraction of crosslinked dimer was calculated using the formula: crosslinking fraction
= dimer / (dimer + monomer). (**C-E**) Functional assays of PhoQ_Cys variants with (blue) and
without (purple) MgrB. *E. coli ΔphoQΔmgrB* cells harboring pUA66 P*mgtA*-*gfp*, pBAD33 *phoQ*
variants, and pTrc99A empty or pTrc99A *mgrB* were grown overnight at 37 °C in LB medium
supplemented with 10 mM MgSO$_4$. Cultures were then diluted 1:100 to fresh LB medium
supplemented with 1 mM MgSO$_4$, 0.008% arabinose and 10 µM IPTG, and grown at 37 °C with
vigorous shaking till the early log phase (OD=0.4-0.5). The fluorescence of cells was monitored
with flow cytometry and normalized to the PhoQ wild type. Each data point in C, D and E is
shown with the calculated average and standard deviation from three biological replicates. The
results are the representative of three independent experiments.
1002

1003
1004
1005 **Fig. S8. MD simulations of PhoQ/MgrB complex models.** Distribution in distances between TM1
1006 – TM1' (left) and TM2 – TM2' (right) at the top, middle, and bottom of PhoQ TM domain.

1007
1008
Fig. S9. The functional assays with fluorescent reporters. PhoQ activity was monitored with a
GFP reporter plasmid (pUA66 P$_{mgtLA}$-*gfp*) with or without MgrB in LB (**A**) and in minimal A
medium (**B**) supplemented with 1 mM magnesium. Breifly, *E. coli ΔmgrB* strain harboring the
GFP reporter plasmid and a pBAD plasmid with or without the wild-type mgrB gene, was grown
in the presence of the indicated PhoQ stimuli to early log phase. The GFP fluorescence of cells
was then measured by flow cytometry. Data are representative of at least three independent
experiments. The function of fusion proteins mCherry-MgrB (**C**) and PhoQ-mneonGreen (**D**)
were anlalyzed with the GFP report and a mCherry reporter (pUA66 P$_{mgtLA}$-*mcherry*). Indicated
deletion strains were complemented with a plasmid harboring the gene of indicated fusion
proteins. The expression of the fusion protein was induced with 10 µM IPTG and 0.008%
arabinose for the pTrc and pBAD plasmids, respectively. Cells were grown to early log phase and
the fluorescence of cells was measured by flow cytometry. Cells transformed with empty
plasmids and plasmids harboring the corresponding wild-type genes were used as negative and
positive controls, respectively. Data are representative of two independent experiments.
1023

1024 **Table S1. List of strains used in this study.**

1025

Strain	Genotype	Source or Reference
JingY11	MG1655	*E. coli* Genetic Stock Center, CGSC# 7740
lptD4213	F-, *lptD4213*, *galK2*(Oc), *λ⁻*, *IN(rrnD-rrnE)1*, *rpsL200*(strR), *rph-1*	Yale Coli Genetic Stock Center
E. coli BL21(DE3)	BL21(DE3)	New England Biolabs
JingY34	MG1655 Δ*mgrB::FRT*	[1]
JingY90	MG1655 Δ*phoQ::FRT* Δ*mgrB::FRT*	[1]

1026
1027

Table S2. List of plasmids used in this study.

Plasmid	Genotype	Source or Reference
pJY1	pBAD33 *phoQ-(his)$_6$*	This study
pJY31	pBAD33RBS	[2]
pJY151	pBAD33RBS *mgrB*	[1]
pOB2	pTrc99A *mcherry*	[1]
pJY569	pTrc99A *mcherry-mgrB*	[1]
pJY573	pBAD33RBS *phoQ-mneongreen*	[1]
pJY317	pTrc99A *flag-mgrB*	This study
pJS1	pET-Duet-1 EC *phoQ-his$_6$, flag-mgrB*	This study
pEVOL	pEVOL-pBpF	A gift from Dr. Hans-Georg Koch
pLS1	pBAD33RBS *phoQ L26C*	This study
pLS2	pBAD33RBS *phoQ V27C*	This study
pLS3	pBAD33RBS *phoQ L28C*	This study
pLS4	pBAD33RBS *phoQ S29C*	This study
pLS5	pBAD33RBS *phoQ L30C*	This study
pLS6	pBAD33RBS *phoQ A31C*	This study
pLS7	pBAD33RBS *phoQ Y32C*	This study
pLS8	pBAD33RBS *phoQ G33C*	This study
pLS9	pBAD33RBS *phoQ M34C*	This study
pLS10	pBAD33RBS *phoQ V35C*	This study
pLS11	pBAD33RBS *phoQ A36C*	This study
pLS12	pBAD33RBS *phoQ L37C*	This study
pLS13	pBAD33RBS *phoQ I38C*	This study

pLS14	pBAD33RBS *phoQ G39C*	This study
pLS15	pBAD33RBS *phoQ Y40C*	This study
pLS16	pBAD33RBS *phoQ S41C*	This study
pLS17	pBAD33RBS *phoQ V42C*	This study
pLS18	pBAD33RBS *phoQ S43C*	This study
pLS19	pBAD33RBS *phoQ F44C*	This study
pJS4	pBAD33RBS *phoQ K46C*	This study
pJS5	pBAD33RBS *phoQ T47C*	This study
pJS6	pBAD33RBS *phoQ T48C*	This study
pJS7	pBAD33RBS *phoQ F49C*	This study
pJS8	pBAD33RBS *phoQ R50C*	This study
pJS9	pBAD33RBS *phoQ L51C*	This study
pJS10	pBAD33RBS *phoQ L52C*	This study
pJS11	pBAD33RBS *phoQ R53C*	This study
pJS12	pBAD33RBS *phoQ G54C*	This study
pJS13	pBAD33RBS *phoQ E55C*	This study
pJS14	pBAD33RBS *phoQ S56C*	This study
pJS15	pBAD33RBS *phoQ N57C*	This study
pJS16	pBAD33RBS *phoQ L58C*	This study
pJS17	pBAD33RBS *phoQ F59C*	This study
pJS18	pBAD33RBS *phoQ Y60C*	This study
pJS19	pBAD33RBS *phoQ T61C*	This study
pJS20	pBAD33RBS *phoQ L62C*	This study
pJS21	pBAD33RBS *phoQ D179C*	This study
pJS22	pBAD33RBS *phoQ T181C*	This study

pJS23	pBAD33RBS *phoQ I183C*	This study
pJS24	pBAD33RBS *phoQ E184C*	This study
pJS25	pBAD33RBS *phoQ L185C*	This study
pJS26	pBAD33RBS *phoQ K186C*	This study
pLS20	pBAD33RBS *phoQ S187C*	This study
pLS21	pBAD33RBS *phoQ S188C*	This study
pLS22	pBAD33RBS *phoQ S43C-S187C*	This study
pLS23	pBAD33RBS *phoQ S43C-S188C*	This study
pCK1	pBAD33RBS *phoQ S193C*	This study
pCK2	pBAD33RBS *phoQ W194C*	This study
pCK3	pBAD33RBS *phoQ F195C*	This study
pCK4	pBAD33RBS *phoQ I196C*	This study
pCK5	pBAD33RBS *phoQ Y197C*	This study
pCK6	pBAD33RBS *phoQ V198C*	This study
pCK7	pBAD33RBS *phoQ L199C*	This study
pCK8	pBAD33RBS *phoQ S200C*	This study
pCK9	pBAD33RBS *phoQ N202C*	This study
pCK10	pBAD33RBS *phoQ L203C*	This study
pCK11	pBAD33RBS *phoQ L204C*	This study
pCK12	pBAD33RBS *phoQ L205C*	This study
pCK13	pBAD33RBS *phoQ V206C*	This study
pCK14	pBAD33RBS *phoQ I207C*	This study
pCK15	pBAD33RBS *phoQ L209C*	This study
pCK16	pBAD33RBS *phoQ L210C*	This study
pCK17	pBAD33RBS *phoQ W211C*	This study

pCK18	pBAD33RBS *phoQ V212C*	This study
pCK19	pBAD33RBS *phoQ A213C*	This study
pCK20	pTrc99A *mgrB C16A*	This study
pCK21	pTrc99A *mgrB C16A W20C*	This study
pCK22	pTrc99A *mgrB C16A Q22C*	This study
pCK23	pTrc99A *mgrB C16A F24C*	This study
pJS27	pTrc99A *mgrB D31C*	This study
pJY268	pBAD33RBS *mgrB W20A*	This study
pJY156	pBAD33RBS *mgrB C28A*	This study
pJS51	pET-Duet-1 *mgrB N25-TAG*	This study
pJS52	pET-Duet-1 *mgrB M26-TAG*	This study
pJS53	pET-Duet-1 *mgrB M27-TAG*	This study
pJS54	pET-Duet-1 *mgrB D29-TAG*	This study
pJS55	pET-Duet-1 *mgrB Q30-TAG*	This study
pJS56	pET-Duet-1 *mgrB D31-TAG*	This study
pJS57	pET-Duet-1 *mgrB V32-TAG*	This study
pJS58	pET-Duet-1 *mgrB Q33-TAG*	This study
pJS59	pET-Duet-1 *mgrB F34-TAG*	This study
pJS60	pET-Duet-1 *mgrB F35-TAG*	This study
pJS61	pET-Duet-1 *mgrB S36-TAG*	This study
pJS62	pET-Duet-1 *mgrB G37-TAG*	This study
pJS63	pET-Duet-1 *mgrB I38-TAG*	This study
pJS64	pET-Duet-1 *mgrB A40-TAG*	This study
pJS65	pET-Duet-1 *mgrB I41-TAG*	This study
pJS66	pET-Duet-1 *mgrB N42-TAG*	This study

pJS67	pET-Duet-1 *mgrB Q43-TAG*	This study
pJS68	pET-Duet-1 *mgrB F44-TAG*	This study
pJS69	pET-Duet-1 *mgrB I45-TAG*	This study
pJS70	pET-Duet-1 *mgrB P46-TAG*	This study
pJS71	pET-Duet-1 *mgrB W47-TAG*	This study

1030
1031

1032   **Table S3. List of primers used in this study.**

1033

Primer	Sequence (5' -> 3')
pBAD33RBS *phoQ L26C* fw	AGCGGTAGTAtgcGTGCTTTCGCTTGC
pBAD33RBS *phoQ L26C* rv	GCCGTTGCCAACAGAAAA
pBAD33RBS *phoQ V27C* fw	GGTAGTACTGtgcCTTTCGCTTGCCTACGGAATG
pBAD33RBS *phoQ V27C* rv	GCTGCCGTTGCCAACAGA
pBAD33RBS *phoQ L28C* fw	AGTACTGGTGtgcTCGCTTGCCTACGGAATGG
pBAD33RBS *phoQ L28C* rv	ACCGCTGCCGTTGCCAAC
pBAD33RBS *phoQ S29C* fw	ACTGGTGCTTtgcCTTGCCTACGGAATGG
pBAD33RBS *phoQ S29C* rv	ACTACCGCTGCCGTTGCC
pBAD33RBS *phoQ L30C* fw	GGTGCTTTCGtgcGCCTACGGAATGG
pBAD33RBS *phoQ L30C* rv	AGTACTACCGCTGCCGTT
pBAD33RBS *phoQ A31C* fw	GCTTTCGCTTtgcTACGGAATGGTCG
pBAD33RBS *phoQ A31C* rv	ACCAGTACTACCGCTGCC
pBAD33RBS *phoQ Y32C* fw	TTCGCTTGCCtgcGGAATGGTCG
pBAD33RBS *phoQ Y32C* rv	AGCACCAGTACTACCGCTG
pBAD33RBS *phoQ G33C* fw	GCTTGCCTACtgcATGGTCGCGC
pBAD33RBS *phoQ G33C* rv	GAAAGCACCAGTACTACCG
pBAD33RBS *phoQ M34C* fw	TGCCTACGGAtgcGTCGCGCTGATC
pBAD33RBS *phoQ M34C* rv	AGCGAAAGCACCAGTACT
pBAD33RBS *phoQ V35C* fw	CTACGGAATGtgcGCGCTGATCG
pBAD33RBS *phoQ V35C* rv	GCAAGCGAAAGCACCAGT
pBAD33RBS *phoQ A36C* fw	CGGAATGGTCtgcCTGATCGGTTATAGCG
pBAD33RBS *phoQ A36C* rv	TAGGCAAGCGAAAGCACC
pBAD33RBS *phoQ L37C* fw	AATGGTCGCGtgcATCGGTTATAGCGTCAGTTTC
pBAD33RBS *phoQ L37C* rv	CCGTAGGCAAGCGAAAGC
pBAD33RBS *phoQ I38C* fw	GGTCGCGCTGtgcGGTTATAGCG
pBAD33RBS *phoQ I38C* rv	ATTCCGTAGGCAAGCGAAAG
pBAD33RBS *phoQ G39C* fw	CGCGCTGATCtgcTATAGCGTCAG 66C
pBAD33RBS *phoQ G39C* rv	ACCATTCCGTAGGCAAGC
pBAD33RBS *phoQ Y40C* fw	GCTGATAGGTtgcAGCGTCAGTTTC
pBAD33RBS *phoQ Y40C* rv	GCGACCATTCCGTAGGCA
pBAD33RBS *phoQ S41C* fw	GATCGGTTATtgcGTCAGTTTCGATAAAAC
pBAD33RBS *phoQ S41C* rv	AGCGCGACCATTCCGTAG
pBAD33RBS *phoQ V42C* fw	CGGTTATAGCtgcAGTTTCGATAAAACTACGTTTC
pBAD33RBS *phoQ V42C* rv	ATCAGCGCGACCATTCCG
pBAD33RBS *phoQ S43C* fw	TTATAGCGTCtgcTTCGATAAAACTACGTTTC
pBAD33RBS *phoQ S43C* rv	CCGATCAGCGCGACCATT
pBAD33RBS *phoQ F44C* fw	TAGCGTCAGTtgcGATAAAACTACG

pBAD33RBS *phoQ F44C* rv	TAACCGATCAGCGCGACC
pBAD33RBS *phoQ K46C* fw	CAGTTTCGATtgcACTACGTTTCGG
pBAD33RBS *phoQ K46C* rv	ACGCTATAACCGATCAGC
pBAD33RBS *phoQ T47C* fw	TTTCGATAAAtgcACGTTTCGGCTGTTAC
pBAD33RBS *phoQ T47C* rv	CTGACGCTATAACCGATC
pBAD33RBS *phoQ T48C* fw	CGATAAAACTtgcTTTCGGCTGTTACGTG
pBAD33RBS *phoQ T48C* rv	AAACTGACGCTATAACCG
pBAD33RBS *phoQ F49C* fw	TAAAACTACGtgcCGGCTGTTAC
pBAD33RBS *phoQ F49C* rv	TCGAAACTGACGCTATAAC
pBAD33RBS *phoQ R50C* fw	AACTACGTTTtgcCTGTTACGTG
pBAD33RBS *phoQ R50C* rv	TTATCGAAACTGACGCTATAAC
pBAD33RBS *phoQ L51C* fw	TACGTTTCGGtgcTTACGTGGCGAG
pBAD33RBS *phoQ L51C* rv	GTTTTATCGAAACTGACG
pBAD33RBS *phoQ L52C* fw	GTTTCGGCTGtgcCGTGGCGAGA
pBAD33RBS *phoQ L52C* rv	GTAGTTTTATCGAAACTGACGC
pBAD33RBS *phoQ R53C* fw	TCGGCTGTTAtgcGGCGAGAGCA
pBAD33RBS *phoQ R53C* rv	AACGTAGTTTTATCGAAACTGACG
pBAD33RBS *phoQ G54C* fw	GCTGTTACGTtgcGAGAGCAATC
pBAD33RBS *phoQ G54C* rv	CGAAACGTAGTTTTATCGAAAC
pBAD33RBS *phoQ E55C* fw	GTTACGTGGCtgcAGCAATCTGTTC
pBAD33RBS *phoQ E55C* rv	AGCCGAAACGTAGTTTTATC
pBAD33RBS *phoQ S56C* fw	ACGTGGCGAGtgcAATCTGTTCT
pBAD33RBS *phoQ S56C* rv	AACAGCCGAAACGTAGTTTTATC
pBAD33RBS *phoQ N57C* fw	TGGCGAGAGCtgcCTGTTCTATACC
pBAD33RBS *phoQ N57C* rv	CGTAACAGCCGAAACGTA
pBAD33RBS *phoQ L58C* fw	CGAGAGCAATtgcTTCTATACCCTTGCGAAGTG
pBAD33RBS *phoQ L58C* rv	CCACGTAACAGCCGAAAC
pBAD33RBS *phoQ F59C* fw	GAGCAATCTGtgcTATACCCTTGCG
pBAD33RBS *phoQ F59C* rv	TCGCCACGTAACAGCCGA
pBAD33RBS *phoQ Y60C* fw	CAATCTGTTCtgcACCCTTGCGAAGTG
pBAD33RBS *phoQ Y60C* rv	CTCTCGCCACGTAACAGC
pBAD33RBS *phoQ T61C* fw	TCTGTTCTATtgcCTTGCGAAGTGGG
pBAD33RBS *phoQ T61C* rv	TTGCTCTCGCCACGTAAC
pBAD33RBS *phoQ L62C* fw	GTTCTATACCtgcGCGAAGTGGGAAAACAATAAG
pBAD33RBS *phoQ L62C* rv	AGATTGCTCTCGCCACGT
pBAD33RBS *phoQ D179C* fw	TGTGGTGGTGtgcACCATTCCGG
pBAD33RBS *phoQ D179C* rv	ATGGTTAATTTTGGCATCC
pBAD33RBS *phoQ T181C* fw	GGTGGATACCtgcCCGGTGGAGC
pBAD33RBS *phoQ T181C* rv	ACCACAATGGTTAATTTTGG
pBAD33RBS *phoQ I183C* fw	TACCATTCCGtgcGAGCTAAAAAGTTCCTATATGG
pBAD33RBS *phoQ I183C* rv	TCCACCACCACAATGGTT

pBAD33RBS *phoQ E184C* fw	CATTCCGGTGtgcCTAAAAAGTTCCTATATG
pBAD33RBS *phoQ E184C* rv	GTATCCACCACCACAATG
pBAD33RBS *phoQ L185C* fw	TCCGGTGGAGtgcAAAAGTTCCTATATG
pBAD33RBS *phoQ L185C* rv	ATGGTATCCACCACCACA
pBAD33RBS *phoQ K186C* fw	GGTGGAGCTAtgcAGTTCCTATATGGTC
pBAD33RBS *phoQ K186C* rv	GGAATGGTATCCACCACC
pBAD33RBS *phoQ S187C* fw	GGAGCTAAAAtgcTCCTATATGGTCTG
pBAD33RBS *phoQ S187C* rv	ACCGGAATGGTATCCACC
pBAD33RBS *phoQ S188C* fw	GCTAAAAGTtgcTATATGGTCTGG
pBAD33RBS *phoQ S188C* rv	TCCACCGGAATGGTATCC
pBAD33RBS *phoQ S193C* fw	TATGGTCTGGtgcTGGTTTATCTATG
pBAD33RBS *phoQ S193C* rv	TAGGAACTTTTTAGCTCCAC
pBAD33RBS *phoQ W194C* fw	GGTCTGGAGCtgcTTTATCTATG
pBAD33RBS *phoQ W194C* rv	ATATAGGAACTTTTTAGCTCC
pBAD33RBS *phoQ F195C* fw	CTGGAGCTGGtgcATCTATGTGC
pBAD33RBS *phoQ F195C* rv	ACCATATAGGAACTTTTTAGC
pBAD33RBS *phoQ I196C* fw	GAGCTGGTTTtgcTATGTGCTCTC
pBAD33RBS *phoQ I196C* rv	CAGACCATATAGGAACTTTTTAG
pBAD33RBS *phoQ Y197C* fw	CTGGTTTATCtgcGTGCTCTCAG
pBAD33RBS *phoQ Y197C* rv	CTCCAGACCATATAGGAAC
pBAD33RBS *phoQ V198C* fw	GTTTATCTATtgcCTCTCAGCCAATCTGC
pBAD33RBS *phoQ V198C* rv	CAGCTCCAGACCATATAG
pBAD33RBS *phoQ L199C* fw	TATCTATGTGtgcTCAGCCAATCTGC
pBAD33RBS *phoQ L199C* rv	AACCAGCTCCAGACCATA
pBAD33RBS *phoQ S200C* fw	CTATGTGCTCtgcGCCAATCTGC
pBAD33RBS *phoQ S200C* rv	ATAAACCAGCTCCAGACC
pBAD33RBS *phoQ N202C* fw	GCTCTCAGCCtgcCTGCTGTTAG
pBAD33RBS *phoQ N202C* rv	ACATAGATAAACCAGCTC
pBAD33RBS *phoQ L203C* fw	CTCAGCCAATtgcCTGTTAGTGATCCC
pBAD33RBS *phoQ L203C* rv	AGCACATAGATAAACCAG
pBAD33RBS *phoQ L204C* fw	AGCCAATCTGtgcTTAGTGATCCCGC
pBAD33RBS *phoQ L204C* rv	GAGAGCACATAGATAAACC
pBAD33RBS *phoQ L205C* fw	CAATCTGCTGtgcGTGATCCCGC
pBAD33RBS *phoQ L205C* rv	GCTGAGAGCACATAGATAAAC
pBAD33RBS *phoQ V206C* fw	TCTGCTGTTAtgcATCCCGCTGCTGTG
pBAD33RBS *phoQ V206C* rv	TTGGCTGAGAGCACATAG

pBAD33RBS *phoQ I207C* fw	GCTGTTAGTGtgcCCGCTGCTGT
pBAD33RBS *phoQ I207C* rv	AGATTGGCTGAGAGCACATAG
pBAD33RBS *phoQ L209C* fw	AGTGATCCCGtgcCTGTGGGTCG
pBAD33RBS *phoQ L209C* rv	AACAGCAGATTGGCTGAG
pBAD33RBS *phoQ L210C* fw	GATCCCGCTGtgcTGGGTCGCCG
pBAD33RBS *phoQ L210C* rv	ACTAACAGCAGATTGGCTG
pBAD33RBS *phoQ W211C* fw	CCCGCTGCTGtgcGTCGCCGCCT
pBAD33RBS *phoQ W211C* rv	ATCACTAACAGCAGATTGGCTGAGAGCACATAG
pBAD33RBS *phoQ V212C* fw	GCTGCTGTGGtgcGCCGCCTGGT
pBAD33RBS *phoQ V212C* rv	GGGATCACTAACAGCAGATTGGC
pBAD33RBS *phoQ A213C* fw	GCTGTGGGTCtgcGCCTGGTGGA
pBAD33RBS *phoQ A213C* rv	AGCGGGATCACTAACAGC
pTrc99A *mgrB C16A*	GGTGTTGGCTgcgTTGCTGCTTTGGG
pTrc99A *mgrB C16A*	ACGACAACCAGAACGACC
pTrc99A *mgrB C16A W20C* fw	GTTGCTGCTTtgcGCGCAGGTATTC
pTrc99A *mgrB C16A W20C* rv	GCAGCCAACACCACGACA
pTrc99A *mgrB C16A Q22C* fw	GCTTTGGGCGtgcGTATTCAACATGATGTGCG
pTrc99A *mgrB C16A Q22C* rv	AGCAACGCAGCCAACACC
pTrc99A *mgrB C16A F24C* fw	GGCGCAGGTAtgcAACATGATGTG
pTrc99A *mgrB C16A F24C* rv	CAAAGCAGCAACGCAGCC
pTrc99A *mgrB D31C*	GTGCGATCAGtgcGTACAATTTTTCAGC
pTrc99A *mgrB D31C*	ATCATGTTGAATACCTGC
pBAD33RBS *mgrB W20A*	CTTGCTGCTTgcgGCGCAGGTATTC
pBAD33RBS *mgrB W20A*	CAAGCCAACACCACGACA
pBAD33RBS *mgrB C28A*	CAACATGATGgcgGATCAGGATGTACAATTTTTCAGC
pBAD33RBS *mgrB C28A*	AATACCTGCGCCCAAAGC
pET-Duet-1 *mgrB N25-TAG* fw	GCAGGTATTCtagATGATGTGCGATCAG
pET-Duet-1 *mgrB N25-TAG* rv	GCCCAAAGCAGCAAGCAA
pET-Duet-1 *mgrB M26-TAG* fw	GGTATTCAACtagATGTGCGATCAGGATGTACAATTTTTC
pET-Duet-1 *mgrB M26-TAG* rv	TGCGCCCAAAGCAGCAAG
pET-Duet-1 *mgrB M27-TAG* fw	ATTCAACATGtagTGCGATCAGGATGTACAATTTTTCAGCG
pET-Duet-1 *mgrB M27-TAG* rv	ACCTGCGCCCAAAGCAGC
pET-Duet-1 *mgrB D29-TAG* fw	CATGATGTGCtagCAGGATGTACAATTTTTC
pET-Duet-1 *mgrB D29-TAG* rv	TTGAATACCTGCGCCCAA
pET-Duet-1 *mgrB Q30-TAG* fw	GATGTGCGATtagGATGTACAATTTTTC
pET-Duet-1 *mgrB Q30-TAG* rv	ATGTTGAATACCTGCGCC
pET-Duet-1 *mgrB D31-TAG* fw	GTGCGATCAGtagGTACAATTTTTC
pET-Duet-1 *mgrB D31-TAG* rv	ATCATGTTGAATACCTGC
pET-Duet-1 *mgrB V32-TAG* fw	CGATCAGGATtagCAATTTTTCAGCGGAATTTG
pET-Duet-1 *mgrB V32-TAG* rv	CACATCATGTTGAATACCTG
pET-Duet-1 *mgrB Q33-TAG* fw	TCAGGATGTAtagTTTTTCAGCG

pET-Duet-1 *mgrB Q33-TAG* rv	TCGCACATCATGTTGAATAC
pET-Duet-1 *mgrB F34-TAG* fw	GGATGTACAAtagTTCAGCGGAATTTG
pET-Duet-1 *mgrB F34-TAG* rv	TGATCGCACATCATGTTG
pET-Duet-1 *mgrB F35-TAG* fw	TGTACAATTTtagAGCGGAATTTGTG
pET-Duet-1 *mgrB F35-TAG* rv	TCCTGATCGCACATCATG
pET-Duet-1 *mgrB S36-TAG* fw	ACAATTTTTCtagGGAATTTGTGCCATTAACCAG
pET-Duet-1 *mgrB S36-TAG* rv	ACATCCTGATCGCACATC
pET-Duet-1 *mgrB G37-TAG* fw	ATTTTTCAGCtagATTTGTGCCATTAACC
pET-Duet-1 *mgrB G37-TAG* rv	TGTACATCCTGATCGCAC
pET-Duet-1 *mgrB I38-TAG* fw	TTTCAGCGGAtagTGTGCCATTAACC
pET-Duet-1 *mgrB I38-TAG* rv	AATTGTACATCCTGATCG
pET-Duet-1 *mgrB A40-TAG* fw	CGGAATTTGTtagATTAACCAGTTTATC
pET-Duet-1 *mgrB A40-TAG* rv	CTGAAAAATTGTACATCCTG
pET-Duet-1 *mgrB I41-TAG* fw	AATTTGTGCCtagAACCAGTTTATCC
pET-Duet-1 *mgrB I41-TAG* rv	CCGCTGAAAAATTGTACATC
pET-Duet-1 *mgrB N42-TAG* fw	TTGTGCCATTtagCAGTTTATCC
pET-Duet-1 *mgrB N42-TAG* rv	ATTCCGCTGAAAAATTGTAC
pET-Duet-1 *mgrB Q43-TAG* fw	TGCCATTAACtagTTTATCCCGTG
pET-Duet-1 *mgrB Q43-TAG* rv	CAAATTCCGCTGAAAAATTGTAC
pET-Duet-1 *mgrB F44-TAG* fw	CATTAACCAGtagATCCCGTGGTG
pET-Duet-1 *mgrB F44-TAG* rv	GCACAAATTCCGCTGAAAAATTG
pET-Duet-1 *mgrB I45-TAG* fw	TAACCAGTTTtagCCGTGGTGACTG
pET-Duet-1 *mgrB I45-TAG* rv	ATGGCACAAATTCCGCTG
pET-Duet-1 *mgrB P46-TAG* fw	CCAGTTTATCtagTGGTGACTGC
pET-Duet-1 *mgrB P46-TAG* rv	TTAATGGCACAAATTCCG
pET-Duet-1 *mgrB W47-TAG* fw	GTTTATCCCGtagTGACTGCAGG
pET-Duet-1 *mgrB W47-TAG* rv	TGGTTAATGGCACAAATTCC
Nde1 *flag-mgrB* fw	aaaCATATGGATTATAAAGATGATGATGATAAAGGTATGGTGAGCAAGGGCGAGGAG
EcoRV *flag-mgrB* rv	aaaGATATCTCACCACGGGATAAACTG
Nco1 *phoQ(his)₆* fw	aaaCCATGGGCAAAAAATTACTGCGTCTTTTTTTCCCGC
BamH1 *phoQ(his)₆* rv	aaaGGATCCTTAATGGTGATGGTGATGGTGGC

1034
1035
1036
1037 **SI References**
1038

1039 1. Yadavalli, S.S., et al., *Functional determinants of a small protein controlling a*
1040   *broadly conserved bacterial sensor kinase.* J Bacteriol, 2020.
1041 2. Yuan, J., et al., *Osmosensing by the bacterial PhoQ/PhoP two-component system.*
1042   Proc Natl Acad Sci U S A, 2017. **114**(50): p. E10792-E10798.