Discovering individual-specific gait signatures from data-driven models of neuromechanical dynamics

[1]Taniel S. Winner, [1]Michael C. Rosenberg, [2]Trisha M. Kesar, [1,2]Lena H. Ting, [3]Gordon J. Berman

[1] W.H. Coulter Dept. Biomedical Engineering, Emory University and Georgia Institute of Technology, Atlanta, GA, USA [2] Department of Rehabilitation Medicine, Division of Physical Therapy, Emory University, Atlanta, GA, USA   [3] Department of Biology, Emory University, Atlanta, GA USA

Abstract:

Locomotion results from the interactions of highly nonlinear neural and biomechanical dynamics. Accordingly, understanding gait dynamics across behavioral conditions and individuals based on detailed modeling of the underlying neuromechanical system has proven difficult. Here, we develop a data-driven and generative modeling approach that recapitulates the dynamical features of gait behaviors to enable more holistic and interpretable characterizations and comparisons of gait dynamics. Specifically, gait dynamics of multiple individuals are predicted by a dynamical model that defines a common, low-dimensional, latent space to compare group and individual differences. We find that highly individualized dynamics – i.e., gait signatures – for healthy older adults and stroke survivors during treadmill walking are conserved across gait speed. Gait signatures further reveal individual differences in gait dynamics, even in individuals with similar functional deficits. Moreover, components of gait signatures can be biomechanically interpreted and manipulated to reveal their relationships to observed spatiotemporal joint coordination patterns. Lastly, the gait dynamics model can predict the time evolution of joint coordination based on an initial static posture. Our gait signatures framework thus provides a generalizable, holistic method for characterizing and predicting cyclic, dynamical motor behavior that may generalize across species, pathologies, and gait perturbations.

Introduction

Locomotion is a ubiquitous, complex, and dynamic behavior that is essential for survival. Using cyclic patterns of joint angles, inter-limb and inter-joint coordination, animals effectively move through their environments: walking, running trotting, swimming, flying, and crawling. Even within species and types of locomotion, variations in locomotor patterns often occur across behavioral contexts, groups, and individuals. Thus, although locomotor patterns can appear highly stereotyped, considerable inter- and intra-individual variability exists. Studies of locomotor behaviors have shown systematic differences in movement patterns based on a wide range of neural (Frisk et al., 2019; Ivanenko et al., 2003; Krogh et al., 2022; Young et al., 2022) and biomechanical perturbations (Genthe et al., 2018; Pan et al., 2022; Reisman et al., 2010; Wang et al., 2022) environmental challenges (D'souza et al., 2020; Larsen et al., 2022), psychological state (Attwood et al., 2021; Elkjær et al., 2022), social status (Steptoe and Zaninotto, 2020; Zaninotto et al., 2013), injury (Garcia et al., 2022; Jang and Wikstrom, 2022; Milner et al., 2022), and disease (Ijmker and Lamoth, 2012; Jonkers et al., 2009; Moura Coelho et al., 2022; Prosser et al., 2022; Russo et al., 2022; Troisi Lopez et al., 2022; Young et al., 2022). Furthermore, locomotor impairments can arise from a wide range of physiological and neurological changes, from the subtle changes that may be indicators of progressive disorders (e.g., aging, cognitive impairments) to profound impairments with brain injury (e.g., stroke, spinal cord injury) that can severely limit locomotor function. Although locomotor deficits are often subjectively visible to a human observer, objectively characterizing and understanding sometimes subtle yet important differences in locomotion from a scientific and mechanistic standpoint has been challenging (Correa et al., 2011; Geyer and Herr, 2010; Kuo and Donelan, 2010). For example, kinematic movement patterns (the continuous motion of joint angles over time) have been collected across a wide range of locomotor modes and species but revealing individual-specific differences in kinematics remains difficult. One barrier to progress is that interpreting individual differences in kinematics without an underlying dynamical model is challenging, as kinematics are the result of the complex neuromechanical dynamics that drive the spatiotemporal dependencies of joint kinematics over time. Thus, capturing these underlying gait dynamics is likely essential for interpreting differences in gait and movement across conditions and individuals.

Traditionally, gait dynamics are modeled using physiologically detailed neuromechanical equations, however making predictive models using this approach has often proved challenging (Falisse et al., 2019; Meyer et al., 2016; Pitto et al., 2019a). Partially, this difficulty arises because in order to understand the dynamics underlying gait, we also need to understand how neural feedback and control shape these dynamics. While many models (e.g., musculoskeletal models) that use principles like optimal control can generate simulations of unimpaired gait, as well as changes in gait due to altered biomechanical or neural constraints, they often fail to predict changes in gait kinematics following neurological injury (Meyer et al., 2016) or more subtle perturbations (De Groote and Falisse, 2021; Pitto et al., 2019b). Progress in the physiological modeling of locomotor circuitry in the spinal cord and brainstem demonstrates the role of neural circuits in gait dynamics. However, these models typically rely on simplified (Dzeladini et al., 2014; Taga et al., 1991) biomechanical properties and cannot yet predict the deficits in gait specific to an individual (Angelidis et al., 2021; Geyer and Herr, 2010; Kuo, 2002; McCrea and Rybak, 2008). More importantly, if a hyper-realistic model of the neural and biomechanical system did exist, the relationships between the high-dimensional parameters and actual movement patterns would not likely be unique, as many parameters would not be identifiable, even given massive amounts of data, as many different parameter choices could lead to the same biomechanical output (Holmes

et al., 2006; Prinz et al., 2004). This non-identifiability limits the predictive power and
generalizability of these models to other interventions and conditions outside of limited contexts,
suggesting a need for a more holistic approach.

Despite these challenges, rich individual-specific information exists in gait data. For
instance, through observation of movement, the human brain can perceive many socially salient
features of an individual's gait, suggesting that it should be possible to infer aspects of gait
dynamics from kinematic data. As an example, humans can derive a host of information about
individuals from movement patterns, including gender (Pollick et al., 2005), body size (Troje,
2002), sexual orientation (Johnson et al., 2007), emotion (Gross et al., 2012), individual
differences in dancing (Brown et al., 2021), perceived affective states (Edey et al., 2017) and
underlying intention (Becchio et al., 2012). Furthermore, judgements based on how individuals
move can drive decisions such as partner desirability or attractiveness (Neave et al., 2011),
disease diagnosis (Habersack et al., 2022; Heinik et al., 2010), and treatment planning (Ferrante
et al., 2016; Mikolajczyk et al., 2018).

Despite the rapid advent of technologies providing kinematic measurements through a
wide range of techniques, from videos to wearable sensors, we are still limited in how kinematic
data can help interpret individual differences in gait (Porciuncula et al., 2018; Stenum et al., 2021).
Current approaches to comparing biomechanical features or kinematic trajectories quantify
between-group differences or inter-individual similarity but lack sufficient sensitivity to reveal
interpretable differences in individuals' gaits (Chen et al., 2005; Kettlety et al., 2022; Ries et al.,
2014). Inter-joint coordination differs across individuals, as muscular coordination patterns vary
across a variety of motor skills and deficits in individual-specific ways. Indeed, metrics of muscle
coordination in children with cerebral palsy are consistent with clinician judgements of motor
control complexity that predict intervention outcomes (Steele et al., 2015). Recently, supervised
machine learning methods have been used to classify differences in a large sets of gait kinematics
that were labeled by groups or individuals (Halilaj et al., 2018; Ries et al., 2014). However, these
approaches have not modeled the underlying gait dynamics, nor can they discover subtle
differences in gait that are not labeled a priori.

Here we develop a data-driven framework for modeling gait dynamics that represents
multiple individuals in the same latent space. This latent space reveals individual- and group-level
differences in the neuromechanical dynamics of gait. We used kinematic data from multiple
healthy and neurologically impaired individuals, each walking at six different speeds, to train a
recurrent neural network (RNN) that learns gait dynamics. This phenomenological approach infers
complex spatiotemporal dynamics and enables future kinematic predictions to be made based on
current and prior kinematic postures. Once trained, differences in gait dynamics across groups,
individuals, and walking speed were projected onto a common, low-dimensional latent space of
the model parameters. The stride-averaged representation of gait dynamics in the latent space
constitutes a "gait signature" that we use to characterize differences across individuals, groups,
gait speed, and impairment severity. To demonstrate the generalizability of gait dynamics, we
show that interpolating gait signatures to predict gait kinematics at new walking speeds is more
accurate than interpolating the kinematics themselves in healthy individuals. Further, we show
that the low-dimensional basis functions we discovered have biomechanical interpretability in
terms of the inter- and intra-limb coordination patterns that they generate. The dynamical
projections onto each basis function for each trial can be independently driven through the trained
gait dynamics model to reconstruct the kinematics associated with that specific basis function.

We generated illustrations of the reconstructed joint angle kinematics to visualize and infer what
aspects of gait coordination each subcomponent influences. These subcomponents of gait
coordination can be manipulated independently (i.e., gait sculpting) to infer the relationships
between specific underlying dynamical components and their corresponding kinematic
phenotypes and to identify what specific gait rehabilitation strategies are likely required for
individuals. Finally, our gait dynamics model is generative; it can predict individual-specific time
evolution of kinematics from an initial arbitrary posture (self-driving) once the network is primed
with several gait cycles of the individual's kinematic data. This study establishes a new data-
driven framework to quantitatively interpret individual-specific differences in gait dynamics with
the potential to enable discovery in a wide range of gait coordination deficits, contexts and
interventions in humans and other animals.

Results

Gait signatures: a low-dimensional representation of gait dynamics

We used motion capture to collect sagittal-plane kinematic data that consisted of 15
seconds of continuous gait kinematics from bilateral, hip, knee, and ankle joints from 5 able-
bodied (AB) participants and 7 stroke survivors (> 6 months post-stroke, gait speeds 0.1 to 0.8
m/s) walking on a treadmill at a range of six different speeds each. Taking inspiration from neural
network models that capture neural dynamics (Pandarinath et al., 2018; Sussillo et al., 2015; Vyas
et al., 2020) and biological systems, we implemented a recurrent neural network (RNN) model to
capture the dynamical properties of gait.

Developing the recurrent neural network (RNN) architecture and training the model

The gait dynamics model was developed in Python using common Python libraries such
as TensorFlow, Keras, Pandas, and NumPy. We developed our code in Google Colab to facilitate
open-source sharing of our dynamic framework, which can be found here:
https://github.com/bermanlabemory/gait_signatures. The model architecture was selected based
on two criteria: 1) minimizing model training and validation loss, and 2) maximizing the similarity
of short-time (single stride) and long-time (multiple strides) self-driven model predictions (termed:
gait signature alignment) (Fig. 1 - figure supplement 1). We evaluated these criteria against
alternative models by varying 2 hyperparameters (number of LSTM units and the lookback time,
see Methods). The selected model architecture is a sequence-to-sequence RNN (Sutskever et
al., 2014) consisting of an input layer, a hidden layer of 512 LSTM units, and an output layer. The
RNN learns a map from time-series kinematic input data (0 to T-1) to kinematics one time-step in
the future (1 to T) for all training trials (Fig. 1A). The model was trained until training and validation
error converged and stabilized around the same point (degrees < 0.75°). Thus, the model
successfully learns the underlying dynamics of gait (Fig. 1 - figure supplement 2). The model's
internal states capture trial-specific dynamics predicting the time evolution of joint kinematics;
activation coefficients (H) and memory cell states (C) and are tuned based on kinematic inputs.
Kinematic data was input in multivariate format, not concatenated (Horst et al., 2019;
Santhiranayagam et al., 2015). In brief, our RNN model was designed to capture short and long-
term gait dependencies in time (Ahamed et al., 2021; Hausdorff et al., 1996) as well as inter-and
intra-limb coordination over time, uncovering features of gait that were not previously targeted or
used in gait analysis. To verify whether our model was generalizable, we conducted leave-one-
out cross validation where 12 different models were trained leaving a single individual's 6 trials
on each model run (Fig. 1 - figure supplement 3). Stroke-survivors are known for having

neurological impairments that result in heterogeneous gait dysfunction that are not fully
understood; thus, we anticipate that our gait dynamics model will capture and shed light on these
individual-specific deficits in gait coordination, identify similar coordination strategies or deficits
amongst our stroke cohort, and allow us to compare these different gait dysfunctions to the able-
bodied 'normative' gait (controls).

Generating gait signatures

To generate gait signatures, kinematic trajectories from each walking speed trial across
participants were fed as input into the trained neural network and the corresponding internal states
(H and C parameters, see above) were extracted (Fig.1A). The internal activations prescribe the
spatial and temporal dependencies generating the input kinematics. The resulting time-series of
1024 internal states (512 H, 512 C parameters) were dimensionally reduced using Principal
Components Analysis (PCA) and phase averaged (Revzen and Guckenheimer, 2008). Phase
averaging is applicable here, as the underlying gait dynamics are periodic, and the translation
from time to a phase between 0 and 2π allows us to describe all internal state dynamics in a
speed-independent manner.

Figure 1: Pipeline figure outlining the steps to generating individual-specific gait signatures. *Continuous, multi-*
joint kinematics from multiple individuals are fed into the RNN model as input data and the model is trained sequence-
to-sequence to predict one-step time shifted output kinematics. High dimensional internal parameter (H and C) time
traces per individual are extracted and principal component analysis was applied to reduce the dimensionality of the
data to form individual gait signatures (A). 3D time trace visualizations of 3 representative individuals (able-bodied
(blue), high-functioning (red), low-functioning stroke (orange)) of the 1st 3 dominant principal component contributions
(B, left). 3D projections of the 6-D gait signatures using multi-dimensional scaling (MDS) reveal different gait dynamics
amongst the three gait groups: able-bodied (blue), high-functioning (red) and low-functioning (orange) stroke survivors
(B, right). The size of the circles represents the individual's trial speed (i.e., the smallest circles represent an individual's
slowest gait speed, and the size of the circles increase with gait speed).

The first 6 Principal Components (PCs) explain ~77% of the variance in gait dynamics
(Fig. 1 - figure supplement 4), allowing us to focus on these modes for our visualization and
analysis. The time-varying contributions of the first 3 dominant PCs were plotted in 3D for 3
representative individuals - able-bodied adults, high-functioning stroke (self-selected (SS) walking
speed > 0.4m/s) and low-functioning stroke (SS speed < 0.4m/s) - highlighting that the gait
dynamics between all 3 individuals are different (Fig. 1B, left). The gait dynamics of the high-
functioning stroke survivor (red), while spatially closer to the able-bodied individual (blue) than
the low-functioning stroke survivor (orange), show observable differences in its dynamical
trajectory between to the two individuals. To determine whether some structure exists amongst

the three different subject groups, all the 6-dimensional gait signatures were projected onto a 3D
map using Multidimensional scaling (MDS) (Cox and Cox, 2008) to visualize relative distances
between all gait signatures (Fig.1B, right). The locations of the 3 MDS projections of the 3
representative individuals are not arbitrary, as they belong to clusters of gait signatures of the
same gait group. Thus, gait signatures preserve key clinically relevant features of the underlying
gait dynamics, independent of the individual or speed.

Gait signatures reveal that individual-specific differences in dynamics are favored in the
gait representation, over differences in gait speed

Gait signatures of individuals' 6 speed trials within both cohorts (healthy and stroke) are
tightly grouped together. Gait signatures represent individual-specific dynamics; the unimpaired
cohort exhibit a stereotyped low-dimensional structure across individuals in the able-bodied
cohort (Fig. 2A, i left) vs. the impaired cohort, which display much more variable (i.e., highly
individualized) low-dimensional representations (Fig. 2A, i right). Because the data are phase
averaged over the gait cycle, we demonstrate that gait signature trajectories are well-aligned with
the four gait phases (leg 1 swing, leg 1 stance, leg 2 swing, leg 2 stance), enabling phase-specific
comparisons of differences in gait dynamics. The unimpaired group showed similar structure
across the four gait events (Fig. 2A, ii, left), whereas there was much more variability within the
impaired group (Fig. 2A, ii, right), revealing individual-specific differences within and across
distinct parts of the gait cycle. The similarity between gait signatures was computed and visualized
in a dimensionally reduced gait map space using MDS and colored according to the different
individuals in the dataset (Fig. 2B, i). The unimpaired group form a cluster in the gait map, showing
that individuals in the unimpaired group are distinct from the impaired group. Stroke-survivors
occupy distinct positions from other impaired individuals' sub-clusters in the gait space that
highlight the well-established but poorly understood heterogeneity in gait deficits in the stroke
cohort. Furthermore, individual-specific gait signatures change slightly as individuals walk faster
than their self-selected pace (Fig. 2B, i). However, these within-subject speed-induced changes
are much smaller than between-individual difference in gait signatures. We calculated the
Euclidean distance between individuals' self-selected speed trial gait signature and the calculated
able-bodied centroid (Fig. 2B, i black square) and the results shown on the plot to the right reveal
that low-functioning stroke survivors (self-selected gait speed < 0.4 m/s) are further away from
the able-bodied cluster than the high-functioning stroke survivors. Showing the validity of our
approach, low-functioning stroke survivors are less dynamically similar to AB than higher
functioning stroke survivors.

Gait speed does not appear to strongly influence the differences in dynamics between
individuals' gaits (note that the range of gait speeds for each participant may not have been wide
enough to elicit major differences in their overall dynamics). Overall, as expected, the unimpaired
group walked at faster speeds than the impaired group (Fig. 2B, ii). Individuals in the able-bodied
cluster walk at a range of different speeds, but individual gait signatures still cluster tightly
together. Post-stroke individuals who walk at similar slower speeds, however, maintain their own
distinct individualized groupings. Thus, individuals' characteristic gait signatures were preserved
across their range of walking speeds and were not grouped based on absolute walking speed.
For example, several clinically similar post-stroke individuals (similar overground walking speed
and Fugl-Meyer score (Fugl-Meyer et al., 1975) have very different gait signatures that remain
recognizable across a range of gait speeds (Fig. 2). Furthermore, when used to distinguish
between gait groups and identify individuals, gait signatures perform similarly to a when using a

set of 26 commonly used discrete variables (Fig. 2 - figure supplement 1). Gait signatures also
perform better than continuous kinematics and joint velocities at these same tasks (Fig. 2 - figure
supplement 1). These results serve as a positive control, as researchers previously could
distinguish gait groups by building a classifier based on important subjectively selected discrete
variables. Here, we have created a dynamical representation that can distinguish groups with
similar accuracy. It is not surprising that the continuous kinematics performed worse than the
RNN gait signatures (which were developed from these very same data), as the RNN model used
the data to encode important time-varying changes in the kinematics, allowing for more
information to be extracted. Thus, parameterizing the evolution of individuals' walking patterns
into a common subspace allows for a more holistic, less biased, and straightforward analysis of
primarily their overall differences in gait dynamics, inter- and intra-limb coordination over any
differences attributed to absolute gait speed. Gait signatures can allow gait researchers to study
or analyze the dynamical differences underlying impairment independently from gait speed,
facilitating analysis of dynamics between individuals who may not be capable of walking at the
same speeds and allowing investigation of changes in the underlying mechanism of gait changes
under different conditions (walking speed, gait rehabilitation intervention, age etc.)

Figure 2: Gait signatures reveal highly individualized low dimensional representations of gait dynamics
irrespective of absolute gait speed. A) 3D unimpaired (left) and impaired (right) gait signatures colored by i) individual
and ii) gait phase. Gait signatures are grouped together according to individuals within both cohorts (same hues of blue
cluster together for unimpaired (i, left) and similarly the same hues of red cluster in the impaired cohort (i, right)). In our
convention the right leg of all unimpaired individuals was assigned to be the paretic leg and left leg the non-paretic leg.
Impaired individuals can have either left or right leg paresis. Unimpaired gait signatures reveal a similar looped
structure across the four gait phases that occur during a gait cycle (leg 1 swing, leg 1 stance, leg 2 swing, leg 2 stance)
(ii, left) whereas impaired signatures showed individual-specific differences across the four phases and were more
variable (ii, right). B) 3D multidimensional scaling applied to all gait signatures shows the pronounced separation
between unimpaired (blue hues in left section of map) and impaired (red hues in right section of map) gait dynamics
(i). Impaired signatures (red hues) are located further away from the centroid of all unimpaired gait signatures (black
square), indicating that they are less dynamically similar to the unimpaired individuals. The smallest circles represent
an individual's self-selected walking speed trial and larger circles correspond to the faster speed trials. Low-functioning
stroke survivors (encapsulated in green; based on self-selected gait speed < 0.4m/s) are located furthest away (largest
Euclidean distances) from the unimpaired centroid (i). Gait speed does not appear to strongly influence the differences
in dynamics between individuals as similar speed gait signatures are in different regions of the gait map (ii). Particularly,
gait speed does not explain the heterogeneity in low-functioning stroke survivors' gait dynamics.

Low-functioning stroke-survivors are less dynamically analogous to able-bodied and more dynamically variable compared to high-functioning stroke-survivors.

Clinically, gait rehabilitation researchers use gait speed as a primary quantitative indicator of gait impairment (Awad et al., 2015; Hornby et al., 2020; Jonkers et al., 2009). While this coarse metric gives an overall value or number to one's overall gait function, it does not identify the specific dysfunctions or impairments underlying the individuals' gait. To derive more precise measures or indicators of gait impairment, we anticipated that utilizing this gait signatures framework, we would be able to capture both subtle and obvious differences in impaired gait. In the clinic, stroke survivors are typically segmented into subgroups according to their self-selected walking speeds: high-functioning stroke survivors who typically maintain a self-selected (SS) walking speed above 0.4m/s and low-functioning stroke survivors who adopt SS walking speeds less than or equal to 0.4m/s (Bowden et al., 2008). It is assumed that low-functioning stroke survivors are more impaired and thus adopt slower walking speeds to be able to navigate the environment safely. Gait deficits of stroke survivors within either sub-group are heterogeneous across individuals and include different impairments such as foot drop, reduced paretic push-off during late stance, limited initial heel contact during early stance, as well as further compensatory gait strategies such as hip circumduction and hip hiking. We expected that higher functioning individuals would have less severe impairments and would be more dynamically analogous to able-bodied individuals, whereas low-functioning stroke survivors would exhibit highly variable impairments from each other and be even less dynamically analogous to able-bodied dynamics compared to higher functioning stroke survivors.

To better visualize all developed individuals' gait signatures across their 6 different speed trials in our dataset, we again used MDS to project the 6D gait signatures to 3D. This mapping allows us to visualize the relative locations of individuals in comparison to all the other gait signatures to gain insights on how dynamically similar they are from one another. A 3D MDS gait map of all gait signatures reveals that able-bodied and high-functioning stroke survivors are located near each other, whereas low-functioning stroke survivors are farther and more dispersed, and form distinct clusters in different regions of the map (Fig. 3A). Sub-group level analysis reveals significant differences in the Euclidean distance metric (distance between each gait signature and the able-bodied centroid) between the able-bodied group and the low- and high-functioning stroke survivor groups, respectively (Fig. 3B). Able-bodied gait signatures are located closest to the centroid, followed by high-functioning and low-functioning stroke survivors (Fig. 3B). The within-group dispersion of gait dynamics for the low- and high- functioning stroke survivors was calculated based on the radius of a hypersphere enclosing 95% of the groups' gait signatures. Using a leave-one-out sample with replacement method, multiple within-group dispersion calculations were conducted for each group and the average within-group dispersion was expressed alongside the standard error in Fig. 3C. The 95th percent radius was significantly higher in the low-functioning stroke-survivors gait signatures compared to the high-functioning, highlighting that low-functioning gait signatures were more dispersed from each other (higher inter-individual variability) and the RNN model can capture these individual-specific gait deficits in individuals with more severe gait impairment.

*Figure 3: **Comparison of gait signatures across three gait subgroups: able-bodied (AB), high functioning (HF)***
*328   **and low functioning (LF)**. A) 3D gait map using multidimensional scaling highlights the relative distances between AB*
329   (blue), HF (red) and LF (orange) stroke survivors. LF stroke survivors are less clustered and occupy distinct regions of
330   the map away from the able-bodied centroid (black square). B) Gait dynamics similarity based on Euclidean distance
331   between AB centroid and each participant, showing larger distances within the low-versus high-functioning groups. C)
332   Within-group dispersion of gait signatures based on the radius of a hypersphere enclosing 95% of each group's gait
333   signature reveals more dispersed gait signatures in low- versus high-functioning stroke survivors, highlighting the
334   potential of gait signatures to capture individual differences in more severe gait impairments.

Gait signatures are biomechanically interpretable

While Principal Component trajectories and low-dimensional maps provide one way to compare the overall dynamics between individuals and groups, it remains to be seen what information the independent components of the 6D gait signature represent biomechanically. The contributions of each principal component (PC) to a gait signature fluctuates over the gait cycle, shown for an exemplar able-bodied, one high-functioning stroke survivor, and one low-functioning stroke survivor in Fig. 4A. Superimposed individual stride-averaged PC projections from these 3 individuals (Fig. 4B) highlight the specific differences in each PC. For PC1, both able-bodied and high-functioning stroke survivor traces are within the able-bodied 95% confidence interval, whereas the low-functioning stroke survivor is outside of these bounds around the middle of the gait cycle. For PC2, some regions of the low and high-functioning stroke survivor can be found outside of the confidence interval, however the entirety of the PC3 projection of the low-functioning stroke survivor is found outside of interval (vertically shifted). Given the generative nature of our RNN-based model, a specified number of the loadings on the PCs can be driven through the trained RNN model to reconstruct the corresponding kinematics. Thus, to interpret the individual PC components, the internal parameters corresponding to each isolated PC were driven through the gait dynamics model, generating gait predictions, i.e., a multi-joint coordination pattern and their temporal evolution over the gait cycle that can be visualized in an animation or gait movie. Stick figure snapshots (7 equally spaced samples of 100 frames) show that PC1 encodes dynamics driving hip flexion and extension, PC2 encodes dynamics driving knee flexion and extension and PC3 encodes dynamics driving primarily postural coordination (trunk location relative to joints) (Fig. 4C, Fig. 4 - video supplements 1-4). This framework can potentially allow for the identification and targeting of individual-specific gait deficits, informing the tailoring of precision rehabilitation strategies.

Figure 4: Biomechanical interpretation of gait signatures. A) Gait signatures reveal different gait dynamics between exemplar AB, low-and high-functioning stroke survivors. B) The loadings on each principal component (PC), e.g., the contributions of each PC vary over the gait cycle and can be compared to the AB 95 % confidence interval (gray). C) Each PC generates specific multi-joint gait coordination patterns when used to drive the gait model, enabling biomechanical interpretation of gait deficits and effects of treatment.

The gait dynamics model generalizes to unmeasured speeds

Our gait signature model can capture and predict nonlinear changes in dynamics in response to speed in cases where interpolation of kinematics may fail. We trained a different gait dynamics model using only 15s of data of the 2 fastest and 2 slowest walking speeds of each subject. Weighted averages of gait signatures from an individual walking at these four different gait speeds can be used to generate multi-joint kinematic trajectories that predict data from a gait speed that was not used to train the model (Fig. 5). Predicted kinematics from interpolation of gait signatures across the four speeds resemble the measured kinematic reference more accurately than do the kinematics generated from interpolating gait kinematics, shown for an exemplary AB individual (Fig. 5A) and low-functioning stroke survivor (Fig. 5B). Kinematic prediction from interpolation of dynamics did considerably better than interpolating kinematics directly for the exemplary low-functioning stroke survivor shown in Fig. 5B, indicating that interpolating gait signatures capture nonlinear (non-monotonic) changes in kinematics between speeds. The kinematic output of the interpolated kinematics follows that of the fast speed in the paretic hip closely but does not resemble the measured kinematic reference waveforms for the paretic knee or ankle angles. In some cases where interpolation of kinematics fails, the averaged dynamics do a better job at predicting kinematic trajectories at unseen speeds. Group level analyses show that the R^2 values between the measured and predicted kinematics from interpolated gait dynamics are significantly higher (Wilcoxon paired signed rank test) than interpolating kinematics within the able-bodied cohort (Fig. 5C), but not for stroke (Fig. 5D). In general, averaging gait dynamics produced less variable R^2 values and less R^2 outliers than averaging kinematics in both the able-bodied (Fig. 5C) and stroke survivors (Fig. 5D). The range of R^2 values in the able-bodied cohort for averaged dynamics was -0.20 to 1.00 compared to –1.30 to 0.98 in averaged kinematics whereas the range of R2 values in the stroke cohort for averaged dynamics was 0.46 to 1.00 compared to -0.50 to 1.00 in averaged kinematics. Two low-functioning stroke survivors show higher R^2 values of their hip, knee and ankle kinematic traces when interpolating kinematics vs. dynamics. Post hoc analysis revealed that these two stroke survivors (ST4 and ST2) were furthest

away from the able-bodied centroid (least dynamically similar to able-bodied) as shown in Fig. 2B, i. These results suggest that the RNN largely captures more stereotyped able-bodied dynamics and has a harder time learning the dynamics from more variable stroke individuals, especially those that deviate furthest from able-bodied. Our small sample size also limits the amount of data the RNN sees for each diverse type of stroke dynamics. Thus, with a larger stroke patient sample size and longer trials, the RNN may be able to make better kinematic predictions of lower-functioning stroke survivors. Moreover, this result highlights the utility in predicting kinematics in unseen conditions which in contrast cannot be made using discrete biomechanical or clinical metrics, nor with current biophysical models.

Figure 5: Data-driven gait dynamics model predicts non-linear changes in joint kinematics with gait speed. Gait predictions of joint kinematics (green) at intermediate gait speeds not used in model training were generated by interpolating gait signatures between slow (dashed grey) and fast speeds (dashed black) lines and using them to drive the gait model. Interpolated kinematics from gait dynamics (green) and interpolated directly from kinematics (blue) were compared to the measured reference kinematics (black solid). A) Predictions in an exemplar AB participant are more accurate when interpolating gait signatures compared to interpolating gait kinematics across speeds. B) In an exemplar low-functioning stroke survivor, interpolated gait signatures predict nonlinear changes in kinematics better at intermediate speeds than interpolated gait kinematics. Averaging the kinematics fail in this case where there are larger differences between the slow and fast speed paretic kinematics; the averaged kinematics (blue) follow the fast speed paretic hip kinematics whereas the other angles do not reflect waveforms that resemble either the fast or slow speed. The gait model can therefore predict movement reasonably well when interpolating between tested speeds. There is a statistically significant difference between group level R^2 comparisons (kinematics generated from interpolated dynamics vs interpolated kinematics) in the able-bodied (C) but not in stroke (D) cohorts. However, the range of R^2 values are larger in both able-bodied and stroke kinematic predictions resulting from interpolated kinematics (-1.30-0.98, -0.50-1.00 respectively) vs. predicted from interpolated gait dynamics (-0.20-1.00, 0.46-1.00 respectively). Thus, while the R^2 values may not improve on average for the stroke survivors, the model's performance is more robust overall.

Gait sculpting: manipulating the PC components of an individual's gait signature identifies specific coordination deficits in stroke survivors

Previously, we showed that we can leverage our model to reconstruct the kinematics of healthy PC projections of the gait signature to gain insight into their independent biomechanical interpretations. However, identifying and interpreting the biomechanics related to impaired PC dynamics of stroke-survivors' gait would prove to be even more beneficial, as these dynamics can potentially serve as rehabilitation targets when designing tailored gait intervention/strategies for

individuals. Here we present an example of how we use gait signatures to identify specific
biomechanical or coordination targets in specific stroke survivors. Specifically, we utilize our
finding that the phase-varying contributions of the 6 principal projections of the gait signature differ
in individual-specific manners (Fig. 6A). For example, AB2's 6 PC contributions all lie within the
95% confidence interval of all able-bodied individuals. ST4 primarily shows major deviation from
AB in PC 3 (located entirely above the AB confidence interval), impaired dynamics during paretic
swing in PC 4 and overall irregular shapes in PC 5 and 6. ST2's PC3 is largely within the AB
confidence interval, however PC 4's paretic swing shows major deviation, their PC 5 contribution
is shifted below the AB confidence interval and PC 6 shows an irregular shape. ST3's PC5
projection lies below the AB confidence interval and PC 6 projection is irregularly shaped
compared to AB. To validate our finding that suggested that PC 3 primarily influences hip flexion
or extension, we exchanged AB2's healthy PC1 with that of ST4 (Fig. 6A, orange boxes and
arrow) and we observed if and how AB2's original hip joint kinematics (Fig. 6B, orange box, black
trace) deviated (Fig. 6B, orange box, red dashed trace, Fig. 6 - video supplement 1). To gain
further insight into how the other PC deviations manifest in movement, we manipulated the PC3
projection of AB2 by replacing it with that of ST4 (Fig. 6A, brown boxes and arrow). The kinematic
reconstruction from this manipulation (Fig. 6B, brown box, red dashed trace, Fig. 6 - video
supplement 2) shows a vertical shift downwards for bilateral hip angles and the non-paretic knee.
The vertical shifts in the hip flexion/extension angles suggest a major difference in this individual's
posture (perhaps stroke individual leaned forward more during gait) compared to able-bodied. We
manipulated AB2's PC4 projection by replacing it with that of ST2 (Fig. 6A, green boxes and
arrow). This manipulation affected specifically the paretic and non-paretic ankle angles and both
knee joints primarily during the period between non-paretic stance and paretic swing (Fig. 6B,
green box, red dashed trace, Fig. 6 - video supplement 3). This result highlights a coordination
deficit between these specific joint angles and, if targeted accurately, may allow for corrected gait
patterns of this stroke survivor. Conversely, we tested the effects of replacing an impaired PC
projection with a healthy one to observe how gait impairments can potentially be improved. We
replaced ST3's PC5 with that of AB2 (Fig. 6A, purple boxes and arrow) and observed a substantial
change in the magnitude and shape of bilateral ankle angle trajectories and slight increase in non-
paretic knee magnitude (Fig. 6B, purple box, red dashed trace, Fig. 6 - video supplement 4). We
can infer that to make improvements to ST3's PC5 towards able-bodied or normative kinematics,
rehabilitation focusing on these specific knee and ankle strategies may prove useful.

Figure 6: Gait sculpting: interpolating between components of able-bodied and stroke gait dynamics to visualize anticipated gait improvement. *The components of individuals' gait signatures can be manipulated (gait sculpting) to understand the relationship between specific underlying dynamics and their corresponding kinematic phenotype. A) The projection on each of the 1st 6 principal components (PCs) can be observed for a representative able-bodied (AB2), two low functioning stroke-survivors each having similar self-selected (SS) speeds and Fugl-Meyer (FM) scores (ST2 & ST4, as denoted in Figure 2) and another low functioning stroke survivor (ST3) who has a higher FM score and faster SS walking speed. The PC projections are colored according to the 4 gait phases (non-paretic swing, non-paretic stance, paretic swing, paretic stance). The right leg of the unimpaired individuals was arbitrarily assigned to be paretic and the left leg, non-paretic for consistency. Colored boxes and arrows (orange, brown, green, purple) show specific, single PC manipulations, for example, the orange boxes and arrow illustrate that the PC 1 projection of AB2 was replaced with the impaired PC 1 projection from ST4. B) The AB2:ST4 manipulation (orange) shows how AB2's original phase averaged kinematics (black trace) was manipulated by ST4's impaired PC 1 projection (red dashed traced). ST4's impaired PC 1 manifests in AB2's healthy kinematics showing deviation primarily in the hip kinematics (as suggested in Figure 4 where healthy PC 1 encodes a kinematic subcomponent corresponding to hip flexion/extension) and some deviation in the ankle angles, especially the paretic ankle. The AB2:ST4 manipulation (brown) shows how ST4's impaired PC3 manifests in AB2's healthy kinematics; we observe a vertical shift downwards (red trace) of the bilateral hip angles as well as the non-paretic knee. This change in hip flexion highlights that this impaired PC3 encodes a reduction in the hip flexion angles; pointing to a more crouched gait (trunk is leaning forward more). The AB2:ST2 manipulation (green) shows replacing AB2's PC4 projection with ST2's impaired PC4 dynamics shows deviation in the knee joints especially during paretic swing, a vertical shift upwards in the paretic ankle angle kinematics and deviations around the middle of the gait cycle (transition between non-paretic stance and paretic swing) in the non-paretic ankle kinematics. Alternatively, the AB2:ST3 manipulation (purple) the impaired PC5 in ST3 is replaced with the healthy PC5 projection from AB2 resulting in slight increase in non-paretic knee magnitude and reduced amplitude of paretic and non-paretic ankle flexion. The result of this manipulation points to potential predicted improvements (or deviations) that can occur when aiming to mimic PC5 healthy dynamics in this stroke survivor allowing offline in-silico testing of potential avenues for gait rehabilitation for this stroke survivor.*

Self-driven signatures: Our gait dynamics model revealed robustness of gait predictions establishing the utility of gait signatures in precision medicine

The ability to predict future kinematics based on measured data is key to rapid, virtual design of personalized interventions. We demonstrate that the recurrent neural network model of gait dynamics, once primed with several gait cycles of data from either able-bodied or stroke participants, can predict future joint angle trajectories (Fig. 7). Once the network is primed, an initial posture is presented (initial condition, denoted by blue vertical bar) after which the model self-drives i.e., predicts the general shape of future kinematics in a feedforward manner (without

referencing previous measured data points) in an able-bodied (Fig. 7A, i, left) and stroke individual
(Fig. 7B, ii, left). A smooth transition is seen between the previously measured gait cycle (green)
and the self-driven cycle (red trace) for both AB and stroke (Fig. 7A, i, right, Fig. 7A, ii, right
respectively).

To verify that the model was not generating a gait cycle prediction entirely by chance, we
calculated the Euclidean distance between the kinematics of the predicted (self-driven) gait cycle
and the kinematics from each of the measured gait cycles. We expect that if the model was indeed
predicting future gait cycles (more accurately than chance), the Euclidean distance between the
predicted and corresponding measured kinematics would be lower than over 50% of the other
distances in the distribution. We found that the distance between the able-bodied predicted gait
cycle and corresponding measured kinematic cycle was lower than 79% of the distances of the
other gait cycles (Fig. 7A, ii) whereas in the stroke survivor the respective distance was only lower
than 40% (Fig. 7B, ii) suggesting the model is less able to accurately predict future kinematics in
stroke gait. To calculate the Euclidean distances between the gait cycles, we need to normalize
each gait cycle to the period of the self-driven cycle. To avoid this potential bias, we also
performed a comparison using a metric that was not manipulated in time – gait cycle duration.
After priming the model, we presented the model with the first posture of the trial and ran the
network forward in self-driving mode for the remainder of the trial length (15 seconds). Able-
bodied self-driven predicted kinematics resembled the reference kinematics closely (Fig. 7C, i,
top plot) whereas stroke self-driven predicted kinematics matched the first gait cycle closely but
soon converged to patterns reflecting able-bodied kinematics (Fig. 7C, ii, top plot). The gait cycle
duration of the first few cycles of the self-driven kinematics match those of the measured
kinematics (blue dots located close to the y=x line) in both the exemplary able-bodied (Fig. 7C, i,
bottom plot) and stroke individual shown (Fig. 7C, ii, bottom plot), however kinematics soon
diverged to shorter and relatively consistent gait cycle durations (blue dots appearing almost
horizontal in the plots) result in both cases. The model may preferentially predict able-bodied
kinematics, which were less variable between individuals than were post-stroke kinematics.
These results highlight that the model encodes gait dynamics that can predict kinematics over
short timescales, but the variability and amount of training data may influence predictive power
over long timescales. The potential of the gait dynamics model as a predictive tool provides a
glimpse into the potential future application of our approach for predicting the effects of gait
perturbations, technologies, and treatments, thus reducing time, cost, and participant burden
while facilitating the development of more effective strategies to improve gait.

Figure 7: Our trained RNN model can predict the time evolution of kinematics from an initial posture. *The trained gait dynamics model can predict individual-specific time-evolution of gait kinematics from an arbitrary initial posture (self-driving) in able-bodied (A, i) and stroke (A, ii) once the network is primed with several gait cycles of an individual's data (gait cycle$_{x-n(measured)}$, black solid). This predictive ability shows that the model encodes the gait dynamics underlying movement. Despite inter-cycle kinematic variability, the gait dynamics model can predict the general shape of the next gait cycle of kinematics (gait cycle$_{x(predicted)}$, red) in an able-bodied individual (A, i) and stroke survivor (A, ii), however, predicted kinematics (red) show larger deviation from the measured reference gait cycle (gait cycle$_{x(measured)}$, black dashed) in the stroke survivor. A smooth transition exists between the measured kinematics from the gait cycle preceding (gait cycle$_{x-1(measured)}$, green) the self-driven predicted cycle (red). For the representative able-bodied individual (B, i), the Euclidean distance (deviation) between the predicted gait cycle of kinematics and its respective measured kinematics (reference) is ~79% lower than the distance between the other gait cycles in the trial; ruling out that the kinematic predictions are attributed to chance. The deviation (Euclidean distance) of the predicted gait cycle of stroke (B, ii) kinematics to its reference gait cycle is ~40% lower than the distance between the other gait cycles in the trial. This suggests that the dynamical model is less able to accurately predict stroke kinematics better than chance. The dynamical model was first initialized with all the trial's kinematics data (15 seconds) (black trace) after which the trial's initial posture was presented to the model to self-drive kinematics (red trace) in feedforward mode for 15 seconds (C, i, top plot). The duration of each gait cycle from the measured kinematics is not well encoded by the dynamical model; gait cycle durations of the predicted kinematics are typically underestimated in both able-bodied (C, i, bottom plot) and stroke (C, ii, bottom plot) (to a larger degree) in self-driving mode and as such deviate from the y=x reference line (black).*

Discussion

Summary

Here we establish a data-driven framework for comparing and predicting individual-specific locomotor patterns without needing to construct physiologically based mechanistic models. As an initial proof of concept, complex neuromechanical gait dynamics were modeled using a relatively simple recurrent neural network that captures the rules by which joint kinematics during gait transition from one time point to the next. Because the network was trained on multiple healthy and impaired individuals walking at several speeds, its internal parameters provide a basis

for comparing, interpreting, and predicting gait dynamics. Gait signatures further capture
coordination between joints and limbs without the need for pre-selecting gait features that may
introduce bias and ignore the continuous nature of gait. We show that individuals have little
variance in gait dynamics across speeds, leading to the individual-specific "gait signature" concept
and enabling comparisons between individuals moving at different speeds. Across stroke
survivors, we found greater heterogeneity in low-functioning individuals who exhibited disparate
gait dynamics despite similar clinical metrics, highlighting the potential utility of gait signatures in
providing more sensitive diagnoses to personalize therapies. Gait signatures provide a predictive
simulation framework for sculpting gait dynamics to understand coordination deficits and predict
kinematics, potentially forecasting the effects of rehabilitative devices or treatments. Finally, the
gait signatures methodology can be readily applied to other periodic motions across species and
across conditions that alter movement and may be a powerful adjunct to modern experimental
methods aimed at understanding the neural mechanisms underlying movement.

Computational framework captures the neuromechanical dynamics of walking

Using a data driven modeling approach enabled us to learn the underlying gait dynamics
based on data rather than constructing a neuromechanical gait model based on first principles.
Data-driven approaches in gait have not focused on gait dynamics but have solved tasks based
on unique features in multi-dimensional gait data such as classifying gait based on pathologies
(Mannini et al., 2016) or conditions such as fatigue and non-fatigue (Zhang et al., 2014);
identifying gait events (e.g., initial contact, loading response (Aung et al., 2013; Castano-Pino et
al., 2020; Chia Bejarano et al., 2015); and discriminating individuals (Barton et al., 2012; Horst et
al., 2019)). Gait dynamics have typically been described though neuromusculoskeletal models
based on physical principles focusing on musculoskeletal mechanics, (De Groote and Falisse,
2021; Hainisch et al., 2021) but they lack adequate representations of the neural systems that
contribute to the resulting movement patterns, particularly in neurological impairments such as
stroke (Pitto et al., 2019b). Machine learning methods to capture dynamics have been used
across physics, engineering, and neuroscience to learn the dynamics underlying complex
systems when the governing equations are unknown (Bongard and Lipson, 2007; Pandarinath et
al., 2018; Sanchez-Gonzalez et al., 2020). Recently, machine learning models have been used
in human gait to predict continuous kinetic variables such as ground-reaction forces (Alcantara et
al., 2022) or joint torque (Camargo et al., 2022; Giarmatzis et al., 2020) based on kinematic data.
Dynamical machine learning models have also been used to encode gait dynamics, including
responses to perturbations or assistive devices, but their model structure did not enable
comparisons between individuals (Berrueta et al., 2019; Drnach et al., 2019; Maus et al., 2015;
Rosenberg et al., 2020; Wang and Srinivasan, 2013). Here, our RNN-based gait dynamics model
provides a means to capture the rules underlying continuous, multi-joint coordination between
bilateral lower limb joints, and how they evolve over time. Accordingly, we do not explicitly capture
mechanical dynamics (i.e., the relationship motion and force), but the effects of force interactions
within the body and environment and implicitly represented in how multi-joint kinematics evolve
over time, with the network parameters and the internal states at each time point determining the
output kinematics.

As gait arises from complex interactions between the nervous system and the
musculoskeletal system that are not easily modeled from first principles, a data-driven approach
provides a powerful framework for capturing and comparing neuromechanical constraints on gait
dynamics. While biomechanical dynamics clearly play a role in movement, the activation of

muscles by the nervous system enables the body to perform a variety of motor behaviors.
However, the governing spatiotemporal dynamics of neuromuscular signals are poorly
understood, especially in neuro-pathologies such as stroke. During behaviors such as locomotion,
motor patterns can be characterized by the number and structure of motor modules, or muscle
synergies, defining groups of co-activated muscles producing a biomechanical function for gait
(Ting et al., 2015). Similar motor modules are used within individuals across different task
conditions (Chvatal and Ting, 2013; Torres-Oviedo et al., 2006; Torres-Oviedo and Ting, 2010),
and are shaped by learning and disease (Payne et al., 2020; Sawers et al., 2017). Particularly in
post-stroke gait, motor modules appear to constrain motor function. Fewer motor modules are
observed post-stroke with the number of modules correlated to reduced walking speed (Clark et
al., 2010; Shin et al., 2021). Further, different patterns of motor module merging are seen in slower
walking stroke survivors, differentially affecting gait biomechanics in a manner that may
necessitate individualized rehabilitation approaches (Allen et al., 2018). Adding neural constraints
such as motor modules on muscle activations in musculoskeletal simulations improve predictions
of key physiological variables such as joint loading in osteoarthritis (Walter et al., 2014). However,
relating motor modules to kinematic gait patterns post-stroke and in other neurological disorders
has been challenging, likely because the neural constraints are underspecified (Allen et al., 2019;
De Groote et al., 2014; Falisse et al., 2020, 2019; Steele et al., 2019). Our data-driven gait
dynamics model phenomenologically captures the net effects of both neural and mechanical
constraints on gait and can further contribute to an understanding of normal and impaired gait.
Corroborating results from motor module analysis, there were greater differences in gait dynamics
amongst the slowest walking stroke survivors. Since the gait signatures capture spatiotemporal
constraints underlying gait dynamics, they provide a complementary approach to musculoskeletal
simulations. Ultimately, gait signatures may play a complementary role to biophysical simulations,
enabling the relationships between biomechanical principles, neural constraints, and the
emergent gait dynamics to be revealed.

Gait signatures enable holistic comparison of gait dynamics across individuals, speed,
and groups

In contrast to other applications of dynamical machine learning models for gait, we capture
multiple individuals within a single network, enabling comparisons of gait dynamics across groups,
individuals, and gait conditions. Rather than using the network as a black box solely to generate
predictions, we explicitly compare and interpret the model's internal parameters to identify low-
dimensional latent variables representing gait dynamics. To encourage a generalizable data-
driven gait dynamics model, we omitted subject and trial condition (gait speed) labels as inputs to
the neural network. Adding input labels might force the RNN to create separable gait models,
whereas our goal was to have the network learn a structure that could be modified parametrically
to represent individual differences in the neuromechanics of walking. Similarly,
neuromusculoskeletal models assume common dynamic principles across individuals, using
parameter variations to represent individual differences (De Groote et al., 2014; Falisse et al.,
2020, 2019; Fregly et al., 2007; Meyer et al., 2016). We intentionally designed a relatively simple
RNN architecture (e.g., single layer, linear input/output) as a starting point to recover as much
interpretability as possible, with the awareness that more complexity could be added to the model
architecture (number of hidden layers, number of neurons, etc.) if required to fit a given data set
robustly. The representations of gait dynamics that emerge from our model holistically capture
the changes underlying measured kinematics, without being attributable to specific neural or

biomechanical constraints. The loss of physiological interpretability is counterbalanced by the
holistic approach to representing gait dynamics and explaining gait kinematics features.

Analogous to written signatures, we find that features of individual-specific gait signatures
are largely preserved across walking speeds. Recognizable qualitative features of handwriting
are preserved even as the size of letters changes quantitatively, or if different limbs, or writing
instruments are used (Bernshteĭn, 1967). Similarly, it is well known that individuals can be
recognized based on how they move or walk (Beer et al., 2000; Meyer et al., 2016; Sánchez et
al., 2018; Troje, 2002; Zajac et al., 2002), even if joint angle excursions are similar. We show that
gait dynamics are more similar within individuals across speeds than between individuals, leading
to the concept of the gait signature. In contrast, gait kinetics and kinematics vary characteristically
across speeds, such that they cannot be directly compared across speeds (van Hedel et al.,
2006). The relatively small changes in gait signatures across speeds suggest that the signatures
reflect changes in the spatiotemporal relationships between joint kinematics, rather than
quantitative changes in their magnitude. As such, gait signatures appear to encode individual-
specific constraints of walking, making it possible to compare gait either within or between
individuals walking at different speeds.

Gait signatures characterize the high inter-individual variability in gait impairment amongst
stroke survivors beyond overall gait function explained by clinical gait metrics. This heterogeneity
is a direct reflection of the wide range of impairments in stroke survivors, including muscle
weakness, impaired coordination, spasticity, abnormal synergistic activation (muscles not
independently coordinated), and compensatory motion (Chen et al., 2005; Jonkers et al., 2009;
Little et al., 2018). We found that higher-functioning stroke survivors were more dynamically
similar to each other, whereas lower functioning stroke survivors were more dispersed. In fact,
two low-functioning stroke survivors with similar clinical metrics (Fugl Meyer score and gait speed)
had quite different gait signatures. As such, gait signatures have the potential to provide insights
into individual differences in gait dynamics that are simply not captured by clinical metric such as
gait speed. Moreover, in contrast to higher-functioning stroke survivors who share similar gait
dynamics, lower-functioning stroke survivors may require more individuals individualized
rehabilitation approaches targeting specific aspects of gait dysfunction. Further gait signatures do
not require a priori selection of which gait variables to compare (Patterson et al., 2008; Rinaldi
and Monaco, 2013; Schutte et al., 2000; Wonsetler and Bowden, 2017). As such gait signatures
provide a powerful, holistic approach to enhance the specificity and precision of gait diagnosis
and treatment. This framework can potentially extend to other diseases, disorders, injury, etc. to
gain further insight into individuals' specific impairments and uncover specific targets towards
developing targeted therapies for individuals.

Gait signatures enable biomechanical interpretation and manipulation

Our gait dynamics model enables biomechanical interpretation of gait signatures and
exploring "what if" scenarios to sculpt desirable gait dynamics. Gait signatures are based on
principal components (PCs) of the gait model internal states, where the weightings on each PC
vary over the gait cycle. The model parameters can be prescribed over the gait cycle, resulting in
the predicted kinematic outputs (i.e., joint angles). The gait signature PCs and their time-varying
weightings can be individually prescribed in the network as a method to reveal the specific inter-
and intra-limb coordination patterns governed by each PC. Further, any combination of PC's can
be combined and reweighted to generate new kinematic output patterns. For example, we
interchanged healthy and impaired PCs to gain deeper insight into how specific impaired PCs

alter healthy gait and vice versa. Further, interpolating gait dynamics can predict gait kinematics
at walking speeds that were not used in the training data. Especially when there was a nonlinear
response in gait kinematics across speeds, interpolation of gait dynamics to predict gait
kinematics performed better than interpolating gait kinematics directly. As such our data driven
gait dynamics model can be used to show how changing select components of the gait signature
alters gait kinematics, providing a potential framework to identify personalized therapeutic targets
for gait rehabilitation.

Gait Signatures enable prediction of future kinematics

Another powerful aspect of our gait signatures framework is its ability to generate future
gait kinematics in the absence of new data. The model is self-driving for able-bodied individuals,
predicting multiple cycles of gait kinematics in the future. However, the ability of the model to
predict future stroke kinematics is limited to approximately one gait cycle in the future; rendering
it promising in applications that provide control signals to rehabilitation devices (e.g.,
exoskeletons). Reduced predictive power for the stroke patients can be attributed to our model
architecture's relative simplicity, a small sample size (7 stroke survivors), and short time series
(15 seconds/ 1500 sample points per trial). These factors should be addressed to improve the
predictive capacity of the model for impaired gait in the future. Additionally, including more
variables besides sagittal plane kinematics (e.g., frontal plane and coronal plane kinematics and
joint forces, may improve learning of the underlying dynamics of gait and increase predictive
capability of our model.

Generalization to other species and rhythmic movements

Because the input to this model are periodic sequences of behaviors, our gait dynamics
framework should generalize to other species that display similar behavioral motions. (e.g., flight,
crawling, walking). Physicists, computational biologists, and other scientists can benefit from this
method by studying the dynamical behavior of species whose neuromechanical models and
physics of complex terrains are difficult to model. This is the first study to our knowledge that uses
a neural network to study the dynamics of gait in an interpretable manner. While much work is left
to be done, we have provided a simplistic, unsupervised framework to discover individual-specific
differences in walking in health and disease in humans. Despite being limited by a small dataset,
we have shown that our model is generalizable to characterizing and predicting kinematics of one
held-out subject using leave one out cross validation (Fig. 1 - figure supplement 3). Key to note,
this methodology relies on having a periodic or quasi-periodic pattern as non-periodic patterns
would not be able to generate a phase and subsequent signature. We also limited our inputs to
gait kinematics, anticipating applications to the proliferation of new measurement modalities for
movement in humans and animals such as wearable sensors and markerless video-based motion
capture (Cao et al., 2019; Mathis et al., 2018; Uhlrich et al., 2022). However, the gait signatures
framework could easily be extended to include other data types (e.g., force, muscle activity, joint
loadings, center of mass dynamics) and experimental conditions (overground walking,
biomechanical constraints, gait interventions, such as exoskeletons, functional electrical
stimulation, or treatment e.g., drugs, optogenetics). Overall, by modeling the dynamics of
individual's gait based on measured data, we uncovered individual-specific representations of
individuals' neuromechanical constraints that allows direct comparisons between individuals who
do not walk at the same speed. The gait signatures framework has implications for the diagnosis
of disease, development of future tailored gait therapies or interventions and tracking meaningful
changes in the fundamental neuromechanical mechanism of walking.

Materials and methods

Human subject participants

To develop dynamical signatures of human gait, we collected data in seven post-stroke individuals (age = 56 ± 12 years; 2 females; 48 ± 25 months post-stroke; Lower Extremity Fugl-Meyer = 20 ± 4) and five able-bodied (AB) controls (age = 24 ± 4 years; 4 female). All post-stroke participants experienced a cortical or subcortical ischemic stroke, were able to walk on a treadmill for one minute without an orthotic device, and exhibited no signs of hemi-neglect, orthopedic conditions limiting walking, or cerebellar dysfunction. All participants provided informed consent prior to study participation approved by institutional IRB.

Experimental Design

Participants completed 15-second walking trials at six different speeds, distributed evenly between and ranging from each participant's self-selected (SS) speed to the fastest safe and comfortable speed. Across stroke participants, gait speeds ranged from 0.3-1.6 m/s. Each participant's fastest walking speed was determined by progressively increasing the treadmill speed from the SS speed until the participant could no longer comfortably or safely maintain the speed for 30 seconds. Participants rested for 1-2 minutes between consecutive gait trials. During data collection, speed increased from the participant's SS to their fastest speed (i.e., not randomized).

Data acquisition

Reflective markers were attached to the trunk, pelvis, and bilateral shank, thigh, and foot segments (Kesar et al., 2011a). We collected marker position data while participants walked on a split-belt instrumented treadmill (Bertec Corp., Ohio, USA) using an 8-camera motion analysis system (Vicon Motion Systems, Ltd., UK). Participants held onto a front handrail and wore an overhead safety harness that did not support body weight. Marker data were collected at 100 Hz, and synchronous ground reaction forces were recorded at 2000 Hz and were down sampled to 100Hz using previously established techniques (Kesar et al., 2011b, 2010, 2009).

Data processing

Raw marker position data were labeled, gap-filled, and low-pass filtered in Vicon Nexus. Labeled marker trajectories and ground reaction force raw analog data were low-pass filtered In Visual 3D. Gait events (bilateral heel contact and toe-off) were determined using a 20-N vertical GRF cutoff, and sagittal-plane hip, knee, and ankle joint kinematics were calculated in Visual 3D (C-Motion Inc., Maryland, USA).

RNN model development

Our goal was to start with a simple RNN to reduce overfitting with too many parameters and deep layers. We wanted the simplest model capable of learning the dynamics underlying gait which also preserved interpretability. The simplest recurrent neural network (RNN) model architecture consisted of one input layer, one hidden layer and one output layer. The hidden layer was composed of long short-term memory (LSTM) units with a lookback parameter that spaned at least one gait cycle. Model hyperparameter selection is described in a later paragraph.

RNN model training

Model fitting on our selected dataset and architecture was executed on the order of minutes to tens of minutes, using Keras 3.7.13 and TensorFlow 2.8.2 on Google Colab's standard GPU with high-RAM runtime (54.8 gigabytes). The RNN model was trained using bilateral, sagittal-plane, lower-limb joint angles from 5 able-bodied (AB) participants and 7 stroke survivors each walking on a treadmill at 6 steady speeds, ranging from each participant's preferred speed to the fastest safe speed. Our training dataset was input to the RNN in multivariate format (not concatenated) (Horst et al., 2019; Santhiranayagam et al., 2015). We trained a sequence-to-sequence RNN with 512 long-short-term memory (LSTM) activations units in the single hidden layer, capable of using 15 seconds (sample rate of 100 Hz) time-series kinematic input data (0 to T-1) to predict kinematics one time-step in the future (1 to T) for all training data across individuals and speeds. Our data was batched according to the number of total trials (N = 72); thus, the LSTM maintains its internal state while a batch is being processed, after which the internal state can be maintained or cleared. Because our network retains its internal state from one time step to the next (i.e., the RNN is *stateful*), we have fine-grained control over when the internal state of the LSTM network is reset. The input data from all trials was 'mini-batched' into 2 training batches and 1 validation batch (499 samples each) that were used to update model weights on each model run (epoch). To format our data into equal length input and output mini batches for training and account for the output data being a one-time step shifted version of our input data, our lookback parameter must be one value less than a divisor of the trial length. For example, in our dataset (1500 sample length trials), a lookback parameter of 499 would result in the first mini batch input of samples [0:499] which will predict our reference output samples [1:500], our 2nd mini batch input data would include samples [501:999] and corresponding output [502:1000] and the last mini-batch input of samples [1001: 1499] predicts samples [1002: 1500]. This lookback parameter of 499 allows us to construct 3 mini-batches of shorter input and output data lengths which would be used to train and validate the RNN model (2:1 training: validation mini batch split). Similarly with the lookback parameter of 499 (2:1 training: validation mini-batch split) and 749 (1:1 training: validation split). Mean squared error was used as the LSTM loss function and ADAM as the optimization algorithm. The model was trained for at least 5000 iterations or until training and validation error converged (< 0.75°). The training resulted in a sample-specific dynamical model structure defined by a single set of LSTM network weights (W). The model's internal states capture trial-specific dynamics predicting the time evolution of joint kinematics; activation coefficients (H) and memory cell states (C) and are tuned based on kinematic input.

Model hyperparameter selection

We selected the hyperparameter values of 512 nodes in the LSTM layer and a 499-sample LSTM lookback length (number of samples preceding the current time point that is used to train the LSTM) were selected based on training and validation loss, as well as the ability to encode dynamics over short and long timescales. In two steps, we evaluated all pairs of the following hyperparameter values: 1024, 512, 256, and 128 LSTM nodes and 749, 499, and 249-sample lookback parameters. Because RNN performance can change with the parameters used to initialize the RNN, we fit an RNN gait dynamics model 20 times using random initial parameters, for each hyperparameter pair. First, we compared model training and validation loss for each hyperparameter pair: the 'best' hyperparameter pair would have low training and validation loss. The following [node-lookback] pairs were considered the *best* hyperparameter pairs: 512-499 (MSE_{train} = 0.010 ± 0.001 deg^2; MSE_{val} = 0.018 ± 0.000 deg^2), 256-749 (MSE_{train} = 0.010 ± 0.002

deg^2; MSE$_{val}$ = 0.015 ± 0.001 deg^2), 256-499 (MSE$_{train}$ = 0.010 ± 0.001 deg^2; MSE$_{val}$ = 0.017 ±
0.000 deg^2). (Fig. 1 - figure supplement 1, A). The training loss was not different between
hyperparameter pairs (p > 0.235). The validation loss differed between all three models (p <
0.001), with the 256-749 model having the lowest validation loss. However, if the differences in
validation loss of less than 0.003 deg^2 corresponded to meaningful differences in performance
was unclear.

Our second analysis was, therefore, used to compare the three hyperparameter pairs
deemed *best* in the prior analysis. Here, we evaluated the models' abilities to encode the average
dynamical behavior over long timescales (*long-time*) and the stride-to-stride behavior (*short-time*).
We defined the *best* model as the one with the highest long- and short-time performance. The
following analysis was performed for 10 of the 20 random initializations. For long- and short-time
analyses, we created a single set of *reference* dynamics as done in the manuscript: we performed
one time-step predictions over the full (1500-sample) time-series. This step provided best-case
predictions of the gait dynamics (Fig. 1 - figure supplement 1, B).

Long-time performance: We generated long-time predictions of each trial's gait signatures (RNN
latent states) by simulating each participant's gait dynamics forward in time, 1500 samples into
the future. Each simulation was initialized by setting the RNNs' latent states to those of the last
sample of the trial's reference dynamics and using the last sample of the trial's kinematics. We
then phase-averaged both the reference dynamics and the long-time predictions using the same
technique as described in the main manuscript. Long-time performance was defined as the
similarity of the phase-averaged latent states (*i.e.*, the gait signatures) between the reference and
the long-time predictions and was quantified using R^2. Note that using R^2 as a similarity metric,
rather than the Euclidean distance metric used in the main manuscript, was needed to compare
models with different numbers of nodes. Unlike R^2, Euclidean distances are sensitive to the
number of samples used to compare models, which would bias short- and long-time performance
towards models with fewer nodes. Low R^2 values between predictions indicates that the learned
dynamics are sufficiently complex to capture instantaneous gait dynamics but can also accurately
generate the time evolution of dynamics over the gait cycle - a major challenge in data-driven
models of locomotion (Maus et al., 2015; Rosenberg et al., 2020).

The 512-node model captured gait dynamics over long time scales significantly better (*i.e.*, more
accurate predictions of the time-varying dynamics) than the 249-node models (Fig. 1 - figure
supplement 1, B). For long-time predictions, the 512-node model predictions (R^2 = 0.50 ± 0.46)
were better than the 249-node 499-sample lookback model (ΔR^2 = 0.27 ± 0.06; p < 0.001;
independent-samples t-tests) and the 249-node 499-sample lookback model (ΔR^2 = 0.31 ± 0.07;
p < 0.001).

Short-time performance: We generated short-time predictions by simulating single strides in
each trial's time-series, initialized from the first sample of each stride. Initialization used the latent
RNN states and kinematics of the reference dynamics at the onset of a new stride (phase = 0
rad). For each initial condition, we integrated the dynamics forward in time, up to the onset of the
next stride. For each stride, we then compare the similarity of the reference dynamics to the
dynamics of the corresponding short-time prediction using R^2. Short-time performance was
quantified as the average R^2 value across trials for a single model and initialization.

The 512-node model captured gait dynamics over short time scales significantly better (*i.e.*, more
accurate predictions of the time-varying dynamics) than the 249-node models (Fig. 1 - figure

 supplement 1, B). For short-time predictions, the 512-node model predictions (R^2 = 0.11 ± 0.51)
were more accurate than the 249-node 499-sample lookback model (ΔR^2 = 0.51 ± 0.13; p = 0.055;
independent-samples t-tests) and the 249-node 499-sample lookback model (ΔR^2 = 0.34 ± 0.09;
p < 0.001). Based on difference in short- and long-time prediction performance, we selected the
512-node, 499-sample lookback hyperparameters for the RNN model.

Leave-one-out subject model evaluation for generalizability

Using the selected hyperparameters, 12 different models were trained where one different
subject (all 6 speed trials per subject) was held out for evaluation on each model run. The same
model architecture, training and validation setup was used as the original model trained using the
full dataset (12 subjects). The minimum training loss, validation loss, and overall evaluated test
loss for each model were extracted and box plots of each generated. The Wilcoxon Rank-Sum
Test statistic was used to compare the means. Each model was evaluated on the 6 held-out speed
trials from training and an average loss was calculated for each model. The reference kinematics,
externally driven and self-driven predictions of each of the 6 held-out trials per model were phase
averaged and R^2 between the phase averaged externally driven and long-time self-driven
predictions (*see Long-Time Performance section, above*) were calculated. Box plots for each
metric across the held-out trials were generated and the Wilcoxon Rank-Sum Test statistic used
to   compare   the   means.
884

Computing gait signatures from RNN internal states

To develop the gait signatures, we extract the activation and cell states from the LSTM
(denoted "H" and "C" respectively) which evolve over time (the course of the gait cycle) as the
kinematics of each trial are fed through the trained RNN. These H and C parameters represent
how the model's internal parameters change as it encodes the prediction of future kinematic
trajectories. The selected 512-node LSTM layer had 512 H parameters and 512 C parameters.
Time-varying gait signatures were computed by identifying dominant modes of variation in the
internal states using principal components analysis (PCA). A single PCA operation was used to
transform the internal states for all participants into a common basis. Consequently, inter-trial
differences in the time-varying activations of the principal components (modes) reflect differences
in the underlying dynamics of the individual(s). These activations constituted the time-varying
(1500 sample) *gait signatures*, which had the same dimension as the RNN's hidden layer (1024
units). However, the first six principal components accounted for 77% of the variance in the
internal states.

To compare gait signatures within and between individuals, we phase averaged each
trial's signatures across strides. Rather than linearly interpolating the data between foot contact
events before averaging, as is common in gait analysis (many refs), we computed a continuous
phase using the first 3 gait signature modes for each trial using Von Mises interpolation (Berman
et al., 2014). Compared to averaging across linearly interpolated strides, phase averaging is
expected to reduce the variance in the data at any point in the stride (Revzen and Guckenheimer,
2008; Rosenberg et al., 2020). As the domain for interpolation, we estimated the time-varying
phase for each trial separately using the Phaser algorithm, using the first 3 principal components
as phase variables (Revzen and Guckenheimer, 2008). To align phase estimates across trials,
we defined zero-phase as the maximum of the first principal component.

Gait event estimation of phase averaged signatures

The force plates embedded in the treadmill captured precise gait event timing information (left heel strike, right toe off, right heel strike, left toe off) across individuals' trials which we represented as a vector of 1's, 2's, 3's and 4's, respectively (ground truth markings for the 4 gait events). We leveraged the Phaser algorithm again (Revzen and Guckenheimer, 2008) to develop a phase estimator to transform these 4 gait events over time into gait events over phase. For each trial, we determined the mode phase that corresponded to each of the 4 gait events to gain a representation of where the 4 gait events occurred during phase averaged dynamics (0 - 2π) for each trial.

Interpolation of unseen speed gait signatures to reconstruct kinematics

To demonstrate the generalizability of gait dynamics, we show that linearly interpolating gait signatures to predict gait kinematics at new walking speeds is more accurate than linearly interpolating the kinematics themselves. We trained another RNN model with the same architecture and hyperparameters to the first model, however using only the 2 slowest and 2 fastest speeds from each participant (i.e., we held out the 2 middle speed trials from each participant). We then linearly interpolated the 2-middle speeds' internal states and ran the data through the trained RNN to reconstruct or predict kinematics. We compared the original phase averaged kinematics to the predicted kinematics resulting from each of the two linear interpolations (dynamics and kinematics) using the coefficient of determination. Furthermore, even when we reduced the dimensionality of the model's internal states from the full 1024 to the first 6 principal components (the selected dimension on the gait signatures), it still performed better than interpolating kinematics (also rank = 6).

Biomechanical interpretation of principal components of the gait signature

To reconstruct kinematics from the corresponding underlying dynamics (internal state representations), we restored our trained model's weights to a new model using the 'model.set' and 'model.get_weights' Keras built in functions. The function 'model.predict' takes in the hidden state values (Hs) only (first 512 of the 1024 internal-state time trajectories) and predicts the corresponding kinematics for the provided internal states. Using this framework, we provided this new model with independent principal component representations of individuals' hidden states and visualized the corresponding kinematics through stick figure movie representations of the resulting kinematics over the walking trials.

Predicting time evolution of kinematics from an initial posture (self-driving)

Our trained generative gait model can take in a single initial posture of size (6,1) corresponding to a single time point representation of each of the 6 joint angles to predict the next time step posture/kinematics using command 'model.predict'. To make further predictions, the predicted value is used as the new initial condition (posture) and predictions are made on a one-time step basis in a similar fashion for a pre-specified prediction length (self-driving).

Data and code availability

To ensure rigor, reproducibility, and promote open science, all software is shared under a *GNU GPL* 3.0 license and on GitHub. Links to Google Colab notebooks enable our software to be run on the cloud for users without computational resources. The gait signature code developed

960 Acknowledgements

We thank Kanishk Jain for providing feedback on the initial implementation of the model. TSW
was supported by the Alfred P. Sloan Foundation's Minority Ph.D. (MPHD) program: G-2019-
11435, NSF GRFP 1937971, and NICHD F31HD107968. MCR was supported by NICHD
F32HD108927. TMK was supported by NICHD R01HD095975. TSW, MCR, and LHT were
supported by the McCamish Foundation. LHT was supported by NSF CMMI 1762211 / 1761679.
TSW and GJB were supported by the Simons-Emory International Consortium on Motor Control
(Simons Foundation, 707102). All authors were supported by an Emory University Nexus/Synergy
II Grant.

970 Supplemental Figures

Figure 1- figure supplement 1: Comparison of model performance on training and validation loss (left), and
long- and short-time prediction performance. In both plots, small dots represent the average values across trials for
each random initialization of each model. Large dots and bars denote the average and standard deviation of model
performance metrics across initializations. Left: Training and validation loss (RMSE) for all 12 hyperparameter pairs.
Models in the lower-left consider are considered better. Right: Long- and short-time prediction performance (R^2) for the
3 hyperparameter pairs with the lowest training and validation loss. Models in the upper-right corner are considered
better.

Figure 1- figure supplement 2: RNN model training (green) and validation (blue) loss curves.

Figure 1- figure supplement 3: RNN dynamic learning generalizes across 12 leave-one-individual-out models.
The minimum train loss (blue) and validation loss (green) was low (< 0.02 degrees^2) for 5 models that were trained
each with a single able-bodied individual held out of the training data (A, i) and 7 models each with a stroke individual
held out of training data (A, ii). The magnitude and range of the test loss (evaluation of model on the held-out data)
(orange) was higher than the respective minimum training and validation losses for both held-out able-bodied (A, i) and
stroke (A, ii) models. The magnitude and range of test losses evaluated on held-out able-bodied individuals, however,
were lower than models evaluated on held-out stroke data. The models generate external predictions (blue) of held-
out test trials with higher R^2 values than that of self-driven predictions (red) in models evaluated on both able-bodied
(B, i) and stroke trials (B, ii). The models can generate external kinematic predictions (blue) of held-out able-bodied (B,
i) trials better than that of stroke (B, ii). Self-driven predictions of stroke kinematics were generally very low (R2 values
below 0.5). Models were incapable of generating self-driven predictions for 5 of 30 able-bodied trials and 16 of 42 stroke
trials (these R^2 values are not shown in B i and ii plots). (C) shows reference (black), externally driven (blue) and self-

driven (red) phase averaged kinematic predictions for an exemplary able-bodied trial (C, i) and exemplary stroke trial
(C, ii). Models can predict kinematics of held-out able-bodied trials better (higher R^2) than that of stroke.

Figure 1- figure supplement 4**: **Cumulative proportion of variance explained by the first 100 principal
components of gait dynamics. *Six (6) principal components (PCs) explained 77% of the variance in the gait*
dynamics. The top 6 dominant PCs were used to develop the gait signature.

Figure 2- figure supplement 1: Support vector machine cross-validation classification accuracy of four
different gait descriptors (discrete variables, gait signatures, kinematics, and a combination of kinematics &
joint velocity) for discrimination between: a) gait group (able-bodied vs. stroke) and B) individuals. *Using k =*
25 folds, RNN gait signatures distinguished between impaired and unimpaired gait with 100% accuracy, along with the
26 discrete variables (100%, p = 1), whereas kinematic (92.67 ± 0.15%, p < 0.05) and kinematics & velocity (88.67 ±
0.17%, p <0.05) discrimination were significantly lower. Using k = 6 folds, SVM classification of individuals was most
accurate using RNN gait signatures and discrete variables (100%), lower using kinematics (88.9 ± 0.13%, p = 0.061)
and significantly lower using a kinematics and velocity (68.10 ± 0.16%, p < 0.05).

1009

1010

1011

1012

References

1013

Ahamed T, Costa AC, Stephens GJ. 2021. Capturing the continuous complexity of behaviour in
Caenorhabditis elegans. *Nat Phys* **17**:275–283. doi:10.1038/s41567-020-01036-8
Alcantara RS, Edwards WB, Millet GY, Grabowski AM. 2022. Predicting continuous ground reaction
forces from accelerometers during uphill and downhill running: a recurrent neural network
solution. *PeerJ* **10**:e12752. doi:10.7717/peerj.12752
Allen JL, Kesar TM, Ting LH. 2019. Motor module generalization across balance and walking is impaired
after stroke. *J Neurophysiol* **122**:277–289. doi:10.1152/jn.00561.2018
Allen JL, Ting LH, Kesar TM. 2018. Gait Rehabilitation Using Functional Electrical Stimulation Induces
Changes in Ankle Muscle Coordination in Stroke Survivors: A Preliminary Study. *Front Neurol*
9:1127. doi:10.3389/fneur.2018.01127
Angelidis E, Buchholz E, Arreguit J, Rougé A, Stewart T, von Arnim A, Knoll A, Ijspeert A. 2021. A spiking
central pattern generator for the control of a simulated lamprey robot running on SpiNNaker
and Loihi neuromorphic boards. *Neuromorph Comput Eng* **1**:014005. doi:10.1088/2634-
4386/ac1b76
Attwood AS, Ludwig CJH, Penton-Voak IS, Poh J, Kwong ASF, Munafò MR. 2021. Effects of state anxiety
on gait: a 7.5% carbon dioxide challenge study. *Psychol Res* **85**:2444–2452. doi:10.1007/s00426-
020-01393-2
Aung MSH, Thies SB, Kenney LPJ, Howard D, Selles RW, Findlow AH, Goulermas JY. 2013. Automated
detection of instantaneous gait events using time frequency analysis and manifold embedding.
IEEE Trans Neural Syst Rehabil Eng **21**:908–916. doi:10.1109/TNSRE.2013.2239313
Awad LN, Palmer JA, Pohlig RT, Binder-Macleod SA, Reisman DS. 2015. Walking speed and step length
asymmetry modify the energy cost of walking after stroke. *Neurorehabil Neural Repair* **29**:416–
423. doi:10.1177/1545968314552528
Barton GJ, Hawken MB, Scott MA, Schwartz MH. 2012. Movement deviation profile: a measure of
distance from normality using a self-organizing neural network. *Hum Mov Sci* **31**:284–294.
doi:10.1016/j.humov.2010.06.003
Becchio C, Manera V, Sartori L, Cavallo A, Castiello U. 2012. Grasping intentions: from thought
experiments to empirical evidence. *Front Hum Neurosci* **6**:117. doi:10.3389/fnhum.2012.00117
Beer RF, Dewald JP, Rymer WZ. 2000. Deficits in the coordination of multijoint arm movements in
patients with hemiparesis: evidence for disturbed control of limb dynamics. *Exp Brain Res*
131:305–319. doi:10.1007/s002219900275
Berman GJ, Choi DM, Bialek W, Shaevitz JW. 2014. Mapping the stereotyped behaviour of freely moving
fruit flies. *Journal of The Royal Society Interface* **11**:20140672. doi:10.1098/rsif.2014.0672
Bernshteĭn NA. 1967. The co-ordination and regulation of movements, 1st English ed. ed. Oxford, New
York: Pergamon Press.
Berrueta TA, Pervan A, Fitzsimons K, Murphey TD. 2019. Dynamical System Segmentation for
Information Measures in Motion. *IEEE Robotics and Automation Letters* **4**:169–176.
doi:10.1109/LRA.2018.2884091
Bongard J, Lipson H. 2007. Automated reverse engineering of nonlinear dynamical systems. *Proc Natl*
Acad Sci U S A **104**:9943–9948. doi:10.1073/pnas.0609476104
Bowden MG, Balasubramanian CK, Behrman AL, Kautz SA. 2008. Validation of a speed-based
classification system using quantitative measures of walking performance poststroke.
Neurorehabil Neural Repair **22**:672–675. doi:10.1177/1545968308318837

Brown DD, Wijffels G, Meulenbroek RGJ. 2021. Individual Differences in Sequential Movement
Coordination in Hip-Hop Dance: Capturing Joint Articulation in Practicing the Wave. *Front*
Psychol **12**:731901. doi:10.3389/fpsyg.2021.731901
Camargo J, Molinaro D, Young A. 2022. Predicting biological joint moment during multiple ambulation
tasks. *J Biomech* **134**:111020. doi:10.1016/j.jbiomech.2022.111020
Cao Z, Hidalgo G, Simon T, Wei S-E, Sheikh Y. 2019. OpenPose: Realtime Multi-Person 2D Pose
Estimation using Part Affinity Fields. doi:10.48550/arXiv.1812.08008
Castano-Pino YJ, Gonzalez MC, Quintana-Pena V, Valderrama J, Munoz B, Orozco J, Navarro A. 2020.
Automatic Gait Phases Detection in Parkinson Disease: A Comparative Study. *Annu Int Conf IEEE*
Eng Med Biol Soc **2020**:798–802. doi:10.1109/EMBC44109.2020.9175268
Chen G, Patten C, Kothari DH, Zajac FE. 2005. Gait differences between individuals with post-stroke
hemiparesis and non-disabled controls at matched speeds. *Gait & Posture* **22**:51–56.
doi:10.1016/j.gaitpost.2004.06.009
Chia Bejarano N, Ambrosini E, Pedrocchi A, Ferrigno G, Monticone M, Ferrante S. 2015. A novel
adaptive, real-time algorithm to detect gait events from wearable sensors. *IEEE Trans Neural*
Syst Rehabil Eng **23**:413–422. doi:10.1109/TNSRE.2014.2337914
Chvatal SA, Ting LH. 2013. Common muscle synergies for balance and walking. *Front Comput Neurosci*
7:48. doi:10.3389/fncom.2013.00048
Clark DJ, Ting LH, Zajac FE, Neptune RR, Kautz SA. 2010. Merging of healthy motor modules predicts
reduced locomotor performance and muscle coordination complexity post-stroke. *J*
Neurophysiol **103**:844–857. doi:10.1152/jn.00825.2009
Correa TA, Baker R, Graham HK, Pandy MG. 2011. Accuracy of generic musculoskeletal models in
predicting the functional roles of muscles in human gait. *J Biomech* **44**:2096–2105.
doi:10.1016/j.jbiomech.2011.05.023
Cox MAA, Cox TF. 2008. Multidimensional Scaling In: Chen C, Härdle W, Unwin A, editors. Handbook of
Data Visualization, Springer Handbooks Comp.Statistics. Berlin, Heidelberg: Springer. pp. 315–
347. doi:10.1007/978-3-540-33037-0_14
De Groote F, Falisse A. 2021. Perspective on musculoskeletal modelling and predictive simulations of
human movement to assess the neuromechanics of gait. *Proceedings of the Royal Society B:*
Biological Sciences **288**:20202432. doi:10.1098/rspb.2020.2432
De Groote F, Jonkers I, Duysens J. 2014. Task constraints and minimization of muscle effort result in a
small number of muscle synergies during gait. *Frontiers in Computational Neuroscience* **8**.
Drnach L, Allen JL, Essa I, Ting LH. 2019. A Data-Driven Predictive Model of Individual-Specific Effects of
FES on Human Gait Dynamics2019 International Conference on Robotics and Automation (ICRA).
Presented at the 2019 International Conference on Robotics and Automation (ICRA). pp. 5090–
5096. doi:10.1109/ICRA.2019.8794304
D'souza J, Natarajan DM, Kumaran D DS. 2020. Does the Environment Cause Changes in Hemiparetic
Lower Limb Muscle Activity and Gait Velocity During Walking in Stroke Survivors? *J Stroke*
Cerebrovasc Dis **29**:105174. doi:10.1016/j.jstrokecerebrovasdis.2020.105174
Dzeladini F, van den Kieboom J, Ijspeert A. 2014. The contribution of a central pattern generator in a
reflex-based neuromuscular model. *Front Hum Neurosci* **8**:371. doi:10.3389/fnhum.2014.00371
Edey R, Yon D, Cook J, Dumontheil I, Press C. 2017. Our own action kinematics predict the perceived
affective states of others. *J Exp Psychol Hum Percept Perform* **43**:1263–1268.
doi:10.1037/xhp0000423
Elkjær E, Mikkelsen MB, Michalak J, Mennin DS, O'Toole MS. 2022. Motor alterations in depression and
anxiety disorders: A systematic review and meta-analysis. *J Affect Disord* **317**:373–387.
doi:10.1016/j.jad.2022.08.060

Falisse A, Pitto L, Kainz H, Hoang H, Wesseling M, Van Rossom S, Papageorgiou E, Bar-On L, Hallemans A,
Desloovere K, Molenaers G, Van Campenhout A, De Groote F, Jonkers I. 2020. Physics-Based
Simulations to Predict the Differential Effects of Motor Control and Musculoskeletal Deficits on
Gait Dysfunction in Cerebral Palsy: A Retrospective Case Study. *Front Hum Neurosci* **14**:40.
doi:10.3389/fnhum.2020.00040

Falisse A, Serrancolí G, Dembia CL, Gillis J, Jonkers I, De Groote F. 2019. Rapid predictive simulations with
complex musculoskeletal models suggest that diverse healthy and pathological human gaits can
emerge from similar control strategies. *Journal of The Royal Society Interface* **16**:20190402.
doi:10.1098/rsif.2019.0402

Ferrante S, Chia Bejarano N, Ambrosini E, Nardone A, Turcato AM, Monticone M, Ferrigno G, Pedrocchi
A. 2016. A Personalized Multi-Channel FES Controller Based on Muscle Synergies to Support Gait
Rehabilitation after Stroke. *Frontiers in Neuroscience* **10**.

Fregly BJ, Reinbolt JA, Rooney KL, Mitchell KH, Chmielewski TL. 2007. Design of patient-specific gait
modifications for knee osteoarthritis rehabilitation. *IEEE Trans Biomed Eng* **54**:1687–1695.
doi:10.1109/tbme.2007.891934

Frisk RF, Lorentzen J, Nielsen JB. 2019. Contribution of corticospinal drive to ankle plantar flexor muscle
activation during gait in adults with cerebral palsy. *Exp Brain Res* **237**:1457–1467.
doi:10.1007/s00221-019-05520-3

Fugl-Meyer AR, Jääskö L, Leyman I, Olsson S, Steglind S. 1975. The post-stroke hemiplegic patient. 1. a
method for evaluation of physical performance. *Scand J Rehabil Med* **7**:13–31.

Garcia SA, Johnson AK, Brown SR, Washabaugh EP, Krishnan C, Palmieri-Smith RM. 2022. Dynamic knee
stiffness during walking is increased in individuals with anterior cruciate ligament
reconstruction. *J Biomech* **146**:111400. doi:10.1016/j.jbiomech.2022.111400

Genthe K, Schenck C, Eicholtz S, Zajac-Cox L, Wolf S, Kesar TM. 2018. Effects of real-time gait
biofeedback on paretic propulsion and gait biomechanics in individuals post-stroke. *Top Stroke
Rehabil* **25**:186–193. doi:10.1080/10749357.2018.1436384

Geyer H, Herr H. 2010. A Muscle-Reflex Model That Encodes Principles of Legged Mechanics Produces
Human Walking Dynamics and Muscle Activities. *IEEE Trans Neural Syst Rehabil Eng* **18**:263–273.
doi:10.1109/TNSRE.2010.2047592

Giarmatzis G, Zacharaki EI, Moustakas K. 2020. Real-Time Prediction of Joint Forces by Motion Capture
and Machine Learning. *Sensors (Basel)* **20**:6933. doi:10.3390/s20236933

Gross MM, Crane EA, Fredrickson BL. 2012. Effort-Shape and kinematic assessment of bodily expression
of emotion during gait. *Human Movement Science* **31**:202–221.
doi:10.1016/j.humov.2011.05.001

Habersack A, Fischerauer SF, Kraus T, Holzer H-P, Svehlik M. 2022. Kinematic and Kinetic Gait
Parameters Can Distinguish between Idiopathic and Neurologic Toe-Walking. *Int J Environ Res
Public Health* **19**:804. doi:10.3390/ijerph19020804

Hainisch R, Kranzl A, Lin Y-C, Pandy MG, Gfoehler M. 2021. A generic musculoskeletal model of the
juvenile lower limb for biomechanical analyses of gait. *Computer Methods in Biomechanics and
Biomedical Engineering* **24**:349–357. doi:10.1080/10255842.2020.1817405

Halilaj E, Rajagopal A, Fiterau M, Hicks JL, Hastie TJ, Delp SL. 2018. Machine learning in human
movement biomechanics: Best practices, common pitfalls, and new opportunities. *J Biomech
* **81**:1–11. doi:10.1016/j.jbiomech.2018.09.009

Hausdorff JM, Purdon PL, Peng CK, Ladin Z, Wei JY, Goldberger AL. 1996. Fractal dynamics of human gait:
stability of long-range correlations in stride interval fluctuations. *J Appl Physiol (1985)* **80**:1448–
1457. doi:10.1152/jappl.1996.80.5.1448

Heinik J, Werner P, Dekel T, Gurevitz I, Rosenblum S. 2010. Computerized kinematic analysis of the clock
drawing task in elderly people with mild major depressive disorder: an exploratory study. *Int*
Psychogeriatr **22**:479–488. doi:10.1017/S1041610209991360

Holmes P, Full RJ, Koditschek D, Guckenheimer J. 2006. The Dynamics of Legged Locomotion: Models,
Analyses, and Challenges. *SIAM Review* **48**:207–304.

Hornby TG, Reisman DS, Ward IG, Scheets PL, Miller A, Haddad D, Fox EJ, Fritz NE, Hawkins K, Henderson
CE, Hendron KL, Holleran CL, Lynskey JE, Walter A, and the Locomotor CPG Appraisal Team.
2020. Clinical Practice Guideline to Improve Locomotor Function Following Chronic Stroke,
Incomplete Spinal Cord Injury, and Brain Injury. *J Neurol Phys Ther* **44**:49–100.
doi:10.1097/NPT.0000000000000303

Horst F, Lapuschkin S, Samek W, Müller K-R, Schöllhorn WI. 2019. Explaining the unique nature of
individual gait patterns with deep learning. *Scientific Reports* **9**:2391. doi:10.1038/s41598-019-
38748-8

Ijmker T, Lamoth CJC. 2012. Gait and cognition: the relationship between gait stability and variability
with executive function in persons with and without dementia. *Gait Posture* **35**:126–130.
doi:10.1016/j.gaitpost.2011.08.022

Ivanenko YP, Grasso R, Zago M, Molinari M, Scivoletto G, Castellano V, Macellari V, Lacquaniti F. 2003.
Temporal components of the motor patterns expressed by the human spinal cord reflect foot
kinematics. *J Neurophysiol* **90**:3555–3565. doi:10.1152/jn.00223.2003

Jang J, Wikstrom EA. 2022. Ankle joint contact force profiles differ between those with and without
chronic ankle instability during walking. *Gait Posture* **100**:1–7.
doi:10.1016/j.gaitpost.2022.11.012

Johnson KL, Gill S, Reichman V, Tassinary LG. 2007. Swagger, sway, and sexuality: Judging sexual
orientation from body motion and morphology. *Journal of Personality and Social Psychology*
93:321. doi:10.1037/0022-3514.93.3.321

Jonkers I, Delp S, Patten C. 2009. Capacity to increase walking speed is limited by impaired hip and ankle
power generation in lower functioning persons post-stroke. *Gait Posture* **29**:129–37.

Kesar TM, Binder-Macleod SA, Hicks GE, Reisman DS. 2011a. Minimal detectable change for gait
variables collected during treadmill walking in individuals post-stroke. *Gait Posture* **33**:314–317.
doi:10.1016/j.gaitpost.2010.11.024

Kesar TM, Perumal R, Jancosko A, Reisman DS, Rudolph KS, Higginson JS, Binder-Macleod SA. 2010.
Novel patterns of functional electrical stimulation have an immediate effect on dorsiflexor
muscle function during gait for people poststroke. *Phys Ther* **90**:55–66.
doi:10.2522/ptj.20090140

Kesar TM, Perumal R, Reisman DS, Jancosko A, Rudolph KS, Higginson JS, Binder-Macleod SA. 2009.
Functional electrical stimulation of ankle plantarflexor and dorsiflexor muscles: effects on
poststroke gait. *Stroke* **40**:3821–3827. doi:10.1161/STROKEAHA.109.560375

Kesar TM, Reisman DS, Perumal R, Jancosko AM, Higginson JS, Rudolph KS, Binder-Macleod SA. 2011b.
Combined effects of fast treadmill walking and functional electrical stimulation on post-stroke
gait. *Gait Posture* **33**:309–313. doi:10.1016/j.gaitpost.2010.11.019

Kettlety SA, Finley JM, Reisman DS, Schweighofer N, Leech KA. 2022. Speed-dependent biomechanical
changes vary across individual gait metrics post-stroke relative to neurotypical adults.
doi:10.1101/2022.04.01.486769

Krogh S, Aagaard P, Jønsson AB, Figlewski K, Kasch H. 2022. Effects of repetitive transcranial magnetic
stimulation on recovery in lower limb muscle strength and gait function following spinal cord
injury: a randomized controlled trial. *Spinal Cord* **60**:135–141. doi:10.1038/s41393-021-00703-8

Kuo AD. 2002. The Relative Roles of Feedforward and Feedback in the Control of Rhythmic Movements.
Motor Control **6**:129–145. doi:10.1123/mcj.6.2.129

Kuo AD, Donelan JM. 2010. Dynamic Principles of Gait and Their Clinical Implications. *Physical Therapy* **90**:157–174. doi:10.2522/ptj.20090125

Larsen RJ, Queen RM, Schmitt D. 2022. Adaptive locomotion: Foot strike pattern and limb mechanical stiffness while running over an obstacle. *J Biomech* **143**:111283. doi:10.1016/j.jbiomech.2022.111283

Little VL, McGuirk TE, Perry LA, Patten C. 2018. Pelvic excursion during walking post-stroke: A novel classification system. *Gait Posture* **62**:395–404. doi:10.1016/j.gaitpost.2018.03.052

Mannini A, Trojaniello D, Cereatti A, Sabatini AM. 2016. A Machine Learning Framework for Gait Classification Using Inertial Sensors: Application to Elderly, Post-Stroke and Huntington's Disease Patients. *Sensors (Basel)* **16**. doi:10.3390/s16010134

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat Neurosci* **21**:1281–1289. doi:10.1038/s41593-018-0209-y

Maus H-M, Revzen S, Guckenheimer J, Ludwig C, Reger J, Seyfarth A. 2015. Constructing predictive models of human running. *J R Soc Interface* **12**:20140899. doi:10.1098/rsif.2014.0899

McCrea DA, Rybak IA. 2008. Organization of mammalian locomotor rhythm and pattern generation. *Brain Research Reviews*, Networks in Motion **57**:134–146. doi:10.1016/j.brainresrev.2007.08.006

Meyer AJ, Eskinazi I, Jackson JN, Rao AV, Patten C, Fregly BJ. 2016. Muscle Synergies Facilitate Computational Prediction of Subject-Specific Walking Motions. *Front Bioeng Biotechnol* **4**:77. doi:10.3389/fbioe.2016.00077

Mikolajczyk T, Ciobanu I, Badea DI, Iliescu A, Pizzamiglio S, Schauer T, Seel T, Seiciu PL, Turner DL, Berteanu M. 2018. Advanced technology for gait rehabilitation: An overview. *Advances in Mechanical Engineering* **10**:1687814018783627. doi:10.1177/1687814018783627

Milner CE, Foch E, Gonzales JM, Petersen D. 2022. Biomechanics associated with tibial stress fracture in runners: A systematic review and meta-analysis. *J Sport Health Sci* S2095-2546(22)00116–8. doi:10.1016/j.jshs.2022.12.002

Moura Coelho R, Hirai H, Martins J, Krebs HI. 2022. Biomarkers for rhythmic and discrete dynamic primitives in locomotion. *Sci Rep* **12**:20165. doi:10.1038/s41598-022-24565-z

Neave N, McCarty K, Freynik J, Caplan N, Hönekopp J, Fink B. 2011. Male dance moves that catch a woman's eye. *Biol Lett* **7**:221–224. doi:10.1098/rsbl.2010.0619

Pan Y-T, Kang I, Joh J, Kim P, Herrin KR, Kesar TM, Sawicki GS, Young AJ. 2022. Effects of Bilateral Assistance for Hemiparetic Gait Post-Stroke Using a Powered Hip Exoskeleton. *Ann Biomed Eng*. doi:10.1007/s10439-022-03041-9

Pandarinath C, O'Shea DJ, Collins J, Jozefowicz R, Stavisky SD, Kao JC, Trautmann EM, Kaufman MT, Ryu SI, Hochberg LR, Henderson JM, Shenoy KV, Abbott LF, Sussillo D. 2018. Inferring single-trial neural population dynamics using sequential auto-encoders. *Nat Methods* **15**:805–815. doi:10.1038/s41592-018-0109-9

Patterson KK, Parafianowicz I, Danells CJ, Closson V, Verrier MC, Staines WR, Black SE, McIlroy WE. 2008. Gait Asymmetry in Community-Ambulating Stroke Survivors. *Archives of Physical Medicine and Rehabilitation* **89**:304–310. doi:10.1016/j.apmr.2007.08.142

Payne AM, Sawers A, Allen JL, Stapley PJ, Macpherson JM, Ting LH. 2020. Reorganization of motor modules for standing reactive balance recovery following pyridoxine-induced large-fiber peripheral sensory neuropathy in cats. *J Neurophysiol* **124**:868–882. doi:10.1152/jn.00739.2019

Pitto L, Kainz H, Falisse A, Wesseling M, Van Rossom S, Hoang H, Papageorgiou E, Hallemans A, Desloovere K, Molenaers G, Van Campenhout A, De Groote F, Jonkers I. 2019a. SimCP: A Simulation Platform to Predict Gait Performance Following Orthopedic Intervention in Children With Cerebral Palsy. *Front Neurorobot* **13**:54. doi:10.3389/fnbot.2019.00054

Pitto L, Kainz H, Falisse A, Wesseling M, Van Rossom S, Hoang H, Papageorgiou E, Hallemans A,
Desloovere K, Molenaers G, Van Campenhout A, De Groote F, Jonkers I. 2019b. SimCP: A
Simulation Platform to Predict Gait Performance Following Orthopedic Intervention in Children
With Cerebral Palsy. *Frontiers in Neurorobotics* **13**.
Pollick FE, Kay JW, Heim K, Stringer R. 2005. Gender recognition from point-light walkers. *Journal of
Experimental Psychology: Human Perception and Performance* **31**:1247. doi:10.1037/0096-
1523.31.6.1247
Porciuncula F, Roto AV, Kumar D, Davis I, Roy S, Walsh CJ, Awad LN. 2018. Wearable Movement Sensors
for Rehabilitation: A Focused Review of Technological and Clinical Advances. *PM R* **10**:S220–
S232. doi:10.1016/j.pmrj.2018.06.013
Prinz AA, Bucher D, Marder E. 2004. Similar network activity from disparate circuit parameters. *Nat
Neurosci* **7**:1345–1352. doi:10.1038/nn1352
Prosser LA, Atkinson HL, Alfano JM, Kessler SK, Ichord RB. 2022. Walking speed influences
spatiotemporal but not symmetry measures of gait in children and adolescents with hemiplegia.
Gait Posture **98**:233–236. doi:10.1016/j.gaitpost.2022.09.088
Reisman DS, Bastian AJ, Morton SM. 2010. Neurophysiologic and rehabilitation insights from the split-
belt and other locomotor adaptation paradigms. *Phys Ther* **90**:187–195.
doi:10.2522/ptj.20090073
Revzen S, Guckenheimer JM. 2008. Estimating the phase of synchronized oscillators. *Phys Rev E Stat
Nonlin Soft Matter Phys* **78**:051907. doi:10.1103/PhysRevE.78.051907
Ries AJ, Novacheck TF, Schwartz MH. 2014. A data driven model for optimal orthosis selection in
children with cerebral palsy. *Gait Posture* **40**:539–544. doi:10.1016/j.gaitpost.2014.06.011
Rinaldi LA, Monaco V. 2013. Spatio-temporal parameters and intralimb coordination patterns describing
hemiparetic locomotion at controlled speed. *J NeuroEngineering Rehabil* **10**:53.
doi:10.1186/1743-0003-10-53
Rosenberg MC, Banjanin BS, Burden SA, Steele KM. 2020. Predicting walking response to ankle
exoskeletons using data-driven models. *J R Soc Interface* **17**:20200487.
doi:10.1098/rsif.2020.0487
Russo Y, Stuart S, Silva-Batista C, Brumbach B, Vannozzi G, Mancini M. 2022. Does visual cueing improve
gait initiation in people with Parkinson's disease? *Hum Mov Sci* **84**:102970.
doi:10.1016/j.humov.2022.102970
Sánchez N, Acosta AM, López-Rosado R, Dewald JPA. 2018. Neural Constraints Affect the Ability to
Generate Hip Abduction Torques When Combined With Hip Extension or Ankle Plantarflexion in
Chronic Hemiparetic Stroke. *Front Neurol* **9**. doi:10.3389/fneur.2018.00564
Sanchez-Gonzalez A, Godwin J, Pfaff T, Ying R, Leskovec J, Battaglia PW. 2020. Learning to Simulate
Complex Physics with Graph Networks. doi:10.48550/arXiv.2002.09405
Santhiranayagam BK, Lai DTH, Sparrow WA, Begg RK. 2015. A machine learning approach to estimate
Minimum Toe Clearance using Inertial Measurement Units. *J Biomech* **48**:4309–4316.
doi:10.1016/j.jbiomech.2015.10.040
Sawers A, Bhattacharjee T, McKay JL, Hackney ME, Kemp CC, Ting LH. 2017. Small forces that differ with
prior motor experience can communicate movement goals during human-human physical
interaction. *J Neuroeng Rehabil* **14**:8. doi:10.1186/s12984-017-0217-2
Schutte LM, Narayanan U, Stout JL, Selber P, Gage JR, Schwartz MH. 2000. An index for quantifying
deviations from normal gait. *Gait and Posture* 7.
Shin SY, Kim Y, Jayaraman A, Park H-S. 2021. Relationship between gait quality measures and modular
neuromuscular control parameters in chronic post-stroke individuals. *J Neuroeng Rehabil* **18**:58.
doi:10.1186/s12984-021-00860-0

Steele KM, Munger ME, Peters KM, Shuman BR, Schwartz MH. 2019. Repeatability of electromyography
recordings and muscle synergies during gait among children with cerebral palsy. *Gait Posture*
67:290–295. doi:10.1016/j.gaitpost.2018.10.009
Steele KM, Rozumalski A, Schwartz MH. 2015. Muscle synergies and complexity of neuromuscular
control during gait in cerebral palsy. *Dev Med Child Neurol* **57**:1176–1182.
doi:10.1111/dmcn.12826
Stenum J, Cherry-Allen KM, Pyles CO, Reetzke RD, Vignos MF, Roemmich RT. 2021. Applications of Pose
Estimation in Human Health and Performance across the Lifespan. *Sensors (Basel)* **21**:7315.
doi:10.3390/s21217315
Steptoe A, Zaninotto P. 2020. Lower socioeconomic status and the acceleration of aging: An outcome-
wide analysis. *Proceedings of the National Academy of Sciences* **117**:14911–14917.
doi:10.1073/pnas.1915741117
Sussillo D, Churchland MM, Kaufman MT, Shenoy KV. 2015. A neural network that finds a naturalistic
solution for the production of muscle activity. *Nat Neurosci* **18**:1025–1033. doi:10.1038/nn.4042
Sutskever I, Vinyals O, Le QV. 2014. Sequence to Sequence Learning with Neural Networks.
arXiv:14093215 [cs].
Taga G, Yamaguchi Y, Shimizu H. 1991. Self-organized control of bipedal locomotion by neural oscillators
in unpredictable environment. *Biol Cybern* **65**:147–159. doi:10.1007/BF00198086
Ting LH, Chiel HJ, Trumbower RD, Allen JL, McKay JL, Hackney ME, Kesar TM. 2015. Neuromechanical
principles underlying movement modularity and their implications for rehabilitation. *Neuron*
86:38–54. doi:10.1016/j.neuron.2015.02.042
Torres-Oviedo G, Macpherson JM, Ting LH. 2006. Muscle synergy organization is robust across a variety
of postural perturbations. *J Neurophysiol* **96**:1530–1546. doi:10.1152/jn.00810.2005
Torres-Oviedo G, Ting LH. 2010. Subject-specific muscle synergies in human balance control are
consistent across different biomechanical contexts. *J Neurophysiol* **103**:3084–3098.
doi:10.1152/jn.00960.2009
Troisi Lopez E, Sorrentino P, Liparoti M, Minino R, Polverino A, Romano A, Carotenuto A, Amico E,
Sorrentino G. 2022. The kinectome: A comprehensive kinematic map of human motion in health
and disease. *Ann N Y Acad Sci* **1516**:247–261. doi:10.1111/nyas.14860
Troje NF. 2002. Decomposing biological motion: A framework for analysis and synthesis of human gait
patterns. *Journal of Vision* **2**:2. doi:10.1167/2.5.2
Uhlrich SD, Falisse A, Kidziński Ł, Muccini J, Ko M, Chaudhari AS, Hicks JL, Delp SL. 2022. OpenCap: 3D
human movement dynamics from smartphone videos (preprint). Bioengineering.
doi:10.1101/2022.07.07.499061
van Hedel HJA, Tomatis L, Müller R. 2006. Modulation of leg muscle activity and gait kinematics by
walking speed and bodyweight unloading. *Gait Posture* **24**:35–45.
doi:10.1016/j.gaitpost.2005.06.015
Vyas S, Golub MD, Sussillo D, Shenoy KV. 2020. Computation Through Neural Population Dynamics.
Annu Rev Neurosci **43**:249–275. doi:10.1146/annurev-neuro-092619-094115
Walter JP, Kinney AL, Banks SA, D'Lima DD, Besier TF, Lloyd DG, Fregly BJ. 2014. Muscle synergies may
improve optimization prediction of knee contact forces during walking. *J Biomech Eng*
136:021031. doi:10.1115/1.4026428
Wang S, Pai Y-C, Bhatt T. 2022. Neuromuscular mechanisms of motor adaptation to repeated gait-slip
perturbations in older adults. *Sci Rep* **12**:19851. doi:10.1038/s41598-022-23051-w
Wang Y, Srinivasan M. 2013. System Identification and Stability Analysis of Steady Human Walking and
the Swing Leg Dynamics. Presented at the ASME 2012 5th Annual Dynamic Systems and Control
Conference joint with the JSME 2012 11th Motion and Vibration Conference. American Society
of Mechanical Engineers Digital Collection. pp. 19–23. doi:10.1115/DSCC2012-MOVIC2012-8663

Wonsetler EC, Bowden MG. 2017. A systematic review of mechanisms of gait speed change post-stroke.
Part 2: Exercise capacity, muscle activation, kinetics, and kinematics. *Top Stroke Rehabil* **24**:394–
403. doi:10.1080/10749357.2017.1282413
Young DR, Banks CL, McGuirk TE, Patten C. 2022. Evidence for shared neural information between
muscle synergies and corticospinal efficacy. *Sci Rep* **12**:8953. doi:10.1038/s41598-022-12225-1
Zajac FE, Neptune RR, Kautz SA. 2002. Biomechanics and muscle coordination of human walking. Part I:
introduction to concepts, power transfer, dynamics and simulations. *Gait Posture* **16**:215–232.
doi:10.1016/s0966-6362(02)00068-1
Zaninotto P, Sacker A, Head J. 2013. Relationship between wealth and age trajectories of walking speed
among older adults: evidence from the English Longitudinal Study of Ageing. *J Gerontol A Biol Sci*
Med Sci **68**:1525–1531. doi:10.1093/gerona/glt058
Zhang J, Lockhart TE, Soangra R. 2014. Classifying lower extremity muscle fatigue during walking using
machine learning and inertial sensors. *Ann Biomed Eng* **42**:600–612. doi:10.1007/s10439-013-
0917-0
1355
1356