Pheromone-based animal communication influences the production of somatic extracellular vesicles

Agata Szczepańska[1#], Katarzyna Olek[2#], Klaudia Kołodziejska[1], Jingfang Yu[3], Abdulrahman Tudu Ibrahim[1,4], Laura Adamkiewicz[1], Frank C. Schroeder[3], Wojciech Pokrzywa[2*], Michał Turek[1*]

Affiliations:

[1]*Laboratory of Animal Molecular Physiology, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland*

[2]*Laboratory of Protein Metabolism, International Institute of Molecular and Cell Biology in Warsaw, Poland.*

[3]*Boyce Thompson Institute and Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853, USA*

[4]*Faculty of Chemistry, Warsaw University of Technology, Warsaw, Poland*

[#] These authors contributed equally

[*] Co-corresponding authors

Correspondence should be directed to:

MT: m.turek@ibb.waw.pl

WP: wpokrzywa@iimcb.gov.pl

Abstract

Extracellular vesicles (EVs) are involved in multiple biological processes; however, there is

limited knowledge of the influence of environmental factors or other individuals in a population

on EV-regulated systems. The largest evolutionarily conserved EVs, exophers, are a

component of the *C. elegans* maternal somatic tissue resource management system induced by

embryos developing *in utero*, and thus progeny of individuals with active exopher biogenesis

(exophergenesis) appear to be privileged. Using this model, we investigated the inter-tissue and

social regulatory mechanisms of exophergenesis. We found that the predominant male

pheromone, ascr#10, increases exopher production in hermaphrodites via the G-protein-

coupled receptor STR-173 in the ASK sensory neurons. In contrast, pheromones from other

hermaphrodites in the population temper exophergenesis. Within the hermaphrodite, an

increase in embryo accumulation drives pro-exopher signals, and the internal sensory neurons

AQR, PQR, and URX play a central role in modulating exopher levels. This intricate process

is regulated partly via the neuropeptides FLP-8 and FLP-21, which originate from the URX

and AQR/PQR/URX neurons, respectively. Our results reveal a regulatory network integrating

internal and external cues, including control of somatic EVs production by the nervous system

in response to social signals.

Keywords

C. elegans, muscles, extracellular vesicles, exophers, pheromones, olfactory neurons

Introduction

Extracellular vesicles (EVs) are lipid-bilayer-enclosed particles released by most cell types. Two major EV types can be distinguished based on their biogenesis; endosome-derived exosomes and membrane-derived ectosomes[1]. EVs can be employed by cells to remove unwanted biological material, such as misfolded proteins and damaged organelles, or to transport cargo, including proteins and nucleic acids, enabling exchange and communication between cells. Therefore, they are critical in multiple physiological processes and pathological states involving disrupted cellular homeostasis[2–5]. A recently discovered class of the largest membrane-derived evolutionarily conserved EVs, termed exophers, mediate both the elimination of waste from cells and the exchange of biological material between tissues - they were shown to play a significant role in cellular stress response, tissue homeostasis, and organismal reproduction[6–9]. Previous studies have spotlighted the role of exophers in safeguarding neuronal activity in *C. elegans* against proteotoxic stress by expelling cellular waste into surrounding tissues[7]. This mechanism is also mirrored in mammalian contexts, where Nicolás-Ávila et al. discovered that mouse cardiomyocytes release exophers loaded with defective mitochondria, thereby preventing extracellular waste accumulation and consequent inflammations, thus upholding metabolic balance within the heart[9]. However, the biological roles of exophers extend beyond the elimination of superfluous cellular components. In our previous work, we showed that the body wall muscles (BWMs) of *C. elegans* release exophers that transport muscle-synthesized yolk proteins to support offspring development, increasing their odds of development and survival[6]. However, how exophergenesis is regulated in response to external factors impacting animal development and reproduction is unclear.

C. elegans exhibits a wide range of social behaviors governed by chemically diverse pheromones. Interwoven with sensory neuron responses, this pheromonal communication influences key developmental factors like growth, generation time, and maternal provisioning

[10–12]. Building on this premise, our present study delves into understanding how these social

cues modulate exophergenesis. Our findings reveal intriguing dynamics – the male-released

pheromone, including ascr#10, promotes exopher formation, whereas hermaphrodite-released

pheromones, including ascr#18, do the opposite, and thus population density plays a vital role

in regulating vesicle production. We determined that various sensory neurons and the G-protein

coupled receptor STR-173 regulate exophergenesis in response to environmental stimuli and

pheromones. We also demonstrated that the AQR/PQR/URX neurons, which are directly

exposed to the body cavity and monitor the worm's body interior, restrict muscle exopher

production. Taken together, we uncovered the basis for exophergenesis regulation to tune its

activity to external conditions. To our knowledge, this study is the first to report how animal

communication influences somatic EVs production. This study provides a deeper

understanding of *C. elegans* physiology and could illuminate physiological mechanisms

preserved across various species.

Results

Male pheromones promote exopher generation in hermaphrodite muscles.

Since muscle exophers mediate the transfer of maternal resources to offspring supporting their

development, we hypothesized that exophergenesis (Fig. 1a) is regulated by metabolites-

derived social cues generated within the worm population. We probed this by investigating

how the presence of males could impact exopher production. To this end, we examined the

number of exophers in *him-5* mutants, characterized by a marked increase in the percentage of

males in the population (approximately 33%, compared to 0.3% in the wild type)[13].

Interestingly, *him-5* mutant hermaphrodites co-cultured with *him-5* mutant males until the L4

stage and then transferred to a male-free plate (Fig. 1b) produce approximately 2.5 times more

exophers than wild-type hermaphrodites grown on a male-free plate (Fig. 1c). This

augmentation could be mediated by the previously postulated embryo-maternal signaling[6] as

him-5 mutant hermaphrodites contain 26% more embryos *in utero* than their wild-type

counterparts (Fig. 1d). To exclude the possibility that the observed increase in exopher numbers

resulted from the *him-5* mutation rather than male presence, we cultured wild-type

hermaphrodites on plates conditioned with *him-5* mutants or wild-type males for 48 hours,

subsequently removing them (Fig. 1e). Growing hermaphrodites on male-conditioned plates

increased exopher production (Fig. 1f) to the same degree as when hermaphrodites were grown

with males until the L4 larvae stage (Fig. 1c). This elevation in exophers production also

coincided with a rise in the number of *in utero* embryos (Fig. 1g), indicating that *C. elegans*

male-secreted compounds, e.g., pheromones, can promote embryo retention in the

hermaphrodite's uterus and increase exophergenesis. Longer exposure to male-conditioned

plates or co-culture with males beyond larval development did not further increase exopher

production in *him-5* mutant hermaphrodites (Supplementary Fig. 1a-c). We next evaluated

whether the rise in exopher production was linked to continuous or stage-specific, acute

exposure to male secretions by growing hermaphrodites on male-conditioned plates during four

24-hour intervals (Fig. 1h). Results show that 24 hours of exposure to male-conditioned plates

is sufficient to increase exophergenesis in hermaphrodite, regardless of the life stage at which

the exposure occurred (Fig. 1h). This implies that exposure to male secretions solely during

worms' larval development can influence exopher level in adult hermaphrodites, with the most

potent effect observed in animals exposed during the L4 larval stage to young adults' day 1

stage, mirroring the effects of continuous exposure to males' secretions (Fig. 1h). In all

experiments, except for exposure during L4 larval stage to young adults' day 1stage, increased

exopher production was observed to coincide with higher embryo retention in the uterus (Fig.

1i). Our data indicate that exophergenesis is finely modulated by perception of male-derived

signals, possibly pheromones, acting through embryo-maternal signaling.

Elevated population density adversely affects exopher production.

In light of our previous findings demonstrating an increase in exopher production in the presence of males within the population, we sought to examine the effect of hermaphrodite population density on this phenomenon. To this end, we established two distinct experimental conditions. In the first condition, hermaphrodites were cultured individually on plates, eliminating potential social cues. In the second condition, worms were raised as groups of 10-hermaphrodites per plate from the beginning of their development (Fig. 2a). In contrast to the effects of the presence of males, our results reveal that hermaphrodites grown at 10 hermaphrodites per plate consistently released fewer exophers compared to those grown as solitary animals (Fig. 2b). Both experimental groups contained an equivalent number of embryos *in utero* (Fig. 2c), suggesting that this modulation of exophergenesis occurs independently of embryo-maternal signaling[6]. Furthermore, we observed that cultivating hermaphrodites in a population as small as five worms per plate was sufficient to decrease exopher production (Fig. 2d). Conversely, escalating the population size to one hundred animals per plate did not cause any further significant reduction in exopher production relative to 10 hermaphrodites per plate (Fig. 2e). Growing a single hermaphrodite on a plate conditioned by other hermaphrodites reduced exopher numbers to a level comparable to that observed at a density of 10 hermaphrodites per plate (Fig. 2f). Furthermore, raising adult hermaphrodites alongside developing larval hermaphrodites significantly decrease exopher production (Fig. 2g), highlighting the importance of population density, as even the presence of immature hermaphrodites can influence exophergenesis. In summary, these findings illuminate a complex interplay between social cues and exopher production in *C. elegans*, with the presence of hermaphrodites leading to a reduction in exopher release. The effect of hermaphrodites contrasts with the observed male-induced increase, highlighting a differential response to social cues from different sexes within the population.

The male pheromone ascr#10 stimulates hermaphrodite exopher production.

Based on our finding that male-conditioned plates are sufficient to increase muscle exopher production in hermaphrodites, we hypothesized that male-secreted pheromones may underlie the observed effects. Males secrete a wide range of sex-specific metabolites[14,15], including several members of the ascaroside family of pheromones, among which ascr#10 is the most abundant[16]. To investigate the hypothesis that muscle exopher production in hermaphrodites may be regulated by male-derived ascaroside pheromones, we took advantage of several worm mutants in which the biosynthesis of the ascaroside side chains via peroxisomal β-oxidation is perturbed[17]. As shown in Figure 3a *acox-1(ok2257)*, *maoc-1(ok2645)*, and *daf-22(ok693)* mutants each exhibit distinct ascaroside profiles, including dramatically different amounts of the male pheromone ascr#10[18]. Similar to the male co-culture experiments, we examined exophergenesis in wild-type animals co-cultured with ascaroside biosynthesis mutants (one wild-type worm with nine mutant worms per plate, Fig. 3b). We found that co-culture with *acox-1(ok2257)* animals boosted wild-type exopher levels, whereas *maoc-1(ok2645)* reduced levels, and *daf-22(ok693)* had no significant effect (Fig. 3c), with embryo counts *in utero* remaining unaffected by all three strains (Fig. 3d), suggesting that regulation of exopher production by hermaphrodite-released pheromones is largely independent of embryo-maternal signaling.

Observing an increased exopher count when wild-type hermaphrodites were exposed to ascarosides released by *acox-1(ok2257)* hermaphrodites, known for their elevated ascr#10 levels[18], and considering the dominance of ascr#10 in male pheromone profiles[16], we assessed the role of this specific ascaroside in exopher upregulation. Remarkably, we found that exposure to low concentrations of ascr#10 (~ 1 nanomolar) led hermaphrodites to produce more exophers than control animals, while the number of embryos in the uterus remained unchanged (Fig. 3e-f). Ascr#10 alone amplifies exophergenesis, yet the *acox-1* mutants also exhibit a

notable presence of ascr#18 alongside it[18]. Prompted by this, we examined ascr#18's role in

exopher modulation. Unexpectedly, exposure of animals to concentrations as low as approx. 1

picomolar of ascr#18 reduced exopher formation in reporter hermaphrodites (Fig. 3g), and at

greater concentrations (~ 1 nanomolar), it also reduced egg retention in utero (Fig. 3h). These

results show that different ascaroside pheromones have starkly different effects on exopher

production, emphasizing the intricate relationship between chemical structure and activity in

ascarosides[17].

Bacterial diet has a limited influence on exopher production

To rule out the possibility that exophergenesis is substantially regulated by the molecules

derived from the bacterial food source and to further explore the relationship between sensory

neurons, pheromones, and bacterial diet in *C. elegans*, we tested various bacterial strains that

might indirectly influence worm-to-worm communication. We directly compared the

Escherichia coli B strain OP50 and K-12 strain HT115, which are prevalently utilized in *C.*

elegans culture and RNAi silencing experiments, respectively[19,20]. Our findings revealed an

increase in the number of exophers when the *E. coli* HT115 diet was applied, as opposed to the

OP50 diet (Supplementary Fig. 2a). However, there was a notable elevation in the number of

eggs present in utero on the second day of adulthood under the HT115 diet (Supplementary

Fig. 2b). Importantly, the effect of the presence of males on the increase in exopher numbers

was observed, regardless of the bacterial strain used for feeding (Supplementary Fig. 2c). The

metabolic activity of the bacteria, whether alive or PFA-killed prior to culture, did not impact

the results (Supplementary Fig. 2a), corroborating that exophergenesis in the worm's muscles

is consistently activated, irrespective of the bacterial type used as a food source.

ASK, AWB, and ADL sensory neurons govern exophergenesis in response to males-social cues

Given the role of many olfactory neurons in detecting ascarosides[16] (Fig. 4a), we examined whether genetic ablation of all ciliated sensory neurons would abolish male pheromone-regulated exopher induction. Indeed, as shown in Fig. 4b, *che-13* mutants, which do not form proper cilia and are incapable of pheromone detection[21], produce a minimal number of exophers (Fig. 4b). To pinpoint the specific sensory neurons mediating pheromone-dependent modulation of exophergenesis, we examined its level in animals with genetically ablated sensory neurons previously identified as capable of pheromone detection[22]. We found that the elimination of ASK, ADL, or AWC suppressed exopher production comparable to that in *che-13* mutants, while ASH ablation led to a modest reduction (Fig. 4c). Notably, worms with genetically ablated neurons, which showed diminished exophergenesis, also displayed fewer eggs *in utero* (Fig. 4d). Considering the prominent effect on exophergenesis observed in genetic ablation of AWC thermo-responsive neurons[23,24], we investigated the influence of temperature on exophergenesis. When growing worms at 15, 20, or 25 °C, we noted a temperature-related rise in exopher formation (Fig. 4e). This increase was inhibited in the absence of AWC, substantiating the temperature-dependent control of exophergenesis by these neurons (Fig. 4e). Interestingly, the introduction of male-specific, pheromone-sensing CEM neurons in hermaphrodites (via *ceh-30* gain-of-function mutation[25]) did not lead to changes in exopher level (Fig. 4c).

Subsequently, we explored which olfactory neurons are essential for detecting male pheromones that enhance exophergenesis. By growing hermaphrodites with genetic ablations of different classes of olfactory neurons on *him-5* mutant male-conditioned plates (as shown in Fig. 1e scheme), we determined that the removal of ASK, AWB, or ADL neurons prevented the increase in the number of embryos *in utero* and nullified the enhancement in exopher

216   production driven by male-emitted pheromones (Fig. 4f; Supplementary Fig. 3a). These results

217   align with previously described roles for ASK, AWB, and ADL neurons in male pheromones

218   sensing[26–29]. This demonstrates the critical role of the neuronal odor recognition system in

219   modulating EV production by muscles and highlights the importance of particular neurons in

220   recognizing distinct pheromone types.

221   ASK, ASI, and ASH olfactory neurons regulate exopher production in hermaphrodite

222   populations

We sought to pinpoint the neurons crucial for downregulating exopher production in response

to the presence of other hermaphrodites. We grew hermaphrodites with genetically ablated

different classes of olfactory neurons either singly or in a 10 hermaphrodite population (Fig.

2a scheme). Our analysis with strains showing impaired olfaction revealed that ASH, ASI, and

ASK neurons are pivotal in this process (Fig. 4g). Remarkably, the masculinization of a

hermaphrodite olfactory circuit through the introduction of CEM male-specific neurons

resulted in the absence of exopher production decrease in ten hermaphrodite population

comparing to solitary worms (Fig. 4g). There was no correlation between exopher numbers and

eggs *in utero* count neither for wild-type animals nor for ablation mutants with (ASI) or without

(AWB) impaired hermaphrodite pheromones recognition (Supplementary Fig. 3b). This further

demonstrates that hermaphrodite pheromones regulate exophergenesis by omitting embryo-

maternal signaling. Finally, the complex interaction between ascaroside synthesis and neural

mechanisms in the regulation of exophergenesis was reinforced by the interplay observed

between ASI and AWB neuronal mutant hermaphrodites with ascaroside mutant

hermaphrodites (Supplementary Fig. 3c-d). In summary, the critical role of ASH, ASI, and

ASK neurons underlines the nuanced regulation of exophergenesis in response to the presence

of other hermaphrodites.

STR-173 receptor mediates pheromone-dependent exophergenesis modulation through ascr#10

Binding signaling molecules to the relevant receptor is the first step in transducing neuron chemosensory signals. More than 1,300 G protein-coupled receptors (GPCRs) mediate this communication in *C. elegans*[30]. Internal states and environmental conditions can modulate GPCRs expression to affect worm behavior[31–34]. To identify the receptor(s) responsible for ascaroside-mediated alterations in exopher formation, we performed RNA-sequencing of animals grown either as a single animal or in a 10-hermaphrodites population (Supplementary Fig. 4a). On the transcriptional level, neither group differed markedly (Supplementary Fig. 4b), and none of the 7TM receptors was significant up- or down-regulated (Supplementary Table 1). However, *str-173* receptor transcript, which shows among all of the G protein-coupled receptors one of the most evident trends between growth conditions (2.3 fold change), is, according to single-cell RNA-seq data[35], expressed almost exclusively in ASK neurons (Supplementary Fig. 4c). Since ASK is crucial for pheromone-mediated modulation of exopher formation, we investigated the *str-173* role in this pathway.

The wrmScarlet CRISPR/Cas9-mediated transcriptional knock-in for the *str-173* gene confirmed its strong expression during the late development and adulthood in ASK neurons and revealed additional expression in OLQ neurons, the pharynx, vulva muscles, and the tail (Fig. 5a-b). Next, we created *str-173* null mutants again using CRISPR/Cas9 editing (Supplementary Fig. 4d). We observed that *str-173* gene knockout neither influence the basal level of exophergenesis in *str-173* mutant hermaphrodites (Fig. 5c) nor robustly impact population density-mediated exophergenesis decrease (Fig. 5d). Interestingly, contrary to wild-type worms, *str-173* mutants did not exhibit an increase in exopher production in response to male pheromones (Fig. 5e). Recognizing the potential function of STR-173 in male pheromone-mediated signaling, we investigated its participation in the ascr#10 recognition.

Contrary to the increase in exopher production observed in wild-type animals, *str-173* mutants

exposed to ascr#10 display a decrease in exopher production, emphasizing STR-173's role in

translating male pheromone cues into physiological responses (Fig. 5f-g). Furthermore, *str-173*

mutants grown on NGM plates supplemented with ascr#18, like wild-type worms, show a

decrease in exopher production (Fig. 5h-i). This emphasizes the specificity of STR-173 binding

to ascr#10 rather than any other type of ascaroside.

The observed decrease in exopher production in *str-173* mutants exposed to ascr#10 suggests

the presence of another ascaroside receptor(s) capable of binding ascr#10 that plays an opposite

role in exopher regulation to STR-173. Exposing worms with ablated ASK neurons to ascr#10

or ascr#18 did not lead to significant changes in exopher counts (Fig. 5j), suggesting that

receptor(s) for ascr#18 and ascr#10, which decrease exopher production, are located in ASK

neurons. In summary, our data indicate a vital role of ASK in exopher regulation, mediated

through various ascaroside receptors expressed in ASK neurons.

AQR, PQR, and URX neurons activity limit exophergenesis

Among the 118 classes of neurons in *C. elegans*, only four are directly exposed to the

pseudocoelomic cavity[36]. Three classes of these neurons, AQR, PQR, and URX, regulate social

feeding in worms[37]. Given that exophers are released to the worm's pseudocoelomic cavity and

are regulated by social cues, we hypothesized that AQR, PQR, and URX might play a role in

exophergenesis. To test for that, we first investigated the effect of genetic ablation of AQR,

PQR, and URX neurons on exopher production. Indeed, the removal of these neurons leads to

a substantial increase in exophergenesis (Fig. 6a). Notably, the increased number of exophers

generated by worms with genetically ablated AQR, PQR, and URX neurons is not the result of

embryo-maternal signaling as these animals have a modestly reduced number of eggs *in utero*

(Fig. 6b) and smaller brood size (Supplementary Fig. 5a-b) than wild-type control.

To further validate the role of AQR, PQR, and URX neurons in the regulation of

exophergenesis, we optogenetically inactivated or activated them using ArchT[38] or ReaChR[39],

respectively, and compared the number of exophers before and after the stimulus. We observed

that 60 min of AQR, PQR, and URX neurons inactivation leads to a significant increase in

exopher release (Fig. 6c and Supplementary Fig. 5c). On the other hand, 60 min of AQR, PQR,

and URX neurons activation resulted in a significant decrease in exopher release after the

stimulus was completed (Fig. 6d and Supplementary Fig. 5d). This modulation was evident at

the adult day 1 (AD1) stage for ReaChr-based activation but not ArchT-based inactivation,

underscoring the potential significance of an optimal embryo count or direct uterus interaction

(Supplementary Fig. 5e-h).

Moreover, our data underscores that the contrasting exophergenesis characteristics observed in

ASK-ablated worms and in worms with genetic elimination of AQR, PQR, and URX neurons

resemble those of wild-type worms in animals lacking all four neuron classes (Fig. 6e). We

further determined that AQR, PQR, and URX neurons are involved in response to

hermaphrodite pheromones (Fig. 6f), but not to male pheromones (Fig. 6g). However, their

heightened activity could not override the indispensable role of fertility in exophergenesis (Fig.

6h-i). In summary, AQR, PQR, and URX neurons distinctly modulate exopher production in

response to hermaphrodite pheromones, underscoring the complexity of neuronally-regulated

reproductive signaling.

URX-expressed FLP-8 and AQR/PQR/URX-expressed FLP-21 neuropeptides inhibit

exophergenesis

To elucidate the mechanism by which AQR/PQR/URX neurons regulate muscle exopher

production, we used single-cell RNA-seq data[31], targeting neuropeptides predominantly

expressed in these neurons with a limited expression in other cells. These criteria were fulfilled

by URX-expressed FLP-8, and AQR, PQR, URX-expressed FLP-21 neuropeptides

(Supplementary Fig. 6 a-c). We could demonstrate that these neuropeptides negatively regulate

exophers production, as evidenced by increased exophers counts in both single and double

mutants (Fig. 7a), independent of embryo-maternal signaling (Fig. 7b). Moreover, FLP-8 and

FLP-21 act downstream of URX and AQR/PQR/URX, respectively, as optogenetical activation

of these neurons in the mutant context failed to suppress exopher release as observed in control

animals (Fig. 7c-e, Supplementary Fig. 7a-c). The results acquired regarding the influence of

the FLP-21 neuropeptide on the regulation of socially-driven exophergenesis align with the

previously established role of FLP-21 in controlling social behavior[40].

Model of exopher regulation mechanism in *C. elegans*

Our investigations have led us to construct the following model detailing the intricate

mechanism of exopher regulation:

1. Male pheromones: Males produce ascarosides, including ascr#10, that increase exopher

production levels. These signals predominantly act through ASK, ADL, and AWB. The

critical role in this pathway is played by ASK-expressed STR-173 G protein-coupled

receptor, which binds ascr#10 to potentiate exopher production.

2. Hermaphrodite pheromones: Hermaphrodites release ascarosides, including ascr#18,

that reduce exopher production levels. These signals predominantly act through ASI,

ASH, and ASK (via ascr#18) sensory neurons.

3. Embryo accumulation effect: The rise in embryos inside the hermaphrodite triggers a

series of pro-exopher signals. Importantly, increased *in utero* embryo accumulation can

be mediated by a blend of male-released pheromones.

4. Neuropeptide control: AQR/PQR/URX pseudocoelomic cavity-opened neurons release

FLP-8 and FLP-21 neuropeptides that negatively regulate exophergenesis. This type of

modulation is important for the decrease in exophergenesis dependent on

hermaphrodite pheromones but not for the increase in reliance on male pheromones.

5. Integrated response: Exopher production is modulated by a blend of internal processes
and external cues, whether from male or hermaphrodite secretions. This balance ensures
that exophergenesis is regulated in tandem with various internal and external
environmental conditions.

In encapsulating our findings, the exopher regulation in *C. elegans* unfolds as a sophisticated

interplay of signals, neuronal circuits, and their associated molecules. These components

collectively respond and adapt to male-derived cues, and hermaphrodite-derived signals,

creating a balanced system for exopher production and release (see Fig. 8 for a detailed

overview).

Discussion

Our study shows that regulation of exophergenesis in *C. elegans* presents a sophisticated

network of neuroendocrine and pheromone signaling pathways that integrate internal and

external environmental cues. These results shed light on the vital roles of male- and

hermaphrodite-derived ascarosides in influencing this complex network. Central to our

findings is the role of male-derived ascaroside pheromones, particularly ascr#10. Detected by

the STR-173 receptor in ASK neurons, this pheromone plays a pivotal role in potentiating

exopher production. On the other end of the spectrum, hermaphrodites produce pheromones,

including ascr#18, which act to temper exopher levels. Our investigations revealed that the

ASH, ASI, and ASK olfactory neurons mediate this response. This interplay between

ascaroside synthesis, chemosensation, and neuronal regulation hints at the sophisticated nature

of *C. elegans*' inter-animal communication. Our research further identified the AQR, PQR, and

URX neurons as pivotal elements in the matrix of exopher regulation. The profound influence

of FLP-8 and FLP-21 neuropeptides, originating from AQR, PQR, and URX neurons, further

consolidated our understanding. The modulation of these peptides, particularly with their

associated neurons, showcases the intricate balance of environmental and social signals with

the activity of endocrine systems within the nematode.

The notable increase in exopher production among hermaphrodites either exposed to male

metabolites or grown with males underscores the significant influence of male pheromones on

cellular processes. This observation aligns well with prior findings suggesting male

pheromones channel resource allocation towards the germline[41]. In this context, it is

conceivable that exophers are a part of this mechanism, possibly playing a pivotal role in

bolstering oocyte and embryo quality. This strategy may enhance reproductive success,

ensuring higher quality offspring even at the potential expense of the individual's somatic

health. However, this reproductive advantage seems to come at a cost. The elevation in

exophergenesis has been linked to a decreased exploratory behavior, which could be associated

with swifter deterioration of muscle function with age[6]. Aprison and Ruvinsky's observations

that hermaphrodites expedite both development and somatic aging in the presence of males[39]

further solidify this notion. This underscores the importance of understanding the role and

regulation of exophers, as they could serve as a nexus between environmental cues (like male

pheromones), reproductive strategies, and age-related degeneration. As such, exophers

probably act as biological executors and carriers of information in inter-animal communication.

This is also consistent with observations that *Drosophila* males secrete EVs that are important

for mating behavior[42], and the exchange of information between male and female flies leads to

increased EVs release from sperm secretory cells, which promotes fertility[43]. Drawing parallels

with mammals, while the signaling molecules (like ascarosides) might differ, the broader

concept of resource allocation and trade-offs between reproduction and somatic well-being

might find relevance, offering avenues for future exploration.

While the proximity of males induces an increase in exophergenesis, a heightened density of

hermaphrodites elicits a contrary response by diminishing it. There are instances where

organisms, in response to densely populated environments, adaptively modulate their

metabolic and reproductive functions to address potential resource scarcity[10,44]. Within the

context of *C. elegans*, an escalation in hermaphrodite density could indicate imminent resource

scarcity. Consequently, this instigates the nematodes to modify survival tactics, encompassing

the reduction of exopher synthesis. Given the complexity of cues—a supportive push from

embryo-maternal signals and a dampening nudge from dense hermaphrodite surroundings—

the resultant exopher production could be a net outcome of these counteracting signals. Future

investigations should unravel these cues' hierarchical dynamics and identify prevailing signals

under diverse conditions. For a comprehensive understanding of this facet, subsequent research

should emphasize alterations in neural activity amidst different social scenarios and discerning

the distinct signaling responses triggered by both male and hermaphrodite stimuli.

Investigating how *C. elegans* balances the demand for supporting offspring development (via

exophers) with the need to adapt to social stresses in crowded conditions can offer deeper

evolutionary insights.

Our study has illuminated the multifaceted nature of the regulatory network involving

ascarosides, their receptors, and olfactory neurons in the environmental adaptations of *C.*

elegans. Specifically, we discovered that ascr#10, which is abundantly excreted upon sexual

maturation by males, amplifies exopher production through the STR-173 receptor, whereas

ascr#18, which is excreted constitutively at much lower rates by hermaphrodites, diminishes

it. While the receptor for ascr#18 remains unidentified, our results indicate it must also be

located within ASK neurons. This intricate interplay between chemoreceptors and pheromones

suggests that exophergenesis is regulated by a complex network influenced by social and

environmental cues. The impact of this network is further highlighted by our observation that

the removal of an olfactory neuron expressing various chemoreceptors or alterations in the

ascaroside biosynthesis pathway can either positively or negatively regulate exophergenesis.

The complexity of the entire regulatory network is escalated by the fact that the lack of activity

of one receptor, achieved either by transcriptional downregulation or post-translational

modification, can drastically alter the biological potential of its ligands. This was demonstrated

in our analysis, where knockdown of *str-173* reversed the pro-exopher potential of ascr#10,

implying that even subtle alterations in receptor activity can have profound consequences on

the physiological responses mediated by the associated ligands. Identifying the anti-exopher

factors will be pivotal in understanding the full spectrum of regulatory effects mediated by

ascarosides and their receptors. Additionally, our study exemplifies the multilayered interplay

between pheromones and cellular and organismal responses. The notable accumulation of

embryos in the hermaphrodite's uterus upon exposure to the male secretome, contrasting with

the lack of egg retention induced by ascr#10, underlines the intricate connection between

external pheromonal cues and internal physiological responses, suggesting a strategic

evolutionary adaptation for resource allocation in response to reproductive efforts. These

findings also highlight that ascr#10 is likely one of several male-excreted small molecule

signals that affect exopher formation and egg laying in hermaphrodites. Moreover, the

interaction of ascr#10- and ascr#18-mediated signaling with each other and additional male-

derived signals can reveal additional pathways and responses. In conclusion, our findings

underscore the need for extensive research into the complex interplay between metabolites

(pheromones/ascarosides), receptors, and intracellular processes and their modulation by social

and environmental cues. Understanding this network will deepen our knowledge of the

mechanisms governing exophergenesis and provide valuable insights into other cellular

processes and environmental adaptations mediated by similar mechanisms in *C. elegans* and

potentially other organisms.

Our research demonstrates the pivotal role of the STR-173 receptor, specifically in ASK

neurons, in orchestrating pheromone-driven exophergenesis. This discovery contributes to the

limited knowledge surrounding 7TM GPCRs as ascaroside/pheromone receptors, considering

the vast number of 7TM GPCRs expressed[45] and metabolites secreted by *C. elegans*[46]. The

association between STR-173 receptors and male ascr#10 suggests a receptor-ligand dynamic,

potentially triggering a series of intracellular processes that modulate exopher production. A

comprehensive exploration of this cascade, spanning receptor engagement to downstream

effects, remains elusive. In the intricate environment of *C. elegans*, our investigation has

provided illuminating insights into the multifaceted roles played by specific sensory neurons

AQR, PQR, and URX and their associated neuropeptides, FLP-8 and FLP-21, in modulating

muscle exopher production. Building on prior studies, our identification of FLP-8 and FLP-21

as modulators of extracellular vesicle release from body wall muscles brings further depth to

our understanding of neuropeptide functionality in *C. elegans*.

Neuropeptides, exemplified by FLP-18, demonstrate intricate roles in regulating muscle

function and behavior in *C. elegans*. Specifically, the co-release of FLP-18 with tyramine from

the RIM neurons and its targeting of body wall muscles via the NPR-5 receptor to modulate

muscle excitability underscores the multifaceted interactions of neuropeptides and

neurotransmitters in these processes[47]. Given this, it is plausible to assume that FLP-8 and FLP-

21 may function along similar pathways. Their involvement might influence muscle

excitability in a manner that subsequently impacts the release dynamics of extracellular

vesicles. Exploring the precise receptors and pathways through which FLP-8 and FLP-21 act

will be instrumental in further elucidating their specific roles in exophers release and muscle

functionality.

In *C. elegans*, specific neuropeptides like FLP-8 and FLP-21, along with distinct neuronal

classes such as AQR, PQR, and URX, appear to play a pivotal role in the modulation of exopher

production. Given that exophers encapsulate damaged or aggregated proteins, understanding

the regulatory mechanisms in this nematode model might offer a novel perspective on protein

aggregation and its clearance. While these specific neuropeptides and neuronal populations are

unique to *C. elegans*, the underlying principles might hold broader biological significance. For

example, if analogous mechanisms exist in more complex organisms, they could potentially

impact our understanding of diseases like Alzheimer's or Parkinson's, where protein

aggregation is a prominent concern. Thus, while direct translational insights may be limited,

C. elegans offers a window into fundamental processes that could shed light on the

sophisticated dance between neurons, protein aggregation, and potential clearance mechanisms

in neurodegenerative conditions.

In conclusion, our research is an initial foray into unraveling the complex regulatory

mechanisms of exopher production in *C. elegans*, suggesting the possibility of universally

conserved physiological systems. The findings hint at a future filled with deeper exploration

into these dynamics, possibly redefining our understanding of intercellular communication

across species. Similar mechanisms modulating cell-to-cell communications via exophers

might exist in humans, as evidenced by studies on other extracellular vesicle classes[43]. Our

study also underscores the significance of the worm's olfactory system in steering exopher

generation within muscles. It is worth noting that human olfactory impairments are linked with

a heightened risk for cardiovascular disease (CVD)[44,45]. Such a correlation might be attributed

to disruptions in exopher-driven homeostatic processes within the heart. Still, it remains

imperative to ascertain if a parallel olfactory-driven exopher regulatory mechanism exists in

mammals.

Figures and Figure legends

Fig. 1

Figure 1. Production of *Caenorhabditis elegans* muscle exophers is regulated by male pheromones.

a   An example of *C. elegans* muscle exopher containing mitochondria. Red fluorescence comes from the RPN-5 proteasome subunit tagged with wrmScarlet fluorescent protein. Green fluorescence shows mitochondrial outer membrane (MOM) tagged with GFP. Arrows point to examples of exophers, asterisks indicate mitochondria-containing exophers.

b   Schematic representation of the experimental setup for investigating the influence of increased male presence in the population. Him-5 mutant hermaphrodites and him-5 mutant males were co-cultured from the L1 stage until the L4 stage and then transferred to a male-free plate. Hermaphrodites were grown on a male-free plate until adulthood day 2 (AD2) when the number of exophers was assessed.

c   Increased percentage of males in the population (via *him-5(e1490)* mutation) leads to a higher exophergenesis level in hermaphrodites. n = 79 and 83; N = 3

d   Increased percentage of males in the population (via *him-5(e1490)* mutation) causes embryo retention in hermaphrodites. n = 79 and 85; N = 3

e   Schematic representation of the experimental setup for investigating the influence of male's secretome on exophergenesis level. 50 *him-5* or wild-type males were grown on a plate for 48 hours before being removed. Next, 10 hermaphrodites were transferred to plates previously occupied by males and grown until adulthood day 2 (AD2), when the number of exophers was assessed.

f   Growing wild-type hermaphrodites on *him-5(e1490)* or wild-type male-conditioned plates increases exophergenesis levels. n = 89 - 106; N = 3.

g   Exposing wild-type hermaphrodites to *him-5(e1490)* or wild-type male secretome causes embryo retention in the uterus. n = 60 - 105; N = 3.

h - i   Exposure to male pheromones for just 24 hours triggers exophergenesis in

hermaphrodites, regardless of the life stage at which the exposure occurs. Higher embryo

retention in the uterus coincided with an increase in exopher production. (h) n = 85 - 90; N =

3, (i) n = 48 – 50; N = 2.

Data information: Scale bars are 20 μm and 5 μm (zoom in). @AD2 means "at adulthood day

2". Data are presented as median with interquartile range; n represents the number of worms;

N represents a number of experimental repeats that were combined into a single value; ns - not

significant (P > 0.05), * P < 0.05, ** P < 0.01 P < 0.0001; (c, d) Mann-Whitney test, (f - i)

Kruskal-Wallis test with Dunn's multiple comparisons test.

Fig. 2

Figure 2. A rise in the population density of hermaphrodites adversely affects exophergenesis.

a Experimental design for determining the effect of the presence of other hermaphrodites on the number of exophers.

b *C. elegans* hermaphrodites grown in the presence of other hermaphrodites produce fewer exophers than solitary animals. n = 88 and 90; N = 3.

c Embryo-maternal signaling is not responsible for the differences in exophergenesis levels as solitary and accompanied hermaphrodites contain the same number of embryos *in utero*. n = 88 and 90; N = 3.

d – e Cultivating as few as five hermaphrodites per plate reduced exopher production, with no significant additional reduction observed when increasing the population to one hundred per plate. (d) n = 70 – 76, (e) n= 100 - 124; N = 3

f A single hermaphrodite on a plate previously occupied by other hermaphrodites yielded exopher numbers comparable to worms in the ten-hermaphrodite population. n = 45 and 60; N = 3 - 4.

g Growing single adult hermaphrodite with developing larval hermaphrodites leads to a significant reduction in exopher production. n = 45 and 61; N = 3 - 4.

Data information: Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant (P > 0.05), * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001; (b, c) Mann-Whitney test, (d - g) Kruskal-Wallis test with Dunn's multiple comparisons test.

a

WT
ascr #10:ascr#18
6:1

ascr #10

ascr #18

acox-1(ok2257)
ascr #10:ascr#18
4:1.9

ascr #18

ascr #10

maoc-1(ok2645)
ascr #10:ascr#18
4:11

ascr #10

ascr #18

daf-22(ok693)
no ascr #10 & #18

- saturated
- α, β-unsaturated
- β-hydroxylated

b

9 asc biosynthesis mutants/N2 worms
1 exopher reporter

exophers scored @AD2

c

No. of exophers/animal

ns
***
**

1♂ WT + 9⚥ N2
1♂ WT + 9⚥ acox-1(ok2257)
1♂ WT + 9⚥ maoc-1(ok2645)
1♂ WT + 9⚥ daf-22(ok693)

d

No. of eggs/animal

ns
ns
ns

WT ♂ + 9⚥ N2
1♂ WT + 9⚥ acox-1(ok2257)
1♂ WT + 9⚥ maoc-1(ok2645)
1♂ WT + 9⚥ daf-22(ok693)

e

No. of exophers/animal

ns
****

WT ⚥
WT ⚥ + ascr#10 1pM
WT ⚥ + ascr#10 1nM

f

No. of eggs/animal

ns
ns

WT ⚥
WT ⚥ + ascr#10 1pM
WT ⚥ + ascr#10 1nM

g

No. of exophers/animal

*
*

WT ⚥
WT ⚥ + ascr#18 1pM
WT ⚥ + ascr#18 1nM

h

No. of eggs/animal

ns
*

WT ⚥
WT ⚥ + ascr#18 1pM
WT ⚥ + ascr#18 1nM

Figure 3. The production of exopher in hermaphrodite is stimulated by ascr#10 and inhibited by ascr#18 pheromones.

a   Mutants with altered very long chain ascarosides biosynthesis exhibit unique ascaroside pheromone profiles, characterized by varying levels of ascr#10 and its ratio to ascr#18. Based on the data from [18]. Circle diameter denotes the proportion of produced ascarosides relative to wild-type.

b   Schematic representation of the experimental setup for investigating the influence of ascaroside side-chain biosynthesis mutants presence on exophergenesis level in wild-type worms.

c   Growing wild-type hermaphrodites in the presence of *acox-1(ok2257)* and *maoc-1(ok2645)* mutant increase and decrease exopher production, respectively. n = 38 - 93; N = 3.

d   Growing wild-type hermaphrodites with ascaroside side-chain biosynthesis mutants does not influence the number of embryos in the uterus. n = 37 - 93; N = 3.

e   Hermaphrodites grown on NGM plates supplemented with 1 ng of ascr#10 have increased levels of exophergenesis. n = 164 - 190; N = 5 - 6.

f   ascr#10 do not influence embryo retention in hermaphrodites. n = 69 - 101; N = 4.

g   Hermaphrodites grown on NGM plates supplemented with 1 pg or 1 ng of ascr#18 have decreased levels of exophergenesis. n = 83 - 90; N = 3.

h   ascr#18 do not influence embryo retention in hermaphrodites. n = 63 - 71; N = 3.

Data information: @AD2 means "at adulthood day 2". Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant ($P > 0.05$), * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$; Kruskal-Wallis test with Dunn's multiple comparisons test.

Fig. 4

Figure 4. Multiple olfactory neurons regulate exophergenesis levels in response to hermaphrodite and male pheromones.

a   Sensory neurons investigated within this study.

b   Hermaphrodites with impaired ciliated sensory neurons have lower exophergenesis levels. n = 66 and 65; N = 3.

c   Genetic ablation of ASK, AWC, ADL, or ASH neurons reduces exopher production. n = 61 - 97; N = 3.

d   Worms with genetically ablated neurons, resulting in reduced exophergenesis, also exhibited fewer eggs *in utero*. n = 42 - 61; N = 2 - 3.

e   AWC neurons regulate the temperature-dependent increase in muscle exopher production. n = 52 - 92; N = 3.

f   Increase in exophergenesis levels due to exposure to male pheromones is mediated by ASK, AWB, and ADL neurons. Schematic representation of the experimental setup is presented in Fig. 1e (without conditioning with WT males). n = 77 - 108; N = 3.

g   Decreased exophergenesis level due to exposure to hermaphrodite's pheromones is mediated by ASH, ASI, and ASK neurons and can be altered by adding CEM male-specific neurons. Schematic representation of the experimental setup is presented in Fig. 2a. n = 49 - 97; N = 3.

Data information: Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant ($P > 0.05$), * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$; (b, f, g) Mann-Whitney test, (c - e) Kruskal-Wallis test with Dunn's multiple comparisons test.

Figure 5. The STR-173 receptor facilitates the pheromone-dependent modulation of exophergenesis via ascr#10.

a - b *str-173* 7TM receptor is expressed in neurons (ASK and probably[#] OLQ) and non-neuronal tissues (pharynx marked with an arrow, vulva, and probably[##] rectal gland marked with circles). Square brackets mark gut autofluorescence. [#]based on the position and scRNAseq data[35] [##]based on the position and the shape.

c Exophergenesis level in *str-173* mutants co-cultured with other hermaphrodites is unchanged compared to wild-type worms. n = 90; N = 6.

d Knocking out the *str-173* gene does not roboustly affect the decrease in exophergenesis mediated by changes in population density. n = 86 - 90; N = 3.

e Increase in exophergenesis levels due to exposure to male pheromones depends on STR-173 7TM receptor. n = 60 - 90; N = 2 - 3.

f - g When exposed to ascr#10, *str-173* mutants exhibit a reduction in exopher production and *in utero* embryo number, underscoring the importance of STR-173 in binding ascr#10. (f) n = 104 – 113, N = 4; (g) n = 60, N = 2

h - i *Str-173* mutants, when cultured on NGM plates supplemented with ascr#18, experience a decrease in exopher production. (h) n = 124 - 134, N = 3; (i) n = 96 - 115, N = 3

j Removal of ASK neurons nullifies ascr#10 and ascr#18 influence on exophergenesis. n = 99 - 135; N = 3.

Data information: Scale bars are (a) 30 μm and (b) 20 μm. Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant ($P > 0.05$), * $P<0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$; (c, d - j) Kruskal-Wallis test with Dunn's multiple comparisons test, (d, e) Mann-Whitney test.

Fig. 6

Figure 6. Pseudocoelom-exposed neurons negatively regulate exopher production.

a   Genetic ablation of pseudocoelom-exposed AQR, PQR, and URX neurons increases exopher production. n = 89 and 100; N = 3.

b   Genetic ablation of pseudocoelom-exposed AQR, PQR, and URX neurons causes a decrease in the number of embryos present in the uterus. n = 89 and 100; N = 3.

c   ArchT-mediated optogenetic inactivation of AQR, PQR, and URX neurons increases exopher production. n = 59; N = 6.

d   ReaChR-mediated optogenetic activation of AQR, PQR, and URX neurons decreases exopher production. n = 59 and 60; N = 6.

e   The opposing exophergenesis levels observed in animals with genetic ablation of ASK neurons (low exophergenesis) and AQR, PQR, and URX neurons (high exophergenesis) converge to an intermediate level in animals with all four neurons removed. n = 89 - 95; N =3.

f   Decreased exophergenesis level due to exposure to hermaphrodite's pheromones is partially mediated by AQR, PQR, and/or URX neurons. n = 51 - 60; N = 3.

g   Increase in exophergenesis levels due to exposure to male pheromones is not altered in animals with genetic ablation of AQR, PQR, and URX neurons. n = 102 - 105; N = 3.

h   Genetic ablation of AQR, PQR, and URX neurons does not rescue the inhibition of exophergenesis caused by FUdR-mediated worm sterility. n = 89 - 90; N = 3.

i   Genetic ablation of AQR, PQR, and URX neurons only partially rescues the inhibition of exophergenesis caused by *rme-2* (yolk receptor) knockdown. n = 90; N = 3.

Data information: +ATR means "with all-trans-retinal". Data are presented as median with interquartile range (a – b, e – i) or mean with SEM (c – d); n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant (P > 0.05), * P < 0.05, ** P < 0.01,*** P < 0.001, **** P < 0.0001; (a - d, f - h) Mann-Whitney test, (e, i) Kruskal-Wallis test with Dunn's multiple comparisons test.

Fig. 7

Figure 7. Neuropeptides FLP-8 released by URX neurons and FLP-21 released by AQR,

PQR, and/or URX inhibit exophergenesis.

a *flp-8(pk360)* and *flp-21(ok889)* single and double mutants have increased

exophergenesis levels. n=180, N=6 for WT; n=90, N=3 for mutants.

b　　　Mutations in *flp-8(pk360)* and *flp-21(ok889)* genes do not influence embryo retention

in utero. n=140, N=5 for WT; n=60-80, N=3 for mutants.

c-e　　Optogenetic activation of AQR, PQR, and URX which do not produce FLP-8 and FLP-

21 neuropeptides do not lead to a decrease in exopher production. n = 65 - 71; N = 6.

Data information: +ATR means "with all-trans-retinal". Data are presented as median with

interquartile range (a – b) or mean with SEM (c – e); n represents the number of worms; N

represents a number of experimental repeats that were combined into a single value; ns - not

significant (P > 0.05), * P < 0.05, **** P < 0.0001; (c - e) Mann-Whitney test, (a, b) Kruskal-

Wallis test with Dunn's multiple comparisons test.

Figure 8. Model

Males produce ascarosides, including ascr#10, which increase exopher production through ASK, ADL, and AWB signaling. The critical player in this pathway is the ASK-expressed STR-173 G protein-coupled receptor, which enhances exopher production in response to ascr#10. In contrast, hermaphrodites release ascarosides like ascr#18, which reduce exopher production, primarily acting via ASI, ASH, and ASK (via ascr#18) sensory neurons. AQR/PQR/URX neurons in the pseudocoelomic cavity release FLP-8 and FLP-21 neuropeptides, negatively regulating exophergenesis. This modulation is crucial for decreasing exophergenesis due to hermaphrodite pheromones but not for the increase driven by male pheromones.

Supplementary Figure 1. Exposing hermaphrodites to males or their scent throughout the entire duration of the experiment do not further elevate exopher production.

a   Schematic representation of the experimental setup for Supplementary Fig. 1b.

b   Exposing hermaphrodites to a male's secretome after the L4 stage does not further increase exopher production. n = 83; N = 3.

c   Co-culturing him-5 hermaphrodites with males increases exophergenesis in hermaphrodites. n = 72 and 90; N = 3.

Data information: Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant (P > 0.05), **** P < 0.0001; Mann-Whitney test.

Supplementary Figure 2. Influence of bacteria diet on muscle exophers formation.

a   Worms fed with metabolically active and inactive bacteria produce a similar number of exophers. Worms grown on *E. coli* HT115 strain produce more exophers than worms grown on *E. coli* OP50 strain. n = 78 - 93; N = 3.

b   Worms grown on *E. coli* HT115 strain contain more eggs *in utero* than worms grown on *E. coli* OP50 strain. n = 46 and 55; N = 3.

c   Growing hermaphrodites on male-conditioned plates increases exophergenesis levels regardless of *E. coli* strain used as a food source. n = 79 and 87; N = 3.

Data information: Data are presented as median with interquartile range; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value; ns - not significant (P > 0.05), * P < 0.05, **** P < 0.0001; (a) Kruskal-Wallis test with Dunn's multiple comparisons test, (b, c) Mann-Whitney test.

Supplementary Figure 3. Influence of pheromones released by ascaroside side-chain biosynthesis mutants on hermaphrodites with genetically ablated sensory neurons.

a   Removal of ASK, AWB, or ADL neurons prevented the increase in the number of embryos in utero after exposing hermaphrodites to male pheromones. n = 55 - 60; N = 3.

b   *In utero* embryo number for wild-type worms and ASI and AWB ablation mutants grown as solitary animals or in ten-hermaphrodites population. n = 55 - 60; N = 3.

c - d   Worms with ablated ASI and AWB neurons grown together with ascaroside side-chain biosynthesis mutants do not exhibit alterations in produced exopher numbers. n = 83 - 120; N = 3.

714   Data information: Data are presented as median with interquartile range; n represents the

715   number of worms; N represents a number of experimental repeats that were combined into a

716   single value; ns - not significant ($P > 0.05$), * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; (a, b) Mann-

717   Whitney test, (c, d) Kruskal-Wallis test with Dunn's multiple comparisons test.

a

b

c

d

...GATTTA————TGCGGT... (wwa1)
...GATTTATTTCTGCGGT... (WT)
...GATTT————GTGCGGT... (wwa2)

Supplementary Figure 4. Worms grown in populations with different densities did not

differ significantly on the transcriptional level.

a   Number of exophers produced by worms that were used for RNAseq analysis. n = 29 -

30; N = 3.

b   Worms grown as a single animal or in a ten-hermaphrodites population did not differ

significantly on the transcriptional level. A transcript with significant change is marked as a

red dot.

c Single-cell RNA-seq data from CeNGENApp[35] show *str-173* strong expression in ASK

neurons and weak expression in OLQ neurons. The circle diameter represents the proportion

of neurons in each cluster that express the *str-173* gene.

d Localization of *str-173* mutations in the gene.

Data information: Data are presented as median with interquartile range; n represents the

number of worms; N represents a number of experimental repeats that were combined into a

single value; * P < 0.05, ** P < 0.01; (**a**) Mann-Whitney test.

Supplementary Figure 5. ReaChr-based activation but not ArchT-based inactivation at

the adult day 1 (AD1) stage influence exopher production in hermaphrodites.

a - **b**   Worms with the genetic ablation of pseudocoelom-exposed AQR, PQR, and URX

neurons have lower brood size than the wild type. n = 34; N = 3.

c - **d**   Control experiment without all-trans retinal (ATR) for ArchT-mediated inactivation

and ReaChR-mediated activation of AQR, PQR, and URX neurons. n = 30; N = 3.

e – **f**   ArchT-mediated inactivation of AQR, PQR, and URX neurons at the adult day 1

(AD1) stage do not modulate exopher production. n = 59 - 72; N = 5 - 6.

g – **h**   ReaChR-mediated activation of AQR, PQR, and URX neurons at the adult day 1 (AD1)

stage significantly decreases exopher production. n = 59 - 62; N = 5 - 6.

Data information: +ATR means "with all-trans-retinal"; no ATR means "without all-trans-

retinal". Data are presented as median with interquartile range (a), mean (b), or mean with SEM

(c – h); n represents the number of worms; N represents a number of experimental repeats that

were combined into a single value; ns - not significant * P < 0.05, **** P < 0.0001; Mann-

Whitney test.

Supplementary Fig. 6

754   **Supplementary Figure 6. FLP-8 and FLP-21 are expressed in URX and AQR, PQR, URX,**

755   **respectively.**

756   **a** - **c**   *flp-8* and *flp-21* are strongly and relatively specifically expressed in AQR, PQR and/or

757   URX neurons. Data taken from[35].

Supplementary Figure 7. Control experiments without all-trans retinal (ATR) ReaChR-mediated activation of AQR, PQR, and URX neurons that do not produce FLP-8 and/or FLP-21 neuropeptides.

a   Control experiment without all-trans retinal (ATR) ReaChR-mediated activation of AQR, PQR, and URX neurons in *flp-8(pk360)* mutant background. n = 60; N = 5.

b   Control experiment without all-trans retinal (ATR) ReaChR-mediated activation of AQR, PQR, and URX neurons in *flp-21(ok889)* mutant background. n = 57 and 58; N = 5.

c   Control experiment without all-trans retinal (ATR) ReaChR-mediated activation of AQR, PQR, and URX neurons in *flp-8(pk360), flp-21(ok889)* double mutant background. n = 59 and 60; N = 5.

Data information: +ATR means "with all-trans-retinal"; no ATR means "without all-trans-retinal". Data are presented as mean with SEM; n represents the number of worms; N represents a number of experimental repeats that were combined into a single value.

Supplementary Table 1.

Gene.name	log2FoldChange	pvalue	padj
srw-86	2.470005259	0.006089231	NA
str-173	**1.210254202**	**0.083872612**	**NA**
sri-39	0.961870706	0.118312493	NA
srm-1	0.74631012	0.197705915	NA
srh-269	0.715357507	0.20128121	NA
sri-36	0.697436016	0.069365061	NA
str-55	0.69275127	0.286227955	NA
srd-64	0.671778641	0.227938718	NA
srsx-36	0.632246484	0.436691291	NA
sri-57	0.609273781	0.193809518	NA
srw-89	0.578848267	0.407242411	NA
srbc-82	0.56597091	0.423856002	NA
srr-2	0.523068614	0.103943944	NA
srx-97	0.505797282	0.424229255	NA
sri-11	0.489747472	0.297473585	NA
srv-15	0.463355568	0.372243301	NA
srj-14	0.460503239	0.065003699	NA
sre-36	0.458443613	0.346744567	NA
srh-217	0.429350684	0.444951979	NA
srh-2	0.409544299	0.241263936	NA
srx-41	0.39088958	0.531069383	NA
srr-10	0.382508507	0.600239609	NA
srz-97	0.370702424	0.498470233	NA

srsx-19	0.353102397	0.54270153	NA
srd-70	0.320458039	0.541451057	NA
srh-71	0.308999771	0.434944825	NA
str-179	0.291901118	0.596460956	NA
str-148	0.267146531	0.651806161	NA
sra-35	0.258242663	0.576292957	NA
srx-45	0.226491874	0.636877507	NA
srsx-27	0.225810039	0.724940673	NA
srr-6	0.223992203	0.491120033	NA
srb-16	0.213930895	0.686266407	NA
sri-55	0.204504014	0.648957813	NA
srx-125	0.196087663	0.632165424	NA
srw-48	0.1760458	0.784611201	NA
sri-35	0.174186697	0.816308334	NA
srsx-25	0.172603337	0.699023544	NA
srj-29	0.145173827	0.82575966	NA
str-176	0.145080793	0.381737839	0.705706926
srv-32	0.140744356	0.836820773	NA
srj-52	0.12758145	0.8360209	NA
srab-4	0.123433522	0.75301419	NA
srxa-14	0.122166676	0.825629763	NA
src-1	0.121226527	0.017785109	0.153059765
srr-4	0.118248088	0.143496801	0.45832863
srsx-34	0.118146738	0.517657949	NA
str-144	0.115566741	0.882522042	NA
sri-8	0.114856655	0.747072742	NA
srh-48	0.107505544	0.732564142	NA

str-245	0.085385814	0.731316789	NA
srv-7	0.081868619	0.676931931	NA
srj-9	0.057842686	0.909552581	NA
sra-33	0.04490344	0.924342693	NA
srsx-18	0.037063376	0.9551973	NA
src-2	0.006822308	0.971252278	NA
srr-1	0.00459541	0.99125387	NA
sri-59	0.000405882	0.999217048	NA
srg-34	-0.030080578	0.947775572	NA
str-172	-0.085472421	0.87286706	NA
srr-3	-0.088516378	0.76868265	NA
srd-32	-0.107551827	0.816267341	NA
srv-1	-0.11334895	0.410156047	0.727497177
srh-237	-0.151085831	0.175026351	0.503664735
str-183	-0.178110179	0.787945578	NA
sra-10	-0.178642194	0.779315606	NA
sra-32	-0.190091468	0.685526539	NA
srd-14	-0.207486411	0.305360875	NA
sre-13	-0.210537797	0.702655411	NA
srd-52	-0.235477439	0.677464298	NA
sri-40	-0.247157736	0.037111528	0.232197335
srz-85	-0.247831631	0.640803343	NA
sra-14	-0.266593158	0.200655026	NA
srh-16	-0.275054658	0.57204149	NA
srh-70	-0.322126344	0.575669232	NA
srab-14	-0.328373116	0.605637275	NA
srx-12	-0.361291214	0.429781376	NA

sre-9	-0.397855081	0.275566108	NA
srx-58	-0.458521562	0.487282023	NA
sri-5	-0.458897325	0.308597251	NA
srt-23	-0.460153207	0.606235802	NA
srm-3	-0.470085367	0.174616055	NA
srh-33	-0.479357546	0.48607886	NA
srh-105	-0.486191119	0.439431381	NA
srw-35	-0.492290252	0.480189518	NA
srx-118	-0.521178132	0.380103143	NA
srm-5	-0.557246057	0.448970879	NA
srd-53	-0.588697815	0.325322641	NA
str-168	-0.611511779	0.293797037	NA
sre-4	-0.659337898	0.279814818	NA
srx-98	-0.789769153	0.126804165	NA
srt-39	-0.803397597	0.204138417	NA
srz-99	-0.866876045	0.048915705	NA
sre-44	-1.105518367	0.091985454	NA

776 **Supplementary Table 2.** List of all exopher-related phenotypes presented in the manuscript.

Figure no.	Condition	Observed phenotype
1c	Exopher level in hermaphrodites in the presence of *him-5(e1490)* males	Increased exopher number in *him-5(e1490)* mutant hermaphrodites
1d	*In utero* embryo retention in the presence of *him-5(e1490)* males	Increased embryo retention in *him-5(e1490)* mutant hermaphrodites
1f	Effect on exopher production in hermaphrodites grown on WT and *him-5* males-conditioned plates	Increased exopher number after exposure to *him-5* and WT males secretome
1g	Effect on *in utero* embryo retention in hermaphrodites grown on WT and *him-5* males-conditioned plates	Increased *in utero* embryo retention after exposure to *him-5* and WT males secretome

1h	Effect of male pheromones presence on exopher production in different developmental time frames	Increased exopher number (strongest effect on exposure during L4 to AD1 stages)
1i	Effect of male pheromones presence on *in utero* embryo retention in different developmental time frames effect	Increased *in utero* embryo retention (apart from exposure during L4 to AD1 stages)
2b	Population size effect on exopher production (single or 10 hermaphrodites)	Single worms produce more exophers than worms cultured in 10 hermaphrodite population
2c	Population size effect on *in utero* embryo retention	No effect
2d	Population size effect on exopher production (single, 5 or 10 hermaphrodites)	Exopher number decreased in 5 and 10 hermaphrodites populations
2e	Population size effect on exopher production (single, 10 or 100 hermaphrodites)	Exopher number decreased in 10 and 100 hermaphrodite populations
2f	Hermaphrodites-conditioned plates effect on exopher production	Hermaphrodite pheromones similarly decrease exopher number as hermaphrodite presence
2g	Larvae presence effect on exopher production	Larvae presence decreases exopher production
3c	Influence of ascaroside side-chain biosynthesis mutants presence on exophergenesis level in wild-type worms	Number of exophers elevated in the presence of *acox-1* mutants; decreased exopher level in the presence of *maoc-1* mutants; *daf-22* mutants presence does not affect exophergenesis
3d	Influence of ascaroside side-chain biosynthesis mutants presence on *in utero* embryo retention	No influence
3e	Effect of ascr#10 exposure on exophergenesis	Dose-dependent effect– increase in exopher production after 1ng ascr#10 exposure; exposure to 1 pg ascr#10 has no effect
3f	Effect of assr#10 on *in utero* embryo retention	No effect
3g	Effect of ascr#18 exposure on exophergenesis	Decrease in exopher number after exposure to 1pg or 1ng of ascr#18
3h	Effect of ascr#18 on *in utero* embryo retention	Decreased in utero retention after exposure to 1ng of ascr#18

4b	Influence of impaired ciliated sensory neurons on exophergenesis in *che-13* mutant	*che-13* mutant produces fewer exophers than wild-type worms
4c	Genetic ablation of chemosensory neurons effect on exopher production	ASK(-), AWC(-), ADL(-), or ASH(-) mutants produce fewer exophers
4d	Genetic ablation of chemosensory neurons effect on *in utero* embryo retention	ASK(-), AWC(-), ADL(-), ASH(-) and CEM(+) have decreased *in utero* embryo retention
4e	Influence of AWC neuron absence on exophergenesis in different temperatures	AWC neurons regulate the temperature-dependent increase in muscle exopher production
4f	Male-specific pheromones effect on exophergenesis in mutants with ablated chemosensory neurons	ASK(-), ADL(-), AWB(-) – no effect WT, AWC(-), ASH(-), ASI(-), CEM(+) – increased exophergenesis after male pheromones exposure
4g	Population size effect on exopher production in mutants with ablated chemosensory neurons	ASH(-), ASI(-), CEM(-), ASK(-) – no effect AWC(-), ADL(-), AWB(-) – exopher number decreased in worms grown in the population of 10 hermaphrodites
5a	*str-173* expression visualization in L1, L4 and AD2 worms	*str-173* is expressed in ASK neurons, vulva, and the tail in L4 and adult worms; L1 larvae express *str-173* in pharynx
5b	*str-173* expression visualization in L4 worms	*str-173* is expressed in ASK neurons
5c	Exopher level in *str-173* mutants	Exopher production is not altered in *str-173* mutants
5d	Population size effect on exopher production in *str-173* mutants	*str-173(wwa1)* mutation does not affect exopher production in worms grown in separation or in 10 hermaphrodites population; *str-173(wwa2)* – higher exopher level in worms grown in separation
5e	Male-specific pheromones effect on exopher production in *str-173* mutants	No effect
5f	Effect of ascr#10 exposure on exophergenesis in *str-173* mutants	Decrease in exopher production
5g	Effect of ascr#10 on eggs in utero retention in *str-173* mutants	Decreased *in utero* embryo retention
5h	Effect of ascr#18 exposure on exophergenesis in *str-173* mutants	Decrease in exopher production

5i	Effect of ascr#18 on eggs in utero retention in *str-173* mutants	Decreased *in utero* embryo retention
5j	Effect of ascr#10 and ascr#18 exposure on exophergenesis in ASK(-) mutant	No effect
6a	Ablation of AQR/PQR/URX neurons effect on exopher level	Increased exopher number in AQR/PQR/URX(-) mutant
6b	Ablation of AQR/PQR/URX neurons effect *in utero* embryo retention	Decreased *in utero* embryo retention in AQR/PQR/URX(-) mutant
6c	ArchT-mediated AQR/PQR/URX inactivation effect on exopher production (AD2)	Increased exopher number after AQR/PQR/URX inactivation
6d	ReaChR-mediated AQR/PQR/URX activation effect on exopher production (AD2)	Decreased exopher number after AQR/PQR/URX activation
6e	Ablation of ASK/AQR/PQR/URX neurons effect on exopher level	Low exophergenesis levels in animals with genetic ablation of ASK neurons and high levels of exophergenesis in animals with genetic ablation of AQR, PQR, and URX neurons are equalized in animals with all four neurons removed
6f	Population size effect on exopher production in AQR/PQR/URX(-) mutant	No effect
6g	Male-specific pheromones effect on exopher production in AQR/PQR/URX(-) mutant	Increase in exopher production after exposure to male secretome
6h	FUdR effect on exophergenesis in AQR/PQR/URX(-) mutant	Decreased exopher production in aminals treated with FUdR is not rescued by AQR/PQR/URX removal
6i	*rme-2* knockdown effect on exophergenesis in AQR/PQR/URX(-) mutant	AQR/PQR/URX removal only partially rescues the inhibition of exophergenesis caused by *rme-2* knockdown
7a	Exopher level in *flp-8* and *flp-21* mutants	Increased exopher production in single *flp-8* and *flp-21* mutants, and in *flp-8; flp-21* double mutant
7b	Number of in utero embryos in *flp-8* and *flp-21* mutants	No change

7c	ReaChR-mediated AQR/PQR/URX; *flp-8(tm741)* activation effect on exopher production (AD2)	Increased exopher production
7d	ReaChR-mediated AQR/PQR/URX; *flp-21(ok889)* activation effect on exopher production (AD2)	No effect
7e	ReaChR-mediated AQR/PQR/URX; *flp-8(tm741)*; *flp-21(ok889)* activation effect on exopher production (AD2)	No effect
Sup. Fig. 1b	*him-5* males secretions effect on *him-5* hermaphrodites' exopher production	No effect
Sup. Fig. 1c	*him-5* males presence effect on *him-5* hermaphrodites eggs in utero retention	Increased number of *in utero* embryo retention when *him-5* males are present
Sup. Fig. 2a	Food source (metabolically active and inactive bacteria) effect on exopher production	Worms fed with metabolically active and inactive bacteria produce a similar number of exophers
Sup. Fig. 2b	Food source (*E. coli* strain OP50 and HT115) effect on *in utero* embryo retention	More embryos were retained in the uterus when grown on HT115
Sup. Fig. 2c	Males-conditioned plates seeded with HT115 bacteria strain effect on exopher production	Increase in exopher production
Sup. Fig. 3a	Male-specific pheromones effect on egg retention in mutants with ablated chemosensory neurons	ASI(-) – increased number of *in utero* embryos after exposure to male pheromones ASK(-), ADL(-), AWB(-) – no change
Sup. Fig. 3b	Population size effect on *in utero* embryos retention in mutants with ablated chemosensory neurons	ASI(-) -more embryos retained in worms grown in the population of 10 hermaphrodites AWB(-) – no change
Sup. Fig. 3c	Influence of ascaroside side-chain biosynthesis mutants presence on exophergenesis level in ASI(-) mutants	No influence
Sup. Fig. 3d	Influence of ascaroside side-chain biosynthesis mutants presence on exophergenesis level in AWB(-) mutants	No influence

Sup. Fig. 4a	Population size effect on the number of exophers produced by worms that were used for RNAseq analysis	Worms cultured in a population of 10 hermaphrodites – fewer exophers produced in each biological repeat
Sup. Fig. 5a	Number of eggs laid by AQR/PQR/URX(-) mutant	Decreased number of laid eggs by AQR/PQR/URX(-) mutant
Sup. Fig. 5b	Number of eggs laid by AQR/PQR/URX(-) mutant on consecutive days of adulthood	No significant change
Sup. Fig. 5c	ArchT-mediated AQR/PQR/URX inactivation effect on exopher production (no ATR control)	No change
Sup. Fig. 5d	ReaChR-mediated AQR/PQR/URX activation effect on exopher production (no ATR control)	No change
Sup. Fig. 5e	ArchT-mediated AQR/PQR/URX inactivation effect on exopher production (AD1)	No change
Sup. Fig. 5f	ArchT-mediated AQR/PQR/URX inactivation effect on exopher production (no ATR control; AD1)	No change
Sup. Fig. 5g	ReaChR-mediated AQR/PQR/URX activation effect on exopher production (AD1)	Decrease in exopher production after 30 min light stimulation
Sup. Fig. 5h	ReaChR-mediated AQR/PQR/URX activation effect on exopher production (no ATR control, AD1)	No change
Sup. Fig. 7a	ReaChR-mediated AQR/PQR/URX; *flp-8(tm741)* activation effect on exopher production (no ATR control)	No change
Sup. Fig. 7b	ReaChR-mediated AQR/PQR/URX; *flp-21(ok889)* activation effect on exopher production (no ATR control)	No change
Sup. Fig. 7c	ReaChR-mediated AQR/PQR/URX; *flp-8(tm741)*; *flp-21(ok889)* activation effect on exopher production (no ATR control)	No change

778 **Materials and Methods**

779 **Reagents and Tools table**

Reagent or Resource	Source	Identifier
Bacterial and Virus Strains		
Escherichia coli RNAi feeding strain	Caenorhabditis Genetics Center	HT115(DE3)
E. coli feeding strain	Caenorhabditis Genetics Center	OP50
Ahringer RNAi library	Source BioScience	*C. elegans* RNAi Collection (Ahringer)
Chemicals, Peptides, and Recombinant Proteins		
All-trans retinal	Sigma-Aldrich	R2500
Bacto Agar	BioShop	AGR001.1
Bacto Peptone	BioShop	PEP403.1
Bacto Tryptone	BioShop	TRP402.1
Sodium Chloride	Chempur	117941206
Streptomycin sulfate salt	Sigma-Aldrich	S6501
Nystatin suspension 10,000 unit/mL	Sigma-Aldrich	N1638
Carbenicillin disodium salt	Roth	6344.2

Magnesium Sulfate heptahydrate	Sigma-Aldrich	M5921
Potassium dihydrogen phosphate	Roth	3904.1
di-potassium hydrogen phosphate	Roth	P749.1
Calcium Chloride (dihydrate)	BioShop	CCL444.500
Cholesterol	BioShop	CHL500.5
Tetracycline HCl	BioShop	TET701
IPTG	BioShop	IPT001
Polystyrene microspheres	Polysciences Europe GmbH	08691-10
Tetramisole hydrochloride	Sigma-Aldrich	L9756
Paraformaldehyde	Sigma-Aldrich	158127
5-Fluoro-2'-deoxyuridine	Sigma-Aldrich	F0503
Sodium hypochlorite solution 14%	VWR	27900
Dimethyl sulfoxide	Sigma-Aldrich	D8418
Ascaroside #10	Synthesized in Frank Schroeder's laboratory	N/A

Ascaroside #18	Synthesized in Frank Schroeder's laboratory	N/A	

Caenorhabditis elegans strains

Genotype	Source	Identifier	Description
wacIs1[myo-3p::rpn-5 CAI = 0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)]	This paper	Strain ACH81	Strain with muscle exopher RFP marker in N2 wild-type background
wacIs1[myo-3p::rpn-5 CAI = 0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1– 50aa::attB5::mGFP::un c-54-3'UTR, unc-119(+)]	Turek *et al.*, 2021	Strain ACH93	Strain with muscle exopher RFP and mitochondrial GFP marker in N2 wild-type background
daf-22(ok693)	CGC	Strain RB859	
wacIs1[myo-3p::rpn-5 CAI = 0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1– 50aa::attB5::mGFP::un c-54-3'UTR, unc-119(+)]; daf-22(ok693)	This paper	Strain WOP456	Strain with muscle exopher RFP and mitochondrial GFP marker in *daf-22(ok693)* mutant background
acox-1(ok2257)	CGC	Strain VC1785	

wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; acox-1(ok2257) I	This paper	Strain TUR7	Strain with muscle exopher RFP and mitochondrial GFP marker in *acox-1(ok2257)* mutant background
maoc-1&E04F6.6(ok2645) II.	CGC	Strain RB2000	
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)]; maoc-1&E04F6.6(ok2645) II	This paper	Strain TUR46	Strain with muscle exopher RFP marker in *maoc-1&E04F6.6(ok2645)* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; che-13(e1805)	This paper	Strain TUR11	Strain with muscle exopher RFP and mitochondrial GFP marker in *che-13(e1805)* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; qrIs2 [sra-9::mCasp1]	This paper	Strain TUR16	Strain with muscle exopher RFP, mitochondrial GFP marker, and ASK neuron genetic ablation[48]

wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; hlh-4(tm604) III	This paper	Strain TUR75	Strain with muscle exopher RFP, mitochondrial GFP marker, and ADL neuron genetic ablation[49]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::Optimal Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)]; oyIs85 [ceh-36p::TU#813 + ceh-36p::TU#814 + srtx-1p::GFP + unc-122p::dsRed]	This paper	Strain WOP499	Strain with muscle exopher RFP, mitochondrial GFP marker, and AWC neuron genetic ablation[50]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; oyIs84 [gpa-4p::TU#813 + gcy-27p::TU#814 + gcy-27p::GFP + unc-122p::DsRed]	This paper	Strain TUR49	Strain with muscle exopher RFP, mitochondrial GFP marker, and ASI neuron genetic ablation[50]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::un	This paper	Strain TUR47	Strain with muscle exopher RFP, mitochondrial GFP marker, and ASH neuron genetic ablation[51]

c-54-3'UTR, unc-119(+)]; peIs1713 [sra-6p::mCasp-1 + unc-122p::mCherry]			
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; peIs1715 [str-1p::mCasp-1 + unc-122p::GFP]	This paper	Strain TUR51	Strain with muscle exopher RFP, mitochondrial GFP marker, and AWB neuron genetic ablation[51]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]nIs133 [pkd-2::GFP] I, ceh-30(n3714) X	This paper	Strain TUR18	Hermaphrodites of this strain have muscle exopher RFP, mitochondrial GFP marker, and contain CEM male-specific ciliated sensory neurons[25]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], qaIs2241[gcy-36::egl-1 + gcy-35::GFP + lin-15(+)]	This paper	Strain TUR37	Strain with muscle exopher RFP marker and AQR/PQR/URX neurons genetic ablation[52]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)],	This paper	Strain TUR59	Strain with muscle exopher RFP and mitochondrial GFP marker in

wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; him-5(e1490) V			*him-5(e1490)* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)]; wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; wwaEx4[gcy-36 promoter::ReAChR::mKate2::unc-54 3'UTR, unc-119(+); myo-2 promoter::mNeonGreen]	This paper	Strain TUR38	Strain with muscle exopher RFP, mitochondrial GFP marker, and ReAChR for optogenetic activation of AQR, PQR, and URX
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)]; wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; wwaEx9[gcy-36 promoter::ArchT::mKate2::unc-54 3'UTR, unc-119(+); myo-2 promoter::mNeonGreen]	This paper	Strain TUR43	Strain with muscle exopher RFP, mitochondrial GFP marker, and ArchT for optogenetic inactivation of AQR, PQR, and URX
str-173(syb6501[str-173::SL2-wrmScarlet]) IV	This paper	PHX6501	Endogenous translational reporter strain for *str-173* expression

str-173(syb6501[str-173::SL2-wrmScarlet]) IV, pkIs591[dpy-20(+) + gap-15::GFP]	This paper	TUR77	Endogenous translational reporter strain for *str-173* expression crossed with strain expressing GFP in ASK, ADL, ASH, PHA, an PHB
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; qrIs2 [sra-9::mCasp1]; qaIs2241[gcy-36::egl-1 + gcy-35::GFP + lin-15(+)]	This paper	Strain TUR61	Strain with muscle exopher RFP, mitochondrial GFP marker, and ASK/AQR/PQR/URX neurons genetic ablation
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; qaIs2241[gcy-36::egl-1 + gcy-35::GFP + lin-15(+)]	This paper	Strain TUR62	Strain with muscle exopher RFP, mitochondrial GFP marker, and AQR/PQR/URX neurons genetic ablation[52]
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3	This paper	Strain WOP558	Strain with muscle exopher RFP and mitochondrial GFP marker in *str-173(wwa1)*

promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; str-173(wwa1) IV			mutant background, backcrossed x2 with N2
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; str-173(wwa2) IV	This paper	Strain WOP559	Strain with muscle exopher RFP and mitochondrial GFP marker in *str-173(wwa2)* mutant background, backcrossed x6 with N2
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)];flp-8(pk360) X	This paper	Strain TUR84	Strain with muscle exopher RFP and mitochondrial GFP marker in *flp-8(pk360) X* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)];flp-21(ok889) V.	This paper	Strain TUR85	Strain with muscle exopher RFP and mitochondrial GFP marker in *flp-21(ok889) V* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-	This paper	Strain TUR93	Strain with muscle exopher RFP, mitochondrial

54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; flp-8(pk360) X; wwaEx4[gcy-36 promoter::ReAChR::mKate2::unc-54 3'UTR, unc-119(+); myo-2 promoter::mNeonGreen]			GFP marker, and ReAChR for optogenetic activation of AQR, PQR, and URX in *flp-8(pk360) X* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; flp-21(ok889) V; wwaEx4[gcy-36 promoter::ReAChR::mKate2::unc-54 3'UTR, unc-119(+); myo-2 promoter::mNeonGreen]	This paper	Strain TUR94	Strain with muscle exopher RFP, mitochondrial GFP marker, and ReAChR for optogenetic activation of AQR, PQR, and URX in *flp-21(ok889) V* mutant background
wacIs1[myo-3 promoter::rpn-5 CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; flp-8(pk360) X; flp-21(ok889) V	This paper	TUR97	Strain with muscle exopher RFP and mitochondrial GFP marker in *flp-8(pk360) X; flp-21(ok889) V* mutant background
wacIs1[myo-3 promoter::rpn-5	This paper	Strain TUR96	Strain with muscle exopher

CAI=0.97::GGGGS Linker-wrmScarlet::unc-54 3'UTR, unc-119(+)], wacIs14[myo-3 promoter::tomm-20_1-50aa::attB5::mGFP::unc-54-3'UTR, unc-119(+)]; flp-8(tm741) X; flp-21(ok889) V; wwaEx4[gcy-36 promoter::ReaChR::mKate2::unc-54 3'UTR, unc-119(+); myo-2 promoter::mNeonGreen]			RFP, mitochondrial GFP marker, and ReaChR for optogenetic activation of AQR, PQR, and URX in *flp-8(pk360) X; flp-21(ok889) V* mutant background
wild type	Caenorhabditis Genetics Center	Strain N2	*C. elegans* wild isolate
Recombinant DNA			
Plasmid: pGLOW77 (myo-2 promoter::mNeonGreen)	Addgene	#177338	
Plasmid: pAS01 (gcy-36 promoter::ReaChR::mKate2-unc-54 3'UTR in pCG150)	This paper	N/A	
Plasmid: pAS02 (gcy-36 promoter::ArchT::mKate2-unc-54 3'UTR in pCG150)	This paper	N/A	
Plasmid: pAZ03 (mKate2-unc-54 3'UTR in pCG150)	This paper	N/A	
Plasmid: ArchT (ArchT template)	gift from Bringmann Lab	N/A	

Plasmid: ReaChR (ReaChR template)	gift from Bringmann Lab	N/A
Plasmid: mKate2-unc-54 3'UTR (mKate2-unc-54 3'UTR template)	gift from Bringmann Lab	N/A
Plasmid: pMH389 (gcy-36 promoter template)	gift from Kapitein Lab	N/A
Plasmid: pCG150	Addgene	#17247
Oligonucleotides		
Used for *acox-1(ok2257)* genotyping	FP1 AGTAGAGGCTGACGGGACTT FP2 CTCGCTCGTTACCTCGTCAA	RP GGGCAAATGGACATCAAGGC
Used for *che-13(e1805)* genotyping	FP1 ATCCGTATTGCTTTGGAGAACGCTGTA FP2 GAGTTCATCTGTTCAAGATGAAGACGCC	RP1 TGATGAGTTTCAGAAGAATCCACGCC RP2 CAGTGGATGAGCAAGATGAAGATGATGA
Used for *daf-22(ok693)* genotyping	FP ACATTTGCCACGTAGCTGGT	RP1 TGGGCCGAGACAAAGCTTAC RP2 GTGGTAAGGGAAAAGCGCAA
Used for *maoc-1(ok2645)* genotyping	FP ACACGTTGT	RP1 AGTCTTGACCCCATGTGTTGA

	ACATCTCTCCCG	RP2 CAGCAATTGAACCACGTCCA
Used for *flp-8(pk360)* genotyping	FP TGCTCTCAAAGACACGTTTATTG	RP1 CACGCCTTCTTGTGCTGAA RP2 CGGTAAGCAGTTTTGAAGCAT
Used for *flp-21(ok889)* genotyping	FP ACAAAGTCATCGAGCAGTAGAT	RP1 TTCCGGCCACCTCAACTATG RP2 ACAACGAGATCCTAAAACGACA
Used for creating pAS01		
-gcy-36	FP TATGACCATGATTACGCCAAGCTATCTGGCACACTTTTTATTTCCATAAACCTGC	RP TTGGCGCGCCCCACTTTGTACAAGAAAGCTGGGTTCCGGATCCTCCTCCGGATC
-ReaChR	FP AAATTCAAACAAGGGCTACCCAACAATGGTCTCCCGTCGTCCATG	RP TTGGCGCGCCCCACTTTGTACAAGAAAGCTGGGTTCCGGATCCTCCTCCGGATC
Used for creating pAS02		
-gcy-36	FP TATGACCATGATTACGCCAAGCTATCTGGCACACTTTTTATTTCCATAAACCTGC	RP TTGGAGGGCGATTGGGTCCATTTTTTGTTGGGTAGCCCTTGTTTGAATTTAC

-ArchT	FP AAATTCAAA CAAGGGCTA CCCAACAAA AAATGGACC CAATCGCCC TC	RP TTGGCGCGCCCCACTTTGTA CAAGAAAGCTGGGTTTTCCT CCTGGCTCTGGGG
Software and Algorithms		
GraphPad Prism 9	GraphPad Software, Inc.	N/A
ZEN	Zeiss	N/A
Office 365	Microsoft	N/A
ImageJ	Wayne Rasband (NIH)	N/A
BioRender	BioRender.com	N/A
Adobe Illustrator CS6	Adobe	N/A

Methods and Protocols

Worm maintenance and strains

Worms were maintained on nematode growth medium (NGM) plates seeded with bacteria

Escherichia coli OP50 or HT115 at 20°C unless stated otherwise[19]. A list of all *C. elegans*

strains used in the study is provided in the table at the beginning of the "Materials and Methods"

section.

Scoring exophers and fluorescence microscopy

For the assessment of exophers number, the protocol described in Turek *et al*. and Banasiak *et*

al. (Bio-Protocol, detailed, step-by-step guide) was applied. Briefly, exophers were scored

using stereomicroscope Zeiss Axio Zoom.V16 with filter sets 63 HE and 38 HE. For each

exopher scoring assay, worms were age-synchronized either from pretzel-stage embryos, or

larval stages L1 and L4. On adulthood day-2, animals were visualized on NGM plates, and the

number of exophers was counted in each freely moving worm.

The representative pictures presented in the manuscript were acquired with an inverted Zeiss

LSM800 laser-scanning confocal microscope with a 40x oil immersion objective. To excite the

GFP and RFP fluorescent proteins, 488- and 561-nm lasers were used. For visualization,

animals were immobilized on 3% agarose pads with 6µl of PolySciences 0.05 µm polystyrene

microspheres or 25 µM tetramisole.

RNA interference assay

RNA interference in *C. elegans* was performed using the standard RNAi feeding method and

RNAi clone[53], previously described in Turek *et al.,* 2021[6]. Briefly, NGM plates, supplemented

with 12,5µg/ml tetracycline, 100µg/ml ampicillin, and 1mM IPTG, were seeded with *E. coli*

HT115 bacteria containing dsRNA against the gene of interest. Control group animals were fed

with bacteria containing the empty vector L4440. Age-synchronized pretzel-stage embryos or

L1 larvae were placed on freshly prepared plates and cultured until day 2 of adulthood when

exophers were scored. The age of the worms was verified at the L4 larvae stage, either younger

or older worms were removed from the experiment.

Pre-conditioning of the plates with *C. elegans* males

50 males from the strain with *him-5(e1490)* mutation were transferred on a fresh, 35 mm NGM

plate with *E. coli* OP50 or HT115 at the L4 developmental stage and older. After 48 hours,

males were removed from the plates.

Culturing worms from L1 to adulthood day 2 on plates pre-conditioned with males

L1 larvae were transferred to plates pre-conditioned with males in a group of 10 hermaphrodites

per plate. For the control group, L1 larvae were transferred to fresh, 35 mm NGM plates seeded

with *E. coli* OP50 or HT115 without pre-conditioning with males. Worms were cultured up to

adulthood day 2 when muscle exophers were counted in each animal.

Culturing worms in 24 hours intervals from L1 to adulthood day 2 on plates pre-conditioned

with males

Worms were kept on plates pre-conditioned with males only for 24 hours at different

developmental stages. All hermaphrodites were picked at L1 stage and grouped into 10 worms

per plate population. The first group was cultured on pre-conditioned plates for 24 hours

starting from hatching from the egg. After 24 hours worms were transferred to a fresh NGM

plate seeded with OP50 bacteria. The second group was grown for 24 hours after hatching from

the egg on the NGM plate, then the worms were moved to pre-conditioned with males plates

for 24 hours, after that time they were transferred to fresh NGM plates seeded with OP50

bacteria. The third group was cultured for the first 48 hours on NGM plates, then transferred

to pre-conditioned with males plates for 24 hours, and then replated on fresh NGM plates. The

fourth group was cultured from the L1 stage for 72 hours on standard NGM plates, after that

time worms were transferred to plates pre-conditioned with males for 24 hours. All

experimental groups were cultured from L1 to adulthood day 2 when muscle exophers were

counted.

Egg retention assay

Firstly, using the stereomicroscope, muscle exophers were counted in each worm. In the following step, hermaphrodites were exposed to a 1.8% hypochlorite solution. When they dissolved, embryos retained in the uterus were counted.

Metabolic inactivation of bacterial food source

The preparation of plates with a metabolically inactive food source for the worms to determine its effects on exophergenesis was done according to the protocol described in Beydoun *et al.*, 2021[54]. Briefly, a single *E. coli* HT115 or OP50 colony was inoculated overnight and then the bacterial culture was split into two flasks. Next, paraformaldehyde (PFA) was added to a final concentration of 0.5% in only one of them, and the flask was placed in the 37°C shaking incubator for 1 h. Afterward, the aliquots were transferred to 50 mL tubes, centrifuged at approximately 3000 × g for 20 min, and washed with 25 mL of LB five times. Control and PFA-treated bacteria were later concentrated accordingly and seeded on the NGM plates.

As control of PFA treatment, new bacterial cultures in LB were set in the 37°C shaking incubator overnight to ensure the replication was blocked and there was no bacterial growth.

Plates were used for experiments at least 5 days after their preparation to ensure no bacteria-derived metabolites were left on plates with PFA-treated *E. coli*. A new batch of bacterial food source was prepared for each biological repetition.

Brood size quantification

Age-synchronized hermaphrodites at the L4 developmental stage were picked for the experiment. Worms were cultured as single worms per 60 mm NGM plate, seeded with *E. coli*

853   OP50 or HT115. Ten worms were counted as one biological repeat. Worms were transferred

854   to fresh plates every day from adulthood day 1 until the end of the egg laying period.

855   The number of eggs laid over the worms' reproductive lifetime was counted manually daily.

856   The data is presented as the total number of eggs laid by each animal.

Quantifying the number of exophers in worms grown in different temperatures

Worms were confronted with a range of physiological temperatures: low 15 $^{\circ}$C, optimal 20 $^{\circ}$C,

and high 25 $^{\circ}$C throughout their development until exophers were scored. To compare the

corresponding stages of development at various temperatures, worms were additionally sorted

at the L4 stage. The exopher number was assessed based on the timepoint of maximal egg

laying, approx. 140 h or 78 h from egg hatching at low or high maintenance temperatures,

respectively. Timing calculations were based on the *C. elegans* development timeline at

different temperatures[55].

Culturing worms in different population sizes

L1 larvae were transferred to 35 mm NGM Petri dishes seeded with *E. coli* OP50 or HT115

bacteria strains (100 µL of bacteria from 10 mL overnight culture grown in 50 mL Erlenmeyer

flask) immediately after hatching as a single larva or in a group of 5, 10 or 100 animals. In

total, thirty plates with single worms, six plates with 5 worms, three plates with 10 worms, and

one plate with 100 worms were used per biological repeat. Exophers were scored on adulthood

day 2 in the worms expressing RFP in BWM.

Quantifying the influence of molecules released by ascaroside biosynthesis mutant on exopher production in wild-type worms

Nine freshly hatched L1 larvae from selected ascaroside biosynthesis mutant strains (*maoc-*

1(ok2645), daf-22(ok693), or *acox-1(ok2257)*) or wild-type hermaphrodites (as a control) were

transferred to fresh NGM plates. Next, one L1 larva of a reporter strain expressing RFP in

BWM was added to each plate and 10 worms in total were grown together on 35 mm Petri dish

seeded with *E. coli* OP50 or HT115 bacteria strains. On adulthood day 2 the number of

exophers was quantified in the worms expressing RFP in BWM, according to description.

Generation of *str-173* mutant strains

The *str-173* gene mutants (*str-173(wwa1)* and *str-173(wwa2)*) were generated using

CRISPR/Cas9 method as previously described[56]. The crRNA sequence used was

ATAATTGGTGGATATACAAATGG. The *str-173* gene locus was sequenced, and deletions

were mapped to the first exon (Extended Data Figure 3d). Both mutations cause frame shifts

and are most likely molecular null alleles. Strains were backcrossed twice (strain *str-*

173(wwa1) and 6 times *(str-173(wwa2))* against N2 wild-type worms.

Generation of optogenetic strains

Optogenetic strains created for this paper contain red-shifted Channel Rhodopsin (ReAChR) or

archaerhodopsin from Halorubrum strain TP009 (ArchT). To generate these strains, firstly,

mKate2-unc-54 3'UTR was amplified from the template and cloned into pCG150 to create a

pAZ03 plasmid. Next, the *gcy-36* promoter was amplified from pMH389, and ReAChR and

ArchT were amplified from respective templates. The *gcy-36* promoter was then cloned into

pAZ03 plasmid with ReAChR and ArchT separately. As a result two plasmids were created:

gcy-36   promoter::ReAChR::mKate2-unc-54   3'UTR   in   pCG150   and   gcy-36

promoter::ArchT::mKate2-unc-54 3'UTR in pCG150. Plasmids were sequenced to verify the

correct sequence of the cloned constructs. All constructs generated for this study were made

using the SLiCE method[57].

Transgenic strains with extrachromosomal arrays were generated by microinjection. DNA was

injected into exopher reporter strain worms with muscle exopher RFP and mitochondrial GFP

marker (ACH93). For injection, DNA was prepared as follows: construct 90 ng/µL and co-

injection marker 10 ng/µL. Positive transformants were selected according to the presence of

co-injection markers (myo-2 promoter::mNeonGreen).

The constructs created for this project and primers used for amplification are listed in the table

at the beginning of the "Materials and Methods" section.

Optogenetics assays

On adulthood day 2

For optogenetic activation or inhibition, 35-mm NGM plates seeded with HT115 *E. coli*

bacteria were covered with 0.2 µM all-trans retinal (ATR). Control plates were not covered

with ATR. Ten to twelve age-synchronized worms were picked per plate from optogenetic

strains (expressing ReAChR or ArchT in AQR/PQR/URX neurons) at adulthood day 1. After

24-hour incubation at 20°C and darkness, muscle exophers extruded by worms were counted

(**1**). Next, experimental plates were placed on the stereomicroscope and illuminated for 1 hour

with green light (HXP 200C illuminator as a light source, band-pass filter Zeiss BP 572/25

(HE), the green light intensity measured at 561 nm = 0.07 mW/mm^2). Immediately after

illumination muscle exophers were counted (**2**). Subsequently, exophers scorings were

performed 15 min (**3**) and 30 min (**4**) after the illumination with a green light was completed.

Worms were kept in the darkness in between counts. The control group was not illuminated.

To provide similar environmental conditions control plates were placed next to the

experimental plate but were shielded from light. Control and treated groups were randomized

before the start of the experiment.

 On adulthood day 1

The assay was conducted similarly as on adulthood day 2 with a difference in time of exposure to the light. Twelve age-synchronized worms from optogenetic strains were picked at the L4 developmental stage and transferred on a 35-mm NGM plate covered with ATR (no ATR for the control group). Worms were then incubated for 24 hours at 20 °C and darkness. Experimental plates were placed on the stereomicroscope and illuminated for 1 hour with green light (HXP 200C illuminator as a light source, band-pass filter Zeiss BP 572/25 (HE), the green light intensity measured at 561 nm = 0.07 mW/mm^2). Exophers extruded from BWM were counted before the exposure to the light (**1**), immediately after the exposure to the light (**2**), and 15 min (**3**), 30 min (**4**), and 24 hours (**5**) after the exposure to the light.

FUdR assay

Age-synchronized young adult animals (day 0) were placed on NGM plates containing 25 μM fluorodeoxyuridine (FUdR) or control NGM plates without FUdR. Exophers number were scored when worms reached adulthood day 2 using a stereomicroscope.

Transcriptome analysis

RNA extractions, library preparations, and sequencing were conducted at Azenta US, Inc (South Plainfield, NJ, USA) as follows:

RNA Extraction

Total RNA was extracted using Qiagen RNeasy Plus mini kit following the manufacturer's instructions (Qiagen, Hilden, Germany).

Library Preparation with polyA selection and Illumina Sequencing

Extracted RNA samples were quantified using Qubit 2.0 Fluorometer (Life Technologies,

Carlsbad, CA, USA) and RNA integrity was checked using Agilent TapeStation 4200 (Agilent

Technologies, Palo Alto, CA, USA).

RNA sequencing libraries were prepared using the NEBNext Ultra II RNA Library Prep Kit

for Illumina following the manufacturer's instructions (NEB, Ipswich, MA, USA). Briefly,

mRNAs were first enriched with Oligo(dT) beads. Enriched mRNAs were fragmented for 15

minutes at 94 °C. First strand and second strand cDNAs were subsequently synthesized. cDNA

fragments were end repaired and adenylated at 3'ends, and universal adapters were ligated to

cDNA fragments, followed by index addition and library enrichment by limited-cycle PCR.

The sequencing libraries were validated on the Agilent TapeStation (Agilent Technologies,

Palo Alto, CA, USA), and quantified by using Qubit 2.0 Fluorometer (Invitrogen, Carlsbad,

CA) as well as by quantitative PCR (KAPA Biosystems, Wilmington, MA, USA).

The sequencing libraries were multiplexed and loaded on the flowcell on the Illumina NovaSeq

6000 instrument according to the manufacturer's instructions. The samples were sequenced

using a 2x150 Pair-End (PE) configuration v1.5. Image analysis and base calling were

conducted by the NovaSeq Control Software v1.7 on the NovaSeq instrument. Raw sequence

data (.bcl files) generated from Illumina NovaSeq was converted into fastq files and de-

multiplexed using Illumina bcl2fastq program version 2.20. One mismatch was allowed for

index sequence identification.

Sequencing Data Analysis

After investigating the quality of the raw data, sequence reads were trimmed to remove possible

adapter sequences and nucleotides with poor quality using Trimmomatic v.0.36. The trimmed

reads were mapped to the *Caenorhabditis elegans* reference genome available on ENSEMBL

using the STAR aligner v.2.5.2b. The STAR aligner is a splice aligner that detects splice

junctions and incorporates them to help align the entire read sequences. BAM files were

generated as a result of this step. Unique gene hit counts were calculated by using feature

Counts from the Subread package v.1.5.2. Only unique reads that fell within exon regions were

counted.

After the extraction of gene hit counts, the gene hit counts table was used for downstream

differential expression analysis. Using DESeq2, a comparison of gene expression between the

groups of samples was performed. The Wald test was used to generate p-values and Log2 fold

changes. Genes with adjusted p-values < 0.05 and absolute log2 fold changes > 1 were called

as differentially expressed genes for each comparison.

RNAseq data was deposited in the GEO database (GSE241786) and can be accessed using the

following links: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE241786

Growing hermaphrodite on plates pre-conditioned with other hermaphrodites

50 wild-type hermaphrodites at the L4 stage were placed on a fresh, 35 mm NGM plate seeded

with *E. coli* OP50 bacteria (100 μL of bacteria from 10 mL overnight culture grown in 50 mL

Erlenmeyer flask). After 12-16 hours, hermaphrodites were removed from the plate and a single

pretzel-stage embryo was transferred to the hermaphrodite-conditioned plate. For the control

groups single pretzel-stage embryos or ten pretzel-stage embryos were situated on a fresh, 35

mm NGM plate with *E. coli* OP50 bacteria without pre-conditioning with hermaphrodites.

When worms reached the second day of adulthood, the number of exophers was scored in each

animal.

Quantifying the influence of larval pheromones on exopher production in

hermaphrodites

A single pretzel-stage embryo was placed on a fresh, 35 mm NGM plate seeded with *E. coli*

OP50 bacteria (100 µL of bacteria from 10 mL overnight culture grown in a 50 mL Erlenmeyer

flask). The worm was cultured until the late L4 stage, when 50 L1 larvae were added to a single

worm. For the control groups, single pretzel-stage embryos or ten pretzel-stage embryos were

placed on a fresh, 35 mm NGM plate with *E. coli* OP50 bacteria. Muscle exophers were scored

in each animal at adulthood day 2.

Quantifying the influence of molecules released by ascaroside biosynthesis mutant on

exopher production

Nine freshly hatched L1 larvae from selected ascaroside biosynthesis mutant strains (*maoc-*

1(ok2645), *daf-22*(ok693), or *acox-1*(ok2257)) or wild-type hermaphrodites (as a control) were

transferred to fresh NGM plates. Next, one L1 larva of a reporter strain expressing RFP in body

wall muscles was added to each plate, and 10 worms in total were grown together up to

adulthood day 2. Muscle exophers were counted in each animal expressing RFP in BWM.

Growing hermaphrodites on plates supplemented with synthetic ascr#10 or ascr#18

Synthetic, concentrated ascr#10 and ascr#18 were stored in DMSO at -80°C. The stocks were

diluted to working solutions with water. A total of 100µL (containing 1ng or 1pg) was applied

and rubbed into the 35-mm NGM plate with a sterile glass rod. Plates were incubated at 20 °C

overnight. The following day plates were seeded with 100µL of *E.coli* OP50 bacteria, and

incubated for an additional 24 hours at 20 °C. Control plates were prepared similarly using

water instead of an ascaroside solution. Pretzel-stage eggs were transferred on plates

supplemented with ascarosides and control plates in groups of twelve eggs per plate. On day 2

adulthood the number of exophers was quantified in the worms expressing RFP in BWM.

Data analysis and visualization tools

The exophers were scored at adulthood day 2 using a stereomicroscope unless stated otherwise.

Data analysis was performed using Microsoft® Excel® and GraphPad Prism 9 software.

Graphical representation of data was depicted using GraphPad Prism 9.

Statistical analysis

No statistical methods were used to predetermine the sample size. Worms were randomly

allocated to the experimental groups for all the data sets and experiments were performed

blinded for the data sets presented in the following figures: Fig. 1; Fig. 3c, d, e, f, g, h; Fig. 4a,

d, g; Fig. 5c, d, e, f, g, h, i, j; Fig. 7a, b; Supplementary Fig. 3a, b. Non-Gaussian distribution

of residuals was assumed; therefore, nonparametric statistical tests were applied: Mann–

Whitney (in comparison between two groups) or Kruskal-Wallis test with Dunn's multiple

comparisons test (in comparison between more than two groups). P-value < 0.05 is considered

significant**.**

1023

References

1. Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* **7**, 1535750 (2018).

2. Colombo, M., Raposo, G. & Théry, C. Biogenesis, Secretion, and Intercellular Interactions of Exosomes and Other Extracellular Vesicles. *Annu. Rev. Cell Dev. Biol.* **30**, 255–289 (2014).

3. Fu, S. *et al.* Extracellular vesicles in cardiovascular diseases. *Cell Death Discov.* **6**, 68 (2020).

4. Hill, A. F. Extracellular Vesicles and Neurodegenerative Diseases. *J. Neurosci.* **39**, 9269 LP – 9273 (2019).

5. Kalluri, R. The biology and function of exosomes in cancer. *J. Clin. Invest.* **126**, 1208–1215 (2016).

6. Turek, M. *et al.* Muscle-derived exophers promote reproductive fitness. *EMBO Rep.* e52071 (2021) doi:10.15252/embr.202052071.

7. Melentijevic, I. *et al.* C. elegans neurons jettison protein aggregates and mitochondria under neurotoxic stress. *Nature* **542**, 367–371 (2017).

8. Cooper, J. F., Guasp, R. J., Arnold, M. L., Grant, B. D. & Driscoll, M. Stress increases in exopher-mediated neuronal extrusion require lipid biosynthesis, FGF, and EGF RAS/MAPK signaling. *Proc. Natl. Acad. Sci.* **118**, e2101410118 (2021).

9. Nicolás-Ávila, J. A. *et al.* A Network of Macrophages Supports Mitochondrial Homeostasis in the Heart. *Cell* **183**, 94-109.e23 (2020).

10. Wong, S. S., Yu, J., Schroeder, F. C. & Kim, D. H. Population Density Modulates the Duration of Reproduction of C. elegans. *Curr. Biol.* **30**, 2602-2607.e2 (2020).

11. Perez, M. F. *et al.* Neuronal perception of the social environment generates an

inherited memory that controls the development and generation time of C. elegans.

Curr. Biol. **31**, 4256-4268.e7 (2021).

12. Wasson, J. A. *et al.* Neuronal control of maternal provisioning in response to social

cues. *Sci. Adv.* **7**, (2021).

13. Hodgkin, J., Horvitz, H. R. & Brenner, S. Nondisjunction Mutants of the Nematode

CAENORHABDITIS ELEGANS. *Genetics* **91**, 67–94 (1979).

14. Burkhardt, R. N. *et al.* Sex-specificity of the C. elegans metabolome. *Nat. Commun.*

14, 320 (2023).

15. Ludewig, A. H. *et al.* An excreted small molecule promotes C. elegans reproductive

development and aging. *Nat. Chem. Biol.* **15**, 838–845 (2019).

16. Izrayelit, Y. *et al.* Targeted Metabolomics Reveals a Male Pheromone and Sex-

Specific Ascaroside Biosynthesis in Caenorhabditis elegans. *ACS Chem. Biol.* **7**, 1321–

1325 (2012).

17. Ludewig, A. H. & Schroeder, F. C. Ascaroside signaling in C. elegans. *WormBook* 1–

22 (2013) doi:10.1895/wormbook.1.155.1.

18. von Reuss, S. H. *et al.* Comparative metabolomics reveals biogenesis of ascarosides, a

modular library of small-molecule signals in C. elegans. *J. Am. Chem. Soc.* **134**,

1817–1824 (2012).

19. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

20. Kamath, R. S., Martinez-Campos, M., Zipperlen, P., Fraser, A. G. & Ahringer, J.

Effectiveness of specific RNA-mediated interference through ingested double-

stranded RNA in Caenorhabditis elegans. *Genome Biol.* **2**, RESEARCH0002 (2001).

21. Perkins, L. A., Hedgecock, E. M., Thomson, J. N. & Culotti, J. G. Mutant sensory cilia

in the nematode Caenorhabditis elegans. *Dev. Biol.* **117**, 456–487 (1986).

22.   Ferkey, D. M., Sengupta, P. & L'Etoile, N. D. Chemosensory signal transduction in

Caenorhabditis elegans. *Genetics* **217**, iyab004 (2021).

23.   Biron, D., Wasserman, S., Thomas, J. H., Samuel, A. D. T. & Sengupta, P. An

olfactory neuron responds stochastically to temperature and modulates Caenorhabditis

elegans thermotactic behavior. *Proc. Natl. Acad. Sci.* **105**, 11002–11007 (2008).

24.   Kuhara, A. *et al.* Temperature Sensing by an Olfactory Neuron in a Circuit Controlling

Behavior of C. elegans. *Science (80-.).* **320**, 803–807 (2008).

25.   Schwartz, H. T. & Horvitz, H. R. The C. elegans protein CEH-30 protects male-

specific neurons from apoptosis independently of the Bcl-2 homolog CED-9. *Genes*

Dev. **21**, 3181–3194 (2007).

26.   Srinivasan, J. *et al.* A blend of small molecules regulates both mating and development

in Caenorhabditis elegans. *Nature* **454**, 1115–1118 (2008).

27.   Macosko, E. Z. *et al.* A hub-and-spoke circuit drives pheromone attraction and social

behaviour in C. elegans. *Nature* **458**, 1171–1175 (2009).

28.   Qian, K.-Y. *et al.* Male pheromones modulate synaptic transmission at the C. elegans

neuromuscular junction in a sexually dimorphic manner. *Elife* **10**, e67170 (2021).

29.   Aprison, E. Z. & Ruvinsky, I. Counteracting Ascarosides Act through Distinct

Neurons to Determine the Sexual Identity of C. elegans

Pheromones. *Curr. Biol.* **27**, 2589-2599.e3 (2017).

30.   Robertson, H. M. & Thomas, J. H. The putative chemoreceptor families of C. elegans.

WormBook 1–12 (2006) doi:10.1895/wormbook.1.66.1.

31.   Gruner, M. *et al.* Feeding state, insulin and NPR-1 modulate chemoreceptor gene

expression via integration of sensory and circuit inputs. *PLoS Genet.* **10**, e1004707

(2014).

32.   Ryan, D. A. *et al.* Sex, Age, and Hunger Regulate Behavioral Prioritization through

Dynamic Modulation of Chemoreceptor Expression. *Curr. Biol.* **24**, 2509–2517

(2014).

33.   Wexler, L. R., Miller, R. M. & Portman, D. S. C. elegans Males Integrate Food Signals

and Biological Sex to Modulate State-Dependent Chemosensation and Behavioral

Prioritization. *Curr. Biol.* **30**, 2695-2706.e4 (2020).

34.   McLachlan, I. G. *et al.* Diverse states and stimuli tune olfactory receptor expression

levels to modulate food-seeking behavior. *Elife* **11**, (2022).

35.   Taylor, S. R. *et al.* Molecular topography of an entire nervous system. *Cell* **184**, 4329-

4347.e23 (2021).

36.   White, J. G., Southgate, E., Thomson, J. N. & Brenner, S. The structure of the nervous

system of the nematode Caenorhabditis elegans. *Philos. Trans. R. Soc. London. Ser. B,*

Biol. Sci. **314**, 1–340 (1986).

37.   Coates, J. C. & de Bono, M. Antagonistic pathways in neurons exposed to body fluid

regulate social feeding in Caenorhabditis elegans. *Nature* **419**, 925–929 (2002).

38.   Okazaki, A. & Takagi, S. An optogenetic application of proton pump ArchT to C.

elegans cells. *Neurosci. Res.* **75**, 29–34 (2013).

39.   Lin, J. Y., Knutsen, P. M., Muller, A., Kleinfeld, D. & Tsien, R. Y. ReaChR: a red-

shifted variant of channelrhodopsin enables deep transcranial optogenetic excitation.

Nat. Neurosci. **16**, 1499–1508 (2013).

40.   Rogers, C. *et al.* Inhibition of Caenorhabditis elegans social feeding by FMRFamide-

related peptide activation of NPR-1. *Nat. Neurosci.* **6**, 1178–1185 (2003).

41.   Aprison, E. Z., Dzitoyeva, S., Angeles-Albores, D. & Ruvinsky, I. A male pheromone

that improves the quality of the oogenic germline. *Proc. Natl. Acad. Sci. U. S. A.* **119**,

e2015576119 (2022).

42.   Corrigan, L. *et al.* BMP-regulated exosomes from Drosophila male reproductive

glands reprogram female behavior. *J. Cell Biol.* **206**, 671–688 (2014).

43.   Sanchez-Lopez, J. A. *et al.* Male-female communication enhances release of

extracellular vesicles leading to high fertility in Drosophila. *Commun. Biol.* **5**, 815

(2022).

44.   DeLong, J. P., Hanley, T. C. & Vasseur, D. A. Competition and the density

dependence of metabolic rates. *J. Anim. Ecol.* **83**, 51–58 (2014).

45.   Thomas, J. H. & Robertson, H. M. The Caenorhabditis chemoreceptor gene families.

BMC Biol. **6**, 42 (2008).

46.   Artyukhin, A. B. *et al.* Metabolomic 'Dark Matter' Dependent on Peroxisomal β-

Oxidation in Caenorhabditis elegans. *J. Am. Chem. Soc.* **140**, 2841–2852 (2018).

47.   Florman, J. T. & Alkema, M. J. Co-transmission of neuropeptides and monoamines

choreograph the C. elegans escape response. *PLoS Genet.* **18**, e1010091 (2022).

48.   Ludewig, A. H. *et al.* Pheromone sensing regulates Caenorhabditis

elegans lifespan and stress resistance via the deacetylase SIR-2.1. *Proc.*

Natl. Acad. Sci. **110**, 5522 LP – 5527 (2013).

49.   Masoudi, N. *et al.* Unconventional function of an Achaete-Scute homolog as a

terminal selector of nociceptive neuron identity. *PLOS Biol.* **16**, e2004979 (2018).

50.   Beverly, M., Anbil, S. & Sengupta, P. Degeneracy and Neuromodulation among

Thermosensory Neurons Contribute to Robust Thermosensory Behaviors in

Caenorhabditis elegans. *J. Neurosci.* **31**, 11718–11727 (2011).

51.   Yoshida, K. *et al.* Odour concentration-dependent olfactory preference change in C.

elegans. *Nat. Commun.* **3**, 739 (2012).

52.   Chang, A. J., Chronis, N., Karow, D. S., Marletta, M. A. & Bargmann, C. I. A

Distributed Chemosensory Circuit for Oxygen Preference in C. elegans. *PLOS Biol.* **4**,

e274 (2006).

53.   Kamath, R. S. & Ahringer, J. Genome-wide RNAi screening in Caenorhabditis

elegans. *Methods* **30**, 313–321 (2003).

54.   Beydoun, S. *et al.* An alternative food source for metabolism and longevity studies in

Caenorhabditis elegans. *Commun. Biol.* **4**, 258 (2021).

55.   Byerly, L., Cassada, R. C. & Russell, R. L. The life cycle of the nematode

Caenorhabditis elegans. I. Wild-type growth and reproduction. *Dev. Biol.* **51**, 23–33

(1976).

56.   Dokshin, G. A., Ghanta, K. S., Piscopo, K. M. & Mello, C. C. Robust Genome Editing

with Short Single-Stranded and Long, Partially Single-Stranded DNA Donors in

Caenorhabditis elegans. *Genetics* **210**, 781–787 (2018).

57.   Zhang, Y., Werling, U. & Edelmann, W. SLiCE: a novel bacterial cell extract-based

DNA cloning method. *Nucleic Acids Res.* **40**, e55–e55 (2012).

1161

1162   **Data availability statement**

1163   The data that support the findings of this study are available from the corresponding authors

1164   upon reasonable request. RNAseq data was deposited in the GEO database (GSE241786) and

1165   can   be   accessed   using   the   following   links:

1166   https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE241786

1167

Acknowledgements

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). We thank Henrik Bringmann and Lukas Kapitein

for plasmids; Peter Askjaer, Henrik Bringmann, and Antonio Miranda Vizuete for discussions

and comments on the manuscript; Zofia Olszewska, Marta Niklewicz, and Monika Woźniak

for assistance with worms maintenance. Work in the M. T. Laboratory was mainly funded by

a National Science Centre SONATA grant (2019/35/D/NZ3/04091) and additionally supported by a National Science Centre SONATA BIS grant (2021/42/E/NZ3/00358). Work in the W. P. Laboratory was funded by the Foundation for Polish Science co-financed by the European Union under the European Regional Development Fund (grant POIR.04.04.00-00-5EAB/18-00 to K.O., and W.P.), and additionally supported by the European Molecular Biology Organization (EMBO Installation Grant No. 3916 to K.O., and W.P.), and the Norwegian Financial Mechanism 2014-2021 operated by the Polish National Science Centre, Poland (project contract number 2019/34/H/NZ3/00691 to W.P.).

Author contributions

Conceptualization: MT, WP; Data curation: MT, WP; Formal analysis: AS, KO, KK, MT, WP, FCS, JY, AIT, LA; Funding acquisition: MT, WP; Investigation: AS, KO, KK, YI, AIT, LA, MT; Methodology: MT, WP, KO, AS, KK, FCS, YI; Project administration: MT, WP; Resources: MT, WP; Supervision: MT, WP; Validation: MT, WP, FCS, AS, KO, KK; Visualization: AS, KO, KK, MT; Writing – original draft: MT, WP; Writing – review & editing: MT, WP, FCS, AS, KO, KK.

Conflict of interest

The authors declare that they have no conflict of interest.