Effects of food supplementation and helminth removal on space use and spatial overlap

in wild rodent populations

Janine Mistrick[1], Jasmine S.M. Veitch[2], Shannon M. Kitchen[3], Samuel Clague[2], Brent C.

Newman[3], Richard J. Hall[4,5,6], Sarah A. Budischak[2], Kristian M. Forbes[3], Meggan E. Craft[1]

[1] Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN, USA

[2] W.M. Keck Science Department, Claremont McKenna, Pitzer, and Scripps Colleges,

Claremont, CA, USA

[3] Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[4] Odum School of Ecology, University of Georgia, Athens, GA, USA

[5] Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia,

Athens, GA, USA

[6] Center for the Ecology of Infectious Diseases, University of Georgia, Athens, GA, USA

Corresponding Author: Janine Mistrick mistr033@umn.edu

ABSTRACT

1. Animal space use and spatial overlap can have important consequences for population-level

processes such as social interactions and pathogen transmission. Identifying how

environmental variability and inter-individual variation affect spatial patterns and in turn influence

interactions in animal populations is a priority for the study of animal behavior and disease

ecology. Environmental food availability and macroparasite infection are common drivers of

variation, but there are few experimental studies investigating how they affect spatial patterns of

wildlife.

2. Bank voles (*Clethrionomys glareolus*) are a tractable study system to investigate spatial

patterns of wildlife and are amenable to experimental manipulations. We conducted a

replicated, factorial field experiment in which we provided supplementary food and removed

helminths in vole populations in natural forest habitat and monitored vole space use and spatial

overlap using capture-mark-recapture methods.

3. Using network analysis, we quantified vole space use and spatial overlap. We compared the

effects of food supplementation and helminth removal and investigated the impact of season,

sex, and reproductive status on space use and spatial overlap.

4. We found that food supplementation decreased vole space use while helminth removal

increased space use. Space use also varied by sex, reproductive status, and season. Spatial

overlap was similar between treatments despite up to three-fold differences in population size.

5. By quantifying the spatial effects of food availability and macroparasite infection on wildlife

populations, we demonstrate the potential for space use and population density to trade off and

maintain consistent spatial overlap in wildlife populations. This has important implications for

spatial processes in wildlife including pathogen transmission.

Keywords: anthelmintic treatment, bank vole, *Clethrionomys glareolus*, food supplementation,

network analysis, space use, spatial overlap

1 INTRODUCTION

How animals use space in their environment and interact with conspecifics is fundamental to

understanding proximity- and contact-driven processes such as social interactions (Kusch &

Lane, 2021) and pathogen transmission (Silk et al., 2018). Interactions with conspecifics are

frequently heterogeneous, reflecting individual variation due to sex and reproductive status and

associated social behavior or, alternatively, can be driven by environmental variability in habitat

quality or population density (Craft, 2015; Tompkins et al., 2011). Since animal space use and

conspecific interactions are crucial determinants of pathogen exposure, identifying how these

are shaped by individual-level variation and environmental variability has important implications

for mitigating effects of novel and zoonotic pathogens on wildlife, domestic animals, and

humans (Silk et al., 2017). However, studies that disentangle the relative contributions of

environmental and individual variation to interactions between animals are rare.

Food availability is a common source of environmental variability that can alter animal space

use and spatial overlap and influence pathogen transmission (Becker et al., 2015). At local

scales, low food availability may increase space use through increased foraging, while areas of

high food availability may decrease space use through increased site fidelity; heterogeneity in

food availability across the landscape may promote movement through increased attraction and

dispersal between patches (Becker et al., 2018). Point food sources (e.g., bird feeders, landfills,

crop fields) can increase spatial overlap of wildlife through aggregation and facilitate both direct

interactions between infected and susceptible animals (Altizer et al., 2018; Forbes et al., 2015)

and indirect interactions through environmentally transmitted pathogens accumulated in the

environment (Cross et al., 2007).

Individual variation among animals can also affect space use and spatial overlap with potential

consequences for pathogen transmission (VanderWaal & Ezenwa, 2016). For example, in

rodents, testosterone-mediated behavior in mature males can increase their contacts with

conspecifics and transmission potential (Grear et al., 2009) and is hypothesized to be a key

driver of commonly observed male bias in infection prevalence (Krasnov et al., 2012). Individual

variation may also arise through differing burdens of macroparasites such as helminths, with

potentially opposing effects on space use and spatial overlap. Helminth infection could increase

feeding behavior, requiring infected hosts to expand their foraging area (Brown et al., 1994).

Conversely, helminths may cause sickness-induced lethargy in infected animals and decrease

their activity, decreasing their space use and interactions with conspecifics (Ghai et al., 2015).

Further, since space use and spatial overlap can influence macroparasite exposure, positive

feedbacks between macroparasite exposure and space use could exacerbate or regulate within-

population variation in macroparasite burden and movement (Hawley et al., 2021).

Much of our knowledge of how food availability and macroparasite infection impact transmission

in wildlife comes from studies of laboratory animals under controlled conditions (Croft et al.,

2011) or observational studies in wildlife (Jolles et al., 2008) and therefore often lack explicit

investigation linking spatial behavior to transmission potential. In natural settings, food

availability and macroparasite infection frequently co-vary and could have balancing or

synergistic effects on animal space use and spatial overlap. For example, abundant food

resources could alleviate increased foraging requirements caused by helminth infection,

resulting in minimal effects of macroparasite burden on space use. Ultimately, rigorous field

experiments in wildlife are necessary to test the independent and synergistic effects of

environmental and individual variation on animal spatial patterns and implications for pathogen

transmission (e.g., Sweeny et al., 2021). Such research is pertinent and timely because land-

use change is altering patterns of food availability and macroparasite burdens that can

exacerbate pathogen emergence and transmission at the human-wildlife interface (Plowright et

al., 2021).

Our goal was to determine the effects of food supplementation and macroparasite removal on

space use and spatial overlap in wildlife populations. We used bank voles (*Clethrionomys*

glareolus, previously *Myodes glareolus*) as our study species as they are amenable to

experimental manipulation, are a common host for macroparasite infection (e.g., helminths like

Heligmosomoides glareoli can have a population-level prevalence of up to 80%; Haukisalmi &

Henttonen, 2000), and have well-documented natural history, particularly with respect to spatial

behavior (Tamarin et al., 1990). Reproductive males hold territories that are approximately

double those of females and that overlap with territories of males and females (reproductive

males average: 1800 m^2; females: 900 m^2; Bujalska & Grüm, 1989). Reproductive females are

territorial toward other reproductive females but may overlap more in the non-breeding season

(Bujalska, 1990). In both sexes, space use of non-reproductive voles is less than that of

reproductive voles (non-reproductive females, males average: 700 and 800 m^2, respectively;

Bujalska & Grüm, 1989). Both food availability and population density are thought to regulate

vole territory size (Bondrup-Nielsen & Karlsson, 1985).

We experimentally manipulated wild bank vole populations via food supplementation and

macroparasite (helminth) removal and monitored individual- and population-level responses.

Specifically, we 1) quantified space use by sex and reproductive status in the breeding and non-

breeding seasons, and 2) quantified spatial overlap between voles each month. Space use and

spatial overlap were compared between treatments to test the effects of food supplementation

and helminth removal on vole spatial patterns. As abundant food often increases small mammal

population density through immigration (Prevedello et al., 2017), we hypothesized that food

supplementation would decrease vole space use and increase spatial overlap. We hypothesized

that helminth infection would increase host movement (due to increased feeding behavior), and

thus helminth removal would decrease space use and spatial overlap.

2 MATERIALS AND METHODS

2.1 Study site and experimental design

We conducted a two-factor field experiment in boreal forests of southern Finland (61.0775°N,

25.0110°E) where bank voles are the dominant rodent species. These forests receive snowfall

from approximately November-April, and absence of snow cover makes monitoring and

experimental manipulation of voles most feasible from May-October. Twelve study sites were

established in old-growth spruce forest patches. All sites were at least two kilometers apart to

prevent dispersal between replicate populations (maximum dispersal distance estimated to be

500 m in heterogeneous habitat, Gliwicz & Ims, 2000). At each site, a standardized trapping grid

(100 m x 100 m, 1 hectare) was established to monitor the vole population, the grid consisting of

61 uniquely identified traps spaced 10 meters between grid rows and columns (Fig. 1).

Sites were randomly assigned to one of four treatment pairings ("treatments"): no manipulation

("unfed-control"; "U-C"); helminth removal only ("unfed-deworm"; "U-D"); food supplementation

only ("fed-control", "F-C"); and both food supplementation and helminth removal ("fed-deworm";

"F-D"). Each treatment was replicated at three sites (Fig. 1). Sites assigned to food

supplementation received a feed mix of mouse chow pellets and sunflower seeds evenly

distributed on the trapping grid every two weeks from May through November (if snowfall had

not yet accumulated). The supplemental feed mix was dosed at 7.54 kcal/m^2 (following Sweeny

et al., 2021) consisting of 7.875 kg of mouse chow pellets (Altromin 1324 10 mm pellets

maintenance diet for rats and mice [3227 kcal/kg]; Altromin Spezialfutter GmbH & Co. KG,

Lage, Germany) and 7.875 kg sunflower seeds (as fed, 6350 kcal/kg). Voles at deworm sites

were given an oral dose of 10 mg/kg Ivermectin and 100 mg/kg Pyrantel on their first capture in

a trapping occasion; this combination of medication is effective at treating larval and adult

helminth infections (Clerc et al., 2019). Voles at control sites received a matching weight-based

control dose *(*17.5% sucrose solution) on their first capture in a trapping occasion.

Figure 1. Field study design and experimental set-up. ("Site Locations") Twelve field sites were

established in southern Finland. ("Treatments") Sites were assigned one of four treatment

pairings: unfed-control, unfed-deworm, fed-control, and fed-deworm and each treatment was

replicated at three sites each. ("Trapping Grid Arrangement") The trapping grid included 61

traps in 11 rows and 11 columns. Rows and columns were spaced 10 meters apart and traps

(dots) were placed in an offset arrangement with 20 meters to the next trap in the same row and

column and 14.14 meters to the next trap on the diagonal.

2.2 Data collection

Vole populations were longitudinally monitored via capture-mark-recapture methods. Every four

weeks from May-October 2021, Ugglan multi-capture live traps (Grahnab, Sweden) were baited

with whole oats and set for 48 hours (a "trapping occasion"). Traps were checked each morning

and evening at approximately 0600 and 1600 (four checks per trapping occasion) and captured

animals were processed and then released at the capture location. Six trapping occasions were

conducted at each site during the study period (total trap events: 17,568). Upon first capture, all

voles were injected with a Passive Integrated Transponder ("PIT tag"; "Skinny" PIT Tag, Oregon

RFID, USA) for unique identification. The trap number, PIT tag number, sex, and reproductive

status were recorded and a fecal sample was collected (to quantify helminth presence and

infection intensity via fecal egg counts) for each vole on their first capture each trapping

occasion. If a vole was recaptured in a trapping occasion, only the trap number and PIT tag

number were recorded, no samples were collected, and the vole was released at the capture

location.

Each capture, voles were categorized as reproductive (perforate vagina, lactating, or pregnant

females; males with scrotal testes) or non-reproductive. Reproductive status per vole was

summarized within the summer breeding season (June-August) and autumn non-breeding

season (September-October in southern Finland; Kaikusalo, 1972) to account for seasonal

differences in vole behavior. If a vole was ever recorded as reproductive during a trapping

occasion within the season, it was recorded as reproductive in that season. In autumn, voles

that were reproductive in summer were also considered reproductive in autumn, even if external

traits were not observed in September or October. Voles were further classified into four

functional groups: reproductive males, reproductive females, non-reproductive males, and non-

reproductive females. Functional groups are population subgroups that are relatively uniform in

their behavior, physiology, and immunology. Conducting epidemiological analysis at this level

pools individuals into biologically meaningful groups and enables more realistic conclusions to

be drawn (Henttonen, 2022).

All trapping, handling, and sampling of wild bank voles was conducted under approval of the

University of Arkansas Institutional Animal Care and Use Committee (IACUC #19105) and the

Finnish Animal Ethics Board (ESAVI-17810-2019). Access to forest sites was provided by

private landowners and by Metsähallitus Metsätalous Oy (MH 6302/2019).

2.3 Data analysis

The efficacy of the deworm treatment was assessed using helminth infection status and

intensity in infected animals (eggs per gram of feces [EPG]), quantified using a salt flotation

method (Pritchard & Kruse, 1982). Helminth infection status was modeled for all sampled

animals using a generalized linear mixed-effects model (binomial family, logit link; "lme4" R

package, Bates et al., 2015). Helminth intensity (natural log-transformed EPG) was modeled for

infected animals using a linear mixed-effects model (lmerTest R package, Kuznetsova et al.,

2017). Both models were parameterized with fixed effects of: treatment group (deworm/control),

treatment stage (pre-treatment [the first capture of a vole] / post-treatment [all subsequent

captures]), and the interaction of treatment group and stage. Vole ID was included as a random

effect. Model interactions were visualized using the 'visreg' R package (Breheny & Burchett,

2017).

To understand general patterns of vole space use and how they were influenced by food

supplementation and helminth removal, the mean space use of voles in each functional group

was characterized in both the breeding and non-breeding seasons for each treatment. Within a

single site, overlapping space use between pairs of voles ("pairwise space-use overlap") was

used to approximate opportunities for direct and indirect interactions (Robert et al., 2012). The

total pairwise space-use overlap of a focal vole with all its neighbors ("individual spatial overlap")

was quantified to investigate how opportunities for interactions varied among individuals. When

interactions like spatial overlap are heterogeneous among individuals, network analysis is a

useful method to visualize their frequency and distribution within a population (Croft et al., 2008;

Krause et al., 2007). We constructed spatial overlap networks to visualize all the pairwise

space-use overlap between voles at each site to investigate how the experimental

manipulations affected opportunities for direct and indirect interactions at the population level.

Vole capture numbers in May were very low (zero to five animals per site), prohibiting network

construction for all sites, therefore May capture data were excluded from downstream analysis.

Additionally, since space use was estimated by functional group (the combination of sex and

reproductive status), only animals with both sex and reproductive status data recorded were

included in the analysis.

Data analysis was conducted in program R version 4.1.2 (R Core Team, 2021).

2.3.1 Factors influencing seasonal space use

Seasonal vole space use was approximated within a season by aggregating capture location

data among individuals by population subgroups to characterize the average space use of an

individual in each group (Wanelik & Farine, 2022). This approach allows space use and spatial

overlap to be estimated for individuals, even if they are caught only once or only in one trap and

helps minimize uncertainty in space-use estimates created by the relatively short observation

periods and limited observations per individual. As such, these methods offer a robust way to

detect biological effects of shared space use when using sparse capture-mark-recapture data

where repeat observations per individual are limited or heterogeneous (Wanelik & Farine,

2022).

Using this approach, we aggregated vole capture locations from each trapping occasion June-

October (hereafter "month") into a summer breeding season (June-August) and an autumn non-

breeding season (September-October). We defined a vole's seasonal "centroid" as the weighted

average of all locations where it was trapped that season, allowing multiple captures in the

same trap to more strongly influence the centroid than a single capture in a trap. For each vole,

the distance from this centroid to every trap in the trapping grid (n=61) was calculated. Then,

distances for all voles (across the entire study) were pooled by season.

We used generalized linear models (GLMs) to characterize the probability of capturing a vole in

a trap a given distance from its seasonal centroid and investigate factors influencing space use

(i.e., sex, reproductive status, treatment), separately in summer and autumn. In each model, the

response variable was whether a vole was caught in a given trap (where 1 indicated it was

caught, 0 indicated it was not) and the explanatory variables were: the natural log distance of

the trap from the vole's seasonal centroid, vole sex, reproductive status (reproductive/non-

reproductive), site-level food treatment (unfed/fed), helminth treatment (control/deworm), and

interaction effects between distance and each of: sex, reproductive status, food treatment, and

helminth treatment, and a three-way interaction between distance, food treatment, and helminth

treatment. The interaction terms were examined to identify parameters that significantly affected

the relationship between distance from the seasonal centroid and capture probability, indicating

differences in space use. Model interactions were visualized using the 'visreg' R package.

2.3.2 Seasonal space use

We then used the significant predictors of capture probability as identified by the seasonal

space use GLMs to divide voles into population subgroups and characterize space use

separately for each group. The distances from a vole's seasonal centroid to every trap in the

trapping grid were pooled across voles, grouped by the combination of season*sex*reproductive

status*food treatment*helminth treatment (n groups = 32). For each group, we fitted a negative

sigmoidal curve (Equation 1; fitted using a Bernoulli GLM, where 1 indicated an animal was

caught in a given trap, 0 indicated it was not; as developed by Wanelik & Farine, 2022) to

characterize the average space use of a vole in that group:

$$P(d) \; = \; \frac{1}{1 + e^{-a-bd}}$$ (Eqn. 1)

where space use was defined as the declining probability P of capturing a vole an increasing

distance (d) from its seasonal centroid as determined by: a (describing the size of the space

use radius), b (describing how steeply the capture probability declines as distance increases),

and d (the logarithmic distance from the centroid). Thus, space use was separately

characterized in the summer and autumn for each treatment for voles of each functional group.

2.3.2 Pairwise space-use overlap and spatial overlap networks

Space-use overlap was estimated between all pairs of voles captured at a site in a given month.

These pairwise interactions were then used to create spatial overlap networks to visualize all

the interactions in the observed population at each site, in each month. First, the monthly

centroid for every vole was calculated as the weighted average of all trap locations where it was

captured that month (1-4 captures/month possible). The appropriate seasonal space use for a

given vole's sex, reproductive status, the site-level food and helminth treatments, and the

season (as calculated above using Eqn. 1) was centered on this monthly centroid.

Overlap between all pairs of voles observed in a month at a site was estimated based on the

overlap in the two voles' space use (as developed by Wanelik & Farine, 2022). Specifically, we

calculated the amount of pairwise space-use overlap by first predicting the (independent)

probabilities of capturing vole 1 and vole 2 (using Eqn. 1) at each grid point (x, y) in a grid of

coordinates X and Y overlapping both of their space use. The amount of pairwise space-use

overlap between the two voles $(E_{1,2})$ was estimated by taking the lower of the two voles' capture

probabilities at each grid point (x, y) and summing them across all grid points and similarly

summing the higher of the two capture probabilities across all grid points. The summed

minimums were then divided by the summed maximums to get a ratio indicating how similar the

two voles' capture probabilities were across space (Equation 2, developed by Wanelik & Farine,

2022):

$$E_{1,2} = \frac{\Sigma_X \, \Sigma_Y \, min \, (P_1(\sqrt{(x_1-x)^2+(y_1-y)^2}), P_2 \, (\sqrt{(x_2-x)^2+(y_2-y)^2})}{\Sigma_X \, \Sigma_Y \, max \, (P_1(\sqrt{(x_1-x)^2+(y_1-y)^2}), P_2(\sqrt{(x_2-x)^2+(y_2-y)^2})}$$ (Eqn. 2)

The more similar the space use of the two voles, the ratio ($E_{1,2}$) approaches 1 (the two voles

have similar capture probabilities at each grid point, indicating more complete overlap), whereas

the more different their space use, the ratio approaches 0 (where one vole has high capture

probability, the other has low capture probability, indicating very little overlap).

We then constructed spatial overlap networks visualizing the pairwise space-use overlap

between all observed voles at each site using the 'igraph' R package (Csardi & Nepusz, 2006).

Separate networks were constructed for each month June-October to capture the temporal

dynamics of network structure and account for conspecific interactions that vary through time

due to changing population size, demographics, or seasonal behaviors (Pinter-Wollman et al.,

2014). In the networks, voles were represented as nodes and only voles captured in a given

month were included in the network. Connections between nodes (edges) were undirected and

were weighted by the value of pairwise space-use overlap between two voles (the edge weight

between voles 1 and 2, $E_{1,2}$) with no thresholding of edge weights.

2.3.3 Individual spatial overlap

We quantified individual spatial overlap within the vole populations using weighted degree.

Biologically, weighted degree represents the sum of all pairwise space-use overlap between a

focal vole and its neighbors ("individual spatial overlap"). Values of weighted degree were

unfiltered, allowing for very small measures of weighted degree indicating a very low probability

of overlap (e.g., weighted degree <0.01). Two additional measures of spatial overlap:

unweighted degree and normalized unweighted degree, are presented in the Supplement:

Individual spatial overlap - Additional metrics (Table S1; Fig. S1, Fig. S2). All network metrics

were calculated using the 'igraph' R package (Csardi & Nepusz, 2006).

Network data are inherently non-independent and degree is influenced by network size (total

number of nodes in the network), limiting the ability to make direct comparisons between

networks of varying size (Farine & Whitehead, 2015). Network size varied between sites and

across months based on the number of voles observed during a trapping occasion. For an

analysis of the effects of treatment on network size, see Supplement: Network size (Table S2).

3 RESULTS

We captured voles and recorded sex and reproductive status data for 742 unique individuals

during our study period (June-October 2021). Of these, 445 voles (60%) were captured at least

twice (mean ± SD captures per vole: 2.75 ± 2.33 [range: 1-17 captures]). We documented 905

captures across the 12 sites in summer (June-August) and 1107 captures in autumn

(September-October). These capture data were used to inform our seasonal space use models.

Fecal egg count data was collected for 1035 captures during the study period. Of these,

helminth eggs were detected in 440 samples (42.5%). Treatment group (control/deworm) had a

significant effect on the relationship between pre- and post-treatment levels of helminth

prevalence (odds ratio=0.51, p=0.022; Table S3) and infection intensity (β=-0.74, p=0.005;

Table S4). Helminth prevalence and the infection intensity were similar pre-treatment and both

were lower in treated voles compared to control voles post-treatment. In dewormed voles,

helminth prevalence remained similar from pre-treatment to post-treatment whereas helminth

prevalence increased in control voles (Fig. S3A). Helminth intensity decreased in dewormed

voles post-treatment while intensity was similar pre- and post-treatment in control voles (Fig.

S3B).

3.1 Seasonal space use

In summer, the seasonal space use model indicated that the effect of distance on capture

probability differed by vole sex, reproductive status, and site-level food supplementation

treatment (GLM interaction terms between each parameter and distance from centroid: all

$p < 0.001$; Table S5; Fig. S4A-C). No effect of helminth removal was detected (GLM interaction

term between helminth treatment and distance from centroid: $p = 0.18$, Table S5; Fig. S4D).

When we estimated summer space use (using Eqn. 1) for each functional group in each

treatment, we found evidence for male voles having greater space use than females and

reproductive voles having greater space use than non-reproductive voles (Fig. 2). Food

supplementation generally decreased vole space use, compared to unfed treatments (Fig. 2).

Figure 2. Space use of bank voles by sex and reproductive status across treatments in the summer breeding season (June-August) and autumn nonbreeding season (September-October). Circle size represents the mean space use area (m^2) where the probability of capture is >0.01.

In autumn, the seasonal space use model indicated that the effect of distance on capture probability differed by vole reproductive status, site-level food treatment, and helminth treatment (GLM interaction terms between each parameter and distance from centroid: all p<0.012, Table S6; Fig. S5B-D). No effect of vole sex was detected in autumn (GLM interaction term between sex and distance from centroid p=0.68, Table S6; Fig. S5A).

When estimating space use, we found that reproductive voles had greater space use than non-reproductive voles, though space use of reproductive and non-reproductive voles was more

similar in autumn than in summer (Fig. 2). The space use of reproductive voles generally

decreased or remained similar from summer to autumn. The space use of non-reproductive

voles increased from summer to autumn. Food supplementation decreased vole space use

compared to unfed treatments (Fig. 2). Helminth removal had the opposite effect: voles in

deworm treatments had increased space use, compared to voles in control treatments (Fig. 2).

3.2 Spatial overlap networks

In every month, all voles overlapped with at least one other vole and populations appeared

densely and homogeneously connected if all edges were included in the network (i.e., no edge

weight thresholding). However, edges of low weight (<0.05) were common and more

heterogeneous network structure appeared with minimal edge weight thresholding (Fig. 3).

Network size was highly variable across months and treatments (range: 2-52 voles). Networks

were smallest in June, peaked in size in August (unfed treatments) or September (fed

treatments), and then decreased slightly though October (Fig. 4A; see also Supplement:

Network Size).

Figure 3. Spatial overlap networks of vole pairwise space-use overlap. Nodes represent individual voles, edges represent pairwise space-use overlap, thicker edges indicate greater space-use overlap between voles. For each treatment, networks from one site are shown. Edge weights are thresholded to a minimum edge weight of 0.05 for visualization purposes.

Mean weighted degree, indicating individual vole spatial overlap (total spatial overlap of a focal vole with its neighbors, weighted by the amount of pairwise space-use overlap with each neighbor) was similar between the unfed-control, fed-control, and fed-deworm treatments in a given month in both summer and autumn. The range of weighted degree values were widest in September when the most voles with high degree (weighted degree >3) were observed (Fig. 4B). Compared to the other treatments, mean weighted degree in the unfed-deworm treatment was smaller with a narrower range in June-September (Fig. 4B). In summer (June-August), mean weighted degree was generally greater for reproductive male voles than reproductive females. In autumn (September-October), weighted degree was more similar between

reproductive male and female voles (Fig. 4B). Similar trends were observed for non-

reproductive voles.

Figure 4. Population-level network size and individual vole spatial overlap. (A) Network size

(number of voles) by month. One data point shown per replicate site. (B) Weighted degree (total

pairwise space-use overlap per vole, weighted by the amount of overlap with each of its

neighbors) by month. Means shown as points and lines indicate the range of values for males

(blue) and females (pink) in each treatment. Data shown for reproductive voles only, trends for

non-reproductive voles were similar.

4 DISCUSSION

Variation in food availability and macroparasite infection can alter animal space use and

intraspecific interactions in ways that influence spatial processes in wildlife. Through

experimental food supplementation and helminth removal in wild bank vole populations, we

quantified the effects of environmental and individual variation on space use and spatial overlap.

Food supplementation decreased vole space use, while helminth removal increased vole space

use, but only in autumn. Space use was also influenced by functional group (combination of sex

and reproductive status) and season. Mean individual vole spatial overlap was similar between

treatments in a given month, despite up to three-fold differences in population density. Our

findings show that both environmental variability and individual variation influence animal space

use, but that space use and population density can trade off and maintain consistent spatial

overlap in wildlife populations under different environmental conditions.

We detected significant effects of both food supplementation and deworming treatment on vole

seasonal space use, though the effect sizes were generally small. The experimental and

replicated design of our study and the large seasonal sample size of captured animals likely

enables the detection of subtle and potentially important differences in space use. Food

supplementation decreased vole space use in both summer and autumn. These findings align

with the general understanding that supplemental food decreases bank vole home range size

(Bondrup-Nielsen & Karlsson, 1985; Mazurkiewicz, 1983). However, food supplementation also

increased population density, introducing a confounding factor which cannot be fully

disentangled from the observed effects on space use. Increased population density can

increase territorial behavior, resulting in smaller, more easily defended home ranges -

potentially independent of food availability (Davis et al., 2015). Decreased space use under food

supplementation is likely a complex response driven by interacting effects of food availability

and population density.

In autumn, deworming treatment increased vole space use as compared to voles with

unmanipulated helminth infection. This effect was not observed in summer, suggesting a time-

lagged effect of deworming treatment. This may be due to individual-level factors; it takes time

for the medication to start working and for behavioral effects of helminth infection to be

impacted, so treatment in one month would not be observed to impact behavior until the

following month. Moreover, the deworming treatment was only applied to captured voles and

thus was heterogeneously applied across the population. As such, several months of trapping

may have been necessary to treat a sufficient proportion of individuals and observe population-

level effects (Pedersen & Fenton, 2015) or repeated treatments per individual were necessary

to decrease the environmental burden and limit infection (Knowles et al., 2013). Increased

space use by voles in the deworm treatments lends support to the hypothesis that animals with

high helminth burdens exhibit sickness behavior and decrease their movements (Ghai et al.,

2015). One potential confounding factor is that population density was very low across the

unfed-deworm treatment, thus the increased space use may be a result of minimal territorial

constraints and not a direct result of helminth removal. However, population sizes were more

similar between the fed treatments (and, in some cases, populations were larger in fed-deworm

than fed-control sites) and space use was still greater with deworming treatment, supporting

helminth removal being a biologically meaningful factor influencing vole space use.

Space use interacted with vole sex and reproductive status as expected based on vole biology.

We found that male space use was greater than that of females during the breeding season and

space use of reproductive voles was greater than that of non-reproductive voles, which is

consistent with the established literature (Bondrup-Nielsen & Karlsson, 1985; Bujalska, 1990).

However, while we are confident in our space use estimates for reproductive voles, the space

use of non-reproductive voles may be more difficult to estimate, particularly in summer. On

average in the summer months, non-reproductive voles were captured less frequently and in

fewer traps than reproductive voles (whereas capture frequency and trap usage were similar in

autumn). We also observed very small space use by non-reproductive voles in summer. This

could be representative of their true space use (e.g., a 'sit and wait' strategy, Bujalska, 1990),

but may also be the result of many young voles who are captured once and then disperse off

the trapping grid and are not captured again, artificially 'shrinking' the observed space use of

non-reproductive voles. While we can not disentangle the relative impacts of these two factors,

our observations still provide valuable comparisons of reproductive vole space use across

seasons and environmental and individual variation by food supplementation and helminth

removal.

It was surprising to us that, despite the differences in space use and population density, we did

not observe clear differences in individual spatial overlap by treatment, even when population

sizes differed. We found that food supplementation increased population density, consistent

with previous manipulations in wild vole populations (Bujalska & Janion, 1981). It has been

hypothesized that territorial behavior among reproductive females should be weakest when food

is abundant and evenly distributed and population density is high (Ostfeld, 1985). Previous

studies in *Clethrionomys* voles support this hypothesis, demonstrating that food addition

decreases reproductive female space use, but increases spatial overlap between females

(Andrzejewski & Mazurkiewicz, 1976; Ims, 1987). We observed decreases in both male and

female space use with food supplementation, but did not observe an increase in spatial overlap

in fed treatments compared to unfed. Similar spatial overlap despite differences in population

density could be explained by a non-linear interaction of space use with population density.

Individuals may move more freely and have larger space use at lower densities (unfed

treatments), but constrain their space use and only contact their nearest neighbors at high

densities (fed treatments; e.g., Davis et al., 2015). Thus, it is possible the effects of food

supplementation on space use and population density may trade off, resulting in similar patterns

of spatial overlap between fed and unfed populations, despite differences in population density.

Our network construction methods were chosen to minimize the effect of infrequent or missing

observations of individuals on network structure. Networks in this study were constructed from

48 hours of observation with a possible maximum of four observations per vole and thus are not

a complete representation of monthly space use. Networks constructed from trapping data often

assign edges between individuals caught in the same or adjacent traps (e.g., Grear et al., 2009;

Perkins et al., 2009), which can bias network measures if capture frequency is heterogeneous.

We combined capture data across individuals and across months to define an average seasonal

space use for each functional group. In this way, the behavior of animals captured many times

fills in the 'missing' data for animals captured only once or twice and minimizes the effect of low

capture frequency on network structure (Wanelik & Farine, 2022). Further, by using overlapping

space-use to define connections in the spatial overlap networks, spatial overlap between voles

is also less dependent on their specific trapped locations. These generalized approaches can

help to minimize the uncertainty associated with network measures when social behavior is

inferred using snapshot observations of individual behavior. One drawback is that aggregating

behavior across functional groups eliminates individual variation within the group. Our objective

was to quantify and compare space use between treatments and populations. By grouping voles

by functional group, we allow behavior to vary across biologically meaningful axes such as sex

and reproductive status (Henttonen, 2022), while capturing the average behavior of each

population subclass which can be more readily generalized to other animal systems.

Spatial overlap can be linked to many direct and indirect interactions between animals including

social interactions and pathogen transmission. When direct observation of interactions between

animals is limited or impossible, thoughtfully aligned measures of spatial overlap can be used to

approximate these interactions and predict transmission opportunities (Craft & Caillaud, 2011;

Mistrick et al., 2022). However, spatial overlap and transmission opportunities may not correlate

across all pathogen types. For instance, direct transmission can be decoupled from spatial

overlap if animals seek out or avoid interactions with conspecifics. Thus, spatial overlap may be

best used to approximate indirect transmission opportunities via the environment where direct

contact between animals is not required (Godfrey et al., 2010). In that context, our finding that

spatial overlap was similar across treatments could suggest that opportunities for environmental

transmission may remain relatively constant even when population size increases–such as

under high resource availability. However, differences in space use by sex and reproductive

status could bias functional groups such as mature males toward greater exposure to

environmental pathogens or greater potential to shed infectious material over larger areas

(Krasnov et al., 2012). As such, the utilities of space use and spatial overlap data can be

extended to various fields, providing insight into animal behavior and its ecological

consequences, even when it cannot be measured directly.

5 CONCLUSIONS

Field studies investigating the effects of environmental and individual variation on animal spatial

patterns are limited, thus this study provides necessary, empirical evidence of the effects of food

supplementation and helminth removal on space use and spatial overlap in wildlife populations.

We demonstrate that environmental variability due to supplemental food and individual variation

in macroparasite infection help shape rodent space use. We also identify the potential for space

use and population size to trade off, maintaining similar levels of spatial overlap between

individuals despite variation in their environment. As such, our research demonstrates that

environmental variability and individual variation drive animal spatial behavior and interact with

population dynamics in wildlife populations and therefore represent important drivers of

population-level contact- and proximity-driven processes such as social interactions and

pathogen transmission.

ACKNOWLEDGEMENTS

This research was funded by the National Science Foundation (DEB-1911925). J.M. was

supported by the National Science Foundation Graduate Research Fellowship Program (award

no. 2237827). Any opinions, findings, and conclusions or recommendations expressed in this

material are those of the authors and do not necessarily reflect the views of the National

Science Foundation. J.M. was also supported by the Dayton Fund of the Bell Museum and the

University of Minnesota Ecology, Evolution, and Behavior Graduate Program and the College of

Biological Sciences.

We are grateful to our collaborators at the Lammi Biological Station and University of Helsinki:

John Loehr, Janne Sundell, Esa-Pekka Tuominen, Tiina Tulonen, Matti Kotakorpi, Riitta Ilola,

Jaakko Vainionpää, Tomas Strandin, and Tarja Sironen, and to the students who helped collect

and process the field data: Alexis Beagle, Muriel Chaudhri, Stephanie Du, Lucie Fornilli,

Mathilde Gaudillère, Teemu Lemola, Nathaniel Mull, and Anni Simonen. Thanks to Vole Fever

project collaborators Clay Cressler, Brendan Haile, and Lexi Beagle as well as Sharon Jansa

and Susan Jones for providing feedback on this work.

Figures were created with BioRender.com. Fig. 1 site map by Anthony Shing.

AUTHOR CONTRIBUTIONS

Kristian Forbes, Sarah Budischak, Meggan Craft, and Richard Hall conceived the ideas,

designed the methodology, and secured the funding; Shannon Kitchen, Janine Mistrick,

Jasmine Veitch, Samuel Clague, Brent Newman, Kristian Forbes, and Sarah Budischak

collected the data; Janine Mistrick analyzed the data, created visualizations, and led the writing

of the manuscript under the mentorship of Meggan Craft. All authors contributed critically to the

drafts and gave final approval for publication.

DATA AVAILABILITY STATEMENT

Data will be made available through the Dryad Digital Repository upon manuscript acceptance.

Code is available in a GitHub repository (github.com/jmistrick/vole-spatial-v2) which will be

archived on Zenodo upon manuscript acceptance.

REFERENCES

Altizer, S., Becker, D. J., Epstein, J. H., Forbes, K. M., Gillespie, T. R., Hall, R. J., Hawley, D.

M., Hernandez, S. M., Martin, L. B., Plowright, R. K., Satterfield, D. A., & Streicker, D. G.

(2018). Food for contagion: Synthesis and future directions for studying host–parasite

responses to resource shifts in anthropogenic environments. *Philosophical Transactions*

of the Royal Society B: Biological Sciences, *373*(1745), 20170102.

https://doi.org/10.1098/rstb.2017.0102

Andrzejewski, R., & Mazurkiewicz, M. (1976). Abundance of food supply and size of the bank

vole's home range. *Acta Theriologica*, *21*(17), 237–253.

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R., Singmann, H., Dai, B.,

Grothendieck, G., Green, P., & Bolker, M. (2015). Package "lme4." *Convergence*, *12*(1),

2.

Becker, D. J., Snedden, C. E., Altizer, S., & Hall, R. J. (2018). Host Dispersal Responses to

Resource Supplementation Determine Pathogen Spread in Wildlife Metapopulations.

The American Naturalist, *192*(4), 503–517. https://doi.org/10.1086/699477

Becker, D. J., Streicker, D. G., & Altizer, S. (2015). Linking anthropogenic resources to wildlife–

pathogen dynamics: A review and meta-analysis. *Ecology Letters*, *18*(5), 483–495.

https://doi.org/10.1111/ele.12428

Bondrup-Nielsen, S., & Karlsson, F. (1985). Movements and spatial patterns in populations of

Clethrionomys species: A review. *Annales Zoologici Fennici*, *22*(3), 385–392. JSTOR.

Breheny, P., & Burchett, W. (2017). Visualization of Regression Models Using visreg. *The R*

Journal, *9*, 56–71.

Brown, E. D., Macdonald, D. W., Tewand, T. E., & Todd, I. A. (1994). *Apodemus sylvaticus*

infected with *Heligmosomoides polygyrus* (Nematoda) in an arable ecosystem:

Epidemiology and effects of infection on the movements of male mice. *Journal of*

Zoology, *234*(4), 623–640. https://doi.org/10.1111/j.1469-7998.1994.tb04869.x

Bujalska, G. (1990). Social System of the Bank Vole, *Clethrionomys Glareolus*. In R. H.

Tamarin, R. S. Ostfeld, S. R. Pugh, & G. Bujalska (Eds.), *Social Systems and Population*

Cycles in Voles (pp. 155–167). Birkhäuser. https://doi.org/10.1007/978-3-0348-6416-

9_15

Bujalska, G., & Grüm, L. (1989). Social Organization of the Bank Vole (*Clethrionomys glareolus*,

Schreber 1780) and Its Demographic Consequences: A Model. *Oecologia*, *80*(1), 70–81.

Bujalska, G., & Janion, S. M. (1981). Bank vole response to an increase in environmental

capacity. *Bulletin of the Academy of Polish Sciences, Ser. Sci. Biol.*, *29*, 129–133.

Clerc, M., Babayan, S. A., Fenton, A., & Pedersen, A. B. (2019). Age affects antibody levels and

anthelmintic treatment efficacy in a wild rodent. *International Journal for Parasitology:*

Parasites and Wildlife, *8*, 240–247. https://doi.org/10.1016/j.ijppaw.2019.03.004

Craft, M. E. (2015). Infectious disease transmission and contact networks in wildlife and

livestock. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

370(1669), 20140107. https://doi.org/10.1098/rstb.2014.0107

Craft, M. E., & Caillaud, D. (2011). Network Models: An Underutilized Tool in Wildlife

Epidemiology? [Review Article]. *Interdisciplinary Perspectives on Infectious Diseases*;

Hindawi. https://doi.org/10.1155/2011/676949

Croft, D. P., Edenbrow, M., Darden, S. K., Ramnarine, I. W., van Oosterhout, C., & Cable, J.

(2011). Effect of gyrodactylid ectoparasites on host behaviour and social network

structure in guppies *Poecilia reticulata*. *Behavioral Ecology and Sociobiology*, *65*(12),

2219–2227. https://doi.org/10.1007/s00265-011-1230-2

Croft, D. P., James, R., & Krause, J. (2008). *Exploring Animal Social Networks*. Princeton

University Press.

Cross, P. C., Edwards, W. H., Scurlock, B. M., Maichak, E. J., & Rogerson, J. D. (2007). Effects

of Management and Climate on Elk Brucellosis in the Greater Yellowstone Ecosystem.

Ecological Applications, *17*(4), 957–964. https://doi.org/10.1890/06-1603

Csardi, G., & Nepusz, T. (2006). The Igraph Software Package for Complex Network Research.

InterJournal Complex Systems, *1695*(5), 1–9.

Davis, S., Abbasi, B., Shah, S., Telfer, S., & Begon, M. (2015). Spatial analyses of wildlife

contact networks. *Journal of The Royal Society Interface*, *12*(102), 20141004.

https://doi.org/10.1098/rsif.2014.1004

Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting animal social

network analysis. *Journal of Animal Ecology*, *84*(5), 1144–1163.

https://doi.org/10.1111/1365-2656.12418

Forbes, K. M., Henttonen, H., Hirvelä-Koski, V., Kipar, A., Mappes, T., Stuart, P., & Huitu, O.

(2015). Food provisioning alters infection dynamics in populations of a wild rodent.

Proceedings of the Royal Society B: Biological Sciences, *282*(1816), 20151939.

https://doi.org/10.1098/rspb.2015.1939

Ghai, R. R., Fugère, V., Chapman, C. A., Goldberg, T. L., & Davies, T. J. (2015). Sickness

behaviour associated with non-lethal infections in wild primates. *Proceedings of the*

Royal Society B: Biological Sciences, *282*(1814), 20151436.

    https://doi.org/10.1098/rspb.2015.1436

Gliwicz, J., & Ims, R. A. (2000). Dispersal in the bank vole. *Polish Journal of Ecology*, 51-61.

    Suppl 48.

Godfrey, S. S., Moore, J. A., Nelson, N. J., & Bull, C. M. (2010). Social network structure and

    parasite infection patterns in a territorial reptile, the tuatara (*Sphenodon punctatus*).

    International Journal for Parasitology, *40*(13), 1575–1585.

    https://doi.org/10.1016/j.ijpara.2010.06.002

Grear, D. A., Perkins, S. E., & Hudson, P. J. (2009). Does elevated testosterone result in

    increased exposure and transmission of parasites? *Ecology Letters*, *12*(6), 528–537.

    https://doi.org/10.1111/j.1461-0248.2009.01306.x

Haukisalmi, V., & Henttonen, H. (2000). Variability of helminth assemblages and populations in

    the bank vole *Clethrionomys glareolus*. *Polish Journal of Ecology*, *48*, 219–231.

Hawley, D. M., Gibson, A. K., Townsend, A. K., Craft, M. E., & Stephenson, J. F. (2021).

    Bidirectional interactions between host social behaviour and parasites arise through

    ecological and evolutionary processes. *Parasitology*, *148*(3), 274–288.

    https://doi.org/10.1017/S0031182020002048

Henttonen, H. (2022). Importance of demography in understanding disease ecology in small

    mammals. *THERYA*, *13*(1), Article 1.

Ims, R. A. (1987). Responses in Spatial Organization and Behaviour to Manipulations of the

    Food Resource in the Vole *Clethrionomys rufocanus*. *Journal of Animal Ecology*, *56*(2),

    585–596. https://doi.org/10.2307/5070

Jolles, A. E., Ezenwa, V. O., Etienne, R. S., Turner, W. C., & Olff, H. (2008). Interactions

    Between Macroparasites and Microparasites Drive Infection Patterns in Free-Ranging

    African Buffalo. *Ecology*, *89*(8), 2239–2250. https://doi.org/10.1890/07-0995.1

Kaikusalo, A. (1972). Population turnover and wintering of the bank vole, *Clethrionomys*

glareolus (Schreb.), in southern and central Finland. *Annales Zoologici Fennici*, *9*(4),

219–224.

Knowles, S. C. L., Fenton, A., Petchey, O. L., Jones, T. R., Barber, R., & Pedersen, A. B.

(2013). Stability of within-host–parasite communities in a wild mammal system.

Proceedings of the Royal Society B: Biological Sciences, *280*(1762), 20130598.

https://doi.org/10.1098/rspb.2013.0598

Krasnov, B. R., Bordes, F., Khokhlova, I. S., & Morand, S. (2012). *Gender-biased parasitism in*

small mammals: Patterns, mechanisms, consequences. 76(1), 1–13.

https://doi.org/10.1515/mammalia-2011-0108

Krause, J., Croft, D. P., & James, R. (2007). Social network theory in the behavioural sciences:

Potential applications. *Behavioral Ecology and Sociobiology*, *62*(1), 15–27.

https://doi.org/10.1007/s00265-007-0445-8

Kusch, J. M., & Lane, J. E. (2021). Determinants of social structure in a northern population of

black-tailed prairie dogs, *Cynomys ludovicianus. Animal Behaviour*, *178*, 1–10.

https://doi.org/10.1016/j.anbehav.2021.05.017

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Mazurkiewicz, M. (1983). Spatial organization of the population. *Acta Theriologica*, *28*(Suppl. 1),

103–144. https://doi.org/10.4098/AT.arch.83-48

Mistrick, J., Gilbertson, M. L. J., White, L. A., & Craft, M. E. (2022). Constructing animal

networks for parasite transmission inference. In V. O. Ezenwa, S. Altizer, & R. J. Hall

(Eds.), *Animal Behavior and Parasitism*. Oxford University Press.

Ostfeld, R. S. (1985). Limiting Resources and Territoriality in Microtine Rodents. *The American*

Naturalist, *126*(1), 1–15. https://doi.org/10.1086/284391

Pedersen, A. B., & Fenton, A. (2015). The role of antiparasite treatment experiments in

assessing the impact of parasites on wildlife. *Trends in Parasitology*, *31*(5), 200–211.

https://doi.org/10.1016/j.pt.2015.02.004

Perkins, S. E., Cagnacci, F., Stradiotto, A., Arnoldi, D., & Hudson, P. J. (2009). Comparison of

social networks derived from ecological data: Implications for inferring infectious disease

dynamics. *The Journal of Animal Ecology*, *78*(5), 1015–1022.

https://doi.org/10.1111/j.1365-2656.2009.01557.x

Pinter-Wollman, N., Hobson, E. A., Smith, J. E., Edelman, A. J., Shizuka, D., de Silva, S.,

Waters, J. S., Prager, S. D., Sasaki, T., Wittemyer, G., Fewell, J., & McDonald, D. B.

(2014). The dynamics of animal social networks: Analytical, conceptual, and theoretical

advances. *Behavioral Ecology*, *25*(2), 242–255. https://doi.org/10.1093/beheco/art047

Plowright, R. K., Reaser, J. K., Locke, H., Woodley, S. J., Patz, J. A., Becker, D. J., Oppler, G.,

Hudson, P. J., & Tabor, G. M. (2021). Land use-induced spillover: A call to action to

safeguard environmental, animal, and human health. *The Lancet. Planetary Health*, *5*(4),

e237–e245. https://doi.org/10.1016/S2542-5196(21)00031-0

Prevedello, J. A., Dickman, C. R., Vieira, M. V., & Vieira, E. M. (2017). Population responses of

small mammals to food supply and predators: A global meta-analysis. *Methods in*

Ecology and Evolution, 927–936. https://doi.org/10.1111/1365-

2656.12072@10.1111/(ISSN)2041-210X.SOUTHAMERICA

Pritchard, M. H., & Kruse, G. (1982). *The collection and preservation of animal parasites. Tech.*

Bull. No. 1. Lincoln, Nebraska: Harold W. Manter Laboratory. University of Nebraska

Press. https://cir.nii.ac.jp/crid/1130000795640263296

R Core Team. (2021). *R: A language and environment for statistical computing.* [Computer

software]. R Foundation for Statistical Computing. Vienna, Austria. URL https://www.R-

project.org/

Robert, K., Garant, D., & Pelletier, F. (2012). Keep in touch: Does spatial overlap correlate with

contact rate frequency? *The Journal of Wildlife Management*, *76*(8), 1670–1675.

https://doi.org/10.1002/jwmg.435

Silk, M. J., Croft, D. P., Delahay, R. J., Hodgson, D. J., Boots, M., Weber, N., & McDonald, R. A.

(2017). Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and

Management. *BioScience*, *67*(3), 245–257. https://doi.org/10.1093/biosci/biw175

Silk, M. J., Drewe, J. A., Delahay, R. J., Weber, N., Steward, L. C., Wilson-Aggarwal, J., Boots,

M., Hodgson, D. J., Croft, D. P., & McDonald, R. A. (2018). Quantifying direct and

indirect contacts for the potential transmission of infection between species using a

multilayer contact network. *Behaviour*, *155*(7–9), 731–757.

https://doi.org/10.1163/1568539X-00003493

Sweeny, A. R., Clerc, M., Pontifes, P. A., Venkatesan, S., Babayan, S. A., & Pedersen, A. B.

(2021). Supplemented nutrition decreases helminth burden and increases drug efficacy

in a natural host–helminth system. *Proceedings of the Royal Society B: Biological

Sciences*, *288*(1943), 20202722. https://doi.org/10.1098/rspb.2020.2722

Tamarin, R. H., Ostfeld, R. S., Pugh, S. R., & Bujalska, G. (1990). *Social Systems and

Population Cycles in Voles*. Birkhäuser Verlag.

Tompkins, D. M., Dunn, A. M., Smith, M. J., & Telfer, S. (2011). Wildlife diseases: From

individuals to ecosystems. *Journal of Animal Ecology*, *80*(1), 19–38.

https://doi.org/10.1111/j.1365-2656.2010.01742.x

VanderWaal, K. L., & Ezenwa, V. O. (2016). Heterogeneity in pathogen transmission:

Mechanisms and methodology. *Functional Ecology*, *30*(10), 1606–1622.

https://doi.org/10.1111/1365-2435.12645

Wanelik, K. M., & Farine, D. R. (2022). A new method for characterising shared space use

networks using animal trapping data. *Behavioral Ecology and Sociobiology*, *76*(9), 127.

https://doi.org/10.1007/s00265-022-03222-5