Effects of Food Addition and Helminth Removal on Space Use and Spatial Overlap in

Wild Rodent Populations

Janine Mistrick[1], Shannon M. Kitchen[2], Jasmine S.M. Veitch[3], Samuel Clague[3], Brent C.

Newman[2], Richard J. Hall[4,5,6], Sarah A. Budischak[3], Kristian M. Forbes[2], Meggan E. Craft[1]

[1] Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN, USA

[2] Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[3] W.M. Keck Science Department, Claremont McKenna, Pitzer, and Scripps Colleges,

Claremont, CA, USA

[4] Odum School of Ecology, University of Georgia, Athens, GA, USA

[5] Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia,

Athens, GA, USA

[6] Center for the Ecology of Infectious Diseases, University of Georgia, Athens, GA, USA

Corresponding Author: Janine Mistrick mistr033@umn.edu

ABSTRACT

1. Animal space use and spatial overlap can have important consequences for pathogen transmission. Identifying how environmental variability and inter-individual variation affect spatial patterns to drive transmission heterogeneity in wildlife is a priority for effective management of wildlife disease. However, there are few experimental studies investigating how food abundance and macroparasite infection affect transmission opportunities in wildlife.

2. Wild bank voles (*Myodes glareolus*) are a useful study system to investigate spatial patterns of wildlife and are amenable to experimental manipulations. We conducted a replicated, factorial field experiment in which we added food and removed helminths in wild vole populations in natural forest habitat and monitored vole space use and spatial overlap using capture-mark-recapture methods.

3. Using a combination of home range and network analyses, we quantified vole space use and spatial overlap. We then compared the effects of food addition and helminth removal and investigated the impact of season and sex on space use and spatial overlap.

4. We found that space use was impacted by breeding season and sex but not by food addition or helminth removal. Sex-based, seasonal changes in shared space use were more notable with food addition. Food addition also increased the frequency of, and variability in, spatial overlap occurring between individual voles at trapping locations, while helminth removal had no effect.

5. Our work provides empirical evidence quantifying the spatial effects of food abundance and macroparasite infection on wildlife populations. We demonstrate the potential for high food abundance to increase spatial overlap relevant for pathogen transmission. Our findings also suggest that individual contributions to transmission may be heterogeneous

within populations and that season and sex are important covariates to consider when

identifying when and with whom transmission is most likely to occur.

KEYWORDS

anthelmintic treatment, disease ecology, *Myodes glareolus*, network analysis, resource

provisioning, spatial overlap, transmission, vole

1 INTRODUCTION

How animals use space in their environment and interact with conspecifics is fundamental to

understanding transmission of pathogens. However, contributions to transmission are frequently

heterogeneous among hosts in a population (Woolhouse et al., 1997). Transmission

heterogeneity can result from both environmental variability and inter-individual variation among

hosts (VanderWaal & Ezenwa, 2016) and may be influenced by host population density or sex-

specific social behavior (Webber & Vander Wal, 2020). Identifying how space use, interactions

between animals, and individual-level traits affect transmission continues to be a key goal in

disease ecology (Silk et al., 2017) and is necessary for conservation, domestic animal health,

and public health, but studies that seek to disentangle the relative contributions of

environmental and individual variation to transmission are rare.

Food abundance is a common source of environmental variability that can influence pathogen

transmission by altering host space use and spatial overlap (Becker et al., 2015). Variation in

food abundance across landscapes can have important impacts on infection prevalence in host

populations, either by promoting animal movement through increased attraction of individuals in

high food areas or by decreasing movement through increased site fidelity at areas of high food

abundance (Becker et al., 2018). When wildlife aggregate around point food sources (e.g. bird

feeders, landfills, crop fields), this facilitates interactions between infected and susceptible hosts

and can lead to increased transmission (Forbes et al., 2015).

Individual variation among hosts (e.g. sex, reproductive status, macroparasite infection) can

also affect animal space use and spatial overlap with potential consequences for pathogen

transmission (VanderWaal & Ezenwa, 2016). For example, testosterone-mediated behavior in

mature males can shape social network structure through increased space use and spatial

overlap (Dakin et al., 2021) and is hypothesized to be a key driver of commonly observed male

bias in infection prevalence (Krasnov et al., 2012). Similarly, frequent interactions among

mature individuals during the breeding season can increase transmission opportunities and

create seasonal peaks in pathogen transmission (Altizer et al., 2006).

The effects of individual variation in macroparasite infection on animal space use and spatial

overlap are not as well understood. Macroparasite infection may require infected hosts to

increase their foraging area and thus potential for interactions with conspecifics (Brown et al.,

1994). Conversely, infected hosts may decrease their activity, thus decreasing space use and

interactions (Lopes et al., 2016). Further, since space use and spatial overlap can influence

macroparasite exposure, positive feedbacks between macroparasite exposure and space use

can exacerbate or regulate within-population variation in macroparasite burden and movement

(Hawley et al., 2021). Despite these findings, further research is needed to understand how

individual parasite-induced behaviors scale up to population-level effects on transmission

(Hawley & Altizer, 2011).

Much of our knowledge of how food abundance and macroparasite infection impact pathogen

transmission in wildlife comes from studies of laboratory animals under highly controlled

conditions (e.g. Budischak et al., 2015; Croft et al., 2011) or observational studies in wildlife

(e.g. Jolles et al., 2008; but see Forbes et al., 2016). However, explicit investigation through

rigorous field experiments on wildlife are necessary to understand the effects of environmental

and individual variation on transmission in their ecological context. Such research is pertinent

and timely because urbanization and other land-use change can influence wildlife disease

dynamics (Bradley & Altizer, 2007; Plowright et al., 2021) and are likely to further increase

heterogeneity in food abundance and macroparasite burdens in wildlife.

Our goal was to determine the effects of food abundance and macroparasite infection on space

use and spatial overlap in wildlife populations. We chose bank voles (*Myodes glareolus*) as a

focal study species as they are amenable to experimental manipulation; are a common host for

macroparasite infection (e.g. helminths like *Heligmosomoides glareoli*; population-level helminth

prevalence approx. 80% in some areas, Haukisalmi & Henttonen, 2000); and have well-

documented natural history, particularly with respect to space use behavior of mature voles.

Mature males hold territories that are larger than those of females and that overlap with

territories of other males and with females. Mature females are territorial toward other mature

females but may overlap more with juvenile females and in the non-breeding season. Both food

availability and population density are thought to regulate vole territory size (Bondrup-Nielsen &

Karlsson, 1985).

Here we experimentally manipulated wild bank vole populations via food addition and removal

of intestinal macroparasites (helminths) and monitored individual- and population-level

responses. Specifically, we 1) quantified space use and shared space use in the breeding and

non-breeding season using home range approximation and 2) quantified spatial overlap

between voles each month based on co-location in capture-mark-recapture trapping grids and

visualized the population-level clustering of these "overlaps" using network approaches. As

abundant food often increases small mammal population density through immigration

(Prevedello et al., 2017), we hypothesized that food addition would result in more spatial overlap

that is highly clustered, corresponding to a decrease in vole space use. We hypothesized that

helminth infection would increase host movement (because of increased foraging

requirements), and thus expected that helminth removal would decrease space use and spatial

overlap, resulting in sparser, less clustered networks. Furthermore, we expected space use and

spatial overlap to differ by season and sex such that heterogeneity in the number of spatial

overlaps per vole would peak during the breeding season when activity is highest and most

different by sex and breeding status.

2 MATERIALS AND METHODS

2.1 Study site and experimental design

We conducted a two-factor field experiment in the boreal forests of southern Finland

(61.0775°N, 25.0110°E) where bank voles are the dominant rodent species. These forests

receive snowfall from approximately November-April, and absence of snow cover makes

monitoring and experimental manipulation of voles most feasible from May-October. Twelve

study sites were established in old-growth spruce forest patches. All sites were at least two

kilometers apart to prevent dispersal between replicate populations. At each site, a standardized

trapping grid (100 m x 100 m, 1 hectare) was established of 61 uniquely identified traps spaced

10 meters between grid rows and columns to monitor the vole population (Figure 1).

Sites were randomly assigned to one of four treatment pairings ("treatments"): no manipulation

("unfed-control"; "U-C"); helminth removal but no food addition ("unfed-deworm"; "U-D"); food

addition but no helminth removal ("fed-control", "F-C"); and both food addition and helminth

removal ("fed-deworm"; "F-D"). Each treatment was replicated at three sites (Figure 1). Sites

assigned to food addition received a feed mix of mouse chow pellets and sunflower seeds

evenly distributed on the trapping grid every two weeks from May through November (if snowfall

had not yet accumulated). The supplemental feed mix was dosed at 7.54 kcal/m^2 (following

Sweeny et al., 2021) consisting of 7.875 kg of mouse chow pellets (Altromin 1324 10 mm pellets

maintenance diet for rats and mice [3227 kcal/kg]; Altromin Spezialfutter GmbH & Co. KG,

Lage, Germany) and 7.875 kg sunflower seeds (as fed, 6350 kcal/kg). Voles at deworm sites

were given an oral dose of 10 mg/kg Ivermectin and 100 mg/kg Pyrantel on their first capture in

a trapping occasion; these doses are known to be effective at treating larval and adult helminth

infections (Clerc et al., 2019). Voles at control sites received a matching weight-based dose of

sugar water *(*17.5% sucrose solution) on their first capture in a trapping occasion.

Figure 1. Field study design and experimental set-up. ("Site Locations") Twelve field sites were

established in southern Finland. ("Treatments") Sites were assigned one of four treatment

pairings: unfed-control, unfed-deworm, fed-control, and fed-deworm and each treatment was

replicated at three sites each. ("Trapping Grid Arrangement") The trapping grid included 61

traps in 11 rows and 11 columns. Rows and columns were spaced 10 meters apart and traps

(dots) were placed in an offset arrangement with 20 meters to the next trap in the same row and

column and 14.14 meters to the next trap on the diagonal.

2.2 Data collection

Vole populations were longitudinally monitored via capture-mark-recapture methods. Every four

weeks from May-October 2021, Ugglan multi-capture live traps (Grahnab, Sweden) were baited

with whole oats and set for 48 hours (a "trapping occasion"). Traps were checked each morning

and evening at approximately 0600 and 1600 (four checks per trapping occasion) and captured

animals were processed and then released at the point of capture. Six trapping occasions were

conducted at each site (total trap events: 17,568). Upon first capture, all voles were injected with

a Passive Integrated Transponder ("PIT tag"; "Skinny" PIT Tag, Oregon RFID, USA) for unique

identification. The trap number, PIT tag number, body mass, sex, and reproductive status were

recorded for each vole on their first capture each trapping occasion. If a vole was recaptured in

a given trapping occasion, only the trap number and PIT tag number were recorded, and the

vole was released at the capture location. Reproductive status was categorized as a binary

measure (reproductive or non-reproductive). Females were considered reproductive if they were

lactating or pregnant. Males were considered reproductive if they had scrotal testes.

Reproductive status within the breeding or non-breeding seasons was summarized on an "all or

nothing" basis: if a vole was ever recorded as reproductive during a trapping occasion within the

season, it was recorded as reproductive in that season. Body mass was used to categorize

voles into two age classes: sub-adult (≤ 17 g) or adult (>17 g) (Balčiauskienė et al., 2009;

Balčiauskienė & Balčiauskas, 2009).

All trapping and handling of wild bank voles was conducted under approval of the University of

Arkansas Institutional Animal Care and Use Committee (IACUC #19105) and the Finnish Animal

Ethics Board (ESAVI-17810-2019). Access to forest sites was provided by private landowners

and by Metsähallitus Metsätalous Oy (MH 6302/2019).

2.3 Data analysis

Quantifying transmission-relevant interactions and visualizing heterogeneity in these

interactions can suggest how individuals in a population may differentially contribute to

pathogen transmission (Craft & Caillaud, 2011). Spatiotemporal overlap is a common and

flexible approach that can approximate transmission opportunities for a range of pathogens (e.g.

Godfrey et al., 2010; Porphyre et al., 2008) under the assumption that spatial overlap and

interaction frequency are directly correlated (Robert et al., 2012). Transmission-relevant overlap

may be approximated finely in small areas and short time windows (e.g. co-occurrence at a

point-location) or more broadly over large areas and long time windows (e.g. home range

overlap), depending on the host-pathogen system and research question of interest (Mistrick et

al., 2022). Bank voles are host to a variety of microparasites with confirmed or suspected

zoonotic potential (e.g. cowpox [*Vaccinia* virus], Puumala hantavirus, Ljungan virus), many of

which are primarily shed in excreta and can remain infectious in the environment for days to

weeks (e.g. Puumala hantavirus, Kallio et al., 2006), providing a biologically relevant temporal

window to approximate transmission via environmental contamination.

To understand general patterns of space use by season and sex and how this might impact

transmission opportunities, a home range approach was used to quantify space use and shared

space use for voles repeatedly captured in the breeding or non-breeding season. To quantify

transmission opportunities more specifically and how these may vary with space use and the

experimental manipulations, co-occurrence at trap locations was used to approximate

transmission-relevant spatial overlap between voles. When interactions like spatial overlap are

heterogeneous among individuals, network analysis is a useful method to visualize their

frequency and distribution within a population and predict how transmission may occur (Bansal

et al., 2007). In spatial overlap networks, points (nodes) represent individual animals and

connections between nodes (edges) signify spatiotemporal overlap. Constructing and

comparing networks for populations experiencing varying ecological conditions enables

researchers to explore how environmental variability affects opportunities for pathogen

transmission.

2.3.1 Space use

Vole space use was quantified by aggregating the data from each trapping occasion into a

summer (May-August) and an autumn season (September-October). These divisions were

chosen to capture changes in vole space use between the summer breeding season and

autumn non-breeding season (breeding season May-September in southern Finland; Kaikusalo,

1972). Home range estimators require repeated observations of individuals; therefore the space

use analysis was conducted with a subset of the full vole capture dataset. Only voles that were

captured at least five times in a given season (hereafter "seasonal residents") were used for this

analysis. Voles that were only captured one to four times in a season were not considered

seasonal residents. However, the autumn and summer seasons were considered separately so

any vole could be considered, for example, a seasonal resident in autumn but not in summer.

The sex, reproductive status, and age class of the seasonal residents in each treatment and

season was noted. A 100% minimum convex polygon ("capture polygon") containing all capture

locations in a given season was constructed for each seasonal resident using the

'adehabitatHR' R package (Calenge, 2006). Space use was quantified as the area of a seasonal

resident's capture polygon in hectares. Shared space use was quantified both as the number of

neighbor voles whose capture polygon overlapped that of a focal vole ("number of capture

polygon overlaps") and the proportion of the focal vole's capture polygon that was shared with

each neighbor vole ("proportion of capture polygon overlap"). Proportion overlap was directed

and was calculated for each individual per pair of voles with overlapping capture polygons and

the sexes of the focal and neighbor voles for each pairing were noted. Space use and shared

space use were compared using nonparametric rank-sum tests (Mann-Whitney U test for two

groups; Kruskal-Wallis test for multiple groups).

2.3.2 Spatial overlap

Spatial overlap networks were constructed for each site and trapping occasion from June-

October (capture numbers in May were very low (zero to five animals per site), prohibiting

network construction for all sites so the May data were dropped from this analysis). Spatial

overlap was visualized as a series of five networks per site, each representing a separate

trapping occasion (hereafter called "month"). This method was chosen to capture the temporal

dynamics of network structure and account for animal interactions that vary through time due to

changing population size, demographics, or seasonal behaviors (Pinter-Wollman et al., 2014).

Networks were constructed using the 'CMRnet' R package (Silk et al., 2021). Spatial overlap

was defined as two voles captured in the same or immediately adjacent trap (within 15 meters)

during a trapping occasion to align with similar studies of rodent spatial overlap (Perkins et al.,

2009) and to align with a biologically relevant temporal window for transmission (48 hours).

Because edges were approximated via spatial overlap, edges were undirected. Edges shown in

the main text are unweighted (edges represent the occurrence of spatial overlap between two

voles). An analysis of degree with edges weighted by the frequency of overlap between pairs of

voles is included in the Supporting Information: Weighted degree.

Network metrics provide quantifiable measures to summarize information at both the node level

(e.g. the number of and heterogeneity in overlaps per individual) and at the network level (e.g.

how the overlaps are clustered in the population) (Croft et al., 2008). Both node- and network-

level metrics (Table 1) were calculated using the 'igraph' R package (Csardi & Nepusz, 2006).

Table 1. Description of network metrics and their biological relevance.

	Network Metric	Network Definition	Biological Relevance
Node Level	Unweighted degree	Number of edges connected to a focal node	Number of unique neighbor voles a focal vole spatially overlapped with
	Weighted degree *(see Supplement: Weighted Degree)*	Number of edges connected to a focal node by the weight of each edge; weight represents strength of interaction	Number of spatial overlaps observed for a focal vole, considering the number of repeat overlaps between each vole pair
Network Level	Degree distribution	Frequency distribution of degree counts in a network (weighted or unweighted)	Quantifies the heterogeneity in spatial overlap in the population, high heterogeneity is often associated with super-spreader events
	Network Size	Total number of nodes in the network	Number of voles observed during a trapping occasion at a given site
	Modularity	Extent to which nodes in the network are organized into highly connected clusters that minimally interact with nodes outside their cluster (value scaled 0-1)	Extent to which spatial overlap is clustered among groups of voles that overlap frequently with each other but minimally with other groups of voles at their site
	Betweenness Centralization	Extent to which nodes are connected linearly in a chain or connected to a central hub node like spokes in a wheel (value scaled 0-1)	Extent to which certain voles may be serving as "hubs" in their network to connect many other voles; high centralization suggests short paths across the network

Network data are inherently non-independent and many network metrics are closely related and

are influenced by network size, limiting the ability to make direct comparisons between networks

of varying size (Farine & Whitehead, 2015). To understand how network size was affected by

treatment and contextualize potential confounding factors influencing network metrics, we

investigated the factors affecting network size using a linear mixed-effects model fit by restricted

maximum likelihood (REML) using the 'lme4' package (Bates et al., 2015). We modeled the

main effects of food addition (unfed/fed), helminth removal (control/deworm), and month. Site

was included as a random effect to account for multiple measures at each site and variation

between replicate sites within a treatment. Network size was log-transformed to normalize

distribution of the data.

We quantified the mean and standard deviation of network clustering metrics (i.e. modularity

and betweenness centralization) by month to explore differences between treatments.

Unweighted degree for all voles at each site in a given month was visualized as a frequency

distribution to identify heterogeneity in the number of spatial overlaps per vole. Differences in

degree and network clustering metrics between sites and treatments were compared using

Kruskal-Wallis tests with Dunn tests (Benjamini-Hochberg correction) for post hoc comparisons.

Heterogeneity in degree is an important feature of networks that can suggest heterogeneity in

transmission, particularly if a small proportion of the population has the largest number of

connections (Lloyd-Smith et al., 2005). We defined "super-overlappers" as voles with an

unweighted degree greater than 1.5 times the interquartile range of degree values of their site in

a given month. As such, the degree value of a super-overlapper was determined in relation to

the population at its site in a given month, not by an arbitrary threshold, and therefore the actual

degree value of super-overlappers differs between populations and months. Super-overlappers

were identified for each site in each month and the sex and reproductive status of these

individuals was noted.

All data analysis was conducted in program R version 4.1.2 (R Core Team, 2021).

3 RESULTS

We captured 770 unique voles totaling 2086 capture events from the 12 sites during the May-

October study period. Of these, 456 individuals (59.2%) were captured at least twice (mean

captures per vole: 2.71 ± 2.30 [range: 1-17]).

3.1 Space use

In total, 95 voles (95/770, 12.3% of all voles captured) were seasonally resident on our trapping

grids (eight voles were resident in both summer and autumn). We documented 314 captures of

48 seasonal residents in the summer breeding season and 316 captures of 55 seasonal

residents in the autumn non-breeding season (Figure S1). An additional 20 animals were

captured at least five times, but the captures were split between the breeding and non-breeding

seasons, so they were not included as seasonal residents. In summer and autumn, seasonal

residents were both males and females (Figure S2). The breeding status of seasonal residents

generally matched that expected in the season (i.e. majority reproductive in summer, non-

reproductive in autumn) and this was consistent between treatments (Figure S3). The age class

of seasonal residents suggested most were adults (>17 g) in summer and sub-adults (≤17 g) in

autumn, and this was similar across treatments (Supporting Information: Age classes of

seasonal residents; Figure S4).

3.1.1 Space use was greatest in males during the breeding season

Space use of seasonal residents varied by sex and season but did not differ by treatment. In the

summer breeding season, male and female space use differed (Mann-Whitney U test, p<0.001)

and male space use was greater and more variable than that of females (male mean

area=0.18±0.12 hectares ([range: 0.02-0.60 ha]; female mean area=0.04±0.04 hectares [range:

0.00-0.15 ha]; Figure 2A; Table S1). Summer space use did not differ between treatments for

males or females (Kruskal-Wallis test, males $p=0.77$; females $p=0.29$). In the autumn non-

breeding season, male and female space use did not differ (Mann-Whitney U test, $p=0.22$; male

mean area = 0.04±0.04 [range: 0.00-0.20]; female mean area=0.03±0.04 [range: 0.00-0.17];

Figure 2A; Table S1). Autumn space use did not differ between treatments for either sex

(Kruskal-Wallis test, males $p=0.16$; females $p=0.51$). Male space use differed between summer

and autumn with mean summer space use over four times greater than autumn space use

(Mann-Whitney U test, p<0.001). Female space use was not different between summer and

autumn (Mann-Whitney U test, $p=0.72$).

3.1.2 Food addition increased female-female shared space use in the non-breeding season

The number of capture polygon overlaps per seasonal resident (indicating shared space use

between pairs of voles) differed by season for males and was higher in summer (Mann-Whitney

U test, p<0.001; summer mean overlaps=2.16±1.61 [range: 0-5]; autumn mean

overlaps=0.59±0.78 [range: 0-2]; Table S2). Females had a similar number of overlaps in

summer and autumn (Mann-Whitney U test, $p=0.64$). In summer, males had more overlaps than

females whereas the two sexes had a similar number in autumn (Mann-Whitney U test, summer

p<0.001; autumn $p=0.91$).

The proportion of capture polygon overlap between seasonal residents did not differ by

treatment in summer or fall (Kruskal-Wallis test, summer $p=0.50$; fall $p=0.13$). In the summer,

the proportion of capture polygon overlap was low to moderate in most treatments and was only

observed for male-male pairings or male-female pairings (Figure 2B). In the autumn, the

proportion of overlap was higher than in summer in all treatments except unfed-deworm where it

was similar (Figure 2B). Female-female overlap was observed in fed treatments in autumn and

the mean proportion of overlap was high (F-C mean=0.54±0.51 [range: 0.18-0.90]; F-D

mean=0.50±0.24 [range: 0.33-0.67]), while low female-female overlap was observed in unfed-

deworm treatments (U-D mean=0.16±0.13 [range: 0.06-0.25]; Table S3). There was no male-

male overlap in fed treatments in autumn.

Figure 2. Summary boxplots of space use and shared space use of seasonal residents by sex

and season, jittered points overlaid to show actual data values. (A) Space use (capture area

size in hectares) in the summer (left panel) and autumn (right panel). Males (blue) show greater

space use compared to females (red) in summer (Kruskal-Wallis test, $p<0.001$). (B) Shared

space use (proportion of capture polygon overlap) in summer (left panel) and autumn (right

panel). Female-female shared space use (red) is rarely observed in summer but is high in fed

treatments in autumn.

3.2 Spatial overlap networks

The June-October data used for this analysis comprised 2052 captures of 759 unique voles. All

sites had at least two animals captured in each of the months in which networks were

constructed, however, two sites had no edges in the June network. Networks frequently had

disconnected nodes and often had multiple components. Generally, we only observed networks

with all edges in a single component in the months of highest population density (Figure S5).

3.2.1 Network size was greatest in fed treatments

Network size was highly variable across months and treatments (range: 2-52 voles). Network

size increased through the year in all treatments. Networks were smallest in June, peaked in

August (unfed treatments) or September (fed treatments), and then decreased slightly though

October (Figure 3A). Network size was moderately affected by food treatment (β=0.50,

p=0.013) and by month (β=0.31, p<0.001; Table S4) with fed treatments and months later in the

year having larger networks. At peak size in September, mean network size in fed treatments

was 1.6-3.1x larger than unfed treatments (Table S5). Deworming treatment had a negligible

effect on network size (β=-0.17, p=0.340; Table S4).

3.2.2 Network clustering metrics were similar between treatments

Network clustering metrics were visualized as a mean and standard deviation per treatment (for

a description of the network metrics and their biological relevance, see Table 1). Modularity

increased and became less variable within treatment from June to October (mean across

treatments: June: 0.27±0.27; Oct: 0.62±0.08, Figure 3B; Table S6), indicating voles overlapped

with more distinct groups of other voles as the year progressed. Mean modularity was similar

between treatments in each month (Kruskal-Wallis test, p>0.12). Betweenness centralization

was low (annual mean across treatments: 0.13±0.11; Figure 3C; Table S7) and similar between

treatments in all months (Kruskal-Wallis test, *p*>0.22), indicating overlap occurred between

many unique pairs of voles and was not focused at a hub individual.

Figure 3. Network size by site and mean values of network clustering metrics by treatment: (A)

Network size (number of voles) by month increased more dramatically in fed treatments (purple)

compared to unfed treatments (green) and fed treatments reached a larger maximum network

size. (B) Mean modularity (the clustering of nodes into highly connected groups) increased from

June to October and became less variable. (C) Mean betweenness centralization (whether the

network is linear or focused at a hub) was low throughout the year. Network metrics shown per

treatment: unfed-control (light green), unfed-deworm (dark green), fed-control (light purple), fed-

deworm (dark purple). Error bars show standard deviation around the mean.

3.2.3 Number of spatial overlaps per vole was greater and more heterogeneous in fed

treatments

Unweighted degree (the number of unique spatial overlaps between a focal vole and its

neighbors) per site and month was visualized as density distributions. In these, the x-axis

indicates the value of interest, and the height of the curve suggests the frequency of occurrence

of that value in the population. Bimodal or multimodal distributions indicate that multiple values

occur in the population at a similar frequency. Narrow distributions suggest low variation in the

population values while wide distributions indicate greater variation.

Mean unweighted degree was similar between treatments in July and August (Kruskal-Wallis

test, July p=0.086; August p=0.056; Figure 4; Table S8). In June, mean degree in the unfed-

control treatments was different from all other treatments (Kruskal-Wallis test $p<0.001$; Dunn

test $p<0.018$) but there was no difference between the other treatments (Dunn test $p>0.08$).

Mean degree differed by treatment in September and October (Kruskal-Wallis test, Sept

$p<0.001$; Oct $p<0.001$) where fed treatments had higher degree than unfed (Dunn test, Sept

$p<0.0013$; Oct $p<0.013$; Figure 4; Table S8). The distribution of degree values per site varied by

treatment and month. Fed treatments tended to have longer tails of degree (i.e. a few voles had

a large number of overlaps) in August-October (Figure 4).

Figure 4. Density distributions of unweighted degree for each of the 12 study sites from June-

October (colored by treatment, means indicated by vertical lines). Fed treatments (purple) had

longer tails of degree distribution in August-October compared to unfed treatments (green).

We also explored weighted degree (edges weighted by the number of overlaps between each

unique pair of voles), but trends were similar to those of unweighted degree (Supporting

Information: Weighted degree; Table S9; Figure S6).

3.2.4 Super-overlappers were most numerous in fed-deworm treatments

Based on unweighted degree values, we identified 31 super-overlappers (i.e. voles that

disproportionately overlapped with other voles) representing 28 unique voles (three voles were

super-overlappers in two separate months) across the 12 sites from June-October (Supporting

Information: Super-overlappers). Super-overlappers were not observed in unfed-control

treatments and only one was observed in unfed-deworm treatments. All other super-overlappers

were observed in fed treatments with 5 times as many super-overlappers in fed-deworm

compared to fed-control (Figure S7). Super-overlappers were present in all months but months

with the largest network sizes had the most super-overlappers (i.e. 14 super-overlappers were

from September and 8 were from August, Figure S8A). Both males and females and

reproductive and non-reproductive voles were super-overlappers. The number of males versus

females and reproductive versus non-reproductive voles that were considered super-

overlappers were similar between fed-control and fed-deworm treatments (Fisher's exact test,

male:female two-tailed p=0.34; reproductive:non-reproductive two-tailed p=0.64; Figure S8B,C;

Table S10).

4 DISCUSSION

Food abundance and macroparasite infection can alter individual space use and intraspecific

interactions in ways that influence pathogen transmission. By experimentally manipulating food

abundance and helminth infection in voles, we quantified drivers of variation in space use and

spatial overlap, and how these were modulated by season and sex. Spatial overlap frequency

and heterogeneity were affected by food addition but not deworming treatment. However, the

population-level clustering of overlaps was similar between treatments. Space use was

influenced by season and sex, but not by food addition or deworming treatment. However, sex-

based, seasonal changes in shared space use were most apparent with food addition. These

findings suggest that while food abundance generally increases transmission opportunities, sex

and season are important factors to determine the "who" and "when" of greatest transmission

potential.

Seasonal resident vole space use was not affected by food addition or helminth removal in

either summer or autumn, which may indicate similar transmission opportunities across a wide

range of habitat conditions. The similarity in space use across treatments is surprising as

supplemental food is generally thought to influence home range size of bank voles (Bondrup-

Nielsen & Karlsson, 1985; Mazurkiewicz, 1983) though studies in other rodents have not found

food to be a major factor determining home range size (Sulok et al., 2004; Wolff, 1985).

However, space use did differ by sex. During the summer breeding season (May-August),

resident male voles had much greater space use than resident females and male space use

was shared with both other males and females. This suggests mature males have a high

potential to contribute to transmission with both male and female conspecifics during the

breeding season: their large space use provides a greater potential to spread infectious material

(urine, feces) across an area that is frequented by other voles. Further, this space use overlap

coincides with the breeding season when activity is naturally high in the population where direct

and indirect interactions with high transmission potential (e.g. mating or fighting, territory

marking - respectively) are more likely to occur. In the autumn non-breeding season

(September-October), both male and female space use was limited - potentially decreasing

transmission opportunities - however, female-female shared space use was high in fed

treatments. High population density associated with abundant food may be an important factor

influencing increased shared space use among females, particularly in the non-breeding season

when mature females are less territorial (Stenseth et al., 1988). This increased overlap could

provide increased transmission opportunities among females. Thus, while we observed males

were important to transmission opportunities during the breeding season regardless of food

abundance, when food abundance is high, females may become more important for

transmission in the non-breeding season. Taken together, our space use findings suggest that

seasonal and sex-based differences in animal space use and shared space use may contribute

to variable patterns of transmission such as seasonal peaks in transmission or changes in which

individuals contribute most to transmission opportunities (Altizer et al., 2006; Hirsch et al.,

2016).

These conclusions about space use were generated from a subset of the vole population that

was captured frequently enough to conduct home range analyses. While the interactions of

seasonal residents with established home ranges likely account for many of the transmission-

relevant interactions in the population (particularly during the breeding season), the role of

juveniles or transient individuals may be overlooked. If, for example, juvenile voles have

markedly different space use than mature adults, this could minimize the difference in vole

space use between the breeding and non-breeding seasons, potentially providing high and

consistent opportunities for environmental transmission throughout the year. Incorporating other

monitoring techniques along with capture-mark-recapture methods (e.g. stationary proximity

loggers placed on the grid to record the PIT tags of animals that pass within a specified

distance) would provide additional locations per individual and would bolster our understanding

of the space use of all animals in the population.

Spatial overlap between voles was strongly affected by food addition but there was no clear

effect of deworming treatment. Food addition resulted in larger networks with more voles than

unfed treatments and voles in fed treatments had more spatial overlaps (higher mean degree)

than voles in unfed treatments. Previous studies have attributed short-term increases in rodent

populations with food addition to increased immigration rather than recruitment (Prevedello et

al., 2017) which aligns with the understanding that abundant food attracts animals to an area.

This influx of animals could lead to an increased rate of pathogen introduction to areas of high

food abundance. Considering heterogeneous environments as a metapopulation model, areas

of high food abundance may play an important role in both concentrating animals, interactions,

and pathogens and also being a source of infected dispersers, potentially spreading pathogens

more broadly on landscape (Becker et al., 2018). Networks in fed treatments also had a greater

number of super-overlappers than unfed treatments, showing that heterogeneity in the number

of overlaps increased in populations with abundant food - and the combination of food addition

and helminth removal may further increase the number of super-overlappers. Heterogeneity in

degree is a common trait of populations and extreme heterogeneity (i.e. the presence of super-

spreaders) has been identified as a key feature creating variability in whether diseases spread

rapidly or fade out in a population (Lloyd-Smith et al., 2005; Meyers et al., 2005). While the

heterogeneity in unweighted degree observed in this study (Supporting Information: Dispersion

parameter (k value) estimates; Table S11) is not as extreme as that of documented

superspreader events (Lloyd-Smith et al., 2005), the observed heterogeneity in spatial overlaps

per vole does suggest that individual voles do not contribute equally to transmission

opportunities and certain individuals could be more important for transmission than others,

particularly when food is abundant.

5 CONCLUSIONS

This study has provided empirical evidence of the effects of food abundance and macroparasite

infection on space use and spatial overlap in wild rodent populations. Field studies investigating

the effects of environmental and individual variation on pathogen transmission are limited, thus

this work addresses a gap in our understanding of factors influencing wildlife disease dynamics.

Our findings suggest increased transmission opportunities with high food abundance, though

which individuals contribute most to transmission may vary by sex and season, particularly

when food abundance and population density are high. We have inferred potential

consequences for transmission, aiding future research that pairs measures of space use and

spatial overlap with infection prevalence to test how these spatial metrics predict transmission

opportunities. As such, this research provides necessary, empirical evidence for the effects of

environmental and individual variation on animal spatial patterns and suggests associated

consequences for pathogen transmission, ultimately helping to better understand and control

infectious diseases in animal populations.

REFERENCES

Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M., & Rohani, P. (2006). Seasonality
and the dynamics of infectious diseases. *Ecology Letters*, *9*(4), 467–484.
https://doi.org/10.1111/j.1461-0248.2005.00879.x

Balčiauskienė, L., & Balčiauskas, L. (2009). Prediction of the body mass of the bank vole
Myodes glareolus from skull measurements: Tava-leethiire (*Myodes glareolus*)
kehamassi hindamine koljumõõtmete põhjal. *Estonian Journal of Ecology*, *58*(2), 77–85.
https://doi.org/10.3176/eco.2009.2.01

Balčiauskienė, L., Balčiauskas, L., & Čepukienė, A. (2009). Growth of the bank vole *Myodes*
glareolus in the non-vegetative period in NE Lithuania: Leethiire (*Myodes glareolus*)
kehamõõtmete kasv talveperioodil Kirde-Leedus. *Estonian Journal of Ecology*, *58*(2),
86–93. https://doi.org/10.3176/eco.2009.2.02

Bansal, S., Grenfell, B. T., & Meyers, L. A. (2007). When individual behaviour matters:
Homogeneous and network models in epidemiology. *Journal of The Royal Society*
Interface, *4*(16), 879–891. https://doi.org/10.1098/rsif.2007.1100

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software*, *67*, 1–48.
https://doi.org/10.18637/jss.v067.i01

Becker, D. J., Snedden, C. E., Altizer, S., & Hall, R. J. (2018). Host Dispersal Responses to
Resource Supplementation Determine Pathogen Spread in Wildlife Metapopulations.
The American Naturalist, *192*(4), 503–517. https://doi.org/10.1086/699477

Becker, D. J., Streicker, D. G., & Altizer, S. (2015). Linking anthropogenic resources to wildlife–
pathogen dynamics: A review and meta-analysis. *Ecology Letters*, *18*(5), 483–495.
https://doi.org/10.1111/ele.12428

Bondrup-Nielsen, S., & Karlsson, F. (1985). Movements and spatial patterns in populations of
Clethrionomys species: A review. *Annales Zoologici Fennici*, *22*(3), 385–392. JSTOR.

Bradley, C. A., & Altizer, S. (2007). Urbanization and the ecology of wildlife diseases. *Trends in*

Ecology & Evolution, *22*(2), 95–102. https://doi.org/10.1016/j.tree.2006.11.001

Brown, E. D., Macdonald, D. W., Tewand, T. E., & Todd, I. A. (1994). *Apodemus sylvaticus*

infected with *Heligmosomoides polygyrus* (Nematoda) in an arable ecosystem:

Epidemiology and effects of infection on the movements of male mice. *Journal of*

Zoology, *234*(4), 623–640. https://doi.org/10.1111/j.1469-7998.1994.tb04869.x

Budischak, S. A., Sakamoto, K., Megow, L. C., Cummings, K. R., Urban, J. F., & Ezenwa, V. O.

(2015). Resource limitation alters the consequences of co-infection for both hosts and

parasites. *International Journal for Parasitology*, *45*(7), 455–463.

https://doi.org/10.1016/j.ijpara.2015.02.005

Calenge, C. (2006). The package "adehabitat" for the R software: A tool for the analysis of

space and habitat use by animals. *Ecological Modelling*, *197*(3), 516–519.

https://doi.org/10.1016/j.ecolmodel.2006.03.017

Clerc, M., Babayan, S. A., Fenton, A., & Pedersen, A. B. (2019). Age affects antibody levels and

anthelmintic treatment efficacy in a wild rodent. *International Journal for Parasitology:*

Parasites and Wildlife, *8*, 240–247. https://doi.org/10.1016/j.ijppaw.2019.03.004

Craft, M. E., & Caillaud, D. (2011). Network Models: An Underutilized Tool in Wildlife

Epidemiology? *Interdisciplinary Perspectives on Infectious Diseases*, 2011, 1-12.

https://doi.org/10.1155/2011/676949

Croft, D. P., Edenbrow, M., Darden, S. K., Ramnarine, I. W., van Oosterhout, C., & Cable, J.

(2011). Effect of gyrodactylid ectoparasites on host behaviour and social network

structure in guppies *Poecilia reticulata*. *Behavioral Ecology and Sociobiology*, *65*(12),

2219–2227. https://doi.org/10.1007/s00265-011-1230-2

Croft, D. P., James, R., & Krause, J. (2008). *Exploring Animal Social Networks*. Princeton

University Press.

Csardi, G., & Nepusz, T. (2006). The Igraph Software Package for Complex Network Research.

589   *InterJournal Complex Systems*, *1695*(5), 1–9.

590 Dakin, R., Moore, I. T., Horton, B. M., Vernasco, B. J., & Ryder, T. B. (2021). Testosterone-

591   mediated behaviour shapes the emergent properties of social networks. *Journal of*

592   *Animal Ecology*, *90*(1), 131–142. https://doi.org/10.1111/1365-2656.13305

593 Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting animal social

594   network analysis. *Journal of Animal Ecology*, *84*(5), 1144–1163.

595   https://doi.org/10.1111/1365-2656.12418

596 Forbes, K. M., Henttonen, H., Hirvelä-Koski, V., Kipar, A., Mappes, T., Stuart, P., & Huitu, O.

597   (2015). Food provisioning alters infection dynamics in populations of a wild rodent.

598   *Proceedings of the Royal Society B: Biological Sciences*, *282*(1816), 20151939.

599   https://doi.org/10.1098/rspb.2015.1939

600 Forbes, K. M., Mappes, T., Sironen, T., Strandin, T., Stuart, P., Meri, S., Vapalahti, O.,

601   Henttonen, H., & Huitu, O. (2016). Food limitation constrains host immune responses to

602   nematode infections. *Biology Letters*, *12*(9), 20160471.

603   https://doi.org/10.1098/rsbl.2016.0471

604 Godfrey, S. S., Moore, J. A., Nelson, N. J., & Bull, C. M. (2010). Social network structure and

605   parasite infection patterns in a territorial reptile, the tuatara (*Sphenodon punctatus*).

606   *International Journal for Parasitology*, *40*(13), 1575–1585.

607   https://doi.org/10.1016/j.ijpara.2010.06.002

608 Haukisalmi, V., & Henttonen, H. (2000). Variability of helminth assemblages and populations in

609   the bank vole *Clethrionomys glareolus*. *Polish Journal of Ecology*, *48*, 219–231.

610 Hawley, D. M., & Altizer, S. M. (2011). Disease ecology meets ecological immunology:

611   Understanding the links between organismal immunity and infection dynamics in natural

612   populations. *Functional Ecology*, *25*(1), 48–60. https://doi.org/10.1111/j.1365-

613   2435.2010.01753.x

614 Hawley, D. M., Gibson, A. K., Townsend, A. K., Craft, M. E., & Stephenson, J. F. (2021).

Bidirectional interactions between host social behaviour and parasites arise through

ecological and evolutionary processes. *Parasitology*, *148*(3), 274–288.

https://doi.org/10.1017/S0031182020002048

Hirsch, B. T., Reynolds, J. J. H., Gehrt, S. D., & Craft, M. E. (2016). Which mechanisms drive

seasonal rabies outbreaks in raccoons? A test using dynamic social network models.

Journal of Applied Ecology, *53*(3), 804–813. https://doi.org/10.1111/1365-2664.12628

Jolles, A. E., Ezenwa, V. O., Etienne, R. S., Turner, W. C., & Olff, H. (2008). Interactions

Between Macroparasites and Microparasites Drive Infection Patterns in Free-Ranging

African Buffalo. *Ecology*, *89*(8), 2239–2250. https://doi.org/10.1890/07-0995.1

Kaikusalo, A. (1972). Population turnover and wintering of the bank vole, *Clethrionomys*

glareolus (Schreb.), in southern and central Finland. *Annales Zoologici Fennici*, *9*(4),

219–224.

Kallio, E. R., Klingström, J., Gustafsson, E., Manni, T., Vaheri, A., Henttonen, H., Vapalahti, O.,

& Lundkvist, Å. (2006). Prolonged survival of Puumala hantavirus outside the host:

Evidence for indirect transmission via the environment. *Journal of General Virology,*

87(8), 2127–2134. https://doi.org/10.1099/vir.0.81643-0

Krasnov, B. R., Bordes, F., Khokhlova, I. S., & Morand, S. (2012). Gender-biased parasitism in

small mammals: Patterns, mechanisms, consequences. *Mammalia*, *76*(1), 1–13.

https://doi.org/10.1515/mammalia-2011-0108

Lloyd-Smith, J. O., Schreiber, S. J., Kopp, P. E., & Getz, W. M. (2005). Superspreading and the

effect of individual variation on disease emergence. *Nature*, *438*(7066), Article 7066.

https://doi.org/10.1038/nature04153

Lopes, P. C., Block, P., & König, B. (2016). Infection-induced behavioural changes reduce

connectivity and the potential for disease spread in wild mice contact networks. *Scientific*

Reports, *6*(1), Article 1. https://doi.org/10.1038/srep31790

Mazurkiewicz, M. (1983). Spatial organization of the population. *Acta Theriologica*, *28*(Suppl. 1),

103–144. https://doi.org/10.4098/AT.arch.83-48

Meyers, L. A., Pourbohloul, B., Newman, M. E. J., Skowronski, D. M., & Brunham, R. C. (2005).

Network theory and SARS: Predicting outbreak diversity. *Journal of Theoretical Biology*,

232(1), 71–81. https://doi.org/10.1016/j.jtbi.2004.07.026

Mistrick, J., Gilbertson, M. L. J., White, L. A., & Craft, M. E. (2022). Constructing animal

networks for parasite transmission inference. In V. Ezenwa, S. M. Altizer, & R. Hall

(Eds.), *Animal Behavior and Parasitism*. Oxford University Press.

Perkins, S. E., Cagnacci, F., Stradiotto, A., Arnoldi, D., & Hudson, P. J. (2009). Comparison of

social networks derived from ecological data: Implications for inferring infectious disease

dynamics. *The Journal of Animal Ecology*, *78*(5), 1015–1022.

https://doi.org/10.1111/j.1365-2656.2009.01557.x

Pinter-Wollman, N., Hobson, E. A., Smith, J. E., Edelman, A. J., Shizuka, D., de Silva, S.,

Waters, J. S., Prager, S. D., Sasaki, T., Wittemyer, G., Fewell, J., & McDonald, D. B.

(2014). The dynamics of animal social networks: Analytical, conceptual, and theoretical

advances. *Behavioral Ecology*, *25*(2), 242–255. https://doi.org/10.1093/beheco/art047

Plowright, R. K., Reaser, J. K., Locke, H., Woodley, S. J., Patz, J. A., Becker, D. J., Oppler, G.,

Hudson, P. J., & Tabor, G. M. (2021). Land use-induced spillover: A call to action to

safeguard environmental, animal, and human health. *The Lancet Planetary Health*, *5*(4),

e237–e245. https://doi.org/10.1016/S2542-5196(21)00031-0

Porphyre, T., Stevenson, M., Jackson, R., & McKenzie, J. (2008). Influence of contact

heterogeneity on TB reproduction ratio R_0 in a free-living brushtail possum *Trichosurus*

vulpecula population. *Veterinary Research*, *39*(3), 1.

https://doi.org/10.1051/vetres:2008007

Prevedello, J. A., Dickman, C. R., Vieira, M. V., & Vieira, E. M. (2017). Population responses of

small mammals to food supply and predators: A global meta-analysis. *Methods in*

Ecology and Evolution, 927–936. https://doi.org/10.1111/1365-2656.12072

R Core Team. (2021). *R: A language and environment for statistical computing.* R Foundation

for Statistical Computing. Vienna, Austria. URL https://www.R-project.org/

Robert, K., Garant, D., & Pelletier, F. (2012). Keep in touch: Does spatial overlap correlate with

contact rate frequency? *The Journal of Wildlife Management*, *76*(8), 1670–1675.

https://doi.org/10.1002/jwmg.435

Silk, M. J., Croft, D. P., Delahay, R. J., Hodgson, D. J., Boots, M., Weber, N., & McDonald, R. A.

(2017). Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and

Management. *BioScience*, *67*(3), 245–257. https://doi.org/10.1093/biosci/biw175

Silk, M. J., McDonald, R. A., Delahay, R. J., Padfield, D., & Hodgson, D. J. (2021). CMRnet: An

r package to derive networks of social interactions and movement from mark–recapture

data. *Methods in Ecology and Evolution*, *12*(1), 70–75. https://doi.org/10.1111/2041-

210X.13502

Stenseth, N. Chr., Bondrup-Nielsen, S., & Ims, R. A. (1988). A Population Dynamics Model for

Clethrionomys: Sexual Maturation, Spacing Behaviour and Dispersal. *Oikos*, *52*(2), 186–

193. JSTOR. https://doi.org/10.2307/3565246

Sulok, M., Slade, N. A., & Doonan, T. J. (2004). Effects of Supplemental Food on Movements of

Cotton Rats (*Sigmodon hispidus*) in Northeastern Kansas. *Journal of Mammalogy*, *85*(6),

1102–1105. https://doi.org/10.1644/BEH-112.1

Sweeny, A. R., Clerc, M., Pontifes, P. A., Venkatesan, S., Babayan, S. A., & Pedersen, A. B.

(2021). Supplemented nutrition decreases helminth burden and increases drug efficacy

in a natural host–helminth system. *Proceedings of the Royal Society B: Biological*

Sciences, *288*(1943), 20202722. https://doi.org/10.1098/rspb.2020.2722

VanderWaal, K. L., & Ezenwa, V. O. (2016). Heterogeneity in pathogen transmission:

Mechanisms and methodology. *Functional Ecology*, *30*(10), 1606–1622.

https://doi.org/10.1111/1365-2435.12645

Webber, Q. M. R., & Vander Wal, E. (2020). Heterogeneity in social network connections is

density-dependent: Implications for disease dynamics in a gregarious ungulate.

Behavioral Ecology and Sociobiology, *74*(6), 77. https://doi.org/10.1007/s00265-020-

02860-x

Wolff, J. O. (1985). The effects of density, food, and interspecific interference on home range

size in *Peromyscus leucopus* and *Peromyscus maniculatus. Canadian Journal of*

Zoology, *63*(11), 2657–2662. https://doi.org/10.1139/z85-397

Woolhouse, M. E. J., Dye, C., Etard, J.-F., Smith, T., Charlwood, J. D., Garnett, G. P., Hagan,

P., Hii, J. L. K., Ndhlovu, P. D., Quinnell, R. J., Watts, C. H., Chandiwana, S. K., &

Anderson, R. M. (1997). Heterogeneities in the transmission of infectious agents:

Implications for the design of control programs. *Proceedings of the National Academy of*

Sciences, *94*(1), 338–342. https://doi.org/10.1073/pnas.94.1.338

ACKNOWLEDGEMENTS

This research was funded by the National Science Foundation (DEB-1911925). J.M. was

supported by the National Science Foundation Graduate Research Fellowship Program under

Grant No. CON-75851, project 00074041. Any opinions, findings, and conclusions or

recommendations expressed in this material are those of the authors and do not necessarily

reflect the views of the National Science Foundation. J.M. was also supported by the Dayton

Fund of the Bell Museum and the University of Minnesota Ecology, Evolution, and Behavior

Graduate Program and the College of Biological Sciences.

We are grateful to our collaborators at the Lammi Biological Station and University of Helsinki

for supporting our in-country work: John Loehr, Janne Sundell, Esa-Pekka Tuominen, Tiina

Tulonen, Matti Kotakorpi, Riitta Ilola, Jaakko Vainionpää, Tomas Strandin, and Tarja Sironen,

and to the students who helped collect and process the field data: Alexis Beagle, Muriel

CONFLICT OF INTEREST

724 The authors claim no conflict of interest.

AUTHOR CONTRIBUTIONS

Kris Forbes and Sarah Budischak (lead), Meggan Craft and Richard Hall (supporting) conceived

the ideas, designed methodology, and secured the funding; Shannon Kitchen, Janine Mistrick,

Jasmine Veitch, Samuel Clague (lead), and Brent Newman (supporting) collected the data;

Janine Mistrick analyzed the data, created visualizations, and led the writing of the manuscript

under the mentorship of Meggan Craft. All authors contributed critically to the drafts and gave

final approval for publication.

DATA AVAILABILITY STATEMENT

735 Data will be made available through the Dryad Digital Repository upon manuscript acceptance.

736 Code is available in a GitHub repository (github.com/jmistrick/vole-spatial) which will be

737 archived on Zenodo upon manuscript acceptance.