TOOLS AND RESOURCES

voyAGEr: free web interface for the analysis of age-related gene expression alterations in

human tissues

Short running title:

Age-related gene changes profiling app

Arthur L. Schneider[1,2,†]
arthur.schneider@centraliens-nantes.org

Rita Martins-Silva[1,†]
rita.silva@medicina.ulisboa.pt

Alexandre Kaizeler[1,†]
alexandre.afonso@medicina.ulisboa.pt

Nuno Saraiva-Agostinho[1,3]
nunodanielagostinho@gmail.com

Nuno L. Barbosa-Morais[1,*]
nmorais@medicina.ulisboa.pt

[1] Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina, Universidade
de Lisboa, Av. Prof. Egas Moniz, 1649-028 Lisbon, Portugal

[2] *Present address:* Sia Partners, 4 Rue Voltaire, 44000 Nantes, France

[3] *Present address:* European Molecular Biology Laboratory, European Bioinformatics
Institute, Wellcome Genome Campus, Hinxton, CB10 1SD, Cambridge, UK

[†] Equal contributions

* Corresponding author

Abstract

We herein introduce voyAGEr, an online graphical interface to explore age-related gene

expression alterations in 49 human tissues. voyAGEr offers a visualisation and statistical

toolkit for the finding and functional exploration of sex- and tissue-specific transcriptomic

changes with age. In its conception, we developed a novel bioinformatics pipeline leveraging

RNA sequencing data, from the GTEx project, encompassing more than 900 individuals.

voyAGEr reveals transcriptomic signatures of the known asynchronous ageing between

tissues, allowing the observation of tissue-specific age-periods of major transcriptional

changes, associated with alterations in different biological pathways, cellular composition,

and disease conditions.

Notably, voyAGEr was created to assist researchers with no expertise in bioinformatics,

providing a supportive framework for elaborating, testing and refining their hypotheses on

the molecular nature of human ageing and its association with pathologies, thereby also

aiding in the discovery of novel therapeutic targets. voyAGEr is freely available at

https://compbio.imm.medicina.ulisboa.pt/app/voyAGEr.

Keywords

47 Humans; Ageing; Computational Biology; Transcriptome; Gene Expression Profiling

Main text

Introduction

The ageing-associated progressive loss of proper tissue homeostasis maintenance makes age a prevalent risk factor for many human pathologies, including cancer, neurodegenerative and cardiovascular diseases [1–3]. A better comprehension of the molecular mechanisms of human ageing is thus required for the development and effective application of therapies targeting its associated morbidities.

Dynamic transcriptional alterations accompany most physiological processes occurring in human tissues [4]. Transcriptomic analyses of tissue samples can thus provide snapshots of cellular states therein and insights into how their modifications over time impact tissue physiology. A small proportion of transcripts has indeed been shown to vary with age in tissue- [5] and sex-specific [6–11] manners. Such variations reflect dysregulations of gene expression that underlie cellular disfunctions [5].

Many studies analysed the age-related changes in gene expression in rodent tissues [12–18], emphasising the role in ageing of genes related to inflammatory responses, cell cycle or the electron transport chain. However, while it is possible to monitor the modifications in gene expression over time in those species by sequencing transcriptomes of organs of littermates at different ages, as a close surrogate of longitudinality, such studies cannot be conducted in humans for ethical reasons. Indeed, most studies aimed at profiling ageing-related gene expression changes in human tissues focused on a single tissue (e.g. muscle [19–21], kidney [22], brain [23–27], skin [28,29], blood [30,31], liver [32], retina [33]) and/or are limited to a comparison between young and old individuals [20,21,28,32,33], failing to fully capture the changes of the tissue-specific gene expression landscape throughout ageing [5]. A few studies were nonetheless led on more than one tissue in humans, from post-mortem samples [34,35] and biopsies [36,37], and in mice [18,36] and rats [38]. The age-related transcriptional profiles derived therein, either from regression [18,34,35,37] or comparison between age groups [36,38], highlight an asynchronous ageing of tissues (discussed in [39]), with some of them more affected by age-related gene expression changes associated with biological mechanisms known to be

impacted by ageing such as mitochondrial activity or metabolic homeostasis. In particular,

tissue-specific periods of major transcriptional changes in the fifth and eighth decades of

human lifespan have been revealed [34], reflecting the so-called digital ageing [39], consistent

with what is observed in mice [17,18]. Furthermore, despite outlining the tissue specificity of

the transcriptomic signatures of human ageing, some synchronisation was found between

tissues like lung, heart and whole blood, which exhibit a co-ageing pattern [35]. Nevertheless,

as each study followed its own specific procedures, from sample collection to data

processing, results from these analyses are hard to compare with one another.

Processed data from those studies have not been made easily accessible and

interpretable to researchers lacking computational proficiency but aiming to use them to

test their novel hypotheses. To fill this void, we have developed voyAGEr, a web application

providing flexible visualisation of comprehensive functional analyses of gene expression

alterations occurring in 49 human tissues with age in each biological sex. We leverage the

large RNA-seq dataset from the Genotype-Tissue Expression (GTEx) project [40], encompassing

tissue samples from hundreds of donors aged from 20 to 70 years, with a pipeline for gene

expression profiling with an optimised temporal resolution. voyAGEr allows to investigate

ageing from two perspectives: (i) gene-centric – how each gene's tissue-specific expression

progresses with age; and (ii) tissue-centric – how tissue-specific transcriptomes change with

age. Additionally, voyAGEr enables the examination of modules of co-expressed genes

altered with age in 4 tissues (brain cortex, skeletal muscle, left ventricle of the heart, whole

blood), namely their enrichment in specific cell types, biological pathways, and association

with diseases. We therefore expect voyAGEr to become a valuable support tool for

researchers aiming to uncover the molecular mechanisms underlying human ageing.

Moreover, being open-source, voyAGEr can be adapted by fellow developers to be used with

alternative datasets (e.g. from other species) or to incorporate other specific functionalities.

voyAGEr is freely available at

https://compbio.imm.medicina.ulisboa.pt/app/voyAGEr.

Results

voyAGEr's interactive exploration of tissue-specific gene expression landscapes in ageing is based on sequential fitting of linear models (v. Materials and Methods) to estimate, for each gene in each tissue:

(i)   the *Age* effect, i.e., how the age-associated changes in gene expression evolve with age itself;

(ii)   the *Sex* effect, i.e., how the differences in gene expression between sexes evolve with age;

(iii)   the *Age&Sex* interaction effect, i.e., how the differences between sexes of age-associated changes in gene expression evolve with age.

We named our approach Shifting Age Range Pipeline for Linear Modelling (ShARP-LM). Briefly, this method consists in performing differential gene expression (with gene expression as a function of the donors' *Age*, *Sex* and *Age&Sex* interaction) in moving age windows spanning 16 years. By considering the percentage of genes altered in each age range, we can highlight age periods of major tissue-specific transcriptomic alterations (**Figure 1**).

Gene-centric analyses of human tissue-specific expression changes across age

The progression of tissue-specific expression of a particular gene across age can be examined in voyAGEr's *Gene* tab. By entering its HGNC symbol in the Gene selector, the user has access to graphical summaries of the gene's tissue-specific expression (sub-tab *Profile*) (**Figure 2A**) and significance of age-related changes in its expression due to the *Age*, *Sex*, and *Age&Sex* effects (sub-tab *Alteration*) (**Figure 2B**) across age. Results can be displayed in a heatmap for all tissues or in a scatter plot for a chosen individual tissue (**Figure 2C**). When the gene is studied in a single tissue, the user can graphically and statistically profile the association of the donors' sex and reported conditions (e.g., history of heart attack or pneumonia) with the gene's expression profile. A table summarizing the donors' metadata is also shown (**Figure 2C**). The user can interactively select donors of interest on the scatter plot and further examine their information in the automatically subsetted table.

An example of a process whose molecular mechanisms are of particular interest to researchers in the ageing field is cellular senescence. Senescence is a stress-induced cell cycle arrest limiting proliferation of potentially oncogenic cells but progressively creating an inflammatory environment in tissues as they age [41,42]. *CDKN2A,* that encodes cell cycle regulatory protein p16[INK4A] known to accumulate in senescent cells [43,44], has its expression increased with age in the vast majority of tissues profiled (**Supplementary Figure 1A**). Similarly, reduced levels of proliferation markers, such as *PCNA* [45] and *MKI67* [46], can be studied as putative markers of ageing of certain tissues. These genes have their expression altered with age in the lung and display a similar expression profile (decreasing from 25 to 30 years old (y.o.), constant between 35 and 50 y.o. and decreasing in older ages) (**Figure 2C**). However, these trends appear to vary according to the donor's history of non-metastatic cancer (**Supplementary Figure 1B**), illustrating voyAGEr's use in helping to find associations between gene expression and age-related diseases.

On a different note, sex biases have been reported in the expression of *SALL1* and *DDX43* in adipose tissue and lung, respectively [10]. voyAGEr allows to not only recapitulate those observations but also assess the temporal window where these changes occur (**Supplementary Figure 2**).

Tissue-specific assessment of gene expression changes across age

Peaks of gene expression alterations

The landscape of global tissue-specific gene expression alteration across age can be examined in voyAGEr's *Tissue* main tab. A heatmap displaying, for all tissues, the statistical significance over age (v. Materials and Methods) of the proportion of genes altered with *Age*, *Sex* or *Age&Sex* (depending on the user's interest) is initially shown (**Figure 3A**), illustrating the aforementioned asynchronous ageing of tissues observed for humans and rodents [18,32–36,39].

The user can then spot the age periods with the most significant gene expression alterations in a selected tissue (**Figure 3B a**), and identify the associated altered genes (**Figure 3B b**). The user can also plot the expression of a given gene of interest across age together with the significance of its expression modification with *Age*, *Sex* or *Age&Sex* (**Figure 3B c**).

An example of voyAGEr's capabilities is illustrated in **Supplementary Figure 3**, showing

substantial transcriptomic alterations in the uterus from the late forties to the early fifties,

overlapping with the age distribution of menopausal onset[47], which could explain the

observed molecular modifications.

It is also possible to visualise tissues with more than one period of transcriptomic changes,

and to individually inspect these periods. As an example, the subcutaneous adipose tissue

appears to go through two main periods of transcriptional changes with age: a major one at

the late 20's (~13 % of altered genes), and a minor one after 45 years (~5% of altered genes)

(**Figure 3B a**). The most altered genes in this first peak appear to have their expression

modified only at this precise age period (e.g. *PRELID1*, *RUNX1T1*, *TUBB4B*, *FGFRL1* and

MALSU1). Similarly, mitochondrial genes (e.g. *MT-CYB*, *MT-ND4*, *MT-ATP6*, *MT-ND2*) (**Figure**

3B b) appear to be the most altered genes in the second peak (**Figure 3B c**). This particularity

suggests that different sets of genes drive the periods of major transcriptional changes,

which begs assessing if they reflect the activation of distinct biological processes.

Gene Set Enrichment

The user can explore the biological functions of the set of genes underlying each peak of

transcriptomic changes by assessing their enrichment in curated pathways from the

Reactome database [48]. voyAGEr performs Gene Set Enrichment Analysis (GSEA) [49] and the

user can visualise heatmaps displaying the evolution over age of the resulting normalised

enrichment score (NES[49], reflecting the degree to which a pathway is over- or under-

represented in a subset of genes) for a given tissue, effect (*Age*, *Sex* or *Age&Sex*) and

Reactome pathway (all, or user-selected) (**Figure 4A a**). To reduce redundancy and facilitate

the understanding of their biological relevance, we clustered those pathways into families

that also include Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways [50] and Gene

Ontology (GO) Biological Processes of level 3 [51]. Briefly, we clustered gene sets from the 3

sources based on the overlap of their genes (v. Materials and Methods), thereby creating

families of highly functionally related pathways. Taking advantage of the complementary

and distinct terminology in Reactome, KEGG and GO, the user can interpret each family's

broad biological function by looking at the word cloud of its most prevalent terms, and

browsing the list of its associated pathways (**Figure 4A b**). For example, although most

pathways enriched in the two aforementioned peaks of altered genes in subcutaneous

adipose tissue were different, there is an overlap of pathways related to metabolism,

including various mitochondrial processes (**Figure 4A**). This highlights the importance of

integrating individual gene data with pathway enrichment analysis to garner more

comprehensive insights on the biological relevance of those changes.

The peaks of transcriptomic changes can also be examined for enrichment in a user-provided

gene set (**Figure 4B**). As expression of senescence-related *CDKN2A* is increased in the

subcutaneous adipose tissue with age (**Supplementary Figure 1A),** we hypothesised that

other senescence-associated genes may be augmented too. Thus we used that voyAGEr

functionality, using the Senequest [42] geneset (supported by at least 4 sources) to test it,

observing no significant alterations (**Figure 4B**).

Modules of co-expressed genes

voyAGEr also allows functional analyses of modules of co-expressed genes (i.e. genes with

highly correlated expression across samples, defined by weighted correlation network

analysis [52]). Genes in the same module are likely to be co-regulated and share biological

functions or associations with phenotypical or pathological traits [53]. Those modules may also

act as markers of core transcriptional features of cellular activity and identity [54].

Concretely, voyAGEr enables the user to visualise how the expression of modules of genes

that are associated with a specific cell type, biological pathway or disease progresses over

age in a specific tissue. After selecting one tissue of interest, the user has, for each module,

access to four levels of information:

(i)   *Expression*: its eigengene expression progression over age (**Figure 5C**);

(ii)   *Cell types*: its enrichment in specific cell types, based on cell type signatures

found in the literature (**Figure 5C**);

(iii) *Pathways*: its enrichment in Reactome pathways;

(iv) *Diseases*: its enrichment in disease markers, based on gene-disease associations

from DisGeNET [55,56], calculated with both the `disgenet2r` package [55] and with

Fisher's tests (**Figure 5B**).

By default, for each tissue, results are displayed in the form of heatmaps of expression

(centred and scaled) illustrating how all modules evolve with age (**Figure 5A**). The user also

has the possibility to select a module of interest and see its eigengene progression over age

in a scatter plot (**Figure 5C**), lists of its association with diseases and pathways ordered by

significance, and a TreeMap for its cell type enrichments (**Figure 5C**). In the example of

Figure 5C, the "MEblue" module, comprising 241 genes co-expressed in the brain cortex,

shows significant enrichment in astrocyte markers. The apparent increase of this module's

expression with age may reflect the known age-related changes in astrocyte activation [57] and

the relative weakening of neuronal activity ("MEyellow" and "MEpink" modules).

As in the *Gene* tab, the user can separate donors based on their sex and associated medical

conditions in the scatter plot of the eigengene expression progression. On the Pathways and

Diseases-Manual tabs below the main plot, the user can also visualise the contingency table

from that specific disease/pathway, on the corresponding column.

Discussion

voyAGEr provides a framework to examine the progression of gene expression over age in

several human tissues, serving as a valuable resource for the ageing research community. In

particular, it helps to identify tissue-specific age periods of major transcriptomic alterations.

The results of our analyses show the complexity of human biological ageing by stressing its

tissue specificity [34] and non-linear transcriptional progression throughout lifetime, consistent

with previous results from both proteomic [58] and transcriptomic [28,34] analyses. By revealing

and annotating the age-specific transcriptional trends in each tissue, voyAGEr aims at

assisting researchers in deciphering the cellular and molecular mechanisms associated with

the age-related physiological decline across the human body.

Due to the tissue-specific nature of the pre-processing steps (v. Read count data pre-

processing in the Methods section), and given that most of the plotted gene expression

distributions are centred and scaled by tissue, it is important to note that voyAGEr may not

be always suited for direct comparisons between different tissues. For instance, it does not

allow to directly ascertain if a gene exhibits different expression levels in different tissues or

if the expression of a particular gene in one tissue changes more drastically with age than in

another tissue. Furthermore, we must emphasise that the majority of GTEx donors

contributed samples to multiple tissues (**Supplementary Figure 4A**), potentially introducing

biases and confounders when comparing gene expression patterns between tissues. Our

analyses of variance (**Supplementary Figure 4B**) and downsampling to control for common

donors (**Supplementary Figures 4C-E**) suggest very limited global confounding between the

impacts of donor and age on gene expression and that any potential cross-tissue bias not to

depend much on the proportion of common donors (**Supplementary Figure 4E**). However,

this effect must be taken into account when comparing specific pairs of tissues (e.g., Colon –

Transverse and Whole Blood, **Supplementary Figure 4D**).

Additionally, voyAGEr allows to scrutinise and visually display the tissue-specific differences

in gene expression between biological sexes across age. Biological sex is an important factor

in the prevalence of ageing-associated diseases, as well as their age of onset, progression

[6,7,59] and sex-related differences in gene expression[8–11]. By profiling the age distribution of

such differences, voyAGEr can lead to a better understanding of their influence in the

aetiology of the sex specificities of human ageing. For instance, we were able to corroborate

findings on the sex-differential transcriptome of adult humans by Gershoni et al. [9], with

voyAGEr emphasising its tissue-specificity and allowing to discriminate the ages at which

sex-related biases appear to be more prevalent (**Supplementary Figure 5**).

Nonetheless, it is essential to interpret sex chromosome-specific gene results in voyAGEr

with caution. For instance, we observe elevated expression of Y-chromosome-specific *DDX3Y*

in males, whilst its female expression (expected to be zero) is very low, in the range of what

can be considered background noise (**Supplementary Figure 6**). Its age-related alterations

exhibit a distinctive peak around the age of 40 apparently driven by subtle changes of gene

expression in female samples, illustrating the need for the abovementioned caution.

One of the limitations of voyAGEr is that most GTEx tissue donors had health conditions and

their frequency increased with age, preventing us from defining a class of healthy individuals

and identifying age-associated transcriptomic changes that could be more confidently

proposed to happen independently of any disease progression. Whilst the large sample sizes

and inherent biological variability among individuals, reflected in the diversity of condition

combinations, are expected to mitigate significant confounding effects, voyAGEr also allows

users to evaluate how tissue-specific gene expression trends vary according to the donors'

diverse conditions (see **Supplementary Figure 1B**).

The development of voyAGEr was accompanied by that of a pipeline, ShARP-LM, that

facilitates the holistic depiction of the transcriptional landscapes of adult human tissues

throughout ageing with a yearly age resolution. We take advantage of the

comprehensiveness of the transcriptome collection from human tissues from the GTEx

project to make our analyses a valid surrogate of a currently undoable longitudinal study. It

confers our method enough statistical robustness to mitigate the inter-individual differences

and deal with the non-uniform distribution of the donors' ages. Nevertheless, it is worth

highlighting that the age distribution of donors does impact the statistical power for

detecting transcriptional changes. Consequently, we are more likely to identify significant

alterations (with p-value < 0.05 in our gene-centric analyses) within age ranges that are

more prevalent in our sample population, often characterised by older individuals

(**Supplementary Figure 7)**. When downsampling to ensure a balanced age distribution, a loss

of statistical power is apparent but a considerable positive correlation with the original

results is maintained and a substantial number of significant alterations remain so

(**Supplementary Figure 8**). This limitation is likely to be overcome by the accumulation of

transcriptomes of human tissues in public databases, promising a gradual increase in

accuracy and age resolution with which human transcriptomic ageing can be profiled.

Similarly, the expanding collection of single-cell transcriptomes in public databases is

yielding improved gene markers for an increasing diversity of human cell types, enhancing

the usefulness of leveraging bulk transcriptomes to study the impact of ageing on the

cellular composition of human tissues, for which the co-expression module approach in

voyAGEr provides a proof-of-concept.

Nonetheless, it is imperative to approach the module-based analysis with caution, as direct

and literal interpretations may be misleading. For instance, it is not uncommon to observe

an enrichment of "Rheumatoid Arthritis" in modules associated with various immune cell

types in anatomical locations, such as the brain cortex, where the disease does not directly

manifest. If a specific module associated with a condition like "Liver Cirrhosis" exhibits an

age-related increase in the brain cortex, of course this does not mean such disease ever

occurs within older brains. Nevertheless, we consider that the module-based approach can

serve as a valuable resource for generating hypotheses.

Given its open-source nature, voyAGEr is envisaged to be a continually evolving resource,

able to accommodate new data and expand its functionalities, namely by incorporating

additional tissues into the modules section and integrating perturbagen data for inference of

molecular causes underlying observed gene expression alterations and small molecules to

target them for therapeutic purposes [60,61].

As an *in silico* approach with no experimental validation for its results, voyAGEr is meant to

be a discovery tool, supporting biologists in the exploration of a large transcriptomic dataset,

thereby generating, refining or preliminary testing hypotheses, laying the groundwork for

subsequent experimental research. It can be an entry point for projects aiming at better

understanding the tissue- and sex-specific transcriptional alterations underlying human

ageing, to be followed by targeted studies focusing on the functional roles of the most

promising markers identified therein in the physiology of ageing. Those marker genes can

contribute to the development of more robust and cross-tissue gene signatures of ageing [62]

and the expansion of age-related gene databases [63,64].

Moreover, the observed diverse and asynchronous changes in gene expression between

tissues over the human adult life provides potentially relevant information for the design of

accurate diagnostic tools and personalised therapies. On one hand, identifying those

changes' association with specific disorders could have a prognostic value by enabling the

identification of their onset before clinical symptoms manifest [36]. On the other hand,

computational screening of databases of genetic and pharmacologically induced human

transcriptomic changes could help to infer their molecular causes and uncover candidate

drugs to delay these effects [60,61,65,66].

Methods

Development platforms

Data analysis was performed in R (version 4.1.2) and the application developed with R package `Shiny` [67]. voyAGEr's outputs are plots and tables, generated with R packages `highcharter` [68] and `DT` [69], respectively, that can easily be downloaded in standard formats (png, jpeg, and pdf for the plots; xls and csv for the tables).

voyAGEr was deployed using Docker Compose and ShinyProxy 2.6.1 in a Linux virtual machine (64GB RAM, 16 CPU threads and 200GB SSD) running in our institutional computing cluster.

Read count data pre-processing

The matrix with the RNA-seq read counts for each gene in each GTEx v8 sample was downloaded from the project's data portal (https://www.gtexportal.org/) [40]. From the 54 tissues available from GTEx v8, five were discarded (kidney medulla, fallopian tube, bladder, ectocervix, endocervix) due to low (<50) numbers of samples.

Read count data for each tissue were then pre-processed separately. We started by filtering out genes deemed uninformative due to their very low expression across samples: only genes with at least 1 CPM in at least 40% of the samples were kept for analysis (the number of genes analysed for each tissue can be found in **Supplementary Table 1**)[70]. Read counts for those kept genes were used to calculate normalisation factors to scale the raw library sizes, using function `calcNormFactors` from `edgeR` [70] that implements the trimmed mean of M-values [71]. Read counts were subsequently normalised and log-transformed with the `voom` function [72] from package `limma` [73].

However, it is well-established that batch effects, which may stem from variations in sample treatment prior to RNA-seq library preparation, can introduce spurious gene expression differences between samples and result in confounding factors[74]. We therefore conducted an impartial and systematic search for potential batch effects. Firstly, we performed a principal component analysis of gene expression for each tissue, using the `prcomp` package.

We quantified the relation between each condition associated with every sample[1] and the

first two principal components, by computing Spearman correlations (for numerical

conditions), t-tests (for binary categorical conditions), or analysis of variance (ANOVA) tests

(for variables with more than two possible values, and, in the case of numerical variables,

fewer than 15 unique values). Conditions that surpassed defined empirical thresholds (p-

value < 0.05, Spearman correlation > 0.3, t-test > 10, and ANOVA > 20) were flagged as

potential batch effects. Except in brain tissues, the COHORT variable (i.e., whether the

samples were collected from organ donors or *post mortem*) appeared to be the main batch

effect, with ripple effects on numerous other related conditions (**Supplementary Figure 9**).

Moreover, SMRIN (sample's RNA integrity number), DTHHRDY (death classification based on

the 4-point Hardy Scale), and MHSMKYRS (smoke years) consistently emerged as conditions

associated with the primary axes of variance. The number of genes detected in each sample,

determined by the filtration step described above, was also identified as a significant

contributor to the primary data variance. We therefore corrected for these five conditions,

on a tissue-by-tissue basis, by adapting the `removeBatchEffect` function from the

`limma` package [73]. Specifically, we employed linear models to estimate the contributions to

gene expression of each of those factors and subtracted such contributions from the original

logCPM matrix. To ensure biological interpretability of the results, we offset the resulting

values to the minimum value in the non-corrected matrix. To prevent sample loss due to

missing values for the aforementioned five conditions and since the number of missing

values was relatively low compared to the total number of samples, imputation was carried

out using the `mice` package[75].

The resulting matrix of logCPM-corrected values was used for all downstream analyses. As an

illustrative example of the importance of batch removal, the expression of surfactant factor

SFTPA2 was found to be associated with donors on a ventilator[76]. Without batch correction,

SFTPA2 expression would have been associated with age due to higher prevalence of such

cases among older individuals (**Supplementary Figure 10**).

---

[1] According to the annotated variables for the dbGaP Study Accession phs000424.v8.p2, listed in
https://www.ncbi.nlm.nih.gov/projects/gap/cgi-
bin/GetListOfAllObjects.cgi?study_id=phs000424.v8.p2&object_type=variable

ShARP-LM

To model the changes in gene expression with age, we developed the Shifting Age Range Pipeline for Linear Modelling (ShARP-LM). For each tissue, we fitted linear models to the gene expression of samples from donors with ages within windows with a range of 16 years shifted through consecutive years of age (i.e. in a sliding window with window size = 16 and step size = 1 years of age). This was the minimum age span needed to guarantee the presence of more than one sample per window, across all considered tissues. As samples at the ends of the dataset's age range would be thereby involved in fewer linear models, we made the window size gradually increase from 11 to 16 years when starting from the "youngest" samples and decrease from 16 to 11 years when reaching the "oldest" (**Supplementary Figure 11)**.

Function `lm` from `limma` was used to fit the following linear model for gene expression (GE):

$$GE \sim E_0 + \alpha.Age + \beta.Sex + \chi.Age\&Sex + \varepsilon$$

For each gene, α, β and χ are the coefficients to be estimated for their respective hypothesised effects. For each sample, *Age* in years and *Sex* in binary were centred and *Age&Sex* interaction given by their product. The coefficients E_0 and ε are thus the expression of the average sample (i.e., with average sex and average age) and the error term. For each gene in each model (i.e., each age window in each tissue), we retrieved the t-statistics of differential expression associated with the three relevant variables and their respective p-values. We considered the average age of the samples' donors within the age window as the representative age of the observed expression changes.

In summary, for a given tissue and variable (*Age*, *Sex* and *Age&Sex*), ShARP-LM yields t-statistics and p-values over age for all genes, reflecting the magnitude and significance of the changes in their expression throughout adult life.

Gene-centric visualisation of tissue-specific expression changes across age

For visualisation purposes, the trend of each gene's expression progression over age in each tissue was derived through Local Polynomial Regression Fitting, using R function `loess` on logCPM values (**Figures 1, 2C, 3B c**). For summarizing in a heatmap a given gene's expression across age in multiple tissues (**Figure 2A**) or the expression of several genes across age in a

426   specific tissue, each gene's regression values are centred and scaled, using R function

427   `scale`.

428   For summarizing in a heatmap the significance of a given gene's expression changes over age

429   in multiple tissues (**Figure 2B**), cubic smoothing splines are fitted to -$\log_{10}(p)$, with p being

430   the t-statistic's p-value, with R function `smooth.spline`.

Tissue-specific quantification of global transcriptomic alterations across age

To assess the global transcriptomic impact of each of the three modelled effects in each

tissue across age, we analysed the progression over age (i.e. over consecutive linear models)

of the percentage of genes whose expression is significantly altered (t-statistic's p-value ≤

0.01) by each effect (**Figure 3B**). To evaluate the significance of each percentage and assess

if high percentages can be confidently associated with major transcriptomic alterations, we

controlled for their false discovery rate (FDR) by randomly permuting the samples' ages and

sexes within each age window fifty thousand times and performing ShARP-LM on each

randomised dataset. We were then able to associate an FDR to each percentage of altered

genes by comparing it with the distribution of those randomly generated (**Figure 3B a**).

We also applied a linear model across the entire age range, thereby providing users with

more insight and supporting evidence into how a specific gene changes with age. For

visualisation purposes, we incorporated a dashed orange line, with the logFC per year for the

Age effect as slope, in the respective scatter plots (**Figure 3B c**). We depict the Sex effect

therein by prominent dots on the average samples, with pink and blue denoting females and

males, respectively.

Gene Set Enrichment Analysis (GSEA)

For each *Peak* of significant gene expression modifications, we performed GSEA [49] on the

ordered (from the most positive to the most negative) t-statistics of differential expression

for the respective tissue and age, using R package `fgsea` [77] and the `Reactome` database [48].

We extracted the GSEA normalised enrichment score (NES), which represents the degree to

which a certain gene set is overrepresented at the extreme ends of the ranked list of genes.

A positive NES corresponds to the gene set's overrepresentation amongst up-regulated

genes within the age window, whereas a negative NES signifies its overrepresentation

amongst down-regulated genes. The NES for each pathway was used in subsequent analyses

as a metric of its up- or down-regulation in the Peak. The resulting normalised enrichment

score (NES) for each pathway was used in subsequent analyses as a metric of its over- or

down-representation in the *Peak*.

To optimise computational efficiency and minimise redundancy in the analysed pathways,

we only considered pathways containing a minimum of 15 genes and up to 500 genes, as

suggested in the `GSEA` User Guide[78]. For the sake of clarity in voyAGEr's visual

representations, we only included pathways with a p-adjusted value less than or equal to

0.05, further narrowing it down to pathways within the top 1% of p-adjusted values.

Additionally, we exclusively featured pathways with at least one significant age Peak (FDR <=

0.05), as illustrated in **Figure 4A**.

Families of pathways

To reduce pathway redundancy and facilitate the assessment of their biological relevance in

results' interpretation, we created an unifying representation of pathways from `Reactome`

[48] ,`KEGG` [50], and level 3 Biological Processes from `GO` [51], by adapting a published pathway

clustering approach [79] to integrate them into families.

The approach relies on the definition of a hierarchy of pathways based on the numbers of

genes they have in common. For each two pathways P_i and P_j, respectively containing sets of

genes G_i and G_j, we computed their overlap index (OI) [79], defined as follows:

$$OI_{ij} = |G_i \cap G_j| / min(|G_i|, |G_j|)$$

Where $|G_i|$ is the number of genes in set G_i and $|G_i \cap G_j|$ is the number of genes in common

between G_i and G_j. $OI_{ij} = 1$ would therefore indicate that P_i and P_j are identical in gene

composition or that one is a subset of the other. On the contrary, $OI_{ij} = 0$ would mean that P_i

and P_j have no genes in common. To ease the computational work, we removed from the

analysis pathways that are subsets of larger pathways (i.e. each pathway whose genes are all

present in another pathway).

From the OI_{ij} matrix, from which each row is a vector with the gene overlaps of pathway *i*

with each of the pathways, we computed the matrix of Pearson's correlation between all

pathways' overlap indexes with R function `cor`. That matrix was finally transformed into

Euclidean distances with R function `dist`, allowing for pathways to be subsequently

clustered with the complete linkage method with R function `hclust`. The final dendrogram
was empirically cut in 10 clusters (**Supplementary Figure 12**). Pathways that were initially
excluded from the computation for being subsets of others were added to the clusters of
their respective *parent* pathways. Each *daughter* pathway with more than one parent was
assigned to the cluster of the parent with the smallest number of genes, thereby maximizing
the daughter-parent similarity. The `data.table` R package, for fast handling of large
matrices, was used in this analysis [80].
495
Gene co-expression modules
Gene co-expression modules were defined with R package `WGCNA`[52]. For each considered
tissue, the process began with the identification of a set of informative genes that exhibit
high variability across samples (referred to as variable A), thus improving module definition.
Next, we calculated the bicorrelation matrix for the expression of all selected genes with the
`bicor` function. We then applied soft thresholding by raising all correlation values to the
power of β, accentuating stronger correlations. The value of β = 12 was chosen in
accordance to the WGCNA FAQs[81], and after confirmation of a free-scale topology using the
`pickSoftThreshold` function. We generated the dissimilarity matrix by subtracting the
output of the `TOMsimilarity` function from 1. Gene co-expression modules were then
defined using a static tree-cutting algorithm, implemented via the `cutreeStaticColor`
function, requiring as parameters a minimum number of genes per module (referred to as
variable B) and the tree-cutting height (variable C).
The empirical determination of parameters A to C was guided by the following principles: (i)
maximizing cell type signature enrichment, (ii) minimizing the number of genes per module,
and (iii) ensuring that the modules' eigengenes exhibited age-related variability. Different
combinations of these variables were exhaustively tested. To maintain biological relevance,
modules consisting of non-assigned genes or genes lacking substantial supporting evidence,
such as pseudogenes, were excluded.
Maximizing cell type enrichment in the modules, with a focus on known markers for specific
cell types, has previously been proven successful in unveiling core transcriptional features of
cell types in the human central nervous system [53]. Cell type enrichment analysis relied on
Fisher tests, providing odds enrichment scores and significance values (p-values). This
involved comparing module genes with the signature, considering as background of genes

those included in the module definition for each tissue. We prioritised modules of genes for

which we obtained at least one significant result for each cell type (odds ratio (OR) > 1 and

p-value < 0.05). The cell type signatures employed in this analysis were sourced from

MSigDB's C8 collection[49], and in the case of Whole Blood, additionally from LM22[82].

Specific variance thresholds (variable A) were employed: 0.5, 0.4, 0.35, and 0.9 for Brain –

Cortex, Muscle – Skeletal, Heart – Left Ventricle, and Whole Blood, respectively. The

minimum number of genes per module (variable B) was set at 15, 20, 20, and 15,

respectively. Tree-cutting heights (variable C) of 0.95, 0.98, 0.99, and 0.97 were respectively

used.

Each module is characterised by a set of genes and an eigengene, represented by the first

principal component obtained through singular-value decomposition of the module's gene

expressions. Subsequently, voyAGEr facilitates an evaluation of cell type enrichment, as

described earlier, and enrichment in biological pathways and diseases. The enrichment of

modules in cell types, Reactome pathways, and diseases (extracted from DisGeNET

[55,56]database version 7.0) was quantified using Fisher's tests. For disease enrichment, the

function `disease_enrichment` from the `disgenet2r` package[55] was employed,

utilising their curated set of diseases. The significance of these enrichments was determined

through p-value/FDR adjustment using Benjamini-Hochberg correction. For visual clarity,

only pathways and diseases displaying significant enrichment (p <= 0.05) in at least one

module were considered.

The four tissues (Brain - Cortex, Muscle - Skeletal, Heart - Left Ventricle, and Whole Blood)

covered by the Module section of voyAGEr were selected due to their relatively high sample

sizes and availability of comprehensive cell type signatures. The increasing availability of

human tissue scRNA-seq datasets (e.g., through the Human Cell Atlas[83]) will allow future

updates of voyAGEr to encompass a wider range of tissues.

Data and Code availability

Processed GTEx v8 RNA-seq data (read count tables) were downloaded from the project's

data portal (https://www.gtexportal.org/). Donor metadata were obtained from

dbGaP - database of Genotypes and Phenotypes (Accession phs000424.v9.p2 project ID

13661). voyAGEr's output tables can be directly downloaded in standard xls and csv formats.

551 The complete source code for voyAGEr, including pre-processing and Shiny app, can be

552 accessed on GitHub at the following repository:

553 https://github.com/DiseaseTranscriptomicsLab/voyAGEr.

Acknowledgements

We thank iMM colleagues Joana Neves and Luísa Lopes, as well as all members from the

Disease Transcriptomics Lab, for providing valuable feedback on the manuscript. We also

thank the very knowledgeable anonymous reviewers designated by *eLife* for their

constructive and insightful suggestions and criticisms to the first version of this manuscript;

their feedback greatly contributed to much improved versions of both the article and

voyAGEr itself.

Funding: This work was supported by European Molecular Biology Organization (EMBO

Installation Grant 3057), FCT - Fundação para a Ciência e a Tecnologia I.P. (FCT Investigator

Starting Grant IF/00595/2014, CEEC Individual Assistant Researcher contract

CEECIND/00436/2018, PhD Studentships SFRH/BD/131312/2017, UI/BD/153368/2022 and

UI/BD/153363/2022, projects LA/P/0082/2020 and UIDP/50005/2020), European Union (EU)

Horizon 2020 Research and Innovation Programme (RiboMed 857119), and GenomePT

project (POCI-01-0145-FEDER-022184), supported by COMPETE 2020 – Operational

Programme for Competitiveness and Internationalization (POCI), Lisboa Portugal Regional

Operational Programme (Lisboa2020), Algarve Portugal Regional Operational Programme

(CRESC Algarve2020), under the PORTUGAL 2020 Partnership Agreement, through the

European Regional Development Fund (ERDF), and by FCT - Fundação para a Ciência e a

Tecnologia.

Authors' contributions

576 A.S. and N.L.B.-M. designed the study, with contributions from R.M.-S. and A.K.; A.S., R.M.-S,

577 and A.K. performed all the development and bioinformatics work, under the supervision of

578 N.L.B.-M.; N.S.-A. set up the web and database implementation; A.S., R.M.-S., A.K., and

579 N.L.B.-M. wrote the manuscript, with contributions from N.S.-A..

Conflict of Interest statement

The authors declare no conflict of interest.

 Figures and Tables

Figure 1: *voyAGEr profiles tissue-specific age-associated changes in gene expression and their differences between sexes.*

For each of the 49 human tissues in GTEx, gene expression was linearly modelled in windows spanning 16 years centred in consecutive years of age, to estimate the effects thereon of *Age*, *Sex* and the interaction between them, i.e., how the *Sex* effect changes with age, equivalent to how the *Age* effect differs between sexes (v. Materials and Methods). In each age window, the percentage of genes with expression significantly altered by of each of those effects gives their respective transcriptomic impact (upper panels). voyAGEr thereby identifies the age periods at which major gene expression changes occur in each tissue. For example, in coronary artery: major age-related transcriptional alterations are found at around 60 years of age (upper left panel), illustrated by the behaviour of *COX11* (bottom left panel); major gene expression differences between males and females happen across the considered age range (upper centre panel), as illustrated by *CD99* (bottom centre panel); major differences between sexes in age-related gene expression alterations happen across the considered age range (upper right panel), as illustrated by *AKT2* (bottom right panel). Solid loess lines in the bottom panels (green for all donors, pink for females, blue for males). Gene expression (GE) in $\log_2$ of counts per million (logCPM).

Figure 2: *Gene-centric analyses of expression alterations across age*

A - Heatmap of *MKI67* expression across tissues over age. **B** - Heatmap of significance of *Age*-associated *MKI67* expression alterations over age. p-values are for the moderated t-statistics of differential gene expression associated with the Age effect (v. the ShARP-LM approach in Materials and Methods). Notably, transcriptional changes are observed in lung (mid 20's, early 30's and after 55). **C** - voyAGEr's *Gene* tab interface. *MKI67* expression in lung is inspected. Donors' information is shown in a table and the scatter plot can be interactively adjusted according to the donors' condition of interest **(Supplementary Figure 1B)**.

Figure 3: *Tissue-specific assessment of gene expression changes across age*

A - Heatmap of significance (FDR based on random permutations of age, v. Materials and

Methods) over age of the proportion of genes with expression significantly altered with *Age*

in the 49 analysed tissues. **B** - Exploration of gene expression changes across age in

Subcutaneous Adipose tissue: (a) Percentage of genes with significantly altered expression

with *Age* over age. Two main peaks of transcriptional changes are noteworthy, a major one

at the late 20's and a minor one after 45; (b) Clicking on the dot of a specific age (29.57 y.o.

in plot *a*) gives access to the list of the most altered genes at that age, ordered by statistical

significance of expression changes (p-value of moderated t-statistic). (c) Plot of expression of

the chosen top gene in the table in *b* across age (bottom) in parallel with the significance of

its expression alterations with *Age*. The expression of *LMO3* significantly increases at around

30 years old, concomitantly with the first peak of transcriptomic changes with *Age*.

Figure 4: *Tissue-specific assessment of pathway expression changes across age in the human*

Subcutaneous Adipose tissue

A - Heatmap depicting the normalised enrichment scores (NES) of Reactome pathways

associated with specific tissues and effects. Pathways are classified into 10 families (a),

which can be characterised by their frequently occurring terms (b), providing insights into

their biological functions. Only pathways significantly linked to gene expression changes in at

least one age window (FDR ≤ 0.01) are displayed. Black squares indicate the two age periods

with prominent transcriptional changes, while yellow squares denote pathways common to

both peaks, primarily belonging to family 2. Word cloud analysis (b) reveals that family 2

pathways are mainly related to metabolism. **B** - Enrichment of a user-provided gene set,

given by the significance of Fisher's tests, in genes altered with *Age* throughout ageing

(based on a user-defined p-value threshold). Here, the given gene set is composed of genes

from Senequest [42] whose link with senescence is supported by at least 4 sources. In this

case, there are no significant peaks.

Figure 5: *Tissue-specific assessment of age-associated progression of modules of co-expressed genes*

A - Heatmap of eigengene expression for all the modules of co-expressed genes in the brain cortex over age. **B** - Heatmap of association of the modules with four selected diseases, computed with the disgenet2r package[55]. **C** - Scatter plot (above) of eigengene expression over age, in all brain cortex samples, for a selected module of 241 genes (MEblue). The eigengene expression is derived from the 1st component of the single value decomposition of its genes' expression. This module was analysed, with Fisher's tests, for enrichment in cell types, based on markers from the literature (Fan[84] and Descartes[85]) and found to be associated with astrocytes, as can be observed by the TreeMap below (where each rectangle's area and darkness are proportional to the significance of its association with a cell type and its colour linked to the markers' source study).

Supplementary Figures and Tables

Supplementary Figure 1: *voyAGEr's Gene tab interface*.

A - Heatmap of *CDKN2A* expression across tissues over age. **B** - *PCNA* expression in lung over age and its association with the donor's history of non-metastatic cancer.

Supplementary Figure 2: *Sex-specific SALL1 and DDX43 expression alterations over age.*

Scatter plots of *SALL1* (**A**) and *DDX43* (**B**) expression over age (above), in adipose tissue and

lung, respectively, coloured based on the donor's biological sex (male in blue and female in

pink). The scatter plots of gene expression are shown together with line plots of the

significance of their difference between sexes (below).

Supplementary Figure 3: *Transcriptomic alterations in the uterus coincident with the onset of*

menopause.

voyAGEr enables the identification of a prominent age period of transcriptional alterations in

the uterus, which coincides with the onset of menopause[47] The genes exhibiting significant

changes (**A**) reflect the transformations observed in biological processes (**B**), marked by a

distinct switch in enrichment scores. Positively enriched pathways become negative, and

vice-versa during this age frame.

Supplementary Figure 4: *Impact of donor overlap between tissues in age-associated trends*

A - Heatmap depicting the percentage of common donors between pairs of tissues. A given

square illustrates the percentage of all samples of tissue in the x axis (Tissue *1*) that is in

common with the tissue in the y axis (Tissue *2*). B - Assessment of the relative contributions

of different sources to the dataset's variance. (Left panel) Tissue accounts for approximately

90% of the total variance, while donor contributes around 10%; age has a minimal impact

(1%), likely due to the relative subtlety of its effects on gene expression and to the tissue

specificity of ageing dynamics. (Right panel) Removal of the donor variable does not transfer

variance to age, suggesting limited confounding between the two variables. **C** - Impact of the

relative proportion of common donors on gene expression correlation between tissue pairs.

Panels A, B, and C showcase the tissue pairs with the highest (Muscle Skeletal / Kidney

Cortex), median (Pancreas / Heart Left Ventricle), and lowest (Small Intestine / Brain

Amygdala) percentages of common donors, respectively. The left panels illustrate gene-by-

gene Pearson's correlations of gene expression between the two tissues, comparing the

scenarios with (x-axis) and without (y-axis) the removal of common donors. The right panels

depict the same comparisons, but with random downsampling (y-axis) in both tissues based

on the proportions defined for common donor removal. The depicted examples show that

the outcomes are comparable when removing common donors or employing random

downsampling. **D** - Comparison of the impacts of removing common donor samples and

random downsampling across tissue pairs. The process of removing common donors

involved the identification and removal of samples from shared donors while maintaining

the original sample size imbalance between tissues. As this process inherently involves

downsampling, which may impact results, we performed additional downsampling by

randomly removing samples from both tissues according to the proportions defined for the

removal of common donors. The heatmap is coloured based on whether the removal of

common donors has a greater (red) or lesser impact (blue) than random downsampling. The

values depicted in the heatmap, denoted as the Impact of Common Donors (ICD), are

computed for each tissue pair. This calculation involves several steps: first, by determining

the absolute difference in Pearson's correlation for each gene's mean expression within

each age window from the ShARP-LM pipeline, between the original data and the subset of

data without common donors (DiffWoCD) or with random downsampling (DiffRD).

Subsequently, the medians of DiffWoCD and DiffRD are computed, and the difference

between these median values provides the ICD for each tissue pair. Due to the unidirectional

nature of correlation (i.e., the results for tissue 1 vs tissue 2 mirror those for tissue 2 vs

tissue 1), the resulting matrix is triangular in form. Grey tiles denote NA values, i.e., where

the tissue-tissue comparison does not have a meaning, namely self-self and between sex-

specific tissues. Top right insert: density line ("smoothed" histogram) of all ICD values. **E** -

Scatter plot relating the Impact of Common Donors (ICD, values in heatmap D) with the

respective percentage of common donors (values in heatmap A).

Supplementary Figure 5: *Tissue-specific sex-differentiated expression.*

A - Heatmap of significance over age, for each tissue, of the proportion of altered genes

between sexes (left), in parallel with the number of sex-differentiated genes highlighted by

Gershoni et al. (v. Figure 1 in [9]) (right). Mammary is corroborated as the tissue with the most

sex-differentiated gene expression. For tissues like Adipose - Subcutaneous, Skin - Not Sun

Exposed, Thyroid, Nerve - Tibial, Muscle - Skeletal, and Spleen, sex-related biases in

expression mainly arise in late life. **B** - Scatter plot relating, for each tissue (each dot), the

number of sex-differentiated genes found by Gershoni et al. with the number of genes found

by voyAGEr to be altered (p-value <=0.05) between sexes in at least 25 age intervals (left).

This correlation is significant (p-value <=0.05) regardless of the considered minimum number

of age intervals a gene must be altered in (right).

Supplementary Figure 6: *Impact of sex-specific genes in the interpretation of voyAGEr*

results.

Representation of the expression of Y-chromosome-specific gene *DDX3Y*, stratified by sex

(**A**), along with the corresponding significance indicating age-related alterations (**B**). Despite

the non-zero expression values for female samples, they fall within a range deemed of

background noise (-2 to -8 logCPM). Notably, when assessing the significance of age-related

changes in the gene, a distinct peak emerges around the age of 40. This peak is spurious, as

it primarily mirrors changes of gene expression in females.

Supplementary Figure 7: *Distributions of GTEx donors' ages and its impact on the statistical power of detection of differential gene expression.*

For thyroid as an example tissue, the sex-differentiated distribution of GTEx samples' ages (v8) (above) are shown together with the percentage of altered genes over age (below) for the three variables (*Age*, *Sex* and *Age&Sex*), as well as their statistical significance, given by the ShARP-LM approach. The statistical power for detecting transcriptional changes is influenced by the age distribution of donors, leading to a higher likelihood of identifying significant alterations (FDR <= 0.05) in age ranges that are more prevalent in the sample population, typically associated with older individuals.

A

B

Supplementary Figure 8: *Effect of downsampling in ShARP-LM results*.

Downsampling was performed by reducing the number of samples in a given window to the

minimum available sample size within all age windows for a given tissue. **A** – Per tissue violin

plots of gene-wide distributions of Pearson's correlation coefficients between original and

downsampled logFC values for the Age variable across age windows, with tissues coloured by

and ordered by increasing percentage of downsampling-associated reduction in the number

of samples. **B** – Density scatter plots of comparison of associated original and downsampled

p-values for each tissue, coloured by the downsampling percentage in each age window,

highlighting the low range of p-values (from 0 to 0.1). Despite changes in logFC with

downsampling, a considerable correlation in significance is maintained, although

downsampling naturally results in a loss of statistical power, evident by the shift of points

towards the first quadrant (dashed lines: p-value = 0.05).

Supplementary Figure 9: *Batch effect correction applied to lung samples.*

(**Top**) Before batch effect correction, 17.7% of the variance between samples appears to be mainly explained by the COHORT variable. (**Bottom**) After batch effect correction, PC1 no longer appears to be explained by the COHORT variable, with its weights haveing a similar distribution between batches.

SFTPA2 (Lung) - AGE

Supplementary Figure 10: *Effects of batch effect correction on gene-centric (SFTPA2) analysis.*

A - Before batch effect correction, *SFTPA2* expression diminishes throughout ageing. This

change is particularly significant after around 55 years of age. **B** - After correction,

expression of this gene shows no significant alterations across age ranges.

Supplementary Figure 11: *Principle of the ShARP-LM method.*

ShARP-LM involves fitting linear models to gene expression data for each tissue, with

samples from donors grouped into age windows of 16 years each, shifted consecutively

through individual years of age using a sliding window approach. As samples at the ends of

the dataset's age range would thereby be involved in fewer linear models, we made the

window size gradually increase from 11 to 16 years when starting from the "youngest"

samples and decrease from 16 to 11 years when reaching the "oldest". The example shows

the expression of an arbitrary gene over age, with 8 age windows shown. Coloured

rectangles represent the age windows and coloured lines the linear modelling of the gene

expression over age in each of them.

Supplementary Figure 12: *Clustering of pathways from Reactome and KEGG, and level 3*

Gene Ontology Biological Processes.

Based on their common genes (v. Materials and Methods), pathways from curated

databases were clustered in 10 families.

Tissue	#Genes
Adipose - Subcutaneous	15519
Adipose - Visceral (Omentum)	15623
Adrenal Gland	15332
Artery - Aorta	15172
Artery - Coronary	15517
Artery - Tibial	14714
Brain - Amygdala	15816
Brain - Anterior cingulate cortex (BA24)	15867
Brain - Caudate (basal ganglia)	16136
Brain - Cerebellar Hemisphere	16207
Brain - Cerebellum	16473
Brain - Cortex	16178
Brain - Frontal Cortex (BA9)	16020
Brain - Hippocampus	15910
Brain - Hypothalamus	16412
Brain - Nucleus accumbens (basal ganglia)	16153
Brain - Putamen (basal ganglia)	15768
Brain - Spinal cord (cervical c-1)	15838
Brain - Substantia nigra	15796
Breast - Mammary Tissue	16311
Cells - Cultured fibroblasts	13737
Cells - EBV-transformed lymphocytes	13743
Colon - Sigmoid	15635
Colon - Transverse	16491
Esophagus - Gastroesophageal Junction	15459
Esophagus - Mucosa	15509
Esophagus - Muscularis	15355
Heart - Atrial Appendage	15003
Heart - Left Ventricle	14036
Kidney - Cortex	16054
Liver	14376
Lung	16616
Minor Salivary Gland	16412
Muscle - Skeletal	13331
Nerve - Tibial	16468
Ovary	15987
Pancreas	14857
Pituitary	17445
Prostate	17065
Skin - Not Sun Exposed (Suprapubic)	16079
Skin - Sun Exposed (Lower leg)	16097
Small Intestine - Terminal Ileum	16901
Spleen	16597
Stomach	15990
Testis	21913
Thyroid	16704
Uterus	15946
Vagina	16538
Whole Blood	13066

769 **Supplementary Table 1**: *Number of studied genes per tissue.*

770 For each tissue, we kept the genes with at least 1 CPM of expression in at least 40% of the

771 samples.

References

1. Campisi, J. Aging, Cellular Senescence, and Cancer. *Annu. Rev. Physiol.* **75**, 685–705 (2013).

2. Niccoli, T. & Partridge, L. Ageing as a Risk Factor for Disease. *Curr. Biol.* **22**, R741–R752 (2012).

3. Wyss-Coray, T. Ageing, neurodegeneration and brain rejuvenation. *Nature* **539**, 180–186 (2016).

4. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The Hallmarks of Aging. *Cell* **153**, 1194–1217 (2013).

5. Stegeman, R. & Weake, V. M. Transcriptional Signatures of Aging. *J. Mol. Biol.* **429**, 2427–2437 (2017).

6. Tower, J. Sex-Specific Gene Expression and Life Span Regulation. *Trends Endocrinol. Metab.* **28**, 735–747 (2017).

7. Austad, S. N. & Fischer, K. E. Sex Differences in Lifespan. *Cell Metab.* **23**, 1022–1033 (2016).

8. Mele, M. *et al.* The human transcriptome across tissues and individuals. *Science* **348**, 660–665 (2015).

9. Gershoni, M. & Pietrokovski, S. The landscape of sex-differential transcriptome and its consequent selection in human adults. *BMC Biol.* **15**, 7 (2017).

10. Kassam, I., Wu, Y., Yang, J., Visscher, P. M. & McRae, A. F. Tissue-specific sex differences in human gene expression. *Hum. Mol. Genet.* **28**, 2976–2986 (2019).

11. Mayne, B. T. *et al.* Large Scale Gene Expression Meta-Analysis Reveals Tissue-Specific, Sex-Biased Gene Expression in Humans. *Front. Genet.* **7**, (2016).

12. Zahn, J. M. *et al.* AGEMAP: A Gene Expression Database for Aging in Mice. *PLoS Genet.* **3**, e201 (2007).

13. Kimmel, J. C. *et al.* Murine single-cell RNA-seq reveals cell-identity- and tissue-specific trajectories of aging. *Genome Res.* **29**, 2088–2103 (2019).

14. Benayoun, B. A. *et al.* Remodeling of epigenome and transcriptome landscapes with aging in mice reveals widespread induction of inflammatory responses. *Genome Res.* **29**, 697–709 (2019).

15. Lui, J. C., Chen, W., Barnes, K. M. & Baron, J. Changes in gene expression associated with

aging commonly originate during juvenile growth. *Mech. Ageing Dev.* **131**, 641–649

(2010).

16. Shavlakadze, T. *et al.* Age-Related Gene Expression Signature in Rats Demonstrate Early,

Late, and Linear Transcriptional Changes from Multiple Tissues. *Cell Rep.* **28**, 3263-

3273.e3 (2019).

17. The Tabula Muris Consortium. A single-cell transcriptomic atlas characterizes ageing

tissues in the mouse. *Nature* **583**, 590–595 (2020).

18. Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature*

583, 596–602 (2020).

19. Zahn, J. M. *et al.* Transcriptional profiling of aging in human muscle reveals a common

aging signature. *PLoS Genet.* **preprint**, e115 (2005).

20. Welle, S., Brooks, A. I., Delehanty, J. M., Needler, N. & Thornton, C. A. Gene expression

profile of aging in human muscle. *Physiol. Genomics* **14**, 149–159 (2003).

21. Jozsi, A. C. *et al.* Aged human muscle demonstrates an altered gene expression profile

consistent with an impaired response to exercise. *Mech. Ageing Dev.* **120**, 45–56 (2000).

22. Rodwell, G. E. J. *et al.* A Transcriptional Profile of Aging in the Human Kidney. *PLoS Biol.*

2, e427 (2004).

23. Galatro, T. F. *et al.* Transcriptomic analysis of purified human cortical microglia reveals

age-associated changes. *Nat. Neurosci.* **20**, 1162–1171 (2017).

24. Olah, M. *et al.* A transcriptomic atlas of aged human microglia. *Nat. Commun.* **9**, 539

(2018).

25. Berchtold, N. C. *et al.* Gene expression changes in the course of normal brain aging are

sexually dimorphic. *Proc. Natl. Acad. Sci.* **105**, 15605–15610 (2008).

26. Lu, T. *et al.* Gene regulation and DNA damage in the ageing human brain. *Nature* **429**,

883–891 (2004).

27. Işıldak, U., Somel, M., Thornton, J. M. & Dönertaş, H. M. Temporal changes in the gene

expression heterogeneity during brain development and aging. *Sci. Rep.* **10**, 4080 (2020).

28. Haustead, D. J. *et al.* Transcriptome analysis of human ageing in male skin shows mid-life

period of variability and central role of NF-κB. *Sci. Rep.* **6**, 26846 (2016).

29. Holzscheck, N. *et al.* Multi-omics network analysis reveals distinct stages in the human

aging progression in epidermal tissue. *Aging* **12**, 12393–12409 (2020).

30. Nakamura, S. *et al.* Identification of blood biomarkers of aging by transcript profiling of

whole blood. *Biochem. Biophys. Res. Commun.* **418**, 313–318 (2012).

31. Harries, L. W. *et al.* Human aging is characterized by focused changes in gene expression

and deregulation of alternative splicing. *Aging Cell* **10**, 868–878 (2011).

32. Thomas, R. Age-Associated Changes in Gene Expression Patterns in the Liver,. *J.*

Gastrointest. Surg. **6**, 445–454 (2002).

33. Yoshida, S., Yashar, B. M., Hiriyanna, S. & Swaroop, A. Microarray analysis of gene

expression in the aging human retina. *Invest. Ophthalmol. Vis. Sci.* **43**, 2554–60 (2002).

34. Gheorghe, M. *et al.* Major aging-associated RNA expressions change at two distinct age-

positions. *BMC Genomics* **15**, 132 (2014).

35. Yang, J. *et al.* Synchronized age-related gene expression changes across multiple tissues

in human and the link to complex diseases. *Sci. Rep.* **5**, 15145 (2015).

36. Aramillo Irizar, P. *et al.* Transcriptomic alterations during ageing reflect the shift from

cancer to degenerative diseases in the elderly. *Nat. Commun.* **9**, 327 (2018).

37. Glass, D. *et al.* Gene expression changes with age in skin, adipose tissue, blood and brain.

Genome Biol. **14**, R75 (2013).

38. Yu, Y. *et al.* A rat RNA-Seq transcriptomic BodyMap across 11 organs and 4

developmental stages. *Nat. Commun.* **5**, 3230 (2014).

39. Rando, T. A. & Wyss-Coray, T. Asynchronous, contagious and digital aging. *Nat. Aging* **1**,

29–35 (2021).

40. Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–

585 (2013).

41. Van Deursen, J. M. The role of senescent cells in ageing. *Nature* **509**, 439–446 (2014).

42. Gorgoulis, V. *et al.* Cellular Senescence: Defining a Path Forward. *Cell* **179**, 813–827

(2019).

43. Gil, J. & Peters, G. Regulation of the INK4b–ARF–INK4a tumour suppressor locus: all for

one or one for all. *Nat. Rev. Mol. Cell Biol.* **7**, 667–677 (2006).

44. Erickson, S. *et al.* Involvement of the Ink4 proteins p16 and p15 in T-lymphocyte

senescence. *Oncogene* **17**, 595–602 (1998).

45. Narita, M. *et al.* Rb-Mediated Heterochromatin Formation and Silencing of E2F Target

Genes during Cellular Senescence. *Cell* **113**, 703–716 (2003).

46. Sun, X. & Kaufman, P. D. Ki-67: more than a proliferation marker. *Chromosoma* **127**,

175–186 (2018).

47. Kaczmarek, M. Variation in Age at Natural Menopause among Polish Women in Relation

to Biological and Social Factors. (2014) doi:10.13140/2.1.2610.1449.

48. Croft, D. *et al.* The Reactome pathway knowledgebase. *Nucleic Acids Res.* **42**, D472–D477

(2014).

49. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–

50 (2005).

50. Kanehisa, M. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.* **28**,

27–30 (2000).

51. Gene Ontology Consortium. The Gene Ontology (GO) database and informatics resource.

Nucleic Acids Res. **32**, 258D – 261 (2004).

52. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

53. van Dam, S., Võsa, U., van der Graaf, A., Franke, L. & de Magalhães, J. P. Gene co-

expression analysis for functional classification and gene–disease predictions. *Brief.*

Bioinform. bbw139 (2017) doi:10.1093/bib/bbw139.

54. Kelley KW, Nakao-Inoue H, Molofsky AV, O. M. Variation among intact tissue samples

reveals the core transcriptional features of human CNS cell classes. *Nat. Neurosci.* **21**,

1171–1184 (2018).

55. Piñero, J. *et al.* The DisGeNET knowledge platform for disease genomics: 2019 update.

Nucleic Acids Res. (2019) doi:10.1093/nar/gkz1021.

56. Piñero, J. *et al.* DisGeNET: a comprehensive platform integrating information on human

disease-associated genes and variants. *Nucleic Acids Res.* **45**, D833–D839 (2017).

57. Palmer, A. L. & Ousman, S. S. Astrocytes and Aging. *Front. Aging Neurosci.* **10**, 337

(2018).

58. Lehallier, B. *et al.* Undulating changes in human plasma proteome profiles across the

lifespan. *Nat. Med.* **25**, 1843–1850 (2019).

59. Khramtsova, E. A., Davis, L. K. & Stranger, B. E. The role of sex in the genomics of human

complex traits. *Nat. Rev. Genet.* **20**, 173–190 (2019).

60. Subramanian, A. *et al.* A Next Generation Connectivity Map: L1000 Platform and the First

1,000,000 Profiles. *Cell* **171**, 1437-1452.e17 (2017).

61. de Almeida BP, Saraiva-Agostinho N, B.-M. N. cTRAP: Identification of candidate causal

perturbations from differential gene expression data. Preprint at (2020).

62. de Magalhães, J. P., Curado, J. & Church, G. M. Meta-analysis of age-related gene

expression profiles identifies common signatures of aging. *Bioinformatics* **25**, 875–881

(2009).

63. Tacutu, R. *et al.* Human Ageing Genomic Resources: new and updated databases. *Nucleic*

Acids Res. **46**, D1083–D1090 (2018).

64. Craig, T. *et al.* The Digital Ageing Atlas: integrating the diversity of age-related changes

into a unified resource. *Nucleic Acids Res.* **43**, D873–D878 (2015).

65. Dönertaş, H. M., Fuentealba Valenzuela, M., Partridge, L. & Thornton, J. M. Gene

expression-based drug repurposing to target aging. *Aging Cell* **17**, e12819 (2018).

66. Janssens, G. E. *et al.* Transcriptomics-Based Screening Identifies Pharmacological

Inhibition of Hsp90 as a Means to Defer Aging. *Cell Rep.* **27**, 467-480.e6 (2019).

67. https://shiny.rstudio.com/gallery/.

68. Kunst, J. highcharter: A Wrapper for the 'Highcharts' Library.

69. Yihui, X., Joe, C. & Xianying, T. DT: A Wrapper of the JavaScript Library 'DataTables'.

70. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–

140 (2010).

71. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

72. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock linear model

analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).

73. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing

and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

74. García-Pérez, R. *et al.* The landscape of expression and alternative splicing variation

across human traits. *Cell Genomics* **3**, 100244 (2023).

75. Buuren, S. V. & Groothuis-Oudshoorn, K. **mice** : Multivariate Imputation by Chained

Equations in *R*. *J. Stat. Softw.* **45**, (2011).

76. McCall, M. N., Illei, P. B. & Halushka, M. K. Complex Sources of Variation in Tissue

Expression Data: Analysis of the GTEx Lung Transcriptome. *Am. J. Hum. Genet.* **99**, 624–

635 (2016).

77. Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2021)

doi:10.1101/060012.

78. GSEA-MSIGDB. Gene Set Enrichment Analysis (GSEA) User Guide.

79. Chen, Y.-A. *et al.* Integrated Pathway Clusters with Coherent Biological Themes for

Target Prioritisation. *PLoS ONE* **9**, e99030 (2014).

80. Dowle, M. & Srinivasan, A. data.table: Extension of 'data.frame'.

81. Langfelder, P. & Horvath, S. WGCNA Frequently Asked Questions.

82. Chen, B., Khodadoust, M. S., Liu, C. L., Newman, A. M. & Alizadeh, A. A. Profiling Tumor

Infiltrating Immune Cells with CIBERSORT. in *Cancer Systems Biology* (ed. Von Stechow,

L.) vol. 1711 243–259 (Springer New York, 2018).

83. Regev, A. *et al.* The Human Cell Atlas. *eLife* **6**, e27041 (2017).

84. Fan, X. *et al.* Spatial transcriptomic survey of human embryonic cerebral cortex by single-

cell RNA-seq analysis. *Cell Res.* **28**, 730–745 (2018).

85. Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**, eaba7721 (2020).