A large-scale serological survey in pets from October 2020 through June 2021 in France

shows significantly higher exposure to SARS-CoV-2 in cats

Matthieu Fritz[a], Eric Elguero[a,1], Pierre Becquart[a,1], Daphné de Riols de Fonclare[a], Déborah Garcia[a],

Stephanie Beurlet [b], Solène Denolly [c], Bertrand Boson [c], Serge G. Rosolen [e, f], François-Loïc Cosset[c],

Alexandra Briend-Marchal [b], Vincent Legros [c, d], Eric M. Leroy[a*]

[a] Maladies Infectieuses et Vecteurs, Ecologie, Génétique, Evolution et Contrôle (MIVEGEC), Univ.

Montpellier, IRD, CNRS, 34394 Montpellier, France

[b] Laboratoire de Biologie vétérinaire VEBIO, Arcueil, France

[c] CIRI – Centre International de Recherche en Infectiologie, Team EVIR, Univ Lyon, Université Claude

Bernard Lyon 1, Inserm, U1111, CNRS, UMR5308, ENS Lyon, Lyon, France.

[d] Université de Lyon, VetAgro Sup, Campus vétérinaire de Lyon, Marcy-l'Etoile, France

[e] Sorbonne Université, INSERM, CNRS, Institut de la Vision, Paris, France

[f] Clinique vétérinaire voltaire, Asnières, France

[1] Authors contributed equally

*Corresponding author:

Eric M.Leroy : UMR IRD-CNRS-UM, 911 avenue Agropolis, 34394 Montpellier, France

Tel : +33 467416107 ; E-mail address : eric.leroy@ird.fr

Keywords: SARS-CoV-2, Pets, Serology, Neutralization assay, Microsphere Immunoassay, One Health

Abstract

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can infect many animals, including

pets such as dogs and cats. Many studies have documented infection in companion animals by bio-

molecular and serological methods. However, only a few have compared seroprevalence in cats and

dogs from the general population, and these studies were limited by small sample sizes and

collections over short periods. Our goal was to obtain a more accurate evaluation of seroprevalence

in companion animals in France and to determine whether cats and dogs differ in their exposure to

SARS-CoV-2. For this purpose, we conducted an extensive SARS-CoV-2 cross-sectional serological

survey of 2036 cats and 3577 dogs sampled by veterinarians during medical examinations in clinics

throughout France. Sampling was carried out from October 2020 through June 2021, a period

encompassing the second and third waves of SARS-CoV-2 infections in humans in the country. Using

a microsphere immunoassay targeting the receptor binding domain and trimeric spike protein, we

found 7.1% seroprevalence in pets. In a subset of 308 seropositive samples, 26.3% had neutralizing

antibodies. We found that cats were significantly more likely to test positive than dogs, with

seropositivity rates of 9.3% and 5.9% in cats and dogs, respectively. Finally, data for both species

showed that seroprevalence was lower in older animals and was not associated with the date of

sampling or the sex of the animal. Our results show that cats are significantly more sensitive to SARS-

CoV-2 than dogs, in line with experimental studies. Our large sample size provides for a reliable,

statistically robust estimate of the frequency of infection of pets from their owners and offers strong

support for the notion that cats are more sensitive to SARS-CoV-2 than dogs. Our findings emphasise

the importance of a One-Health approach to the SARS-CoV-2 pandemic and raise the question of

whether companion animals in close contact with humans should be vaccinated.

Introduction

Two months after the onset of SARS-CoV-2 circulation in humans, two dogs in Hong Kong were

reported to have naturally acquired the virus (1). Since then, many studies have reported viral RNA

and SARS-CoV-2 antibodies in dogs and cats—mostly belonging to COVID-19-infected owners (2-5).

Furthermore, several studies demonstrated that the risk of pets testing seropositive was higher in

COVID-19+ households than for pets from households of unknown status (6-10).

Definitive examples of pet-to-human transmission are scarce. A recent study from Thailand reported

a suspected case of SARS-CoV-2 transmission from a cat to a human (11), and dog-to-human

transmission has yet to be described. However, given that 200 million cats and dogs live in close

proximity to humans in Europe (12), there is ample opportunity for such transmission, and the

potential risks need to be carefully considered.

Several population-based serological studies have reported SARS-CoV-2 antibodies in dogs and cats.

In dogs, estimates of seroprevalence have ranged from 0% to 14.5% (6, 10, 13-21). While in cats,

estimates have ranged from 0% to 21.7% (6, 14, 15, 20, 22-25). For both species, seroprevalence was

highly dependent on the period of sampling (first, second wave *etc.*), the assay used (ELISA,

seroneutralization, etc.), and the country of sampling (China, Croatia, Germany, Italy, the

Netherlands, Poland, Portugal, Spain, Thailand, United-Kingdom, USA). Among these studies, five

directly compared cats and dogs. There is some experimental and epidemiological evidence

suggesting that cats are more susceptible to infection than dogs (6, 7, 15, 26, 27). However,

significant species differences have not always been observed in population-based studies (13, 17,

19). This is perhaps because of significant study limitations—a low number of enrolled animals, a

short sampling period, etc.—that have curtailed robust estimates of infection rate in pets with

enough statistical power to recognize differences in COVID-19 epidemiology.

Here we report estimates of the frequency of SARS-CoV-2 infection from 2036 cats and 3577 dogs

sampled at veterinary clinics from October 2020 through June 2021 throughout France—the largest

serological survey of SARS-CoV-2 infections in companion animals to date. The study allows for

robust estimates of pet infection rate and provides strong support for the hypothesis that species

differences in susceptibility observed in experimental studies translate into a significant increase in

infection rate in cats.

Materials and Methods

Sampling

The cross-sectional, nationwide sampling was possible thanks to a network of veterinary clinics

across France working with VEBIO. VEBIO is a veterinary diagnostic laboratory which performed all

categories of medical analyses, including infectious diseases, hematology, endocrinology, oncology …

(see more details in https://www.vebio.fr/). No inclusion and exclusion criteria were applied for the

selection of blood samples, except that they came only from veterinary clinics working with VEBIO.

VEBIO notified the veterinary clinics that following requested biomedical analyses, the remaining

serum could be used in a SARS-CoV-2 research project. No specific request for samples was

addressed to the vets. Thus, the SARS-CoV-2 analysis is based on samples collected during the regular

activities of the vets.

Blood samples were collected in dry/EDTA tubes from dogs and cats during routine healthcare visits

or for diagnostic purposes at veterinary clinics. After centrifugation, the serum/plasma was kept at

+4 °C until sent to VEBIO. Rapid and safe shipping practices were used to avoid contamination and

ensure samples reached VEBIO within 48 h. At the VEBIO facility, an aliquot was taken from the

sample to perform the requested biomedical analyses. Another aliquot was then stored at +4 °C until

sent to the MIVEGEC lab, Montpellier, where serological analyses were performed. Safe shipping

practices with an approved professional carrier were also used for shipment to the MIVEGEC lab.

Finally, the samples were stored at the MIVEGEC lab at −20 °C until testing (Figure 1). For shipping to

the CIRI lab, SARS-CoV-2-positive samples detected by MIA were transported by an approved

professional carrier at -20°C to ensure optimal safety conditions. Data (age, sex, clinical history, and

region localization, when available) from dogs and cats were provided anonymized by VEBIO to the

MIVEGEC lab.

Figure 1. Logistics of sample collection and distribution. Sera collected during routine healthcare

visits by veterinarians throughout France were first sent to VEBIO in Ile-de-France. Aliquots of the

samples were then made and sent to the IRD in Montpellier (Hérault) via an approved carrier.

Ethics

According to the act governing the "use of live animals for scientific purposes" effective in France on

14 January 2022, ethical approval was not sought or required since all pets were sampled by a

veterinarian during a health care visit. All applicable international and national guidelines for the care

of pets were followed.

Microsphere Immunoassay (MIA)

Dog and cat serum samples were tested using a multiplex microsphere immunoassay (MIA). Ten µg

of two recombinant SARS-CoV-2 antigens, receptor-binding domain (RBD) and trimeric spike (tri-S),

both derived from the Whuhan-Hu-1 strain (The Native Antigen Company, Kidlington United-

Kingdom), were used to capture specific serum antibodies. Distinct MagPlex microsphere sets

(Luminex Corp, Austin, TX, USA) were respectively coupled to viral antigens using the amine coupling

kit (Bio-Rad Laboratories, Marnes-la-Coquette, France) according to the manufacturer's instructions.

Microsphere mixtures were successively incubated with serum samples (1:400), biotinylated protein

A and biotinylated protein G (4 µg/mL each) (Thermo Fisher Scientific, Illkirch, France), and

streptavidin-R-phycoerythrin (4 µg/mL) (Life technologies, Illkirch, France) on an orbital shaker and

protected from light. Measurements were performed using a Luminex 200 instrument (Luminex

Corp, Austin, TX, USA), and at least 100 events were read for each bead set. Binding events were

displayed as median fluorescence intensities (MFI). Specific seropositivity cut-off values for each

antigen were set at three standard deviations above the mean MFI of pre-pandemic serum from 53

dogs and 30 cats sampled before 2019. These samples were stored in biobanks at the IRD and

VetAgro Sup. MIA specificity was set for each antigen at 96.2% for dogs and 100% for cats based on

the pre-pandemic populations. MIA was first validated using sera from two COVID-19 PCR+ humans,

kindly provided by Meriadeg Ar Gouilh, and then with sera from SARS-CoV-2 PCR+ cats and dogs,

provided by several veterinarians.

Because of the excellent specificity observed for both antigens and to account for any isotypic

variability, an animal was deemed positive for SARS-CoV-2 antibodies following a positive result in at

least one of the two tests.

Neutralization activity measurement

An MLV-based pseudoparticle carrying a GFP reporter pseudotyped with SARS-CoV-2 spike protein

(Wuhan-Hu-1 strain) (SARS-CoV-2pp) was used to measure neutralizing antibody activity in cat and

dog sera. Each SARS-CoV-2-positive sample detected by MIA was processed according to a

neutralization procedure as previously described (28) . Briefly, for neutralization assays, a sample of

~1×10^3 pseudoparticles was incubated with a 100-fold dilution of sera or control antibodies for

1 h at 37 °C before infection of Vero-E6R cells. At 72 h post-transduction, the percentage of GFP-

positive cells was determined by flow cytometry (at least 10 000 events recorded). The level of

infectivity is expressed as the percentage of GFP-positive cells and compared to cells infected with

SARS-CoV-2pp incubated without serum. As a control, the same procedure was done using RD114

pseudoparticles to identify sera with aspecific neutralization. Sera exhibiting more than 30% SARS-

CoV-2pp neutralization were considered positive. Pre-pandemic serum from France was used as a

negative control, and an anti-SARS-CoV-2 RBD antibody was used as a positive control.

Statistical analyses

Associations between SARS-CoV2 infection status (positive or negative) and the covariates region,

age, and sex were assessed using binomial (logistic) generalized linear models. The region was

defined by where the animal lived at the time of sampling. Age was that recorded by the

veterinarian, with variable precision, generally in months for young animals and whole years for

older animals. Its accuracy is unknown. The associated statistical tests were likelihood ratio tests. All

analyses were performed using R software (29).

Results

Blood collection

Blood samples from 2036 cats and 3577 dogs were collected during routine healthcare visits by

veterinarians from October 2020 through June 2021 (Table 1). Samples came from all 13 regions of

metropolitan France. Corsica was excluded due to too few samples. Almost half of the samples came

from Ile-de-France, the region including Paris, reflecting population density and proximity to the

Veterinary diagnostic laboratory (VEBIO), where all samples for biomedical analyses requested by the

veterinarians were handled (Materials and Methods). The number of samples received from other

regions largely depended on the number of veterinarians working with VEBIO in those regions (Figure

2). Unfortunately, we could not study the clinical history of the animals due to variability in how each

veterinarian reported this information.

	October 2020	November 2020	December 2023	January 2021	February 2021	March 2021	April 2021	May 2021	June 2021
Cats	49	256	275	291	225	296	305	166	173
Dogs	84	428	543	475	403	474	597	282	291
Total	133	684	818	766	628	770	902	448	464

160 **Table 1** : numbers of samples collected by month and by species.

Figure 2. (a). Map of France showing the number of SARS-CoV-2-positive cat sera per region. The total number of sera samples collected per region is indicated. Seroprevalence in each region is indicated as a percentage. Regions are shaded in green according to seroprevalence. The total number of sera samples and global seroprevalence for France is in the top left corner. **(b). Map of France showing the number of SARS-CoV-2-positive dog sera per region.** The total number of sera samples collected per region is indicated. Seroprevalence in each region is indicated as a percentage. Regions are shaded in orange according to seroprevalence. The total number of sera samples and global seroprevalence for France is in the top left corner.

Global seroprevalence

For the sera samples, 401 (7.1%) showed a positive result either against RBD, tri-S, or both (Supplementary Table 1). We next determined the presence of antibodies with neutralizing activity among the positive sera. To save time, we randomly tested approximately 75% (308) of positive sera samples. Seroneutralizing activity was detected in 81 (26.3%) of the 308 pet sera samples. Among these positive samples, 39 (48%) were positive for both RBD and tri-S, 39 (48%) were positive only for tri-s and 3 (4%) were only positive for RBD. Only the seroprevalence from MIA assays was analyzed in the remainder of the study.

Seroprevalence in cats and dogs

We observed that a significantly greater proportion of cats were positive (189/2036, 9.3%) than dogs (212/3577, 5.9%); OR = 1.62, 95% c.i. [1.32 - 1.99], P-value = 3.8e-06, Table 2. In addition, sera from MIA-positive cats were more likely to show neutralizing activity (49/144, 34%) than dogs (32/164, 19.5%); OR = 2.12, 95% c.i. [1.27 - 3.57], P-value = 0.0039). Species differences were not always significant within each region, likely due to reduced statistical power. However, when differences were significant, it was always the case that cats were more likely to be positive than dogs. (Table 2).

Table 2. Seroprevalence of IgG SARS-CoV-2 antibodies detected in blood samples from cats and dogs

collected in different French regions from October 2020 through June 2021. Data are presented as

No. positive, percentage (95 % exact binomial confidence intervals). Odds ratios were computed by

fitting binomial models region by region; an OR > 1 indicates cats were more likely to be positive than

dogs. In this analysis, individual data on pet age and sex were not considered, as age was not

available for all animals. P-values were computed by the likelihood ratio test.

Region	Cats	Cats seroprevalence		Dogs	Dogs seroprevalence		OR (95% c.i.)	P-value
Auvergne-Rhône-Alpes	4/92	4.3%	(1.2-10.8)	25/298	8.4%	(5.5-12.1)	0.50 (0.17 - 1.47)	0.17
Bourgogne-Franche-Comté	3/49	6.1%	(1.3-16.9)	4/91	4.4%	(1.2-10.9)	1.42 (0.30 - 6.61)	0.66
Bretagne	3/38	7.9%	(1.7-21.4)	5/113	4.4%	(1.5-10.0)	1.85 (0.42 - 8.14)	0.43
Centre-Val de Loire	7/26	26.9%	(11.6-47.8)	3/53	5.7%	(1.2-15.7)	6.14 (1.44 - 26.2)	0.0098
Grand Est	8/42	19.0%	(8.6-34.1)	5/172	2.9%	(1.0-6.7)	7.86 (2.42 - 25.5)	0.00057
Hauts-de-France	6/91	6.6%	(2.5-13.8)	16/211	7.6%	(4.4-12.0)	0.86 (0.33 - 2.27)	0.76
Île-de-France	104/1193	8.7%	(7.2-10.5)	98/1248	7.9%	(6.4-9.5)	1.12 (0.84 - 1.49)	0.44
Normandie	3/62	4.8%	(1.0-13.5)	8/144	5.6%	(2.4-10.7)	0.86 (0.22 - 3.37)	0.83
Nouvelle-Aquitaine	21/213	9.9%	(6.2-14.7)	20/611	3.3%	(2.0-5.0)	3.23 (1.72 - 6.09)	0.00037
Occitanie	8/92	8.7%	(3.8-16.4)	12/386	3.1%	(1.6-5.4)	2.97 (1.18 - 7.49)	0.028
Provence-Alpes-Côte d'Azur	19/86	22.1%	(13.9-32.3)	12/182	6.6%	(3.5-11.2)	4.02 (1.85 - 8.73)	0.00036
Pays de la Loire	3/52	5.8%	(1.2-15.9)	4/68	5.9%	(1.6-14.4)	0.98 (0.21 - 4.58)	0.98
Total	**189/2036**	**9.3%**	**(8.0 - 10.6)**	**212/3577**	**5.9%**	**(5.2 - 6.8)**	**1.62 (1.32 - 1.99)**	**3.8×10^{-6}**

Seroprevalence by sex

We found no significant sex differences in seropositivity rates, either for all animals (females: 6.9%;

163/2361; males 7.5%; 212/2842; p = 0.24) or among cats (females 9.4%; 78/827, males 9.8%;

99/1009, p = 0.68) and dogs (females: 5.5%; 85/1534, males: 6.2%; 113/1833; p = 0.27) tested

separately (Table 3).

Table 3. Seroprevalence of IgG SARS-CoV-2 antibodies in blood samples from cats and dogs by sex

from October 2020 through June 2021 Data are presented as No. positive, percentage (95 % exact

binomial confidence intervals). Odds ratios > 1 indicate males are more likely to be positive than

females and were computed by fitting binomial generalized linear models, with age as a controlling

factor. P-values correspond to likelihood ratio tests.

Sex	Cats	Cat seroprevalence	Dogs	Dog seroprevalence	Cats + Dogs	Cats + Dogs seroprevalence
Female	78/827	9.4% (7.5-11.5)	85/1534	5.5% (4.4-6.8)	163/2361	6.9% (5.9-8.0)
Male	99/1009	9.8% (8.0-11.8)	113/1833	6.2% (5.1-7.4)	212/2842	7.5% (6.5-8.5)
Total	**177/1836**	**9.6% (8.3-11.1)**	**198/3367**	**5.9% (5.1-6.7)**	**375/5203**	**7.2% (6.5-7.9)**
OR	1.08 (0.75 - 1.54)		1.20 (0.87 - 1.67)		1.15 (0.91 - 1.47)	
P-value	0.68		0.27		0.24	

Seroprevalence by age

Age was reported for 1657 cats (range: 0.2 – 22yr) and 2781 dogs (range: 0.1 – 18.5yr). Among cats,

18.4% aged [0-3] years, 10.4% aged]3-9] years, and 6.4% aged over 9 years tested positive.

Among dogs, 8.8% aged [0-3] years, 5.8% aged]3-9] years, and 5.1% aged over 9 years tested

positive. Using a binomial model with age entered as a continuous variable, we observed a significant

decrease in seroprevalence with age in cats (OR for a one-year increase in age = 0.91, 95% c.i. [0.88 -

0.94], p-value = 3.7e-08) and dogs (OR = 0.95, 95% c.i. [0.92 - 0.99], p-value = 0.016) (Figure 3)

(Supplementary Table 2).

Figure 3. (a). The number of cat blood samples tested by age group for anti-SARS-CoV-2 antibodies by MIA from October 2020 through June 2021. Samples testing negative are shaded grey, and seropositive samples are in orange. Seroprevalence is represented by black dots, with 95 % binomial confidence interval. The red line represents the linear regression. **(b). The number of dog blood**

samples tested by age group for anti-SARS-CoV-2 antibodies by MIA from October 2020 through

June 2021. Samples testing negative are shaded grey, and seropositive samples are in orange.

Seroprevalence is represented by black dots, with 95 % binomial confidence intervals. The red line

represents the linear regression.

Seroprevalence over time

We next examined whether seroprevalence was associated with the time of sampling. For this

analysis, we selected animals at least one year old at the date of sampling (Figure 4). Seroprevalence

was not associated with the time of sampling for cats: p-value = 0.41. However, seroprevalence

among dogs increased over the 9 months of the study (OR = 3.47, 95% c.i. [1.47 - 8.23], p-value =

0.0045).

(a)

(b)

234   **Figure**

4. (a). The number of cat blood samples tested each month for anti-SARS-CoV-2 antibodies by MIA

from October 2020 through June 2021. Samples testing negative are shaded grey, and seropositive

samples are in orange. Seroprevalence is represented by black dots, with 95 % binomial confidence

interval. The red line represents the linear regression. (b). The number of dog blood samples tested

each month for anti-SARS-CoV-2 antibodies by MIA from October 2020 through June 2021. Samples

testing negative are shaded grey, and seropositive samples are in orange. Seroprevalence is

represented by black dots, with 95 % binomial confidence interval. The red line represents the linear

regression. Notice that in this figure, dates have been pooled by calendar month for illustrative

purposes but that in the statistical analysis exact dates were used. Likewise, the regression lines are

similarly illustrative as the statistical tests were based on logistic (binomial) regression.

Discussion

This study reports a large-scale serological survey of pet (cats and dogs) to detect anti-SARS-CoV-2

IgG antibodies. The samples were collected in metropolitan France from October 2020 through June

2021, a period including peaks of the second and third waves of SARS-CoV-2 infections in France.

From a sample of 5613 pets, we reported a seroprevalence of anti-SARS-CoV-2 antibodies of 7.1%.

We observed that only a small percentage of samples (48%) were positive for both tri-s and RBD,

indicating that the RBD assay may be less sensitive than the tri-s assay. This may be explained by the

fact that the full trimeric spike antigen may bind a broader range of antibodies than the receptor-

binding domain, which includes only a small part of the spike protein. We found neutralizing

antibody activity in the sera of only 26% of seropositive pets. Previous studies have shown that some

pets do not develop neutralizing antibodies (5, 30). Cats were more likely to produce neutralizing

antibodies than dogs, which is likely associated with a more prolonged and intense immune

stimulation in cats. In humans, disease severity is positively correlated with neutralizing antibody

levels(31).

In cats, we found a higher seroprevalence (9.3%) than previously observed in other European

countries, which ranged from 0% to 6.4% (13, 17-19, 21-24). However, most of these studies were

done before the second wave, during a period of relatively lower viral circulation than our sampling

period. In addition, most of these studies used a seroneutralisation assay.

In dogs, the observed seroprevalence (5.9%) is in accord with a previous study in France showing a

prevalence of 4.8% in companion and military working dogs sampled between February 2020 and

February 2021 (16). Other studies looking for SARS-CoV-2 antibodies in dogs have reported

seroprevalences ranging from 0% to 14.5% (10, 13, 17-19, 21).

Importantly, we observed a significantly higher seroprevalence of anti-SARS-CoV-2 antibodies in cats

than in dogs (p = 4.2e-08). The statistical significance of this difference varied among regions, likely

due to the reduced power and perhaps some unintended sampling bias by veterinarians. For

example, the smallest sample size was in the Bourgogne-Franche-Comté region, where we observed

no significant difference between dogs and cats. Furthermore, for a region like Ile-de-France, where

people live mostly in apartments, we can also hypothesize that dogs live in closer contact with

owners than in the rest of France. Previous studies with fewer samples have either found no

significant difference between species (8, 9, 13, 17, 19, 32) or that cats have significantly higher

seroprevalence than dogs (3, 6, 7).

Our study of a very large population of dogs and cats in natural conditions provides some evidence

that cats are more susceptible to SARS-CoV-2 infection than dogs, at least during the time frame of

our sampling period. Potential causes of species differences in susceptibility between cats and dogs

are numerous, but likely include a variety of biological and behavioural factors, as well as differences

in exposure. Intererestingly, ACE-2 shows greater sequence similarity between cat and human

orthologs than observed between dogs and humans (33). The absence of data such as the pet

lifestyle (Indoor/Outdoor), or the frequency and nature of contacts with humans and other animals

restricts our ability to identify a potential cause of the observed difference. In previous studies, most

infected pets were epidemiologically linked to humans who had tested positive for COVID-19 (34).

We did not observe significant sex differences in seroprevalence in either species (p = 0.45). Our

findings are consistent with most previous studies also reporting an absence of sex differences in

dogs and cats (6, 17, 32, 35). A smaller study of 188 dogs and 61 cats found higher seropositivity in

male dogs and an absence of a sex difference in cats (9). Another study found that male dogs

sampled from the general population were more likely to test positive than females, but this

difference was not observed in dogs from COVID-19+ households (10). There is little evidence of a

significant sex difference in susceptibility in humans. However, men are more likely to be affected by

severe forms of COVID than women for a variety of reasons (36).

In terms of age, we observed a higher seroprevalence among younger animals (between 0-3 years)

for both species that then decreased with age. A study of dogs sampled from the general population

found seroprevalence was highest in animals aged 5-6 years and that in COVID-19+ households,

seroprevalence peaked in slightly younger dogs, aged between one and five years (10). Other studies

have reported no significant associations with age in cats and dogs (6, 17). An experimental study in

cats found that juveniles appear more vulnerable than subadults (27). The decreasing seroprevalence

we observed with age could also arise from age-dependent behavioural changes. For example, young

animals (< 3 years old) are more active and curious and may be in greater contact with their owners

than older animals that prefer to remain quieter. The decrease could also reflect immunosenescence

in older animals, as observed in humans.

Interestingly, we observed a slight increase in seroprevalence in dogs during the study's nine months

of sampling, a trend not observed among cats. We expected an increase because antibodies have a

longer persistence in the organism than viral RNA; thus, animals sampled at later dates would

represent an accumulation of cases. The absence of a positive association between seroprevalence

and the time of sampling in cats has been reported in two other studies in Europe, but conclusions

were limited by the small number of samples collected over just a few months (24, 37). The absence

of an association in cats suggests a limited persistence of antibodies in cats than dogs. Few studies

have investigated variation in the persistence of antibodies in animals. For example, a study carried

out on seven dogs and two cats infected in natural conditions showed persistence of neutralizing

antibodies up to 10 months after infection in four of the dogs and the two cats, but also that

persistence was markedly reduced in two of the dogs after three months (38). Moreover, a study of

two cats found that neutralizing antibodies had disappeared by 110 days (25). Based on these data,

one possible reason for the lack of increase in seroprevalence during our study period could be a

progressive seroreversion of infected cats that is equally compensated by the number of new

infections, i.e. seroconversion. If so, this would mean that the observed seroprevalence is not an

accurate reflection of the total number of infections, at least in cats, during the whole epidemic.

Instead, seroprevalence provides a snapshot if infections acquired during a time period that remains

to be defined by longitudinal serological studies of cats and dogs. This also suggests that the

seroprevalence observed in our study may underestimate the actual proportion of cats infected

during the entirety of the epidemic.

Human-to-pet transmission may promote viral adaptation facilitating re-infection with novel viral

strains in humans (39). While one case of infection from cat to human has recently been reported,

the large number of pet cats and their frequent close interaction with humans provides ample

opportunity. This possibility raises the question of a vaccination strategy for animals susceptible to

SARS-CoV-2 infection. While pets do not currently seem to play a role in the ongoing pandemic, our

results emphasize the magnitude of SARS-CoV-2 infection in pets is not trivial. Combined with the

size of domestic cat and dog populations and the close contact with their human companions, our

results highlight the importance of collecting more data on SARS-CoV-2 transmissibility and

pathogenicity in companion animals, especially with the emergence of new variants. Also, when a

SARS-CoV-2 infection is suspected in a pet, we suggest collecting a sample for RT-qPCR confirmation

of infection, followed by whole-genome sequencing to identify new mutations, particularly in

antigenic sites targeted by the immune system. Finally, similar public health recommendations

applied to humans should also be implemented for animals to prevent human-to-animal

transmission, such as not having contact with animals when a household member is COVID-19

positive.

Acknowledgements: We are grateful to the pet owners for giving us permission to sample their pets.

We thank the veterinarians that helped us with sampling. We thank Simon Thierry and Trung Thanh

Nguyen for technical support and assistance. We also thank Kurt McKean

(https://octopusediting.com/) for the English editing of the manuscript.

Funding: The study was funded by the French National Agency for Research (ANR-RA-COVID-19;

Geographical and temporal serological investigation of companion animal infection with SARS-CoV-2

during the second wave of COVID-19 in France, CoVet). This study received financial support through

a research grant from IDEXLYON project of Université de Lyon as part of the "Programme

Investissements d'Avenir" (ANR-16-IDEX-0005). The study was funded by OIE through the European

Union "EBO-SURSY" project.

Conflicts of Interest: The authors declare no conflicts of interest.

Institutional Review Board Statement: According to the act of "use of live animals for scientific

purposes" effective in France on 14 January 2022, ethical approval was not sought or required since

all pets were sampled by a veterinarian during a health care visit. All applicable international and

national guidelines for the care of pets were followed.

Informed Consent Statement: Not applicable.

Data Availability Statement: Raw data and scripts are available on zenodo:

https://doi.org/10.5281/zenodo.7646559

Author Contributions:

M.F., P.B., S.G.R., A.B.-M., V.L., and E.M.L. conceived and designed the study.

M.F., E.E., D.d.R.d.F., D.G., S.D., B.B., and V.L designed and performed the experiments.

All authors analyzed the data and interpreted and discussed the results.

M.F. and E.L. wrote the manuscript with input from all authors.

All authors have read and agreed to the published version of the manuscript.

References

1.   Sit THC, Brackman CJ, Ip SM, Tam KWS, Law PYT, To EMW, et al. Infection of dogs with SARS-
CoV-2. Nature. 2020;586(7831):776-8.
2.   Barrs V, Peiris M, Tam KWS, Law PYT, Brackman C, To EMW, et al. SARS-CoV-2 in Quarantined
Domestic Cats from COVID-19 Households or Close Contacts, Hong Kong, China. Emerging Infectious
Disease journal. 2020;26(12):3071.
3.   Hamer SA, Pauvolid-Corrêa A, Zecca IB, Davila E, Auckland LD, Roundy CM, et al. SARS-CoV-2
Infections and Viral Isolations among Serially Tested Cats and Dogs in Households with Infected
Owners in Texas, USA. Viruses. 2021;13(5):938.
4.   Jairak W, Charoenkul K, Chamsai E, Udom K, Chaiyawong S, Bunpapong N, et al. First cases of
SARS-CoV-2 infection in dogs and cats in Thailand. Transboundary and emerging diseases. 2021.
5.   Krafft E, Denolly S, Boson B, Angelloz-Pessey S, Levaltier S, Nesi N, et al. Report of One-Year
Prospective Surveillance of SARS-CoV-2 in Dogs and Cats in France with Various Exposure Risks:
Confirmation of a Low Prevalence of Shedding, Detection and Complete Sequencing of an Alpha
Variant in a Cat. Viruses. 2021;13(9):1759.
6.   Barroso R, Vieira-Pires A, Antunes A, Fidalgo-Carvalho I. Susceptibility of Pets to SARS-CoV-2
Infection: Lessons from a Seroepidemiologic Survey of Cats and Dogs in Portugal. Microorganisms.
2022;10(2):345.
7.   Colitti B, Bertolotti L, Mannelli A, Ferrara G, Vercelli A, Grassi A, et al. Cross-Sectional
Serosurvey of Companion Animals Housed with SARS-CoV-2–Infected Owners, Italy. Emerging
Infectious Disease journal. 2021;27(7).
8.   Fritz M, Rosolen B, Krafft E, Becquart P, Elguero E, Vratskikh O, et al. High prevalence of
SARS-CoV-2 antibodies in pets from COVID-19+ households. One Health. 2021;11:100192.
9.   Patterson EI, Elia G, Grassi A, Giordano A, Desario C, Medardo M, et al. Evidence of exposure
to SARS-CoV-2 in cats and dogs from households in Italy. Nature Communications. 2020;11(1):6231.
10.   Stevanovic V, Tabain I, Vilibic-Cavlek T, Mauric Maljkovic M, Benvin I, Hruskar Z, et al. The
Emergence of SARS-CoV-2 within the Dog Population in Croatia: Host Factors and Clinical Outcome.
Viruses. 2021;13(8):1430.
11.   Sila T, Sunghan J, Laochareonsuk W, Surasombatpattana S, Kongkamol C, Ingviya T, et al.
Suspected Cat-to-Human Transmission of SARS-CoV-2, Thailand, July-September 2021. Emerg Infect
Dis. 2022;28(7):1485-8.
12.   The European Pet Food Industry. FACTS & FIGURES 2020 European Overview. 2020.
13.   Barroso-Arévalo S, Barneto A, Ramos ÁM, Rivera B, Sánchez R, Sánchez-Morales L, et al.
Large-scale study on virological and serological prevalence of SARS-CoV-2 in cats and dogs in Spain.
Transboundary and emerging diseases. 2021.
14.   Barua S, Hoque M, Adekanmbi F, Kelly P, Jenkins-Moore M, Torchetti MK, et al. Antibodies to
SARS-CoV-2 in dogs and cats, USA. Emerging Microbes & Infections. 2021:1-20.
15.   Dileepan M, Di D, Huang Q, Ahmed S, Heinrich D, Ly H, et al. Seroprevalence of SARS-CoV-2
(COVID-19) exposure in pet cats and dogs in Minnesota, USA. Virulence. 2021;12(1):1597-609.

16. Laidoudi Y, Sereme Y, Medkour H, Watier-Grillot S, Scandola P, Ginesta J, et al. SARS-CoV-2
antibodies seroprevalence in dogs from France using ELISA and an automated western blotting assay.
One Health. 2021;13:100293.
17. Pomorska-Mól M, Turlewicz-Podbielska H, Gogulski M, Ruszkowski JJ, Kubiak M, Kuriga A, et
al. A cross-sectional retrospective study of SARS-CoV-2 seroprevalence in domestic cats, dogs and
rabbits in Poland. BMC Veterinary Research. 2021;17(1):322.
18. Smith SL, Anderson ER, Cansado-Utrilla C, Prince T, Farrell S, Brant B, et al. SARS-CoV-2
neutralising antibodies in dogs and cats in the United Kingdom. Current Research in Virological
Science. 2021;2:100011.
19. Stevanovic V, Vilibic-Cavlek T, Tabain I, Benvin I, Kovac S, Hruskar Z, et al. Seroprevalence of
SARS-CoV-2 infection among pet animals in Croatia and potential public health impact.
Transboundary and emerging diseases. 2020;n/a(n/a).
20. Udom K, Jairak W, Chamsai E, Charoenkul K, Boonyapisitsopa S, Bunpapong N, et al.
Serological survey of antibodies against SARS-CoV-2 in dogs and cats, Thailand. Transboundary and
emerging diseases. 2021.
21. Zhao S, Schuurman N, Li W, Wang C, Smit LAM, Broens E, et al. Serologic Screening of Severe
Acute Respiratory Syndrome Coronavirus 2 Infection in Cats and Dogs during First Coronavirus
Disease Wave, the Netherlands. Emerging Infectious Disease journal. 2021;27(5):1362.
22. Michelitsch A, Hoffmann D, Wernike K, Beer M. Occurrence of Antibodies against SARS-CoV-2
in the Domestic Cat Population of Germany. Vaccines (Basel). 2020;8(4).
23. Michelitsch A, Schön J, Hoffmann D, Beer M, Wernike K. The Second Wave of SARS-CoV-2
Circulation—Antibody Detection in the Domestic Cat Population in Germany. Viruses.
2021;13(6):1009.
24. Schulz C, Martina B, Mirolo M, Müller E, Klein R, Volk H, et al. SARS-CoV-2–Specific Antibodies
in Domestic Cats during First COVID-19 Wave, Europe. Emerging Infectious Disease journal.
2021;27(12).
25. Zhang Q, Zhang H, Gao J, Huang K, Yang Y, Hui X, et al. A serological survey of SARS-CoV-2 in
cat in Wuhan. Emerg Microbes Infect. 2020;9(1):2013-9.
26. Bosco-Lauth AM, Hartwig AE, Porter SM, Gordy PW, Nehring M, Byas AD, et al. Experimental
infection of domestic dogs and cats with SARS-CoV-2: Pathogenesis, transmission, and response to
reexposure in cats. Proceedings of the National Academy of Sciences. 2020;117(42):26382-8.
27. Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, et al. Susceptibility of ferrets, cats, dogs,
and other domesticated animals to SARS-coronavirus 2. Science. 2020;368(6494):1016-20.
28. Legros V, Denolly S, Vogrig M, Boson B, Siret E, Rigaill J, et al. A longitudinal study of SARS-
CoV-2-infected patients reveals a high correlation between neutralizing antibodies and COVID-19
severity. Cellular & Molecular Immunology. 2021;18(2):318-27.
29. R Development Core Team. R: A language and environment for statistical computing. Vienna,
Austria: R Foundation for Statistical Computing; 2018.
30. Ferasin L, Fritz M, Ferasin H, Becquart P, Corbet S, Ar Gouilh M, et al. Infection with SARS-
CoV-2 variant B.1.1.7 detected in a group of dogs and cats with suspected myocarditis. Vet Rec.
2021;189(9):e944.
31. Garcia-Beltran WF, Lam EC, Astudillo MG, Yang D, Miller TE, Feldman J, et al. COVID-19-
neutralizing antibodies predict disease severity and survival. Cell. 2021;184(2):476-88.e11.
32. Calvet GA, Pereira SA, Ogrzewalska M, Pauvolid-Corrêa A, Resende PC, Tassinari WdS, et al.
Investigation of SARS-CoV-2 infection in dogs and cats of humans diagnosed with COVID-19 in Rio de
Janeiro, Brazil. PLOS ONE. 2021;16(4):e0250853.
33. Damas J, Hughes Graham M, Keough Kathleen C, Painter Corrie A, Persky Nicole S, Corbo M,
et al. Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in
vertebrates. Proceedings of the National Academy of Sciences. 2020;117(36):22311-22.
34. Maurin M, Fenollar F, Mediannikov O, Davoust B, Devaux C, Raoult D. Current Status of
Putative Animal Sources of SARS-CoV-2 Infection in Humans: Wildlife, Domestic Animals and Pets.
Microorganisms. 2021;9(4).

35.   Jairak W, Charoenkul K, Chamsai E, Udom K, Chaiyawong S, Hangsawek A, et al. Survey of
SARS-CoV-2 in dogs and cats in high-risk areas during the second wave of COVID-19 outbreak,
Thailand. Zoonoses and Public Health. 2022.

36.   Bechmann N, Barthel A, Schedl A, Herzig S, Varga Z, Gebhard C, et al. Sexual dimorphism in
COVID-19: potential clinical and public health implications. The Lancet Diabetes & Endocrinology.
2022.

37.   Adler JM, Weber C, Wernike K, Michelitsch A, Friedrich K, Trimpert J, et al. Prevalence of anti-
severe acute respiratory syndrome coronavirus 2 antibodies in cats in Germany and other European
countries in the early phase of the coronavirus disease-19 pandemic. Zoonoses and Public
Health.n/a(n/a).

38.   Decaro N, Grassi A, Lorusso E, Patterson EI, Lorusso A, Desario C, et al. Long-term persistence
of neutralizing SARS-CoV-2 antibodies in pets. Transboundary and emerging diseases. 2021.

39.   Bashor L, Gagne RB, Bosco-Lauth AM, Bowen RA, Stenglein M, VandeWoude S. SARS-CoV-2
evolution in animals suggests mechanisms for rapid variant selection. Proceedings of the National
Academy of Sciences. 2021;118(44):e2105253118.

469