A large-scale serological survey in pets from October 2020 through June 2021 in France

shows significantly higher exposure to SARS-CoV-2 in cats

Matthieu Fritz[a], Eric Elguero[a,1], Pierre Becquart[a,1], Daphné de Riols de Fonclare[a], Déborah Garcia[a],

Stephanie Beurlet [b], Solène Denolly [c], Bertrand Boson [c], Serge G. Rosolen [e, f], François-Loïc Cosset[c],

Alexandra Briend-Marchal[b], Vincent Legros[c, d], Eric M. Leroy[a*]

[a] Maladies Infectieuses et Vecteurs, Ecologie, Génétique, Evolution et Contrôle (MIVEGEC), Univ.

Montpellier, IRD, CNRS, 34394 Montpellier, France

[b] Laboratoire de Biologie vétérinaire VEBIO, Arcueil, France

[c] CIRI – Centre International de Recherche en Infectiologie, Team EVIR, Univ Lyon, Université Claude

Bernard Lyon 1, Inserm, U1111, CNRS, UMR5308, ENS Lyon, Lyon, France.

[d] Université de Lyon, VetAgro Sup, Campus vétérinaire de Lyon, Marcy-l'Etoile, France

[e] Sorbonne Université, INSERM, CNRS, Institut de la Vision, Paris, France

[f] Clinique vétérinaire voltaire, Asnières, France

[1] Authors contributed equally

*Corresponding author:

Eric M.Leroy : UMR IRD-CNRS-UM, 911 avenue Agropolis, 34394 Montpellier, France

Tel : +33 467416107 ; E-mail address : eric.leroy@ird.fr

Keywords: SARS-CoV-2, Pets, Serology, Neutralization assay, Microsphere Immunoassay, One Health

Abstract

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can infect many animals, including

pets such as dogs and cats. Many studies have documented infection in companion animals by bio-

molecular and serological methods. However, only a few have compared seroprevalence in cats and

dogs from the general population, and these studies were limited by small sample sizes and

collections over short periods. Our goal was to obtain a more accurate evaluation of seroprevalence

in companion animals in France and to determine whether cats and dogs differ in their exposure to

SARS-CoV-2. For this purpose, we conducted an extensive SARS-CoV-2 serological survey of 2036 cats

and 3577 dogs sampled by veterinarians during medical examinations in clinics throughout France.

Sampling was carried out from October 2020 through June 2021, a period encompassing the second

and third waves of SARS-CoV-2 infections in humans in the country. Using a microsphere

immunoassay targeting the receptor binding domain and trimeric spike protein, we found 7.1%

seroprevalence in pets. In a subset of 308 seropositive samples, 26.3% had neutralizing antibodies.

We found that cats were significantly more likely to test positive than dogs, with seropositivity rates

of 9.3% and 5.9% in cats and dogs, respectively. Finally, data for both species showed that

seroprevalence was lower in older animals and was not associated with the date of sampling or the

sex of the animal. Our results show that cats are significantly more sensitive to SARS-CoV-2 than

dogs, in line with experimental studies. Our large sample size provides for a reliable, statistically

robust estimate of the frequency of infection of pets from their owners and offers strong support for

the notion that cats are more sensitive to SARS-CoV-2 than dogs. Our findings emphasise the

importance of a One-Health approach to the SARS-CoV-2 pandemic and raise the question of

whether companion animals in close contact with humans should be vaccinated.

Introduction

Two months after the onset of SARS-CoV-2 circulation in humans, two dogs in Hong Kong were

reported to have naturally acquired the virus (Sit et al., 2020). Since then, many studies have

reported viral RNA and SARS-CoV-2 antibodies in dogs and cats—mostly belonging to COVID-19-

infected owners (Barrs et al., 2020; Hamer et al., 2021; Jairak et al., 2021; Krafft et al., 2021).

Furthermore, several studies demonstrated that the risk of pets testing seropositive was higher in

COVID-19+ households than for pets from households of unknown status (Patterson et al., 2020;

Colitti et al., 2021; Fritz et al., 2021; Stevanovic et al., 2021; Barroso et al., 2022).

Definitive examples of pet-to-human transmission are scarce. A recent study from Thailand reported

a suspected case of SARS-CoV-2 transmission from a cat to a human (Sila et al., 2022), and dog-to-

human transmission has yet to be described. However, given that 200 million cats and dogs live in

close proximity to humans in Europe (The European Pet Food Industry, 2020), there is ample

opportunity for such transmission, and the potential risks need to be carefully considered.

Several population-based serological studies have reported SARS-CoV-2 antibodies in dogs and cats.

In dogs, estimates of seroprevalence have ranged from 0% to 14.5% (Stevanovic et al., 2020;

Barroso-Arévalo et al., 2021; Barua et al., 2021; Dileepan et al., 2021; Laidoudi et al., 2021;

Pomorska-Mól et al., 2021; Smith et al., 2021; Stevanovic et al., 2021; Udom et al., 2021; Zhao et al.,

2021; Barroso et al., 2022). While in cats, estimates have ranged from 0% to 21.7% (Michelitsch et

al., 2020; Zhang et al., 2020; Barua et al., 2021; Dileepan et al., 2021; Michelitsch et al., 2021; Schulz

et al., 2021; Udom et al., 2021; Barroso et al., 2022). For both species, seroprevalence was highly

dependent on the period of sampling (first, second wave *etc.*), the assay used (ELISA,

seroneutralization, etc.), and the country of sampling (China, Croatia, Germany, Italy, the

Netherlands, Poland, Portugal, Spain, Thailand, United-Kingdom, USA). Among these studies, five

directly compared cats and dogs. There is some experimental and epidemiological evidence

suggesting that cats are more susceptible to infection than dogs (Bosco-Lauth et al., 2020; Shi et al.,

2020; Colitti et al., 2021; Dileepan et al., 2021; Barroso et al., 2022). However, significant species

differences have not always been observed in population-based studies (Stevanovic et al., 2020;

Barroso-Arévalo et al., 2021; Pomorska-Mól et al., 2021). This is perhaps because of significant study

limitations—a low number of enrolled animals, a short sampling period, etc.—that have curtailed

robust estimates of infection frequency in pets with enough statistical power to recognize

differences in COVID-19 epidemiology.

Here we report estimates of the frequency of SARS-CoV-2 infection from 2036 cats and 3577 dogs

sampled at veterinary clinics from October 2020 through June 2021 throughout France—the largest

serological survey of SARS-CoV-2 infections in companion animals to date. The study allows for

robust estimates of pet infection frequency and provides strong support for the hypothesis that

species differences in susceptibility observed in experimental studies translate into a significant

increase in infection frequency in cats.

Materials and Methods

Sampling

The cross-sectional, nationwide sampling was possible thanks to a network of veterinary clinics

across France working with VEBIO. VEBIO is a veterinary diagnostic laboratory which performed all

categories of medical analyses, including infectious diseases, hematology, endocrinology, oncology …

(see more details in https://www.vebio.fr/). No inclusion and exclusion criteria were applied for the

selection of blood samples, except that they came only from veterinary clinics working with VEBIO.

VEBIO notified the veterinary clinics that following requested biomedical analyses, the remaining

serum could be used in a SARS-CoV-2 research project. No specific request for samples was

addressed to the vets. Thus, the SARS-CoV-2 analysis is based on samples collected during the regular

activities of the vets.

Blood samples were collected in dry/EDTA tubes from dogs and cats during routine healthcare visits

or for diagnostic purposes at veterinary clinics. After centrifugation, the serum/plasma was kept at

+4 °C until sent to VEBIO. Rapid and safe shipping practices were used to avoid contamination and

ensure samples reached VEBIO within 48 h. At the VEBIO facility, an aliquot was taken from the

sample to perform the requested biomedical analyses. Another aliquot was then stored at +4 °C until

sent to the MIVEGEC lab, Montpellier, where serological analyses were performed. Safe shipping

practices with an approved professional carrier were also used for shipment to the MIVEGEC lab.

Finally, the samples were stored at the MIVEGEC lab at −20 °C until testing (Figure 1). For shipping to

the CIRI lab, SARS-CoV-2-positive samples detected by MIA were transported by an approved

professional carrier at -20°C to ensure optimal safety conditions. Data (age, sex, clinical history, and

region localization, when available) from dogs and cats were provided anonymized by VEBIO to the

MIVEGEC lab.

Figure 1. Logistics of sample collection and distribution. Sera collected during routine healthcare

visits by veterinarians throughout France were first sent to VEBIO in Ile-de-France. Aliquots of the

samples were then made and sent to the IRD in Montpellier (Hérault) via an approved carrier.

Ethics

According to the act governing the "use of live animals for scientific purposes" effective in France on

14 January 2022, ethical approval was not sought or required since all pets were sampled by a

veterinarian during a health care visit. All applicable international and national guidelines for the care

of pets were followed.

Microsphere Immunoassay (MIA)

Dog and cat serum samples were tested using a multiplex microsphere immunoassay (MIA). Ten µg

of two recombinant SARS-CoV-2 antigens, receptor-binding domain (RBD) and trimeric spike (tri-S),

both derived from the Whuhan-Hu-1 strain (The Native Antigen Company, Kidlington United-

Kingdom), were used to capture specific serum antibodies. Distinct MagPlex microsphere sets

(Luminex Corp, Austin, TX, USA) were respectively coupled to viral antigens using the amine coupling

kit (Bio-Rad Laboratories, Marnes-la-Coquette, France) according to the manufacturer's instructions.

Microsphere mixtures were successively incubated with serum samples (1:400), biotinylated protein

A and biotinylated protein G (4 µg/mL each) (Thermo Fisher Scientific, Illkirch, France), and

streptavidin-R-phycoerythrin (4 µg/mL) (Life technologies, Illkirch, France) on an orbital shaker and

protected from light. Measurements were performed using a Luminex 200 instrument (Luminex

Corp, Austin, TX, USA), and at least 100 events were read for each bead set. Binding events were

displayed as median fluorescence intensities (MFI). Specific seropositivity cut-off values for each

antigen were set at three standard deviations above the mean MFI of pre-pandemic serum from 53

dogs and 30 cats sampled before 2019. These samples were stored in biobanks at the IRD and

VetAgro Sup. MIA specificity was set for each antigen at 96.2% for dogs and 100% for cats based on

the pre-pandemic populations. MIA was first validated using sera from two COVID-19 PCR+ humans,

kindly provided by Meriadeg Ar Gouilh, and then with sera from SARS-CoV-2 PCR+ cats and dogs,

provided by several veterinarians.

Because of the excellent specificity observed for both antigens and to account for any isotypic

variability, an animal was deemed positive for SARS-CoV-2 antibodies following a positive result in at

least one of the two tests.

Neutralization activity measurement

An MLV-based pseudoparticle carrying a GFP reporter pseudotyped with SARS-CoV-2 spike protein

(Wuhan-Hu-1 strain) (SARS-CoV-2pp) was used to measure neutralizing antibody activity in cat and

dog sera. Each SARS-CoV-2-positive sample detected by MIA was processed according to a

neutralization procedure as previously described (Legros et al., 2021). Briefly, for neutralization

assays, a sample of ~1×10^3 pseudoparticles was incubated with a 100-fold dilution of sera or

control antibodies for 1 h at 37 °C before infection of Vero-E6R cells. At 72 h post-transduction, the

percentage of GFP-positive cells was determined by flow cytometry (at least 10 000 events

recorded). The level of infectivity is expressed as the percentage of GFP-positive cells and compared

to cells infected with SARS-CoV-2pp incubated without serum. As a control, the same procedure was

done using RD114 pseudoparticles to identify sera with aspecific neutralization. Sera exhibiting more

than 30% SARS-CoV-2pp neutralization were considered positive. Pre-pandemic serum from France

was used as a negative control, and an anti-SARS-CoV-2 RBD antibody was used as a positive control.

Statistical analyses

Associations between SARS-CoV2 infection status (positive or negative) and the covariates region,

age, and sex were assessed using binomial (logistic) generalized linear models. The region was

defined by where the animal lived at the time of sampling. Age was that recorded by the

veterinarian, with variable precision, generally in months for young animals and whole years for

older animals. Its accuracy is unknown. The associated statistical tests were likelihood ratio tests. All

analyses were performed using R software (R Development Core Team, 2018).

Results

Blood collection

Blood samples from 2036 cats and 3577 dogs were collected during routine healthcare visits by

veterinarians from October 2020 through June 2021 (Table 1). Samples came from all 13 regions of

metropolitan France. Corsica was excluded due to too few samples. Almost half of the samples came

from Ile-de-France, the region including Paris, reflecting population density and proximity to the

Veterinary diagnostic laboratory (VEBIO), where all samples for biomedical analyses requested by the

veterinarians were handled (Materials and Methods). The number of samples received from other

regions largely depended on the number of veterinarians working with VEBIO in those regions (Figure

2). Unfortunately, we could not study the clinical history of the animals due to variability in how each

veterinarian reported this information.

Table 1 : numbers of samples collected by month and by species.

	10/2020	11/2020	12/2020	01/2021	02/2021	03/2021	04/2021	05/2021	06/2021
cats	49	256	275	291	225	296	305	166	173
dogs	84	428	543	475	403	474	597	282	291
all	133	684	818	766	628	770	902	448	464

Figure 2. (a). Map of France showing the number of SARS-CoV-2-positive cat sera per region. The

total number of sera samples collected per region is indicated. Seroprevalence in each region is

indicated as a percentage. Regions are shaded in green according to seroprevalence. The total

number of sera samples and global seroprevalence for France is in the top left corner. **(b). Map of**

France showing the number of SARS-CoV-2-positive dog sera per region. The total number of sera

samples collected per region is indicated. Seroprevalence in each region is indicated as a percentage.

Regions are shaded in orange according to seroprevalence. The total number of sera samples and

global seroprevalence for France is in the top left corner.

Global seroprevalence

For the sera samples, 401 (7.1%) showed a positive result either against RBD, tri-S, or both

(Supplementary Table 1). We next determined the presence of antibodies with neutralizing activity

among the positive sera. To save time, we randomly tested approximately 75% (308) of positive sera

samples using a retrovirus-based pseudoparticle assay detecting SARS-CoV-2 neutralizing antibodies

(Materials and Methods). Pets were considered positive if they showed at least 30% neutralization.

Seroneutralizing activity was detected in 81 (26.3%) of the 308 pet sera samples. Among these

positive samples, 39 (48%) were positive for both RBD and tri-S, 39 (48%) were positive only for tri-s

and 3 (4%) were only positive for RBD. Previous studies have shown that some pets do not develop

neutralizing antibodies (Ferasin et al., 2021; Krafft et al., 2021); therefore, only the seroprevalence

from MIA assays was analyzed in the remainder of the study.

Seroprevalence in cats and dogs

We observed that a significantly greater proportion of cats were positive (189/2036, 9.3%) than dogs

(212/3577, 5.9%); OR = 1.62, 95% c.i. [1.32 - 1.99], P-value = 3.8e-06, Table 1. In addition, sera from

MIA-positive cats were more likely to show neutralizing activity (49/144, 34%) than dogs (32/164,

19.5%); OR = 2.12, 95% c.i. [1.27 - 3.57], P-value = 0.0039). Species differences were not always

significant within each region, likely due to reduced statistical power. However, when differences

were significant, it was always the case that cats were more likely to be positive than dogs. (Table 2).

Table 2. Seroprevalence of IgG SARS-CoV-2 antibodies detected in blood samples from cats and dogs

collected in different French regions from October 2020 through June 2021. Data are presented as

No. positive, percentage (95 % exact binomial confidence intervals). Odds ratios were computed by

fitting binomial models region by region; an OR > 1 indicates cats were more likely to be positive than

dogs. In this analysis, individual data on pet age and sex were not considered, as age was not

available for all animals. P-values were computed by the likelihood ratio test.

Region	Cats	Cats seroprevalence		Dogs	Dogs seroprevalence		OR (95% c.i.)	P-value
Auvergne-Rhône-Alpes	4/92	4.3%	(1.2-10.8)	25/298	8.4%	(5.5-12.1)	0.50 (0.17 - 1.47)	0.17
Bourgogne-Franche-Comté	3/49	6.1%	(1.3-16.9)	4/91	4.4%	(1.2-10.9)	1.42 (0.30 - 6.61)	0.66
Bretagne	3/38	7.9%	(1.7-21.4)	5/113	4.4%	(1.5-10.0)	1.85 (0.42 - 8.14)	0.43
Centre-Val de Loire	7/26	26.9%	(11.6-47.8)	3/53	5.7%	(1.2-15.7)	6.14 (1.44 - 26.2)	0.0098
Grand Est	8/42	19.0%	(8.6-34.1)	5/172	2.9%	(1.0-6.7)	7.86 (2.42 - 25.5)	0.00057
Hauts-de-France	6/91	6.6%	(2.5-13.8)	16/211	7.6%	(4.4-12.0)	0.86 (0.33 - 2.27)	0.76
Île-de-France	104/1193	8.7%	(7.2-10.5)	98/1248	7.9%	(6.4-9.5)	1.12 (0.84 - 1.49)	0.44
Normandie	3/62	4.8%	(1.0-13.5)	8/144	5.6%	(2.4-10.7)	0.86 (0.22 - 3.37)	0.83
Nouvelle-Aquitaine	21/213	9.9%	(6.2-14.7)	20/611	3.3%	(2.0-5.0)	3.23 (1.72 - 6.09)	0.00037
Occitanie	8/92	8.7%	(3.8-16.4)	12/386	3.1%	(1.6-5.4)	2.97 (1.18 - 7.49)	0.028
Provence-Alpes-Côte d'Azur	19/86	22.1%	(13.9-32.3)	12/182	6.6%	(3.5-11.2)	4.02 (1.85 - 8.73)	0.00036
Pays de la Loire	3/52	5.8%	(1.2-15.9)	4/68	5.9%	(1.6-14.4)	0.98 (0.21 - 4.58)	0.98
Total	**189/2036**	**9.3%**	**(8.0 - 10.6)**	**212/3577**	**5.9%**	**(5.2 - 6.8)**	**1.62 (1.32 - 1.99)**	**3.8×10^{-6}**

Seroprevalence by sex

We found no significant sex differences in seropositivity rates, either for all animals (females: 6.9%;

163/2361; males 7.5%; 212/2842; p = 0.24) or among cats (females 9.4%; 78/827, males 9.8%;

99/1009, p = 0.68) and dogs (females: 5.5%; 85/1534, males: 6.2%; 113/1833; p = 0.27) tested

separately (Table 3).

Table 3. Seroprevalence of IgG SARS-CoV-2 antibodies in blood samples from cats and dogs by sex

from October 2020 through June 2021 Data are presented as No. positive, percentage (95 % exact

binomial confidence intervals). Odds ratios > 1 indicate males are more likely to be positive than

females and were computed by fitting binomial generalized linear models, with age as a controlling

factor. P-values correspond to likelihood ratio tests.

Sex	Cats	Cat seroprevalence	Dogs	Dog seroprevalence	Cats + Dogs	Cats + Dogs seroprevalence
Female	78/827	9.4% (7.5-11.5)	85/1534	5.5% (4.4-6.8)	163/2361	6.9% (5.9-8.0)
Male	99/1009	9.8% (8.0-11.8)	113/1833	6.2% (5.1-7.4)	212/2842	7.5% (6.5-8.5)
Total	**177/1836**	**9.6% (8.3-11.1)**	**198/3367**	**5.9% (5.1-6.7)**	**375/5203**	**7.2% (6.5-7.9)**
OR	1.08 (0.75 - 1.54)		1.20 (0.87 - 1.67)		1.15 (0.91 - 1.47)	
P-value	0.68		0.27		0.24	

Seroprevalence by age

Age was reported for 1657 cats (range: 0.2 – 22yr) and 2781 dogs (range: 0.1 – 18.5yr). Among cats,

18.4% aged 0-3 years, 10.4% aged 3-9 years, and 6.4% aged over nine years tested positive.

Among dogs, 8.8% aged 0-3 years, 5.8% aged 3-9 years, and 5.1% aged over 9 years tested positive.

Using a binomial model with age entered as a continuous variable, we observed a significant

decrease in seroprevalence with age in cats (OR for a one-year increase in age = 0.91, 95% c.i. [0.88 -

0.94], p-value = 3.7e-08) and dogs (OR = 0.95, 95% c.i. [0.92 - 0.99], p-value = 0.016) (Figure 3)

(Supplementary Table 2).

Figure 3. (a). The number of cat blood samples tested by age group for anti-SARS-CoV-2 antibodies

by MIA from October 2020 through June 2021. Samples testing negative are shaded grey, and

seropositive samples are in orange. Seroprevalence is represented by black dots, with 95 % binomial

confidence interval. The red line represents the linear regression. **(b). The number of dog blood**

samples tested by age group for anti-SARS-CoV-2 antibodies by MIA from October 2020 through

June 2021. Samples testing negative are shaded grey, and seropositive samples are in orange.

Seroprevalence is represented by black dots, with 95 % binomial confidence intervals. The red line

represents the linear regression.

Seroprevalence over time

We next examined whether seroprevalence was associated with the time of sampling. For this

analysis, we selected animals at least one year old at the date of sampling (Figure 4). Seroprevalence

was not associated with the time of sampling for cats: p-value = 0.41. However, seroprevalence

among dogs increased over the 9 months of the study (OR = 3.47, 95% c.i. [1.47 - 8.23], p-value =

0.0045).

(a)

(b)

Figure

4. (a). The number of cat blood samples tested each month for anti-SARS-CoV-2 antibodies by MIA

from October 2020 through June 2021. Samples testing negative are shaded grey, and seropositive

samples are in orange. Seroprevalence is represented by black dots, with 95 % binomial confidence

interval. The red line represents the linear regression. **(b). The number of dog blood samples tested**

each month for anti-SARS-CoV-2 antibodies by MIA from October 2020 through June 2021. Samples

testing negative are shaded grey, and seropositive samples are in orange. Seroprevalence is

represented by black dots, with 95 % binomial confidence interval. The red line represents the linear

regression. Notice that in this figure, dates have been pooled by calendar month for illustrative

purposes but that in the statistical analysis exact dates were used. Likewise, the regression lines are

similarly illustrative as the statistical tests were based on logistic (binomial) regression.

Discussion

This study reports a large-scale serological survey of pet (cats and dogs) to detect anti-SARS-CoV-2

IgG antibodies. The samples were collected in metropolitan France from October 2020 through June

2021, a period including peaks of the second and third waves of SARS-CoV-2 infections in France.

From a sample of 5613 pets, we report a seroprevalence of anti-SARS-CoV-2 antibodies of 7.1%. We

observed that only a small percentage of samples (48%) were positive for both tri-s and RBD,

indicating that the RBD assay may be less sensitive than the tri-s assay. This may be explained by the

fact that the full trimeric spike antigen may bind a broader range of antibodies than the receptor-

binding domain, which includes only a small part of the spike protein. We found neutralizing

antibody activity in the sera of only 26% of seropositive pets. Cats were more likely to produce

neutralizing antibodies than dogs, which is likely associated with a more prolonged and intense

immune stimulation in cats. In humans, disease severity is positively correlated with neutralizing

antibody levels (Garcia-Beltran et al., 2021).

In cats, we found a higher seroprevalence (9.3%) than previously observed in other European

countries, which ranged from 0% to 6.4% (Michelitsch et al., 2020; Stevanovic et al., 2020;

Barroso-Arévalo et al., 2021; Michelitsch et al., 2021; Pomorska-Mól et al., 2021; Schulz et al., 2021;

Smith et al., 2021; Zhao et al., 2021). However, most of these studies were done before the second

wave, during a period of relatively lower viral circulation than our sampling period. In addition, most

of these studies used a seroneutralisation assay.

In dogs, the observed seroprevalence (5.9%) is in accord with a previous study in France showing a

prevalence of 4.8% in companion and military working dogs sampled between February 2020 and

February 2021 (Laidoudi et al., 2021). Other studies looking for SARS-CoV-2 antibodies in dogs have

reported seroprevalences ranging from 0% to 14.5% (Stevanovic et al., 2020; Barroso-Arévalo et al.,

2021; Pomorska-Mól et al., 2021; Smith et al., 2021; Stevanovic et al., 2021; Zhao et al., 2021).

Importantly, we observed a significantly higher seroprevalence of anti-SARS-CoV-2 antibodies in cats

than in dogs (p = 4.2e-08). The statistical significance of this difference varied among regions, likely

due to the reduced power and perhaps some unintended sampling bias by veterinarians. For

example, the smallest sample size was in the Bourgogne-Franche-Comté region, where we observed

no significant difference between dogs and cats. Furthermore, for a region like Ile-de-France, where

people live mostly in apartments, we can also hypothesize that dogs live in closer contact with

owners than in the rest of France. Previous studies with fewer samples have either found no

significant difference between species (Patterson et al., 2020; Stevanovic et al., 2020;

Barroso-Arévalo et al., 2021; Calvet et al., 2021; Fritz et al., 2021; Pomorska-Mól et al., 2021) or that

cats have significantly higher seroprevalence than dogs (Colitti et al., 2021; Hamer et al., 2021;

Barroso et al., 2022).

Our study of a very large population of dogs and cats in natural conditions provides some evidence

that cats are more susceptible to SARS-CoV-2 infection than dogs, at least during the time frame of

our sampling period. Potential causes of species differences in susceptibility between cats and dogs

are numerous, but likely include a variety of biological and behavioural factors, as well as differences

in exposure. Intererestingly, ACE-2 shows greater sequence similarity between cat and human

orthologs than observed between dogs and humans (Damas et al., 2020). The absence of data such

as the pet lifestyle (Indoor/Outdoor), or the frequency and nature of contacts with humans and other

animals restricts our ability to identify a potential cause of the observed difference. In previously

studies, most infected pets were epidemiologically linked to humans who had tested positive for

COVID-19 (Maurin et al., 2021).

We did not observe significant sex differences in seroprevalence in either species (p = 0.45). Our

findings are consistent with most previous studies also reporting an absence of sex differences in

dogs and cats (Calvet et al., 2021; Pomorska-Mól et al., 2021; Barroso et al., 2022; Jairak et al., 2022).

A smaller study of 188 dogs and 61 cats found higher seropositivity in male dogs and an absence of a

sex difference in cats (Patterson et al., 2020). Another study found that male dogs sampled from the

general population were more likely to test positive than females, but this difference was not

observed in dogs from COVID-19+ households (Stevanovic et al., 2021). There is little evidence of a

significant sex difference in susceptibility in humans. However, men are more likely to be affected by

severe forms of COVID than women for a variety of reasons (Bechmann et al., 2022).

In terms of age, we observed a higher seroprevalence among younger animals (between 0-3 years)

for both species that then decreased with age. A study of dogs sampled from the general population

found seroprevalence was highest in animals aged 5-6 years and that in COVID-19+ households,

seroprevalence peaked in slightly younger dogs, aged between one and five years (Stevanovic et al.,

2021). Other studies have reported no significant associations with age in cats and dogs (Pomorska-

Mól et al., 2021; Barroso et al., 2022). An experimental study in cats found that juveniles appear

more vulnerable than subadults (Shi et al., 2020). The decreasing seroprevalence we observed with

age could also arise from age-dependent behavioural changes. For example, young animals (< 3 years

old) are more active and curious and may be in greater contact with their owners than older animals

that prefer to remain quieter. The decrease could also reflect immunosenescence in older animals, as
observed in humans.

Interestingly, we observed a slight increase in seroprevalence in dogs during the study's nine months
of sampling, a trend not observed among cats. We expected an increase because antibodies have a
longer persistence in the organism than viral RNA; thus, animals sampled at later dates would
represent an accumulation of cases. The absence of a positive association between seroprevalence
and the time of sampling in cats has been reported in two other studies in Europe, but conclusions
were limited by the small number of samples collected over just a few months (Adler et al.; Schulz et
al., 2021). The absence of an association in cats suggests a limited persistence of antibodies in cats
than dogs. Few studies have investigated variation in the persistence of antibodies in animals. For
example, a study carried out on seven dogs and two cats infected in natural conditions showed
persistence of neutralizing antibodies up to 10 months after infection in four of the dogs and the two
cats, but also that persistence was markedly reduced in two of the dogs after three months (Decaro
et al., 2021). Moreover, a study of two cats found that neutralizing antibodies had disappeared by
110 days (Zhang et al., 2020). Based on these data, one possible reason for the lack of increase in
seroprevalence during our study period could be a progressive seroreversion of infected cats that is
equally compensated by the number of new infections, i.e. seroconversion. If so, this would mean
that the observed seroprevalence is not an accurate reflection of the total number of infections, at
least in cats, during the whole epidemic. Instead, seroprevalence provides a snapshot if infections
acquired during a time period. This also suggests that the seroprevalence observed in our study may
underestimate the actual proportion of cats infected during the entirety of the epidemic. In order to
enhance our comprehension of antibody persistence in cats and dogs, a longitudinal serological study
involving a larger number of cats and dogs has to be conducted.

Human-to-pet transmission may promote viral adaptation facilitating re-infection with novel viral
strains in humans (Bashor et al., 2021). While one case of infection from cat to human has recently

been reported, the large number of pet cats and their frequent close interaction with humans

provides ample opportunity. This possibility raises the question of a vaccination strategy for animals

susceptible to SARS-CoV-2 infection. While pets do not currently seem to play a role in the ongoing

pandemic, our results emphasize the magnitude of SARS-CoV-2 infection in pets is not trivial.

Combined with the size of domestic cat and dog populations and the close contact with their human

companions, our results highlight the importance of collecting more data on SARS-CoV-2

transmissibility and pathogenicity in companion animals, especially with the emergence of new

variants. Also, when a SARS-CoV-2 infection is suspected in a pet, we suggest collecting a sample for

RT-qPCR confirmation of infection, followed by whole-genome sequencing to identify new mutations,

particularly in antigenic sites targeted by the immune system. Finally, similar public health

recommendations applied to humans should also be implemented for animals to prevent human-to-

animal transmission, such as not having contact with animals when a household member is COVID-19

positive.

Acknowledgements: We are grateful to the pet owners for giving us permission to sample their pets.

We thank the veterinarians that helped us with sampling. We thank Simon Thierry and Trung Thanh

Nguyen for technical support and assistance. We also thank Kurt McKean

(https://octopusediting.com/) for the English editing of the manuscript.

Funding: The study was funded by the French National Agency for Research (ANR-RA-COVID-19;

Geographical and temporal serological investigation of companion animal infection with SARS-CoV-2

during the second wave of COVID-19 in France, CoVet). This study received financial support through

a research grant from IDEXLYON project of Université de Lyon as part of the "Programme

Investissements d'Avenir" (ANR-16-IDEX-0005). The study was funded by OIE through the European

Union "EBO-SURSY" project.

Conflicts of Interest: The authors declare no conflicts of interest.

Institutional Review Board Statement: According to the act of "use of live animals for scientific

purposes" effective in France on 14 January 2022, ethical approval was not sought or required since

all pets were sampled by a veterinarian during a health care visit. All applicable international and

national guidelines for the care of pets were followed.

Informed Consent Statement: Not applicable.

Data Availability Statement: Raw data and scripts are available on zenodo:

https://doi.org/10.5281/zenodo.8125085

Author Contributions:

M.F., P.B., S.G.R., A.B.-M., V.L., and E.M.L. conceived and designed the study.

M.F., E.E., D.d.R.d.F., D.G., S.D., B.B., and V.L designed and performed the experiments.

All authors analyzed the data and interpreted and discussed the results.

M.F. and E.L. wrote the manuscript with input from all authors.

All authors have read and agreed to the published version of the manuscript.

References

Adler, J. M., Weber, C., Wernike, K., Michelitsch, A., Friedrich, K., Trimpert, J., Beer, M., Kohn, B.,
Osterrieder, K., & Müller, E. Prevalence of anti-severe acute respiratory syndrome
coronavirus 2 antibodies in cats in Germany and other European countries in the early phase
of the coronavirus disease-19 pandemic. *Zoonoses and Public Health*, *n/a*(n/a).
https://doi.org/https://doi.org/10.1111/zph.12932
Barroso, R., Vieira-Pires, A., Antunes, A., & Fidalgo-Carvalho, I. (2022). Susceptibility of Pets to SARS-
CoV-2 Infection: Lessons from a Seroepidemiologic Survey of Cats and Dogs in Portugal.
Microorganisms, *10*(2), 345. https://www.mdpi.com/2076-2607/10/2/345
Barroso-Arévalo, S., Barneto, A., Ramos, Á. M., Rivera, B., Sánchez, R., Sánchez-Morales, L.,
Pérez-Sancho, M., Buendía, A., Ferreras, E., Ortiz-Menéndez, J. C., Moreno, I., Serres, C., Vela,
C., Risalde, M. Á., Domínguez, L., & Sánchez-Vizcaíno, J. M. (2021). Large-scale study on
virological and serological prevalence of SARS-CoV-2 in cats and dogs in Spain. *Transbound
Emerg Dis*. https://doi.org/10.1111/tbed.14366
Barrs, V., Peiris, M., Tam, K. W. S., Law, P. Y. T., Brackman, C., To, E. M. W., Yu, V. Y. T., Chu, D. K. W.,
Perera, R. A. P. M., & Sit, T. H. C. (2020). SARS-CoV-2 in Quarantined Domestic Cats from
COVID-19 Households or Close Contacts, Hong Kong, China. *Emerging Infectious Disease
journal*, *26*(12), 3071. https://doi.org/10.3201/eid2612.202786

Barua, S., Hoque, M., Adekanmbi, F., Kelly, P., Jenkins-Moore, M., Torchetti, M. K., Chenoweth, K., Wood, T., & Wang, C. (2021). Antibodies to SARS-CoV-2 in dogs and cats, USA. *Emerging Microbes & Infections*, 1-20. https://doi.org/10.1080/22221751.2021.1967101

Bashor, L., Gagne, R. B., Bosco-Lauth, A. M., Bowen, R. A., Stenglein, M., & VandeWoude, S. (2021). SARS-CoV-2 evolution in animals suggests mechanisms for rapid variant selection. *Proceedings of the National Academy of Sciences*, *118*(44), e2105253118. https://doi.org/doi:10.1073/pnas.2105253118

Bechmann, N., Barthel, A., Schedl, A., Herzig, S., Varga, Z., Gebhard, C., Mayr, M., Hantel, C., Beuschlein, F., Wolfrum, C., Perakakis, N., Poston, L., Andoniadou, C. L., Siow, R., Gainetdinov, R. R., Dotan, A., Shoenfeld, Y., Mingrone, G., & Bornstein, S. R. (2022). Sexual dimorphism in COVID-19: potential clinical and public health implications. *The Lancet Diabetes & Endocrinology*. https://doi.org/10.1016/s2213-8587(21)00346-6

Bosco-Lauth, A. M., Hartwig, A. E., Porter, S. M., Gordy, P. W., Nehring, M., Byas, A. D., VandeWoude, S., Ragan, I. K., Maison, R. M., & Bowen, R. A. (2020). Experimental infection of domestic dogs and cats with SARS-CoV-2: Pathogenesis, transmission, and response to reexposure in cats. *Proceedings of the National Academy of Sciences*, *117*(42), 26382-26388. https://doi.org/10.1073/pnas.2013102117

Calvet, G. A., Pereira, S. A., Ogrzewalska, M., Pauvolid-Corrêa, A., Resende, P. C., Tassinari, W. d. S., Costa, A. d. P., Keidel, L. O., da Rocha, A. S. B., da Silva, M. F. B., dos Santos, S. A., Lima, A. B. M., de Moraes, I. C. V., Mendes Junior, A. A. V., Souza, T. d. C., Martins, E. B., Ornellas, R. O., Corrêa, M. L., Antonio, I. M. d. S., . . . Menezes, R. C. (2021). Investigation of SARS-CoV-2 infection in dogs and cats of humans diagnosed with COVID-19 in Rio de Janeiro, Brazil. *PLoS One*, *16*(4), e0250853. https://doi.org/10.1371/journal.pone.0250853

Colitti, B., Bertolotti, L., Mannelli, A., Ferrara, G., Vercelli, A., Grassi, A., Trentin, C., Paltrinieri, S., Nogarol, C., Decaro, N., Brocchi, E., & Rosati, S. (2021). Cross-Sectional Serosurvey of Companion Animals Housed with SARS-CoV-2–Infected Owners, Italy. *Emerging Infectious Disease journal*, *27*(7). https://doi.org/10.3201/eid2707.203314

Damas, J., Hughes Graham, M., Keough Kathleen, C., Painter Corrie, A., Persky Nicole, S., Corbo, M., Hiller, M., Koepfli, K.-P., Pfenning Andreas, R., Zhao, H., Genereux Diane, P., Swofford, R., Pollard Katherine, S., Ryder Oliver, A., Nweeia Martin, T., Lindblad-Toh, K., Teeling Emma, C., Karlsson Elinor, K., & Lewin Harris, A. (2020). Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates. *Proceedings of the National Academy of Sciences*, *117*(36), 22311-22322. https://doi.org/10.1073/pnas.2010146117

Decaro, N., Grassi, A., Lorusso, E., Patterson, E. I., Lorusso, A., Desario, C., Anderson, E. R., Vasinioti, V., Wastika, C. E., Hughes, G. L., Valleriani, F., Colitti, B., Ricci, D., Buonavoglia, D., Rosati, S., Cavaliere, N., Paltrinieri, S., Lauzi, S., Elia, G., & Buonavoglia, C. (2021). Long-term persistence of neutralizing SARS-CoV-2 antibodies in pets. *Transbound Emerg Dis*. https://doi.org/10.1111/tbed.14308

Dileepan, M., Di, D., Huang, Q., Ahmed, S., Heinrich, D., Ly, H., & Liang, Y. (2021). Seroprevalence of SARS-CoV-2 (COVID-19) exposure in pet cats and dogs in Minnesota, USA. *Virulence*, *12*(1), 1597-1609. https://doi.org/10.1080/21505594.2021.1936433

Ferasin, L., Fritz, M., Ferasin, H., Becquart, P., Corbet, S., Ar Gouilh, M., Legros, V., & Leroy, E. M. (2021). Infection with SARS-CoV-2 variant B.1.1.7 detected in a group of dogs and cats with suspected myocarditis. *Vet Rec*, *189*(9), e944. https://doi.org/https://doi.org/10.1002/vetr.944

Fritz, M., Rosolen, B., Krafft, E., Becquart, P., Elguero, E., Vratskikh, O., Denolly, S., Boson, B., Vanhomwegen, J., Gouilh, M. A., Kodjo, A., Chirouze, C., Rosolen, S. G., Legros, V., & Leroy, E. M. (2021). High prevalence of SARS-CoV-2 antibodies in pets from COVID-19+ households. *One health (Amsterdam, Netherlands)*, *11*, 100192. https://doi.org/10.1016/j.onehlt.2020.100192

Garcia-Beltran, W. F., Lam, E. C., Astudillo, M. G., Yang, D., Miller, T. E., Feldman, J., Hauser, B. M., Caradonna, T. M., Clayton, K. L., Nitido, A. D., Murali, M. R., Alter, G., Charles, R. C., Dighe, A.,

Branda, J. A., Lennerz, J. K., Lingwood, D., Schmidt, A. G., Iafrate, A. J., & Balazs, A. B. (2021).
COVID-19-neutralizing antibodies predict disease severity and survival. *Cell*, *184*(2), 476-
488.e411. https://doi.org/10.1016/j.cell.2020.12.015

Hamer, S. A., Pauvolid-Corrêa, A., Zecca, I. B., Davila, E., Auckland, L. D., Roundy, C. M., Tang, W.,
Torchetti, M. K., Killian, M. L., Jenkins-Moore, M., Mozingo, K., Akpalu, Y., Ghai, R. R.,
Spengler, J. R., Barton Behravesh, C., Fischer, R. S. B., & Hamer, G. L. (2021). SARS-CoV-2
Infections and Viral Isolations among Serially Tested Cats and Dogs in Households with
Infected Owners in Texas, USA. *Viruses*, *13*(5), 938. https://www.mdpi.com/1999-
4915/13/5/938

Jairak, W., Charoenkul, K., Chamsai, E., Udom, K., Chaiyawong, S., Bunpapong, N., Boonyapisitsopa,
S., Tantilertcharoen, R., Techakriengkrai, N., Surachetpong, S., Tangwangvivat, R.,
Suwannakarn, K., & Amonsin, A. (2021). First cases of SARS-CoV-2 infection in dogs and cats
in Thailand. *Transbound Emerg Dis*. https://doi.org/10.1111/tbed.14383

Jairak, W., Charoenkul, K., Chamsai, E., Udom, K., Chaiyawong, S., Hangsawek, A., Waenkaew, S.,
Mungaomklang, A., Tangwangvivat, R., & Amonsin, A. (2022). Survey of SARS-CoV-2 in dogs
and cats in high-risk areas during the second wave of COVID-19 outbreak, Thailand. *Zoonoses*
and Public Health. https://doi.org/10.1111/zph.12907

Krafft, E., Denolly, S., Boson, B., Angelloz-Pessey, S., Levaltier, S., Nesi, N., Corbet, S., Leterrier, B.,
Fritz, M., Leroy, E. M., Gouilh, M. A., Cosset, F.-L., Kodjo, A., & Legros, V. (2021). Report of
One-Year Prospective Surveillance of SARS-CoV-2 in Dogs and Cats in France with Various
Exposure Risks: Confirmation of a Low Prevalence of Shedding, Detection and Complete
Sequencing of an Alpha Variant in a Cat. *Viruses*, *13*(9), 1759.
https://doi.org/https://doi.org/10.3390/v13091759

Laidoudi, Y., Sereme, Y., Medkour, H., Watier-Grillot, S., Scandola, P., Ginesta, J., Andréo, V., Labarde,
C., Comtet, L., Pourquier, P., Raoult, D., Marié, J. L., & Davoust, B. (2021). SARS-CoV-2
antibodies seroprevalence in dogs from France using ELISA and an automated western
blotting assay. *One health (Amsterdam, Netherlands)*, *13*, 100293.
https://doi.org/10.1016/j.onehlt.2021.100293

Legros, V., Denolly, S., Vogrig, M., Boson, B., Siret, E., Rigaill, J., Pillet, S., Grattard, F., Gonzalo, S.,
Verhoeven, P., Allatif, O., Berthelot, P., Pélissier, C., Thiery, G., Botelho-Nevers, E., Millet, G.,
Morel, J., Paul, S., Walzer, T., . . . Pozzetto, B. (2021). A longitudinal study of SARS-CoV-2-
infected patients reveals a high correlation between neutralizing antibodies and COVID-19
severity. *Cellular & Molecular Immunology*, *18*(2), 318-327. https://doi.org/10.1038/s41423-
020-00588-2

Maurin, M., Fenollar, F., Mediannikov, O., Davoust, B., Devaux, C., & Raoult, D. (2021). Current Status
of Putative Animal Sources of SARS-CoV-2 Infection in Humans: Wildlife, Domestic Animals
and Pets. *Microorganisms*, *9*(4). https://doi.org/10.3390/microorganisms9040868

Michelitsch, A., Hoffmann, D., Wernike, K., & Beer, M. (2020). Occurrence of Antibodies against
SARS-CoV-2 in the Domestic Cat Population of Germany. *Vaccines (Basel)*, *8*(4).
https://doi.org/10.3390/vaccines8040772

Michelitsch, A., Schön, J., Hoffmann, D., Beer, M., & Wernike, K. (2021). The Second Wave of SARS-
CoV-2 Circulation—Antibody Detection in the Domestic Cat Population in Germany. *Viruses*,
13(6), 1009. https://www.mdpi.com/1999-4915/13/6/1009

Patterson, E. I., Elia, G., Grassi, A., Giordano, A., Desario, C., Medardo, M., Smith, S. L., Anderson, E.
R., Prince, T., Patterson, G. T., Lorusso, E., Lucente, M. S., Lanave, G., Lauzi, S., Bonfanti, U.,
Stranieri, A., Martella, V., Solari Basano, F., Barrs, V. R., . . . Decaro, N. (2020). Evidence of
exposure to SARS-CoV-2 in cats and dogs from households in Italy. *Nature Communications*,
11(1), 6231. https://doi.org/10.1038/s41467-020-20097-0

Pomorska-Mól, M., Turlewicz-Podbielska, H., Gogulski, M., Ruszkowski, J. J., Kubiak, M., Kuriga, A.,
Barket, P., & Postrzech, M. (2021). A cross-sectional retrospective study of SARS-CoV-2
seroprevalence in domestic cats, dogs and rabbits in Poland. *BMC Veterinary Research*,
17(1), 322. https://doi.org/10.1186/s12917-021-03033-2

R Development Core Team. (2018). *R: A language and environment for statistical computing*. In R
Foundation for Statistical Computing. http://www.R-project.org
Schulz, C., Martina, B., Mirolo, M., Müller, E., Klein, R., Volk, H., Egberink, H., Gonzalez-Hernandez,
M., Kaiser, F., von Köckritz-Blickwede, M., & Osterhaus, A. (2021). SARS-CoV-2–Specific
Antibodies in Domestic Cats during First COVID-19 Wave, Europe. *Emerging Infectious
Disease journal*, *27*(12). https://doi.org/10.3201/eid2712.211252
Shi, J., Wen, Z., Zhong, G., Yang, H., Wang, C., Huang, B., Liu, R., He, X., Shuai, L., Sun, Z., Zhao, Y., Liu,
P., Liang, L., Cui, P., Wang, J., Zhang, X., Guan, Y., Tan, W., Wu, G., . . . Bu, Z. (2020).
Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2.
Science, *368*(6494), 1016-1020. https://doi.org/10.1126/science.abb7015
Sila, T., Sunghan, J., Laochareonsuk, W., Surasombatpattana, S., Kongkamol, C., Ingviya, T.,
Siripaitoon, P., Kositpantawong, N., Kanchanasuwan, S., Hortiwakul, T., Charernmak, B.,
Nwabor, O. F., Silpapojakul, K., & Chusri, S. (2022). Suspected Cat-to-Human Transmission of
SARS-CoV-2, Thailand, July-September 2021. *Emerg Infect Dis*, *28*(7), 1485-1488.
https://doi.org/10.3201/eid2807.212605
Sit, T. H. C., Brackman, C. J., Ip, S. M., Tam, K. W. S., Law, P. Y. T., To, E. M. W., Yu, V. Y. T., Sims, L. D.,
Tsang, D. N. C., Chu, D. K. W., Perera, R., Poon, L. L. M., & Peiris, M. (2020). Infection of dogs
with SARS-CoV-2. *Nature*, *586*(7831), 776-778. https://doi.org/10.1038/s41586-020-2334-5
Smith, S. L., Anderson, E. R., Cansado-Utrilla, C., Prince, T., Farrell, S., Brant, B., Smyth, S., Noble, P.-J.
M., Pinchbeck, G. L., Marshall, N., Roberts, L., Hughes, G. L., Radford, A. D., & Patterson, E. I.
(2021). SARS-CoV-2 neutralising antibodies in dogs and cats in the United Kingdom. *Current
Research in Virological Science*, *2*, 100011.
https://doi.org/https://doi.org/10.1016/j.crviro.2021.100011
Stevanovic, V., Tabain, I., Vilibic-Cavlek, T., Mauric Maljkovic, M., Benvin, I., Hruskar, Z., Kovac, S.,
Smit, I., Miletic, G., Hadina, S., Staresina, V., Radin, L., Plichta, V., Skrlin, B., Vrbanac, Z.,
Brkljacic, M., Cvetnic, M., Habus, J., Martinkovic, K., . . . Barbic, L. (2021). The Emergence of
SARS-CoV-2 within the Dog Population in Croatia: Host Factors and Clinical Outcome. *Viruses*,
13(8), 1430. https://www.mdpi.com/1999-4915/13/8/1430
Stevanovic, V., Vilibic-Cavlek, T., Tabain, I., Benvin, I., Kovac, S., Hruskar, Z., Mauric, M., Milasincic, L.,
Antolasic, L., Skrinjaric, A., Staresina, V., & Barbic, L. (2020). Seroprevalence of SARS-CoV-2
infection among pet animals in Croatia and potential public health impact. *Transbound
Emerg Dis*, *n/a*(n/a). https://doi.org/https://doi.org/10.1111/tbed.13924
The European Pet Food Industry. (2020). FACTS & FIGURES 2020 European Overview.
https://doi.org/https://fediaf.org/images/FEDIAF_Facts_and_Figures_2020.pdf
Udom, K., Jairak, W., Chamsai, E., Charoenkul, K., Boonyapisitsopa, S., Bunpapong, N.,
Techakriengkrai, N., & Amonsin, A. (2021). Serological survey of antibodies against
SARS-CoV-2 in dogs and cats, Thailand. *Transbound Emerg Dis*.
https://doi.org/10.1111/tbed.14208
Zhang, Q., Zhang, H., Gao, J., Huang, K., Yang, Y., Hui, X., He, X., Li, C., Gong, W., Zhang, Y., Zhao, Y.,
Peng, C., Gao, X., Chen, H., Zou, Z., Shi, Z. L., & Jin, M. (2020). A serological survey of SARS-
CoV-2 in cat in Wuhan. *Emerg Microbes Infect*, *9*(1), 2013-2019.
https://doi.org/10.1080/22221751.2020.1817796
Zhao, S., Schuurman, N., Li, W., Wang, C., Smit, L. A. M., Broens, E., Wagenaar, J., van Kuppeveld, F. J.
M., Bosch, B.-J., & Egberink, H. (2021). Serologic Screening of Severe Acute Respiratory
Syndrome Coronavirus 2 Infection in Cats and Dogs during First Coronavirus Disease Wave,
the Netherlands. *Emerging Infectious Disease journal*, *27*(5), 1362.
https://doi.org/10.3201/eid2705.204055

551