1 **Title:** plastiC: A pipeline for recovery and characterization of plastid genomes from
2 metagenomic datasets
3

4 **Authors**: Ellen S. Cameron[a,b,*], Mark L. Blaxter[b], Robert D. Finn[a]
5
6 [a] EMBL-EBI European Bioinformatics Institute, Wellcome Genome Campus, Hinxton,
7 Cambridge CB10 1SD, United Kingdom
8
9 [b] Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, CB10 1SA,
10 United Kingdom
11
12 * Corresponding Author:
13 E-mail: escameron@ebi.ac.uk (E.S. Cameron)
14

Abstract

Summary
Most metagenomic workflows have been developed for identification of prokaryotic and
eukaryotic community constituents, but tools for identification of plastid genomes are
lacking. *plastiC* is a workflow that allows users to identify plastid genomes in metagenome
assemblies, assess completeness, and predict taxonomic association from diverse
environmental sources.
22
23 *Availability and Implementation*
24 *plastiC* is a Snakemake workflow available at https://github.com/Finn-Lab/plastiC.
25
26 *Supplementary information*
27 Supplementary information available online.

1. Introduction

Culture-independent sequencing of environmental samples, often referred to as metagenomics, has revolutionised the field of microbial ecology, allowing taxonomic and functional characterization of microbial communities from diverse environmental sources. A variety of bioinformatic tools and pipelines are available for the generation and quality estimation of prokaryotic and eukaryotic metagenomic assembled genomes (MAGs) (Parks *et al.*, 2015; Saary *et al.*, 2020). Organellar genomes are essential to the functioning of eukaryotic organisms but are poorly assessed in current workflows. Plastid (or chloroplast) genomes are of particular interest because of their roles in primary production, yet the diversity of plastids present in metagenomic data is usually ignored and constrains the full understanding of community structure and function. Most existing MAG methods are rooted in single-copy marker gene (SCMG) approaches. However, SCMG approaches are not easily transferred to the genomes of plastid organelles due to reduced genome size, variable gene content (especially driven by organellar-nuclear gene transfer) and large repetitive regions. Thus, alternative methods for quality evaluation are required. Here, we present *plastiC*, a Snakemake (Mölder *et al.*, 2021) workflow for the identification and evaluation of plastids in metagenomic samples available at www.github.com/Finn-Lab/plastiC.

2. Workflow

plastiC is designed to analyse contigs generated from metagenomic assembly workflows (Figure 1). Contigs from these assemblies can be sorted into bins using intrinsic sequence and coverage features, and these bins processed to identify those that putatively correspond to prokaryotic and eukaryotic MAGs. These MAGs are typically processed to identify likely taxonomic affiliation derived from marker gene placement, and their quality (*i.e.* completeness and contamination) assessed. While *plastiC* can be run on raw assemblies, we recommend removing contigs from bins/MAGs determined as "high quality" before analysis. Putative plastid contigs are identified using *Tiara* (Karlicki *et al.*, 2022). The assembly is subsequently (re-)binned using *metaBAT2* (Kang *et al.*, 2019) with a reduced bin size of 50 kb to ensure that putative plastid bins are retained. The resulting bins subsequently are scanned for plastid contigs as identified by *Tiara*, and bins containing more than a user-specified threshold (default: 90%) of plastid sequences are retained as probable plastid bins.

Machine learning algorithms trained on KEGG module completeness were recently demonstrated as an effective approach for estimating metagenomic bin completeness (Chklovski *et al.*, 2022) and we adapted this approach in *plastiC*. *Prodigal* (Hyatt *et al.*, 2010) is used to predict genes in the probable plastid bins, and these predictions are compared to the UniRef100 database (Bateman *et al.*, 2021) using *diamond blastp* (Buchfink *et al.*, 2021) to obtain KEGG annotations (Kanehisa *et al.*, 2016). plastiC calculates KEGG module coverage to estimate completeness (see Supplementary Methods). Due to the presence of shared metabolic pathways and genes between plastid and bacterial chromosomes, bacterial bins may score as highly complete, and this finding supports the requirement to filter assemblies to remove bacterial MAGs before running *plastiC*.

Importantly, *plastiC* also identifies the potential eukaryotic source of the plastid. *plastiC* initially performs a taxonomic classification on the plastid contigs using *CAT* (von Meijenfeldt *et al.*, 2019) and additionally searches for taxonomic marker genes using *hmmer* (for *rbcL*) (Eddy, 2022) and *barrnap* (for rRNA loci) (Seeman, 2022). These loci can be used for further taxonomic analyses. As outputs, *plastiC* produces report files summarising bin

74 span, estimated completeness, and taxonomic allocation for each putative plastid bin, as
75 well as FASTA files of the marker loci identified.

76
Figure 1: Data preparation and overall workflow for *plastic.* Users provide metagenomic
assemblies from which putative plastid bins are identified. A summary report and FASTA
files of identified marker loci are generated for each sample.

3. Discussion
Currently, metagenomic characterization of plastid genomes is limited to the
identification of plastid contigs using *Tiara*. However, only identifying plastid contigs prevent
full disentanglement of metagenomic datasets, including the potential presence of multiple
plastid genomes arising from different sources. *plastiC* uniquely provides users with the
opportunity to characterize plastid genomes including separation into independent plastid
genomes and source classification to inform on present eukaryotic photosynthetic
organisms. Notably, this is the first tool to provide a completeness estimate for plastid
genomes from metagenomic sources. The high copy number and small genome size of
plastids may permit easier recovery and serve as a proxy for the identification of the
presence of eukaryotes which may not have sufficient genomic signal to be recovered as a
high-quality eukaryotic MAG, as well as the potential to expand reference databases with
metagenomic derived plastids. *plastiC* will support characterization of plastids in diverse
ecological contexts including, for example, symbiotic interactions (e.g., lichens; see
Supplementary Case Study), complex marine ecosystems and characterisation of
components of herbivore diets.

4. References

Bateman,A. *et al.* (2021) UniProt: the universal protein knowledgebase in 2021. *Nucleic*
Acids Res, **49**, D480–D489.

Buchfink,B. *et al.* (2021) Sensitive protein alignments at tree-of-life scale using DIAMOND.
Nat Methods, **18**, 366–368.

Chklovski,A. *et al.* (2022) CheckM2: a rapid, scalable and accurate tool for assessing
microbial genome quality using 1 machine learning 2 3 4. *bioRxiv*.

Eddy, S. (2022) HMMER: Biosequence analysis using profile hidden Markov models.
http://hmmer.org

Hyatt,D. *et al.* (2010) Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics*.

Kanehisa,M. *et al.* (2016) KEGG as a reference resource for gene and protein annotation.
Nucleic Acids Res, **44**, D457–D462.

Kang,D.D. *et al.* (2019) MetaBAT 2: An adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. *PeerJ*, **2019**.

Karlicki,M. *et al.* (2022) Tiara: deep learning-based classification system for eukaryotic
sequences. *Bioinformatics*, **38**, 344–350.

von Meijenfeldt,F.A.B. *et al.* (2019) Robust taxonomic classification of uncharted microbial
sequences and bins with CAT and BAT. *Genome Biol*, **20**.

Mölder,F. *et al.* (2021) Sustainable data analysis with Snakemake. *F1000Res*, **10**, 33.

Parks,D.H. *et al.* (2015) CheckM: Assessing the quality of microbial genomes recovered from
isolates, single cells, and metagenomes. *Genome Res*, **25**, 1043–1055.

Saary,P. *et al.* (2020) Estimating the quality of eukaryotic genomes recovered from
metagenomic analysis with EukCC. *Genome Biol*, **21**.

Seeman, T. (2022). barrnap 0.9: rapid ribosomal RNA prediction.
https://github.com/tseemann/barrnap