1

Variables associated with owner perceptions of the health of their dog: further analysis of data from a large international survey

Richard Barrett-Jolley, Alexander J. German[*]

Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool, UK

Running Title: Variables associated with dog health

* Corresponding author

E-mail: ajgerman@liverpool.ac.uk

Abstract

In a recent study (doi: 10.1371/journal.pone.0265662), associations were identified between owner-reported dog health status and diet, whereby those fed a vegan diet were perceived to be healthier. However, the study was limited because it did not consider possible confounding from variables not included in the analysis. The aim of the current study was to extend these earlier findings, using different modelling techniques and including multiple variables, to identify the most important predictors of owner perceptions of dog health.

From the original dataset, two binary outcome variables were created: the *'any health problem'* distinguished dogs that owners perceived to be healthy ("no") from those perceived to have illness of any severity; the *'significant illness'* variable distinguished dogs that owners perceived to be either healthy or having mild illness ("no") from those perceived to have significant or serious illness ("yes"). Associations between these health outcomes and both owner-animal metadata and healthcare variables were assessed using logistic regression and machine learning predictive modelling using XGBoost.

For the *any health problem* outcome, best-fit models for both logistic regression (area under curve [AUC] 0.842) and XGBoost (AUC 0.836) contained the variables dog age, veterinary visits and received medication, whilst owner age and breed size category also featured. For the *significant illness* outcome, received medication, veterinary visits, dog age and were again the most important predictors for both logistic regression (AUC 0.903) and XGBoost (AUC 0.887), whilst breed size category, education and owner age also featured in the latter. Any contribution from the dog vegan diet variable was negligible.

The results of the current study extend the previous research using the same dataset and suggest that diet has limited impact on owner-perceived dog health status; instead, dog age, frequency of veterinary visits and receiving medication are most important.

Introduction

There is an increasing interest from owners in feeding unconventional diets, such as those utilising uncooked (so-called 'raw' diets) or plant-based (either vegetarian or vegan diets) ingredients. In a recent UK survey, 7% and approaching 1% of pet owners reportedly fed their dog either a raw or vegan diet, respectively [1]. Veganism is an increasingly popular food choice in humans, with recent surveys indicating 2-3% [2], 4% [3] and 3% [4] of people in the UK, Europe and USA, respectively, to be vegan. Although still uncommon (<1% of owners), use of plant-based (vegan or vegetarian) diets for dogs is increasing in popularity [1], not least amongst owners who are vegan themselves [5]. Interestingly, the popularity of feeding raw diets is also increasing, with the same survey suggesting that 7% of the UK dog-owning population use this method [1], a proportion similar to that reported in recent international study, where 9% of dog owners exclusively fed raw food [6]. The main reason that owners cite for considering a plant-based diet is concern about animal welfare [5], whilst owners who feed raw diets believe these to be more 'natural' and healthier than commercial food despite an absence of evidence to support this [7]. Despite their increasing popularity, concerns remain regarding the safety of feeding unconventional foods; for both vegan [5,8] and raw [9-11] food, in terms of nutritional adequacy. For raw food, there are additional concerns about contamination, both with pathogenic bacteria and bacteria resistant to multiple antimicrobials [11-13].

In recent studies, the safety of feeding unconventional diets for dogs has been examined, albeit using owner reports of health from questionnaires or surveys. In one study of 16,475 households that fed raw food, only 39 (0.2%) recalled a member of the household acquiring a pathogenic infection during the time they were using the food [14], perhaps, suggesting that risks to owners from bacterial contamination of raw food are uncommon. The health of dogs being fed unconventional diets has also been assessed. In one survey of 1,413 owners

conducted across North America, those feeding plant-based diets reported fewer health

conditions in their dogs and a longer lifespan, compared with owners of dogs fed a meat-

based diet [6]. In a second study [15], owner-reported health of dogs being fed various diet

types was assessed. Compared with dogs fed a conventional diet, owners that fed either raw

food or a vegan diet reported better health. However, besides being reliant on owner-

reported health, rather than more objective health measures, no account was taken of possible

confounding from other variables. To the authors' credit, the original study data were made

available for use by other researchers (https://rebrand.ly/2020/pet-food-consumers-study),

which not only included the diet and health data they analysed, but also other metadata

including owner (e.g., country, career, education, age and gender) and animal (e.g., age, sex,

neuter status and breed) variables, and healthcare-related variables (e.g., veterinary visits,

receiving medication, use of therapeutic diet). Therefore, the aim of the current study was to

extend the earlier findings using different statistical techniques to create models that best

predicted owner perceptions of health, and to identify the relative importance of the variables

that contributed to the final model. This included the use of binary logistic regression and

machine learning predictive modelling using a scalable tree boosting system (XGBoost [16]).

Materials and Methods

Study design

A secondary analysis was conducted using data from a recent international survey of owner opinions on the health of their dog [15]. The open-access dataset from this study is accessible at https://rebrand.ly/2020/pet-food-consumers-study. The reason for conducting this further analysis was that many variables gathered were not analysed statistically, and that the effect of possible confounding was not adequately investigated, both of which were limitations acknowledged by the study authors [15]. Further, we were particularly interested in exploring associations between a wider range of variables and owner perceptions of the health of their dog.

Dataset

Full details of survey design, methodology, information collected and category definitions have been reported previously [15]. Briefly, the original survey was designed using an online survey platform (https://www.onlinesurveys.ac.uk), with owners being asked to record details about their dog (including demographic data, main diet type and health characteristics) and themselves. Owner information used for the current study included location (country or continental region), setting (urban, rural or a mix of urban and rural), education level (qualifications achieved), household income, whether their occupation was in an animal-related career (including veterinarian, veterinary technician or nurse, animal breeder, animal trainer or pet industry worker), age, gender and diet. Animal information used for the current study included: age, sex, neuter status, breed size category (e.g., toy, small, medium, large and giant) and main diet type (including conventional, raw, vegetarian, vegan, *in vitro* meat, insect, fungal or algal). We also used responses from owners about the

healthcare their dog received including number of veterinary visits in the last 12 months,

whether the dog had received medication (besides those used in preventive care e.g. routine

vaccinations and endo- or ecto-parasite treatments) in the last 12 months, and whether the

dog was on a therapeutic diet; when dogs were fed a therapeutic diet, owners were asked to

base the information on their dog's diet on their previous diet [15]. Finally, in the original

survey, owners were also asked their opinion about the health of their dog in the last 12

months, with possible responses including: 'healthy', 'generally healthy with minor or

infrequent problems', 'significant or frequent problems', 'seriously ill' or 'unsure'. Since we

were primarily interested in owner perceptions of health, we used this information to create

our outcome variables, as described below.

128 Statistical methods

129 Statistical software

All data pre-processing and statistical analyses were conducted using an online open-

access statistical language and environment (R, version 4+) [17], with several packages

including: 'aod' version 1.3.2 [18], 'broom' version 0.8.0 [19], 'car' version 3.0.13 [20],

'caret' version 6.0.93 [21], 'cluster' version 2.1.4 [22], 'clusterSim' version 0.51.3 [23],

'corrplot' version 0.92 [24], 'data.table' version 1.14.2 [25], 'dbscan' version 1.1-11 [26],

'DescTools' version 0.99.45 [27], 'dplyr' version 1.0.9 [28], 'factoextra' version 1.0.7 [29],

'fmsb' version 0.7.3 [30], 'foreign' version 8.82 [31], 'ggforce' version 0.4.1 [32], 'ggplot2'

version 3.3.6 [33], 'ggthemes' version 4.2.4 [34], 'glmtoolbox' version 0.1.3 [35],

'heatmaply' version 1.4.0 [36], 'Hmisc' version 4.7-0 [37], 'IHW' version 1.26.0 [38].

'mltools' version 0.3.5 [39], 'pROC' version 1.18.0 [40], 'RColorBrewer' version 1.1-3 [41],

'readxl' version 1.4.0 [42], 'reshape2' version 0.8.9 [43], 'ROCR' version 1.0-11 [44],

'rsample' version 1.1.0 [45], 'rstatix' version 0.7.0 [46], 'showtext' version 0.9-5 [47];

'smotefamily' version 1.3.1 [48], 'stringr' version 1.4.0 [49], 'tidyverse' version 1.3.1 [50],

'uwot' version 0.1.16 [51] and xgboost version 1.6.0.1 [16]. Statistical reports of all data pre-

processing and analysis are included in the supplementary information (S1-S9 Files), whilst

all code is available online (https://github.com/RichardBJ/CanineHealth23).

Initial dataset pre-processing

Initially, data were imported into the statistical environment using the 'readxl' package

[42], and then manipulated using the 'dplyr' package [28]. Columns were renamed for

clarity and to improve compatibility with use in R (see below). Counts of all categorical data

were examined, and categories with small groups were either removed or combined with

others to exclude missing data or to reduce dataset imbalance, as described below. The

variables included in the dataset were owner variables and animal variables (collectively

owner-animal metadata), as well as healthcare variables and health outcomes variables.

Data pre-processing occurred in two stages: in the first stage (prior to Uniform

Manifold Approximation and Projection; UMAP; see below), the original dog sex variable

(male sexually intact, male castrated, female sexually intact, female spayed) was separated

into two binary variables, *dog sex* (male, female) and *neuter status* (yes, no). In addition,

data were excluded from dogs where age was reported as 'unsure' (n=3) or were aged <1

year (n=26), to ensure a focus on adult dogs.

To facilitate the modelling studies (logistic regression and XGBoost), further data pre-

processing steps were undertaken after UMAP. For owner variables, pre-processing included

combining categories with small numbers in the *location* variable (e.g., 'Africa' [n=7], 'Asia'

[n=25], 'other' [n=11] and 'South America' [n=47]) into a new 'other' category, and also

removing the 'other' category (n=15) from the *settings* variable. The *owner education*

variable was simplified by combining the 'basic' (n=15) and 'high school' (n=478) categories

and combining the 'PG' (n=515) and 'PhD' (n=90) categories. A binary *animal career*

variable was created from the *career* variable by combining the 'veterinarian' (n=125),

'veterinary technician or nurse' (n=54), 'animal breeder' (n=17), 'animal trainer' (n=140)

and 'pet industry worker' (n=205) categories into a 'yes' category, whilst the remaining

option ('none of the above' [n=2,137]) formed the 'no' category. The *owner income* variable

was simplified by removing the 'prefer not to answer' category (n=184) and ordered, whilst

the *owner age* variable was simplified (by combining the '18-19y' [n=19] and '20-29y'

[n=391] categories into a new '<30y' category and combining the '60-69y' [n=364] and

'>70y' [n=85] categories into a new '>60y' category) and then ordered. Given small

numbers, the 'prefer not to answer' (n=17) and 'other '(n=4) categories were removed from

owner gender variable, and the 'other' category (n=18) was removed from the *owner diet*

variable. Further, binary variables were created for owners who reported consuming a vegan

diet (yes vs. no) and owners who reported consuming either a vegan or vegetarian diet (yes vs.

no).

For animal variables, in addition to having a continuous *dog age* variable, an ordinal

variable was also created with data grouped into quintiles (1-3y, 3-5y, 5-7y, 7-9y, 9-20y).

For the *dog diet* variable, the 'unsure' (n=8), 'insect-based' (n=5), 'meat-based (lab-grown)'

(n=6) and 'mixture' (n=13) categories were all removed since numbers were again small.

Further, binary variables were created for dogs consuming a *vegan* (yes vs. no), *vegan or*

vegetarian (yes or no) or a *raw* (yes vs. no) diet, and for dogs in the 'giant breed' category

(yes vs. no) of the *breed size category* variable.

For healthcare variables, the 'unsure' category (n=13) was removed from the *veterinary*

visits variable, and two ordinal variables were created; the first comprised 5 categories ('0

visits', '1 visit', '2 visits', '3 visits', '4+ visits'), whilst the second comprised 4 categories ('0

visits', '1 visit', '2 visits', '3+ visits' [n.b., '3 visits' and '4+ visits' categories combined]).

For our outcome variables (owner's perception of the health of their dog), we first

removed data from owners who reported that they were 'unsure' (n=5). From this, two

binary health status variables were created: the first was an *any health problem* variable,

where dogs classified as 'healthy' were classified as 'no', and all remaining categories

('generally healthy with minor or infrequent problems', 'significant or frequent problems'

and 'seriously ill') were classified as 'yes'; the second was a *significant illness* variable,

whereby dogs classified either as 'healthy' or 'generally healthy with minor or infrequent

problems' were classified as 'no', and those classified as having 'significant or frequent

problems' or being 'seriously ill' were classified as 'yes'.

As a final pre-processing step, a *decision-maker* (primary vs. other) variable was

created from data about the role owners played choosing a diet for their dog; for this, the

'primary decision-maker' category was classified as 'primary', whilst the other two

categories ('play no role', n=15; 'play some lesser role', n=96) were classified as 'other'.

Given concerns about the reliability of information from owners who were not primary

decision-makers and healthcare received, its effect on the outcome variables used in

modelling studies (both binary illness variables) was first assessed by logistic regression,

before the other modelling studies, as described below.

Owner-dog metadata visualisation with uniform manifold approximation and projection with density-based spatial clustering

Unsupervised uniform manifold approximation and projection (UMAP) [52] was used

to reduce the multivariate owner and dog metadata data to two dimensions for visualisation

purposes (using the 'uwot' package [51]). After this, density-based spatial clustering

(DBSCAN, using the 'dbscan' package [26]) was applied to the UMAP outputs to identify

potential groupings within these owner-animal metadata, prior to the data pre-processing

necessary for the statistical and machine learning modelling, as explained above. The UMAP

approach was specifically developed to be applicable to mixed datasets with categorical,

ordinal and numeric data [52]. It is a potent, non-linear dimension-reduction technique that

outperforms principal component analysis (PCA) and t-distributed stochastic neighbour

embedding (TSNE) in its ability to preserve the global and local structure of data [52]. It

employs a specific type of manifold projection known as Riemannian geometry, which has

the strength of enabling meaningful clusters to be identified and visualised in a reduced

dimensional space. This approach differs from PCA in that it typically reduces multivariate

data to two parameters (arbitrarily called x and y); further, rather than calculating variability,

UMAP instead non-linearly transforms the data to create a lower-dimensional representation,

allowing more complex data patterns to be captured, compared with linear methods like PCA.

The axes in UMAP do not represent 'principal components' and cannot be interpreted in the

same way; instead, they help visualise high-dimensional data in a lower-dimensional space.

Each record (individual row of the dataset with one owner-dog combination)

contributes one point to the UMAP graph. The following animal variables were included:

dog age, *dog sex*, *neuter status*, *diet* and *breed size category*. In addition, the following

owner variables were included: *owner age*, *owner gender*, *owner diet*, *location*, *setting*,

education, *income* and *decision-maker status*. Following best practise [53], different distance

metrics were used including Euclidian (for continuous data e.g., *dog age*), Manhattan (for

ordered categorical data e.g., *owner age*, *education*) and Hamming (for non-ordered

categorical data e.g. *dog diet*, *dog sex* etc)

The quality of DBSCAN clustering was assessed with a silhouette plot along with

silhouette scores [53] and also by calculating Davies-Bouldin's cluster separation measure

[54]. The silhouette plot provides a visual representation of how well points within the

UMAP have been classified into their respective clusters, whilst average silhouette widths

(silhouette scores) quantify how similar the point is to its own cluster relative to other

clusters; silhouette scores can range from +1 to -1, with positive and negative values

indicating good and poor matching, respectively; further, average cluster silhouette scores of

0.26-0.50, 0.51-0.70 and 0.71-1.00 represent weak, reasonable and strong clustering,

respectively [55]. Davies-Bouldin's cluster separation measure is determined from the ratio

between the scatter within and the separation between clusters, with smaller values

suggesting better clustering and values <1 representing very good clustering.

250 Visualising associations amongst owner-animal metadata and healthcare
251 variables

After the second data pre-processing stage, Kendall's *tau* [56] was used to conduct

exploratory correlation analysis using the R package 'rstatix' [46], and a correlation matrix

[57] was plotted with the 'corrplot' package [24]. Data were first coerced into a numerical

ordinal form, except for the *location* variable because there was no ordinal relationship

amongst categories; instead, this variable was one-hot encoded, whereby a new binary

variable was added for each category. *P* values are expressed as their negative logarithm (-

Log*P*), whilst false discovery rate (FDR) was calculated by the Benjamini-Hochberg

procedure using the 'IHW' package [38]. Hierarchical clustering was added with R

stats::hclust using the Ward method [58]. Strengths of association were reported according to

Cohen [59], whereby correlation coefficients of 0.1-0.3, 0.3-0.5 and 0.5-1.0 represented small,

medium and large correlation, respectively.

264 Binary logistic regression

Binary logistic regression models were created using the 'glm' function in R. Both

binary illness variables (*any health problem* and *significant illness*) were used separately as

outcome variables, whilst all owner, animal and healthcare variables were tested as predictor

variables. We used a combination of simple and multiple logistic regression. Simple logistic

regression analyses were initially performed using data from all owners (irrespective of

decision-maker status), with results shown in the supplementary information (S1 and S2

Tables). However, these models revealed concerns over the reliability of data from owners

who were not primary decision-makers (see below) and, as a result, all remaining logistic

regression analyses were instead conducted with the primary decision-maker dataset. For

these analyses, the dataset was first randomly divided into training (75%) and test (25%)

datasets, using the 'sample' function in R, a step that was necessary to facilitate model

validation by receiver operating characteristic (ROC) curves (see below).

Simple logistic regression models created with the primary decision-maker dataset

enabled unadjusted associations between each outcome variable and single predictor

variables to be determined. Some variables had been coded in different ways, as described

above; for example, *dog age* was coded as both a continuous variable and an ordinal variable

with 5 categories, with the continuous *dog age* variable also being tested with and without

basis (b-)splines; further, veterinary visits were coded as both a 4- or 5-category variable,

whilst *dog diet* was coded both as a 4-category variable [conventional, raw, vegetarian and

vegan] or as separate binary variables for *raw diet*, *vegan diet* or a combined *vegan-

vegetarian* binary variable [see above]. To determine the coding approach that was most

appropriate for such variables, performance of competing models was assessed using the

Bayesian Information Criterion (BIC, see below) [60]; the coding approach that fitted the

data best was selected for use in the initial multiple regression modelling as long as model

assumptions were met (e.g., independence of errors, the requirement for a linear relationship

between the logit of the outcome and continuous predictor variables). For example, the

vegan diet variable performed better than the *vegan-vegetarian diet* variable, the continuous

dog age variable performed better than the ordinal *dog age* variable, albeit that b-splines

(utilising boundary knots and an internal knot at the median value [6 years]) were required to

ensure that model assumptions were met. Further, for the *significant illness* outcome variable,

the *giant breed* predictor variable performed better than the *breed size category* variable, and

there were also some issues with model convergence for the latter. Models containing two

related predictor variables, with and without interaction terms, were also tested when these

were clinically relevant (e.g., between dog sex and neuter status, between owner and dog diet

categories and between location and setting).

In the first step of the multiple logistic regression modelling, two multiple regression

models were created for each outcome variable (*any health problem* and *significant illness*);

all predictor variables were included in the first of these models (*all-variable* model), whilst

only owner and animal variables were included in the second model (*owner-animal metadata*

model). The performance of *all-variable* and *owner-animal metadata* models were then

compared with each other and with the *best-fit* models that were created subsequently. For

these *best-fit* models, we used a supervised, stepwise approach starting with all variables.

This initial model was manually refined in both a backwards and forwards stepwise fashion,

with the BIC being used to select the model within the same family with the best

generalisability (a measure of its goodness of fit compared with its complexity) [60]. Using

this approach, variables could be added or removed until the model with the smallest BIC

was found, according to previously published rules [61], whereby model generalisability was

deemed to be superior if the BIC of the new model was less than the previous model by at

least 2 units.

After selecting a *best-fit* model and, given the importance of diet in the original work

[15], both the *dog diet vegan* and *owner diet vegan* variables were separately added back, to

determine their effect on model performance.

Results are reported as estimates of the regression coefficients (β) with the associated

standard error (SE), and with odds ratios (OR) with the associated 99% confidence intervals

(99%-CI). Influential datapoints were identified and assessed using Cook's distance and, in

most models, none were identified. In the occasional cases where outliers were identified, we

checked them in the original data to make sure there were no obvious errors (e.g., unrealistic

results such as dog age >20 years); no such errors were found. Given that we were using data

from a secondary source, had no further means of verifying them (e.g., by cross-checking

against the original questionnaire response), we decided not to remove them as there was no

valid justification to do so (e.g., a typographical error created when the original data were

entered). Possible multicollinearity in all models was assessed using the generalised variance

inflation factor (GVIF) and GVIF($1/(2 \times Df)$); these were deemed to be acceptable when all

values were <4 and <2 for GVIF and GVIF($1/(2 \times Df)$), respectively [62]. If necessary,

multicollinearity was resolved by removing the variable with the greatest GVIF. Goodness

of fit was tested by a visual inspection of observed and expected results and the Hosmer and

Lemeshow test for large datasets. In addition, the proportion of variance explained in the

overall model was assessed by the coefficient of determination (pseudo-R^2) based on the

method reported by Nagelkerke [63]. Further, the log-odds of the outcome variable and dog

age was plotted to enable visual assessment of whether the relationship was linear, and this

was further assessed with the Box-Tidwell test. Finally, prediction accuracy was assessed by

creating ROC curves, using the ROCR package [44], and then calculating their area under the

curve (AUC). Values could range from 0 to 1; a model that performed no better than chance

would have an AUC of 0.5, and models predicting better than by chance would have AUC

>0.5, with an AUC of 1.0 suggesting perfect prediction. The test dataset was used to

generate the ROC curves and AUC for models using data from owners who were primary

decision-makers; however, because the *all-owner* dataset was not subdivided, AUC are based

in ROC curves from the original dataset.

Machine learning predictive modelling

A scalable tree boosting system (XGBoost; "eXtreme Gradient Boosted" tree) was

implemented in the statistical environment [16]. Data were pre-processed as described above

and, to facilitate XGBoost, ranked variables used for the Kendall's tau correlation (as

described above) were coerced to a numeric datatype, using one-hot encoding where

necessary, and then subdivided 70:30 into training and test sets. A model was then built on

the training data without any involvement of the test set. After pre-processing but before

training, training data were further divided into training and validation subsets (70:30), and

the training subset augmented as per standard practice for machine learning [64]. We trialled

two augmentation methods: SMOTE [48] and a similar method where underrepresented

classes were duplicated, but with simple addition of signal noise. Final models used the latter

given that it performed better than SMOTE. Test data were left untouched. The model was

then trained using the augmented training data with area-under the curve (AUC) as the

evaluation metric, against the validation training subset. Finally, the model was built and

saved to file. Receiver operating characteristic curves were calculated and plotted with the

saved model against the previously unseen test dataset using the packages 'pROC' [40] and

'ROCR' [44]. Prediction accuracy was estimated by calculating the ROC AUC in a similar

manner to that described above for logistic regression. Different measures of the contribution

of the variables ('features') included in the final models were also calculated and depicted

graphically. These included 'feature importance' (a metric of the fractional contribution of

each feature to the model, based on the total gain from including each feature), 'cover' (a

metric of the number of observations related to this feature in the model) and 'frequency' (a

metric that represents the relative number of times a feature has been used in trees). Since

these metrics are normalised, the sum of scores for all features in the model is 1.0.

Results

Owner-dog metadata visualised by Uniform Manifold Approximation and Projection

The original dataset comprised information on 2,639 owner-dog pairs [15]. After the initial pre-processing stage to create separate dog sex and neuter status variables, and also removing dogs without age data (n=3) or that were <1 year of age (n=27), the number of owner-dog pairs with available data was 2,609. We first used a UMAP projection and density-based clustering to create a simple overall visualisation of relationships in these owner-animal metadata (Fig 1; S1 and S2 Figs). Six distinct clusters were identified, with a Davies-Bouldin's cluster separation measure of 0.652 (good). Good clustering was evident within 3 clusters (average silhouette scores: cluster 2, 0.86; cluster 5, 0.84; cluster 6, 0.80), there was reasonable clustering in two others (average silhouette scores; cluster 3, 0.62; cluster 4, 0.59), but clustering was poor in the final cluster (cluster 1, average silhouette score 0.36).

Examining the owner-animal metadata, cluster 2 was most distinctive, almost-exclusively comprising dogs fed either a vegan (328/374 [87.7%] points in the cluster, 97.6% of all vegan dogs) or vegetarian (31/374 [8.3%] points in the cluster, 88.6% of all vegetarian dogs) diet, and also containing a predominance of owners consuming a vegan diet (332/374 [88.8%] points in the cluster, 57.1% of all vegan owners). In contrast, cluster 3 comprised owners who consumed either a vegan (249/608, 41.0%), vegetarian (240/608, 39.5%) or pescatarian (119/608, 19.6%) diet, but fed their dog either a conventional meat-based (423/608, 69.6%) or raw (175/608, 28.8%) diet. Cluster 6 was also distinctive in that it mostly comprised owners who consumed an omnivore (214/316, 67.7%) or omnivore reducing (98/316, 31.0%) diet and fed their dogs a raw diet (299/316, 94.6%). In the

remaining 3 clusters (clusters 1, 4 and 5), most owners consumed either an omnivore or

omnivore-reducing diet and usually fed their dog either a conventional meat-based diet or

raw diet. Besides owner and dog diet, there were no major differences amongst clusters for

the other variables, although some variables did appear to separate some clusters (e.g., cluster

4 for the setting variable; cluster 5 for the location variable; S1 and S2 Figs).

Fig 1. Visualisation of owner metadata by Uniform Manifold Approximation and Projection

with Density-Based Spatial Clustering. Metadata for 2,609 owner-dog pairs were pre-processed

with all factors as numeric and subject to dimension reduction with the UMAP projection technique.

Owner variables included in this visualisation were *diet*, *sex*, *location*, *education* and *income*; animal

variables included were *age*, *sex*, *neuter status*, *diet* and *breed size category*. Healthcare variables

were not included. Each individual row of the data (owner-dog combination) contributes one UMAP

x and y coordinate. Points colour-coded by *dog diet* (a) or *owner diet* (b), where 5 distinct clusters are

evident. (c) Silhouette plot depicting how well clusters were separated within the UMAP; silhouette

values measure how similar the point is to its own cluster relative to other clusters, with positive and

negative values indicating good and poor matching, respectively. The numbers in circles are the

average silhouette score for each cluster, whereby values of 0.26-0.50, 0.51-0.70 and 0.71-1.00

representing weak, reasonable and strong clustering, respectively.

Summary of owner-animal metadata

After the further data pre-processing stage, as described above, the number of owner-

dog pairs with available data was 2,322, and details are shown in Table 1. This population

was similar to the data used in the previous study [15]. Briefly, owners represented a broad

age range with a relatively even distribution in education level, from high school to

postgraduate. However, most (2,146, 92.4%) respondents were female, whilst most

responses were from the United Kingdom (1,659, 71.4%), with other European countries the

next most common region represented. Almost a quarter of the respondents (532, 22.9%)

defined themselves as vegan with a further 235 (10.1%) reporting to be vegetarian.

Regarding the dogs in this dataset (Table 1), a range of ages and breed sizes were included

with the majority of male (294/1169, 74.9%) and female (192/1042, 81.6%) dogs being

neutered. Approximately a third (744, 32.0%) of dogs were fed a raw meat diet, whilst 313

(13.5%) and were 35 (1.5%) fed vegan or vegetarian diets, respectively. Finally, most

owners (2,211; 95.2%) reported that they were the primary decision-makers for their dog's

diet (Table 1).

Table 1. Comparison of summary information from the original dataset of 2,322 owner-dog pairs, and the dataset used for statistical analyses comprising 2,211 owner-dog pairs where the owner was the primary decision-maker.

Variable [1]	Categories	All owners		Primary decision-makers	
		Number	Percentage	Number	Percentage
Owner variables					
Location	United Kingdom	1659	71.4%	1609	72.8%
	Other European country	358	15.4%	318	14.4%
	North America	128	5.5%	119	5.4%
	Australia / New Zealand / Oceania	100	3.4%	97	4.4%
	Other region	77	3.3%	68	3.1%
Setting	Urban	824	35.5%	780	35.3%
	Equally urban and rural	740	31.9%	703	31.8%
	Rural	758	32.6%	728	32.9%
Owner age (years)	<30	385	16.6%	340	15.4%
	30-39	507	21.8%	482	21.8%
	40-49	474	20.4%	452	20.4%
	50-59	528	22.7%	515	23.3%
	≥60	428	18.4%	422	19.1%
Owner gender	Female	2146	92.4%%	2055	92.9%
	Male	176	7.6%	156	7.1%
Education	Basic or high school	439	18.9%	414	18.7%
	College	662	28.5%	636	28.8%
	Graduate	682	29.4%	645	29.2%
	Postgraduate	539	23.2%	516	23.3%

Income	Low	364	15.7%	352	15.9%
	Medium	1611	69.4%	1530	69.2%
	High	347	14.9%	329	14.9%
Animal-related Career	Yes	435	18.9%	425	19.2%
	No	1887	81.3%	1786	80.8%
Owner diet [2]	Omnivore	931	40.1%	901	40.7%
	Omnivore (restricted)	500	21.5%	479	21.7%
	Pescatarian	124	5.3%	117	5.3%
	Vegan	532	22.9%	493	22.3%
	Vegetarian	235	10.1%	221	10.0%
Decision-maker status[3]	Primary	2211	95.2%	2211	100.0%
	Some lesser role	96	4.1%	---	---
	No role	15	0.06%	---	---
Animal Variables					
Age (years)	1 to 5	1144	49.3%	1089	49.3%
	6 to 20	1178	50.7%	1122	50.7%
Breed size category					
	Medium	896	38.6%	859	38.9%
	Toy	56	2.4%	52	2.4%
	Small	467	20.1%	439	19.8%
	Large	803	34.6%	764	34.5%
	Giant	100	4.3%	97	4.4%
Sex	Female	1095	47.2%	1042	47.1%
	Male	1227	52.8%	1169	52.9%
Neuter status	Sexually intact	512	22.0%	486	22.0%

	Neutered	1810	78.0%	1725	78.0%
Dog diet [2]	Conventional	1230	53.0%	1152	52.1%
	Raw	744	32.0%	732	33.1%
	Vegan	313	13.5%	295	13.3%
	Vegetarian	35	1.5%	32	1.5%
Healthcare variables					
Veterinary visits	None	408	17.6%	399	18.0%
	1	907	39.1%	873	39.5%
	2	506	21.8%	473	21.4%
	3	200	8.6%	185	8.4%
	4 or more	301	13.0%	281	12.7%
Received medication	No	1386	59.7%	1331	60.2%
	Yes	936	40.3%	880	39.8%
Switched to a therapeutic diet	No	2211	95.2%	2111	95.5%
	Yes	111	4.8%	100	4.5%

[1] Definitions of the different categories are given in the original study [15]. [2] Owners were asked to choose a single diet classification based on
the main diet ingredients and whether these were conventional, raw, vegetarian or vegan in origin [15]; vegetarian diets could not include meat
but could include eggs or milk, whilst vegan diets could not include any [15]. [3] Owners were asked "When choosing your animal's diet, are you
the: primary decision-maker, play some lesser role or play no role" [15].

We were concerned with the reliability of data from the 111 owners (4.8%) who were

not the primary decision-makers, because their knowledge about the health of their dog could

well be less reliable and their beliefs and behaviours might also be different, all of which

would likely impact perceptions about the health of their dog. To examine this, we created

separate binary logistic regression models for the *any health problem* and *significant illness*

outcome variables, with *decision-maker status* as the sole predictor variable. There was no

difference in the odds of an owner recording their dog as having a *serious illness*, whether

they were a primary decision-maker or not (OR 0.662: 99%-CI 0.286, 1.896; *P*=0.252,

pseudo-R^2 0.0015). However, the odds of primary decision-makers recording their dog as

having *any health problem* were less than for owners who were not primary decision-makers

(OR 0.576: 99%-CI 0.347, 0.958; *P*=0.005, pseudo-R^2 0.0046). As a result of this difference,

we decided to focus further analyses on data from the 2,211 owners who were primary

decision-makers. However, to determine the effect of this decision, simple logistic regression

was also performed on the data from all owners (irrespective of decision-maker status; S1

Table; S2 Table), enabling comparisons to be made with simple logistic regression performed

on the data from primary decision-makers only (see below).

Summary of healthcare variables

Details of healthcare variables are also given in Table 1. Most dogs had visited their

veterinarian at least once in the last year (all owner dataset 1,914 [82.4%]; primary decision-

maker dataset 1,812 [81.9%]), whilst smaller proportions of dogs had received medication

(all owner dataset 936 [40.3%]; primary decision-maker dataset 880 [39.8%]) and or had

been switched to a therapeutic diet (all owner dataset 111 [4.8%]; primary decision-maker

dataset 100 [4.5%]).

Associations amongst owner-animal metadata and healthcare variables

Results of Kendall's tau correlations are shown in Fig 2, whilst datasets of all correlation coefficients and FDR-corrected -Log$P$ are presented in the supplementary information (S3 and S4 Tables, respectively). Several significant associations were identified including a large positive association between owners and dogs consuming *vegan diets* (Kendall's tau 0.66, -Log$P$ 212.0 [P<0.001]), a small positive association between UK location and dogs being fed a *raw diet* (Kendall's tau 0.24, -Log$P$ 29.5 0 [P<0.001]), as well as positive associations amongst healthcare variables (*received medication* and *veterinary visits*, Kendall's tau 0.51, -Log$P$ 154.8 0 [P<0.001]; *received medication* and *switched to therapeutic diet*, Kendall's tau 0.17, -Log$P$ 14.9 0 [P<0.001]; *switched to a therapeutic diet* and *veterinary visits*, Kendall's tau 0.16, -Log$P$ 15.3 0 [P<0.001]). In contrast, weak associations were observed between the *dog vegan diet* variable and the healthcare variables (*dog vegan diet* and *veterinary visits*, Kendall's tau -0.02, -Log$P$ 0.5 [P=0.340]; *dog vegan diet* and *received medication*, Kendall's tau -0.11, -Log$P$ 2.7 0 [P=0.002]; *dog vegan diet* and *switched to a therapeutic diet*, Kendall's tau 0.00, -Log$P$ 0.1 0 [P=0.843]).

Fig 2. Naïve correlation amongst owner-animal metadata variables. Data were first coerced into a numerical ordinal form, except for the Location variable which was not ordinal and, therefore, one-hot encoding was instead used. Correlations between variables were calculated using Kendall's tau, with positive and negative correlations coloured in blue and red, respectively. Both the size and intensity of the colour are proportional to the size of Kendall's tau. Not surprisingly, a large, dark blue circle is evident when variables are correlated with themselves. Hierarchical clustering, using the Ward method [58], is shown on the left side of the figure. All pairs are shown whether statistically significant or not; however, correlation coefficients and associated log P-values, corrected

488 for false-discovery rate using the Benjamini-Hochberg procedure, are shown in full in the

489 supplementary tables (S3 and S4 Tables).

491 **Owner, animal and healthcare variables associated with the *any***

492 ***health problem* binary variable**

493 **Simple regression**

Within the *any health problem* binary variable, 894 (38.5%) and 1429 (61.5%) dogs

were classified as 'yes' and 'no', respectively. Simple binary logistic regression was

performed first to assess the unadjusted effects of each predictor variable separately using

data from primary decision-makers only (Table 2). Broadly similar results were obtained

when logistical regression was performed on the dataset from all owners (irrespective of

decision-maker status; S1 Table). For example, estimates of the regression coefficients (β)

for the *dog diet vegan* variable was -0.43 (SE 0.131) using data from all owners and β -0.30

(SE 0.159) using data from primary decision-makers only.

Table 2. Results of simple (i.e., univariable) binary logistic regression analyses examining associations between owner, animal and veterinary variables and the *any health problem* binary (the presence of any illness as reported by the owner).

Variable [1]	Estimate	Odds ratio	99%-CI	*P*-value	Pseudo-R^2	BIC	AUC
Owner variables							
Location					0.0048	2231	0.491
United Kingdom	Ref	---	---	---			
Other European country	0.20 (0.147)	1.226	0.834, 1.787	0.165			
North America	0.30 (0.219)	1.355	0.765, 2.373	0.165			
Australia / New Zealand / Oceania	0.40 (0.241)	1.497	0.798, 2.783	0.094			
Other region	0.23 (0.299)	1.258	0.569, 2.698	0.442			
Setting [2]							
Urban	Ref	---	---	---	0.0034	2211	0.516
Rural	0.22 (0.106)	1.241	0.945, 1.629	0.041			
Owner age (years)							
<30	Ref	---	---	---	0.0025	2234	0.556
30-39	-0.01 (0.165)	0.989	0.646, 1.515	0.945			
40-49	-0.04 (1.66)	0.957	0.624, 1.470	0.794			
50-59	-0.24 (0.165)	0.789	0.516, 1.208	0.151			
≥60	-0.12 (0.169)	0.889	0.575, 1.375	0.488			
Owner gender					0.0029	2211	0.511
Female	Ref	---	---	---			
Male	-0.39 (0.211)	0.680	0.386, 1.154	0.068			
Education					0.0010	2228	0.444
Basic or high school	Ref	---	---	---			
College	0.00 (0.153)	1.002	0.676, 1.489	0.990			
Graduate	-0.11 (0.153)	0.893	0.602, 1.326	0.458			

Postgraduate	0.02 (0.159)	1.021	0.678, 1.541	0.897			
Income					0.0004	2222	0.538
Low	Ref	---	---	---			
Medium	-0.06 (0.140)	0.941	0.657, 1.355	0.666			
High	-0.13 (0.185)	0.879	0.555, 1.415	0.487			
Animal-related career					0.0005	2214	0.542
No	Ref	---	---	---			
Yes	0.10 (0.125)	1.108	0.800, 1.528	0.413			
Owner diet [3]					0.0089	2226	0.525
Omnivore	Ref	---	---	---			
Omnivore (restricted)	0.21 (0.134)	1.231	0.871, 1.735	0.120			
Pescatarian	0.67 (0.223)	1.947	1.094, 3.468	0.003			
Vegetarian	0.18 (0.177)	1.988	0.757, 1.883	0.305			
Vegan	0.02 (0.137)	1.013	0.710, 1.440	0.925			
Owner on vegan diet [3]					0.0007	2214	0.503
No	Ref	---	---	---			
Yes	-0.12 (0.125)	0.888	0.642, 1.221	0.341			
Animal Variables							
Age (per year) [4]					0.0842	2116	0.634
1 to 5	0.30 (0.192)	1.350	0.826, 2.229	0.119			
6 to 20	2.83 (0.285)	16.943	8.199, 35.639	<0.001			
Breed size category					0.0041	2232	0.533
Medium	Ref	---	---	---			
Toy	0.04 (0.345)	1.036	0.408, 2.460	0.919			
Small	0.30 (0.139)	1.350	0.943, 1.932	0.031			
Large	0.14 (0.119)	1.149	0.846, 1.563	0.242			

	Coef (SE)	OR	95% CI	p		n	
Giant	0.23 (0.251)	1.261	0.652, 2.391	0.355			
Sex					0.0002	2215	0.471
Female	Ref	---	---	---			
Male	0.06 (0.101)	0.946	0.729, 1.228	0.585			
Neuter status					0.0267	2182	0.522
Sexually intact	Ref	---	---	---			
Neutered	0.73 (0.132)	2.079	1.488, 2,943	<0.001			
Dog diet [3]					0.0270	2196	0.536
Conventional	Ref	---	---	---			
Raw	-0.61 (0.115)	0.544	0.404, 0.729	<0.001			
Vegetarian	0.08 (0.399)	1.083	0.374, 3.053	0.842			
Vegan	-0.53 (0.164)	0.591	0.384, 0.897	0.001			
Dog on vegan diet [2]					0.0030	2211	0.535
No	Ref	---	---	---			
Yes	-0.30 (0.159)	0.740	0.488, 1.106	0.058			

Healthcare variables

	Coef (SE)	OR	95% CI	p		n	
Veterinary visits					0.228	1933	0.770
None	Ref	---	---	---			
1	0.64 (0.186)	1.889	1.184, 3.053	<0.001			
2	1.94 (0.194)	6.940	4.266, 11.635	<0.001			
3	2.09 (0.234)	8.064	4.413, 15.114	<0.001			
4 or more	2.84 (0.232)	17.098	9.555, 31.709	<0.001			
Received medication					0.319	1772	0.768
No	Ref	---	---	---			
Yes	2.30 (0.118)	10.016	7.422, 13.622	<0.001			
Switched to therapeutic diet					0.0446	2160	0.542
No	Ref	---	---	---			

Yes	1.92 (0.294)	6,885	3.372, 15.657	<0.001	

Results presented are from simple (i.e., univariable) binary logistic regression, whereby each independent predictor variable is tested separately in a logistic regression model. These results were then used to determine the variables to include in subsequent multiple regression analysis, as shown in Fig 4 and S4 Table. Results are reported as estimated (β) with its standard error in brackets, odds ratios and 99% confidence intervals (99%-CI). Model performance assessed using the coefficient of determination (pseudo-R^2) based on the method reported by Nagelkerke [63], with pseudo-R^2, the Bayesian information criterion (BIC [60,61]) and area under the receiver operating characteristic curve (AUC) for the test dataset. For BIC, models having the best fit have lower BIC values; n.b., BIC can only be compared within the same family of models. For AUC, values can range from 0 to 1; a model that performed no better than chance would have an AUC of 0.5, and models predicting better than by chance would have AUC >0.5, with an AUC of 1.0 suggesting perfect prediction. [1] Definitions of the different categories are given in the original study [15]. [2] Please note that the urban category in *setting* variable combines the 'urban' and 'equally urban and rural categories'. [3] Please see the footnote to Table 1 for details of how owner and dog diets were assigned. [4] Dog age analysed as a continuous variable with B-splines, utilising boundary knots and an internal knot at the median value (6 years); therefore, odds ratios represent are the effect per year for each side of that knot.

Multiple regression

Next, we created two multiple regression models, again on data from primary decision-makers, with *any health problem* binary as the outcome variable and that either contained all predictor variables or just the owner-animal metadata. For the *all-variable* model (S3a Fig), the variables with the strongest associations with the outcome variable were *dog age* (both the '1-5y' and '6-20y' categories), *received medication*, *veterinary visits* (the '1 visit', '2 visits', '3 visits' and ≥4 visits' categories) and *switched to a therapeutic diet*. In terms of model performance, pseudo-R^2 was 0.439, BIC was 1836 and AUC on the test dataset was 0.852 (S5 Fig). For the *owner-animal metadata* model (S3b Fig), the variables with the strongest associations with the outcome variable were *owner age* ('50-59y' and '≥60y' categories), *dog age* ('6-20y' category), *breed size category* ('giant breed' category) and *dog diet* ('vegan diet' category). However, performance of this model was considerably worse in terms of generalisability (BIC 2244), the proportion of variance explained (pseudo-R^2 0.150) and prediction accuracy (AUC on the test dataset: 0.668).

Supervised backwards and forwards stepwise regression was then performed to create a best-fit multiple regression model for the *any health problem* binary outcome variable, starting with the model containing all variables. After refinement by backwards and forwards stepwise elimination, the best-fit model contained 4 variables: *dog age*, *veterinary visits*, *received medication* and *switched to a therapeutic diet* (Fig 3a; model 1, S5 Table). Compared with the all-variable model, generalisability was markedly improved (BIC 1656), with only slight reductions in both the amount of variability explained (pseudo-R^2 0.420) and prediction accuracy (AUC for the test dataset:0.842).

Fig 3. Final binary logistic regression model of variables associated with the *any health problem* binary in owners who were primary decision-makers. Results of the best-fit binary logistic

regression model of the *any health problem* binary (a) and the effect of adding either the *dog diet*

vegan (b) or *owner diet vegan* (c) to this best-fit model. The dots represent the odds ratio for each

variable, whilst the bars represent 99% confidence intervals (99%-CI). Variables where the 99%-CI

range does not include 1.0 (vertical dotted line) are depicted in red, whilst those that include 1.0 are

depicted in blue. Note the logarithmic scale for the X-axis in each graph.

As an additional multiple logistic regression step, the effect of adding the *dog diet*

vegan variable to this best-fit model was assessed (Fig 3b; model 2; S4 Table);

generalisability was worse (BIC 1660), whilst the amount of variance explained (pseudo-R^2

0.422) and prediction accuracy (AUC for the test dataset 0.845) were similar. Equivalent

results were obtained when the *owner diet vegan diet* variable was instead added to the final

best-fit model (BIC 1660, pseudo-R^2 0.421, AUC 0.844; Fig 3c; model 2; S4 Table).

555 Machine learning predictive modelling with XGBoost

The ability of owner, animal and healthcare variables to predict the *any health problem*

variable was further assessed using machine learning predictive modelling using the

XGBoost algorithm. Two separate models were created using data from primary decision-

makers; the first contained all variables, whilst the second was a reduced model containing

just the owner-animal metadata (i.e., without healthcare variables). Based on ROC analysis,

the prediction accuracy of the all-variable model was reasonable (AUC 0.836; Fig 4a). The

XGBoost procedure also enables the most important variables contributing to the prediction

to be identified, for the all-variable model (Fig 4b); the variables that most strongly predicted

the *any health problem* outcome variable were (in order, based on the sum of the 3 metrics):

received medication (importance 0.393, cover 0.129, frequency 0.065), *dog age* (importance

0.175, cover 0.182, frequency 0.188) and *veterinary visits* (importance 0.153, cover 0.181,

frequency 0.155), followed by *owner age* (importance 0.048, cover 0.083, frequency 0.106)

and then *breed size category* (importance 0.046, cover 0.070, frequency 0.087). Other

variables followed these including *education*, *dog sex*, *location*, and *dog diet*; for the latter

variable, *dog diet raw* was 10[th] in order of importance (importance 0.015, cover 0.026,

frequency 0.032), whilst *dog diet vegan* was 15[th] in order of importance (importance 0.008,

cover 0.021, frequency 0.013).

Fig 4. All-variable XGBoost model on the *any health problem* binary using the primary

decision-maker dataset. (a) Receiver operating characteristic curve of a prediction model containing

all variables (owner, animal and healthcare). This shows the increasing true positive and false

positive rates, with decrease of the threshold probability for prediction of *any health problem*.

Prediction accuracy was good as assessed by ROC analysis (area under curve 0.836, 99%-CI: 0.794-

0.878). Acceptance threshold is indicated by the colour bar on the right-hand side and shown at

discrete points on the curve. For example, with a threshold probability of 0.2, the model correctly

predicts approximately 80% of dogs where the *any health problem* binary was classified as 'yes', but

the false-positive rate is approximately 30%. (b) Graph depicting the relative contribution of

predictor variables to the all-variable XGBoost model for the *any health problem* binary. Importance'

represents fractional contribution of each feature to the model, based on the total gain from including

each feature; 'cover' represents the number of observations related to this feature in the model);

'frequency' represents the relative number of times a feature has been used in trees). Variables are

organised in order of importance left to right, based on the sum of the 3 metrics, whilst variables that

are not shown did not contribute to the final model.

The order of importance of predictor variables was broadly similar in the reduced

(owner-animal metadata) model although, based on ROC analysis, prediction accuracy was

poor (AUC 0.674, Fig 5a). In this reduced model, *dog age* was overwhelmingly most

important (importance 0.261, cover 0.196, frequency 0.157), followed by *owner age*

(importance 0.115, cover 0.123, frequency 0.141, *education level* (importance 0.095, cover

0.089, frequency 0.124), *breed size category* (importance 0.091, cover 0.099, frequency

0.108) and *neuter status* (importance 0.060, cover 0.062, frequency 0.047). Other variables

including *dog diet*, *urban setting*, *location* and *owner income* and followed these (Fig 5b); for

the dog diet variable, *dog diet raw* was 7th in order of importance (importance 0.054, cover

0.048, frequency 0.060), whilst *dog diet vegan* was 12th in order of importance (importance

0.039, cover 0.045, frequency 0.031).

Fig 5. Reduced (owner-animal metadata) XGBoost model on the *any health problem* binary

using the primary decision-maker dataset. (a) Receiver operating characteristic curve of a reduced

prediction model only containing owner and animal variables. This shows the increasing true positive

and false positive rates, with decrease of the threshold probability for prediction of *any health*

problem. Prediction accuracy was moderate, as assessed by ROC analysis (AUC 0.674, 99%-CI:

0.617-0.731). Acceptance threshold is indicated by the colour bar on the right-hand side and shown at

discrete points on the curve. For example, with a threshold probability of 0.2, the model correctly

predicts almost 80% of dogs where the *any health problem* binary was classified as 'yes', but the

false-positive rate is over 60%. The fact that the prediction thresholds are all low (< 0.5) shows that

this model struggles to predict health issues. (b) Graph depicting the relative contribution of predictor

variables to the all-variable XGBoost model for the *any health problem* binary. 'Importance'

represents fractional contribution of each feature to the model, based on the total gain from including

each feature; 'cover' represents the number of observations related to this feature in the model);

'frequency' represents the relative number of times a feature has been used in trees). Variables are

organised from left to right, based on the sum of the 3 metrics, whilst variables that are not shown did

not contribute to the final model.

Results were broadly similar when machine learning predictive modelling with

XGBoost was used to create both all-variable and reduced (owner-animal metadata) models

for the *any health problem* outcome measure using the all-owner dataset (S5 and S6 Figs).

Owner, animal and healthcare variables associated with the

significant illness binary variable

Simple regression

Within the *significant illness* binary variable, 122 (5.5%) and 2089 (94.5%) dogs were

classified as 'yes' and 'no', respectively. Results of the simple binary logistic regression

stage, to assess the unadjusted effects of each predictor variable separately using data from

primary decision-makers only, are shown in Table 3, whilst the results of equivalent analyses

on the dataset from all owners (irrespective of decision-maker status) are shown in S2 Table.

Once again, estimates of the regression coefficients were broadly similar from both data sets.

For example, estimates of the regression coefficients (β) for the *dog diet vegan* variable was -

0.46 (SE 0.309) using data from all owners and β -0.44 (SE 0.402) using data from primary

decision-makers only.

Table 3. Results of simple (i.e., univariable) binary logistic regression analyses examining associations between owner, animal and healthcare variables and the *significant illness* binary variable ('significant' or 'serious illness' as reported by owners who were primary decision-makers).

Variable [1]	Estimate	Odds ratio	99%-CI	*P*-value	Pseudo-R^2	BIC	AUC
Owner variables							
Location					0.0171	681	0.481
United Kingdom	Ref	---	---	---			
Other European country	0.58 (0.297)	1.787	0.788, 3.712	0.051			
North America	0.31 (0.483)	1.368	0.310, 4.083	0.516			
Australia / New Zealand / Oceania	1.06 (0.402)	2.884	0.901, 7.475	0.008			
Other region	0.77 (0.542)	2.163	0.394, 7.311	0.155			
Setting [2]					0.0007	667	0.516
Rural	Ref	---	---	---			
Urban	0.14 (0.234)	1.153	0.621, 2.086	0.542			
Owner age (years)					0.0060	687	0.488
<30	Ref	---	---	---			
30-39	0.43 (0.366)	1.539	0.617, 4.186	0.238			
40-49	0.07 (0.391)	1.067	0.392, 3.045	0.867			
50-59	0.16 (0.377)	1.171	0.450, 3.257	0.676			
≥60	-0.18 (0.417)	0.839	0.280, 2.513	0.674			
Owner gender					0.0011	667	0.501
Female	Ref	---	---	---			
Male	-0.38 (0.522)	0.683	0.129, 2.145	0.466			
Education					0.0035	681	0.530
Basic or high school	Ref	---	---	---			
College	-0.22 (0.360)	0.803	0.319, 2.105	0.543			

	Coef (SE)	OR	95% CI	p			
Graduate	0.79 (0.340)	1.083	0.461, 2.728	0.815			
Postgraduate	0.20 (0.348)	1.220	0.506, 3.123	0.586			
Income					0.0001	675	0.514
Low	Ref	---	---	---			
Medium	0.01 (0.314)	1.009	0.473, 2.447	0.977			
High	-0.07 (0.420)	0.932	0.304, 2.771	0.867			
Animal-related career					0.0041	666	0.533
No	Ref	---	---	---			
Yes	0.40 (0.257)	1.485	0.739, 2.811	0.124			
Owner diet [3]					0.0068	686	0.578
Omnivore	Ref	---	---	---			
Omnivore (restricted)	0.35 (0.296)	1.417	0.647, 3.026	0.239			
Pescatarian	-0.26 (0.618)	0.774	0.102, 2.994	0.678			
Vegetarian	0.57 (0.356)	1.774	0.664, 4.287	0.108			
Vegan	0.22 (0.310)	1.247	0.545, 2.736	0.475			
Owner on vegan diet [3]					0.000	668	0.485
No	Ref	---	---	---			
Yes	0.04 (0.274)	1.042	0.490, 2.036	0.881			
Animal Variables							
Age (per year) [4]					0.0592	643	0.704
1 to 5	0.10 (0.382)	1.104	0.333, 4.239	0.840			
6 to 20	2.98 (0.527)	19.652	5.124, 79.645	<0.001			
Giant breed [5]					0.0081	663	0.505
No	Ref	---	---	---			
Yes	0.91 (0.393)	2.473	0.786, 6.223	0.021			

Sex						0.0023	667	0.498
	Female	Ref	---	---	---			
	Male	-0.25 (0.227)	0.778	0.430, 1.340	0.270			
Neuter status						0.0081	663	0.546
	Sexually intact	Ref	---	---	---			
	Neutered	0.64 (0.330)	1.904	0.873, 4.906	0.051			
Dog diet [3]						0.0124	676	0.596
	Conventional	Ref	---	---	---			
	Raw	-0.58 (0.268)	0.559	0.269, 1.086	0.030			
	Vegetarian	0.24 (0.750)	1.269	0.091, 6.167	0.751			
	Vegan	-0.63 (0.410)	0.533	0.157, 1.375	0.125			
Dog on vegan diet [2]						0.0025	666	0.518
	No	Ref	---	---	---			
	Yes	-0.44 (0.402)	0.642	0.192, 1.615	0.271			
Healthcare variables								
Veterinary visits						0.2417	554	0.841
	None	Ref	---	---	---			
	1	0.31 (0.670)	1.358	0.279, 11.118	0.648			
	2	1.30 (0.645)	3.667	0.842, 28.960	0.044			
	3	1.68 (0.698)	5.372	0.983, 45.803	0.016			
	4 or more	3.54 (0.603)	34.615	9.942, 256.974	<0.001			
Received medication						0.197	558	0.777
	No	Ref	---	---	---			
	Yes	3.09 (0.427)	21.931	8.374, 81.002	<0.001			
Switched to therapeutic diet						0.050	640	0.555
	No	Ref	---	---	---			

	Yes	1.87 (0.310)	6.460	2.773, 13.900	<0.001

Results presented are from simple (i.e., univariable) binary logistic regression, whereby each independent predictor variable is tested separately in a logistic regression model. These results were then used to determine the variables to include in subsequent multiple regression analysis, as shown in Fig 4 and S4 Table. Results are reported as estimated (β) with its standard error in brackets, odds ratios and 99% confidence intervals (99%-CI). Model performance assessed using the coefficient of determination (pseudo-R^2) based on the method reported by Nagelkerke [63], with pseudo-R^2, the Bayesian information criterion (BIC [60,61]) and area under the receiver operating characteristic curve (AUC) for the test dataset. For BIC, models having the best fit have lower BIC values; n.b., BIC can only be compared within the same family of models. For AUC, values can range from 0 to 1; a model that performed no better than chance would have an AUC of 0.5, and models predicting better than by chance would have AUC >0.5, with an AUC of 1.0 suggesting perfect prediction. [1] Definitions of the different categories are given in the original study [15]. [2] Please note that the urban category in *setting* variable combines the 'urban' and 'equally urban and rural categories'. [3] Please see the footnote to Table 1 for details of how owner and dog diets were assigned. [4] Dog age analysed as a continuous variable with B-splines, utilising boundary knots and an internal knot at the median value (6 years); therefore, odds ratios represent are the effect per year for each side of that knot. [5] Please note that breed was better categorised using the *breed size category* variable for regression models using the *significant illness* outcome.

Multiple regression

Next, we created both an *all-variable* and an *owner-animal metadata* model, using data

from primary decision-makers, with *significant illness* binary as the outcome variable. For

the *all-variable* model (S4a Fig), the variables with the strongest associations with the

outcome variable were *dog age* ('6-20y' category), *giant breed*, and *received medication*. In

terms of model performance, pseudo-R^2 was 0.403, BIC was 679 and AUC on the test dataset

was 0.890. For the *owner-animal metadata* model (S4b Fig), the variables with the strongest

associations with the outcome variable were *dog age* ('6-20y' category) and *giant breed*.

Once again, the performance of this model was considerably worse in terms of

generalisability (BIC 799), the proportion of variance explained (pseudo-R^2 0.124) and

prediction accuracy (AUC on the test dataset: 0.692).

Supervised backwards and forwards stepwise logistic regression was again used to

create a best-fit multiple regression model for the *significant illness* outcome variable,

starting with all variables. After refinement by backwards and forwards stepwise elimination,

the best-fit model contained 3 variables: *dog age*, *veterinary visits* and *received medication*

(Fig 6a; model 1, S5 Table). Compared with the *all-variable* model, generalisability was

markedly improved BIC 517), with better prediction accuracy (AUC for the test dataset:

0.903), and only a small decreased in the amount of variability explained (pseudo-R^2 0.342).

Fig 6. Final best-fit binary logistic regression model of variables associated with the *significant*

***illness* binary in owners who were primary decision-makers.** Results of the best-fit binary logistic

regression model of the *significant illness* binary (a) and the effect of adding either the *dog diet vegan*

(b) or *owner diet vegan* (c) to this best-fit model. The dots represent the odds ratio for each variable,

whilst the bars represent 99% confidence intervals (99%-CI). Variables where the 99%-CI range does

not include 1.0 (vertical dotted line) are depicted in red, whilst those that include 1.0 are depicted in

blue. Note the logarithmic scale for the X-axis in each graph.

As a further multiple logistic regression step, the effect of adding the *dog diet vegan*

diet variable to this best-fit model was assessed (Fig 6b; model 2; S6 Table); generalisability

was worse (BIC 524), whilst the amount of variance explained (pseudo-R^2 0.342) and

prediction accuracy (AUC for the test dataset 0.904) were similar. Equivalent results were

obtained when the *owner diet vegan diet* variable was instead added to the final best-fit

model (BIC 524, pseudo-R^2 0.342, AUC 0.903; Fig 6c; model 2; S6 Table).

Machine learning predictive modelling with XGBoost

Using machine learning predictive modelling with XGBoost, and data from primary

decision-makers, all-variable and reduced (owner-animal metadata) models were created (S2

File) for the *serious illness* outcome variable. Based on ROC analysis, the prediction

accuracy of the all-variable model was reasonable (AUC 0.887; Fig 7a). In order (Fig 7b),

the variables that most strongly predicted the *significant illness* outcome variable were:

veterinary visits (importance 0.530, cover 0.473, frequency 0.285), *dog age* (importance

0.182, cover 0.219, frequency 0.302) and *received medication* (importance 0.175, cover

0.190, frequency 0.143), followed by *breed size category* (importance 0.050, cover 0.034,

frequency 0.079), *education* (importance 0.016, cover 0.020, frequency 0.050), *owner age*

(importance 0.012, cover 0.014, frequency 0.036) and *dog sex* (importance 0.009, cover

0.010, frequency 0.034). Other variables followed these including *location, urban setting*,

owner sex, *animal career* and *therapeutic food*. For the *dog diet* variable, *dog diet raw* was

17th in order of importance (importance 6.14 x10^{-5}, cover 4.09 x10^{-5}, frequency 1.55 x10^{-4})

relative importance), whilst the *dog diet vegan* variable was of insufficient importance to be

included in the model (i.e., relative importance less than that of the *dog diet raw*).

Fig 7. All-variable XGBoost model on the 'significant illness' binary using the primary

decision-maker dataset..   (a) Receiver operating characteristic curve of a prediction model

containing all variables (owner, animal and healthcare). This shows the increasing true positive and

false positive rates, with decrease of the threshold probability for prediction of *any health problem*.

Prediction accuracy was good, as assessed by ROC analysis (area under curve 0.887, 99%-CI: 0.816-

0.959). Acceptance threshold is indicated by the colour bar on the right-hand side and shown at

discrete points on the curve. For example, with a threshold probability of 0.14, the model correctly

predicts over 80% of dogs where the *any health problem* outcome was classified as 'yes', and the

false-positive rate is less than 15%. (b) Graph depicting the relative contribution of predictor

variables to the all-variable XGBoost model for the *any health problem* binary. 'Importance'

represents fractional contribution of each feature to the model, based on the total gain from including

each feature; 'cover' represents the number of observations related to this feature in the model);

'frequency' represents the relative number of times a feature has been used in trees). Variables are

organised from left to right, based on the sum of the 3 metrics, whilst variables that are not shown did

not contribute to the final model.

As with the *any health problem* binary, ROC analysis suggested that prediction

accuracy of the reduced (owner-animal metadata) model was poor (AUC 0.689, Fig 8a).

Once again, *dog age* (importance 0.304, cover 0.362, frequency 0.224) was the most

important predictor, followed by *breed size category* (importance 0.195, cover 0.226,

frequency 0.186), *owner age* (importance 0.091, cover 0.082, frequency 0.125), *urban setting*

(importance 0.077, cover 0.066, frequency 0.086) and *education* (importance 0.072, cover

0.060, frequency 0.094); other variables followed these including *dog sex*, *dog diet*, *location*,

animal career and *neuter status* (Fig 8b); for the dog diet variable, *dog diet raw* was 7[th] in

order of importance (importance 0.059, cover 0.042, frequency 0.047), whilst *dog diet vegan*

was 12[th] in order of importance (importance 0.018, cover 0.014, frequency 0.016).

Fig 8. Reduced (owner-animal metadata) XGBoost model on the *significant illness* binary

using the primary decision-maker dataset.. (a) Receiver operating characteristic curve of a reduced

prediction model only containing owner and animal variables. This shows the increasing true positive

and false positive rates, with decrease of the threshold probability for prediction of *significant illness*.

Prediction accuracy was moderate, as assessed by ROC analysis (area under curve 0.689, 99%-CI:

0.572-0.805). Acceptance threshold is indicated by the colour bar on the right-hand side, and also

shown at discrete points on the curve. For example, with a threshold probability of 0.05, the model

correctly predicts approximately 73% of dogs where the *significant illness* outcome was classified as

'yes', but the false-positive rate is approximately 45%. (b) Graph depicting the relative contribution

of predictor variables to the all-variable XGBoost model for the *any health problem* binary.

Importance' represents fractional contribution of each feature to the model, based on the total gain

from including each feature; 'cover' represents the number of observations related to this feature in

the model); 'frequency' represents the relative number of times a feature has been used in trees).

Variables are organised from left to right, based on the sum of the 3 metrics, whilst variables that are

not shown did not contribute to the final model.

Results were broadly similar when machine learning predictive modelling with

XGBoost was used to create both all-variable and reduced (owner-animal metadata) models

for the *significant illness* outcome variable using the all-owner dataset (S7 and S8 Figs).

Discussion

In the current study, we examined associations between owner perceptions of the health of their dog and a range of owner-related, animal-related and healthcare variables. Predictor variables that were most strongly associated with our outcome variables were dog age and healthcare variables (including number of veterinary visits and receiving medication), whilst the contribution from other variables was more limited, and the effect of dog diet was negligible. We utilised data from a previous study, where the opinions of owners about the health of their dog was gathered by questionnaire [15]. The primary aim of that study was to examine associations between owner opinions of dog health status and the type of diet that owners predominantly fed, with the key finding being that health status was positively associated with feeding either a vegan or raw diet, compared with other diet types (including conventional and vegetarian). Given the focus of that study, other information gathered in the questionnaire was not assessed and no account was taken of possible confounding amongst variables. Therefore, to extend the findings of the previous work, additional owner, animal and healthcare variables were studied. Previous work has reported associations between such variables and owner decisions about feeding. For example, animal variables, such as age and neuter status, were associated with the owner feeding choice in the previous study of Knight et al. [15]. In the same study, maintenance of pet health was the most-common reason cited by owners for choosing a particular food [15] whilst, in other research, owner characteristics (such as geographic location) are also associated with food choice [65]. Finally, owner characteristics can also be associated with aspects of dog health; for example, both owner age and income are associated with the prevalence of obesity in dogs [66]. Considering these previous findings, our decision to examine variables beyond those of diet type was justified.

To ensure that owner perceptions of dog health were examined, we created two

outcome variables from a question in the original survey question where owners were asked

to rate their dog's health over the previous 12 months [15]. Such responses are not objective

measures of health; rather, they require owners to formulate a subjective holistic opinion,

which will necessarily require them to take many factors into account (Fig 9a). There will be

a contribution from the actual health of their dog (which in turn is associated with other

factors), but this will be influenced by their knowledge and recollection, whereas general

opinions on what constitutes 'health' will be affected by owner attitudes, beliefs and

behaviours.

Fig 9. Flow diagrams illustrating possible pathways for associations between owner perceptions

of dog health and the owner, animal and healthcare variables. The green box depicts the outcome

variable (owner perception of health), blue boxes depict the predictor variables and yellow boxes

depict possible mechanisms through which predictor variables may act. Arrows represent the most

likely direction of associations, which are unidirectional except for associations between actual dog

health and healthcare variables (red double-headed arrow), because these might plausibly act in both

directions. (a) Possible associations amongst all available variables within the all-owner dataset; as

illustrated, owner perceptions of health will be associated with the actual health of the dog, but the

effect could be modified by owner knowledge and recollection, as well as being influenced by

attitudes, beliefs and behaviours; decision-maker status has the potential to influence both of these.

(b) Possible associations amongst all available variables for the primary decision-maker dataset,

where any variance from the decision-maker variable is eliminated by removal of owners who are not

primary decision-makers; this has the effect of simplifying possible pathways through which the

variables act. (c) Possible associations amongst variables in models on the primary decision-maker

dataset where the healthcare variables are not included; as illustrated, the impact of this is different

from (b) because no data or variance are removed; therefore, although the healthcare variables are

ignored in the modelling, the variance associated within them remains within the dataset, with the

potential to affect poor performance.

This study also differed from the original study because we excluded dogs <1y because

numbers were small (n=26) and, arguably, insufficient to reveal insights into how the

perceptions of owners with dogs in their growth phase might vary. We were also concerned

about the small numbers (n=111) of respondents who were not make the primary decision

about their dog's diet. Some of these owners reported playing "some lesser role", whilst

others reported playing "no role" in diet decision-making [15]. Not only would this raise

concerns about the accuracy of the diet information, but these owners might also be less

involved in other aspects of the dog's life, including health and veterinary matters, which

might adversely affect the accuracy of other information provided (Fig 9a). There might also

be differences in their opinions and attitudes towards health matters in general. Analyses

performed using *decision-maker status* as an outcome variable supported these concerns, not

least since the odds of a dog being recorded as having a health problem were less for owners

who were primary decision-makers than for those who were not. One way to account for this

would be to include *decision-maker status* as a covariate in all analyses but the marked

imbalance in group size (2,211 vs. 111) could adversely affect the performance of procedures

such as logistic regression. Therefore, we instead decided to limit further analyses to owners

who were primary decision-makers only. Decisions about excluding observations from a

dataset should be made with caution, properly justified and consider the possible impact on

results. Based upon our flow diagram of associations within in the dataset (Fig 9b), we

determined that it would be possible to eliminate all pathways associated with the *decision-*

maker status variable, thereby simplifying the overall structure without affecting any other

associations amongst variables. This was because removing all data from owners who are

not primary decision-makers effectively 'collapses' the *decision-maker status* variable into a

single category such that variance is no longer associated with it. Estimates of regression

coefficients and odds ratios were broadly similar in logistic regression models using the all-

owner dataset (S1 Table, S2 Table), compared with models using the primary decision-maker

dataset (Tables 2 and 3). Any differences are likely to be the result of differences in the

characteristics, attitudes and beliefs of the owners that were not primary decision-makers. Of

course, one main disadvantage of focusing on data from owners who are primary decision-

makers is that it limits the generalisability of findings to the wider pet-owning public, not

least to other members of a dog-owning family. However, given the original concern about

data reliability, further studies involving all family members would be required to explore

such opinions properly.

The approach of the current study also differed by our decision to include healthcare

variables (*received medication*, *veterinary visits*, *switched to a therapeutic diet*) as predictors.

In the previous study [15], these variables were instead used as proxy measures for dog health

and compared amongst dogs fed different diet types. Whilst both owner perceptions of health

and healthcare variables will be associated with actual dog health, they are materially

different. As already discussed, owner perceptions of health were derived from a question

where owners subjectively rated their dog's health [15], and these perceptions will be

influenced by other factors including owner knowledge and recollections, as well as their

beliefs and attitudes (Fig 9). Conversely, the healthcare variables were derived from survey

questions that required factual recall rather than the formulation of a subjective, holistic

opinion about health. For example, owners were asked whether their dog "had received

medications in the last year" and how many times the dog had "visited a veterinarian or

received a home visit" [15]. Given that such healthcare information will contribute to the

knowledge an owner utilises when constructing opinions about the health of their dog, we

reasoned that they were valid predictor variables in our statistical modelling. Further,

excluding them from statistical models could be problematic as illustrated in Fig 9c; unlike

removing observations from owners who were not primary decision-makers (as discussed

above), where any variance associated with the variable is removed, simply excluding a

variable from a statistical model will not eliminate any associated variance. The 'hidden'

variance associated with the healthcare variables might then be incorrectly attributed to other

variables leading to erroneous estimates of their regression coefficients. Any unassigned

variance also adversely affects model performance, which likely explains why the models

that only included owner-animal metadata performed poorly as discussed below.

Using simple logistic regression, we identified associations between owner perceptions

of dog health and several owner (e.g., *location*, *diet*), animal (e.g., *age*, *breed*, *neuter status*)

and healthcare (e.g., *veterinary visits*, *received medication*, *switched to a therapeutic food*)

variables. Further, owner perceptions of dog health were associated with the *dog diet*

variable in simple logistic regression (both *raw diet* and *vegan diet* associated with the *any*

health problem variable; *raw diet* associated with the *serious illness* variable). These

findings are not surprising, and would be expected because they were the result of univariable

analyses (simple logistic regression) and, therefore, were broadly similar to the univariable

analyses (e.g., odds ratios, one-way ANOVA) conducted in the previous study [15]. In the

current study, and as described in the methods section, data pre-processing was necessary to

ensure that our statistical analyses were valid. The approach taken was to remove or combine

groups where numbers of dogs was small, to ensure better balance amongst categories within

a predictor variable. The fact that the results obtained from our univariable analyses (simple

logistic regression) were similar to those of Knight et al. [15] suggests that, despite this pre-

processing, the final dataset remained representative of the dataset from which it was drawn.

We next used a combination of multiple logistic regression and machine learning

predictive modelling so that multiple variables could be analysed concurrently. This enabled

the creation of models that best predicted owner perceptions of dog health, and to determine

the relative importance of variables contributing to the final models. We chose this combined

methods approach to maximise the benefits of each whilst minimising their disadvantages.

Multiple logistic regression is a well-known, widely available technique that is familiar to

most people and has been extensively used in veterinary studies including those in dogs [67-

70]. Procedures for supervised model selection are well established, such as those based on

BIC [60,61], which can reduce the risk of model overfitting and can take relevant prior

knowledge into account. For example, in our exploratory analyses, we tested interactions

with a possible clinical (e.g., between the *veterinary visit* and *received medication* variables),

biological (between the *dog sex* and *neuter status* variables) or epidemiological (between the

location and *setting* variables) and psychological (between *owner diet* and *dog diet* variables).

One disadvantage of our manual selection method is that it could introduce bias into the

selection process. Other disadvantages include the fact that logistic regression makes certain

statistical assumptions (e.g., independence of errors, the requirement for a linear relationship

between the logit of the outcome and predictors, absence of multicollinearity and lack of

strong influential outliers; [71]) which, if not met, could lead to poor prediction accuracy.

Key advantages of machine learning methods are the fact that they make fewer prior

assumptions about data distribution and, therefore, can better model complex, nonlinear

relationships between the outcome and predictors even when there is significant noise [72].

However, there are disadvantages including requiring a large dataset, the time and resources

needed for computation, challenges with interpretation of the models produced and

susceptibility to errors [73]. In this respect, any machine learning procedure is susceptible to

overfitting, although this depends upon context; models are often intended to be generalisable

to a wide range of datasets beyond the dataset that they were trained on. Arguably, this

would be less of an issue for the current study because many of the variables of significance

were identified using both multiple logistic regression and machine learning predictive

modelling, whilst their relative contribution to model prediction were broadly similar (see

below). Therefore, despite the limitations of the different modelling techniques, our findings

should be valid.

Using multiple logistic regression, the main variables associated with dog health in the

best-fit models were *dog age*, *veterinary visits*, and whether the dog had been prescribed

medication; in some models, whether the dog had been switched to a therapeutic food was

also of significance, especially when owners described their dogs as "generally healthy, but

with minor or infrequent problems" (e.g., binary logistic regression on the *any health*

problem variable). Depending on the model used, these variables explained between 34%

and 52% of variance within the dataset, with very little additional variance (2% to 6%,

depending on the model) gained by the addition of any or all other predictor variables. Many

of the same variables were also identified using XGBoost, and prediction accuracy (as

measured by AUC) was similar, but other predictor variables were also identified including

owner age and *education*. However, in contrast to the findings of the previous study [15], the

diet of the dog had little to no effect in all models, except for those that only included owner-

animal metadata, albeit that these models performed poorly. Therefore, we could not find

convincing evidence that particular food types (conventional, vegan, vegetarian or raw) have

any clinically meaningful positive or negative association with dog health status. As with the

original study [15], these analyses were limited by the dataset used, including its cross-

sectional design, meaning that causality of associations cannot be assumed, and its reliance

on subjective owner perceptions of health status, which might not reflect the actual health of

the dogs. Future studies should be considered, for example, cohort studies or randomised

trials utilising objective measures of health such as information from electronic patient record

as or formal diagnoses of disease by a veterinarian.

A key limitation of using questionnaires to ascertain the perceptions of owners about

dog health is that such information might be affected by owner recollection leading to errors

in classification of disease status, so-called response bias [74]. Responses might be biased

for various reasons including acquiescence, courtesy, the order of questions or even the

perceived social desirability of the responses [74]. Recall bias is a form of response bias

whereby the recollections of the owner are affected by their past experiences [75]. Owners of

dogs with health problems might search their memories more thoroughly for possible risk

factors (such as diet choice, number of visits to the vet, use of medication etc, diet changes)

than owners of dogs that do not have health problems, with the potential to exaggerate any

associations identified.

As discussed above, it is also feasible that the attitudes and beliefs of owners might

either consciously or unconsciously have influenced responses about the health of their dog, a

point that the authors of the previous study emphasised [15]. For example, owners who

believe a particular type of diet to be optimal for dog health, might be more likely to perceive

their dog to be healthy, whether or not this was objectively true. Whilst such a bias would

feasibly affect any diet type, a greater bias might be expected with diets perceived to be

'unconventional', again, as acknowledged in the previous study [15]. Although increasing in

popularity [1], both vegan diets and raw meat diets are still uncommon choices for owners,

and many veterinary professionals either do not recommend them or might even advise

against their use. Current evidence suggests that both such diet types are often not

formulated appropriately to be complete and balanced for essential nutrients [5,8,9-11].

Further, there are also concerns with feeding raw food given the potential for contamination

either with bacteria of possible pathogenic potential or bacteria resistant to antimicrobials,

both of which might pose a risk to the health of owners and other in-contact people [11-13].

Given such health concerns, owners who make an active choice to feed either a raw or vegan

diet could be more defensive about this diet choice, compared with owners who feed

conventional diets, and this attitude which might unconsciously have biased responses about

the health of their dog, for example, minimising the significance of any illnesses.   In the

original study questionnaire, the authors did attempt to minimise the influence of canine diet

variables on reported health status, by placing the health-related questions before most other

variables [15].   However, it is unclear as to whether this would adequately eliminate pre-

existing unconscious biases resulting from diet choice.

One variable that featured consistently in all models was dog age, whereby illness

severity worsened as age increased, especially when in dogs that were ≥6 years. Perceptions

of health in senior dogs are likely to be associated with quality of life, which is known to be

negatively associated with age [76].   However, it might also be because many chronic

diseases have associations with increasing age, including osteoarthritis [68], obesity [77],

diabetes mellitus [70] and many forms of cancer [78]].   It is noteworthy that such chronic

diseases are also negatively associated with quality of life [79-82].   An alternative

explanation for the negative association between age and health in the current study is the

fact that owners of older dogs perceive their health to be poorer simply because they expect

older dogs to be less healthy [83].

Owner perceptions about dog health were also positively associated with them

reporting that their dog had received medication in the last 12 months. Once again, the

observational study design means that causality cannot be assumed. It is unclear whether

such dogs are receiving these medications because they are genuinely less healthy (which is

why they were prescribed), whether the knowledge that their dog is receiving medication

means the owner perceives their dog to be less healthy or both (Fig 9).   Finally, the

association might be indirect, if both the outcome and predictor variable were associated with

a third variable. In this regard, there was also a strong positive association between owner

perception of health and the frequency of veterinary visits, whilst a positive correlation

between frequency of veterinary visits and receiving medication was also evident, meaning

that all three variables are likely to be inter-related. Further, in some models, another

healthcare variable (switching to a therapeutic diet) potentially associated with frequency of

veterinary visits was also included. Besides these, several additional associations were

identified with XGBoost, but not with logistic regression, the most important of which were

owner age and *education*. Owners that were older and those with a higher level of education

perceived their dogs to be healthier than did other owners. Further studies would be required

to explore all such associations further and determine their true significance to canine health.

Although the study was large, the population studied was not representative of the

typical dog-owning public. In this respect, 33% and 13% of owners reportedly fed their dog

raw and vegan diets, respectively, which is greater than expected; for example, in a recent

UK survey, the proportion of owners feeding raw and vegan diets were 7% and <1%,

respectively [1]. Further, the proportion of the UK human population reportedly consuming a

vegan diet is estimated to be between 2 and 3% [2] whilst, in the current study, 23% of

owners reported consuming a vegan diet. This raises a concern about how representative this

study population as a whole, and is of particular significance given the strong clustering

based on owner diet. Although the reasons for such an imbalance is not clear, it might well

have resulted from the method of owner recruitment. In this respect, the study was widely

advertised via social media but, given concerns that owners feeding dogs unconventional

diets might be under-represented, relevant interest groups were actively targeted and invited

to participate. This strategy was successful in ensuring that adequate numbers of dog owners

feeding unconventional diets were included, ensuring that statistical comparisons were

meaningful in both the previous [15] and current studies. However, the recruitment strategy

might have inadvertently generated an unrepresentative study population which, in turn, may

have influenced the results obtained, as discussed above.

In addition to the unrepresentative study population, multiple associations amongst

groups were identified with the potential for confounding amongst variables. For example,

feeding a raw diet was more commonly reported by owners from the UK, whilst feeding a

vegan diet was more common in owners from Europe. Further, there were positive

associations between owner income and education, and between dog age and neuter status.

Including healthcare variables in multiple logistic regression analyses might also have hidden

the associations seen with the *owner-animal metadata*, including diet type. In logistic

regression analyses, we accounted for these associations by determining multicollinearity

using VIF, refining models where this might be a problem. No significant issues of multi-

collinearity amongst predictor variables were identified in any of the final models. Whilst

this does not necessarily guarantee that variance from each factor in the models was correctly

assigned, any errors are likely to be small. The use of XGBoost would further address these

concerns given that these techniques do not make prior assumptions about data distribution

and can better take account of complexity in interactions amongst variables [69].

Nonetheless, to ensure any *owner-animal metadata* effects were not unfairly penalised by the

inclusion of the healthcare variables, we also examined models containing only the *owner-*

animal metadata. In these models, *dog age* remained as the variable of most importance, but

other variables of lesser importance were identified that had not been seen on the original

models (including *breed size category*, *owner age*, *education level* and *urban setting*), whilst

a limited effect of diet remained (i.e., the *vegan diet* variable was only the 12th most

important predictor variable). The significance of any associations between these variables

owner perceptions of health are not known. Nevertheless, since predictive accuracy of all

models (both multiple logistic regression and XGBoost) utilising just the *owner-animal*

metadata was poor, these analyses should be considered exploratory and any conclusions

drawn should be made with caution.

A final approach we took, to ensure that the we fully accounted for any possible effect

of the *dog diet vegan* variable, was to add it back into the best-fit models for both the *any*

health problem and *serious illness* outcome variables. In both cases, this additional predictor

variable did not improve the performance of either model and, interestingly, models with

equivalent performance could be created when the *owner diet vegan* was instead added.

The main limitations to the current study have already been discussed above, including

the fact that owner-reported health was the main outcome measure and the fact that the study

population was not representative of the general dog-owning population. Further, significant

data pre-processing was required to ensure adequate group sizes for statistical analyses. The

disadvantage of such an approach was that many smaller groups needed to be excluded,

meaning that we might have missed some variables with a potential impact on owner-

reported canine health. That said, in preliminary testing, we coded several of our variables in

different ways (e.g., *dog age* as a continuous or ordinal 5-category variable; *veterinary visits*

as a 4- or 5-category ordinal variable), and selected the approach that produced the best

model fit. Therefore, whilst some genuine effects from small categories might still have been

missed, the final best-fit models we selected were those that most closely fitted the data.

A third limitation was the fact that we did not contact the authors of the original study

before conducting our data analyses, to clarify any uncertainties with the dataset and seek

guidance on our planned approach. We chose not to do so because the original paper was

well written, with clearly presented study methodology and results. Further, since

independent replication is critical to the scientific method, it was arguably better not to have

direct contact with the primary study authors when conducting statistical analyses.

A fourth limitation was that the outcome variables tested in the current study were

derived from only one of the health metrics examined in the original study, namely owner-

reported dog health [15]. Other health metrics reported by Knight *et al*. included asking

owners "to report what they believed their veterinarian's assessment [of their dog's health] to

be" [15] and owner reports of 22 possible health disorders, grouped into broad categories

(e.g., allergy, cancer/tumours, hormonal, skin/coat) [15]. Our conclusions might have

differed if one or more of these alternative health metrics had been analysed, but we decided

against this because, in our opinion, all were inferior measures. Not only were these metrics

also based on subjective owner reports, but owners were required to speculate on what their

veterinarian's opinion about the health of their dog would be. Further, there were no clear

definitions of the health disorder categories, meaning that these would likely be broad,

comprising many different diseases with differing aetiology and severity. Finally, the

number of dogs included in many categories was small, being ≤50 in 12 of the 22 [15].

Whilst further work could be considered, where these additional metrics are analysed using

the techniques of the current study, it is questionable as to whether anything further would be

gained. Instead, it would be preferable to conduct a prospective study utilising objective,

veterinary-derived measures of health, as discussed above.

A final limitation was the fact that there might be confounding between healthcare

variables (e.g., *veterinary visits* and *received medication*) and dietary choices. For example,

owners who are sceptical about conventional diets and instead choose to feed either vegan or

raw food, might also be less willing either visiting their veterinarian or allowing their dog to

receive medication. In support of this, negative associations were identified in the current

study between feeding a raw diet and both veterinary visit frequency and the use of

medications; further, in the previous study, dogs fed raw diets were less likely to be neutered

than dogs fed conventional diets [15]. However, results were inconsistent for dogs fed vegan

diets; whilst there was a negative association between feeding a vegan diet and the odds of a

dog receiving medications, there was no association with frequency of veterinary visits.

Further, in the previous study, dogs fed vegan diets were more likely to be neutered than

those fed conventional diets [15]. Given these complexities, further research would be

required to investigate associations between different diet choices and healthcare variables.

1078

1079 Conclusions

1080 Variables most strongly associated with owner perceptions of health are dog age and

1081 healthcare variables including number of veterinary visits, receiving medication and being

1082 switched to a therapeutic diet. These results extend the findings of previously-published

1083 research [15] and suggest that, when other variables are accounted for, the association

1084 between dog diet choice and owner-perceived canine health is negligible.

1085

1086 Acknowledgements

1087 The authors wish to acknowledge the authors of the original paper [15] for providing

1088 making their data freely available to other researchers for further analysis.

1089

1090 Funding statement

1091 This study was not funded by any research grant. The costs of open access publishing

1092 were covered by the University of Liverpool Institutional Open Access Fund

1093 (https://www.liverpool.ac.uk/open-research/open-access/open-access-funding/).

1094

1095 Competing interests statement

AJG is an employee of the University of Liverpool, but his position is financially-supported by Royal Canin. AJG has also received financial remuneration and gifts for providing educational material, speaking at conferences, and consultancy work, all unrelated to the current study. Royal Canin had no involvement in any aspect of the current work including study design, data analysis, drafting the manuscript or the decision to submit the work for publication.

RBJ is an employee of the University of Liverpool, whose salary is financially-supported by the Higher Education Funding Council for England. RBJ also receives a stipend from United Kingdom Research and Innovation for work as a panellist for the Biotechnology and Biological Sciences Research Council.

None of these disclosures alter our adherence to PLOS ONE policies on sharing data and materials.

Research data for this article

The original research data are accessible at https://rebrand.ly/2020/pet-food-consumers-study. All statistical reports and codes are available in the supplementary information (S1-S5 Files) and online (https://github.com/RichardBJ/CanineHealth22).

References

1. PDSA. PAW Report 2022. [cited 2022 Dec 19]. Available from: https://www.pdsa.org.uk/media/12965/pdsa-paw-report-2022.pdf.

2. YouGov 2022. Dietary choices of Brits (e.g. vegetarian, flexitarian, meat-eater etc). 2022 [cited 2022 Oct 20]. Available from:

https://yougov.co.uk/topics/lifestyle/trackers/dietery-choices-of-brits-eg-

vegeterian-flexitarian-meat-eater-etc.

3.   New Nutrition Business. 10 key trends in food, nutrition & health 2020. 2022

[cited   2022   Dec   19].   Available   from: https://www.new-

nutrition.com/keytrend?id=135.

4.   Gallup. Snapshot: few Americans vegetarian or vegan [internet]. 2018 [cited

2022 Oct 20]. Available from: https://news.gallup.com/poll/238328/snapshot-few-

americans-vegetarian-

vegan.aspx?g_source=link_NEWSV9&g_medium=NEWSFEED&g_campaign=it

em_&g_content=Snapshot%3a%2520Few%2520Americans%2520Vegetarian%25

20or%2520Vegan.

5.   Dodd SAS, Adolphe JL, Verbrugghe A. Plant-based diets for dogs. J Am Vet Med

Assoc. 2018 Dec 1;253(11):1425-1432. doi: 10.2460/javma.253.11.1425

6.   Dodd S, Khosa D, Dewey C, Verbrugghe A. Owner perception of health of North

American dogs fed meat- or plant-based diets. Res Vet Sci. 2022 Dec;149:36-46.

doi: 10.1016/j.rvsc.2022.06.002

7.   Morelli G, Bastianello S, Catellani P, Ricci R. Raw meat-based diets for dogs:

survey of o'ners' motivations, attitudes and practices. BMC Vet Res. 2019 Mar

4;15(1):74. doi: 10.1186/s12917-019-1824-x

8.   Dodd SAS, Shoveller AK, Fascetti AJ, Yu ZZ, Ma DWL, Verbrugghe A. A

Comparison of Key Essential Nutrients in Commercial Plant-Based Pet Foods

Sold in Canada to American and European Canine and Feline Dietary

Recommendations.   Animals   (Basel).   2021   Aug   9;11(8):2348.   doi:

10.3390/ani11082348

9.   Larsen JA, Parks EM, Heinze CR, Fascetti AJ. Evaluation of recipes for home-

1144   prepared diets for dogs and cats with chronic kidney disease. J Am Vet Med Assoc.

1145   2012 Mar 1;240(5):532-8. doi: 10.2460/javma.240.5.532

1146   10.   Stockman J, Fascetti AJ, Kass PH, Larsen JA. Evaluation of recipes of home-

1147   prepared maintenance diets for dogs. J Am Vet Med Assoc. 2013 Jun

1148   1;242(11):1500-5. doi: 10.2460/javma.242.11.1500. Erratum in: J Am Vet Med

1149   Assoc. 2014 Jul 15;245(2):177

1150   11.   Vecchiato CG, Schwaiger K, Biagi G, Dobenecker B. From nutritional adequacy

1151   to hygiene quality: a detailed assessment of commercial raw pet-food for dogs and

1152   cats. Animals (Basel). 2022 Sep 13;12(18):2395. doi: 10.3390/ani12182395

1153   12.   Groat EF, Williams NJ, Pinchbeck G, Warner B, Simpson A, Schmidt VM. UK

1154   dogs eating raw meat diets have higher risk of Salmonella and antimicrobial-

1155   resistant Escherichia coli faecal carriage. J Small Anim Pract. 2022 Jun;63(6):435-

1156   441. doi: 10.1111/jsap.13488

1157   13.   Sealey JE, Hammond A, Mounsey O, Gould VC, Reyher KK, Avison MB.

1158   Molecular ecology and risk factors for third-generation cephalosporin-resistant

1159   Escherichia coli carriage by dogs living in urban and nearby rural settings. J

1160   Antimicrob Chemother. 2022 Aug 25;77(9):2399-2405. doi: 10.1093/jac/dkac208

1161   14.   Anturaniemi J, Barrouin-Melo SM, Zaldivar-López S, Sinkko H, Hielm-Björkman

1162   A. Owners' perception of acquiring infections through raw pet food: a

1163   comprehensive internet-based survey. Vet Rec. 2019 Nov 30;185(21):658. doi:

1164   10.1136/vr.105122

1165   15.   Knight A, Huang E, Rai N, Brown H. Vegan versus meat-based dog food:

1166   Guardian-reported indicators of health. PLoS One. 2022 Apr 13;17(4):e0265662.

1167   doi: 10.1371/journal.pone.0265662.

16. Chen, T., Guestrin, C. XGBoost: a scalable tree boosting system [cited 2022 Nov 8]. Available from: https://arxiv.org/abs/1603.02754#.

17. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2022 [cited 2022 Nov 30]. Available from: https://www.R-project.org/

18. Lesnoff, M., Lancelot, R. (2012). aod: analysis of overdispersed data. R package version 1.3.2. 2012 [cited 2022 Dec 19]. Available from: http://cran.r-project.org/package=aod

19. Robinson D, Hayes A, Couch S. broom: convert statistical objects into tidy tibbles; R package version 0.8.0. 2022 [cited 2022 Dec 19]. Available from: https://CRAN.R-project.org/package=broom.

20. Fox J, Weisberg S. An R Companion to Applied Regression, Third Edition; 2019 [cited 2022 Dec 19]. Available from: https://socialsciences.mcmaster.ca/jfox/Books/Companion/

21. Kuhn M. caret: classification and regression training; R package version 6.0-93. 2022 [cited 2022 Dec 19]. Available from: https://CRAN.R-project.org/package=caret.

22. Maechler M, Rousseeuw P, Struyf A, Hubert M, Hornik K (2022). cluster: cluster analysis basics and extensions. R package version 2.1.4, 2022 [cited 2023 Nov 30]. Available from https://CRAN.R-project.org/package=cluster

23. Walesiak M, Dudek A (2020). The choice of variable normalization method in cluster analysis. In Soliman KS (ed.) Education excellence and innovation management: a 2025 vision to sustain economic development during global challenges. pp 325-340. ISBN 978-0-9998551-4-1.

1192   24.   Wei T, Simko V (2021). R package 'corrplot': visualization of a correlation matrix

1193   (version 0.92). 2021 [cited 2022 Nov 8]. Available from:

1194   https://github.com/taiyun/corrplot.

1195

25.   Dowle M, Srinivasan A. data.table: extension of `data.frame`; R package version

1.14.6.   2022 [cited 2022 Dec 19].   Available from: https://CRAN.R-

project.org/package=data.table.

26.   Hahsler M, Piekenbrock M (2022). dbscan: density-based spatial clustering of

applications with noise (DBSCAN) and related algorithms; R package version 1.1-

11.   2022 [cited 2022 Dec 19].   Available from: https://CRAN.R-

project.org/package=dbscan.

27.   Signorell A, Aho K, Alfons A, Anderegg N,Aragon T, Arachchige A, et al.

DescTools: tools for descriptive statistics; R package version 0.99.47. 2022 [cited

2022 Dec 19]. Available from: https://cran.r-project.org/package=DescTools

28.   Wickham H, François R, Henry L, Müller K. dplyr: A Grammar of Data

Manipulation. R package version 1.0.9. 2022 [cited 2022 Dec 19]. Available

from: https://CRAN.R-project.org/package=dplyr.

29.   Kassambara A, Mundt F (2020). factoextra: extract and visualize the results of

multivariate data analyses. R package version 1.0.7., 2020 [cited 2023 Nov 30].

Available from: https://CRAN.R-project.org/package=factoextra

30.   Nakazawa M. fmsb: functions for medical statistics book with some demographic

data. R   package version 0.7.3. 2022 [cited 2022 Dec 19].   Available from:

https://CRAN.R-project.org/package=fmsb

31.   R Core Team. foreign: read data stored by 'Minitab', 'S', 'SAS', 'SPSS', 'Stata',
1216   'Systat', 'Weka', 'dBase', ..; R package version 0.8-82. 2022 [cited 2022 Dec 19].
1217   Available from: https://CRAN.R-project.org/package=foreign.

1218   32.   Pedersen T. ggforce: Accelerating 'ggplot2'_. R package version 0.4.1. 2020 [cited
1219   2023 Dec 6]. Available from : https://CRAN.R-project.org/package=ggforce

1220

1221   33.   Wickham H. ggplot2: elegant graphics for data analysis. 3rd ed. New York:
1222   Springer-Verlag New York, 2016.

1223   34.   Arnold J. ggthemes: extra themes, scales and geoms'for 'gg'lot2'; R package
1224   version 4.2.4. 2021 [cited 2022 Dec 19]. Available from: https://CRAN.R-
1225   project.org/package=ggthemes.

1226   35.   Vanegas L, Rondón L, Paula G. (2022). glmtoolbox: set of tools to data analysis
1227   using generalized linear models; R package version 0.1.4. 2022 [cited 2022 Dec
1228   19]. Available from: https://CRAN.R-project.org/package=glmtoolbox.

1229   36.   Galili'T, O'Callaghan A, Sidi J, Sievert C. heatmaply: an R package for creating
1230   interactive cluster heatmaps for online publishing. Bioinformatics. 2018 May
1231   1;34(9):1600-1602. doi: 10.1093/bioinformatics/btx657

1232   37.   Harrell Jr F. Hmisc: Harrell miscellaneous. R package version 4.7-0 [internet].
1233   2022   [cited   2022   Dec   12].   Available   from: https://CRAN.R-
1234   project.org/package=Hmisc

1235   38.   Ignatiadis N, Klaus B, Zaugg JB, Huber W. Data-driven hypothesis weighting
1236   increases detection power in genome-scale multiple testing. Nat Methods. 2016
1237   Jul;13(7):577-80. doi: 10.1038/nmeth.3885

1238   39.   Gorman B. mltools: machine learning tools. R package version 0.3.5, 2018 [cited
1239   2022 Dec 12]. Available from: https://CRAN.R-project.org/package=mltools.

1240   40.   Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, et al. pROC: an

1241   open-source package for R and S+ to analyze and compare ROC curves. BMC

1242   Bioinformatics. 2011 Mar 17;12:77. doi: 10.1186/1471-2105-12-77

1243   41.   Neuwirth E (2022). _RColorBrewer: ColorBrewer Palettes_. R package version

1244   1.1-3.   2022   [cited   2023   Dec   6].   Available   from: https://CRAN.R-

1245   project.org/package=RColorBrewer

1246   42.   Wickham H, Bryan J. readxl: read Excel files; R package version 1.4.0, 2022

1247   [cited 2022 Dec 19]. Available from: https://CRAN.R-project.org/package=readxl.

1248   43.   Wickham H. Reshaping data with the reshape package. J Stat Soft 2007;21(12):1-

1249   20. doi: 10.18637/jss.v021.i12

1250   44.   Sing T, Sander O, Beerenwinkel N, Lengauer T. ROCR: visualizing classifier

1251   performance   in   R.   Bioinformatics.   2005   Oct   15;21(20):3940-1.   doi:

1252   10.1093/bioinformatics/bti623

1253   45.   Silge J, Chow F, Kuhn M, Wickham H. rsample: General Resampling

1254   Infrastructure; R package version 1.1.0, 2022 [cited 2022 Dec 19]. Available

1255   from: https://CRAN.R-project.org/package=rsample.

1256   46.   Kassambara A. rstatix: pipe-friendly framework for basic statistical tests. R

1257   package   version   0.7.0.   2021   [cited   2022   Nov   8].   Available   from:

1258   https://CRAN.R-project.org/package=rstatix.

1259   47.   Qiu Y, details. aotisSfAf. showtext: using fonts more easily in R graphs; R

1260   package   version   0.9-5.   2022   [cited   2022   Dec   19].   Available   from:

1261   https://CRAN.R-project.org/package=showtext.

1262   48.   Siriseriwan W. smotefamily: A collection of oversampling techniques for class

1263   imbalance problem based on SMOTE; R package version 1.3.1. 2019 [cited 2022

1264   Dec 19]. Available from: https://CRAN.R-project.org/package=smotefamily.

1265   49.   Wickham H. stringr: simple, consistent wrappers for common string pperations; R

1266   package version 1.4. 2019 [cited 2022 Dec 19]. Available from: https://CRAN.R-

1267   project.org/package=stringr.

1268   50.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, Francois R, et al.

1269   Welcome to the tidyverse. J Open Source Software. 2019;4(43):1686. doi:

1270   10.21105/joss.01686

1271   51.   Melville J (2023). _uwot: the uniform manifold approximation and projection

1272   (UMAP) method for dimensionality reduction_. R package version 0.1.16. 2023

1273   [cited 2023 Dec 6]. Available from: https://CRAN.R-project.org/package=uwot

1274   52.   McInnes L, Healy J, Melville J. UMAP: uniform manifold approximation and

1275   projection for dimension reduction. arXiv:1802.03426v3. 2018 [cited 2022 Dec

1276   19]. doi: 10.48550/arXiv.1802.03426

1277   53.   Rousseeuw PJ. Silhouettes: a graphical aid to the interpretation and validation of

1278   cluster analysis. J Comput Appl Math. 1987;20:53-65. Journal of Computational

1279   and Applied Mathematics. 20. 53-65. doi: 10.1016/0377-0427(87)90125-7

1280   54.   Davies DL, Bouldin DW. A cluster separation measure. IEEE Trans Pattern Anal

1281   Machine Intel. 1979;1:224-227. doi: 10.1109/TPAMI.1979.4766909

1282   55.   Kaufmann L, Rousseeuw PJ. Finding groups in data: an introduction to cluster

1283   analysis. Hoboken, NJ: Wiley-Interscience. p.87. 1990.

1284   doi:10.1002/9780470316801. ISBN 9780471878766.

1285   56.   Kendall MG. The treatment of ties in ranking problems. Biometrika. 1945

1286   Nov;33:239-51. doi: 10.1093/biomet/33.3.239

1287   57.   Friendly M. Corrgrams: exploratory displays for correlation matrices. Am Stat.

1288   2002;56:316–324. doi: 10.1198/000313002533

1289   58.   Murtagh, F., Legendre, P. Ward's Hierarchical Agglomerative Clustering Method:

1290   Which Algorithms Implement Ward's Criterion?. J Classif. 2014;31:274–295. doi:

1291   10.1007/s00357-014-9161-z

1292   59.   Cohen J. A power primer. Psychol Bull. 1992 Jul;112(1):155-159. doi:

1293   10.1037//0033-2909.112.1.155

1294   60.   Schwartz, G. Estimating the dimension of a model. Ann Stat. 1978;6:461-464.

1295   61.   Neath, A.A., Cavanaugh, J.E. The Bayesian information criterion: background,

1296   derivation, and applications.   WIREs Comput Stat 2012;4:199-203. doi:

1297   10.1002/wics.199

1298   62.   Fox J, Monette G. Generalized collinearity diagnostics.   J Am Stat Assoc.

1299   1992;87:417, 178-183, DOI: 10.1080/01621459.1992.10475190

1300   63.   Nagelkerke NJD.   A note on a general definition of the coefficient of

1301   determination. Biometrika. 1991;78:691-692. doi: 10.1093/biomet/78.3.691

1302   64.   Wong SC, Gatt A, Stamatescu V, McDonnell MD.   Understanding data

1303   augmentation for classification: when to warp? 2016 [cited 22 Nov 8]. Available

1304   from: https://arxiv.org/abs/1609.08764.

1305   65.   Hoummady S, Fantinati M, Maso D, Bynens A, Banuls D, Santos NR, Roche M,

1306   Priymenko N. Comparison of canine owner profile according to food choice: an

1307   online preliminary survey in France. BMC Vet Res. 2022 May 4;18(1):163. doi:

1308   10.1186/s12917-022-03258-9

1309   66.   Courcier EA, Thomson RM, Mellor DJ, Yam PS. An epidemiological study of

1310   environmental factors associated with canine obesity. J Small Anim Pract. 2010

1311   Jul;51(7):362-7. doi: 10.1111/j.1748-5827.2010.00933.x

67.   Westgarth C, Pinchbeck GL, Bradshaw JW, Dawson S, Gaskell RM, Christley RM.

Factors associated with dog ownership and contact with dogs in a UK community.

BMC Vet Res. 2007 Apr 3;3:5. doi: 10.1186/1746-6148-3-5

68.   Anderson 'L, O'Neill DG, Brodbelt DC, Church DB, Meeson RL, Sargan D, et al.

Prevalence, duration and risk factors for appendicular osteoarthritis in a UK dog

population under primary veterinary care. Sci Rep. 2018 Apr 4;8(1):5641. doi:

10.1038/s41598-018-23940-z

69.   Zapata I, Moraes LE, Fiala EM, Zaldivar-Lopez S, Couto CG, Rowell JL, et al.

Risk-modeling of dog osteosarcoma genome scans shows individuals with

Mendelian-level polygenic risk are common. BMC Genomics. 2019 Mar

19;20(1):226. doi: 10.1186/s12864-019-5531-6

70.   Heeley AM, O'Neill DG, Davison LJ, Church DB, Corless EK, Brodbelt D.

Diabetes mellitus in dogs attending UK primary-care practices: frequency, risk

factors and survival. Canine Genet Epidemiol. 2020;7: 6. doi: 10.1186/s40575-

020-00087-7

71.   Stoltzfus JC. Logistic regression: a brief primer. Acad Emerg Med. 2011

Oct;18(10):1099-104. doi: 10.1111/j.1553-2712.2011.01185.x

72.   Gravesteijn BY, Nieboer D, Ercole A, Lingsma HF, Nelson D, van Calster B,

Steyerberg EW; CENTER-TBI collaborators. Machine learning algorithms

performed no better than regression models for prognostication in traumatic brain

injury.   J   Clin   Epidemiol.   2020   Jun;122:95-107.   doi:

10.1016/j.jclinepi.2020.03.005

73.   Data-Flair. Advantages and disadvantages of machine learning language [internet].

[cited 2022 Dec 9]. Available from: https://data-flair.training/blogs/advantages-

and-disadvantages-of-machine-learning/.

74.   Villar A. Response Bias. Encyclopedia of Survey Research Methods. Lavrakas

1338   PJ, editor. Thousand Oaks, Sage Publications Ltd. p. 752-753.

1339   75.   Grimes DA, Schulz KF. Bias and causal associations in observational research.

1340   Lancet. 2002 Jan 19;359(9302):248-52. doi: 10.1016/S0140-6736(02)07451-2

1341   76.   Rodger S, Scott EM, Nolan A, Wright AK, Reid J. Effect of age, breed, and sex on

1342   the health-related quality of life of owner assessed healthy dogs. Front Vet Sci.

1343   2021 Feb 5;8:603139. doi: 10.3389/fvets.2021.603139

1344   77.   Lund EM, Armstrong PA, Kirk CA, Klausner JS. Prevalence and risk factors for

1345   obesity in adult dogs from private US veterinary practices. Intern J Appl Res Vet

1346   Med. 2006;4[2]:177-186.

1347   78.   Merlo DF, Rossi L, Pellegrino C, Ceppi M, Cardellino U, Capurro C, et al. Cancer

1348   incidence in pet dogs: findings of the Animal Tumor Registry of Genoa, Italy. J

1349   Vet   Intern   Med.   2008   Jul-Aug;22(4):976-84.   doi: 10.1111/j.1939-

1350   1676.2008.0133.x

1351   79.   German AJ, Holden SL, Wiseman-Orr ML, Reid J, Nolan AM, Biourge V, Morris

1352   PJ, Scott EM. Quality of life is reduced in obese dogs but improves after

1353   successful   weight   loss.   Vet   J.   2012   Jun;192(3):428-34.   doi:

1354   10.1016/j.tvjl.2011.09.015

1355   80.   Niessen SJ, Powney S, Guitian J, Niessen AP, Pion PD, Shaw JA, et al. Evaluation

1356   of a quality-of-life tool for dogs with diabetes mellitus. J Vet Intern Med. 2012

1357   Jul-Aug;26(4):953-61. doi: 10.1111/j.1939-1676.2012.00947.x

1358   81.   Iliopoulou MA, Kitchell BE, Yuzbasiyan-Gurkan V. Development of a survey

1359   instrument to assess health-related quality of life in small animal cancer patients

1360   treated with chemotherapy. J Am Vet Med Assoc. 2013 Jun 15;242(12):1679-87.

1361   doi: 10.2460/javma.242.12.1679

1362   82.   Roberts C, Armson B, Bartram D, Belshaw Z, Capon H, Cherry G, et al.

1363   Construction of a conceptual framework for assessment of health-related quality of

1364   life in dogs with osteoarthritis. Front Vet Sci. 2021 Sep 24;8:741864. doi:

1365   10.3389/fvets.2021.741864

1366   83.   Wallis LJ, Radford AD, German AJ, Jackson J, Belshaw Z, Kubinyi E, et al. Just

1367   old age" - a qualitative investigation of UK owner experiences and attitudes to

1368   ageing of senior and geriatric dogs. Proceedings of the 30[th] International Society

1369   for Anthrozoology Conference; 2021 June 22-24; online. p81.

1370

1371

Supplementary material

S1 File. Statistical report containing details of all data pre-processing steps to create the dataset for primary decision-makers, as well as initial data visualisation using UMAP.

S2 File. Statistical report containing details of all data pre-processing steps to create the dataset for all owners.

S3 File. Statistical report containing details of all statistical analyses for the Kendall's rank correlation, and also XGBoost modelling for the *any health problem* outcome using data from primary decision makers.

S4 File. Statistical report containing details of all statistical analyses for the Kendall's rank correlation, and also XGBoost modelling for the *any health problem* outcome using using data from all owners.

S5 File. Statistical report containing details of XGBoost modelling for the *significant illness* outcome using data from primary decision makers.

S6 File. Statistical report containing details of XGBoost modelling for the *significant illness* outcome using data from all owners.

S7 File. Statistical report containing details of binary logistic regression modelling for the *any health problem* variable using data from primary decision makers.

1397   **S8 File. Statistical report containing details of binary logistic regression modelling for**

1398   **the *any health problem* outcome using data from all owners.**

1399

1400   **S9 File. Statistical report containing details of binary logistic regression modelling for**

1401   **the *significant illness* variable using data from primary decision makers.**

1402

1403   **S10 File. Statistical report containing details of binary logistic regression modelling for**

1404   **the *significant illness* variable using data from all owners.**

1405

1406   **S1 Table.   Results of simple (i.e., univariable) binary logistic regression analyses**

1407   **examining associations between owner, animal and veterinary variables and the *any***

1408   ***health problem binary* for all owners.**

1409

1410   **S2 Table.   Results of simple (i.e., univariable) binary logistic regression analyses**

1411   **examining associations between owner, animal and veterinary variables and the**

1412   ***significant illness* binary for all owners.**

1413

1414   **S3 Table.   Kendall's tau correlation coefficients for the naïve associations amongst**

1415   **owner-animal metadata variables in Fig 2.**

1416

1417   **S4 Table.   Respective log *P*-values for the Kendall's tau correlations amongst owner-**

1418   **animal metadata variables in Fig 2 and S1 Table.**

1419

S5 Table. Final multiple binary logistic regression model examining associations between owner, animal and veterinary variables and the presence of *any health problem* as reported by the owner.

S6 Table. Final multiple binary logistic regression model examining associations between owner, animal and veterinary variables and the presence of a significant or serious illness as reported by the owner.

S1 Fig. Owner-dog metadata visualised by Uniform Manifold Approximation and Projection with Density-Based Spatial Clustering. Owner-pet metadata were pre-processed with all factors as numeric and subject to dimension reduction with the UMAP projection technique. Owner variables included in this visualisation were *diet*, *sex*, *location*, *education* and *income*; animal variables included were *age*, *sex*, *neuter status*, *diet* and *breed size category*. Healthcare variables were not included. Each individual row of the data (owner-dog combination) contributes one UMAP x and y coordinate, analogous to PC1 and PC2 in a principal component analysis. Four or five distinct clusters were evident. Points are colour-coded by *owner gender* (a), *income* (b), *location* (c), *setting* (d), *education* € and *decision-maker status* (f) as indicated in the legend.

S2 Fig. Owner-dog metadata visualised by Uniform Manifold Approximation and Projection with Density-Based Spatial Clustering. Owner-pet metadata were pre-processed with all factors as numeric and subject to dimension reduction with the UMAP projection technique. Owner variables included in this visualisation were *diet*, *sex*, *location*, *education* and *income*; animal variables included were *age*, *sex*, *neuter status*, *diet* and *breed size category*. Healthcare variables were not included. Each individual row of the data

1445 (owner-dog combination) contributes one UMAP x and y coordinate. Four or five distinct

1446 clusters were evident. Points are colour-coded by *owner age* (a), *animal career* (b), *dog age*

1447 (c), *breed size category* (d), *dog sex* € and *neuter status* (f) as indicated in the legend.

1448

S3 Fig. Multiple binary logistic regression model, on data from all owners, with *any*

***health problem* as the outcome variable and including either all owner, animal and**

including either all healthcare variables (a) or only the owner-animal metadata (b).

The dots represent the odds ratio for each variable, whilst the bars represent 99% confidence

intervals (99%-CI). Variables where the 99%-CI range does not include 1.0 (vertical dotted

line) are depicted in red, whilst those that include 1.0 are depicted in blue. Note the

logarithmic scale for the X-axis.

1456

S4 Fig. Multiple binary logistic regression model with *significant illness* as the outcome

variable, on data from owners who were primary carers, either all owner, animal and

including either all healthcare variables (a) or only the owner-animal metadata (b).

The dots represent the odds ratio for each variable, whilst the bars represent 99% confidence

intervals (99%-CI). Variables where the 99%-CI range does not include 1.0 (vertical dotted

line) are depicted in red, whilst those that include 1.0 are depicted in blue. Note the

logarithmic scale for the X-axis.

1464

S5 Fig. **All-variable XGBoost model on the *any health problem* binary using the all-**

owner dataset. (a) Receiver operating characteristic curve of a prediction model containing

all variables (owner, animal and healthcare). This shows the increasing true positive and

false positive rates, with decrease of the threshold probability for prediction of *any health*

problem. Prediction accuracy was good as assessed by ROC analysis (area under curve 0.838,

99%-CI: 0.797-0.879). Acceptance threshold is indicated by the colour bar on the right-hand

side and shown at discrete points on the curve. (b) Graph depicting the relative contribution

of predictor variables to the all-variable XGBoost model for the *any health problem* binary.

'Importance' represents fractional contribution of each feature to the model, based on the

total gain from including each feature; 'cover' represents the number of observations related

to this feature in the model); 'frequency' represents the relative number of times a feature has

been used in trees). Variables are organised in order of importance left to right, based on the

sum of the 3 metrics, whilst variables that are not shown did not contribute to the final model.

1478

S6 Fig. **Reduced (owner-animal metadata) XGBoost model on the *any health problem***

binary using the all-owner dataset. (a) Receiver operating characteristic curve of a reduced

prediction model only containing owner and animal variables. This shows the increasing true

positive and false positive rates, with decrease of the threshold probability for prediction of

any health problem. Prediction accuracy was moderate, as assessed by ROC analysis (AUC

0.682, 99%-CI: 0.628-0.737). Acceptance threshold is indicated by the colour bar on the

right-hand side and shown at discrete points on the curve. The fact that the prediction

thresholds are all low (< 0.5) shows that this model struggles to predict health issues. (b)

Graph depicting the relative contribution of predictor variables to the all-variable XGBoost

model for the *any health problem* binary. 'Importance' represents fractional contribution of

each feature to the model, based on the total gain from including each feature; 'cover'

represents the number of observations related to this feature in the model); 'frequency'

represents the relative number of times a feature has been used in trees). Variables are

organised in order of importance left to right, based on the sum of the 3 metrics, whilst

variables that are not shown did not contribute to the final model.

1494

S7 Fig. **All-variable XGBoost model on the** *significant illness* **binary using the all-owner**

dataset. (a) Receiver operating characteristic curve of a prediction model containing all

variables (owner, animal and healthcare). This shows the increasing true positive and false

positive rates, with decrease of the threshold probability for prediction of *significant illness*.

Prediction accuracy was good, as assessed by ROC analysis (area under curve 0.884, 99%-

CI: 0.796-0.972). Acceptance threshold is indicated by the colour bar on the right-hand side

and shown at discrete points on the curve. (b) Graph depicting the relative contribution of

predictor variables to the all-variable XGBoost model for the *significant illness* binary.

'Importance' represents fractional contribution of each feature to the model, based on the

total gain from including each feature; 'cover' represents the number of observations related

to this feature in the model); 'frequency' represents the relative number of times a feature has

been used in trees). Variables are organised in order of importance left to right, based on the

sum of the 3 metrics, whilst variables that are not shown did not contribute to the final model.

1508

S8 Fig. **Reduced (owner-animal metadata) XGBoost model on the** *significant illness*

binary using the all-owner dataset. (a) Receiver operating characteristic curve of a reduced

prediction model only containing owner and animal variables. This shows the increasing true

positive and false positive rates, with decrease of the threshold probability for prediction of

significant illness. Prediction accuracy was moderate, as assessed by ROC analysis (area

under curve 0.664, 99%-CI: 0.535-0.693). Acceptance threshold is indicated by the colour

bar on the right-hand side, and also shown at discrete points on the curve. (b) Graph

depicting the relative contribution of predictor variables to the all-variable XGBoost model

for the *significant illness* binary. 'Importance' represents fractional contribution of each

feature to the model, based on the total gain from including each feature; 'cover' represents

the number of observations related to this feature in the model); 'frequency' represents the

relative number of times a feature has been used in trees). Variables are organised in order of

importance left to right, based on the sum of the 3 metrics, whilst variables that are not shown

did not contribute to the final model.

1523

Binary logistic regression: any health problem

Odds ratio (log scale)

Model P < 0.001
Pseudo R^2 = 0.420
AUC = 0.842

Binary logistic regression: any health problem

Model P < 0.001
Pseudo R^2 = 0.421
AUC = 0.844

Odds ratio (log scale)