Whole-genome sequencing reveals diverse mechanisms underlying quantitative

pathogenicity and host adaptation in a fungal plant pathogen

Reda AMEZROU*[1], Aurélie DUCASSE[1], Jérôme COMPAIN[2], Nicolas LAPALU[1,2], Anais PITARCH[1],

Laetitia DUPONT[1], Johann CONFAIS[1], Henriette GOYEAU[1], Gert GHJ KEMA[3], Daniel CROLL[4],

Joëlle AMSELEM[2], Andrea SANCHEZ-VALLET[5], Thierry C MARCEL*[1]

[1] Université Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, Thiverval-Grignon, France

[2] UR URGI, INRA, Université Paris-Saclay, Versailles, France

[3] Plant Research International B.V., Wageningen, The Netherlands

[4] Université de Neuchâtel, Department of Ecology and Evolution, Neuchâtel, Switzerland

[5] CBGP, INIA, Campus de Montegancedo UPM, Pozuelo de Alarcón, Madrid, Spain

* Corresponding authors: thierry.marcel@inrae.fr; reddamez@gmail.com

Word count (main text): 4,932

Abstract

Knowledge of genetic determinism and evolutionary dynamics mediating host-pathogen interactions is

essential to manage fungal plant diseases. However, the genetic architecture of fungal pathogenicity

remains poorly understood, and studies often focus on large-effect effector genes triggering strong,

qualitative resistance. It is not clear how this translates to predominately quantitative interactions. Here,

we used the *Zymoseptoria tritici*-wheat model to elucidate the genetic architecture of quantitative

pathogenicity and mechanisms mediating host adaptation. *Z. tritici* is a globally occurring pathogen that

causes severe yield losses on wheat. We performed whole-genome sequencing of 103 isolates and

quantified pathogenicity traits on 12 cultivars harbouring different resistance genes. A multi-host

genome-wide association studies identified 58 candidate genes associated with pathogenicity, of which

nineteen were highly expressed and/or differentially expressed *in planta*. Two of these had large effects

and three were shared among more than one cultivar, suggesting that *Z. tritici* pathogenicity is often

quantitative and host-specific. Analysis of genetic diversity revealed that sequence polymorphism is the

main evolutionary process mediating differences in quantitative pathogenicity, a process that is likely

facilitated by genetic recombination and transposable element dynamics. We found signatures of

positive diversifying selection in ~68% of the candidate genes acting on specific amino acid

substitutions, likely responsible for evasion of host recognition. Finally, we used functional approaches

to confirm the role of an effector-like gene and a methyltransferase in quantitative pathogenicity. This

study highlights the complex genetic architecture of quantitative pathogenicity, extensive diversifying

selection and plausible mechanisms facilitating pathogen adaptation.

Introduction

Fungal diseases cause major damage to crop production and threaten food security worldwide[1]. Understanding the molecular dialogue between pathogens and their hosts is essential to design durable and effective control strategies. Major molecular factors in the ability of fungal pathogens to cause disease are effectors. Those are proteins delivered into the host apoplast or translocated inside cells to manipulate host physiology or suppress its immunity to favour infection[2]. In turn, plants respond to pathogen invasion by activating defence mechanisms based on recognition of pathogen-associated molecular patterns and effectors by cell surface and intracellular receptors. The discovery of resistance genes (*R*) that encode receptors has provided potential means to control fungal diseases through resistance breeding[3]. However, pathogens have repeatedly overcome host resistance by evading recognition or by suppression of host immunity. These rapid adaptations are mainly driven by genetic variation at pathogenicity genes and genome evolution[4]. Gene diversification, deletions or horizontal acquisitions have generated adaptive variants at loci encoding effectors. Transposable elements (TEs), high mutation and recombination rates are thought to contribute considerably to the extensive genome variation in many fungal species[5-8].

In the case of antagonistic host-pathogen interactions, the fate of genetic variation is determined partly by the selection pressure exerted by host *R* proteins. This is especially true when *R* genes and effectors are involved in a gene-for-gene (GFG) relationship[9-12]. Moreover, the diversity of host genotypes in *R*-gene content exerts heterogeneous selection pressure on pathogen populations, giving rise to the co-maintenance of multiple pathogen variants better adapted to specific subsets of hosts[3,13]. Hence, in addition to the identification of effector genes, elucidating the mechanisms contributing to gene diversification provides critical information about their evolvability. This knowledge has tremendous applications for the durability of crop protection strategies[14].

The first fungal effectors to be identified were small secreted proteins (SSP). SSPs trigger strong resistance (*i.e.* encoded by avirulence genes) as a result of their recognition by plant immune receptors following a typical GFG relationship[15]. The increasing availability of fungal genomes over the past decade allowed significant advances in investigating molecular plant–fungal interactions and facilitated effector discovery. However, effector biology has focused almost exclusively on SSPs despite the growing evidence that other genetic factors are also involved in host colonization[16]. For instance, small noncoding RNAs and fungal secondary metabolites play similar effector roles in various plant pathogenic fungi[17]. Other fungal proteins were shown to interact with the host immune system or act to promote pathogenicity; these include for example cell wall–degrading enzymes[18], protease inhibitors[19] or disruptors of the hormone-signalling pathway[20].

Effector discovery also led to the systematic search for host genes conferring resistance[15]. Resistance to many fungal diseases can be divided into two categories: (*i*) qualitative (or race-specific) resistance, which is usually controlled by large-effect genes and is effective against avirulent strains; and (*ii*)

quantitative (or non–race-specific) resistance, usually controlled by genes with small-to-moderate
effects and is thought to be less specific. While cloning plant immune receptors remains challenging,
the discovery of the interacting fungal effectors is considered more straightforward[14]. However, fungal
pathogenicity studies often focus on large-effect genes, whose interaction with genetic factors from the
host trigger a strong immune response, leading to qualitative resistance. It remains unclear how these
findings translate to quantitative models, where the interaction of quantitative components of
pathogenicity (also called aggressiveness in the plant pathology literature) and plant resistance, trigger
a partial/incomplete immune response. By contrast to quantitative resistance in plants, the genetic basis
of quantitative pathogenicity in pathogens and the process of host adaptation have received relatively
little attention. Thus, elucidating the biology of quantitative pathogenicity factors and their evolutionary
dynamics is timely and may require the adoption of complementary experimental approaches. To
address this, we used the *Zymoseptoria tritici* – wheat pathosystem, which is known to be largely
quantitative *in natura*.

Zymoseptoria tritici is an ascomycete filamentous fungus that causes the Septoria tritici blotch (STB)
disease on wheat. *STB* is a major threat to wheat crops in Europe and in temperate regions around the
world[21]. *Z. tritici* is hemibiotrophic and remains exclusively apoplastic through its infection cycle, which
is characterized by an asymptomatic phase of 7 to 14 days, followed by an induction of leaf necrosis[22].
This stage is accompanied by the development of pycnidia containing asexual spores that further spread
the disease[23,24]. To date, three *Z. tritici* pathogenicity genes that encode avirulence factors have been
cloned. These include *AvrStb6* and *AvrStb9* that are recognized by the *R* genes *Stb6* and *Stb9*,
respectively, and lead to a strong qualitative resistance response[12,25,26], and *Avr3D1* which triggers
quantitative resistance on cultivars harbouring *Stb7* or *Stb12*[10]. In addition to the above-mentioned *R*
genes, 18 other race-specific *Stb* resistance genes and numerous quantitative resistance loci have been
genetically mapped in wheat[27], but only *Stb6* and *Stb16q* have been cloned to date[28,29]. An ideal approach
to identify genetic determinants of complex traits is through Genome-Wide Association Studies
(GWAS). GWAS aims to associate sequence polymorphisms to phenotypic variation, and relies on
linkage disequilibrium (LD) –the non-random association of alleles at different loci– to identify causal
variants. The approach has been used extensively in plants but remains at its infancy on fungal
populations[30,31].

The aim of the present work was to characterize the genetic determinants underlying quantitative
pathogenicity in *Z. tritici* and to investigate the evolutionary processes that could facilitate pathogen
adaptation to host resistances. We hypothesized that quantitative pathogenicity relies mostly on gene-
for-gene interactions between host and pathogen, and that the genes involved are not all encoding SSPs
but also genetic factors from the non-secretory pathway. To test our hypotheses, we performed an
extensive genome wide association study combined with transcriptomics and population genomics. We
identified candidate genes putatively mediating quantitative pathogenicity and confirmed the role of two
of these genes using functional approaches. Together, our results generate a comprehensive view on the

complex genetic architecture of quantitative pathogenicity and pathogen adaptation to host resistances.

Beyond this model, our findings may have broad implications for sustainable disease management.

Results

Characterization of the fungal collection. To efficiently perform GWAS and reduce confounding

effects, it is important to select a population that is genetically and phenotypically diverse with little

stratification. Following these guidelines, we focused on a subset of 103 French *Z. tritici* isolates selected

from a larger population for their genetic diversity (data not shown). We sequenced their genomes and

generated a total of 247 Gb of sequence data with an average depth of 61x. After quality assessment, we

obtained a total of 1,463,638 Single Nucleotide Polymorphisms (SNPs) that we further filtered for Minor

Allele Frequency (MAF) and missing rates, resulting in a final set of 718,810 SNPs. Analysis of

functional variant effect revealed that 59,318 SNPs (8.25%) induced missense mutations and 2,493

(0.3%) resulted in a loss of function such as premature stop codons, loss of stop or start codons (Table

1). As expected, principal component analysis (PCA) using full polymorphism data did not show any

genetic structure; the first three PCs explained less than 4% of the total variance (Fig. S1). We further

confirmed this with STUCTURE using a subset of evenly distributed SNPs (1 SNP per 5 kb) and like

the PCA analysis, Bayesian clustering failed to infer sub-populations (Fig. 1A). Similarly, individual

isolates showed very low genetic relatedness (Fig. S2). Genome-wide LD decayed (r^2<0.2) over physical

distance at ~0.5 kb (Fig. 1B). Given our high SNP density (1 SNP every 55 bp) we expected marker-

trait associations to be resolved to the individual gene level. When pathotyped on wheat differential

cultivars, isolates showed a wide virulence spectrum depending on the host cultivar, and strong isolate

× cultivar interactions (Tables S1, S2). Overall, strains isolated from the cultivar 'Premio' were more

aggressive compared to those isolated from the susceptible cultivar 'Apache', even when tested on the

cultivar 'Apache' (Fig. S3).

Table 1 | Functional variants identified in a population of 103 *Z. tritici* isolates

Functional variant effect	Count	Percent
5' UTR	10431	1.45
Gain of start codon	1084	0.15
Loss of start codon	127	0.02
Synonymous coding	162938	22.67
Non-synonymous coding	59318	8.25
Intron	58217	8.10
Gain of stop codon	1109	0.15
Loss of stop codon	173	0.02
3' UTR	8794	1.22

A **B**

Fig. 1 | Population structure and Linkage Disequilibrium (LD) in the *Z. tritici* population. A. Admixture plot
of the fungal population. Each vertical bar represents a single isolate and is colored according to the
membership coefficient (Q_i) to the three sub-population clusters from STRUCTURE. **B.** LD decay over
physical distance. The r^2 values were calculated between pairs of SNPs up to a physical distance of 20 kb
and fitted against the physical distance using a non-linear model. The dotted orange line shows the distance
(0.5 kb) at which LD decays at the $r^2 < 0.2$ level.

A complex genetic architecture and host-specificity underlies pathogenicity. To elucidate the
genetic basis of the observed pathogenicity differences, we investigated two key quantitative phenotypes
(PLACN and PLACP) using mixed linear models for marker-trait association tests, performed
independently on each of the studied cultivars. Collectively, we identified 94 genome-wide significant
SNPs and an additional 53 at the 10% False Discovery Rate (FDR) threshold, defining 58 distinct genes
(Fig. 2, Table S4). The strongest associations were at the *AvrStb6* and *AvrStb9* loci and were detected
in all *Stb6-* and *Stb9-* containing cultivars, respectively. Initially not thought to harbour *Stb9*, PLACN
on cultivar 'Cadenza' was highly associated (*P*=3.57e-13) with polymorphisms in *AvrStb9*, suggesting
that the cultivar possesses the corresponding *R* gene. The presence of *Stb9* in 'Cadenza' was recently
confirmed by its tightly linked genetic markers (data not shown). Interestingly, the single-locus model
failed to detect *AvrStb9* and demonstrates the usefulness of the multi-locus mixed model in detecting
loci with larger effects. However, if multiple SNPs were in strong LD, biases may be created while
estimating the genetic variance. Nevertheless, the multi-locus approach in our study generally
outperformed the traditional single-locus tests (Fig. S4). Candidate pathogenicity loci were identified in
the thirteen *Z. tritici* core chromosomes. Despite recent suggestions of accessory chromosomes being
enriched in putative effectors[32], we could not map any gene residing on accessory chromosomes in this
multi-host GWAS (Fig. 2, Table S4). Only eleven genes were shared among PLACP and PLACN (Fig.
2, Table S4), which was expected, since necrosis coverage and pycnidia production levels were highly
variable (Fig. S3). Hence, considering both traits in association analyses could reveal genetic
determinants influencing both, the conidial production and host susceptibility. The phenotypic variance
explained by the identified loci associated with pathogenicity traits on each cultivar ranged from 9.1%

to 65.7% (Table S4). With the exception of *AvrStb6* and *AvrStb9*, all of the detected genes accounted for small-to-moderate effects (R^2<30%) and similarly, analysis of virulent/avirulent alleles showed quantitative differences (Table S4, Fig. 3B). Further, the number of associated genes varied greatly between cultivars (three candidates for 'Bulgaria-88' and 'Toisondor' *vs* ten for 'Israel-493') and ~76% were specific to a single host cultivar (Table S4, Fig. 3A). This weak overlap observed between genes is a strong indication of host-specificity, in consistence with the different *Stb* genes or *Stb* combinations harboured by the cultivar set used here.

Fig. 2 | Overview of the identified candidate pathogenicity genes from the multi-host GWAS. **A.** Number of identified genes per cultivar. **B.** Gene IDs and their significant associations. Black dots correspond to PLACP (percentage of leaf area covered by pycnidia) and red dots to PLACN (percentage of leaf area covered by necrosis) marker-trait associations. The larger the dot the stronger the association (*P*-value). Radial lines connect dots to the cultivar where it was detected and to the gene ID. **C.** Nucleotide diversity (π) estimates in a 50 kb sliding window with a step size of 25 kb. Genomic positions are displayed on the x-axis and π values are displayed on the y-axis. The y-axis ranges from 0 to 0.045. **D.** Tajima's D estimates in 50 kb

sliding windows with a step size of 25 kb. Genomic positions are displayed on the x-axis. Tajima's D values are displayed on the y-axis. The y-axis ranges from -2.5 to 2. **E.** The number of associations for each gene (PLACN/PLACP per cultivar).

***In planta* gene expression and functional annotation.** To reveal the expression profile of the identified candidate pathogenicity genes using GWAS, we calculated the relative abundance of *Z. tritici* transcripts, using RNAseq data generated during different *Z. tritici* infection time courses[22]. Of the 58 GWAS candidates, ten genes exhibited no expression (FPKM = 0) or very low expression (FPKM < 5) throughout the infection cycle, while another ten genes were found to be highly expressed (FPKM above the 95th percentile of the mean expression value) either at specific time points or throughout the infection cycle (Fig. 3C, Table S5). Interestingly, *AvrStb6*, *Zt_6_00224* and *Zt_3_00467* were among the most highly expressed *Z. tritici* genes at the disease transitional phase (9 dpi; Table S5). Differentially expressed genes *in planta* were identified by examining the ratio of expression between the Czapek-Dox broth (CDB) culture medium and the five time points of wheat infection. In our set of candidate genes, sixteen were differentially expressed with a FDR below 0.10 and a log2-fold change greater than 2. Among them, eleven were up-regulated at specific time points of the infection (Fig. 3C; Table S5). For subsequent investigations, we considered a set of nineteen genes as high-confidence candidates, whose expression profiles supported their potential role in pathogenicity; this included the sixteen differentially expressed genes and three additional ones that were highly expressed during the infection (Fig. 3; Table S5).

Fig. 3 | The quantitative nature and host-specificity of *Z. tritici* pathogenicity. **A.** An upset plot of candidate pathogenicity genes across host cultivars. **B.** Boxplots showing variation in quantitative pathogenicity (PLACP) of the isolates carrying different lead SNP alleles of the candidate pathogenicity genes. **C.** Differential gene expression shown as log2-fold change compared to the culture medium. Red shades indicate up-regulated genes and blue shades indicate down-regulated genes during the infection cycle.

Eleven protein products out of the nineteen retained as best candidates were predicted to be secreted (Table S6). These secreted proteins could be effectors promoting disease or interacting with the cognate *Stb* resistance genes analysed in this study. We also found three proteins with predicted transmembrane helices (THMs), a class that is systematically discarded in effector discovery studies. EffectorP predicted seven as putative effectors, with five (excluding *AvrStb6*) lacking homology with other fungal species and might be considered as *Z. tritici*-specific candidate effectors (Table S6). The identified conserved protein domains included functions such as hydrolase (*e.g.*, Glycosyl hydrolase), oxidoreductase (*e.g.*, Thioredoxin reductase) or transferase activities (*e.g.*, Methyltransferase; Table S6). Using PHIB-Blast, we found that seven candidates shared homology with known pathogenicity factors from other pathogenic fungi and therefore constitute promising candidates mediating quantitative pathogenicity in *Z. tritici* (Table S6).

Factors of genetic diversity in pathogenicity genes. Gain in virulence and/or escape from host recognition is often mediated by polymorphisms in pathogenicity genes. We investigated sequence variation and diversity in the identified genes and detailed results are summarized in Table S7. In particular, we compared patterns of nucleotide diversity in candidate pathogenicity genes with the genome-wide levels. We observed higher genetic diversity in pathogenicity genes compared to other protein-coding genes (median $\pi_{Pathogenicity}$=0.0212 vs. $\pi_{Genome-wide}$=0.0079; $P < 0.01$; Fig. 4A). We found no significant impact of loss of function mutations or presence/absence polymorphisms, except for *Mycgr3G110052*. In addition, we wanted to examine which mechanisms were likely to contribute to the high sequence diversity in our candidates. As TE dynamics have been observed to contribute to fungal genome plasticity and diversity, we calculated the proximity of candidate pathogenicity genes to the nearest TEs. 52.6% of these genes were flanked by or adjacent to TEs (*i.e.* at less than 5 kb distance; Table S7). Comparison with genome-wide genes with regards to TE proximity revealed that candidate pathogenicity genes were significantly closer to TEs ($P < 0.05$; Fig. 4B). Gene diversification is also accelerated by meiotic recombination, which influences both natural population dynamics and selection. We used a combination of algorithms in two independent software (RDP4 and GARD) to detect signatures of recombination. We found signatures of recombination supported by both detection methods in a significant proportion (63.2%) of the putative pathogenicity genes (Table S8). Interestingly, genes with low or no signatures of recombination were in close proximity to TEs, whereas those with higher recombination events were found at greater distances from TEs (Fig. 4C). We hypothesize that recombination in particular and proximity to TEs promotes allelic diversity in genes that are putatively involved in quantitative pathogenicity.

Fig. 4 | Genetic diversity of quantitative pathogenicity genes in *Z. tritici* and factors contributing to gene diversification. **A.** Distribution of nucleotide diversity (π) among candidate pathogenicity genes and the rest of the protein-coding genes. Statistical significance was determined using a permutation test with 1000 iterations; *P*<0.01**. **B.** TE distance compared between pathogenicity genes and the rest of the protein-coding genes in *Z. tritici*. Statistical significance was determined using a permutation test with 1000 iterations; *P*<0.05*. **C.** the relationship between TE distance and recombination events in candidate pathogenicity genes.

Pathogenicity genes are hotspots of positive diversifying selection. We assumed that candidate pathogenicity genes may have evolved under *R*-genes selective pressure on pathogen populations to escape recognition. Therefore, we searched for episodes of positive diversifying selection on protein-coding sequences by using the ω (d_N/d_S) parameter. Codon sites under positive selection are defined as those with elevated nonsynonymous/synonymous substitution ratios (ω > 1) compared to the expectation under neutral evolution, ω = 1. The proportion of positively-selected sites ($\omega > 1$) varied greatly among genes and ranged from 0.42% in the case of *Zt_1_01987* to 10.97% for *AvrStb6* (Table S7). We also found four genes with a d_N/d_S > 1 (*Zt_3_00467*, ω=7.03; *AvrStb6*, ω=2.84; *Mycgr3G106276*, ω=1.18 and *AvrStb9*, ω=1.04), that could be considered as genes under strong selection pressure. Since ω averaged across all gene-tree branches rarely surpasses 1, and as the selective pressure could affect codon sites differently, we tested different codeML models to detect positively-selected codons. For all model comparisons, we found signatures of positive diversifying selection in 68.4% of candidate genes using both, the more conservative M1a vs. M2a and the M7 vs. M8 model

pairs (Table S7). Together, this shows that pathogenicity genes are hotspots of positive diversifying
selection, likely as a result of a selection pressure exerted by host *R* genes.

Two functionally distinct genes mediate quantitative pathogenicity. To determine the role of the
candidate genes in quantitative pathogenicity, we selected two distinct genes for functional validation:
Zt_3_00467 (Fig. 5A), which encodes a SSP that is highly expressed *in planta*, with a peak at the disease
transition phase (Fig. 5B); and *Zt_6_00682* (Fig. 6A), which encodes a S-adenosylmethionine-
dependent methyltransferase (SAM-MTase) that is differentially expressed during the disease
asymptomatic phase (Fig. 6B). For *Zt_3_00467*, we ectopically expressed the avirulent (IPO10273) and
virulent (3D7) alleles in the 3D7 strain background, and assessed quantitative pathogenicity traits on the
host cultivars 'Arina' and 'Riband'. Overall, Mutants expressing the avirulent allele (*3D7 +*
Zt_3_00467_{IPO10273}) showed a slight reduction of PLACP levels compared to those expressing the
virulent allele (*3D7 + Zt_3_00467_{3D7}*; Fig. 5C). When compared individually, one mutant expressing
the avirulent allele showed a phenotype similar to the wild-type, while another expressing the virulent
allele had higher PLACP compared to the wild-type (Fig. S7). The difference in phenotypic responses
between ectopic mutants is a well-known phenomenon that may be caused by distinct allele integration
sites in the different mutant lines. Testing transformants expressing the avirulent allele on the cultivar
'Riband' led to lower development of pycnidia compared to the virulent and the wild-type strains (Fig.
S8). 'Arina' and 'Riband' share a common resistance gene known as *Stb15*. These phenotypic
observations suggest that *Zt_3_00467* recognition leads to quantitative resistance which could be
attributed to *Stb15*. Comparison of the avirulent and virulent protein sequences showed seven amino
acid substitutions, all positively selected (Fig. 5D). Analysis of *Zt_3_00467* in the global *Z. tritici*
collection showed that the gene was conserved and that most protein variants were population specific
and segregating by geographical origin (Fig. S9).

Fig. 5 | An effector gene encoding a positively-selected small secreted protein is involved in quantitative pathogenicity of *Z. tritici*. **A.** Regional association plot of a candidate effector protein detected on cultivar ''Arina''. X and Y axes indicate positions on chromosome 6 and −log10 (*P*-value) for associations with PLACP, respectively. Annotations on the reference genome are shown at the bottom with the candidate gene represented by a green box, other genes by blue boxes and transposable elements by grey boxes. A linkage disequilibrium plot is shown below the annotation bar. **B.** log2-fold expression changes of *Zt_3_00467* based on RNAseq data collected throughout the time course of *Z. tritici* infection on wheat. **C.** percentage of leaf area covered by pycnidia (PLACP) produced by the wild-type strain 3D7 and the ectopic transformants expressing the virulent (3D7) and avirulent (IPO10273) alleles. Statistical significance was determined using a Tukey's HSD test. **D.** Alignment of the virulent and avirulent protein sequences. Residues in yellow constitute the predicted signal peptide, grey residues show protein mutations and red asterisks show positively selected sites with a Bayes Empirical Bayes posterior probability > 0.95.

For *Zt_6_00682*, we generated knockout (KO) mutants of the 3D7 strain by targeted gene disruption and ectopic transformants with the virulent (3D7) and avirulent (3D1) alleles in the KO background. This allowed us to assess allele-dependent effects of *Zt_6_00682*. Three independent KO and ectopic transformants for each allele were tested on the cultivar 'Shafir', where the association was detected. Compared to the wild-type, KO transformants resulted in unexpectedly high PLACP values ranging from 67% to 100% (Fig. 6C). Ectopic transformation of the 3D1 allele in the KO background (3D7Δ*Zt_6_00682*+ *Zt_6_00682₃D₁*) resulted in reduced levels of PLACP, while ectopic lines

transformed with the 3D7 allele (3D7Δ*Zt_ 6_00682*+ *Zt_ 6_00682₃ᴅ₇*) showed a slightly higher PLACP than the wild-type (Fig. 6C). Besides 'Shafir', the deletion mutants were aggressive on another four cultivars, but phenotypic differences between ectopic transformants and wild-type were not significant (Fig. S6). Five amino acid differences were found between the avirulent and virulent protein sequences, of which three were under positive selection (Fig. 6D). Similar to *Zt_3_00467, Zt_6_00682* was conserved in the global population but was highly polymorphic (Fig. S9).

Fig. 6 | A gene encoding a methyltransferase mediates quantitative pathogenicity in *Z. tritici*. **A.** Regional association plot of the pathogenicity gene detected on cultivar ''Shafir''. **B.** log2-fold expression changes of *Zt_6_00682* based on RNAseq transcription data collected throughout the time course of *Z. tritici* infection on wheat. **C.** Percentage of leaf area covered by pycnidia (PLACP) produced by the wild-type strain 3D7, knockout mutants and the complementation mutants carrying the virulent (3D7) and avirulent (3D1) alleles. Statistical significance was determined using a Tukey's HSD test. **D.** Alignment of the virulent and avirulent protein sequence. Residues in orange constitute the methyltransferase domain, grey residues show protein mutations and red asterisks show positively selected sites with a Bayes Empirical Bayes posterior probability > 0.95.

Discussion

Our results point to a complex genetic architecture of pathogenicity in *Z. tritici*, involving many genes with variable additive effects, resulting in quantitative phenotypes on the host cultivar. These findings are consistent with previous studies[33,34] and provide further evidence that pathogenicity in this pathosystem is complex and results from the contribution of both large- and small- effect genes. We highlighted nineteen differentially and highly expressed genes *in planta* and focused on their features for subsequent investigations. The vast majority of these genes had a peak expression in the phases corresponding to the colonization and the pre-pycnidia formation, before the appearance of symptoms. During colonization, hyphae penetrate into host tissue *via* stomata and colonize the substomatal cavity, but remain exclusively in the intercellular space. This is followed by hyphal invasion of the apoplast leading to pre-pycnidia formation in substomatal cavities[23]. It was shown that these are the key phases of effector gene deployment by plant pathogens[35]. These generally encode small, secreted, cysteine-rich proteins with no sequence homology in other species and play a vital role in successful pathogen invasion, suppression of host immune perception and manipulation of its physiology[36]. Here, we mapped a set of diverse secreted proteins. While effector-like proteins may initiate an effector-triggered immune response, as was shown for the gene *Zt_3_00467*, which triggered quantitative resistance in cultivars 'Riband' and 'Arina', secreted molecules with a known functional domain could also promote disease.

Of the functionally annotated secreted proteins, we identified domains associated with macromolecule degradation, fungal metabolism or stress response, among others. Some genes encoded proteins that show strong homology with functionally characterized virulence factors from other pathogenic fungi. For instance, *Zt_1_01674* encodes a Thioredoxin reductase (*Trr*), a flavoenzyme that plays a central role in the thioredoxin oxidative stress resistance pathway and is involved in regulating gene transcription, cell growth and apoptosis[37]. Deletion of the *Trr* ortholog resulted in reduced hyphal growth, sexual reproduction and virulence in *Fusarium graminearum*[38] and profoundly affected pathogenicity of the entomopathogenic fungus *Beauveria bassiana*[39]. We hypothesize that *Zt_1_01674* may play similar roles and could be an important factor in *Z. tritici* development and pathogenicity. We also identified a cell-wall degrading enzyme (*Zt_2_00163*) belonging to the Glycoside Hydrolase protein family. Glycoside hydrolases are known to play important roles in fungal invasion and nutrition acquisition via the degradation or modification of plant cell wall components, such as cellulose or hemicellulose[40].

Genes encoding proteins predicted to be non-secreted were common in our candidate list, highlighting not only the polygenic nature of pathogenicity, but also the potential role of the non-secretory pathway in fungal pathogenicity. This was demonstrated by the functional studies of *Zt_6_00682*, which encodes a SAM-dependent Methyltransferase (SAM-MTase). SAM-MTases catalyse the transfer of methyl groups from SAM to a large variety of substrates, ranging from metabolites to macromolecules, including proteins[41]. They are involved in epigenetic regulation with functions ranging from the control of gene expression, silencing or activation of transposon control[42]. *Zt_6_00682* shares some sequence

similarities with the putative SAM-MTase gene *LaeA*, a component of the velvet transcription factor complex that functions as a global regulator of fungal secondary metabolism and development for a handful of fungi[43]. Recent work showed that mutants lacking the velvet gene *velB* in *Z. tritici* have defects in vegetative growth and asexual reproduction[44], while the loss of *LaeA* attenuates pathogenicity in a variety of filamentous fungi[45]. However, the inactivation of *Zt_6_00682* promoted pycnidia formation, suggesting that it functions differently from *LaeA* and may negatively regulate the transcriptional activity of genes or metabolites associated with host colonization. However, how this SAM-MTase affects the regulation of pathogenicity factors needs to be investigated as a key step towards understanding the evolutionary mechanisms associated with the emergence of highly aggressive strains.

In addition, we identified three genes with predicted transmembrane domains. For instance, the *DPM2* ortholog (*Zt_1_01325*) was highly expressed throughout the infection cycle, highlighting its potential role in *Z. tritici* pathogenicity. *DPM2* is a regulatory subunit of the dolichol-phosphate mannose synthase complex, key enzymes for GPI anchor biosynthesis[46]. The alteration of the *DPM* complex affected cell wall integrity, fungal growth and increased chitin levels in *Candida albicans*[46]. Traditionally, transmembrane proteins are systematically discarded in effector gene prediction studies. There is however a growing evidence that they could play key roles in pathogenicity of filamentous fungi and our results support this hypothesis[47,48].

We found substantial evidence for host specificity demonstrated by a highly significant isolate × cultivar interaction and few shared associations between host cultivars. It is also probable that the weak overlap in candidate pathogenicity genes among host cultivars is partly due to a lack of statistical power to detect variants of small effect[49]. However, besides *Stb6*- and *Stb9*-carrying cultivars, there was no meaningful correlation among phenotypes on different hosts, which further supports host specificity. In several well studied host-pathogen systems, specificity is determined mainly by the matching of resistance or susceptibility genes and their corresponding effectors[50-52], following a GFG relationship (or inverse GFG in the case of necrotrophic fungal pathogens).

In *Z. tritici*, pathogenicity can be both qualitative[26,53,54] and quantitative[10,55]. Here, we showed that pathogenicity is predominantly quantitative based on multiple genes of small-to-moderate effects. This was demonstrated with the allele analysis of candidate genes and functional studies of *Zt_3_00467* and *Zt_6_00682*. In contrast to what has been shown for most GFG interactions, the recognition of the *Zt_3_00467* and *Zt_6_00682* avirulent alleles does not trigger strong resistance, but rather a reduction at the onset of disease symptoms. Furthermore, the fact that avirulent alleles are only recognized by certain wheat cultivars suggests that the interaction also follows a GFG model and demonstrates that *Zt_3_00467* and *Zt_6_00682* are likely quantitative avirulence factors interacting with specific host resistance genes. This is further supported by recent work in *Z. tritici* as shown for *Avr3D1* and a recently reported effector gene (*G_07189*) that triggers quantitative *Stb20q* resistance[11,56]. This is not uncommon

in other filamentous fungi. For instance, the late effector LmSTEE98 from *Leptosphaeria maculans*

triggers quantitative resistance following a GFG model[57], similarly, silencing three *Blumeria graminis*

f sp. *tritici* effectors lead to quantitative gain of resistance[58]. Since *Zt_3_00467* encodes a typical

effector protein and is recognized by hosts harbouring the *Stb15* resistance gene, we suggest that

Zt_3_00467 could be *AvrStb15* and trigger an effector-induced defence. We hypothesise that *Stb15*

resistance is not strong enough to block pathogen infection, although we cannot completely rule out the

existence of a *Zt_3_00467* haplotype that triggers qualitative resistance. However, additional genetic

factors might also contribute to differences in pathogenicity between avirulent and virulent strains. On

the other hand, *Zt_6_00682* encodes an atypical avirulence factor and may interact indirectly with the

host immune system. The fact that *Zt_6_00682* resembles genes involved in transcriptional regulation

and that its deletion mutants are more aggressive on some cultivars, suggests that *Zt_6_00682* negatively

regulates pathogenicity factors that are specifically recognized by one or more (unknown) host

quantitative resistance gene(s). Complementation mutants further support host specificity, where

recognition of the avirulent allele leads to a dramatic reduction in pycnidia formation. Unlike deletion

mutants, this was only observed on the cultivar where the gene was detected initially ('Shafir').

However, the resistance gene involved and the underlying mechanisms through which the avirulent

allele is recognized remain unknown.

In addition to *AvrStb6* and *AvrStb9*, we found the SSP *Zt_2_01230* to be significantly associated with

variation in quantitative pathogenicity on two cultivars, suggesting a common yet unknown *Stb*

resistance gene. We also identified a number of secreted and species specific proteins. Since canonical

fungal effectors are generally thought to be secreted, species-specific and without conserved protein

domain[59], we propose that the ones identified here constitute a set of candidate effectors interacting with

their cognate host *Stb* resistance products, and should be a priority target for functional studies.

Unlocking the molecular basis underpinning these interactions and how it could impact host defence

would lead to a better exploitation of quantitative resistance, which is thought to be more durable[60].

Evolution and diversification of pathogenicity genes enables adaptation to new host cultivars and, in

response to host population changes, they undergo changes in rapid allele frequency[61]. We observed

high levels of genetic diversity in our candidate pathogenicity genes when compared to the genome-

wide gene content. We found a single case of gene deletion and no significant impact of loss of function.

Because of the intact and functional coding sequences for nearly all of the identified genes, we suggest

that sequence polymorphism is the primary mechanism behind the rapid adaptation of *Z. tritici* to host

populations, permitting the pathogen to circumvent host resistance while conserving the presumable

function of the pathogenicity protein. We next investigated what could contribute to this extensive

genetic diversity. Recombination is a central source of diversity in sexual fungal pathogens[62]. Here, we

detected recombination events for a significant number of candidates, which suggests a role in the

observed genetic diversity of pathogenicity genes. In *Z. tritici*, it was reported that recombination is a

major driver of its evolution and is highly heterogeneous, concentrated in narrow genomic regions that are enriched in putative effector genes[8,63]. TEs are also thought to contribute to gene diversification by creating fast-evolving genomic regions through non-homologous recombination or repeat-induced point mutations[64]. Here, we found that candidate pathogenicity genes are indeed at closer distances to TEs, suggesting their potential role in sequence diversity. We also observed that TE distance was correlated with recombination events in putative pathogenicity genes, and that the levels of genetic diversity were high for genes with the highest recombination events, regardless of their distances to TEs. We hypothesize that recombination is likely the major driver underlying the diversification of pathogenicity genes. A recent study supports this hypothesis where, unlike distance to TEs, the recombination rate was strongly associated with adaptive substitutions in *Z. tritici* genomes[4].

A key determinant of genetic diversity at pathogenicity genes is selection pressure. Understanding the evolution of pathogenicity in fungal populations with respect to the selection pressure imposed by host populations is necessary for the sustainable management of *R* genes in the field. This can be achieved by searching for signatures of positive diversifying selection at the risk of recovering some false positive signals. In this study, we used conservative maximum likelihood models and models that have been shown to be robust to recombination and significantly reduce false positives[65]. More than two thirds of candidate genes harbour signatures of positive selection acting on specific codons, suggesting that the diversifying selection is likely the primary evolutionary force maintaining non-synonymous substitutions relevant for *Z. tritici*-wheat quantitative interactions. For instance, the vast majority of different mutations between the avirulent and virulent versions of *Zt_3_00467* and *Zt_6_00682* have been identified as positively selected. Such selection pressure could be the result of host *R* gene recognition, and may have greatly contributed to the accumulation of beneficial mutations. This agrees with a previous study, which demonstrated that rapidly evolving genes play a prominent role in host-specific disease development[66]. Together, these results suggest that host–pathogen coevolution drives the emergence and maintenance of allelic diversity at loci likely involved in direct (or indirect) interactions with host *R* genes. Future surveys of the diversity and structure of *R* genes in wheat populations will provide a deeper insight into reciprocal patterns of selection, coevolution, and the maintenance of diversity in resistance and pathogenicity in natural populations.

Methods

Fungal material. We analysed 103 *Z. tritici* isolates in this study deriving predominantly from a large collection of French isolates collected in 2009 and 2010. This genetically diverse sample was mainly isolated from the French cultivars 'Apache' (n=44) and 'Premio' (n=42). The remaining were isolated from cultivars 'Soissons' (n=9), 'Caphorn' (n=3), 'Alixan' (n=1), 'Bermude' (n=1), 'Dinosor' (n=1), and 'Garcia' (n=1). One isolate was collected in the United Kingdom in 2009 on cultivar 'Humbert'. All strains are monospore non-clonal isolates were originally isolated either at the Plant Research International in the Netherlands or at INRAE UR BIOGER in France. For the functional validation of

candidate pathogenicity genes, we used the two strains 3D1 (*i.e.*, ST99CH3D1) and 3D7 (*i.e.*, ST99CH3D7) collected in 1999 from a Swiss wheat field and previously used for the functional validation of *Avr3D1*[10].

Plant material. The pathogenicity of the 103 isolates was initially evaluated on 29 wheat cultivars, including 15 carrying known *Stb* resistance genes. After the first replicate, only cultivars on which we observed phenotypic variability were kept for second and third replicates. These were the cultivars 'Bulgaria-88' (*Stb1* and *Stb6*), 'Israel-493' (*Stb3* and *Stb6*), 'Tadinia' (*Stb4* and *Stb6*), 'Shafir' and 'Cadenza' (*Stb6*), 'Estanzuela Federal' (*Stb7*), 'Courtot' (*Stb9*), 'Apache' (*Stb4/5* and *Stb11*), 'Salamouni' (*Stb13*, *Stb14* and *StbSm3*) and 'Arina' (*Stb6* and *Stb15*); the resistance background of cultivars 'Premio' and 'Toisondor' was not known. Details on the differential lines are available in Table S3. This genetic material is available from the INRAE Small grain cereals Biological Resources Centre in Clermont-Ferrand (<https://www6.clermont.inrae.fr/umr1095_eng/Organisation>).

Quantitative pathogenicity assays. For inoculum preparation, blastospores of *Z. tritici* stored at -80°C were used to inoculate a liquid growth medium (Difco™ YPD Broth Cat. No. 242820) agitated at 140 rpm in a 17°C climate cell for 144 h and transferred to agar growth medium (Difco™ YPD Agar Cat. No. 242720) for 96 h. Fresh blastospores were recovered with loops, resuspended in sterile water, and the inoculum was adjusted to a concentration of 10^6 spores.L^{-1}, to which one drop of Tween®20 (Sigma Aldrich® Cat. No. 9005-64-5) was added per 15 mL before inoculation. Five seeds from each cultivar were sown in 90*90*95 mm pots containing Floradur B soil (80% blond peat / 20% black peat, with 1.2 kg.m^{-3} fertilizer 18/10/20). One day before inoculation, we kept three homogeneous seedlings on which a 75-mm-length portion was delimited on the first true leaf using a black marker. Sixteen days old seedlings were inoculated by applying the inoculum six times on the marked leaves with a flat tipped paintbrush (synthetic bristles, 12 mm width), depositing approximately 30,000 spores.cm^{-2} on the leaf surface. After inoculation, plants were watered and enclosed in transparent polyethylene high-density bags for 72 h to initiate the infection. Ten days post-inoculation, leaves were removed keeping only the marked leaf to ensure light homogeneity and longevity of the inoculated leaf. From sowing to disease scoring, plants were kept in growth chambers regulated at 80% relative humidity, 16 h of light per 24 h at a photon flux of 300 µmol.m^{-2}.s^{-1}, 22°C temperature under light and 18°C in the darkness. Quantitative pathogenicity was assessed as percentages of leaf area covered by pycnidia (PLACP) and by necrosis (PLACN) at 21 days post inoculation, averaged over three leaves and three replicates (nine leaves per interaction). Phenotypic characterization of the generated deletion and ectopic transformants was done following the same procedure described above in two replicates.

Whole-genome resequencing. Blastospores were harvested after seven days growth in liquid cultures on yeast extract-peptone-dextrose Broth (Difco™ YPD Broth), and then lyophilised for 24 h. High

502 molecular weight DNA was extracted using a phenol-chloroform-based procedure[67]. Illumina paired-
503 end libraries were prepared for each sample and DNA fragments were re-sequenced with 2x100 bp reads
504 on a HiSeq-2000 sequencing system at Genewiz, Inc. (formerly Beckman Coulter Genomics). Depth of
505 sequence coverage was variable depending on the isolate, ranging from 16 to 303 genome equivalents
506 with an average of 61 genome equivalents.
507
SNP calling and quality filtering. For each isolate, reads were trimmed using trimmomatic v0.32[68] and
then mapped to the reference genome IPO323[69] using the mem algorithm from BWA v0.7.7 with default
settings[70]. Samtools v0.1.19 and Picard tools v1.106 were used to filter reads[71]
(<http://broadinstitute.github.io/picard/>); to remove optical duplicates, secondary alignments, reads
with a mapping quality below 30 and to keep only pairs in which both reads met quality checks. SNP
calling was done with Freebayes v0.9 using the option --report-monomorphic[72] and all positions
detected as low complexity regions or TEs were excluded. Low-complexity regions and TEs were
detected using RepeatMasker with default settings (RepeatMasker Open-4.0
<http://www.repeatmasker.org>) and REPET[73], respectively. Quality filters were applied to VCF files
using in-house scripts. Depth of coverage (DP) calculated at each position followed a normal distribution
around the depth main peak. Hence, a DP filter was set at twice the standard deviation of the mean. All
VCFs were stored as a matrix reduced to keep only positions where at least one isolate carries an
alternative allele at a non-filtered position. The SNP matrix was parsed as a .tped format, converted to
.ped/.map. The final matrix was filtered for less than 50% missing data and more than 10% Minor Allele
Frequency (MAF) using PLINK v.1.9[74] <https://cog-genomics.org/plink/>.

Linkage disequilibrium (LD). To assess the mapping power and resolution of GWAS, we estimated
genome-wide LD decay using PopLDdecay[75]. The average squared Pearson's correlation coefficient (r^2)
was computed between pairs of SNPs up to 20 kb, and plotted over physical distance by fitting a
nonlinear regression model.

Population structure analyses. Two approaches were used to investigate the fungal population
structure: (1) we performed a principal component analysis (PCA) using independent SNPs ($r^2 < 0.2$)
with the --pca command PLINK v.1.9 and (2) the admixture model in STRUCTURE v.2.2 software[76]
with a 10^5 burn-in period, 5×10^5 MCMC steps and 5 iterations for each k value from 1 to 10. The optimal
number of sub-populations was determined with the Evanno method[77].

GWAS. For each quantitative trait, we conducted GWAS using single- and multi-locus linear mixed
models based on the EMMA algorithm implemented in GAPIT v.3.0[78-80]. We used a VanRaden derived
kinship matrix as a random effect, and set the PCA option to 'true' to determine the optimum number
of PCs to be included in the model, following the Bayesian Information Criterion (BIC). The most robust
model was chosen by visualizing Quantile-Quantile plots that illustrate the observed vs expected P
values. A uniform threshold of $P=1/n$ (n being the effective number of independent SNPs) was used as

the genome-wide significance threshold. The effective number of independent SNPs was calculated using the Genetic type 1 Error Calculator; <http://grass.cgs.hku.hk/gec/download.php>. The suggestive genome-wide threshold for the 718,810 SNPs was 3.96×10^{-6} and the significant P value was 1.98×10^{-7}. We also adjusted SNP P-values to account for multiple testing correction using the false discovery rate (FDR) procedure. SNPs with an FDR-adjusted P value < 0.10 were deemed significant. For LD analysis, we first calculated the pairwise LD with the lead SNP in a window size of 1 Mb using the --r^2 command in PLINK v.1.9. We set a cut-off of $r^2 > 0.5$ to determine candidate genomic regions and assigned predicted genes using a custom Perl script. In case more than one gene is assigned to the candidate region, we picked the closest one from the lead SNP having pathogenicity-like properties (*e.g.*, secretion signal, differential expression *in planta*). If no gene fell in the LD region, we selected the closest gene to the lead SNP. LD heatmaps were constructed using Haploview v4.2[81].

Functional annotation. We inferred gene models from the *ab initio Z. tritici* annotation[82] in the candidate genomic regions identified by GWAS. We used RNAseq datasets of the reference isolate IPO323[22] to check for missing genes or manually refine gene models (*e.g.*, splice junctions, missing exons). N-terminal secretion signals were predicted using the Hidden Markov Model (HMM) scoring method implemented in SignalP 3.0[83]. Transmembrane domains were identified with TMHMM 2.0[84]. Effector probabilities were predicted through EffectorP 3.0[85]. Conserved protein domains were assigned using HMM scan against the PFAM 35.0 database and blastp was used to search for homologs from other fungi. Finally, we used PHIB-BLAST, a blast server for PHI-base (Pathogen-Host Interactions), to search for homologs against functionally characterized pathogenicity factors.

Gene expression analysis. RNAseq data from *in vitro* and *in planta* (1-, 4-, 9-, 14- and 21-days post infection; dpi) were obtained from a previous study[22]Cliquez ou appuyez ici pour entrer du texte.. We computed gene expression quantified as fragments per kilobase of exon per million fragments mapped (FPKM) using the improved gene models with Cuffdiff v.2.2.1[86]. We inferred pairwise log2-fold expression changes and P value adjustment for *in planta* RNAseq samples compared to the Czapek-Dox broth (CDB) culture medium. Genes with an FDR-adjusted P-value < 0.10 and a log2|fold change|> 2 were considered as differentially expressed during the infection.

***De novo* assembly and sequence diversity.** We used SPAdes v.3.14.1 to produce draft assemblies for each isolate[87]. We ran the tool with the following settings: -k 21,33,55,75,95 --careful to reduce mismatches and indel errors. We performed blastn[88] searches using ~1000 bp of the target locus as a query to retrieve matching contig segments in each assembly. High-confidence matches were first extracted from the contig set using samtools v1.10 and subsequently aligned with MAFFT v7.464 using the --auto option and 1,000 iterative refinement cycles[89]. Alignments were inspected using JalView[90] and ambiguous regions were removed with trimAL[91] with less stringent parameters. We used the batch

mode of DNAsp v.6[92] to calculate summary statistics of population genetic parameters associated with

candidate genes such as nucleotide diversities (π)[93] and Tajima's D[94]. TE locations for *Z. tritici* were

retrieved from a previous study[82] and we parsed gene distances to TEs using the command 'closest'

from bedtools[95].

Evolutionary analysis. Before testing for positive diversifying selection, we checked for evidence of

recombination, as this can cause false positives. We used the software packages RDP4[96] and GARD[97]

for recombination detection. We considered genes with recombination events if both detection methods

found evidence of recombination. We then constructed non-recombinant phylogenetic trees of gene

coding sequences using the RAxML algorithm[98] implemented in RDP4. Statistical support for the nodes

was obtained after 100 bootstrap replicates. We used codeml of the PAML package[99] to estimate the

evolutionary rates of candidate genes, based on Markov models of codon evolution. Branch lengths of

the inferred ML trees were estimated by using the model M0 and then fixed in further analyses. The

model M0 assumes constant selective pressure across codon sites and over time. The selective pressure

at the coding-sequence level was measured by ω, the ratio of non-synonymous to synonymous rates

(d_N/d_S). A d_N/d_S ratio of 1 ($\omega = 1$) indicates neutrality, whereas $\omega < 1$ and $\omega > 1$ suggests purifying and

diversifying selection, respectively. Models M1a, M2a, M7 and M8 of variable selective pressure across

codon sites were used to estimate selective pressure and test for positive selection[100,101]. For each

candidate gene, we performed pairwise likelihood ratio tests (LRTs) for positive selection, comparing

models that allow sites with $\omega > 1$ (M2a and M8) with models that do not (M1a and M7). We considered

genes under positive diversifying selection if both model comparisons in LRT were significant. Finally,

positively selected sites were inferred with a Bayes Empirical Bayes (BEB) probability[101] and codons

with $\omega > 1$ at a probability > 0.95 were considered positively-selected.

Functional validation of candidate pathogenicity genes. For *Zt_6_00682*, we generated knockout

mutants in the 3D7 isolate and ectopic transformants expressing the virulent and avirulent alleles in the

knock-out background. To create plasmid constructs for gene disruption, flanking regions (~1 kb before

start and stop codons) of *Zt_6_00682* were amplified from gDNA using primers listed in Table S9. The

hygromycin resistance gene cassette was amplified from the plasmid pNOV2114_HygR_attP1_attP2

and cloned into a pNOV2214 vector. The vector was then digested with *Hind*III restriction enzyme (New

England Biolabs), amplified and used for the cloning of PCR fragments with a Gibson Assembly cloning

kit (New England Biolabs) following the manufacturer's instructions. Constructs were transformed into

Escherichia coli NEB 5-α using a heat shock transformation, screened by PCR and verified by Sanger

sequencing using primers listed in Table S9. Confirmed plasmids were transformed into *Agrobacterium*

tumefaciens cells by heat shock and validated by PCR. The vector was used to transform *Z. tritici* via *A.*

tumefaciens-mediated transformation (ATMT) following a procedure adapted from Zwiers & de

Waard[102]. Transformed *Z. tritici* colonies growing in the presence of hygromycin were tested by PCR to

confirm the successful deletion of the gene. To validate the allele-dependent effect of *Zt_6_00682*, we generated and validated ectopic transformants by placing both the virulent (3D7) and the avirulent (3D1) alleles into the 3D7 KO background, following the same procedure described above with the following modifications: (*i*) we used the vector pNOV_3Gate_SUL with sulfonylurea as a resistance marker and the digestion was achieved with *Hind*III and *Kpn*I restriction enzymes. The resulting vectors, pNOV_sulf_3D7g.7232 and pNOV_sulf_3D1g.7155, were used to transform *Z. tritici* via ATMT following the procedure described above. Primers used to verify gene disruptions and ectopic integrations are listed in Table S9. Finally, three independent transformants carrying the different virulence alleles and three knock-out mutants were used for phenotypic characterization *in planta*. For *Zt_3_00467*, we ectopically expressed the virulent (3D7) and avirulent (IPO10273) alleles in the 3D7 background. A fragment containing the *Zt_3_00467* gene, including 1800 bp upstream of the start codon and 463 bp downstream of the stop codon was amplified using Phusion DNA polymerase (NEB) and the primers included in table S9. This fragment was cloned into a pCGEN vector[103] previously digested with *Kpn*I (New England Biolabs), using the In-Fusion HD Cloning Kit (Takara Bio) and following the manufacturer's instructions. Vectors were transformed using heat-shocked *E. coli* Stellar cells and verified using Sanger sequencing. Transformation via ATMT was performed as described above and selection for the *Z. tritici* mutants was performed on geneticin (150 μg.mL^{-1})-containing plates.

Data availability

Sequence data of the 103 French isolates are deposited at the NCBI Sequence Read Archive under the accession number PRJNA777581.

Acknowledgements

We thank David Gouache (ARVALIS Institut du Végétal, Boigneville, France), Marc-Henri Lebrun, Henriette Goyeau, Frédéric Suffert and Anne-Sophie Walker (INRAE UR BIOGER, Palaiseau, France) for their help in establishing the fungal collection; Robert King and Jason Rudd (Rothamsted Research, Harpenden, United Kingdom) for providing RNAseq data; Gabriel Scalliet (Syngenta Crop Protection AG, Stein, Switzerland) for providing the vectors pNOV2114, pNOV_3Gate_SUL and pNOV_3Gate_HYG used to clone the candidate pathogenicity genes; Martin Willigsecker and Béatrice Beauzoone (INRAE UR BIOGER) for their help in the pathology assays; and Thierry Rouxel (INRAE UR BIOGER) for critical reading of the manuscript and valuable feedback.

Financial Disclosure

This work was supported by the French National Research Agency (program BIOADAPT, ANR-12-ADAP-0009-04), the FSOV 2008 B project and BASF France SAS for R.A. postdoctoral fellowship. T.C.M. was supported by a Short-Term Scientific Mission granted by the COST SUSTAIN action FA1208. A.S.V. was supported by the Ministerio de Ciencia e Innovación (Ramón y Cajal program,

grant no. RYC2018-025530-I). INRAE BIOGER benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007).

Author contributions

T.C.M and R.A. conceived and designed the study. G.H.J.K. and H.G. isolated and provided most of the natural *Z. tritici* isolates constituting the GWAS panel. A.D., T.C.M. and J.C. carried out pathogenicity assays on the natural isolates, and R.A. on the transformant isolates. A.D. and T.C.M. prepared the DNA samples for whole-genome resequencing. J.C., N.L. and J.A. performed the bioinformatics analyses, including preparation of the SNP matrix and *de novo* genome assemblies. D.C. provided sequence data of the global *Z. tritici* collection. R.A., L.D., A.P. and A.S.V. performed the cloning and genetic transformations. R.A. performed the genome-wide association, gene expression and evolutionary sequence analyses. R.A. and T.C.M. wrote the manuscript. All authors have reviewed the manuscript.

Competing Interests

The authors declare they have no competing interests.

References

1. Savary, S., et al. The global burden of pathogens and pests on major food crops. *Nat. Ecol. Evol.* **3**, 430–439 (2019). https://doi.org/10.1038/s41559-018-0793-y

2. Franceschetti, M., et al. Effectors of filamentous plant pathogens: commonalities amid diversity. *Microbiol. Mol. Biol. Rev.* **81**, e00066-16 (2022). https://doi.org/10.1128/MMBR.00066-16

3. lo Presti, L., et al. Fungal effectors and plant susceptibility. *Annu. Rev. Plant Biol.* **66**, 513–545 (2015). https://doi.org/10.1146/annurev-arplant-043014-114623

4. Grandaubert, J., Dutheil, J. Y., & Stukenbrock, E. H. The genomic determinants of adaptive evolution in a fungal pathogen. *Evol. Lett.* **3**, 299–312 (2019). https://doi.org/10.1002/evl3.117

5. Möller, M., & Stukenbrock, E. H. Evolution and genome architecture in fungal plant pathogens. *Nat. Rev. Microbiol.* **15**, 756–771 (2017). https://doi.org/10.1038/nrmicro.2017.76

6. Raffaele, S., & Kamoun, S. Genome evolution in filamentous plant pathogens: why bigger can be better. *Nat. Rev. Microbiol.* **10**, 417–430 (2012). https://doi.org/10.1038/nrmicro2790

7. Rouxel, T., et al. Effector diversification within compartments of the *Leptosphaeria maculans* genome affected by repeat-induced point mutations. *Nat. Commun.* **2**, 202 (2011). https://doi.org/10.1038/ncomms1189

8. Stukenbrock, E. H., & Dutheil, J. Y. Fine-scale recombination maps of fungal plant pathogens reveal dynamic recombination landscapes and intragenic hotspots. *Genetics* **208**, 1209–1229 (2018). https://doi.org/10.1534/genetics.117.300502

9. Singh, P.K., et al. Co-evolutionary interactions between host resistance and pathogen avirulence genes in rice-*Magnaporthe oryzae* pathosystem, *Fungal Genet. Biol.* **115**, 9-19 (2018). https://doi.org/10.1016/j.fgb.2018.04.005

10. Meile, L., et al. A fungal avirulence factor encoded in a highly plastic genomic region triggers partial resistance to septoria tritici blotch. *New Phytol.* **219**, 1048–1061 (2018). https://doi.org/10.1111/nph.15180

11. Langlands-Perry, C., et al. Quantitative and qualitative plant-pathogen interactions call upon similar
pathogenicity genes with a spectrum of effects. *Front. Plant Sci.* **14**, 1128546 (2023).
https://doi.org/10.3389/fpls.2023.1128546

12. Amezrou, R., et al. A secreted protease-like protein in *Zymoseptoria tritici* is responsible for
avirulence on *Stb9* resistance gene in wheat. *PLoS Pathog.* **19**, e1011376 (2023).
https://doi.org/10.1371/journal.ppat.1011376

13. Stukenbrock, E. H., & McDonald, B. A. Population genetics of fungal and oomycete effectors
involved in gene-for-gene interactions. *Mol. Plant-Microbe Interact.* **22**, 371–380 (2009).
https://doi.org/10.1094/MPMI-22-4-0371

14. van de Wouw, A. P., & Idnurm, A. Biotechnological potential of engineering pathogen effector
proteins for use in plant disease management. *Biotechnol. Adv.* **37**, 107387 (2019).
https://doi.org/10.1016/j.biotechadv.2019.04.009

15. Domazakis, E., et al. Effectoromics-based identification of cell surface receptors in potato. *Methods
mol. biol.* **1578**, 337–353 (2017). https://doi.org/10.1007/978-1-4939-6859-6_29

16. Varden, F. A., de la Concepcion, J. C., Maidment, J. H. R., & Banfield, M. J. Taking the stage:
effectors in the spotlight. *Curr. Opin. Plant Biol.* **38**, 25–33 (2017).
https://doi.org/10.1016/j.pbi.2017.04.013

17. Collemare, J., O'Connell, R., & Lebrun, M.-H. Nonproteinaceous effectors: the terra incognita of
plant–fungal interactions. *New Phytol.* **223**, 590–596 (2019). https://doi.org/10.1111/nph.15785

18. Kubicek, C. P., Starr, T. L., & Glass, N. L. Plant cell wall–degrading enzymes and their secretion
in plant-pathogenic fungi. *Annu. Rev. Phytopathol.* **52**, 427–451 (2014).
https://doi.org/10.1146/annurev-phyto-102313-045831

19. van der Hoorn, R. A. L. Plant proteases: from phenotypes to molecular mechanisms. *Annu. Rev.
Plant Biol.* **59**, 191–223 (2008). https://doi.org/10.1146/annurev.arplant.59.032607.092835

20. Ma, K.-W., & Ma, W. Phytohormone pathways as targets of pathogens to facilitate infection. *Plant
Molecular Biology* **91**, 713–725 (2016). https://doi.org/10.1007/s11103-016-0452-0

21. O'Driscoll, A., Kildea, S., Doohan, F., Spink, J., & Mullins, E. The wheat Septoria conflict: a new
front opening up? *Trends Plant Sci.* **19**, 602–610 (2014).
https://doi.org/10.1016/j.tplants.2014.04.011

22. Rudd, J. J., et al. Transcriptome and metabolite profiling of the infection cycle of *Zymoseptoria
tritici* on wheat reveals a biphasic interaction with plant immunity involving differential pathogen
chromosomal contributions and a variation on the hemibiobiotrophic lifestyle definition. *Plant
Physiol.* **167**, 1158–1185 (2015). https://doi.org/10.1104/pp.114.255927

23. Duncan, K. E., & Howard, R. J. Cytological analysis of wheat infection by the leaf blotch pathogen
Mycosphaerella graminicola. *Mycol. Res.* **104**, 1074–1082 (2000).
https://doi.org/10.1017/S0953756299002294

24. Kema, G. H. J., Sayoud, R., Annone, J. G., & van Silfhout, C. H. Genetic variation for virulence
and resistance in the wheat-*Mycosphaerella graminicola* pathosystem II. Analysis of interactions
between pathogen isolates and host cultivars. *Phytopathology* **86**, 213–220 (1996).
https://doi.org/10.1094/Phyto-86-213

25. Kema, G. H. J., et al. Stress and sexual reproduction affect the dynamics of the wheat pathogen
effector AvrStb6 and strobilurin resistance. *Nat. Genet.* **50**, 375–380 (2018).
https://doi.org/10.1038/s41588-018-0052-9

26. Zhong, Z., et al. A small secreted protein in *Zymoseptoria tritici* is responsible for avirulence on wheat cultivars carrying the *Stb6* resistance gene. *New Phytol.* **214**, 619–631 (2017). https://doi.org/10.1111/nph.14434

27. Brown, J. K. M., Chartrain, L., Lasserre-Zuber, P., & Saintenac, C. Genetics of resistance to *Zymoseptoria tritici* and applications to wheat breeding. *Fungal Genet. Biol.* **79**, 33–41 (2015). https://doi.org/10.1016/j.fgb.2015.04.017

28. Saintenac, C., et al. Wheat receptor-kinase-like protein Stb6 controls gene-for-gene resistance to fungal pathogen *Zymoseptoria tritici*. *Nat. Genet.* **50**, 368–374 (2018). https://doi.org/10.1038/s41588-018-0051-x

29. Saintenac, C., et al. A wheat cysteine-rich receptor-like kinase confers broad-spectrum resistance against Septoria tritici blotch. *Nat. Commun.* **12**, 433 (2021). https://doi.org/10.1038/s41467-020-20685-0

30. Bartoli, C., & Roux, F. Genome-wide association studies in plant pathosystems: toward an ecological genomics approach. *Front. Plant Sci.* **8**, 763 (2017). https://www.frontiersin.org/article/10.3389/fpls.2017.00763

31. Sánchez-Vallet, A., Hartmann, F. E., Marcel, T. C., & Croll, D. Nature's genetic screens: using genome-wide association studies for effector discovery. *Mol. Plant Pathol.* **19**, 3–6 (2018). https://doi.org/10.1111/mpp.12592

32. Habig, M., Quade, J., & Stukenbrock, E. H. Forward genetics approach reveals host genotype-dependent importance of accessory chromosomes in the fungal wheat pathogen *Zymoseptoria tritici*. *MBio* **8**, e01919–17 (2017). https://doi.org/10.1128/mBio.01919-17

33. Hartmann, F. E., Sánchez-Vallet, A., McDonald, B. A., & Croll, D. A fungal wheat pathogen evolved host specialization by extensive chromosomal rearrangements. *ISME J.* **11**, 1189–1204 (2017). https://doi.org/10.1038/ismej.2016.196

34. Stewart, E. L., et al. Quantitative trait locus mapping reveals complex genetic architecture of quantitative virulence in the wheat pathogen *Zymoseptoria tritici*. *Mol. Plant Pathol.* **19**, 201–216 (2018). https://doi.org/10.1111/mpp.12515

35. Gohari, A.M., et al. Effector discovery in the fungal wheat pathogen *Zymoseptoria tritici*. *Mol. Plant Pathol.* **16**, 931–945 (2015). https://doi.org/10.1111/mpp.12251

36. Toruño, T. Y., Stergiopoulos, I., & Coaker, G. Plant-pathogen effectors: cellular probes interfering with plant defenses in spatial and temporal manners. *Annu. Rev. Phytopathol.* **54**, 419–441 (2016). https://doi.org/10.1146/annurev-phyto-080615-100204

37. Arnér, E. S. J., & Holmgren, A. Physiological functions of thioredoxin and thioredoxin reductase. *Eur. J. Biochem.* **267**, 6102–6109 (2000). https://doi.org/10.1046/j.1432-1327.2000.01701.x

38. Fan, X., et al. Thioredoxin reductase is involved in development and pathogenicity in *Fusarium graminearum*. *Front. Microbiol.* **10**, 393 (2019). https://www.frontiersin.org/article/10.3389/fmicb.2019.00393

39. Zhang, L.-B., Tang, L., Ying, S.-H., & Feng, M.-G. Distinct roles of two cytoplasmic thioredoxin reductases (Trr1/2) in the redox system involving cysteine synthesis and host infection of *Beauveria bassiana*. *Appl. Microbiol. Biotechnol.* **100**, 10363–10374 (2016). https://doi.org/10.1007/s00253-016-7688-0

40. Esquerré-Tugayé, M.-T., Boudart, G., & Dumas, B. Cell wall degrading enzymes, inhibitory proteins, and oligosaccharides participate in the molecular dialogue between plants and pathogens. *Plant Physiology and Biochemistry*, *38*(1), 157–163 (2000). https://doi.org/10.1016/S0981-9428(00)00161-3

41. Zhang, J., & Zheng, Y. G. SAM/SAH analogs as versatile tools for SAM-dependent methyltransferases. *ACS Chem. Biol.* **11**, 583–597 (2016). https://doi.org/10.1021/acschembio.5b00812

42. Martienssen, R. A., & Vincent, C. DNA methylation and epigenetic inheritance in plants and filamentous fungi. *Science* **293**, 1070–1074 (2001). https://doi.org/10.1126/science.293.5532.1070

43. Bayram, Ö., & Braus, G. H. Coordination of secondary metabolism and development in fungi: the velvet family of regulatory proteins. *FEMS Microbiol. Rev.* **36**, 1–24 (2012). https://doi.org/10.1111/j.1574-6976.2011.00285.x

44. Tiley, A.M.M., White, H.J., Foster, G.D., & Bailey, A.M. The *ZtvelB* gene is required for vegetative growth and sporulation in the wheat pathogen *Zymoseptoria tritici*. *Front. Microbiol.* **10,** 2210 (2019). https://doi.org/10.3389/fmicb.2019.02210

45. Jain, S., & Keller, N. Insights to fungal biology through LaeA sleuthing. *Fungal Biol. Rev.* **27**, 51–59 (2013). https://doi.org/10.1016/j.fbr.2013.05.004

46. Juchimiuk, M., Kruszewska, J., & Palamarczyk, G. Dolichol phosphate mannose synthase from the pathogenic yeast *Candida albicans* is a multimeric enzyme. *Biochim. Biophys. Acta Gen. Subj.* **1850**, 2265–2275 (2015). https://doi.org/10.1016/j.bbagen.2015.08.012

47. Weiland, P., & Altegoer, F. Identification and characterization of two transmembrane proteins required for virulence of *Ustilago maydis*. *Front. Plant Sci.* **12**, 669835 (2021). https://doi.org/10.3389/fpls.2021.669835

48. Gu, Q., Chen, Y., Liu, Y., Zhang, C., & Ma, Z. The transmembrane protein FgSho1 regulates fungal development and pathogenicity via the MAPK module Ste50-Ste11-Ste7 in *Fusarium graminearum*. *New Phytol.* **206**, 315–328 (2015). https://doi.org/10.1111/nph.13158

49. Korte, A., & Farlow, A. The advantages and limitations of trait analysis with GWAS: a review. *Plant Methods* **9**, 29 (2013). https://doi.org/10.1186/1746-4811-9-29

50. Bourras, S., et al. Multiple avirulence loci and allele-specific effector recognition control the *Pm3* race-specific resistance of wheat to powdery mildew. *The Plant Cell* **27**, 2991–3012 (2015). https://doi.org/10.1105/tpc.15.00171

51. Faris, J. D., et al. A unique wheat disease resistance-like gene governs effector-triggered susceptibility to necrotrophic pathogens. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 13544–13549 (2010). https://doi.org/10.1073/pnas.1004090107

52. Liu, W., et al. Recent progress in understanding PAMP- and effector-triggered immunity against the rice blast fungus *Magnaporthe oryzae*. *Mol. Plant* **6**, 605–620 (2013). https://doi.org/10.1093/mp/sst015

53. Brading, P. A., Verstappen, E. C. P., Kema, G. H. J., & Brown, J. K. M. A gene-for-gene relationship between wheat and *Mycosphaerella graminicola*, the Septoria tritici blotch pathogen. *Phytopathology* **92**, 439–445 (2012). https://doi.org/10.1094/PHYTO.2002.92.4.439

54. Kema, G. H. J., Verstappen, E. C. P., & Waalwijk, C. Avirulence in the wheat Septoria tritici leaf blotch fungus *Mycosphaerella graminicola* is controlled by a single locus. *Mol. Plant-Microbe Interact.* **13**, 1375–1379 (2000). https://doi.org/10.1094/MPMI.2000.13.12.1375

55. Zhan, F., et al. Linear correlation analysis of *Zymoseptoria tritici* aggressiveness with *in vitro* growth rate. *Phytopathology* **106**, 1255–1261 (2016). https://doi.org/10.1094/PHYTO-12-15-0338-R

56. Meile L, et al. Natural variation in Avr3D1 from *Zymoseptoria* sp. contributes to quantitative gene-for-gene resistance and to host specificity. *New Phytol.* **238**, 1562-1577 (2023). https://doi.org/10.1111/nph.18690

57. Jiquel, A., et al. A gene-for-gene interaction involving a 'late' effector contributes to quantitative resistance to the stem canker disease in *Brassica napus*. *New Phytol.* **231**, 1510–1524 (2021). https://doi.org/10.1111/nph.17292

58. Schaefer, L.K., et al. Cross-kingdom RNAi of pathogen effectors leads to quantitative adult plant resistance in wheat. *Front. Plant Sci.* **11**, 253 (2020). https://doi.org/10.3389/fpls.2020.00253

59. Kim, K.-T., et al. Kingdom-wide analysis of fungal small secreted proteins (SSPs) reveals their potential role in host association. *Front. Plant Sci.* **7**, 186 (2016). https://www.frontiersin.org/article/10.3389/fpls.2016.00186

60. Kettles, G. J., & Kanyuka, K. Dissecting the molecular interactions between wheat and the fungal pathogen *Zymoseptoria tritici*. *Front. Plant Sci.* **7**, 508 (2016). https://www.frontiersin.org/article/10.3389/fpls.2016.00508

61. Barrett, L. G., et al. Diversity and evolution of effector loci in natural populations of the plant pathogen *Melampsora lini*. *Mol. Biol. Evol.* **26**, 2499–2513 (2009). https://doi.org/10.1093/molbev/msp166

62. Burdon, J. J., & Silk, J. Sources and patterns of diversity in plant-pathogenic fungi. *Phytopathology* **87**, 664–669 (1997). https://doi.org/10.1094/PHYTO.1997.87.7.664

63. Croll, D., Lendenmann, M. H., Stewart, E., & McDonald, B. A. The impact of recombination hotspots on genome evolution of a fungal plant pathogen. *Genetics* **201**, 1213–1228 (2015). https://doi.org/10.1534/genetics.115.180968

64. Seidl, M. F., & Thomma, B. P. H. J. Transposable elements direct the coevolution between plants and microbes. *Trends Genet.* **33**, 842–851 (2017). https://doi.org/10.1016/j.tig.2017.07.003

65. Anisimova, M., Nielsen, R., & Yang, Z.. Effect of recombination on the accuracy of the likelihood method for detecting positive selection at amino acid sites. *Genetics* **164**, 1229–1236 (2003). https://doi.org/10.1093/genetics/164.3.1229

66. Poppe, S., Dorsheimer, L., Happel, P., & Stukenbrock, E. H. Rapidly evolving genes are key players in host specialization and virulence of the fungal wheat pathogen *Zymoseptoria tritici* (*Mycosphaerella graminicola*). *PLoS Pathog.* **11**, e1005055 (2015). https://doi.org/10.1371/journal.ppat.1005055

67. Fagundes, W. C., Haueisen, J., & Stukenbrock, E. H. Dissecting the biology of the fungal wheat pathogen *Zymoseptoria tritici*: A laboratory workflow. *Curr. Protoc. Microbiol.* **59**, e128 (2020). https://doi:10.1002/cpmc.128

68. Bolger, A. M., Lohse, M., & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014). https://doi.org/10.1093/bioinformatics/btu170

69. Goodwin, S. B., et al. Finished genome of the fungal wheat pathogen *Mycosphaerella graminicola* reveals dispensome structure, chromosome plasticity, and stealth pathogenesis. *PLoS Genet.* **7**, e1002070 (2011). https://doi.org/10.1371/journal.pgen.1002070

70. Li, H., & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009). https://doi.org/10.1093/bioinformatics/btp324

71. Li, H., et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009). https://doi.org/10.1093/bioinformatics/btp352

72. Garrison, E., & Marth, G. Haplotype-based variant detection from short-read sequencing. *arXiv*, 1207.3907 (2012). http://arxiv.org/abs/1207.3907

73. Flutre, T., Duprat, E., Feuillet, C., & Quesneville, H. Considering transposable element diversification in *de novo* annotation approaches. *PLoS ONE* **6**, e16526 (2011). https://doi.org/10.1371/journal.pone.0016526

74. Chang, C. C., et al. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* **4** (2015). https://doi.org/10.1186/s13742-015-0047-8

75. Zhang, C., Dong, S.-S., Xu, J.-Y., He, W.-M., & Yang, T.-L. PopLDdecay: a fast and effective tool for linkage disequilibrium decay analysis based on variant call format files. *Bioinformatics* **35**, 1786–1788 (2019). https://doi.org/10.1093/bioinformatics/bty875

76. Pritchard, J. K., Stephens, M., & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics* **155**, 945–959 (2000). https://doi.org/10.1093/genetics/155.2.945

77. Evanno, G., Regnaut, S., & Gouget, J. Detecting the number of clusters of individuals using the software structure: a simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005). https://doi.org/10.1111/j.1365-294X.2005.02553.x

78. Kang, H. M., et al. Efficient control of population structure in model organism association mapping. *Genetics* **178**, 1709–1723 (2008). https://doi.org/10.1534/genetics.107.080101

79. Lipka, A. E., et al. GAPIT: genome association and prediction integrated tool. *Bioinformatics* **28**, 2397–2399 (2012). https://doi.org/10.1093/bioinformatics/bts444

80. Segura, V., et al. An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations. *Nat. Genet.* **44**, 825–830 (2012). https://doi.org/10.1038/ng.2314

81. Barrett, J. C., Fry, B., Maller, J., & Daly, M. J. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* **21**, 263–265 (2005). https://doi.org/10.1093/bioinformatics/bth457

82. Grandaubert, J., Bhattacharyya, A., & Stukenbrock, E. H. RNA-seq based gene annotation and comparative genomics of four fungal grass pathogens in the genus Zymoseptoria identify novel orphan genes and species-specific invasions of transposable elements. *G3-Genes Genom. Genet.* **5**, 1323-1333 (2015). https://doi.org/10.1534/g3.115.017731

83. Dyrløv Bendtsen, J., Nielsen, H., von Heijne, G., & Brunak, S. Improved prediction of signal peptides: SignalP 3.0. *J. Mol. Biol.* **340**, 783–795 (2004). https://doi.org/10.1016/j.jmb.2004.05.028

84. Krogh, A., Larsson, B., von Heijne, G., & Sonnhammer, E. L. L. Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J. Mol. Biol.* **305**, 567–580 (2001). https://doi.org/10.1006/jmbi.2000.4315

85. Sperschneider, J., & Dodds, P. N. EffectorP 3.0: prediction of apoplastic and cytoplasmic effectors in fungi and oomycetes. *Mol. Plant-Microbe Interact.* **35**, 146–156 (2021). https://doi.org/10.1094/MPMI-08-21-0201-R

86. Trapnell, C., et al. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat. Protoc.* **7**, 562–578 (2012). https://doi.org/10.1038/nprot.2012.016

87. Bankevich, A., et al. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012). https://doi.org/10.1089/cmb.2012.0021

88. Camacho, C., et al. BLAST+: architecture and applications. *BMC Bioinform.* **10**, 421 (2009). https://doi.org/10.1186/1471-2105-10-421

89. Katoh, K., & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013). https://doi.org/10.1093/molbev/mst010

90. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M., & Barton, G. J. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009). https://doi.org/10.1093/bioinformatics/btp033

91. Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009). https://doi.org/10.1093/bioinformatics/btp348

92. Rozas, J., et al. DnaSP 6: DNA Sequence polymorphism analysis of large data sets. *Mol. Biol. Evol.*
34, 3299–3302 (2017). https://doi.org/10.1093/molbev/msx248

93. Nei, M. *Molecular Evolutionary Genetics* (Columbia University Press, New York, 1987).
https://doi.org/10.7312/nei-92038

94. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism.
Genetics **123**, 585–595 (1989). https://doi.org/10.1093/genetics/123.3.585

95. Quinlan, A. R., & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010). https://doi.org/10.1093/bioinformatics/btq033

96. Martin, D. P., Murrell, B., Golden, M., Khoosal, A., & Muhire, B. RDP4: Detection and analysis of
recombination patterns in virus genomes. *Virus Evol.* **1** (2015). https://doi.org/10.1093/ve/vev003

97. Kosakovsky Pond, S. L., Posada, D., Gravenor, M. B., Woelk, C. H., & Frost, S. D. W. GARD: a
genetic algorithm for recombination detection. *Bioinformatics* **22**, 3096–3098 (2006).
https://doi.org/10.1093/bioinformatics/btl474

98. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312–1313 (2014). https://doi.org/10.1093/bioinformatics/btu033

99. Yang, Z., & Bielawski, J. P. Statistical methods for detecting molecular adaptation. *Trends Ecol.*
Evol. **15**, 496–503 (2000). https://doi.org/10.1016/S0169-5347(00)01994-7

100.   Nielsen, R., & Yang, Z. Likelihood models for detecting positively selected amino acid sites
and applications to the HIV-1 envelope gene. *Genetics* **148**, 929–936 (1998).
http://www.genetics.org/content/148/3/929.abstract

101.   Yang, Z., Wong, W. S. W., & Nielsen, R. Bayes empirical bayes inference of amino acid sites
under positive selection. *Mol. Biol. Evol.* **22**, 1107–1118 (2005).
https://doi.org/10.1093/molbev/msi097

102.   Zwiers, L.-H., & de Waard, M. A. Efficient Agrobacterium tumefaciens-mediated gene
disruption in the phytopathogen *Mycosphaerella graminicola*. *Curr. Genet.* **39**, 388–393 (2001).
https://doi.org/10.1007/s002940100216

103.   Motteram, J., et al. Aberrant protein N-glycosylation impacts upon infection-related growth
transitions of the haploid plant-pathogenic fungus *Mycosphaerella graminicola*. *Mol. Microbiol.* **81**,
415–433 (2011). https://doi.org/10.1111/j.1365-2958.2011.07701.x