Individual Astrocyte Morphology in the Collagenous Lamina Cribrosa Revealed by Multicolor DiOlistic Labeling

Susannah Waxman[1], Marissa Quinn[1], Cara Donahue[2], Louis D. Falo Jr.[2,3],
Daniel Sun[4], Tatjana C. Jakobs[4], Ian A. Sigal[1,3] *

1) Department of Ophthalmology, University of Pittsburgh, Pittsburgh PA, USA;

2) Department of Dermatology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA;

3) Department of Bioengineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA, USA;

4) Department of Ophthalmology, Massachusetts Eye and Ear, Harvard Medical School, Boston, MA;

Short title: MuDi labeling individual collagenous LC astrocytes

* **Correspondence:**

Ian A. Sigal, Ph.D.

Laboratory of Ocular Biomechanics

Department of Ophthalmology, University of Pittsburgh School of Medicine

203 Lothrop Street, Eye and Ear Institute, Rm. 930, Pittsburgh, PA 15213

Phone: (412) 864-2220; Fax: (412) 647-5880

Email: ian@OcularBiomechanics.com

www.OcularBiomechanics.com

Keywords: astrocyte, lamina cribrosa, microscopy, morphology, optic nerve head

Disclosures: Nothing to disclose.

Funding: Supported in part by National Institutes of Health grants R01-EY023966, R01-EY031708, R01-HD045590, R01-HD083383, 1S10RR028478-01, P30-EY008098, and T32-EY017271; Eye and Ear Foundation (Pittsburgh, PA); Research to Prevent Blindness (unrestricted grant to UPMC Ophthalmology and Stein Innovation Award to Sigal IA).

Acknowledgements: Dr. Martin Oberbarnscheidt, for generously sharing vibratome equipment and training in its proper use

<u>Abstract</u>

Astrocytes in the lamina region of the optic nerve head play vital roles in supporting retinal
ganglion cell axon health. In glaucoma, these astrocytes are implicated as early responders to
stressors, undergoing characteristic changes in cell function as well as cell morphology. Much of
what is currently known about individual lamina astrocyte morphology has been learned from
rodent models which lack a defining feature of the human optic nerve head, the collagenous
lamina cribrosa (LC). Current methods available for evaluation of collagenous LC astrocyte
morphology have significant shortcomings. We aimed to evaluate Multicolor DiOlistic labeling
(MuDi) as an approach to reveal individual astrocyte morphologies across the collagenous LC.

Gold microcarriers were coated with all combinations of three fluorescent cell membrane dyes,
DiI, DiD, and DiO, for a total of seven dye combinations. Microcarriers were delivered to 150µm-
thick coronal vibratome slices through the LC of pig, sheep, goat, and monkey eyes via MuDi.
Labeled tissues were imaged with confocal and second harmonic generation microscopy to
visualize dyed cells and LC collagenous beams, respectively. GFAP labeling of DiOlistically-
labeled cells with astrocyte morphologies was used to investigate cell identity. 3D models of
astrocytes were created from confocal image stacks for quantification of morphological features.

DiOlistic labeling revealed fine details of LC astrocyte morphologies including somas, primary
branches, higher-order branches, and end-feet. Labeled cells with astrocyte morphologies were
GFAP$^+$. Astrocytes were visible across seven distinct color channels, allowing high labeling
density while still distinguishing individual cells from their neighbors. MuDi was capable of
revealing tens to hundreds of collagenous LC astrocytes, in situ, with a single application. 3D
astrocyte models allowed automated quantification of morphological features including branch
number, length, thickness, hierarchy, and straightness as well as Sholl analysis.

MuDi labeling provides an opportunity to investigate morphologies of collagenous LC astrocytes,
providing both qualitative and quantitative detail, in healthy tissues. This approach may open
doors for research of glaucoma, where astrocyte morphological alterations are thought to coincide
with key functional changes related to disease progression.

1. Introduction

1.1. **Background**

The lamina cribrosa (LC) is an early site of retinal ganglion cell (RGC) injury in glaucoma, a leading cause of irreversible blindness worldwide. The LC is densely populated by astrocytes which, in health, play vital roles in maintaining RGC homeostasis and healthy vision(J. Salazar et al., 2019; Prada et al., 2016). Growing evidence supports an early role of lamina astrocyte functional and morphological changes as factors that can be protective against and can contribute to glaucomatous neurodegeneration.(Hernandez et al., 2008; Liu et al., 2022; Morgan, 2000; Shinozaki and Koizumi, 2021; Sun et al., 2017) Changes in astrocyte function are known to impact several key processes involved in optic nerve head (ONH) health, such as hemodynamic regulation,(Attwell et al., 2010; Dong et al., 2022) blood-brain barrier maintenance,(Heithoff et al., 2021; Liu et al., 2018) and extracellular matrix remodeling.(Hernandez, 2000; Schneider and Fuchshofer, 2016) These astrocyte functional changes are associated with changes in cell morphology. Astrocytes undergo extensive reorganization in the transition between health and glaucoma. Characteristic "reactive" lamina astrocytes exhibit thickened and shortened cell processes, loss of higher-order branches, and reduced spatial coverage of individual astrocytes(Lye-Barthel et al., 2013; Sun et al., 2010; Sun and Jakobs, 2012; Wang et al., 2017). The morphology and arrangement of astrocytes across the lamina can provide important insights into their roles in health and in disease progression.

1.2. **Why is a new method to label astrocytes needed?**

Established methods to analyze LC astrocyte morphology have important limitations. Despite the recognized importance of LC astrocytes in health and glaucoma, the methods used to investigate their morphology have critical constraints that inhibit effective large-scale analysis of individual cells. Mouse lines expressing fluorescent reporters under the control of astrocyte-specific promoters(Lye-Barthel et al., 2013; Nguyen et al., 2017; Wang et al., 2017) allow for high-quality visualization of these cells. However, mice lack a true collagenous LC like that found in human eyes. Collagen beams and neural tissue pores of the collagenous LC define crucial biological and mechanical properties of the tissue(May and Lütjen-Drecoll, 2002; Midgett et al., 2020; Oikawa et al., 2021; Sigal and Ethier, 2009) that the analogous structure in mice, the glial lamina, does not approximate. To visualize astrocytes of the collagenous LC, the research model must have a true collagenous LC.

A common method used to visualize and identify astrocytes in LC tissue sections is

immunohistochemistry of the GFAP protein. GFAP expression of astrocytes in the brain tends to
be low in health and is upregulated in reactivity.(Liddelow and Barres, 2017) In the glial lamina,
GFAP is strongly expressed by astrocytes, even in the absence of damage or reactive
gliosis.(Lye-Barthel et al., 2013; Sun et al., 2009) Immunohistochemical labeling of GFAP reveals
the location of the GFAP[+] intermediate filaments found in most astrocytes. However, when using
methods that label all or most astrocytes in the LC, it is often not possible to distinguish one cell
from its neighbor due to the high density of astrocytes in the region and their complex and
branched organizations. Complexity is further compounded by the fact that GFAP labeling does
not reveal the full extent of astrocyte morphology, as has been shown in the brain, retina, and
mouse glial lamina.(Bushong et al., 2002; Escartin et al., 2021; Luna et al., 2016; Stokum et al.,
2018; Sun and Jakobs, 2012) GFAP can be visualized in many thicker primary processes, but
often does not label finer higher-order processes, missing important morphological detail.
Astrocyte GFAP content and organization changes substantially in glaucoma,(Chaudhary et al.,
2022; Ling et al., 2020) but reveals only a fraction of astrocyte morphology. Serial block-face
scanning electron microscopy can produce high enough resolution images for users to reconstruct
the entirety of individual astrocytes and separate them from their neighbors with excellent
detail.(Bonney et al., 2022; Calì et al., 2019) However, understanding the diversity of astrocyte
morphologies and spatial relationships to other structures across the LC of several individuals
with this approach would be prohibitively time- and resource-intensive.
85
In lieu of large animal astrocyte reporter lines, astrocyte gene transfer approaches(Ge et
al., 2020; Merienne et al., 2013) could be attempted in the collagenous LC of research animals
such as non-human primates, *in vivo*. This would require considerable time, cost, and risk. The
LC can be difficult to target in live animals both due to selectivity of the blood-brain barrier and
the poorly accessible location of the structure in the eye. Single-cell dye injections can bypass
many challenges of gene transfer approaches and similarly reveal astrocyte morphology. This
approach has been successfully implemented in astrocytes of the brain(Bushong et al., 2002;
Moye et al., 2019) and the mouse glial lamina(Butt et al., 1994; Sun, 2018) specifically. However,
as this technique requires cannulating cells with a micropipette one at a time, it can be time and
labor-intensive to visualize multiple cells across the tissue at once.
96
1.3. **How does Multicolor DiOlistic labeling fit in as a logical next step?**
DiOlistic labeling is a technique that utilizes a gene gun to propel gold or tungsten
microcarriers coated in cell membrane dyes into individual cells within tissues. With this

approach, tens to hundreds of dye-coated microcarriers can be delivered to tissues
simultaneously.   This technique has been implemented in the brain and retina to visualize neuron
morphologies(O'Brien and Lummis, 2006; Santina and Ou, 2018; Seabold et al., 2010; Staffend
and Meisel, 2011) and in the mouse glial lamina(Jakobs, 2014; Sun et al., 2010) to visualize
astrocyte morphologies. By delivering microcarriers coated in all combinations of 3 spectrally
distinct fluorescent cell membrane dyes (DiI, DiD, and DiO), a multicolor DiOlistic (MuDi) labeling
approach has been accomplished in the mouse brain and retina.(Gan et al., 2000) These multiple
separate color channels enabled labeling and visualization of a large number of cells in close
proximity to each other while still allowing to distinguish cells from their neighbors.

In this work, we aimed to determine the suitability of MuDi labeling(Gan et al., 2000) as an
approach to visualize collagenous LC astrocyte morphology, *in situ*. The scope of this work is
focused on testing the ability of MuDi labeling to reveal individual astrocyte morphologies in the
collagenous LC. We demonstrate that this technique reveals morphological detail of individual
astrocytes across the LC with a single labeling application. Astrocyte morphological features can
be semi-automatically quantified and extracted from images on a cell-by-cell and branch-by-
branch basis for detailed analysis. This labeling method rendered samples compatible with other
labeling and imaging modalities such as immunolabeling for GFAP, lectin-labeling of blood
vessels, and second harmonic generation (SHG) imaging of LC collagen beams. The biological
relevance of findings from LC astrocyte models will be discussed in detail in future work. MuDi
labeling of the collagenous LC can open new avenues for qualitative and quantitative investigation
of astrocyte morphology in health, and in the future, in glaucoma.

<u>2. Methods</u>

2.1. **Tissue acquisition and preparation**
Pig, sheep, goat, and monkey eyes were utilized for experiments (N = 2, 4, 16, and 12
eyes, respectively). Pig, sheep, and goat eyes were acquired within 1 hour of sacrifice. Rhesus
macaque monkey eyes were obtained within 30 minutes of sacrifice. ONHs were isolated and
fixed in methanol-free 4% PFA for 2-24 hours and held in PBS at 4°C. All procedures were
approved by the University of Pittsburgh's Institutional Animal Care and Use Committee (IACUC)
and adhered to both the guidelines set forth in the National Institute of Health's Guide for the Care
and Use of Laboratory Animals and the Association of Research in Vision and Ophthalmology

(ARVO) statement for the use of animals in ophthalmic and vision research. No live animals were
used in this work.

2.2. **Vibratome sectioning**

ONHs were embedded in low melting temperature agarose (Sigma, A9045), secured to
the stage of a vibratome (Leica, VT1200S) with cyanoacrylate glue, and immersed in ice-cold
PBS. Coronal slices were cut at 150µm-thickness. As the collagen of the ONH is highly elastic,
high blade advancement speeds can cause deformation of the tissue before cutting, resulting in
uneven slices, increased disruption of cell membranes, and in some instances, dislodging of the
tissue from its agarose block, resulting in loss of tissue orientation. Low blade advancement
speed (0.01-0.05 mm/second) helped ensure even tissue cutting. Slices were transferred from
the vibratome, gently with a paintbrush, to fresh PBS. Slices were screened under polarized light
and brightfield microscopy to visualize the collagen beams and surrounding anatomical structures
indicative of the LC region. Slices were stored in PBS at 4°C for up to 48 hours prior to DiOlistic
labeling.

2.3. **Multicolor DiOlistics bullet preparation**

DiOlistics bullets were prepared similarly to as done previously by others.(Benediktsson
et al., 2005; Gan et al., 2000; Jakobs, 2014; O'Brien and Lummis, 2006; Sun et al., 2010)   For
multicolor bullets, 70mg of 1.0µm diameter gold microparticles (BioRad, 1652263) were divided
equally into 7 1.5mL tubes and each mixed in 100µL ethanol. 4mg of each DiI (Sigma-Aldrich
42364, 549/567nm), DiO (Invitrogen D275, 484/501nm), and DiD (Invitrogen D7757, 644/665nm
excitation/emission) were used to make separate stock solutions in 500µL ethanol. Dye solutions
were added to tubes containing gold microparticles in all combinations and in equal proportions,
such that each tube received a total of 1.7mg of dye. For example, DiI-coated microparticles
received 1.7mg of DiI. DiI + DiO-coated microparticles received 0.85mg of DiI and 0.85mg of
DiO. DiI + DiO + DiD-coated microcarriers received 0.57mg of each dye.

Gold-dye mixtures were vortexed for 30 seconds, pipetted onto separate glass microscope
slides (Fisher, 12550A3), and spread evenly across slides with a pipette tip. Ethanol was allowed
to fully evaporate for at least 20 minutes, rendering dry microcarriers coated in dyes. The dry
microcarriers were scraped off their respective slides with a razor and collected together.
Microcarriers were suspended in 1mL deionized water and sonicated for 10 minutes to minimize
clumping. Sonicated microcarriers were filtered through a 20µm filter (Millipore, NY2004700) to

remove large clumps and resuspended in 3mL deionized water. 30µL of 0.5 mg/ml
polyvinylpyrrolidone in isopropanol was added to this mixture.

Microcarrier mixture was loaded into Tefzel tubing (Bio-Rad, 1652441) in a tubing prep
station (Bio-Rad, 1652418). Microcarriers were allowed to sediment to the bottom of the tube for
5-10 minutes before liquid was slowly withdrawn from the tube. Tubing was rotated for 5 minutes
in the tubing prep station to distribute the microcarriers along the interior of the tubing. Nitrogen
was run through the tubing for 30 minutes at 2 PSI to remove the remaining water. Tubing was
cut into 1.3cm lengths with a tubing cutter (Bio-Rad, 1652422) and stored in an airtight container
with desiccant, protected from light, for as long as 6 months. Single-color bullets were prepared
the same way as multicolor bullets, but using 4mg of DiI or DiD as the only dye and 23mg of gold
microparticles instead of 70mg.

2.4. **DiOlistic labeling**

A Helios Gene Gun (BioRad, 1652411) was loaded with bullets, connected to a
pressurized nitrogen tank, and used to deliver dye-coated microcarriers into vibratome slices at
150-200 PSI. Vibratome slices were gently removed from PBS with a paintbrush and placed in a
60mm dish. Slices were covered with a 20µm filter (Millipore, NY2004700) and the spacer of the
gene gun was positioned flush with the filter. The gene gun was oriented perpendicular to the
sample as in Figure 2, and each sample was shot once with the gene gun. Samples were returned
to PBS within 30 seconds to prevent sample dehydration and tissue damage. Samples
counterstained with DAPI were incubated in 1:1000 DAPI (ThermoFisher, 62248) for 20 minutes
and washed 3 times with PBS. Slices were mounted labeled-side-up on glass slides in water-
based mounting medium (Shandon, 9990402) for microscopy.

2.5. **Additional non-DiOlistic labeling**

For samples labeled immunohistochemically and/or with lectin (non-DiOlistic labeling), this
was done either before, after, or without DiOlistic labeling. Each method has specific benefits
and limitations, detailed below.

Before DiOlistic labeling

Samples with non-DiOlistic labeling completed before DiOlistic labeling, such as those in
Figure 1C-**E**, allowed imaging of DiOlistically-labeled cells and other labels in the same imaging
session. This minimizes the risk of interpretation errors, such as false negative and false positive
colocalization of GFAP and DiOlistics signals, caused by potential image registration
imperfections or sample distortion between imaging sessions. These samples however, cannot
be extensively permeabilized with detergents to enhance non-DiOlistic label penetration through
sample thicknesses. Detergents, such as the commonly used Triton X-100, create holes in cell
membranes that can result in substantial leakage of DiOlistically-delivered dyes from cells into
extracellular spaces and onto neighboring cells, preventing clear visualization of individual cell
morphologies. This method is also not readily compatible with the use of multicolor DiOlistic
bullets, as many fluorescent probes used in immunohistochemistry have similar fluorescent
profiles to DiD, DiO, or DiI. Therefore, single-color bullets were used in these samples and less
cells were labeled at once to ensure clear spatial separation of labeled cells.

Three vibratome slices from three goat eyes were incubated for 4 hours in 5% BSA
blocking solution without permeabilization agent, overnight in GFAP primary antibody (rabbit
origin, (Invitrogen, PA5-16291) at 1:200 in blocking solution, and washed 3x10min in PBS.
Samples were incubated with secondary antibody at 1:500 dilution (488 goat-anti-rabbit, Abcam,
ab150077) for 6 hours at room temperature, washed 3x10min in PBS, incubated in 1:1000 DAPI
for 20 minutes, and washed 3x10 minutes in PBS. Immediately after non-DiOlistic labeling was
completed, samples were labeled with single-color DiD DiOlistics.

After DiOlistic labeling

Samples labeled with non-DiOlistic labels after DiOlistic labeling, such as those in **Figure
8**, were permeabilized in order to enhance non-DiOlistic label penetration, without the same
concern necessary for cell membrane integrity as samples treated with non-DiOlistic labels before
DiOlistic labeling. Samples treated with non-DiOlistic labels after DiOlistic labeling must however
be removed from their slides after imaging of DiOlistics signal for this additional labeling and
washing. Images from the first round of imaging (to visualize DiOlistic labeling) and second round
of imaging (to visualize non-DiOlistic labeling) must be carefully registered to each other based
on shared information between the two, such as DAPI signal and/or ECM autofluorescence. The
process of lipophilic dye removal and re-labeling caused minimal tissue distortion but given the

scale of astrocytes and their fine processes, interpretation of colocalized signals should be done
with caution and/or verified with samples labeled before to DiOlistics.

After imaging of 4 DiOlistically-labeled slices was completed, one slice was introduced to
an ethanol gradient (50%, 70%, 90% for 1 hour each) and held in 98% ethanol for two additional
days to test the ability of ethanol to remove dyes from the tissue. When this sample was observed
post-ethanol wash with the same imaging parameters, no DiOlistically-labeled cells or appreciable
MuDi dye signal was observed in the tissue. The remaining three samples were blocked, labeled
with non-DiOlistics labels, and washed in PBS prior to ethanol-mediated MuDi dye removal.
These three samples were blocked and permeabilized thoroughly in 5% BSA 0.05% Triton X-100
(block/perm) for 24 hours, incubated with GFAP primary antibody diluted 1:200 and 1:200 DyLight
Tomato lectin (VectorLabs, DL-1178) in block/perm for 48 hours, washed for 1 day in PBS,
incubated for 24 hours in 1:500 secondary in block/perm, and washed 3x10 minutes in PBS.
Samples were then introduced to the same ethanol gradient as above. Samples were rehydrated
for imaging and mounted in water-based mounting medium with the previously imaged side facing
up.

Without DiOlistic labeling

Tissue labeled for GFAP and without DiOlistic labeling, to visualize GFAP alone (**Figure
1A, B,** N = 1 eye), was prepared the same way as samples labeled after DiOlistic labeling, but
without lectin or ethanol washes.

2.6. **Imaging**

An upright confocal microscope (BX61; Olympus) was used to screen each DiOlistically-
labeled sample at low magnification (4x, 10x, and/or 20x). In samples with sufficient density and
quality of labeling in screening, tissues were observed under high magnification (40x). Dyes must
distribute across the cell membrane of individual cells in order to reveal morphology. Cells were
screened at 40x magnification for extent of dye spread prior to extensive imaging. Segmentation-
quality images were collected after astrocyte end-feet and high-order processes could be
visualized across the LC. This occurred within 1 hour of labeling at room temperature. Imaging
was completed within 24-48 hours of DiOlistic labeling. 40x images were collected at an XY
image size of 1024x1024 or 800x800 pixels, and with a 0.5-2μm Z step throughout the visible
thickness of the sample. Imaging of non-DiOlistics labels (GFAP, lectin) was completed with the
same confocal microscope at 20 and 40x magnifications, with the same X, Y, and Z image sizes
as used for imaging of DiOlistics. SHG imaging was conducted with a multiphoton microscope

(A1; Nikon) at 25x. SHG images were collected at a 1024x1024 or 512 x 512 pixel XY image size
at a 2-5µm Z step throughout the visible thickness of the sample.

2.7. **Astrocyte segmentation and 3D model construction**

Images were imported into Imaris (Bitplane, version 9.9) software for 3D visualization and
model construction of labeled astrocytes. The Filaments tool and the "AutoPath-no loops"
algorithm were used to automatically segment individual astrocytes. Each astrocyte was created
with a single starting-point at its soma and seed-points along its branches. Imaris-generated
models were checked to ensure that they faithfully represented image volume information.
Aberrant branches were removed, and missing branches were added manually. Quantitative
branch-by-branch level information about morphological features, such as branch number, length,
thickness, hierarchy, straightness, and Sholl intersections was extracted from the Statistics panel.
Sholl analysis was conducted with the Filament Sholl Analysis XTension.

3. Results

3.1. **DiOlistically-labeled cells with astrocyte morphologies are GFAP[+]**

GFAP labeling revealed astrocyte intermediate filaments in the collagenous LC. However,
GFAP labeling was insufficient to understand the morphology of individual astrocytes within LC
tissues (**Figure 1A, B**). The high density and structural complexity of GFAP filaments rendered
discerning one astrocyte from its neighbor not possible. Additionally, GFAP does not reveal the
full extent of the spatial territory occupied by astrocytes.(Bushong et al., 2002; Escartin et al.,
2021; Luna et al., 2016; Stokum et al., 2018; Sun and Jakobs, 2012) GFAP filaments are often
found within thicker primary processes but can be absent in higher-order processes. Conclusions
about the morphology of astrocytes labeled with GFAP alone are therefore subject to critical
misinterpretations.

DiOlistic delivery of dye-coated microcarriers to GFAP-labeled goat LC tissues resulted in
stochastic cell membrane labeling of individual cells. The density of single-color dye-coated
microcarriers delivered to LC tissues was optimized to ensure clear spatial separation of labeled
cells. DiOlistically-labeled cells with astrocyte morphologies colocalized partially with GFAP,
demonstrating astrocyte identity. **Figure 1C & D** show instances of DiOlistically-labeled cells of
the goat LC with astrocyte morphologies. These cells had a single soma, typically 3-8 major
processes, and multiple fine higher-order processes. Processes extended within neural tissue

pores and contacted LC beams. In samples counterstained with DAPI, these cells contained a
single nucleus (**Figure 1E**).

3.2. **Multicolor DiOlistic labeling of the LC: workflow**

To visualize multiple collagenous LC astrocytes in close proximity to each other while still
discerning one cell from another, we employed a MuDi labeling approach, as done by Gan et
al(Gan et al., 2000) to visualize cells in the brain and retina of mice. ONHs from pig, sheep, goat,
and monkey eyes were sectioned into 150 µm-thick coronal slices with a vibratome (**Figure 2A**).
Groups of microcarriers were coated with all combinations of 3 spectrally distinct fluorescent cell
membrane dyes to create a total of seven color combinations (**Figure 2B**). Groups of tens to
hundreds of multicolor dye-coated microcarriers were loaded into "bullets" for simultaneous
delivery to tissues with a gene gun. These groups of microcarriers were propelled into tissue
slices containing collagenous LC (**Figure 2C**). Microcarriers were embedded within cells and the
dyes they carried were distributed across individual cell membranes. DiOlistically-labeled tissues
were imaged with confocal microscopy to visualize labeled cells and DAPI-labeled cell nuclei
and/or ECM autofluorescence. Select tissues were imaged with second harmonic generation
(SHG) microscopy to visualize LC collagen (**Figure 2D**).

3.3. **Multicolor DiOlistic labeling allows LC astrocyte visualization across seven distinct**
color channels

MuDi allowed for LC astrocyte visualization across the seven distinct color channels
created by all combinations of the fluorescent dyes, DiD, DiO, and DiI (**Figure 3**). The seven
color channels shown in **Figure 3** were the most commonly observed colors for labeled
astrocytes. Example cells in **Figure 3** were of goat origin. As noted in Gan et al,(Gan et al., 2000)
additional colors were observed in cells labeled by multiple microcarriers. For example, a cell in
this study shown in cyan received one part DiI and one part DiO. If this cyan cell received an
additional microcarrier containing one part DiI and one part DiD, it would appear as an
intermediate color not included in the original seven categories. At the density of microcarriers
included in each bullet, the gas pressure used for each gene gun shot, and the distance the gene
gun was set from the tissues in this study, most labeled cells received only one microcarrier. The
distribution of microcarriers delivered to tissue samples could be observed under transmitted light
(**Supplemental Figure 1**) as dark puncta approximately 1µm in diameter.

3.4. **Multicolor DiOlistic labeling reveals 3D morphologies of individual astrocytes across**
the LC

MuDi-labeled coronal slices of the LC allowed for visualization of tens to hundreds of
individual astrocytes across the tissue, *in situ* (**Figure 4A**). Confocal images collected at high
magnification revealed cell bodies and fine processes of individual cells in close proximity to each
other (**Figure 4B**). Most astrocyte branches were oriented along the coronal plane while some
spanned the anterior-posterior direction, as seen in mice.(Lye-Barthel et al., 2013; Wang et al.,
2017) Example tissue shown in **Figure 4** was of goat origin. Optimal labeling density was
approximately 140 cells per mm^2 of canal section area. At this labeling density, we were able to
visualize a large number of cells across the LC with minimal instances of two or more cells labeled
with the same dye(s) in close enough proximity as to obscure individual cell morphologies.
Collagen beams characteristic of the collagenous LC were visualized with SHG, revealing the
relationship of DiOlistically-labeled astrocytes to surrounding collagen supports. SHG signal
colocalized with tissue autofluorescence in the 405nm channel observed in confocal microscopy
and aided in registration of SHG images to confocal images.

Z-stack images collected at 40x magnification allowed for creation of DiOlistically-labeled
astrocyte 3D models (**Figure 4C**). Example astrocyte models are shown in relation to their
respective confocal volume information in **Figure 5** and **Video 1**. These models granted
automated extraction of quantitative morphological features such as branch length, thickness,
hierarchy, and straightness (**Figure 6**). Three example astrocyte models from the goat LC are
included in **Figure 6**. For 3D views of these astrocytes see **Video 2**. The same methodology
was used to create 50 astrocyte models, supporting the robustness of this method for systematic
analysis. Sholl analysis (**Figure 7**), convex hull area, number of branching points, number of
terminal points, cell volume, and a variety of other quantitative metrics were also readily captured.
Sholl data from 3 example collagenous LC astrocytes demonstrates a similar average distribution
of crossing counts to mouse glial lamina astrocytes from Lye-Barthel et al, (Lye-Barthel et al.,
2013) with higher crossing counts in collagenous LC astrocytes. The ability to extract these
quantitative metrics can be helpful in future studies designed to characterize and/or compare
healthy and glaucomatous collagenous LC astrocyte morphologies.

The morphologies of collagenous LC astrocytes were in line with those of other fibrous
astrocytes from large mammals, just as glial lamina astrocyte morphologies are in line with fibrous
astrocyte morphologies of the mouse.(Oberheim et al., 2009; Sun et al., 2010) Protoplasmic
astrocytes found in other areas of the central nervous system have substantially bushier
morphologies and higher branch densities than do fibrous astrocytes.(Oberheim et al., 2009)

3.5. **Labeling and visualization of LC astrocytes is effective across species**

MuDi allowed visualization of LC astrocytes across all species with a collagenous LC

tested (**Figure 8**). Astrocytes were visualized in the LC of monkey (**Figure 8A**), goat (**Figure 8B**),

pig (**Figure 8C**), and sheep (**Figure 8D**). Detail of astrocyte morphologies was visible across

species, including thin higher-order branches and end-feet (A'-H'). End-feet were observed in

contact with collagenous LC beams (**Figure 4B, Figure 8**).

3.6. **Multicolor DiOlistics is compatible with labeling and imaging of other structures**

MuDi was compatible with labeling and imaging of other collagenous LC structures. We

have shown that MuDi in the LC allows for seven color channels with which astrocytes can be

visualized. Paired SHG, DAPI labeling, and additional post-DiOlistics markers, this provides the

potential for 12 distinct channels for visualization of LC structures in the same tissue.

SHG imaging and visualization of collagen was possible in DiOlistically-labeled LC tissues

without additional sample labeling or preparation. The presence of LC collagen beams is a

defining feature of the anatomic region and is a key difference between the collagenous LC, found

in humans and the species used in this study, and the glial lamina of mice and rats. Visualizing

LC collagen in tandem with astrocyte morphology and spatial arrangement can provide

verification of tissue location and a better understanding of individual astrocyte relationships to

surrounding collagen beams.

With many fluorophores available being in the DiO, DiI, and DiD ranges (similar to FITC,

TRITC, and Cy5), the use of MuDi could preclude the use of other common labeling options.

However, due to the solubility of DiO, DiI, and DiD in ethanol, these dyes could be washed from

tissue samples in an ethanol gradient. Samples labeled for GFAP (FITC-range secondary) and

lectin (Cy5-range conjugate) that were then washed of DiOlistics dyes allowed specific

visualization of GFAP intermediate filaments and blood vessels without appreciable DiOlistics

signal (**Figure 9**). This provided an additional two distinct channels for visualization of LC

structures in these tissues, with potential for an additional third post-DiOlistics channel in the

TRITC/DiI-range, not used here. A total of 12 distinct channels for visualization of LC structures

could be possible in the same sample with the use of MuDi (7), DAPI (1), SHG (1), and additional

labels after washing away DiOlistics dyes in the FITC, TRITC, Cy5 ranges (3).

3.7. **Multicolor DiOlistic labeling can reveal axons and other cell types**

396   MuDi preferentially labels astrocytes in LC coronal slices due to their abundance, size,

and general orientation along the coronal plane. However, this labeling technique is not inherently

discriminatory of cell type. RGC axons, LC cells, and scleral fibroblasts can also be labeled by

this technique but have markedly different labeling frequencies and cell morphologies. Examples

of labeled cells with fibroblast-like morphologies along LC beams and in the peripapillary sclera

region are included in **Figure 10**. Because of the largely longitudinal orientation of RGC axons,

individual axons will occupy little space in the coronal plane. When axons in coronal slices are

labeled with DiOlistics, they appear as small puncta or hollow tubes in projection, depending upon

magnification power. With increasing deviation of sample orientation from the coronal plane, each

axon occupies more space within a tissue slice. Axons are more likely to be stochastically hit by

a DiOlistics microcarrier when they occupy more space. Axon labeling in slices cut at an angle

that deviated from the coronal plane is shown in **Figure 11**. Axons are visualized in the nerve

fiber layer and prelamina (**Figure 11A**) as well as the LC, retrolamina, and optic nerve (**Figure**

11B, C).   In samples sectioned longitudinally, axons were frequently labeled, apparent as

filamentous structures oriented along the sagittal plane (**Figure 11C**).

<u>4. Discussion</u>

414   In this work, we investigated the capability of MuDi labeling to allow visualization of

astrocytes in the collagenous LC. Analysis of tissues treated with this approach resulted in three

key findings. First, DiOlistics allowed visualization of individual astrocytes and fine details of their

morphologies across the collagenous LC. Second, multicolor labeling granted visualization of

multiple cells in close proximity at once while still allowing to discern individual cells from their

neighbors. Third, 3D astrocyte models made from confocal image volumes facilitated automated

quantification of morphological features. Below we discuss the significance and implications of

these findings.

4.1.   **DiOlistics allowed visualization of individual astrocytes and fine details of their**

morphologies across the collagenous LC. As shown in earlier works from other groups,

DiOlistic labeling is capable of revealing complex morphologies of neurons in brain

tissues(O'Brien and Lummis, 2006; Seabold et al., 2010; Staffend and Meisel, 2011) down to the

detail of their dendritic spines. DiOlistics is also capable of labeling astrocytes in complex tissue

environments, as shown in the mouse, chimpanzee, and human brains.(Oberheim et al., 2009,

2008) In mice, DiOlistics has been used to reveal individual astrocyte morphologies in the glial
lamina specifically.(Gan et al., 2000; Oberheim et al., 2008)   Individual lamina astrocyte
morphologies in species with a collagenous LC, like humans, are less well-described. We applied
DiOlistics to the collagenous LC of pig, sheep, goat, and monkey to visualize resident astrocyte
morphologies. We demonstrated that DiOlistics allowed visualization of individual astrocytes and
microscale detail of their morphologies in the context of their complex LC tissue environments.

As shown in **Figure 4A**, DiOlistic labeling resulted in transfer of dye-coated microcarriers
to cells across coronal slices through the LC. The same sample imaged at higher magnification
(**Figure 4B**) reveals morphologies of individual astrocytes within LC pores.   In samples
immunostained for GFAP, a common astrocyte marker, DiOlistically-labeled cells with astrocyte
morphologies were GFAP[+]. Detailed structural analysis of GFAP, F-actin, nuclei, and collagen
beams in the healthy and glaucomatous human LC(Guan et al., 2022) has provided important
insights into the differences of these elements between health and advanced disease. However,
detail of individual LC astrocyte morphologies cannot be determined through visualization of these
elements alone. Consistent with other descriptions of ONH astrocytes observed with different
methods,(Ju et al., 2015; Lye-Barthel et al., 2013; Nguyen et al., 2017; Wang et al., 2017) the LC
astrocytes visualized in this study had multiple ramified processes. Astrocyte end-feet were often
observed in contact with LC beams, shown in detail in **Figure 7**.   To our knowledge, individual
astrocyte morphologies in the collagenous LC, labeled through a method that can reveal the
entirety of the cell plasmalemma, have not been previously reported.

Primary culture of astrocytes from models that have a collagenous LC has been
accomplished(Hernandez et al., 1988; Lopez et al., 2020) and provides an excellent opportunity
for *in vitro* visualization. However, the process of primary culture requires removal of cells from
the context of surrounding collagen beams, retinal ganglion cells, blood vessels, and other
astrocytes. This, along with maintenance on a 2D surface rather than a 3D tissue environment,
can result in substantial alteration of astrocyte morphologies. Visualization of LC astrocytes within
the context of their environment, as we have done here, with preserved spatial relationships to
surrounding structures, is optimal to understand their original morphologies.

4.2. **The multicolor approach allows for labeling of many cells at once while still allowing**
separation of cells from their neighbors. Multicolor DiOlistics, developed by Gan et. al, was
demonstrated to allow visualization of and spatial separation of neighboring neurons and
astrocytes in the mouse brain and retina.(Gan et al., 2000) We applied this multicolor approach
to label and visualize tens to hundreds of cells across the pig, goat, sheep, and monkey LC at

once while allowing separation of individual cells from their neighbors. Due to the density of
astrocytes in the LC and their complex, highly branched morphologies, techniques using a single
fluorescent label to visualize astrocytes would require a relatively low labeling density in order to
effectively distinguish between neighboring cells. Manual injection of different dyes into individual
cells is possible,(Ju et al., 2015) but can be time and labor-intensive. After a single gene gun
shot, MuDi allows spatial separation of neighboring astrocytes to be visualized through multiple
distinct color channels.

MuDi is similar in concept to Brainbow small research animal reporter lines(Livet et al.,
2007; Pan et al., 2011; Richier and Salecker, 2015) which have been used to help distinguish
individual neurons and astrocytes from their neighbors in the brain. As in the *Thy1-Brainbow-1.0*
mouse line, a greater number of color channels than seven can be created from 3 fluorescent
reporters if more intermediate colors are utilized (ex. two parts green and one part red). With
combinatorial expression of three fluorescent reporters (GFP, RFP, and CFP), Livet et al could
distinguish neural cells in approximately 100 distinct color groups.(Livet et al., 2007) MuDi, and
variations of it, can be useful to investigate questions similar to those allowed by Brainbow tissues,
but in large animal or human donor eyes where gene modification is not as easily achieved. For
example, the spatial territories of individual collagenous LC astrocytes and the corresponding
regions of RGC axons over which they have direct physiologic influence could be observed in
health in disease. This has potential to inform our understanding of how individual astrocyte
territories and their zones of physiologic influence may change in stress states, informing our
understanding of disease progression and pathways for neuroprotective strategies.
484

4.3. **Automated quantification of morphological features was facilitated by DiOlistics-**
derived 3D astrocyte models. 3D astrocyte models were readily constructed from confocal
image volumes for automated quantification of morphological features. Sharing representative
images and qualitative interpretations based on large image sets can be a valuable way to
communicate summaries of scientific findings. However, quantifying morphological features from
large image datasets in addition to sharing qualitative interpretations can allow for a clearer,
broader, and more replicable picture of LC astrocyte morphology. Additionally, morphologies of
astrocytes are often categorized into two broad, qualitative groups: healthy / "non-reactive" and
pathologic / "reactive." As the understanding of astrocyte structure and function has become
more nuanced, it is important to supplement qualitative and categorical observations with detailed
quantitative ones that can be evaluated and analyzed statistically, as is allowed by the approach
described in this work.

The quantification of morphological features that this approach allows opens doors to ask

quantitative questions about the heterogeneity of astrocyte morphologies within a single

individual, among different individuals, and between different species, as we aim to do in our

future work. The technique can allow for investigation of quantitative differences and similarities

in astrocyte morphologies between LC regions, such as central vs. peripheral LC astrocytes or

nasal vs. temporal ones. It can additionally allow for quantification of astrocyte spatial

relationships with surrounding collagenous LC elements, such as collagen beams, blood vessels,

and other astrocytes. Importantly, it may provide clearer insights into astrocyte morphology

throughout the progression of aging and glaucoma in experimental models or in human donor

eyes. The ability to label tens to hundreds of cells at once in a single tissue and quantify their

morphologies can allow us to discern patterns that may be missed in smaller sample sizes or with

qualitative interpretation alone.

Important limitations of this approach must be acknowledged. DiOlistics relies on

stochastic labeling of cell membranes and is not inherently discriminant of cell type. Astrocytes

are abundant in the LC, accounting for over 90% of nuclei in the region,(Guan et al., 2022) with

cell bodies largely oriented along the coronal plane. Therefore, many cells labeled in these

coronal slices were astrocytes. MuDi will label larger cells or more coronally-oriented cells more

frequently than smaller and more longitudinally-oriented ones. This approach may therefore

oversample larger astrocytes oriented along the coronal plane and undersample smaller

astrocytes with a smaller footprint along the coronal plane. Despite the lack of complete cell-type

specificity, astrocyte morphologies were distinctly discernible from those of other LC cellular

elements, such as retinal ganglion cell axons and LC cells. The technique's compatibility with

other more cell-type-specific labeling methods, such as the GFAP labeling in this study, allowed

for additional validation of the identity of DiOlistically-labeled cells.

The penetration of microcarriers in tissues is determined by the biomechanical properties

of the tissue, the pressure at which microcarriers are propelled into the tissue, and the distance

the gene gun is positioned from the sample. At 150-200 PSI and gene gun distance of 30mm,

we observed microcarriers up to 80μm deep within the samples. Higher pressures can increase

microcarrier penetration depth, but risks damage to tissues. The limited penetration of

microcarriers necessitates sectioning of tissues prior to labeling. The process of sectioning can

result in severing cells and incomplete visualization of cell morphology. The collection of thick

tissue slabs with a vibratome minimizes the number of cells in samples with severed processes.
However, severed cells are likely to leak dye into surrounding tissues more quickly than intact
cells. Cells with poorly defined morphologies due to dye leakage were not suitable for 3D model
construction and can be excluded from morphological analyses. Gene gun modifications have
been shown to allow for deeper bullet penetration at lower gas pressures.(O'Brien et al., 2001) A
combination of gene gun modification and dye-preserving tissue clearing methods(Ke et al., 2016;
Li et al., 2019) could allow for visualization of labeled astrocytes in whole LCs.
538

The time-sensitive nature of imaging samples labeled with MuDi is an important
consideration for study design in which large high-resolution confocal image sets can take several
hours to collect. LC slices labeled with MuDi must be imaged within 24-48 hours of labeling, as
dyes can leak out of cells and into surrounding space over time. Dye leakage and redistribution
obscures the morphology of individual cells, preventing morphological analysis. Examples of
tissues with dye leakage preventing high quality visualization of individual cell morphologies are
included in **Supplemental Figure 2**. The use of fixable dyes such as CM-DiI may help mitigate
this issue.
547

The tissues used in this study were fixed prior to labeling and imaging. This prevents
investigation of cell function in conjunction with morphology and tracking any changes over time.
DiOlistic labeling has been accomplished in organotypic slice cultures of brain tissues.
Preliminary data from our lab demonstrates that MuDi labeling can be accomplished in the non-
fixed LC vibratome slices as well. This technique, or similar techniques using fluorescent dextrans
instead of cell membrane dyes,(Jakobs, 2014) may allow investigation of LC astrocyte structure
and function in *ex vivo* slice culture. *Ex vivo* culture and drug delivery systems of eyes with a
collagenous LC(Chan et al., 2022) combined with DiOlistics may be able to relate structural
astrocyte features to functional ones and track changes over time.
557

Overall, MuDi labeling of the collagenous LC provides a way to visualize morphological
detail of individual LC astrocytes at a scale not provided by other techniques. This approach
revealed LC astrocyte morphologies, allowed cells in close proximity to be distinguished from one
another, and facilitated quantification of morphological features. It is compatible with other
labeling and imaging techniques such as immunohistochemistry and SHG. MuDi labeling
provides the potential for future large-scale qualitative and quantitative analysis of astrocyte
morphological features and spatial relationships with interacting elements, such other astrocytes,

collagen beams, blood vessels, and axons, across the collagenous LC. Investigation was
possible across different healthy animal models. In future studies, this technique may allow for
investigation of LC astrocyte morphology in different glaucoma model progression states or in
human donor tissues as well. This approach can help provide important insights into collagenous
LC astrocyte physiology in health and disease.

<u>Acknowledgements:</u>

Disclosures: Nothing to disclose.

Funding: Supported in part by National Institutes of Health grants R01-EY023966, R01-
EY031708, R01-HD045590, R01-HD083383, 1S10RR028478-01, P30-EY008098, and T32-
EY017271; Eye and Ear Foundation (Pittsburgh, PA); Research to Prevent Blindness
(unrestricted grant to UPMC Ophthalmology and Stein Innovation Award to Sigal IA).
576
577 We would like to acknowledge Dr. Martin Oberbarnscheidt for generously sharing vibratome
578 equipment and training in its proper use.

References

Attwell, D., Buchan, A.M., Charpak, S., Lauritzen, M., Macvicar, B.A., Newman, E.A., 2010. Glial and neuronal control of brain blood flow. Nature 468, 232–243.

Benediktsson, A.M., Schachtele, S.J., Green, S.H., Dailey, M.E., 2005. Ballistic labeling and dynamic imaging of astrocytes in organotypic hippocampal slice cultures. J. Neurosci. Methods 141, 41–53.

Bonney, S.K., Coelho-Santos, V., Huang, S.-F., Takeno, M., Kornfeld, J., Keller, A., Shih, A.Y., 2022. Public Volume Electron Microscopy Data: An Essential Resource to Study the Brain Microvasculature. Front Cell Dev Biol 10, 849469.

Bushong, E.A., Martone, M.E., Jones, Y.Z., Ellisman, M.H., 2002. Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. J. Neurosci. 22, 183–192.

Butt, A.M., Colquhoun, K., Tutton, M., Berry, M., 1994. Three-dimensional morphology of astrocytes and oligodendrocytes in the intact mouse optic nerve. J. Neurocytol. 23, 469–485.

Calì, C., Agus, M., Kare, K., Boges, D.J., Lehväslaiho, H., Hadwiger, M., Magistretti, P.J., 2019. 3D cellular reconstruction of cortical glia and parenchymal morphometric analysis from Serial Block-Face Electron Microscopy of juvenile rat. Prog. Neurobiol. 183, 101696.

Chan, D., Won, G.J., Read, A.T., Ethier, C.R., Thackaberry, E., Crowell, S.R., Booler, H., Bantseev, V., Sivak, J.M., 2022. Application of an organotypic ocular perfusion model to assess intravitreal drug distribution in human and animal eyes. J. R. Soc. Interface 19, 20210734.

Chaudhary, P., Stowell, C., Reynaud, J., Gardiner, S.K., Yang, H., Williams, G., Williams, I., Marsh-Armstrong, N., Burgoyne, C.F., 2022. Optic Nerve Head Myelin-Related Protein, GFAP, and Iba1 Alterations in Non-Human Primates With Early to Moderate Experimental Glaucoma. Invest. Ophthalmol. Vis. Sci. 63, 9.

Dong, Y., Fu, Y., Qian, X., Lin, L., Yuan, Y., Li, Y., Shao, W., Gao, Q., 2022. Optic nerve head astrocytes contribute to vascular associated effects. Front. Med. 9, 943986.

Escartin, C., Galea, E., Lakatos, A., O'Callaghan, J.P., Petzold, G.C., Serrano-Pozo, A., Steinhäuser, C., Volterra, A., Carmignoto, G., Agarwal, A., Allen, N.J., Araque, A., Barbeito, L., Barzilai, A., Bergles, D.E., Bonvento, G., Butt, A.M., Chen, W.-T., Cohen-Salmon, M., Cunningham, C., Deneen, B., De Strooper, B., Díaz-Castro, B., Farina, C., Freeman, M., Gallo, V., Goldman, J.E., Goldman, S.A., Götz, M., Gutiérrez, A., Haydon, P.G., Heiland, D.H., Hol, E.M., Holt, M.G., Iino, M., Kastanenka, K.V., Kettenmann, H., Khakh, B.S.,

Koizumi, S., Lee, C.J., Liddelow, S.A., MacVicar, B.A., Magistretti, P., Messing, A., Mishra,

A., Molofsky, A.V., Murai, K.K., Norris, C.M., Okada, S., Oliet, S.H.R., Oliveira, J.F.,

Panatier, A., Parpura, V., Pekna, M., Pekny, M., Pellerin, L., Perea, G., Pérez-Nievas, B.G.,

Pfrieger, F.W., Poskanzer, K.E., Quintana, F.J., Ransohoff, R.M., Riquelme-Perez, M.,

Robel, S., Rose, C.R., Rothstein, J.D., Rouach, N., Rowitch, D.H., Semyanov, A., Sirko, S.,

Sontheimer, H., Swanson, R.A., Vitorica, J., Wanner, I.-B., Wood, L.B., Wu, J., Zheng, B.,

Zimmer, E.R., Zorec, R., Sofroniew, M.V., Verkhratsky, A., 2021. Reactive astrocyte

nomenclature, definitions, and future directions. Nat. Neurosci. 24, 312–325.

Gan, W.B., Grutzendler, J., Wong, W.T., Wong, R.O.L., Lichtman, J.W., 2000. Multicolor

"DiOlistic" Labeling of the Nervous System Using Lipophilic Dye Combinations. Neuron 27,

219–225.

Ge, L.-J., Yang, F.-H., Li, W., Wang, T., Lin, Y., Feng, J., Chen, N.-H., Jiang, M., Wang, J.-H.,

Hu, X.-T., Chen, G., 2020. In vivo Neuroregeneration to Treat Ischemic Stroke Through

NeuroD1 AAV-Based Gene Therapy in Adult Non-human Primates. Front Cell Dev Biol 8,

590008.

Guan, C., Pease, M.E., Quillen, S., Ling, Y.T.T., Li, X., Kimball, E., Johnson, T.V., Nguyen, T.D.,

Quigley, H.A., 2022. Quantitative Microstructural Analysis of Cellular and Tissue

Remodeling in Human Glaucoma Optic Nerve Head. Invest. Ophthalmol. Vis. Sci. 63, 18.

Heithoff, B.P., George, K.K., Phares, A.N., Zuidhoek, I.A., Munoz-Ballester, C., Robel, S., 2021.

Astrocytes are necessary for blood--brain barrier maintenance in the adult mouse brain.

Glia 69, 436–472.

Hernandez, M.R., 2000. The optic nerve head in glaucoma: role of astrocytes in tissue

remodeling. Prog. Retin. Eye Res. 19, 297–321.

Hernandez, M.R., Igoe, F., Neufeld, A.H., 1988. Cell culture of the human lamina cribrosa.

Invest. Ophthalmol. Vis. Sci. 29, 78–89.

Hernandez, M.R., Miao, H., Lukas, T., 2008. Astrocytes in glaucomatous optic neuropathy.

Prog. Brain Res. 173, 353–373.

Jakobs, T.C., 2014. Analysis of Morphology and Structural Remodeling of Astrocytes, in:

Bakota, L., Brandt, R. (Eds.), Laser Scanning Microscopy and Quantitative Image Analysis

of Neuronal Tissue. Springer New York, New York, NY, pp. 129–143.

J. Salazar, J., I. Ramírez, A., De Hoz, R., Salobrar-Garcia, E., Rojas, P., A. Fernández-Albarral,

J., López-Cuenca, I., Rojas, B., Triviño, A., M. Ramírez, J., 2019. Anatomy of the Human

Optic Nerve: Structure and Function, in: M. Ferreri, F. (Ed.), Optic Nerve. IntechOpen.

Ju, W.-K., Kim, K.-Y., Noh, Y.H., Hoshijima, M., Lukas, T.J., Ellisman, M.H., Weinreb, R.N.,

Perkins, G.A., 2015. Increased mitochondrial fission and volume density by blocking

glutamate excitotoxicity protect glaucomatous optic nerve head astrocytes. Glia 63, 736–

753.

Ke, M.-T., Nakai, Y., Fujimoto, S., Takayama, R., Yoshida, S., Kitajima, T.S., Sato, M., Imai, T.,

2016. Super-Resolution Mapping of Neuronal Circuitry With an Index-Optimized Clearing

Agent. Cell Rep. 14, 2718–2732.

Liddelow, S.A., Barres, B.A., 2017. Reactive Astrocytes: Production, Function, and Therapeutic

Potential. Immunity 46, 957–967.

Ling, Y.T.T., Pease, M.E., Jefferys, J.L., Kimball, E.C., Quigley, H.A., Nguyen, T.D., 2020.

Pressure-Induced Changes in Astrocyte GFAP, Actin, and Nuclear Morphology in Mouse

Optic Nerve. Invest. Ophthalmol. Vis. Sci. 61, 14.

Liu, C.-Y., Yang, Y., Ju, W.-N., Wang, X., Zhang, H.-L., 2018. Emerging Roles of Astrocytes in

Neuro-Vascular Unit and the Tripartite Synapse With Emphasis on Reactive Gliosis in the

Context of Alzheimer's Disease. Front. Cell. Neurosci. 12, 193.

Liu, Y.-X., Sun, H., Guo, W.-Y., 2022. Astrocyte polarization in glaucoma: a new opportunity.

Neural Regeneration Res. 17, 2582–2588.

Livet, J., Weissman, T.A., Kang, H., Draft, R.W., Lu, J., Bennis, R.A., Sanes, J.R., Lichtman,

J.W., 2007. Transgenic strategies for combinatorial expression of fluorescent proteins in the

nervous system. Nature 450, 56–62.

Li, W., Germain, R.N., Gerner, M.Y., 2019. High-dimensional cell-level analysis of tissues with

Ce3D multiplex volume imaging. Nat. Protoc. 14, 1708–1733.

Lopez, N.N., Clark, A.F., Tovar-Vidales, T., 2020. Isolation and characterization of human optic

nerve head astrocytes and lamina cribrosa cells. Exp. Eye Res. 197, 108103.

Luna, G., Keeley, P.W., Reese, B.E., Linberg, K.A., Lewis, G.P., Fisher, S.K., 2016. Astrocyte

structural reactivity and plasticity in models of retinal detachment. Exp. Eye Res. 150, 4–21.

Lye-Barthel, M., Sun, D., Jakobs, T.C., 2013. Morphology of astrocytes in a glaucomatous optic

nerve. Invest. Ophthalmol. Vis. Sci. 54, 909–917.

May, C.A., Lütjen-Drecoll, E., 2002. Morphology of the murine optic nerve. Invest. Ophthalmol.

Vis. Sci. 43, 2206–2212.

Merienne, N., Le Douce, J., Faivre, E., Déglon, N., Bonvento, G., 2013. Efficient gene delivery

and selective transduction of astrocytes in the mammalian brain using viral vectors. Front.

Cell. Neurosci. 7, 106.

Midgett, D., Liu, B., Ling, Y.T.T., Jefferys, J.L., Quigley, H.A., Nguyen, T.D., 2020. The Effects

of Glaucoma on the Pressure-Induced Strain Response of the Human Lamina Cribrosa.

Invest. Ophthalmol. Vis. Sci. 61, 41.

Morgan, J.E., 2000. Optic nerve head structure in glaucoma: astrocytes as mediators of axonal
damage. Eye 14 (Pt 3B), 437–444.

Moye, S.L., Diaz-Castro, B., Gangwani, M.R., Khakh, B.S., 2019. Visualizing Astrocyte
Morphology Using Lucifer Yellow Iontophoresis. J. Vis. Exp. https://doi.org/10.3791/60225

Nguyen, C., Midgett, D., Kimball, E.C., Steinhart, M.R., Nguyen, T.D., Pease, M.E., Oglesby,
E.N., Jefferys, J.L., Quigley, H.A., 2017. Measuring Deformation in the Mouse Optic Nerve
Head and Peripapillary Sclera. Invest. Ophthalmol. Vis. Sci. 58, 721–733.

Oberheim, N.A., Takano, T., Han, X., He, W., Lin, J.H.C., Wang, F., Xu, Q., Wyatt, J.D., Pilcher,
W., Ojemann, J.G., Ransom, B.R., Goldman, S.A., Nedergaard, M., 2009. Uniquely
hominid features of adult human astrocytes. J. Neurosci. 29, 3276–3287.

Oberheim, N.A., Tian, G.-F., Han, X., Peng, W., Takano, T., Ransom, B., Nedergaard, M., 2008.
Loss of astrocytic domain organization in the epileptic brain. J. Neurosci. 28, 3264–3276.

O'Brien, J.A., Holt, M., Whiteside, G., Lummis, S.C., Hastings, M.H., 2001. Modifications to the
hand-held Gene Gun: improvements for in vitro biolistic transfection of organotypic
neuronal tissue. J. Neurosci. Methods 112, 57–64.

O'Brien, J.A., Lummis, S.C.R., 2006. Diolistic labeling of neuronal cultures and intact tissue
using a hand-held gene gun. Nat. Protoc. 1, 1517–1521.

Oikawa, K., Teixeira, L.B.C., Keikhosravi, A., Eliceiri, K.W., McLellan, G.J., 2021. Microstructure
and resident cell-types of the feline optic nerve head resemble that of humans. Exp. Eye
Res. 202, 108315.

Pan, Y.A., Livet, J., Sanes, J.R., Lichtman, J.W., Schier, A.F., 2011. Multicolor Brainbow
imaging in zebrafish. Cold Spring Harb. Protoc. 2011, db.prot5546.

Prada, D., Harris, A., Guidoboni, G., Siesky, B., Huang, A.M., Arciero, J., 2016. Autoregulation
and neurovascular coupling in the optic nerve head. Surv. Ophthalmol. 61, 164–186.

Richier, B., Salecker, I., 2015. Versatile genetic paintbrushes: Brainbow technologies. Wiley
Interdiscip. Rev. Dev. Biol. 4, 161–180.

Santina, L.D., Ou, Y., 2018. Biolistic Labeling of Retinal Ganglion Cells. Methods Mol. Biol.
1695, 161–170.

Schneider, M., Fuchshofer, R., 2016. The role of astrocytes in optic nerve head fibrosis in
glaucoma. Exp. Eye Res. 142, 49–55.

Seabold, G.K., Daunais, J.B., Rau, A., Grant, K.A., Alvarez, V.A., 2010. DiOLISTIC labeling of
neurons from rodent and non-human primate brain slices. J. Vis. Exp.
https://doi.org/10.3791/2081

Shinozaki, Y., Koizumi, S., 2021. Potential roles of astrocytes and Müller cells in the

pathogenesis of glaucoma. J. Pharmacol. Sci. 145, 262–267.

Sigal, I.A., Ethier, C.R., 2009. Biomechanics of the optic nerve head. Exp. Eye Res. 88, 799–

807.

Staffend, N.A., Meisel, R.L., 2011. DiOlistic labeling in fixed brain slices: phenotype,

morphology, and dendritic spines. Curr. Protoc. Neurosci. Chapter 2, Unit 2.13.

Stokum, J.A., Kwon, M.S., Woo, S.K., Tsymbalyuk, O., Vennekens, R., Gerzanich, V., Simard,

J.M., 2018. SUR1-TRPM4 and AQP4 form a heteromultimeric complex that amplifies

ion/water osmotic coupling and drives astrocyte swelling. Glia 66, 108–125.

Sun, D., 2018. Visualizing Astrocytes of the Optic Nerve. Glaucoma.

https://doi.org/10.1007/978-1-4939-7407-8_18

Sun, D., Jakobs, T.C., 2012. Structural remodeling of astrocytes in the injured CNS.

Neuroscientist 18, 567–588.

Sun, D., Lye-Barthel, M., Masland, R.H., Jakobs, T.C., 2010. Structural remodeling of fibrous

astrocytes after axonal injury. J. Neurosci. 30, 14008–14019.

Sun, D., Lye-Barthel, M., Masland, R.H., Jakobs, T.C., 2009. The morphology and spatial

arrangement of astrocytes in the optic nerve head of the mouse. J. Comp. Neurol. 516, 1–

19.

Sun, D., Moore, S., Jakobs, T.C., 2017. Optic nerve astrocyte reactivity protects function in

experimental glaucoma and other nerve injuries. J. Exp. Med. 214, 1411–1430.

Wang, R., Seifert, P., Jakobs, T.C., 2017. Astrocytes in the Optic Nerve Head of Glaucomatous

Mice Display a Characteristic Reactive Phenotype. Invest. Ophthalmol. Vis. Sci. 58, 924–

932.

5. Figures

Figure 1) DiOlistically-labeled cells with astrocyte morphologies are GFAP+

Figure 1) DiOlistically-labeled cells with astrocyte morphologies were GFAP$^+$. **A**) GFAP-
immunolabeled vibratome slice through the ONH. Astrocytes and their GFAP$^+$ intermediate
filaments densely populate neural tissue pores and line extracellular matrix beams in the LC. **B**)

Detail of the white box in **A** shows the complex, filamentous, and tightly packed arrangement of
GFAP filaments within the LC. IHC of GFAP reveals a limited amount of astrocyte morphology,
labeling only a portion of the cell. This, coupled with the complexity, scale, and density of GFAP$^+$
filaments renders questions about individual astrocyte morphologies difficult to investigate. **C**)
Fluorescent cell membrane tracers, such as DiD, can be introduced into individual cells through
single-color DiOlistic labeling. This revealed the morphology of individual GFAP$^+$ astrocytes
within the context of surrounding tissues. **D**) Images from C shown with LUTs adjusted to improve
visibility of GFAP colocalization in regions with strong DiOlistics signal and of fine astrocyte
branches with less intense DiOlistics signal. **E**) Astrocyte within the white box in **D** with DiD,
GFAP, and DAPI channels shown separately and merged. Yellow arrows indicate location of the
DiOlistically-labeled cell's nucleus. Representative images were from goat LC. Images are
maximum intensity projections of confocal Z stacks. **A**: 20x, **B-E**: 40x.

Figure 2) Multicolor DiOlistic labeling of the lamina cribrosa: workflow

Figure 2) Multicolor DiOlistic labeling of the lamina cribrosa: workflow. **A)** Optic nerve heads from

goat, sheep, pig, and monkey eyes were sectioned coronally at 150µm-thickness to collect slices

containing LC (red). **B)** Gold microcarriers 1µm in diameter were coated with all combinations of

3 fluorescent cell membrane dyes, DiD, DiO, and DiI, to create a total of 7 distinct color channels.

C) A gene gun was used to propel groups of dye-coated microcarriers into vibratome slices

containing LC. **D)** Labeled tissues were imaged with confocal microscopy, and/or second

harmonic generation microscopy.

Figure 3) Multicolor DiOlistic labeling allows individual LC astrocyte visualization across seven

distinct color channels

Figure 3) Individual LC astrocytes of the goat labeled with MuDi at 40x magnification. All

combinations of the fluorescent dyes, DiD, DiO, and DiI, allowed cells to be visualized in distinct

color channels depending upon the dye(s) they received. Images are maximum intensity

projections of confocal stacks. DiD: red, DiO: green, DiI: blue. Scale bar: 25 µm.

Figure 4) Multicolor DiOlistic labeling reveals 3D morphologies of individual astrocytes across the LC

Figure 4) MuDi reveals 3D morphologies of individual astrocytes across the LC. **A**) A coronal section through the goat LC labeled with MuDi. Dye-coated microcarriers labeled hundreds of cells across the sample. Image collected at 10x magnification. **B**) High magnification of the white box within **A** reveals fine details of individual cells within LC pores. DAPI-labeled nuclei and autofluorescence from LC beams (left). Collagen imaged via SHG overlaid verifies the location and appearance of LC beams and neural tissue pores (middle). The MuDi channels are shown alone (right). **C**) Detail of the astrocyte within the white box in **B**. 3D segmentation of the same cell is shown at anterior, sagittal, and posterior viewpoints. Confocal and SHG images are shown as maximum intensity projections. DiD + DiO + DiI = MuDi

784   **Figure 5**) 3D astrocyte models reflect respective confocal images.

785
Figure 5) 3D astrocyte models reflect respective confocal images. Example goat astrocytes in
A-C are shown as 3D models (left), model outlines (yellow) overlaid on their respective confocal
images (middle), and as confocal images alone (right). Confocal images here were each
pseudocolored green to facilitate uniform visualization.

Figure 6) 3D astrocyte segmentations allow extraction of morphological features for quantitative

analysis

Figure 6) 3D astrocyte segmentations allow extraction of morphological features for quantitative

analysis. Example astrocyte 3D segmentations (shown in white) were used for automated

extraction of morphological features including astrocyte branch length, thickness, hierarchy, and

straightness. These morphological features were readily visualized on their respective 3D models

(colormaps) and extracted for future statistical analysis at a cell-by-cell and branch-by-branch

level (histograms). Example cells were of goat origin.

 Figure 7) Astrocyte models allow 3D Sholl analysis

Figure 7) Astrocyte models allow 3D Sholl analysis. **A)** Three representative goat astrocytes

were analyzed in 3D. Colored points indicate branch crossings with Sholl spheres created at 10

μm increments from the astrocyte soma center-point. **B)** Crossing counts of the three example

cells and the average. **C)** Astrocyte branch crossings at Sholl spheres in mouse eyes from Lye-

Barthel et al. 2013.(Lye-Barthel et al., 2013) The crossing counts were higher in the collagenous

LC, although the pattern of distribution was remarkably consistent with that of the control mouse.

Figure 8) Labeling and visualization of LC astrocytes are effective across species.

Figure 8) Labeling and visualization of LC astrocytes are effective across species. LC astrocytes
were successfully MuDi labeled and visualized within the goat **(A, E)**, monkey **(B, F)**, pig **(C, G)**,
and sheep **(D, H)** LC. Representative example images demonstrate labeling of the soma, fine
processes, and astrocyte end-feet. Detail within the white boxes of **A-D and E-H** is shown in **A'-**
D' and E'-H'. Sheep and pig samples in these images were counterstained with DAPI. The
grayscale channel in goat and monkey images show tissue autofluorescence only. Scale bars:
25 μm.

Figure 9) DiOlistic labeling is compatible with downstream labeling and imaging of other

structures

Figure 9) DiOlistic labeling is compatible with downstream labeling and imaging of other

structures. **A**) A slice through the monkey ONH containing collagenous LC labeled with MuDi.

The gray 405nm channel shows tissue autofluorescence. **B**) The same sample was imaged via

second harmonic generation (SHG) microscopy to visualize collagen. **C**) Cell membrane dyes

were washed from the sample with ethanol prior to re-imaging. The sample was labeled with

GFAP and tomato lectin to visualize astrocyte intermediate filaments and blood vessels. Signal

from GFAP, lectin, and tissue autofluorescence are shown independently in **D**, **E**, and **F**,

respectively. LUTs in **E** and **F** were adjusted to aid visualization. Example tissues were of

monkey origin.

Figure 10) Multicolor DiOlistic labeling can reveal cells with fibroblast-like morphologies.

Figure 10: Multicolor DiOlistic labeling can reveal cells with fibroblast-like morphologies. **A)**
MuDi-labeled sheep LC with extracellular matrix autofluorescence shown in grayscale. **B)** The
same image as above, showing MuDi signal only. White boxes within **A** and **B** indicate regions
containing presumptive fibroblasts along collagen beams and in the peripapillary sclera. **C)**
Detail of MuDi-labeled cells with fibroblast-like morphologies (arrows).

Figure 11) Multicolor DiOlistic labeling can reveal axons

Figure 11) DiOlistic labeling can reveal axons. In tissues anterior **(A)** and posterior **(B)** to the LC,
DiOlistically-labeled RGC axons can be seen running largely parallel to each other. Axons are
most readily labeled and visualized in longitudinal sections **(C)**, where single axons occupy the
greatest amount of area, increasing the probability of DiOlistic axon labeling and dye diffusion
across greater length of the axon. Samples in **A** and **B** were from coronal slices of monkey,
labeled with MuDi. Grayscale in **A** and **B** shows tissue autofluorescence from the 405nm channel
only. The longitudinal slice in **C** was from goat, labeled with single-color DiI DiOlistics and
counterstained in DAPI. Although axons are more readily visualized in these samples than in
coronal slices through the LC, not all structures labeled via DiOlistics here are axons. For
example, the coronally-oriented DiI-labeled branches in **C2** may belong to LC astrocytes. NFL:
nerve fiber layer, PL: prelamina, LC: lamina cribrosa, RLR: retrolaminar region, ON: optic nerve.

Supplemental Materials

Supplemental Figure 1) Visualization of microcarrier distribution

Supplemental Figure 1) Visualization of microcarrier distribution. An image of the region shown
in **Figure 4B** (left) is provided as a reference for the respective brightfield image (middle). The
brightfield image shows boundaries of the same collagen beams and neural tissue pores. Detail
of the region within the black box in the middle panel shows bullets from multicolor DiOlistics
visible as dark puncta approximately 1µm in diameter. Optimal labeling density for revealing a
large number of cells while minimizing spatial overlap among those in the same color channel
was approximately 140 cells per mm^2 of canal section area. SHG = second harmonic generation.

Supplemental Figure 2) Dye leakage obscures visualization of morphology.

Supplemental Figure 2: Dye leakage obscures visualization of morphology. Example images
include tissues labeled with MuDi in which individual astrocyte morphologies were not
distinguishable. Dyes can diffuse from individual labeled cells into surrounding areas if cell
membrane integrity is sufficiently compromised. Diffuse dye-labeled regions with multiple nuclei
and without clear astrocyte morphologies, apparent in **A**, were not considered astrocytes and
were excluded from morphological analysis. Microcarriers can be aberrantly delivered in clumps
if not properly sonicated, filtered, or stored with desiccant. Delivery of microcarrier clumps can
result in large patches of tissue being labeled with the same dye (such as the green region in **B**)
and potential lysis of cells bombarded with microcarrier clumps larger than 1µm in diameter.
Some dye diffusion was confined to individual pores, as in **C**, labeling parts of several cells within
a pore. Other dye spread, particularly in samples treated with fixative that contained methanol or
labeled at pressures higher than 200 PSI, was more diffuse across the sample, such as the blue
and red signal in **B**.

Videos:

Video 1:

Example 3D DiOlistics confocal volume information shown in relation to a 3D astrocyte model.

Video 2:

Astrocyte models from **Figures 5**-**7** shown in 3D.