Research Articles

Abrupt shifts in the plant species richness and community composition along an aridity

gradient in the Mongolian grasslands

Naohiro I. Ishii[1][*], Issei Nishimura[1], Yulan Qi[1], Gantsetseg Batdelger[2], Maiko Kagami[1], Gaku

Takimoto[3], Takehiro Sasaki[1]

[1]Graduate School of Environment and Information Sciences, Yokohama National University,

79-7 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

[2]Information and Research Institute of Meteorology, Hydrology, and Environment (IRIMHE)

of Mongolia, Ulaanbaatar, 15160, Mongolia

[3]Graduate School of Agricultural and Life Sciences, University of Tokyo, 1-1-1 Yayoi, Bunkyo-

ku, Tokyo 113-8657, Japan

[*]**Correspondence to:**

Naohiro I. Ishii

Graduate School of Environment and Information Sciences, Yokohama National University, 79-

7 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

E-mail: na0hiro1shi1@gmail.com

Highlights

•   Patterns in plant species richness and composition in Mongolian steppe were studied

•   Shifts in species richness and community composition were nonlinear along an aridity

gradient abrupt at high aridity

•   Nonlinear shift indicates the vulnerability of plant communities to changes in highly

arid regions

Abstract

Aridification is a major facet of climate changes in drylands. However, little is known about the

shifts in the plant community composition to aridity. Dryland grasslands are widely used for

livestock grazing. Although grazing impacts can be relatively small in arid regions, overgrazing

has great impacts on vegetation, regardless of aridity. Thus, investigations under long-term

grazing exclusion are suitable to clarify the response of composition to aridity without

confounding effect of grazing. We investigated the shifting patterns in species richness and

composition with grazing exclosure along a broad range of aridity (0.6–0.95) in Mongolia. The

shifts in the species richness and composition (the axis of NMDS) were investigated by model

selection for GLMs, including the linear and quadratic terms of aridity, and soil environments.

The best models included both the linear and quadratic terms of aridity, which showed nonlinear

and unimodal responses to aridity. Particularly, their shifts were relatively abrupt at high aridity,

which might be attributed to environmental filtering by aridity and competitive exclusion by

aridity specialists. Our study firstly demonstrated the nonlinear shift of plant community

composition to aridity, indicating the vulnerability of plant communities to press changes of

aridity, especially in highly arid regions.

Keywords

Aridity, community composition, dissimilarity, Mongolia, species richness, steppe.

1. Introduction

Shifts in community composition along climatic gradients can help predict the impacts of climate change on an ecosystem's structure and functioning (Chapin et al., 2000). Changes in species richness and community composition caused by environmental changes also have substantial effects (Sasaki et al., 2022; Spaak et al., 2017; Wang and Ali, 2021). Aridification, which increases aridity (calculated as [1 – precipitation/potential evapotranspiration]), is one of the major facets of ongoing climate change in terrestrial ecosystems (Huang et al., 2016). Drylands are particularly vulnerable to aridification (Huang et al., 2016), resulting in shifts in multiple structural and functional attributes in the communities and ecosystems (Berdugo et al., 2022, 2020; Maestre et al., 2016). (Berdugo et al., 2020) reported abrupt changes in vegetation cover and plant species richness along aridity gradients (Berdugo et al., 2020). However, little is known about the shift patterns in the plant community composition along an aridity gradient. Previous studies have suggested that shifts in the composition of soil bacterial and nematode communities in China exhibit nonlinear responses to changing aridity gradients (Pan et al., 2022; Xiong et al., 2020). Understanding whether the shifting pattern in the plant community composition along an aridity gradient is consistent with that in other taxa can provide implications for the impacts of aridity shifts under future climate change conditions on an ecosystem's functioning and inter-trophic interactions.

Grassland vegetation is widely used for livestock grazing to support secondary

production in drylands. Numerous studies have revealed the impact of livestock grazing on

plant community composition mediated by selective herbivory, depending on the livestock

species (Díaz et al., 2007; Li et al., 2022; Sasaki et al., 2008a). The grazing impacts on plant

species richness and community composition differed according to changes in water availability

among the sites. Previous studies have demonstrated that the relative magnitude of grazing

impacts on the vegetation is relatively low in high-aridity regions (Herrero-Jáuregui and

Oesterheld, 2018; Price et al., 2022). However, species richness and composition of plant

communities can drastically be altered and homogenized by overgrazing, regardless of aridity

(Ahlborn et al., 2020; Sasaki et al., 2008a, 2009; Wang et al., 2017). Thus, the spatial patterns

of the plant community composition in grazed grasslands can be confounded by grazing impacts

across a broad range of aridity levels. To infer how future changes in the plant community

composition will be driven by aridity, investigations of plant communities under long-term

grazing exclusion are essential. Notably, shifts in the plant community composition along a

wide range of aridity gradients remain poorly understood in the Eurasian steppe.

Here, we investigated the shifts in species richness and composition of plant

communities in the long-term grazing exclosure along an aridity gradient in Mongolian

grasslands. Mongolia covers a vast area of the eastern part of the Eurasian Steppe. In Mongolia,

the climate is spatially characterized by broad variations in temperature and precipitation

(Vandandorj et al., 2017). In this study, we aimed to determine whether the response patterns of

the plant species richness and community composition to aridity are linear or convex to provide

insights into the vulnerability of plant communities to future aridification.

2. Materials and methods

2.1. Study sites and survey design

Across central Mongolia, ranging from semi-arid to arid regions, over 600 km, 12 study

sites were selected at the meteorological sites of the Mongolian Information and Research

Institute of Meteorology, Hydrology, and Environment (IRIMHE) (Figure 1). We aimed to

ensure that the study area covered a range of vegetation types within the rangeland ecosystems

of Mongolia. Central Mongolia has the typical climate of Central Asia, which is characterized

by hot and short summers with approximately 70 % annual precipitation and cold winters,

including extreme climate events such as drought. Based on the climate data between 1978 and

2017 provided by IRIMHE, the mean annual precipitation ranged from 97 mm at Tsogt-Ovoo

(site 11) to 333 mm at Darkhan (site 05); the mean annual temperature ranged from -0.8 ℃ at

Ugtaal (site 06) to 7.6 ℃ at Khanbogd (site 12). Thus, there is a wide gradient of aridity from

north to south (Figure 1) that supports three main vegetation types: forest steppe, steppe, and

desert steppe (Table 1).

We set three 5 × 5 m blocks, which included three 1 × 1 m plots, at each meteorological

site (50 × 50 m), to observe the heterogeneity at both block- and site-level scales. All the sites

were enclosed for at least 50 years to avoid livestock grazing. In each plot, vegetation was

sampled by identifying all the species present from July to August 2022, and the vegetation data

were pooled for each site.

2.2. Environmental variables

Climatic and soil environmental data were collected across the study sites. The aridity,

calculated as [1 – precipitation/potential evapotranspiration], was obtained from the Global

Aridity Index and Potential Evapotranspiration Climate Database v3 (Trabucco and Zomer,

2022). Since the soil environmental variables can explain the large-scale patterns of the plant

community composition across the Mongolian grasslands (Sasaki et al., 2008b), we derived the

soil environmental variables based on information from SoilGrids at a resolution of 250 m

(Hengl et al., 2017): N, total nitrogen content in cg/kg; SOC, soil organic carbon content in

dg/kg; pH, soil pH in H_2O; clay content mass fraction in %; sand, sand content mass fraction

in %; silt, silt content mass fraction in %. Because of the high correlations between aridity and

N, SOC, and pH, and between the sand and silt content (Figure S1), we used the aridity, clay,

and silt content as the environmental variables in the subsequent analysis. The former

correlations, which resulted from the long-term feedback between plant biomass production and

edaphic variables, were consistent with the patterns observed in previous studies (Berdugo et

al., 2020; Delgado-Baquerizo et al., 2013; Moreno-Jiménez et al., 2019).

2.3. Data analysis

The species richness and community composition were quantified based on a species-by-

site matrix. To compare the community composition among the different sites, nonmetric multi-

dimensional scaling (NMDS) based on Sorensen dissimilarity was conducted using the function

'metaMDS' of the R package 'vegan.' Sorensen dissimilarity was used to focus on the shift in

community composition via species replacement and changes in species richness rather than

changes in abundance. In this study, the dimensional reduction to one dimension was performed

because the stress value (0.077) was lower than 0.2 (Kruskal, 1964).

To investigate the linear and quadratic relationships between the species richness,

community composition (axis of NMDS), and environmental variables (aridity, clay, and silt

content), we performed a generalized linear model (GLM) analysis. In the species richness

models, a Poisson distribution with a log link was used. A Gaussian GLM was used in the

models for community composition. The model selection was based on the Akaike Information

Criterion (AIC) using the function 'dredge' of the R package 'MuMIn.'

3. Results

As a result of the model selection for species richness and community composition (the axis

of NMDS), both the best models with the lowest AIC included the linear and quadratic terms

of aridity (Table 2 and Table S2); this indicated that aridity induces nonlinear responses in

species richness and community composition and relatively abrupt shifts in high aridity levels

(Figure 2). In both of the best models, which included edaphic variables, the linear and quadratic

terms of aridity were significant (Table 2). The best model for species richness included linear

or quadratic terms of clay and silt content, and the coefficients were not significant. In the best

models for community composition, the quadratic term of silt content was significant, which

showed a unimodal relationship, but a lower effect size than that of aridity. Visual inspection of

the reductions in species richness and the axis of NMDS with aridity showed that species

richness and composition changed only slightly at low aridity, but abruptly at high aridity

(Figure 2).

4. Discussion

In this study, we first demonstrated nonlinear shifts in species richness and community

composition of exclosed plant communities along the aridity gradient across the Mongolian

steppes. Notably, the decline in species richness and the shift in community composition along

the broad spatial gradient of aridity were more abrupt at high aridity levels (aridity > 0.8)

(Figure 2). Previous research has shown that aridity is a critical determinant of community

composition throughout the entire area of grazed rangelands in Mongolia (Suzuki et al., 2022).

Our study is novel because we demonstrated a nonlinear response in plant community

composition to spatial changes in aridity, by eliminating the confounding effects of grazing.

The nonlinear shift in the plant community composition along the aridity gradient was

mainly attributed to the relatively high level of environmental filtering as a result of abiotic and

biotic factors in the high-aridity regions. In addition to the ungrazed plant communities

investigated in our study, a larger variation in plant species composition was observed in the

desert steppe and the transition zone between the desert steppe and steppe compared to the

steppe and forest steppe across the grazed rangelands in Mongolia (Ahlborn et al., 2020). The

relatively high level of variation in community composition of both the ungrazed and grazed

communities, under high aridity, might be a result of an intensifying reduction in species

richness and species turnover.

Because of environmental filtering by aridity, the dominance of stress-tolerant species

and exclusion of non-tolerant species can often be observed in arid and semi-arid ecosystems

with high aridity (Berdugo et al., 2019; Grime, 1977; Le Bagousse-Pinguet et al., 2017). A

global meta-analysis demonstrated the nonlinear and threshold responses of plant species

richness to aridity, and the relatively steep decrease at aridity > 0.78 (Berdugo et al., 2020),

which was roughly consistent with the extreme decline in species richness at aridity > 0.8 in

this study (Figure 2a). This result repeatedly indicates that aridity is an important determinant

of community composition in high-aridity regions. In addition, previous studies have

demonstrated that high aridity contributes to the severe limitations of not only water

unavailability but also nutrient unavailability via long-term feedback from the vegetation (e.g.,

nitrogen, carbon, micronutrients, and soil pH) (Berdugo et al., 2020; Delgado-Baquerizo et al.,

2013; Moreno-Jiménez et al., 2019), resulting in severe environmental filtering by abiotic

factors indirectly associated with aridity.

The relatively abrupt shift in community composition under high aridity conditions

may also be attributed to species turnover via competitive exclusion by highly adapted species

in arid environments (Berdugo et al., 2019; Grime, 1977; Macarthur and Levins, 1967).

Contrary to the stress gradient hypothesis (Bertness and Callaway, 1994), increasing aridity can

lead to the dominance of aridity specialists and an increase in the relative importance of

competitive exclusion by the superior specialists, rather than facilitation (Berdugo et al., 2019).

Previous studies have also indicated that negative effects on water availability by aridity

specialists (e.g., shrub species) can cause competitive exclusion in dryland ecosystems

(Butterfield et al., 2016; O'Brien et al., 2017). The decrease in niche breadth due to aridity

might cause ecological divergence (limiting similarity) in functional strategies to minimize

fitness differences and mediate interspecific competition at local scales (Macarthur and Levins,

1967), although ecological convergence is pronounced with an increase in aridity at regional

scales (Le Bagousse-Pinguet et al., 2017).

Across the study sites with high aridity (sites 09–12) in this study, one or two shrub

species were dominant or predominant in each site (e.g., *Zygophyllum kaschgaricum*, *Potaninia*

mongolica, *Anabasis brevifolia*, and *Salsola passerina*; Table 1), and the turnover of these shrub

species among the sites was relatively large. Additionally, the spatial distributions of grass and

forb species were allopatric to the shrub species in each plot across the sites (N.I. Ishii, personal

observation). Although we could not empirically evaluate the impact of competition and

facilitation on the community composition, the limiting similarity among the shrub species and

the absence of facilitative effects by the same on composition were partially explained in the

context of the previous paragraph.

The present study demonstrated the relatively abrupt response of the plant species

richness and community composition in the high aridity levels across Mongolia, indicating the

vulnerability of the plant communities to changes in aridity due to future climate change,

especially in a desert steppe. Although accurate threshold detection for species richness and

community composition along the aridity gradient (Berdugo et al., 2020) could not be

performed in this study because of the small sample size of the study sites, future studies should

use continuous and intensive vegetation sampling across the whole area of the Eurasian steppe.

Although space-for-time substitution is a promising method for predicting community

responses to climate change from snapshots of spatial patterns (Blois et al., 2013; Fukami and

Wardle, 2005), the spatial and temporal shifts of community and functional compositions to

environmental changes do not necessarily coincide (Sandel et al., 2010). Therefore, long-term

monitoring of the plant community composition across large environmental gradients is crucial

to validate the results based on spatial data such as ours.

219   While the relative magnitude of grazing impacts on vegetation is lower in highly arid

regions (Herrero-Jáuregui and Oesterheld, 2018; Price et al., 2022), species richness and

composition of plant communities can be drastically altered and homogenized by overgrazing

regardless of the climatic context (Ahlborn et al., 2020; Sasaki et al., 2009, 2008a; Wang et al.,

2017). Recently, livestock numbers have rapidly increased across the Eurasian steppe

(Kakinuma et al., 2019), leading to the expansion of overgrazing impacts on vegetation.

Therefore, comparisons of species richness and community composition between grazed and

ungrazed communities will become increasingly important to clarify the pure effects of aridity

and the interactive effects of aridity and grazing on vegetation. Assessing the differentiation of

grazed communities from communities with grazing exclosures as a 'reference community' will

be a powerful tool for predicting the response of not only community composition, but also

ecosystem functioning and services in drylands, to the interaction of aridification and grazing

(Maestre et al., 2022).

Ethics approval

Not applicable.

Consent to participate

All the authors approved participation in this study.

Consent for publication

All the authors approved the publication of this study.

Data availability

Data are available from the corresponding author upon reasonable request.

Code availability

Codes are available from the corresponding author upon reasonable request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have influenced the work reported in this study.

Author Contributions

N.I.I., G.B., G.T., M.K., and T.S. designed the study. All the authors contributed to the

field surveys. N.I.I. performed the analyses and wrote the first draft of the manuscript. All the

authors have contributed to the revisions.

Acknowledgments

We would like to thank our laboratory colleagues for their help with the analyses.

Funding

This work was financially supported by a fund for Fostering Joint International

Research A (no. 19KK0393) and a grant-in-aid for Scientific Research B (no. 22H03791)

presented to T.S. from the Ministry of Education, Culture, Sports, Science, and Technology of

Japan.

References

Ahlborn, J., von Wehrden, H., Lang, B., 2020. Climate–grazing interactions in Mongolian rangelands: Effects of grazing change along a large-scale environmental gradient. journal of arid 173, 104043.

Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán, J.J., Gross, N., Saiz, H., Maire, V., Lehman, A., Rillig, M.C., Solé, R.V., Maestre, F.T., 2020. Global ecosystem thresholds driven by aridity. Science 367, 787–790.

Berdugo, M., Maestre, F.T., Kéfi, S., Gross, N., Le Bagousse-Pinguet, Y., Soliveres, S., 2019. Aridity preferences alter the relative importance of abiotic and biotic drivers on plant species abundance in global drylands. J. Ecol. 107, 190–202.

Berdugo, M., Vidiella, B., Solé, R.V., Maestre, F.T., 2022. Ecological mechanisms underlying aridity thresholds in global drylands. Funct. Ecol. 36, 4–23.

Bertness, M.D., Callaway, R., 1994. Positive interactions in communities. Trends Ecol. Evol. 9, 191–193.

Blois, J.L., Williams, J.W., Fitzpatrick, M.C., Jackson, S.T., Ferrier, S., 2013. Space can substitute for time in predicting climate-change effects on biodiversity. Proc. Natl. Acad. Sci. U. S. A. 110, 9374–9379.

283   Butterfield, B.J., Bradford, J.B., Armas, C., Prieto, I., Pugnaire, F.I., 2016. Does the stress-

284   gradient hypothesis hold water? Disentangling spatial and temporal variation in plant

285   effects on soil moisture in dryland systems. Funct. Ecol. 30, 10–19.

286   Chapin, F.S., 3rd, Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L.,

287   Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C., Díaz, S., 2000.

288   Consequences of changing biodiversity. Nature 405, 234–242.

289   Delgado-Baquerizo, M., Maestre, F.T., Gallardo, A., Bowker, M.A., Wallenstein, M.D., Quero,

290   J.L., Ochoa, V., Gozalo, B., García-Gómez, M., Soliveres, S., García-Palacios, P.,

291   Berdugo, M., Valencia, E., Escolar, C., Arredondo, T., Barraza-Zepeda, C., Bran, D.,

292   Carreira, J.A., Chaieb, M., Conceição, A.A., Derak, M., Eldridge, D.J., Escudero, A.,

293   Espinosa, C.I., Gaitán, J., Gatica, M.G., Gómez-González, S., Guzman, E., Gutiérrez,

294   J.R., Florentino, A., Hepper, E., Hernández, R.M., Huber-Sannwald, E., Jankju, M., Liu,

295   J., Mau, R.L., Miriti, M., Monerris, J., Naseri, K., Noumi, Z., Polo, V., Prina, A., Pucheta,

296   E., Ramírez, E., Ramírez-Collantes, D.A., Romão, R., Tighe, M., Torres, D., Torres-

297   Díaz, C., Ungar, E.D., Val, J., Wamiti, W., Wang, D., Zaady, E., 2013. Decoupling of

298   soil nutrient cycles as a function of aridity in global drylands. Nature 502, 672–676.

299   Díaz, S., Lavorel, S., McINTYRE, S.U.E., Falczuk, V., Casanoves, F., Milchunas, D.G., Skarpe,

300   C., Rusch, G., Sternberg, M., Noy-Meir, I., Landsberg, J., Zhang, W., Clark, H.,

Campbell, B.D., 2007. Plant trait responses to grazing ? a global synthesis. Glob. Chang.

Biol. 13, 313–341.

Fukami, T., Wardle, D.A., 2005. Long-term ecological dynamics: Reciprocal insights from

natural and anthropogenic gradients. Proceedings of the Royal Society B: Biological

Sciences 272, 2105–2115.

Grime, J.P., 1977. Evidence for the existence of three primary strategies in plants and its

relevance to ecological and evolutionary theory. Am. Nat.

Hengl, T., Mendes de Jesus, J., Heuvelink, G.B.M., Ruiperez Gonzalez, M., Kilibarda, M.,

Blagotić, A., Shangguan, W., Wright, M.N., Geng, X., Bauer-Marschallinger, B.,

Guevara, M.A., Vargas, R., MacMillan, R.A., Batjes, N.H., Leenaars, J.G.B., Ribeiro,

E., Wheeler, I., Mantel, S., Kempen, B., 2017. SoilGrids250m: Global gridded soil

information based on machine learning. PLoS One 12, e0169748.

Herrero-Jáuregui, C., Oesterheld, M., 2018. Effects of grazing intensity on plant richness and

diversity: a meta-analysis. Oikos 127, 757–766.

Huang, J., Yu, H., Guan, X., Wang, G., Guo, R., 2016. Accelerated dryland expansion under

climate change. Nat. Clim. Chang. 6, 166–171.

Kakinuma, K., Yanagawa, A., Sasaki, T., Rao, M.P., Kanae, S., 2019. Socio-ecological interactions in a changing climate: A review of the Mongolian pastoral system. Sustain. Sci. Pract. Policy 11, 5883.

Kruskal, J.B., 1964. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. Psychometrika 29, 1–27.

Le Bagousse-Pinguet, Y., Gross, N., Maestre, F.T., Maire, V., de Bello, F., Fonseca, C.R., Kattge, J., Valencia, E., Leps, J., Liancourt, P., 2017. Testing the environmental filtering concept in global drylands. J. Ecol. 105, 1058–1069.

Li, Y., Shi, C., Zhang, T., Wang, H., Yang, Z., Tang, S., Li, F.Y., 2022. Summer grazing by three livestock species at moderate intensity enhances primary productivity and drives community divergence in a semi‐arid steppe. Appl. Veg. Sci. 25. https://doi.org/10.1111/avsc.12683

Macarthur, R., Levins, R., 1967. The Limiting Similarity, Convergence, and Divergence of Coexisting Species. Am. Nat. 101, 377–385.

Maestre, F.T., Bagousse-Pinguet, Y.L., Delgado-Baquerizo, M., Eldridge, D.J., Saiz, H., Berdugo, M., Gozalo, B., Ochoa, V., Guirado, E., García-Gómez, M., Valencia, E., Gaitán, J.J., Asensio, S., Mendoza, B.J., Plaza, C., Díaz-Martínez, P., Rey, A., Hu, H.-W., He, J.-Z., Wang, J.-T., Lehmann, A., Rillig, M.C., Cesarz, S., Eisenhauer, N.,

Martínez-Valderrama, J., Moreno-Jiménez, E., Sala, O., Abedi, M., Ahmadian, N.,

Alados, C.L., Aramayo, V., Amghar, F., Arredondo, T., Ahumada, R.J., Bahalkeh, K.,

Salem, F.B., Blaum, N., Boldgiv, B., Bowker, M.A., Bran, D., Bu, C., Canessa, R.,

Castillo-Monroy, A.P., Castro, H., Castro, I., Castro-Quezada, P., Chibani, R., Conceição,

A.A., Currier, C.M., Darrouzet-Nardi, A., Deák, B., Donoso, D.A., Dougill, A.J., Durán,

J., Erdenetsetseg, B., Espinosa, C.I., Fajardo, A., Farzam, M., Ferrante, D., Frank,

A.S.K., Fraser, L.H., Gherardi, L.A., Greenville, A.C., Guerra, C.A., Gusmán-

Montalvan, E., Hernández-Hernández, R.M., Hölzel, N., Huber-Sannwald, E., Hughes,

F.M., Jadán-Maza, O., Jeltsch, F., Jentsch, A., Kaseke, K.F., Köbel, M., Koopman, J.E.,

Leder, C.V., Linstädter, A., le Roux, P.C., Li, X., Liancourt, P., Liu, J., Louw, M.A.,

Maggs-Kölling, G., Makhalanyane, T.P., Issa, O.M., Manzaneda, A.J., Marais, E., Mora,

J.P., Moreno, G., Munson, S.M., Nunes, A., Oliva, G., Oñatibia, G.R., Peter, G., Pivari,

M.O.D., Pueyo, Y., Quiroga, R.E., Rahmanian, S., Reed, S.C., Rey, P.J., Richard, B.,

Rodríguez, A., Rolo, V., Rubalcaba, J.G., Ruppert, J.C., Salah, A., Schuchardt, M.A.,

Spann, S., Stavi, I., Stephens, C.R.A., Swemmer, A.M., Teixido, A.L., Thomas, A.D.,

Throop, H.L., Tielbörger, K., Travers, S., Val, J., Valkó, O., van den Brink, L., Ayuso,

S.V., Velbert, F., Wamiti, W., Wang, D., Wang, L., Wardle, G.M., Yahdjian, L., Zaady,

E., Zhang, Y., Zhou, X., Singh, B.K., Gross, N., 2022. Grazing and ecosystem service

delivery in global drylands. Science 378, 915–920.

Maestre, F.T., Eldridge, D.J., Soliveres, S., Kéfi, S., Delgado-Baquerizo, M., Bowker, M.A.,

García-Palacios, P., Gaitán, J., Gallardo, A., Lázaro, R., Berdugo, M., 2016. Structure

and functioning of dryland ecosystems in a changing world. Annu. Rev. Ecol. Evol. Syst.

47, 215–237.

Moreno-Jiménez, E., Plaza, C., Saiz, H., Manzano, R., Flagmeier, M., Maestre, F.T., 2019.

Aridity and reduced soil micronutrient availability in global drylands. Nat Sustain 2,

371–377.

O'Brien, M.J., Pugnaire, F.I., Armas, C., Rodríguez-Echeverría, S., Schöb, C., 2017. The shift

from plant-plant facilitation to competition under severe water deficit is spatially

explicit. Ecol. Evol. 7, 2441–2448.

Pan, H., Gao, H., Peng, Z., Chen, B., Chen, S., Liu, Y., Gu, J., Wei, X., Chen, W., Wei, G., Jiao,

S., 2022. Aridity threshold induces abrupt change of soil abundant and rare bacterial

biogeography in dryland ecosystems. mSystems 7, e0130921.

Price, J.N., Sitters, J., Ohlert, T., Tognetti, P.M., Brown, C.S., Seabloom, E.W., Borer, E.T.,

Prober, S.M., Bakker, E.S., MacDougall, A.S., Yahdjian, L., Gruner, D.S., Olde

Venterink, H., Barrio, I.C., Graff, P., Bagchi, S., Arnillas, C.A., Bakker, J.D., Blumenthal,

D.M., Boughton, E.H., Brudvig, L.A., Bugalho, M.N., Cadotte, M.W., Caldeira, M.C., Dickman, C.R., Donohue, I., Grégory, S., Hautier, Y., Jónsdóttir, I.S., Lannes, L.S., McCulley, R.L., Moore, J.L., Power, S.A., Risch, A.C., Schütz, M., Standish, R., Stevens, C.J., Veen, G.F., Virtanen, R., Wardle, G.M., 2022. Evolutionary history of grazing and resources determine herbivore exclusion effects on plant diversity. Nat Ecol Evol 6, 1290–1298.

Sandel, B., Goldstein, L.J., Kraft, N.J.B., Okie, J.G., Shuldman, M.I., Ackerly, D.D., Cleland, E.E., Suding, K.N., 2010. Contrasting trait responses in plant communities to experimental and geographic variation in precipitation. New Phytol. 188, 565–575.

Sasaki, T., Ishii, N.I., Makishima, D., Sutou, R., Goto, A., Kawai, Y., Taniguchi, H., Okano, K., Matsuo, A., Lochner, A., Cesarz, S., Suyama, Y., Hikosaka, K., Eisenhauer, N., 2022. Plant and microbial community composition jointly determine moorland multifunctionality. J. Ecol. 110, 2507−2521.

Sasaki, T., Okayasu, T., Jamsran, U., Takeuchi, K., 2008a. Threshold changes in vegetation along a grazing gradient in Mongolian rangelands. J. Ecol. 96, 145–154.

Sasaki, T., Okayasu, T., Shirato, Y., Jamsran, U., Okubo, S., Takeuchi, K., 2008b. Can edaphic factors demonstrate landscape-scale differences in vegetation responses to grazing? Plant Ecol. 194, 51–66.

Sasaki, T., Okubo, S., Okayasu, T., Jamsran, U., Ohkuro, T., Takeuchi, K., 2009. Two-phase

functional redundancy in plant communities along a grazing gradient in Mongolian

rangelands. Ecology 90, 2598–2608.

Spaak, J.W., Baert, J.M., Baird, D.J., Eisenhauer, N., Maltby, L., Pomati, F., Radchuk, V., Rohr,

J.R., Van den Brink, P.J., De Laender, F., 2017. Shifts of community composition and

population density substantially affect ecosystem function despite invariant richness.

Ecol. Lett. 20, 1315–1324.

Suzuki, K., Tungalag, R., Narantsetseg, A., Tsendeekhuu, T., Masato, S., Yamanaka, N., Takashi,

K., 2022. Composition, distribution, and environmental drivers of Mongolian rangeland

plant communities. J. plant ecol. https://doi.org/10.1093/jpe/rtac100

Trabucco, A., Zomer, R.J., 2022. Version 3 of the "Global Aridity Index and Potential

Evapotranspiration (ET0) Database": Estimation of Penman-Monteith Reference

Evapotranspiration. figshare. https://doi.org/10.6084/m9.figshare.7504448.v3

Vandandorj, S., Munkhjargal, E., Boldgiv, B., Gantsetseg, B., 2017. Changes in event number

and duration of rain types over Mongolia from 1981 to 2014. Environ. Earth Sci. 76, 1–

12.

Wang, L.-Q., Ali, A., 2021. Climate regulates the functional traits-aboveground biomass

relationships at a community-level in forests: A global meta-analysis. Sci. Total Environ.

761, 143238.

Wang, Y., Heberling, G., Görzen, E., Miehe, G., Seeber, E., Wesche, K., 2017. Combined effects

of livestock grazing and abiotic environment on vegetation and soils of grasslands

across Tibet. Appl. Veg. Sci. 20, 327–339.

Xiong, D., Wei, C., Wubs, E.R.J., Veen, G.F., Liang, W., Wang, X., Li, Q., Putten, W.H., Han,

X., 2020. Nonlinear responses of soil nematode community composition to increasing

aridity. Glob. Ecol. Biogeogr. 29, 117–126.

Tables and figures

Table 1. Locations, aridity, vegetation type, and dominant species of the 12 study sites. Aridity

is calculated as 1 – (precipitation/potential evapotranspiration).

Table 1.

Site	ID	Latitude	Longitude	Aridity	Vegetation	Dominant species
Erdenet	01	49°02'07.08"N	106°56'20.74"E	0.62	Forest steppe	*Elymus chinensis, Koeleria macrantha, Medicago ruthenica*
Hutag	02	49°23'37.32"N	106°50'50.38"E	0.64	Forest steppe	*Artemisia frigida, Stipa krylovii, Medicago ruthenica*
Bulgan	03	48°49'05.52"N	106°54'04.54"E	0.64	Forest steppe	*Agropyron cristatum, Stipa krylovii, Cleistogenes squarrosa*
Tarialan	04	49°36'33.12"N	106°47'56.30"E	0.65	Forest steppe	*Artemisia frigida, Cleistogenes squarrosa, Heteropapus hispidus*
Darkhan	05	49°29'25.44"N	107°03'45.58"E	0.69	Steppe	*Carex pediformis, Artemisia scoparia, Heteropapus hispidus*
Ugtaal	06	48°15'29.16"N	107°01'37.27"E	0.72	Steppe	*Potentilla acaulis, Elymus chinensis, Stipa krylovii*
Zuunmod	07	47°42'47.52"N	107°07'48.46"E	0.72	Steppe	*Artemisia frigida, Stipa krylovii, Arenaria capillaris*
Mandalgobi	08	45°44'35.52"N	107°05'03.43"E	0.87	Desert steppe	*Carex stenophylloides, Stipa glareosa, Allium polyrrhizm*
Dalanzadgad	09	43°34'37.92"N	106°57'40.39"E	0.92	Desert steppe	*Stipa gobica, Artemisia pectinata, Convolvulus amanii*

Saikhan	10	44°05'42.00"N	106°54'09.12"E	0.92	Desert steppe	*Convolvulus amanii, Stipa gobica, Cleistogenes songorica*
Tsogt-Ovoo	11	44°25'23.52"N	107°01'16.39"E	0.93	Desert steppe	*Allium polyrrhizm, Anabasis brevifolia, Salsola passerina*
Khanbogd	12	43°11'44.52"N	107°08'46.46"E	0.94	Desert steppe	*Zygophyllum kaschgaricum, Cleistogenes songorica, Potaninia mongolica*

Table 2. Results of the generalized linear models (GLM) for the environmental variables on

species richness and community composition (the axis of nonmetric multi-dimensional scaling,

NMDS). The full models included the linear and quadratic terms of aridity (shown as aridity

and aridity^2, respectively) and the clay and silt content as the independent variables. The

standardized regression coefficients and the standard error of the best model, based on AIC,

(Figure S2) were shown.

Dependent variable	Independent variable	Estimate	SE	z value	P value
Species richness	Aridity	**-0.422**	0.087	-4.827	<0.001
	Aridity^2	**-0.384**	0.140	-2.751	0.006
	Clay^2	-0.150	0.080	-1.866	0.062
	Silt	-0.230	0.127	-1.807	0.071
	Silt^2	-0.164	0.086	-1.907	0.057
MDS	Aridity	**-0.796**	0.039	-20.168	<0.001
	Aridity^2	**-0.361**	0.043	-8.320	<0.001
	Silt^2	**-0.119**	0.040	-2.988	0.017

429   **Figure 1.** Spatial distribution of the 12 study sites and aridity in Mongolia. Aridity is calculated

430   as 1 – (precipitation/potential evapotranspiration).

433   **Figure 2.** Relationship between the community composition (the axis of nonmetric multi-

434   dimensional scaling) and environmental variables. The regression line and its confidence

435   interval, based on the regression models shown in Table 2, were denoted.

Supplementary information

Table S1. Pairwise correlation matrix of the environmental variables. Aridity, $1 - (\text{precipitation/potential evapotranspiration})$; N, total nitrogen content in cg/kg; SOC, soil organic carbon content in dg/kg; pH, soil pH in H_2O; clay content mass fraction in %; sand, sand content mass fraction in %; silt, silt content mass fraction in %.

	Aridity	N	SOC	pH	Clay	Sand	Silt
Aridity	-	-	-	-	-	-	-
N	-0.988	-	-	-	-	-	-
SOC	-0.971	0.956	-	-	-	-	-
pH	0.758	-0.720	-0.762	-	-	-	-
Clay	-0.443	0.480	0.313	-0.150	-	-	-
Sand	0.716	-0.744	-0.738	0.526	-0.352	-	-
Silt	-0.451	0.456	0.557	-0.441	-0.278	-0.801	-

445   **Table S2.** Model selection of generalized linear models (GLM) for species richness and

446   community composition (the axis of non-metric multidimensional scaling (NMDS)) using the

447   Akaike Information Criterion (AIC). The models included the linear and quadratic terms of

448   aridity and the clay and silt content as the independent variables. The models were shown to

449   have ΔAIC less than 2.

Dependent variable	Model	Rank	df	AIC	ΔAIC
	Aridity+Aridity^2+Clay^2+Silt+Silt^{2h}	1	6	74.57	0.00
	Aridity+Aridity^2+Silt^2	2	4	75.70	1.14
	Aridity+Aridity^2	3	3	75.79	1.23
	Aridity+Aridity^2+Clay^2+Silt^2	4	5	75.87	1.31
	Aridity+Aridity^2+Clay^2+Silt	5	5	76.16	1.59
Species richness	Aridity+Aridity^2+Silt+Silt^2	6	5	76.21	1.64
	Aridity+Aridity^2+Clay+Clay^2+Silt^2	7	6	76.25	1.69
	Aridity+Aridity^2+Clay+Clay^2+Silt+Silt^2	8	7	76.30	1.73
	Aridity+Clay^2	9	3	76.46	1.89
	Aridity+Aridity^2+Clay^2	10	4	76.47	1.90
	Aridity	11	2	76.55	1.98
	Aridity+Aridity^2+Silt^2	1	5	-12.93	0.00
MDS	Aridity+Aridity^2+Clay+Silt^2	2	6	-12.61	0.32
	Aridity+Aridity^2+Silt+Silt^2	3	6	-11.06	1.87
	Aridity+Aridity^2+Clay^2+Silt^2	4	6	-10.98	1.95