Original Article

Difference in determinants and changing patterns of plant community composition with

and without grazing across semi-arid and arid regions in Mongolian steppe

Naohiro I. Ishii[1,2*], Issei Nishimura[1], Yulan Qi[1], Gantsetseg Batdelger[3], Maiko Kagami[1], Gaku

Takimoto[4], Takehiro Sasaki[1]

[1]Graduate School of Environment and Information Sciences, Yokohama National University,

79-7 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

[2]Arid Land Research Center, Tottori University, 1390 Hamasaka, Tottori 680-0001, Japan

[3]Information and Research Institute of Meteorology, Hydrology, and Environment (IRIMHE)

of Mongolia, Ulaanbaatar, 15160, Mongolia

[4]Graduate School of Agricultural and Life Sciences, University of Tokyo, 1-1-1 Yayoi, Bunkyo-

ku, Tokyo 113-8657, Japan

[*]**Correspondence to:**

Naohiro I. Ishii

Graduate School of Environment and Information Sciences, Yokohama National University, 79-

7 Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

E-mail: na0hiro1shi1@gmail.com

Highlights

•   Determinants and changing patterns in plant community composition of Mongolian

steppe along aridity gradients in semi-arid and arid regions were investigated with the

consideration of the effects of aridity and edaphic variables for communities with the

absence/presence of livestock grazing, respectively.

•   Although aridity had significant impacts on community composition regardless of the

regions and the absence/presence of grazing, the difference in response patterns was observed

between the regions, which did not depend on grazing.

•   Regardless of the aridity levels, the effects of soil pH on community composition were

eliminated by grazing.

Abstract

Aridity, edaphic variables related to aridity, and livestock grazing are major drivers of plant

community composition across dryland grasslands. With accounting these factors, little is

known about differences in determinants and changing patterns along aridity gradients with

different average of aridity. Thus, the comparative investigations of communities without and

with grazing in semi-arid and arid regions are suitable to clarify the difference in compositional

responses along aridity gradients between potential vegetations and degraded ones with grazing.

We investigated compositional changes of communities without/with grazing in semi-arid

(north) /arid (south) regions across Mongolia. The compositional changes based on Bray-Curtis

dissimilarity were investigated by generalized dissimilarity modeling, including geographic

distance, aridity, soil pH, and sand and clay contents as independent variables. The determinants

and changing patterns of community composition were compared among the four groups.

Aridity had significant impacts on community composition regardless of the regions and the

absence/presence of grazing. However, without the dependency on grazing, the difference in

response patterns was observed between the regions. The compositional change was steeper

especially at the upper edge of aridity rather than the lower edge in the arid region. This

indicates the vulnerability of plant communities to aridity shifts due to future climate change in

desert steppe of Mongolia. In addition, regardless of the regions, the effects of soil pH on

community composition were eliminated by grazing. Because soil pH indirectly affected by

aridity shifts can have impacts on community composition of potential vegetations without

grazing, the long-term monitoring of vegetation dynamics needs observations of both of

communities without and with grazing.

Keywords

dryland, Eurasia steppe, generalized dissimilarity modeling, grassland, soil pH.

Introduction

Plant community composition is a central factor determining the impacts of climate change and anthropogenic disturbances on ecosystem structures and functioning (Chapin et al., 2000). Across dryland ecosystems, aridity and livestock grazing are the most influential factors on these facets via the changes in plant community composition (Maestre et al., 2016).

Berdugo et al. (2020) reported abrupt shifts in the ecosystem structures (e.g. plant cover and species richness) along a broad gradient of aridity. Additionally, aridity affects not only water availability but also edaphic variables through long-term feedback from vegetations (Berdugo et al., 2020; Delgado-Baquerizo et al., 2013; Moreno-Jiménez et al., 2019) Because edaphic variables can have impacts on the patterns of community composition across dryland grasslands (Sasaki et al., 2008b; Zemmrich et al., 2010), the consideration of both of aridity and edaphic variables associated with aridity is required to clarify community assembly in drylands. However, little is known about the consistency in patterns of changes in community composition among different aridity gradients with this consideration. Understanding consistency or inconsistency in compositional changes across different gradients of aridity can provide implications for the impacts of aridity shifts under climate change on plant communities and their functioning in dryland ecosystems.

Grassland vegetations are widely used for livestock grazing to support secondary

production in drylands. Numerous studies have revealed the impact of grazing on community

composition mediated by selective herbivory (Díaz et al., 2007; Li et al., 2022; Sasaki et al.,

2008a). The grazing impacts on community composition can differ according to differences in

water availability. Previous studies have demonstrated that the relative magnitude of grazing

impacts on vegetation is relatively low in more arid regions (Herrero-Jáuregui and Oesterheld,

2018; Price et al., 2022). However, regardless of aridity levels, community composition can be

drastically altered and homogenized by high intensity of grazing, namely overgrazing (Ahlborn

et al., 2020; Sasaki et al., 2009b, 2008a; Wang et al., 2017). Thus, the patterns of changes in

community composition along aridity gradients in grazed grasslands can be confounded

depending on grazing. To infer how the changes of community composition along different

aridity gradients will be driven by climate change, the comparisons of compositional changes

are essential with the consideration of the interaction between aridity levels and grazing.

Notably, these comparisons have not been conducted.

Here, we investigated the patterns of changes in plant community composition along

aridity gradients with different levels of average aridity (semi-arid and arid regions) in the

Mongolian steppe depending on the absence/presence of grazing (without/with the long-term

exclosure of livestock). Mongolia covers a vast area of the eastern part of the Eurasian Steppe.

In Mongolia, the climate is spatially characterized by broad variations in aridity associated with

temperature and precipitation (Vandandorj et al., 2017). Suzuki et al. (2022) demonstrated that

aridity was the important driver of plant community composition across the whole range of

Mongolia. Thus, we tested the research questions below mentioned to provide implications into

the vulnerability of the Mongolian steppe communities and the long-term monitoring of

vegetation dynamics to climate change: 1) whether and how the effects of aridity and edaphic

variables on community composition differ between semi-arid and arid grasslands, and 2) does

grazing alter these effects.

Materials and methods

Across Central Mongolia, ranging from semi-arid to arid regions, over 600 km, 12 study sites

were selected at the meteorological sites of the Mongolian Information and Research Institute

of Meteorology, Hydrology, and Environment (IRIMHE) (Table 1, and Figure S1). We aimed

to ensure that the study area covered a range of vegetation types within the rangeland

ecosystems of Mongolia. Central Mongolia has the typical climate of Central Asia, which is

characterized by hot and short summers with approximately 70 % annual precipitation and cold

winters, including extreme climate events such as drought. Based on the climate data between

1978 and 2017 provided by IRIMHE, the mean annual precipitation ranged from 97 mm at

Tsogt-Ovoo to 333 mm at Darkhan; the mean annual temperature ranged from -0.8 °C at Ugtaal

to 7.6 °C at Khanbogd. Thus, there is a wide gradient of aridity from north to south regions

(Figure S1), which supports three main vegetation types: forest steppe, steppe, and desert steppe

(Table 1).

We set three 5 × 5 m blocks, which included three 1 × 1 m plots, at inside and outside

of meteorological sites (50 × 50 m), to observe the heterogeneity at both block- and site-level

scales. All the sites had ungrazed and grazed replicates. Ungrazed ones were exclosed for at

least 50 years to avoid livestock grazing. Because the distances between ungrazed and grazed

replicates in each site varies among sites, from a minimum of 25 m to a maximum of 3 km, the

comparison of community composition between ungrazed and grazed replicates in each site was

not performed in this study. In each plot, vegetation was sampled by identifying all species

rooting in each plot, and absolute cover (%) of each species was recorded from July to August

2022. Then, the vegetation data were pooled for each block. Simpson diversity index, which

reflects the diversity index mainly based on relative abundance of dominant species, was almost

saturated when the vegetation data of more than two plots was used for the calculation of

Simpson index in each site (Figure S2).

Aridity, which is calculated as [1–precipitation/potential evapotranspiration], based on

climatic data for the 1970–2000 period was obtained from the Global Aridity Index and

Potential Evapotranspiration Climate Database v3 (Trabucco and Zomer, 2022). The edaphic

variables were investigated based on the measurement of a soil sample in each block: total

nitrogen content in cg/kg; soil organic carbon content in dg/kg; soil pH in H_2O; clay, sand, and

silt content mass fraction in %. Soil cores with 5 cm in diameter and 10 cm in length in three

plots were pooled by each block, and used for the measurement of edaphic variables. Because

of the high level of correlations (absolute value of a correlation > 0.6) between aridity, and total

nitrogen and soil organic carbon, and between sand and silt content, we used the aridity, soil

pH, sand content, and clay content as the environmental variables in the subsequent analysis.

Nonmetric multi-dimensional scaling (NMDS) based on Bray-Curtis dissimilarity was

performed for the whole dataset of vegetation using the function 'metaMDS' of the R package

'vegan.' The dimensional reduction to two dimensions was performed because the stress value

(0.093) was lower than 0.2 (Kruskal, 1964). In addition, the clustering based on Ward's

algorithm was conducted using the Bray-Curtis dissimilarity matrix.

We performed the analysis independently for the following four groups: the groups

from the north (semi-arid) and south (arid) regions in Mongolia without/with grazing (north-

ungrazed, north-grazed, south-ungrazed, and south-grazed). To assess the difference in

community composition among the study sites, Bray-Curtis dissimilarity was calculated for

each group. The abundance-based dissimilarity was then partitioned into balanced variation in

abundance and abundance gradient using the function 'beta.multi.abund' of the R package

'betapart' (Baselga, 2017). This partitioning informs relative contributions of species turnover

(balanced variation) and nestedness (abundance gradient) to community composition difference.

Moveorver, to identify factors influencing community compositions, we applied generalized

dissimilarity modeling (GDM; Ferrier et al. (2007)). GDM is a matrix regression technique that

permits non-linear responses of compositional dissimilarity to spatial or environmental

gradients. This approach has an additional advantage to depict the rate of compositional shift at

any point along the gradient of a variable because GDM fits monotonically increasing flexible

I-spline functions for each independent variable. When a *I*-spline was plotted, the maximum

height depicts the magnitude in an effect of a given variable on compositional dissimilarity

(partial dissimilarity) with holding all the other independent variables constant. The first

derivatives (slopes) of a spline indicates the rate of compositional shift at any point along the

gradient of a variable. Using the function 'gdm' of the R package 'gdm', we fitted full GDMs

with pairwise geographical distance and the selected environmental variables as independent

variables for each group. These models were fitted with the three *I*-spline basis functions per

variable, and regression coefficient at each point (lower edge, mid-point, and upper edge) along

an independent variable was estimated. The uncertainty for each GDM plot (standard deviation)

was calculated using the function 'plotUncertainty' with 1,000 bootstrap iterations. Additionally,

we fitted GDMs with a single independent variable. From these models, the significance of each

variable was calculated using Monte Carlo permutation by the function 'gdm.varImp.' We

investigated 1) whether and how the effects of aridity and edaphic variables on community

composition differ between semi-arid and arid grasslands, based on the comparisons in the

results of GDMs between the north and south regions. In addition, the comparisons in the results

of GDMs between the ungrazed and grazed communities assessed that 2) does grazing alter

these effects. To check correlations between the matrices of distance (geographical distance and

differences in the selected environmental variables), the calculation of Mantel's correlation

coefficient based on Spearman correlation coefficient and Mantel test for each pair of the

variables were performed.

Results

The result of NMDS and clustering showed that there were mainly two groups of community

composition along the first axis (Figure S3–S5). These two groups corresponded to the

communities in the north (forest steppe) and south (steppe and desert steppe) regions in Central

Mongolia (Figure S5). The partitioning of Bray-Curtis dissimilarity exhibited the large

contribution with more than 90% of balanced variation in abundance (species turnover) rather

than abundance gradient (nestedness) (Table S1). GDMs revealed that the significant effects of

geographic distance and aridity on the variation of community composition across all the groups

(Table 2). On the other hand, only in the ungrazed communities of the both regions, the impacts

of soil pH on community composition were significant. We observed the similar response

patterns of community composition (partial Bray-Curtis dissimilarity) to geographic distance

and soil pH in the both regions (Table 2, and Figure 1). However, the different response patterns

to aridity were visually observed between the north and south regions: the steeper slope in the

lower edge of aridity and saturated response in the upper edge in the north region, and the

steeper response in the mid-point and upper edge in the south region. This pattern was not

depended on the absence/presence of grazing. Mantel correlation coefficients between

geographic distance and aridity difference in the north and south regions were similarly high,

0.748 and 0.790, respectively (Table S2).

Discussions

Although aridity had significant impacts on community compositions in both of the semi-arid

and arid regions of Mongolia (Table 2), we firstly demonstrated the difference in response

patterns along the aridity gradients between these regions regardless of grazing (Figure 1). Thus,

grazing did not confound the aridity impacts on community composition. The changes in

community composition in the south and arid region was abrupt especially at the upper edge of

aridity (Figure 1). This pattern might be mainly attributed to the relatively high level of

environmental filtering and species turnover because of abiotic and biotic factors by high level

of aridity. Because of the physiological limitations associated with aridity, the dominance of

stress-tolerant species and exclusion of non-tolerant species can often be observed in dryland

ecosystems (Berdugo et al., 2019; Grime, 1977; Le Bagousse-Pinguet et al., 2017). Previous

studies have also demonstrated that high aridity contributes to the severe limitations of not only

water availability but also soil nutrient availability, such as nitrogen, carbon, and micronutrients,

via long-term feedback from vegetations (Berdugo et al., 2020; Delgado-Baquerizo et al., 2013;

Moreno-Jiménez et al., 2019), resulting in severe environmental filtering by the abiotic factors

associated with aridity. As a biotic factor, species turnover via competitive exclusion by highly

adapted species in severely arid environments (Berdugo et al., 2019; Grime, 1977; Macarthur

and Levins, 1967; Michalet et al., 2014) might attribute to the observed response patterns of

community composition in this study. Increasing aridity can lead to the dominance of aridity

specialists and an increase in the relative importance of competitive exclusion by the superior

specialists, rather than facilitation (Berdugo et al., 2019). Some of previous studies have

indicated that negative effects on water availability by aridity specialists (e.g. shrub species)

can cause competitive exclusion in dryland ecosystems (Butterfield et al., 2016; O'Brien et al.,

2017). In this study, one or two shrub species were dominant or predominant in the sites with

the highest aridity (e.g., *Zygophyllum kaschgaricum*, *Potaninia mongolica*, *Anabasis brevifolia*,

and *Salsola passerina*; Table 1) In addition, the relative abundance of these shrub species was

positively correlated with aridity (Table S3), which might support the competitive exclusions

by shrub species.

223   Our study also emphasized that the elimination of the effects of soil pH on grassland

community composition across the semi-arid and arid regions by grazing (Table 2, and Figure

1). Although the relationships between species richness and soil pH, which is related to nutrient

availability, were negative or absent in semi-arid and arid regions (Palpurina et al., 2017), the

community composition including relative abundance of component species only without

grazing was significantly affected by soil pH in both of the semi-arid and arid regions. This

indicated that the grazed communities lost some variations of community composition

associated with spatial heterogeneity of soil pH. In a coastal dune community, grazing removed

the impact of soil pH on community composition (Tahmasebi Kohyani et al., 2008). In addition,

grazing drastically decreased the number of species of which abundance were strongly affected

by soil pH in a limestone plant community (Amezaga et al., 2004). Selective herbivory by

livestock can lead to communities with biased life history and functional traits (e.g. annual and

small-height plants) (Díaz et al., 2007), resulting in community composition less sensitive to

soil pH via the reduction in diversity of responses to nutrient heterogeneity among species.

237   Our study indicated the vulnerability of plant communities to aridity shifts due to future

climate change, especially in the upper edge of aridity range of Mongolia, namely desert steppes.

Because accurate threshold detections for community composition along the aridity gradients

(Berdugo et al., 2020) could not be performed in this study, future studies should use continuous

and intensive vegetation sampling across the whole range of the Eurasian steppe and other

regions. As a limitation in this study, there is no observation of temporal variability of vegetation

and aridity. For example, Sasaki et al. (2009a) observed the inter-annual variations of plant

community composition in the Mongolian steppe. Then, we propose the temporal observations

of the relationships between community composition, aridity, and edaphic variables along

multiple aridity gradients as a next step for understanding community assembly related to aridity,

and its spatial and temporal variability. As another important aspect of this study, we raised the

elimination of the impacts of soil pH, which is positively correlated with aridity globally

(Moreno-Jiménez et al., 2019), on community composition by grazing. Because soil pH can

have impacts on community composition of potential vegetations via indirect effects of aridity

shifts, our study empathized the importance of not only grazed communities but also ungrazed

ones for the long-term monitoring regarding vegetation dynamics exposed to climate change.

Ethics approval

Not applicable.

Consent to participate

All the authors approved participation in this study.

Consent for publication

All the authors approved the publication of this study.

Data availability

Data are available from the corresponding author upon reasonable request.

Code availability

Codes are available from the corresponding author upon reasonable request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have influenced the work reported in this study.

Author Contributions

274   N.I.I., G.B., G.T., M.K., and T.S. designed the study. All the authors contributed to the

field surveys. N.I.I. performed the analyses and wrote the first draft of the manuscript. All the

authors have contributed to the revisions.

Acknowledgments

279   We would like to thank our laboratory colleagues for their help with the analyses.

Funding

282   This work was financially supported by a fund for Fostering Joint International

Research A (no. 19KK0393) and a grant-in-aid for Scientific Research B (no. 22H03791)

presented to T.S. from the Ministry of Education, Culture, Sports, Science, and Technology of

Japan.

References

Ahlborn, J., von Wehrden, H., Lang, B., 2020. Climate–grazing interactions in Mongolian
rangelands: Effects of grazing change along a large-scale environmental gradient. J.
Arid Environ. 173, 104043.

Amezaga, I., Mendarte, S., Albizu, I., Besga, G., Garbisu, C., Onaindia, M., 2004. Grazing
intensity, aspect, and slope effects on limestone grassland structure. J. Range. Manag.
57, 606–612.

Baselga, A., 2017. Partitioning abundance‐based multiple‐site dissimilarity into
components: balanced variation in abundance and abundance gradients. Methods Ecol.
Evol. 8, 799–808.

Berdugo, M., Delgado-Baquerizo, M., Soliveres, S., Hernández-Clemente, R., Zhao, Y., Gaitán,
J.J., Gross, N., Saiz, H., Maire, V., Lehman, A., Rillig, M.C., Solé, R.V., Maestre, F.T.,
2020. Global ecosystem thresholds driven by aridity. Science 367, 787–790.

Berdugo, M., Maestre, F.T., Kéfi, S., Gross, N., Le Bagousse-Pinguet, Y., Soliveres, S., 2019.
Aridity preferences alter the relative importance of abiotic and biotic drivers on plant
species abundance in global drylands. J. Ecol. 107, 190–202.

Butterfield, B.J., Bradford, J.B., Armas, C., Prieto, I., Pugnaire, F.I., 2016. Does the stress-
gradient hypothesis hold water? Disentangling spatial and temporal variation in plant
effects on soil moisture in dryland systems. Funct. Ecol. 30, 10–19.

Chapin, F.S., 3rd, Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L.,
Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C., Díaz, S., 2000.
Consequences of changing biodiversity. Nature 405, 234–242.

Delgado-Baquerizo, M., Maestre, F.T., Gallardo, A., Bowker, M.A., Wallenstein, M.D., Quero,
J.L., Ochoa, V., Gozalo, B., García-Gómez, M., Soliveres, S., García-Palacios, P.,
Berdugo, M., Valencia, E., Escolar, C., Arredondo, T., Barraza-Zepeda, C., Bran, D.,
Carreira, J.A., Chaieb, M., Conceição, A.A., Derak, M., Eldridge, D.J., Escudero, A.,
Espinosa, C.I., Gaitán, J., Gatica, M.G., Gómez-González, S., Guzman, E., Gutiérrez,
J.R., Florentino, A., Hepper, E., Hernández, R.M., Huber-Sannwald, E., Jankju, M., Liu,
J., Mau, R.L., Miriti, M., Monerris, J., Naseri, K., Noumi, Z., Polo, V., Prina, A., Pucheta,
E., Ramírez, E., Ramírez-Collantes, D.A., Romão, R., Tighe, M., Torres, D., Torres-
Díaz, C., Ungar, E.D., Val, J., Wamiti, W., Wang, D., Zaady, E., 2013. Decoupling of
soil nutrient cycles as a function of aridity in global drylands. Nature 502, 672–676.

Díaz, S., Lavorel, S., McINTYRE, S.U.E., Falczuk, V., Casanoves, F., Milchunas, D.G., Skarpe,
C., Rusch, G., Sternberg, M., Noy-Meir, I., Landsberg, J., Zhang, W., Clark, H.,

Campbell, B.D., 2007. Plant trait responses to grazing? A global synthesis. Glob. Chang.
Biol. 13, 313–341.

Ferrier, S., Manion, G., Elith, J., Richardson, K., 2007. Using generalized dissimilarity
modelling to analyse and predict patterns of beta diversity in regional biodiversity
assessment. Divers. Distrib. 13, 252–264.

Grime, J.P., 1977. Evidence for the existence of three primary strategies in plants and its
relevance to ecological and evolutionary theory. Am. Nat. 111, 1169–1194.

Herrero-Jáuregui, C., Oesterheld, M., 2018. Effects of grazing intensity on plant richness and
diversity: a meta-analysis. Oikos 127, 757–766.

Kruskal, J.B., 1964. Multidimensional scaling by optimizing goodness of fit to a nonmetric
hypothesis. Psychometrika 29, 1–27.

Le Bagousse-Pinguet, Y., Gross, N., Maestre, F.T., Maire, V., de Bello, F., Fonseca, C.R., Kattge,
J., Valencia, E., Leps, J., Liancourt, P., 2017. Testing the environmental filtering concept
in global drylands. J. Ecol. 105, 1058–1069.

Li, Y., Shi, C., Zhang, T., Wang, H., Yang, Z., Tang, S., Li, F.Y., 2022. Summer grazing by three
livestock species at moderate intensity enhances primary productivity and drives

community divergence in a semi‐arid steppe. Appl. Veg. Sci. 25, e12683.

Macarthur, R., Levins, R., 1967. The limiting similarity, convergence, and divergence of
coexisting species. Am. Nat. 101, 377–385.

Maestre, F.T., Eldridge, D.J., Soliveres, S., Kéfi, S., Delgado-Baquerizo, M., Bowker, M.A.,
García-Palacios, P., Gaitán, J., Gallardo, A., Lázaro, R., Berdugo, M., 2016. Structure
and functioning of dryland ecosystems in a changing world. Annu. Rev. Ecol. Evol. Syst.
47, 215–237.

Michalet, R., Le Bagousse-Pinguet, Y., Maalouf, J.-P., Lortie, C.J., 2014. Two alternatives to
the stress-gradient hypothesis at the edge of life: the collapse of facilitation and the
switch from facilitation to competition. J. Veg. Sci. 25, 609–613.

Moreno-Jiménez, E., Plaza, C., Saiz, H., Manzano, R., Flagmeier, M., Maestre, F.T., 2019.
Aridity and reduced soil micronutrient availability in global drylands. Nat Sustain 2,
371–377.

O'Brien, M.J., Pugnaire, F.I., Armas, C., Rodríguez-Echeverría, S., Schöb, C., 2017. The shift
from plant-plant facilitation to competition under severe water deficit is spatially
explicit. Ecol. Evol. 7, 2441–2448.

Palpurina, S., Wagner, V., von Wehrden, H., Hájek, M., Horsák, M., Brinkert, A., Hölzel, N.,
Wesche, K., Kamp, J., Hájková, P., Danihelka, J., Lustyk, P., Merunková, K., Preislerová,
Z., Kočí, M., Kubešová, S., Cherosov, M., Ermakov, N., German, D., Gogoleva, P.,
Lashchinsky, N., Martynenko, V., Chytrý, M., 2017. The relationship between plant

species richness and soil pH vanishes with increasing aridity across Eurasian dry grasslands. Glob. Ecol. Biogeogr. 26, 425–434.

Price, J.N., Sitters, J., Ohlert, T., Tognetti, P.M., Brown, C.S., Seabloom, E.W., Borer, E.T., Prober, S.M., Bakker, E.S., MacDougall, A.S., Yahdjian, L., Gruner, D.S., Olde Venterink, H., Barrio, I.C., Graff, P., Bagchi, S., Arnillas, C.A., Bakker, J.D., Blumenthal, D.M., Boughton, E.H., Brudvig, L.A., Bugalho, M.N., Cadotte, M.W., Caldeira, M.C., Dickman, C.R., Donohue, I., Grégory, S., Hautier, Y., Jónsdóttir, I.S., Lannes, L.S., McCulley, R.L., Moore, J.L., Power, S.A., Risch, A.C., Schütz, M., Standish, R., Stevens, C.J., Veen, G.F., Virtanen, R., Wardle, G.M., 2022. Evolutionary history of grazing and resources determine herbivore exclusion effects on plant diversity. Nat Ecol Evol 6, 1290–1298.

Sasaki, T., Okayasu, T., Jamsran, U., Takeuchi, K., 2008a. Threshold changes in vegetation along a grazing gradient in Mongolian rangelands. J. Ecol. 96, 145–154.

Sasaki, T., Okayasu, T., Ohkuro, T., Shirato, Y., Jamsran, U., Takeuchi, K., 2009a. Rainfall variability may modify the effects of long-term exclosure on vegetation in Mandalgobi, Mongolia. J. Arid Environ. 73, 949–954.

Sasaki, T., Okayasu, T., Shirato, Y., Jamsran, U., Okubo, S., Takeuchi, K., 2008b. Can edaphic factors demonstrate landscape-scale differences in vegetation responses to grazing? Plant Ecol. 194, 51–66.

Sasaki, T., Okubo, S., Okayasu, T., Jamsran, U., Ohkuro, T., Takeuchi, K., 2009b. Two-phase functional redundancy in plant communities along a grazing gradient in Mongolian rangelands. Ecology 90, 2598–2608.

Suzuki, K., Tungalag, R., Narantsetseg, A., Tsendeekhuu, T., Masato, S., Yamanaka, N., Takashi, K., 2022. Composition, distribution, and environmental drivers of Mongolian rangeland plant communities. J. plant ecol. https://doi.org/10.1093/jpe/rtac100

Tahmasebi Kohyani, P., Bossuyt, B., Bonte, D., Hoffmann, M., 2008. Importance of grazing and soil acidity for plant community composition and trait characterisation in coastal dune grasslands. Appl. Veg. Sci. 11, 179–186.

Trabucco, A., Zomer, R.J., 2022. Version 3 of the "Global aridity index and potential evapotranspiration (ET0) database": Estimation of Penman-Monteith reference evapotranspiration. figshare. https://doi.org/10.6084/m9.figshare.7504448.v3

Vandandorj, S., Munkhjargal, E., Boldgiv, B., Gantsetseg, B., 2017. Changes in event number and duration of rain types over Mongolia from 1981 to 2014. Environ. Earth Sci. 76, 1–12.

Wang, Y., Heberling, G., Görzen, E., Miehe, G., Seeber, E., Wesche, K., 2017. Combined effects of livestock grazing and abiotic environment on vegetation and soils of grasslands across Tibet. Appl. Veg. Sci. 20, 327–339.

394 Zemmrich, A., Manthey, M., Zerbe, S., Oyunchimeg, D., 2010. Driving environmental factors
395   and the role of grazing in grassland communities: A comparative study along an
396   altitudinal gradient in Western Mongolia. J. Arid Environ. 74, 1271–1280.

Tables and figures

Table 1. Locations, aridity, vegetation type, and dominant species of the 12 study sites. Aridity is calculated as 1 − (precipitation/potential evapotranspiration).

Region	Site	ID	Latitude	Longitude	Aridity	Vegetation	Dominant species
North (semi-arid) region	Erdenet	ERD	49°02'07.08"N	106°56'20.74"E	0.62	Forest steppe	*Elymus chinensis, Koeleria macrantha, Medicago ruthenica*
	Hutag	HUT	49°23'37.32"N	106°50'50.38"E	0.64	Forest steppe	*Artemisia frigida, Stipa krylovii, Medicago ruthenica*
	Bulgan	BUL	48°49'05.52"N	106°54'04.54"E	0.64	Forest steppe	*Agropyron cristatum, Stipa krylovii, Cleistogenes squarrosa*
	Tarialan	TAR	49°36'33.12"N	106°47'56.30"E	0.65	Forest steppe	*Artemisia frigida, Cleistogenes squarrosa, Heteropapus hispidus*
	Darkhan	DAR	49°29'25.44"N	107°03'45.58"E	0.69	Steppe	*Carex pediformis, Artemisia scoparia, Heteropapus hispidus*
	Ugtaal	UGT	48°15'29.16"N	107°01'37.27"E	0.72	Steppe	*Potentilla acaulis, Elymus chinensis, Stipa krylovii*
	Zuunmod	ZUU	47°42'47.52"N	107°07'48.46"E	0.72	Steppe	*Artemisia frigida, Stipa krylovii, Arenaria capillaris*

	Mandalgobi	MAN	45°44'35.52"N	107°05'03.43"E	0.87	Desert steppe	*Carex stenophylloides, Stipa glareosa, Allium polyrrhizm*
South (arid) region	Dalanzadgad	DAL	43°34'37.92"N	106°57'40.39"E	0.92	Desert steppe	*Stipa gobica, Artemisia pectinata, Convolvulus amanii*
	Saikhan	SAI	44°05'42.00"N	106°54'09.12"E	0.92	Desert steppe	*Convolvulus amanii, Stipa gobica, Cleistogenes songorica*
	Tsogt-Ovoo	TSO	44°25'23.52"N	107°01'16.39"E	0.93	Desert steppe	*Allium polyrrhizm, Anabasis brevifolia, Salsola passerina*
	Khanbogd	KHA	43°11'44.52"N	107°08'46.46"E	0.94	Desert steppe	*Zygophyllum kaschgaricum, Cleistogenes songorica, Potaninia mongolica*

403   **Table 2.** Regression coefficients and their significance of geographic distance, aridity, and soil pH based on generalized dissimilarity

404   modeling (GDM) in ungrazed and grazed communities of the north and south regions. Regression coefficients on a spline (in Figure 1) at the

405   lower edge, mid-point, and upper edge of each independent variable in each group (north-ungrazed, north-grazed, south-ungrazed, and south-

406   grazed) are shown. Bold characters denote statistically significant variables.

Region	Ungrazed/grazed community	Proportion of variation explained (%)	Independent variable	Regression coefficient		*P* value
North (semi-arid) region	Ungrazed community	66.3	**Geographic distance**	lower edge	0.705	
				mid-point	0.000	<0.001
				upper edge	0.000	
			Aridity	lower edge	0.626	
				mid-point	0.000	0.003
				upper edge	0.000	
			pH	lower edge	0.141	
				mid-point	0.000	0.010
				upper edge	0.909	
			Sand	lower edge	0.000	
				mid-point	0.000	0.231
				upper edge	0.012	

			Clay	lower edge	0.000	
				mid-point	0.000	0.633
				upper edge	0.000	
			Geographic distance	lower edge	0.651	
				mid-point	0.000	<0.001
				upper edge	0.000	
			Aridity	lower edge	0.618	
				mid-point	0.000	0.004
				upper edge	0.000	
	Grazed community	55.9	pH	lower edge	0.000	
				mid-point	0.000	0.337
				upper edge	0.053	
			Sand	lower edge	0.000	
				mid-point	0.000	1.000
				upper edge	0.000	
			Clay	lower edge	0.137	
				mid-point	0.000	0.328
				upper edge	0.000	
South (arid) region	Ungrazed community	77.6	**Geographic distance**	lower edge	0.588	
				mid-point	0.765	0.008
				upper edge	0.000	
			Aridity	lower edge	0.000	
				mid-point	0.828	0.001
				upper edge	0.841	

		pH	lower edge	0.000	0.007
			mid-point	1.066	
			upper edge	0.000	
		Sand	lower edge	0.000	0.423
			mid-point	0.000	
			upper edge	0.000	
		Clay	lower edge	0.000	1.000
			mid-point	0.000	
			upper edge	0.032	
Grazed community	78.0	**Geographic distance**	lower edge	1.572	<0.001
			mid-point	0.019	
			upper edge	0.000	
		Aridity	lower edge	0.000	<0.001
			mid-point	0.753	
			upper edge	0.599	
		pH	lower edge	0.000	1.000
			mid-point	0.000	
			upper edge	0.000	
		Sand	lower edge	0.000	0.317
			mid-point	0.000	
			upper edge	0.447	
		Clay	lower edge	0.000	0.741
			mid-point	0.000	

upper edge	0.051	

409 **Figure 1.** Relationships between partial Bray-Curtis dissimilarity and geographic distance,

410 aridity, and soil pH based on generalized dissimilarity modeling (GDM). The estimated splines

411 and their confidence intervals are denoted.

Supplementary information

Table S1. Partitioning of Bray-Curtis dissimilarity to balanced variation in abundance (species turnover) and abundance gradient (nestedness).

Region	Ungrazed/grazed community	Balanced variation in abundance (%)	Abundance gradient (%)
North (semi-arid) region	Ungrazed community	97.6	2.4
	Grazed community	97.2	2.8
South (arid) region	Ungrazed community	95.4	4.6
	Grazed community	91.4	8.6

7   **Table S2.** Mantel correlation coefficients based on Spearman's ρ between variables in ungrazed and grazed communities of the north and

8   south regions. Bold characters denote statistically significant correlations.

Variable	Variable	North_Ungrazed	North_Grazed	South_Ungrazed	South_Grazed
Geographic distance	Aridity	**0.748**	**0.748**	**0.790**	**0.790**
Geographic distance	pH	0.128	**0.236**	0.117	**0.327**
Geographic distance	Sand	**0.188**	**0.275**	0.137	**0.460**
Geographic distance	Clay	0.072	-0.036	-0.010	-0.074
Aridity	pH	0.079	**0.174**	0.008	**0.472**
Aridity	Sand	**0.336**	**0.237**	0.171	**0.255**
Aridity	Clay	-0.050	0.050	0.089	-0.088
pH	Sand	0.037	**0.176**	-0.009	-0.030
pH	Clay	**0.289**	-0.006	-0.019	0.024
Sand	Clay	**0.261**	0.069	**0.423**	0.132

11 **Table S3.** Spearman's ρ between aridity and relative abundance of shrub species in ungrazed
12 and grazed communities of the north and south regions. Bold characters denote statistically
13 significant correlations.

Region	Fence	ρ
North (semi-arid) region	Ungrazed community	0.295
	Grazed community	0.106
South (arid) region	Ungrazed community	**0.622**
	Grazed community	**0.644**

16 **Figure S1.** Spatial distribution of the 12 study sites and aridity in Mongolia. Aridity is
17 calculated as 1 – (precipitation/potential evapotranspiration).
18

Figure S2. Relationships between number of plots and Simpson diversity index of ungrazed

 and grazed communities in each site.

24   **Figure S3.** Result of nonmetric multi-dimensional scaling (NMDS) based on Bray-Curtis
25   dissimilarity of ungrazed/grazed communities in the north and south regions.
26

Figure S4. Result of loading values of species in nonmetric multi-dimensional scaling (NMDS)

based on Bray-Curtis dissimilarity of ungrazed/grazed communities in the north and south

regions.

31   **Figure S5.** Result of clustering based on Ward's algorithm of ungrazed/grazed communities in
32   the north and south regions. Bray-Curtis dissimilarity matrix was used as a distance matrix.
33