The SLAP$_{TAG}$: A new molecular tag adapted for the development of a high-performance, low-cost, affinity chromatography system

Emanuel J. Muruaga[1,2], Paula J. Uriza[1,2], Gonzalo A. K. Eckert[1,2], María V. Pepe[1,2], Cecilia M. Duarte[1,2], Mara S. Roset[1,2] * and Gabriel Briones[1,2] *.

1. Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín (UNSAM)- Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). 2. Escuela de Bio y Nanotecnologías (EByN), Universidad Nacional de San Martín.

Running title: The SLAP$_{TAG}$ affinity chromatography (SAC)

Keywords: Affinity chromatography-SLAP$_{TAG}$.

*Corresponding author

Mailing address: Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín (IIB-UNSAM), Av. 25 de Mayo y Francia, Campus Migueletes, CP1650, San Martín, Buenos Aires, Argentina.

Tel.: +54 11 4006-1500 (Int 2139); Fax: + 54 11 (5411) 4006-1559.

E-mail: gbriones@iib.unsam.edu.ar, mroset@iib.unsam.edu.ar

Abstract

The $SLAP_{TAG}$ is a novel molecular TAG derived from a protein domain present in the sequence of

Lactobacillus acidophilus SlpA ($SlpA^{284-444}$). Proteins from different biological sources, with different

molecular weights or biochemical functions, can be fused in frame to the $SLAP_{TAG}$ and efficiently purified

by the specific binding to a bacterial-derived chromatographic matrix named here Bio-Matrix (BM). To set

an optimized protocol for the $SLAP_{TAG}$-based affinity chromatography (SAC) different binding and elution

conditions were evaluated. The binding equilibrium between $SLAP_{TAG}$ and BM was reached after a few

minutes at $4^{\circ}C$, being the apparent dissociation constant (K_D) of 4.3 uM, a value similar to the one

determined for other S-layer proteins and their respective bacterial cell-walls. A reporter protein was

generated (H_6-GFP-$SLAP_{TAG}$) to compare the efficiency of SAC against a commercial system based on a

Ni^{2+}-charged agarose matrix, observing no differences in the H_6-GFP-$SLAP_{TAG}$ purification performance.

The stability and reusability of the BM were evaluated, and it was determined that the matrix was stable

for more than a year, being possible to reuse it five times without a significant loss in the efficiency for

protein purification. Alternatively, we explored the recovery of bound SLAP-tagged proteins by proteolysis

using the $SLAP_{ASE}$ (a SLAP-tagged version of the HRV-3c protease) that released a tag-less GFP ($SLAP_{TAG}$-

less). Additionally, iron nanoparticles were linked to the BM and the resulting BM_{mag} was successfully

adapted for a magnetic SAC, a technique that can be potentially applied for high-throughput-out protein

production and purification.

Main

The amalgamation of genomics, transcriptomics, proteomics, metabolomics, synthetic biology, and artificial

intelligence has become a powerful approach to understand the complexity of any biological process,

uncovering uncharacterized molecular mechanisms [1] [2] [3].

In addition to being central for scientific research, OMICs also have become the core of the

biomanufacturing platform oriented to the generation of bioproducts of commercial value (either new

products or biosimilars) [4]. Amidst the universe of bioproducts, therapeutic proteins are a growing class of

biopharmaceutics that are being used for cancer, autoimmunity, and inflammatory diseases [5].

Because of their clinical efficiency and low toxicity, therapeutic proteins (monoclonal antibodies, cytokines,

antigens, hormones, vaccines, etc.) have become an excellent option for different ailment treatments [6] [7].

However, one of the limitations these treatment face is the high cost of manufacturing, which has

bottlenecked therapeutic protein production and promoted the development of several strategies to

improve the upstream and downstream processes [7].

The downstream process includes all the purification steps oriented to capture and eliminate protein

aggregates and any remanent impurities derived from the host cell (e.g., DNA), preserving the biological

activity of the purified bioproduct. At the core of the downstream process, chromatography is the critical

step but also the most expensive one [8] [9]. To reduce critical costs associated with this step different

approaches have been evaluated including the adoption of single-use technologies [10].

Affinity chromatography is a special type of liquid chromatography that exploits the existence of natural [11]

or artificial [12] affinities between two moieties: the molecular target (or tag) and its ligand (the affinity ligand)

which is usually immobilized onto a chromatographic stationary phase to generate a chromatography

matrix. Thus, proteins of interest can be fused in frame to different molecular tags (such as His-6X, GST,

MBP, etc.) expressed and purified ideally in a single step from a complex mixture of proteins.

Since it was developed by Starkestein in 1910 [13], affinity chromatography was gaining popularity and

centrality for many industrial processes such as the purification of proteins for diagnosis, research, and

therapeutic purposes [14]. Regulatory requirements for the purity and quality of the proteins varies greatly

depending on the area of application. For instance, for industrial use, bio-products can be used with little

purification. Also, recombinant protein produced for research, diagnosis, or non-clinical purposes have a

less stringent regulatory approval process than proteins designed as biopharmaceuticals.

Here, we explored the adaptation of a protein domain present in the carboxy terminus of the *Lactobacillus*

acidophilus protein SlpA as a molecular tag. *In silico* analysis of this region (SlpA$^{284-444}$) allowed us to identify

a tandem of two copies of a 60-aminoacid protein domain named SLP-A (pfam03217) which is necessary

for the association. Here, this dual SLP-A domain was named SLAP$_{TAG,}$ and the adaptation as a molecular tag

was evaluated. Recently, we have characterized the binding properties of the SLAP$_{TAG}$ and characterized

its association to live *Lactobacillus* wall for vaccine purposes [15]. Thus, the SLAP$_{TAG}$ was fused in frame to a

chimeric antigen derived from the Shiga toxin-producing *Escherichia coli* (STEC) formed by the peptides

EspA$^{36-192}$, Intimin$^{653-953}$ and Tir$^{240-378}$ (or EIT). The resulting chimeric antigen (EIT-SLAP$_{TAG}$) recombinantly

expressed and purified was able to associate to the bacterial cell-wall of *L. acidophilus*, a process that we

named decoration. Thus, EIT-decorated *L. acidophilus* after oral administration was able to deliver the EIT

antigen to the intestinal mucosa eliciting a protective immune response that controls an experimental STEC

infection in mice [15]. Remarkably, the decoration process does not modify genetically the *Lactobacillus*

genome preserving its GRAS (Generally Recognized as Safe) status, a trait that is important for vaccine

purposes.

Here, we explore the adaptation of the SLAP$_{TAG}$ for the development of a novel affinity chromatography,

the SLAP affinity chromatography (SAC), and a comprehensive protocol is presented.

Results

SLAP$_{TAG}$ binds rapidly and efficiently to the Bio-matrix (BM)

As described in Materials and Methods the SLAP$_{TAG}$ was fused in-frame to the carboxy-terminus of the green

fluorescent protein (GFP) to generate the chimeric protein (H$_6$)-GFP-SLAP$_{TAG}$ adapted here as a SLAP$_{TAG}$

reporter protein. As shown in Fig. 1A the structure of the chimeric protein (H$_6$)-GFP-SLAP$_{TAG}$ was predicted

based on alphaFold-2 method [16]. The beta-barrel corresponding to GFP and the two globular domains

corresponding to SLAP$_{TAG}$ were predicted with high performance and as expected, linkers and 6xHis-tag,

which are flexible regions, showed low prediction values.

As a working chromatographic matrix, a culture of *Bacillus subtilis* natto was processed as described in

Materials and Methods to generate a bacterial-derived affinity chromatography matrix, named here Bio-

Matrix (BM). Interestingly, although *B. subtilis* has no S-layer, we have demonstrated previously that this

bacterium can be externally covered with SLAP-tagged proteins to generate a recombinant S-layer on its

bacterial cell wall, a process that we called decoration [15]. Decoration of *B. subtilis* with SLAP-tagged proteins

is possible because teichoic acid and lipoteichoic acid (which are the molecules responsible for SLAP$_{TAG}$

association) have the same chemical composition than *Lactobacillus acidophilus*. Interestingly, in *B. subtilis*

and *L. acidophilus* teichoic acid and lipoteichoic acid are distributed homogeneously on their cell wall.

To characterize the binding properties of the SLAP$_{TAG}$, the reporter protein (H$_6$)-GFP-SLAP$_{TAG}$, was

recombinantly expressed in *Escherichia coli*, purified by affinity chromatography mediated by its Hisx6 tag

(H$_6$), and further incubated with BM under a variety of conditions. Initially, as shown in Fig. 1B, the optimal

binding temperature of (H$_6$)-GFP-SLAP$_{TAG}$ to the BM was evaluated, finding that the best binding efficiency

was reached when the incubation was performed at 4^{o}C. To get insights into the SLAP$_{TAG}$ association

dynamics, (H$_6$)-GFP-SLAP$_{TAG}$ was incubated for 5, 30 and 60 minutes with BM and as shown in the Fig. 1C,

the maximal binding of (H$_6$)-GFP-SLAP$_{TAG}$ to BM was reached very rapidly (already at 5 minutes), showing

no significant increments in its association at longer time points. These results indicated that the SLAP$_{TAG}$

binds very rapidly and with an apparent high affinity to the BM. To quantify the affinity of the interaction

between the SLAP$_{TAG}$ with the BM, an adsorption isotherm was performed to determine the equilibrium

apparent dissociation constant as described in Materials and Methods. As shown in Fig. 1D the apparent K$_D$

was estimated as 4.7 uM. Interestingly, this dissociation constant value was in the same order than other

K$_D$ described for different microbial S-layer proteins with their respective bacterial walls [17–19]. In addition,

the maximum adsorption capacity (Bmax) of BM was estimated as 1.152 mmols of (H$_6$)-GFP-SLAP$_{TAG}$ or 53.9

mg of protein per milliliter of BM or 1.03 g of SLAP$_{TAG}$ protein/g of dry BM.

Elution of (H$_6$)-GFP-SLAP$_{TAG}$ protein can be performed with different buffers

As it was mentioned, the SLAP$_{TAG}$ contains the protein region responsible for the association of the protein

SlpA to the *L. acidophilus* cell wall. As reported, *L. acidophilus* SlpA has the natural ability to self-assemble

on the bacterial surface to generate a proteinaceous layer named S-layer [20], a highly ordered wall structure

that functions as a protective barrier against bacteriophages, resistance to low pH and proteases, and

bacterial adhesion. It is well characterized that remotion of the S-layer can be performed efficiently by the

addition of chaotropic agents like LiCl, a compound that disrupts hydrogen bonds leading to a partial

denaturation of proteins and the consequent detachment of the S-layer [20]. Thus, initially, to study elution

conditions for (H$_6$)-GFP-SLAP$_{TAG}$, LiCl solution was selected as the positive control for elution and as shown

in Fig. 2A detachment of BM-associated (H$_6$)-GFP-SLAP$_{TAG}$ was achieved with high efficiency with a 2M LiCl

solution at room temperature, in a short lapse of five minutes (Fig. 2B). As it was mentioned above, lithium

solutions have deleterious effects on proteins, which is an undesired effect for protein purification,

especially for enzyme purification. Therefore, different milder alternatives of buffers were explored to elute

SLAP$_{TAG}$-tagged proteins, conditions that can preserve the protein structure and therefore their function.

As shown in Fig. 2C, a series of sugars (monosaccharide and disaccharides) described previously [21] were

tested for the elution of the reporter SLAP observing a partial elution efficiency when compared with the

LiCl elution control (Fig. 2C). Next, the cationic nature of the SLAP$_{TAG}$ at neutral pH was considered. Taking

in consideration the importance of this positive charge in the interaction of SLAP$_{TAG}$ with the membrane-

bound compound teichoic acid, a set of cationic compounds was evaluated for the elution. As shown in Fig.

2C, 0.3M of CTAB was able to elute (H_6)-GFP-SLAP$_{TAG}$ with a similar efficiency as the LiCl solutions. Although

CTAB was very efficient in this step, we found that this compound was difficult to remove downstream from

the elution fraction. Then, considering that SLAP$_{TAG}$ has a theoretical isoelectric point (pI) value of 9.92, we

explored if the modification of pH can be adapted as an elution method. As shown in Fig. 2C, when the pH

of the bicarbonate buffer was close to the theoretical SLAP$_{TAG}$ isoelectric point (pI), (H_6)-GFP-SLAP$_{TAG}$ was

efficiently eluted from BM. In addition, after establishing the bicarbonate buffer at pH 10 as an elution

buffer, we explore if the addition of NaCl can improve the elution process. As shown in Fig. 2C, the addition

of 200 mM of NaCl was able to maximize the yield of recovery of (H_6)-GFP-SLAP$_{TAG}$ in the eluate.

Optimized SAC protocol allowed the efficient purification of proteins with similar efficiency to the high-

performance Ni^{2+}-charged agarose matrix (IMAC).

With all the experimental information obtained, an optimized SAC protocol for the purification of SLAP-

tagged proteins was established as shown in Fig. 3A. Remarkably, the entire process of protein purification

can be performed in 15 minutes. To have a direct observation of the purification process of (H_6)-GFP-

SLAP$_{TAG}$, the BM was fixed and immobilized on coverslips at different steps of the process to be observed

by confocal microscopy. As shown in Fig. 3B, before the incubation of BM with (H_6)-GFP-SLAP$_{TAG}$, only a red

native autofluorescence can be observed from fixed *Bacillus subtilis* natto cells present in BM. Remarkably,

after the incubation of BM with (H_6)-GFP-SLAP$_{TAG,}$ it was possible to detect the adhesion of the reporter

protein to the BM forming a recombinant fluorescent S-layer that covers completely the bacterial surface

(Fig. 3C). As expected, after the addition of the elution buffer the (H_6)-GFP-SLAP$_{TAG}$ was completely removed

from the bacterial surface (Fig. 3D). In addition, the whole process of purification can also be monitored by

direct observation under UV light (Fig. 3E)

Since SAC was efficient in protein purification, we compared this new technique against an established

affinity purification protocol. For this, we took advantage that (H_6)-GFP-SLAP$_{TAG}$ also has an His-tag that can

be purified by a Metal-Affinity-Chromatography or IMAC. As described in Materials and Methods, a

bacterial lysate (H_6)-GFP-SLAP$_{TAG}$ was divided into two fractions and both protocols were performed

accordingly, confirming that SAC optimized protocol was able to purify (H$_6$)-GFP-SLAP$_{TAG}$ with high efficiency

(Table Suppl. 1 and Fig. 3F) and with a similar yield to the one obtained with the commercial IMAC (Fig. 3G).

Proteins of different biological sources, molecular weights, biochemical functions, or expressed by

procaryotic or eukaryotic expression systems can be efficiently purified by the SAC.

To evaluate if SLAP$_{TAG}$ and SAC can be adapted as a universal protein purification system, a set of the

selected proteins were fused to the SLAP$_{TAG}$ (Fig. 4). Thus, bacterial proteins like *Salmonella* flagellin

(Fig.5A), the STEC proteins, EspA (Fig. 5B), the chimeric EIT (Fig. 5C) , the *B. abortus* Omp19 (Fig. 5D), the

bacteriophage protein T7-lysozyme (Fig. 5E), the human viral proteins Rhinovirus 3C Protease (Fig. 5E), the

mouse Galectin-8 (Fig. 5G) and the commercial molecular tag GST (Fig. 5H), were fused in frame with the

SLAP$_{TAG}$. As shown in Fig. 5, all the selected proteins were purified efficiently with SAC. In addition to the

bacterial expression system, different protein expressions systems (yeast and mammalian cells) were also

evaluated (Fig. 6). As shown in Fig. 6A the viral HRV 3C protease fused to the SLAP$_{TAG}$ was expressed and

purified from *Pichia pastoris* supernatant. Also, the SARS-CoV2 SPIKE fused to the SLAP$_{TAG}$ was purified

from the supernatant of transfected HEK293 (Fig. 6B). These results confirmed that the SLAP$_{TAG}$ can be

widely adopted for affinity chromatography purification.

BM is a reusable chromatography matrix with long-term stability.

As described in Materials and Methods a batch of BM was produced, and several aliquots were frozen at -

20^{o}C to study time stability. As shown in Fig. 7A, at different times, a few aliquots were unfrozen and tested

for protein purification of the reporter protein (H$_6$)-GFP-SLAP$_{TAG}$ being the BM stable for more than a year

(14 months). In addition, BM reusability capacity was evaluated determining that the matrix can be reused

five times with no modification of the protein purification yield (Fig. 7B).

Use of the Human Rhinovirus 3C protease fused to the SLAP$_{TAG}$ (SLAP$_{ASE}$) to release the SLAP-tagged proteins from BM.

As shown in Fig. 4 and 5 sequence of the viral protease HRV-3C was fused to the SLAP$_{TAG}$ (named SLAP$_{ASE}$), recombinantly expressed, and purified (Fig. 8B and 8D, lane SLAPase) to evaluate its activity. A reporter protein for SLAP$_{ASE}$ activity was generated (GFP-LEVLFQGP-SLAP$_{TAG}$) (Table 1) and a protocol for tag-remotion by SLAPase was set as described in Materials and methods. As shown in Fig. 8A the GFP-LEVLFQGP-SLAP$_{TAG}$ was expressed (Fig 8B and Fig. 8C , lane Input or IN) and mixed with the BM for 5 minutes allowing the binding process. After binding, a purified SLAP$_{ASE}$ was added to the mix and incubated for 60 minutes releasing a tag-less GFP (Fig. 8B and Fig 8C, Fig 8E, lanes flow-through or FT) by proteolysis. Cut SLAP$_{TAG}$ is not observed in the flow-through indicating that was retained by the BM along with the SLAPase. To confirm these steps, post-proteolysis BM was incubated with LiCl solution to remove any residual bound SLAP-tagged protein. As shown in Fig. 8B and 8D (lane Elution or E), the LiCl solution released the SLAPase and the cut SLAP$_{TAG}$.

Adaptation of SAC to a magnetic affinity chromatography.

Since we confirmed the efficiency and universality of SAC for the purification of recombinant proteins, we explored a magnetic affinity chromatography SAC alternative (Fig. 9A). As shown in Fig. 9B, 9C and Supplementary Video-1 the (H$_6$)-GFP-SLAP$_{TAG}$ was purified in a few and easy steps. As shown in the SDS-PAGE (Fig. 9B) the adapted SAC protocol for magnetic chromatography (BM$_{mag}$) was able to purify the reporter protein (H$_6$)-GFP-SLAP$_{TAG}$ very efficiently (Fig. 9B , lane Elution or E). Interestingly, this protocol can be adapted to commercial devices that use magnetic force for protein or DNA purification (like King Fisher Flex from ThermoFisher). In Fig. 9C the entire process of purification was monitored by direct observation under UV light (Supp. video 1). Noteworthy, the binding of the BM$_{mag}$ to GFP quenched the fluorescence of this protein, an effect that was described for transition metal binding to GFP or binding of iron cations to fluorescent proteins (Fig. 9C, tube IN+BM$_{mag}$) [22,23].

DISCUSSION

Proteins are macromolecules formed by a particular amino acid sequence that determines a given atomic spatial distribution, also known as protein conformation. Far from being a rigid structure, proteins behave as nano-machines performing precise molecular activities because of the internal movements of protein parts or protein domains. Proteins can interact intramolecularly or intermolecularly with different macromolecules such as proteins, DNA, RNA, polysaccharides, or with small compounds [24]. Some proteins degrade aliments to produce essential nutrients (like the digestive enzymes), proteins that transport critical compounds for supporting life (like hemoglobin that transports oxygen to the cells), proteins that shape the cellular structure (like actin, tubulin or keratin), proteins that function as hormones (like insulin), proteins that defense the organisms against pathogens (like the antibodies) [24]. In addition to the multiplicity of existent polypeptides in nature, molecular biology and biotechnology have generated a myriad of chimeric proteins by combining different protein domains [25]. These artificial proteins can be used as therapeutic tools to treat cancer, autoimmunity, or different medical conditions [26].

The SARS-CoV-2 pandemic has generated a multidimensional global crisis with devastating effects on human health, also having a massive impact on the economics everywhere. As a direct consequence of this global disaster, national governments were thrown into a global raging competition to acquire all the required medical supplies. In addition to the shortage of masks or ventilators, the huge demand for diagnostic reagents (such as recombinant proteins) has bottlenecked the development of new viral diagnosis kits based on ELISA or Real-time PCR tests.

Among all other things, the pandemic crisis had taught us that the availability of low-cost affinity chromatography might be a helpful tool for producing pure recombinant proteins to develop low-cost diagnostics tests.

Here we propose the adaptation of the binding domain of the *Lactobacillus acidophilus* SlpA as a molecular tag. As it was shown here, SAC is a useful tool for protein purification because i) SAC protocol uses only simple and low-cost reagents, ii) the chromatography matrix is ¨grown¨ instead of being chemically

synthetized, iii) ligands (LTA and teichoic acid) are naturally integrated to surface of the BM with no need

of chemical reactions to crosslink them to the matrix.

In summary, we propose that SLAP affinity chromatography can be adapted as an in-house protein

purification system, available for any Laboratory around the globe, to produce pure recombinant proteins

for research, diagnosis, and for the food industry. Although so far regulatory issues might preclude the use

of the SAC for proteins used as biotherapeutics, new efforts have been performed to develop a new version

of the matrix chromatography suitable for more stringent industrial or clinical purposes.

METHODS

Strains, plasmids, and transformation

Bacterial strains and plasmids used here are summarized in Table 1. *Escherichia coli* and *Bacillus subtilis*

strains were grown in Luria Bertani (LB) medium (Sigma, St. Louis, MO, United States) at 37ºC 180 rpm.

Bacterial plasmid vectors were transformed in *E. coli* DH5α or in *E. coli* BL21 (DE3) for protein expression.

Pichia pastoris was grown in Yeast Extract–Peptone–Dextrose (YPD) medium at 28ºC 180 rpm. HEK293F cells

were grown and maintained at 0.5–3.0 × 10^6 cells/mL in a humidified CO_2 platform shaker incubator at 37 ºC

in Dulbecco's Modified Eagle's Medium (DMEM - ThermoFisher). SARS-CoV-2 was a gift from Jason McLellan

(Addgene # 154754; http://n2t.net/addgene:154754;RRID: addgene_154754) [27].

Cloning

Lysozyme gene was amplified from pLysS plasmid (Millipore Sigma-Novagen) with oligonucleotide primers

Fw-lys-*Nde*I   (CCCATATGGCTCGTGTACAGTTTAAACAACGTG)   and   Rv-lys-SalI

(CGGTCGACTCCACGGTCAGAAGTGACCAGTTCG). The PCR product was digested with *Nde*I and *Sal*I and then

cloned into the pLC3-EITH7-SlpA vector at the *Nde*I and *Sal*I restriction sites. The recombinant plasmid was

amplified through transformation into *E. coli* DH5α. Finally, the lysozyme gene was subcloned into pET28-e

GFP-SlpA at the *Nde*I and *Not*I restriction sites and transformed into *E. coli* BL21 by electroporation.

EspA gene was amplified with the following primers: *pRvEspAXbaI* (GCTCTAGATTTACCAAGGGATATTGCTG)

and pFwEITSalI (ACGCGTCGACGATATGAATGAGGCATCTAAA). After restriction with *Xba*I and *Sal*I, the gene

was cloned into *pGEX-SlpA,* which was previously digested with the same enzymes. The plasmid was

introduced by electroporation into *E. coli* BL21.

The *hrv-3c-slap$_{tag}$* gene was synthesized and cloned in pMAL-c5X and pPICZalfa B for protein expression in *E.*

coli and *P. pastoris* respectively (Gene Universal, Delaware-USA).

For *P. pastoris* GS115 transformation, 5 ug of pPICZalfa B – *hrv-3c* plasmid was linearized with *Sac*I restriction

enzyme and transformed into cells through electroporation using 2 mm gap cuvettes (1500 V, 125 X, 50 lF).

The transformed cells were selected by seeded on a YPD medium containing Zeocin 100ug/ml for resistance

selection. Colonies were further grown in test tubes containing YPD broth and after 24 h, cultures were

induced by adding 1% (v/v) pure methanol every 24 h (1). Supernatants were sampled after 72 h of induction

and the best producers were chosen for further experiments.

pET28-eGFP-SLAP was amplified and the SLAP$_{TAG}$ sequence subcloned into SARS-CoV-2 S HexaPro plasmid

(Addgene#154754) between *BamH*I and *Xho*I restriction sites.

Protein expression

E. coli BL21 were grown at 37ºC 180 rpm in liquid LB supplemented with antibiotics until they reached OD$_{600}$

= 0.6. Then, 0.1 mM IPTG was added, and bacteria were grown for another 20 hours at 18ºC 180 rpm. Finally,

bacteria were harvested and lyzed by sonication. The bacterial lysates were centrifuged for clarification.

Pichia pastoris protein expression was induced with methanol. Briefly, *P. pastoris* cultures were grown in 50

ml of YPD for 48 h. until dextrose was consumed. Then, 200 ul methanol pulses were supplied every 24 h for

5 days. Finally, *P. pastoris* were harvested, and supernatants were collected.

HEK cells were transfected using polyethyleneimine (PEI) for protein expression. Briefly, 30000 cells per well

were seeded in 24 well plates and incubated at 37 °C, 5% CO_2 for 24 h. For transfection, 4 ul of PEI were

diluted in 40 ul of DMEM. 500 ng of DNA was added, and the transfection mix was incubated for 20 min at

room temperature. Then, the transfection mix was transferred to the cells and incubated for 48 h. Finally,

the supernatant was collected to check protein expression.

6xHis-Tag purification method

Purification of 6xHis-tagged proteins was carried out according to the manufacturer's protocol. Briefly,

benchtop columns were equilibrated with buffer (200 mM NaCl, 50 mM Tris-HCl pH 7.5). Columns were

loaded with clarifies of bacterial extracts. After being washed, columns were eluted in steps with 10, 50, 100,

300, and 500 mM imidazole in the equilibration buffer.

To be compared with Bio-Matrix system, the IMAC batch protocol was performed. Ni-NTA superflow resin

(Qiagen) was equilibrated with buffer (200 mM NaCl, 50 mM Tris-HCl pH 7.5) in 1,5 ml tubes. Clarifies of

bacterial lysate were loaded and mixed in an orbital shaker for 1 hour at 4ºC. Samples were centrifugated

and, after being washed, the resin was eluted with 500-mM imidazole in equilibration buffer.

Bio-Matrix preparation

B. subtilis were grown in 200 mL of Luria-Bertani broth at 37ºC 180 rpm for 48 hrs. Then, the culture was

centrifugated and the bacteria were washed twice with PBS. The culture was resuspended in PBS with 2%

glutaraldehyde and incubated overnight with soft agitation. Next, fixed bacteria were washed twice with PBS

and stored in 20 % ethanol.

BM was weighted in a drying scale (KERN MLS-D) and a calibration curve for dry weight vs optical density was

performed.

SLAP$_{TAG}$ purification method

To purify the SLAP-tagged proteins, Bio-Matrix was equilibrated in binding buffer (200 mM NaCl, 50 mM Tris-

HCl pH 7.5). Then, samples were incubated with the Bio-Matrix at different times and temperatures. Samples

were centrifuged (8000 rpm) and the Bio-Matrix was washed 3 times with binding buffer. Finally, samples

were incubated at different times and temperatures with the corresponding solution for elution. Proteins

expressed in HEK293 and *P. pastoris* were purified from the supernatant with the final protocol described in

this paper.

Protein analysis

Protein samples were dissolved in a cracking buffer and incubated for 5 min at 100ºC. Protein electrophoresis

was performed at 120V on 12% SDS-PAGE gel. Gels were stained in a Coomassie-Blue solution (20%

methanol, 10% acetic acid).

For Western Blot analysis, proteins were transferred to a nitrocellulose membrane for 55 min at 15V using a

semi-dry electroblotting transfer unit (Bio-Rad, Hercules, CA, United States). Membranes were incubated for

1 hour with blocking buffer (1% dry skim milk, 0,1% Tween in PBS). Then, membranes were incubated for 1

hour with primary antibody diluted in blocking buffer (1/500). After washing with PBS-0,1% Tween,

membranes were incubated for 1 hour with IRDye fluorophore-labeled secondary antibodies (LI-COR, Lincoln,

NE, United States) diluted in a blocking buffer (1/20000). Finally, membranes were scanned using the Odyssey

Imaging System (LI-COR).

Fluorescence measurements

GFP fluorescence measurements were performed at 485/535nm excitation-emission wavelength

respectively using FilterMax F5 Microplate Reader in Black 96 Well Plates (Thermo).

Protein modeling

(H_6)-GFP-SLAP$_{TAG}$ structure was predicted by AlphaFold[28]. ColabFold web interface was employed using

standard settings (five models and no templates).

Adsorption isotherm

Adsorption isotherms for (H_6)-GFP-SLAP$_{TAG}$ on Bio-Matrix were performed using batch experiments. BM was

equilibrated with binding buffer (50 mM Tris-HCl, 200 mM NaCl, and pH 7.6). Purified (H_6)-GFP-SLAP$_{TAG}$

protein at 3 mg/ml in binding buffer was used as a stock solution. Different concentrations of protein were

incubated with 10 ul of BM. After reaching equilibrium, samples were centrifugated and GFP fluorescence

from the supernatant was measured. Unbound protein in equilibrium with the BM was calculated using GFP

fluorescence. Bound protein was estimated by the difference between input and unbound protein. The

adsorption isotherm data were then fitted to the Langmuir isotherm equation to calculate the parameters

Qmax and K_D.

Confocal Fluorescence Microscopy

Samples were incubated for 30 min in plates treated with poly-L-Lysine (Sigma, St. Louis, MO, United State).

After treatment with PFA (4% in PBS), samples were washed twice with PBS. Finally, samples were observed

with a confocal laser-scanning microscope Olympus FV1000 using a PlanApo N (60 × 1.42 NA) oil objective.

Cleavage of SLAP$_{TAG}$ with SLAP$_{ASE}$ protease

Cleavage buffer recommended for commercial HRV3c protease was prepared: 50 mM Tris-HCl, pH 7.0, 150

mM NaCl, 1 mM EDTA, and 1 mM dithiothreitol. GFP-LEVLFQGP-SLAP$_{TAG}$ protein was bound to the Bio-Matrix.

The sample was washed with cleavage buffer at 4ºC. The protease SLAP$_{ASE}$ was added and incubated at 4ºC.

Percolate containing GFP protein without SLAP$_{TAG}$ was recovered. As a control, the protease and SLAP$_{TAG}$ were

eluted with Bio-Matrix elution buffer after cleavage.

Synthesis of iron nanoparticles

Iron nanoparticles were synthesized by reverse co-precipitation as described by Nadi et al [29] . In summary,

the precursor was prepared by dissolving 0.89 g of FeCl2.4H2O in 90 mL of water. The sample was incubated

in stirring for 15 min and sonicated for 10 min to ensure complete dissolution of the salt. The solution was

then poured over a solution of ammonium hydroxide diluted 1:2 with water and stirred for an hour. Finally,

the sample was washed repeatedly with deionized water.

Magnetic Bio-Matrix generation

Briefly, 5 ml of Bio-Matrix were resuspended in PBS and iron nanoparticles were added at a final

concentration of 40 g/L. The sample was mixed with gentle stirring for 30 minutes. Then, it was washed

repeatedly with PBS 1X and preserved in 20% ethanol.

Antibody generation

Mice were intraperitoneally injected with 10 ug of purified GST-SLAP$_{TAG}$ or GFP in aluminum hydroxide (Imject

Alum ThermoFisher). Two and four weeks later, booster doses of 5 ug of protein were administrated.

Statistical analysis

Statistical analyses were performed using GraphPad Prism 9 software. Statistical significance was analyzed

by one-way ANOVA with Bonferroni.

References

1.   Cai, Z., Poulos, R. C., Liu, J. & Zhong, Q. Machine learning for multi-omics data integration in cancer.

iScience **25**, 103798 (2022).

2.   Hassan, M. *et al.* Innovations in Genomics and Big Data Analytics for Personalized Medicine and

Health Care: A Review. *Int. J. Mol. Sci.* **23**, (2022).

3.   Cong, Y. & Endo, T. Multi-Omics and Artificial Intelligence-Guided Drug Repositioning: Prospects,

Challenges, and Lessons Learned from COVID-19. *Omi. A J. Integr. Biol.* **26**, 361–371 (2022).

4.   Amer, B. & Baidoo, E. E. K. Omics-Driven Biotechnology for Industrial Applications. *Frontiers in*

Bioengineering and Biotechnology vol. 9 at https://doi.org/10.3389/fbioe.2021.613307 (2021).

5.   Zhu, M. M., Mollet, M., Hubert, R. S., Kyung, Y. S. & Zhang, G. G. Industrial Production of Therapeutic

Proteins: Cell Lines, Cell Culture, and Purification. in *Handbook of Industrial Chemistry and*

Biotechnology 1639–1669 (Springer International Publishing, 2017). doi:10.1007/978-3-319-52287-

6_29.

6.   Lu, R. M. *et al.* Development of therapeutic antibodies for the treatment of diseases. *Journal of*

Biomedical Science vol. 27 at https://doi.org/10.1186/s12929-019-0592-z (2020).

7.   Narayanan, H., Sponchioni, M. & Morbidelli, M. Integration and digitalization in the manufacturing

of therapeutic proteins. *Chem. Eng. Sci.* **248**, 117159 (2022).

8.   Khoury, G. El *et al.* Bespoke affinity ligands for the purification of therapeutic proteins. *Pharm.*

Bioprocess. **3**, 139–152 (2015).

9.   Owczarek, B., Gerszberg, A. & Hnatuszko-Konka, K. A Brief Reminder of Systems of Production and

Chromatography-Based Recovery of Recombinant Protein Biopharmaceuticals. *BioMed Research*

International vol. 2019 at https://doi.org/10.1155/2019/4216060 (2019).

10.   Samaras, J. J., Micheletti, M. & Ding, W. Transformation of Biopharmaceutical Manufacturing

Through Single-Use Technologies: Current State, Remaining Challenges, and Future Development.

Annu. Rev. Chem. Biomol. Eng. **13**, 73–97 (2022).

11.   Kuntz, I. D., Chen, K., Sharp, K. A. & Kollman, P. A. The maximal affinity of ligands. *Proc. Natl. Acad.*

Sci. U. S. A. **96**, 9997–10002 (1999).

12.   Mouratou, B., Béhar, G. & Pecorari, F. Artificial affinity proteins as ligands of immunoglobulins.

Biomolecules **5**, 60–75 (2015).

13.   Hais, I. M. Biospecific sorption, Prague, 1910: Emil Starkenstein (1884-1942). *J. Chromatogr. B*

Biomed. Sci. Appl. **376**, 5–9 (1986).

14.   Rodriguez, E. L. *et al.* Affinity chromatography: A review of trends and developments over the past

50 years. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences*

vol. 1157 at https://doi.org/10.1016/j.jchromb.2020.122332 (2020).

15.   Uriza, P. J. *et al.* Development of an Antigen Delivery Platform Using Lactobacillus acidophilus

Decorated With Heterologous Proteins: A Sheep in Wolf's Clothing Story. *Front. Microbiol.* **11**,

(2020).

16.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583 (2021).

17.   Garduno, R. A., Phipps, B. M. & Kay, W. W. Physical and functional S-layer reconstitution in

Aeromonas salmonicida. *J. Bacteriol.* **177**, 2684–2694 (1995).

18.   Mader, C., Huber, C., Moll, D., Sleytr, U. B. & Sára, M. Interaction of the Crystalline Bacterial Cell

Surface Layer Protein SbsB and the Secondary Cell Wall Polymer of Geobacillus stearothermophilus

PV72 Assessed by Real-Time Surface Plasmon Resonance Biosensor Technology. *J. Bacteriol.* **186**,

1758–1768 (2004).

19.   Li, J., Hu, X., Yan, J. & Yuan, Z. Species-specific cell wall binding affinity of the S-layer proteins of

mosquitocidal bacterium Bacillus sphaericus C3-41. *Appl. Environ. Microbiol.* **75**, 3891–3895 (2009).

20.   Lortal, S. *et al. S-layer of Lactobacillus helveticus atcc 12046: isolation, chemical characterization and*

405   *re-formation after extraction with lithium chloride Slayer of Lactobacillus helveticus ATCC 12046:*

406   *isolation, chemical characterization and re-formation after extraction with lithium chloride*. *Society*

407   *for General Microbiology* vol. 138 https://hal.inrae.fr/hal-02713428 (1992).

408   21.   Fina Martin, J. *et al.* Exploring lectin-like activity of the S-layer protein of Lactobacillus acidophilus

409   ATCC 4356. *Appl. Microbiol. Biotechnol.* **103**, 4839–4857 (2019).

410   22.   Kim, I. J., Xu, Y. & Nam, K. H. Spectroscopic Analysis of Fe Ion-Induced Fluorescence Quenching of

411   the Green Fluorescent Protein ZsGreen. *J. Fluoresc.* **31**, 307–314 (2021).

412   23.   Richmond, T. A., Takahashi, T. T., Shimkhada, R. & Bernsdorf, J. Engineered metal binding sites on

413   green fluorescence protein. *Biochem. Biophys. Res. Commun.* **268**, 462–465 (2000).

414   24.   Buxbaum, E. *Enzymes Are Biocatalysts*. *Fundamentals of Protein Structure and Function* (2015).

415   doi:10.1007/978-3-319-19920-7_4.

416   25.   Baldo, B. A. Chimeric Fusion Proteins Used for Therapy: Indications, Mechanisms, and Safety. *Drug*

417   *Saf.* **38**, 455–479 (2015).

418   26.   Fei Wen, Sheryl B. Rubin-Pitel, and H. Z. Engineering of Therapeutic Proteins.Chp7. 153–177 (2009).

419   27.   Hsieh, C.-L. *et al.* Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. *Science (80-.).*

420   **369**, 1501–1505 (2020).

421   28.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589

422   (2021).

423   29.   Nadi, A. *et al.* Immobilisation of bacteria onto magnetic nanoparticles for the decolorisation and

424   degradation of azo dyes. *IET Nanobiotechnology* **13**, 144–149 (2019).

Figures

Figure 1: Characterization of (H$_6$)-GFP-SLAP$_{TAG}$ binding to Bio-matrix. (A) The AlphaFold2 model of (H$_6$)-GFP-

SLAP$_{TAG}$ fusion protein. AlphaFold2 predicted structure was automatically colored by the pLDDT confidence

measure. High accuracy is colored in blue while low accuracy in red. Despite SLAP$_{TAG}$ has not been crystallized,

structure prediction showed a good performance. (B) Figure compares (H$_6$)-GFP-SLAP$_{TAG}$ purification yield of

SLAP$_{TAG}$-based affinity chromatography for different temperatures of the binding process. (C) Figure

compares (H$_6$)-GFP-SLAP$_{TAG}$ purification yield of SLAP$_{TAG}$-based affinity chromatography for different times of

binding incubation. (D) Adsorption isotherm of (H$_6$)-GFP-SLAP$_{TAG}$ onto Bio-Matrix. Asterisk (****) denotes

significant difference using the ANOVA method, Bonferroni test ($p < 0.0001$).

Figure 2: Characterization of (H$_6$)-GFP-SLAP$_{TAG}$ elution. (A) Figure compares (H$_6$)-GFP-SLAP$_{TAG}$ purification

yield of SLAP$_{TAG}$-based affinity chromatography for different temperatures of elution. (B) Figure compares

(H$_6$)-GFP-SLAP$_{TAG}$ purification yield of SLAP$_{TAG}$-based affinity chromatography for different times of elution

incubation. (C) Figure compares (H$_6$)-GFP-SLAP$_{TAG}$ purification yield of SLAP$_{TAG}$-based affinity chromatography

system for different eluents. Asterisk denotes significant difference using the ANOVA method, Bonferroni

test (*** $p = 0,0003$; **** $p < 0.0001$).

Figure 3: Optimized protocol for (H$_6$)-GFP-SLAP$_{TAG}$ purification using the Bio-matrix and comparison with

the high-performance Ni^{2+}-charged agarose matrix. (A) Graphical description of the SLAP$_{TAG}$-based affinity

chromatography protocol. (B) Confocal microscopy of the Bio-Matrix. *Bacillus subtilis* red native

autofluorescence is observed. (C) Confocal microscopy of (H$_6$)-GFP-SLAP$_{TAG}$ bound to the Bio-Matrix. GFP is

visualized on the *Bacillus* surface. (D) Confocal microscopy of the Bio-Matrix after elution of (H$_6$)-GFP-SLAP$_{TAG}$.

(E) Tubes containing purification fractions seen under UV light. GFP input fluorescence is recovered in the

eluate. (F) Coomassie blue staining analysis of (H$_6$)-GFP-SLAP$_{TAG}$ scaled purification. BM = Bio-Matrix; MK =

protein marker (KDa); IN = input; FT = flowthrough; W = wash; E = elution. (G) Immobilized metal affinity

chromatography (IMAC) and SLAP$_{TAG}$-based affinity chromatography system are compared in their capacity

to purify (H$_6$)-GFP-SLAP$_{TAG}$. Relative fluorescence of total (H$_6$)-GFP-SLAP$_{TAG}$ recovered in elution fraction is

shown.

Figure 4: Representation of SLAP-tagged recombinant protein structure. Lys = bacteriophage T4 lysozyme;

GFP = *Aequorea victoria* green fluorescent protein; GST = glutathione-s-transferase; HRV-3c = human

Rhinovirus 3C Protease; EspA = *E. coli* EspA protein; Omp19 = *B. abortus* Omp19 protein; MBP = maltose-

binding protein; Gal8 = mouse Galectin-8; EITH7 = EspA, Intimin and Tir fusion protein from the Shiga-toxin

producing *E. coli*; Spike = SARS-CoV2 Spike protein.

Figure 5: Analysis of different SLAP-tagged protein purifications expressed in *E. coli*. Coomassie Blue stained

SDS-PAGE of purification fractions of SLAP$_{TAG}$-based affinity chromatography for (A) FliC-SLAP$_{TAG}$ (B) EspA-

SLAP$_{TAG}$ (C) Gal8-SLAP$_{TAG}$ (D) GST-SLAP$_{TAG}$ (E) Lys -SLAP$_{TAG}$ (F) Omp19-SLAP$_{TAG}$ (G) EITH7-SLAP$_{TAG}$ (H) SLAP$_{ASE}$

(human rhinovirus 3c fused to SLAP$_{TAG}$). MK = protein marker (KDa); IN = input; FT = flowthrough; W = wash;

E = elution.

Figure 6: Western blot analysis of purification of SLAP-tagged proteins using different expression systems.

(A) Purification of SLAP$_{ASE}$ (human rhinovirus 3c fused to SLAP$_{TAG}$) protease expressed in *Pichia pastoris*. (B)

Purification of SARS CoV2 Spike-SLAP$_{TAG}$ protein expressed in HEK293 cells. MK = protein marker (KDa); IN =

input; FT = flowthrough; W = wash; E = elution.

Figure 7: Analysis of Bio-Matrix stability in time and reuse. (A) The figure compares (H$_6$)-GFP-SLAP$_{TAG}$

purification yield of SLAP$_{TAG}$-based affinity chromatography at different times when it is conserved at -20ºC.

(B) Figure compares (H$_6$)-GFP-SLAP$_{TAG}$ purification yield of SLAP$_{TAG}$-based affinity chromatography for

different cycles of reuse. Asterisk denotes significant difference using the ANOVA method, Bonferroni test

(** p = 0,0059; *** p = 0,0002; **** p < 0.0001)

Figure 8: SLAP$_{TAG}$ removal. (A) Graphical description of the protocol to remove the SLAP$_{TAG}$ using SLAP$_{TAG}$-

based affinity chromatography. SDS-PAGE of purification fractions of SLAP$_{TAG}$ removal protocol stained with

Coomassie blue solution (B) or under UV light (C). Western Blot of purification fractions of SLAP$_{TAG}$ removal

protocol revealed with anti- SLAP$_{TAG}$ (D) or anti-GFP (E) antibodies. SLAP$_{ASE}$ = purified protease; IN = input; FT

= flowthrough; E = elution; MK = protein marker (KDa)

Figure 9: GFP-SLAP$_{TAG}$ purification using Magnetic Bio-Matrix. (A) Graphical description of the purification

protocol for magnetic Bio-Matrix. (B) Photo of 1,5 tubes ml containing fractions of GFP-SLAP$_{TAG}$ magnetic

purification under UV light. (BM$_{mag}$ = magnetic Bio-Matrix; IN = input; FT = flowthrough; W = wash; E =

elution)

TABLE 1

Strain		
E. coli DH5α	F−φ80*lac*ZΔM15 Δ(*lac*ZYA-*arg*F) U169 *rec*A1 *end*A1 *hsd*R17(rK−, mK+) *pho*A *sup*E44λ- *thi*-1 *gyr*A96 *rel*A1	Invitrogen
E. coli BL21 Codon plus	[*omp*T *hsd*S(rB− mB–) *dcm* + Tcr *gal* λ (DE3) *end*A Hte Cmr]	Stratagen
B. subtilis	Wild type strain var. *natto*	ATCC 15245
Pichia pastoris	*P. pastoris* strain GS115	BioRad, USA
Plasmid		
pET28-eGFP-SlpA	A vector containing GFP gene from *Aequorea victoria* fused to SLAP$_{TAG}$	[21]
pET22-eGFP-SlpA	A vector containing GFP gene from *Aequorea victoria* fused to LEVLFQGP sequence and SLAP$_{TAG}$	This study
pGEX-SlpA	A vector containing GST form *Shitosoma japonicum* fused to SLAP$_{TAG}$	[15]
pLC3-EITH7-SlpA	A vector containing EspA-Intimin-Tir fusioned genes from STEC *E. coli* fused to SLAP$_{TAG}$	[15]
pLC3-Omp19-SlpA	A vector containing Omp19 gene from *Brucella abortus* fused to SLAP$_{TAG}$	[15]
pLC3-FliC-SlpA	A vector containing FliC gene from *Salmonella enterica* fused to SLAP$_{TAG}$	[15]
pLC3-Gal8-SlpA	A vector containing Gal8 gene from *Mus musculus* fused to SLAP$_{TAG}$	[15]
pET28-Lys-SlpA	A vector containing Lysozyme gene from *Gallus gallus* fused to SLAP$_{TAG}$	This study
pMAL-c5X-HRV3c-SlpA	A vector containing human rhinovirus (HRV) type 14 3C protease gene fused to SLAP$_{TAG}$ (SLAPase)	This study
pGEX-2T-EspA-SlpA	A vector containing the EspA gene from STEC *E. coli* fused to SLAP$_{TAG}$	This study
pPICZalphaB-HRV3c-SlpA	A vector containing human rhinovirus (HRV) type 14 3C protease gene fused to SLAP$_{TAG}$ (SLAPase)	This study
Addgene #154754	A vector containing the sequence of SARS CoV2 Spike ectodomain Hexa-pro	[27]
pSpike-SLAP$_{TAG}$	Plasmid Addgene #154754 fused with the SLAP$_{TAG}$	This study
pLysS	Vector for expression of T7 lysozyme.	Novagene

Figure 1

Figure 2

Figure 3

A

1 Equilibrate BM in Binding Buffer

2 Load sample, mix and incubate
5 min
4 °C
Sample

3 Centrifugate and wash pellet twice with Washing Buffer. Discard supernatant

4 Resuspend pellet in Elution Buffer to elute purified protein
5 min
24 °C

5 Centrifugate and keep supernatant

B

10 µm

C

10 µm

D

10 µm

E

BM
IN
FT
W
E

F

MK   IN   FT   W   E
100
70
50
40 ← 46 KDa
30
25

G

Relative fluorescence units (RFU)

5×10^7
4×10^7
3×10^7
2×10^7
1×10^7
0

IMAC   SAC

Figure 4

Figure 5

Figure 6

A: IN FT W E MK — 100, 70, 50, 40, 30, 25; 49 KDa

B: IN FT W E MK — 161 KDa; 100, 70, 50, 40, 30

Figure 7

Figure 8

A

1. Equilibrate BM in Binding Buffer

2. Load sample, mix and incubate — 5 min, 4 °C — Sample

3. Centrifugate and wash pellet twice with Washing Buffer. Discard supernatant

4. Resuspend pellet in SLAP$_{ASE}$ Buffer and add SLAP$_{ASE}$ protease to elute purified protein — 60 min, 4 °C

5. Centrifugate and keep supernatant

B

SLAP$_{ASE}$ IN FT E MK

C

SLAP$_{ASE}$ IN FT E MK

D

SLAP$_{ASE}$ IN FT E MK

ANTI SLAP

E

SLAP$_{ASE}$ IN FT E MK

ANTI GFP

Figure 9

A

1. Equilibrate BM$_{mag}$ in Binding Buffer
2. Load samble, mix and incubate — 5 min, 4 °C, Sample
3. Transfer to magnete and wash pellet twice. Discard supernatant
4. Resuspend pellet in Elution Buffer to elute purified protein — 5 min, 24 °C
5. Transfer to magnete and keep supernatant

B

IN FT W E MK

C

BM$_{mag}$ IN IN + BM$_{mag}$ FT + BM$_{mag}$ W + BM$_{mag}$ E + BM$_{mag}$ E