1

Longitudinal monitoring of maternal care and maternal neglect

Luisa Schuster[1-3], Renee Henderson[1-3], Shreya Sankar[1-3], Deepasri Ananth[1-3], Matilda Kirk[1-3],

Paola Leone[1-3], Karen E. Adolph[4,5], Robert C. Froemke[1-4]*, and Adam Mar[1,2]*

[1] Neuroscience Institute,

[2] Department of Neuroscience and Physiology,

[3] Department of Otolaryngology, New York University Grossman School of Medicine, New

York, NY, USA.

[4] Center for Neural Science,

[5] Department of Psychology, New York University, New York, NY, USA.

.

* Correspondence to: A. Mar, phone: 212-263-9295; email: adam.mar@nyulangone.org

and: R.C. Froemke, phone: 212-263-4082; email: robert.froemke@med.nyu.edu

Parental care is required to ensure the health and survival of offspring. Caretaking in mammals is challenging as infants are largely helpless and require near-continual oversight over prolonged periods. Mouse pups in particular cannot thermoregulate, and will succumb to hypothermia unless kept warm by an adult caretaker or an insulating nest. Parents must balance caretaking behaviors with other activities important for survival such as regulating their own temperature and foraging for food or water, which necessarily involve periods of pup neglect. To understand in high detail the consequential decision-making involved in care of infant pups, we built a new system for long-term 24/7 continuous monitoring of mouse homecage behaviors, including thermal imaging, video recording from above, side and directly under the nest, with environmental controls to standardize housing conditions across animals. We monitored single dams and their offspring, before, during and after parturition, from initial mating over four consecutive litters. We found that about half of the animals had high litter survival rates ('high-pup-survival' dams) but the other half had little to no pups survive ('low-pup-survival' dams). The relative amount of litter survival did not change or improve across litters in absence of intervention: Low-pup-survival dams continued to neglect pups and nest-building across litters, losing nearly all offspring within a day after birth. We could predict which animals would become low-pup-survival dams by inspecting the nests they built a few days before parturition. Low-pup-survival dams minimally engaged with nest building, and after birth, generally avoided pups and the nest. Nest quality deteriorated due to movement of the dam in and out of the nest, exacerbating temperature loss and mortality of the litter. In contrast, high-pup-survival dams spent a considerable amount of time rebuilding and adjusting the nest over the first postnatal day and beyond. Remarkably, if we co-housed low-pup-survival dams with a high-pup-survival dam and her

litter, when the low-pup-survival dam was bred again and singly-housed, litter survival rates thereafter were consistently much higher along with increased attention toward pups and nest. Thus, even under well-controlled and ideal lab housing conditions, mouse maternal care can be remarkably variable, with continued nest rebuilding as the most important factor in ensuring high litter survival rates. Furthermore, despite some animals perseverating in maternal neglect and losing their pups, caretaking and nest-building can improve after interactions with experienced and successful parents.

Main Text

Parental behavior is essential for the well-being of offspring and the survival of the species (Rilling and Young, 2014). Adult animals can become highly sensitive to the needs of their infants to adequately care for them, as infants require intense continuous protection for days to years depending on species, and single parenting is especially challenging. Mouse pups in particular rely on their mother for sustenance, thermoregulation, and protection, supported by successful nest-building (Lynch and Possidente, 1978). However, pup mortality can result if dams are neglectful or provide inadequate care during the postnatal period. Previous studies of rodents and other species report a remarkably high degree of infant mortality, both outdoors in native habitats and animals housed in laboratory conditions (Brajon et al., 2021). We aimed to determine when and how pup mortality occurs, if it is due to parental neglect or other factors, and if mouse mothers might learn from certain experiences to improve maternal care over subsequent litters.

We designed a high-resolution system to monitor parental care and family life in mice with multiple cameras, recording from homecages continuously 24 hours/day for weeks to months

across consecutive litters (**Fig. 1A**). Individual cages were contained within custom-designed

environmental control boxes, containing at least two camera mounts to record the nest from the

side, underneath, and overhead views, including infrared cameras to record in darkness and a

thermal imaging camera to monitor temperature. Video was combined with an inexpensive, open-

source system to control LED light-strips to provide a naturalistic light/dark cycle, together with

temperature and humidity sensors connected to an Arduino circuit board that controls two fans,

and adjusts their speed to maintain the enclosure temperature at 26-28 °C (**Fig. 1B**), a level close

to adult mouse thermoneutrality of 30°C (Król & Speakman, 2003; Gaskill et al., 2012). Data were

saved to local servers and streamed remotely for machine learning-based quantitative behavioral

analysis along with annotations by several independent experienced human raters.

We monitored a total of 38 single-housed female mice continuously over 3-4 months, from first

mating and first litter through subsequent breeding and litters (**Fig. 1C**, top). Although many

different maternal behaviors were documented and scored over the postnatal period, we focused

on interactions with pups and the nest in the first few days of each litter (**Fig. 1C**, bottom). This

was due largely to surprising high pup mortality even in the protected conditions of the lab

vivarium. We quantified litter survival from the moment of birth over the first 24 postnatal hours,

in which most mortality was observed. For the first cohort of 15 mice, the average survival of the

first litter was 62.2±12.1%. However, litter survival was clearly bimodal—7/15 first-time dams

had 100% pup survival, and 5/15 dams had 0% pup survival (**Fig. 1D**, Litter 1). We refer to dams

with >50% litter survival as 'high-pup-survival' dams, and dams with <50% pups surviving as

'low-pup-survival' dams.

We hypothesized that for low-pup-survival dams, litter survival rates might improve over

subsequent litters as dams learned from experience or had other forms of maternal adaptation in

the service of improved caregiving. However, this was not the case. As shown in **Figure 1D**, litter

survival of singly-housed C57Bl/6 mice continued to be bimodal, with 9/15 high-pup-survival

dams and 6/15 low-pup-survival dams for litter 2, 5/15 low-pup-survival dams for litter 3, and 4/8

low-pup-survival dams for litter 4 (**Fig. 1D**, left). With the exception of a single animal, high-pup-

survival dams continued to have high litter survival rates across all 3-4 monitored litters, and low-

pup-survival dams consistently lost most or all of their litters across all monitored litters (**Fig. 1D**,

right). This meant that the litter survival rate of the first litter predicted with high accuracy the

survival of subsequent litters (**Fig. 1E**; $N=15$, $r \geq 0.8$, $p<0.001$).

Inspection of video recordings showed that almost none of the litter mortality observed in either

high- or low-pup-survival dams was due to aggression or mauling of otherwise healthy pups

(observed in 3/38 dams). Instead, low-pup-survival dams neglected pups and nests after initial

nest-building was complete. In contrast, high-pup-survival dams consistently re-engaged in nest-

building, especially because entering and leaving the nest sometimes disrupted nest integrity. We

found that hand-scoring nest quality 1-3 days prior to parturition also predicted litter survival

across dams (**Fig. 2A**).

The importance of nest quality likely reflects the function of nests to keep pups insulated especially

when the dam is not present in the nest, because pups do not reliably thermoregulate their body

temperature and can perish from hypothermia (Weber and Olsson, 2008). However, it proved

challenging to reliably hand-score nests over extended periods, because nest structure was dynamic

and the visual features of a nest might not perfectly relate to the thermal insulative qualities that

nests must provide. Therefore, we used thermal imaging to monitor pup temperature directly. This

required us to segment the nest from the video recording and use it to isolate the temperature of

the pups within the nest (**Fig. 2B**).

We monitored pup temperature during periods when dams made excursions from the nest. This

happened frequently as mother animals left to forage for food and water, and to regulate their own

body temperature as they can overheat if they stay too long on the nest (Leon et al., 1978; Gamo

et al., 2016; Król and Speakman, 2019). We focused on moderate-duration excursions of >23

seconds, averaging pup temperature and heat loss over this period. We found that both the initial

temperature and the temperature drop predicted pup mortality: higher tonic temperatures and less

average heat loss were correlated with higher pup survival (**Fig. 2C,D**).

On average the total time spent on the nest was unrelated to litter survival rates (**Fig. 2E**). Initial

time spent nest-building also was uncorrelated with litter survival (**Fig. 2F**). Instead, the main post-

partum predictor of pup survival was amount of time the dam spent rebuilding the nest (**Fig. 2G**).

Low-pup-survival dams typically spent some time nest-building initially, but then gave up, and

subsequently lost most or all of their offspring within hours. High-pup-survival dams continued to

adjust and rebuild the nest especially after leaving the nest and disrupting nest integrity. We also

noticed that many high-pup-survival dams stocked food pellets near the nest (**Fig. 2H**). This

seemed less related to shorter foraging times and more related to structural bolstering of nest wall

integrity due to the weight of the food pellets.

It was surprising that over multiple rounds of breeding, low-pup-survival dams did not increase

attention to pups and time spent rebuilding their nests. We wondered if low-pup-survival dams

somehow could not provide adequate caregiving under any circumstance, or if instead perhaps

with different experiences these animals could become less neglectful and have higher survival

rates. Previously we documented the emergence of alloparenting in initially-neglectful and pup

averse virgin female C57Bl/6 mice, after being co-housed with experienced dams and litters

(Marlin et al. 2015; Carcea et al. 2021). In this case for alloparenting, behavioral changes in the

virgin were due to encouragement or coercion by the experienced dam, who shepherded or forced

the virgin to spend time in the nest until eventually the virgin became a successful alloparent.

We reasoned that similar parental behavior might emerge if we co-housed an initially low-pup-

survival dam with an experienced dam and her litter. We identified a separate cohort of low-pup-

survival dams with 0% litter survival at litter 1 and <50% litter survival for litter 2. Prior to mating

these animals for litter 3, we co-housed some of these animals for 16-20 days with an experienced

high-pup-survival dam and litter; other animals were kept isolated for a similar duration. We then

bred these animals and monitored pup survival for litters 3 and 4 when they were returned to being

singly-housed in our behavioral monitoring chambers (**Fig. 3A**).

We found that temporarily co-housing 'low-pup-survival' dams with another experienced mother

led to much higher survival rates of litters 3 and 4, compared to the other animals who were just

kept isolated for 16-20 days (**Fig. 3B**; red, co-housed dams, litter 3 survival: 88.5±4.8%, N=9;

black, isolate dams, litter 3 survival: 2.5±5.6%, N=5, $p<10^{-4}$ compared to litter 3 survival of co-

housed animals, Student's unpaired two-tailed t-test). Inspection of video recorded during the co-

housing period revealed that while the low-pup-survival dams were initially pup-averse, within

hours to days they began interacting with pups and helped with childcare. This seemed to be due

to behaviors of the other experienced dam, who shepherded the low-pup-survival dam into the nest

and seemingly demonstrated aspects of maternal care (Carcea et al. 2021). This improvement of

maternal care in the low-pup-survival dams did not seem to be due to, e.g., social buffering effects

from co-housing with any other animal. Co-housing the low-pup-survival dam between litters 2

and 3 with either a pup-naïve virgin female or an experienced paternal male did improve pup

survival for litter 3 (**Fig. 3C**); instead, there was a significant reduction in litter survival after co-

housing with a naïve virgin female. Co-housing in these cases occurred without pups, as none of

the adult animals would be lactating.

Our study provides insights into mouse maternal behavior over time and across consecutive litters,

the impact of the nest and its insulating capacity on pup survival, and the role of socially-

transmitted parental behavior via co-housing with an experienced mother animal. These

observations resulted from the 'documentary' approach to monitoring animal behavior

continuously over extended periods of days to months, with multiple cameras including under the

nest. This was essential to accurately capture moments of parturition, to identify the number of

pups delivered by each dam and the caregiving behaviors or neglect that resulted.

Our findings indicate that even under the controlled and protected conditions of a laboratory

vivarium, childcare can be immensely challenging and taxing, requiring continued effort on the

part of caregivers to ensure pups are fed and warm. Under the conditions of our environmentally-

controlled home cages, approximately a third to a half of C57Bl/6 mouse mothers were observed

to lose most to all of their litters due to neglect of pups and nest. The nest structure was quite

dynamic, needing to be repeatedly adjusted or rebuilt due to the movement of the mother in and

out of the nest, and the motion of the pups. Thermal imaging was important to relate this aspect of

parental behavior to litter survival, as pups rapidly dropped in temperature without a well-

constructed insulating nest and died within minutes.

One potential concern is that the extent of pup mortality and relatively high prevalence of low-

pup-survival dams is a consequence of sustained single-adult housing especially combined with

the stress of childcare. A related caveat is that our studies were performed exclusively in C57Bl/6

mice, and it is known that there are differences in parental behaviors across rodent strains (Ward,

1980). Even accounting for the genetic background, within the C57Bl/6 strain there was

considerable bi-modality in childcare abilities, which might reflect other individual differences in

gene expression and/or neural responses to infant distress cues that will need to be further

documented. Most wild species of mice are believed to live in multi-female colonies (Ferrari et

al., 2019) where perhaps many different adults can participate in childcare, tending to pups and

nests, and shepherd or help other inexperienced adults take care of offspring. However, little is

known about the fine-scale adult social dynamics of colony organization and how this relates to

communal child-care. The system we have developed enables a systematic and parametric

exploration of different variables related to social structure and housing conditions, and promises

to reveal much more about the family life of lab mice and other species.

References and Notes

Brajon, S., Morello, G. M., Capas-Peneda, S., Hultgren, J., Gilbert, C., & Olsson, A. (2021). All
the pups we cannot see: Cannibalism masks perinatal death in laboratory mouse breeding
but infanticide is rare. *Animals*, *11*(8), 2327. https://doi.org/10.3390/ani11082327

Carcea, I., Caraballo, N. L., Marlin, B. J., Ooyama, R., Riceberg, J. S., Mendoza Navarro, J. M.,
Opendak, M., Diaz, V. E., Schuster, L., Alvarado Torres, M. I., Lethin, H., Ramos, D.,
Minder, J., Mendoza, S. L., Bair-Marshall, C. J., Samadjopoulos, G. H., Hidema, S.,
Falkner, A., Lin, D., … Froemke, R. C. (2021). Oxytocin neurons enable social
transmission of maternal behaviour. *Nature*, *596*(7873), 553–557.
https://doi.org/10.1038/s41586-021-03814-7

Ferrari, M., Lindholm, A. K., & König, B. (2019). Fitness consequences of female alternative
reproductive tactics in house mice (*mus musculus domesticus*). *The American Naturalist*,
193(1), 106–124. https://doi.org/10.1086/700567

Gamo, Y., Bernard, A., Troup, C., Munro, F., Derrer, K., Jeannesson, N., Campbell, A., Gray,
H., Miller, J., Dixon, J., Mitchell, S. E., Hambly, C., Vaanholt, L. M., & Speakman, J. R.
(2016). Limits to sustained energy intake XXIV: Impact of suckling behaviour on the body
temperatures of lactating female mice. *Scientific Reports*, *6*(1).
https://doi.org/10.1038/srep25665

Gaskill, B. N., Gordon, C. J., Pajor, E. A., Lucas, J. R., Davis, J. K., & Garner, J. P. (2012). Heat
or insulation: Behavioral titration of mouse preference for warmth or access to a nest. *PLoS
ONE*, *7*(3). https://doi.org/10.1371/journal.pone.0032799

Król E., & Speakman, J. R. (2003). Limits to sustained energy intake VI. energetics of lactation
in laboratory mice at thermoneutrality. *Journal of Experimental Biology*, *206*(23), 4255–
4266. https://doi.org/10.1242/jeb.00674

Król, E., & Speakman, J. R. (2019). Switching off the furnace: Brown Adipose Tissue and
lactation. *Molecular Aspects of Medicine*, *68*, 18–41.
https://doi.org/10.1016/j.mam.2019.06.003

Leon, M., Croskerry, P. G., & Smith, G. K. (1978). Thermal control of mother-young contact in
rats. *Physiology & Behavior*, *21*(5), 793–811. https://doi.org/10.1016/0031-
9384(78)90021-5

Lynch, C. B., & Possidente, B. P. (1978). Relationships of maternal nesting to thermoregulatory
nesting in house mice (mus musculus) at warm and cold temperatures. *Animal Behaviour*,
26, 1136–1143. https://doi.org/10.1016/0003-3472(78)90103-3

Marlin, B. J., Mitre, M., D'amour, J. A., Chao, M. V., & Froemke, R. C. (2015). Oxytocin
enables maternal behaviour by balancing cortical inhibition. *Nature*, *520*(7548), 499–504.
https://doi.org/10.1038/nature14402

Rilling, J. K., & Young, L. J. (2014). The biology of mammalian parenting and its effect on
offspring social development. *Science*, *345*(6198), 771-776.
https://doi.org/10.1126/science.1252723

Ward, R. (1980). Some effects of strain differences in the maternal behavior of inbred mice.
Developmental Psychobiology, *13*(2), 181–190. https://doi.org/10.1002/dev.420130209

Weber, E. M., & Olsson, I. A. (2008). Maternal behaviour in mus musculus sp.: An ethological
review. *Applied Animal Behaviour Science*, *114*(1-2), 1–22.
https://doi.org/10.1016/j.applanim.2008.06.006

Acknowledgements: We thank D. Lin and R.M. Sullivan for comments, discussions, and

technical assistance. This work was funded by NICHD (HD088411 to R.C.F.), the BRAIN

Initiative (NS107616 to R.C.F. and A.M.), and NSF (a graduate research fellowship to L.S.).

Author contributions: L.S. and A.M. built the 24/7 monitoring system. L.S. performed behavioral

experiments. L.S., R.H., S.S., D.A., M.K., and P.L. performed behavioral analysis. K.E.A.

provided analysis tools and data archiving. L.S., R.C.F., K.E.A., and A.M. designed the study and

wrote the paper.

Materials and Methods

All procedures were approved under NYU Grossman School of Medicine IACUC protocols.

System for Long-Term 24/7 Behavioral Analysis. For long-term monitoring of the acquisition and expression of maternal behavior in single-housed animals, we used 8–10-weeks-old C57BL/6 naïve virgin female mice. No statistical methods were used to predetermine sample size. The experiments were not randomized. The behavioral raters were blinded to litter survival rate, allocation during co-housing, and outcome assessment.

A virgin female mouse was housed alone in a 27.94 × 27.94 × 27.94 cm plexiglass homecage covered with plentiful bedding material, food pellets, hydrogel and/or a water bottle (refreshed every 3-5 days), and one Nestlet. We began by recording continuous video of the naïve virgin behavior while housed alone for up to 5 days, before introducing an experienced C57BL/6 male mouse for breeding afterwards. The male mouse was removed from the enclosure after successful intercourse, confirmed through visual examination of video footage, typically after 24-48 hours. For further confirmation of mating, when removing the male mouse from the enclosure, we also checked the virgin for a vaginal plug and we closely monitored the female mouse for physical and behavioral changes typical of a gravid animal. If by day 10 post-mating, the female mouse did not show weight-gain or weight-redistribution, a new experienced male was introduced in the homecage for mating.

Dams were allowed to stay with the pups until weaning on P21 in the same homecage. After that, the female was placed back in the enclosure for a few days and mated again, following the same procedure as with the first litter. This was performed for four consecutive litters.

Video Acquisition. Video recordings were acquired from 1-3 cameras (OV5647 sensor) sensitive

to infrared light and outfitted with 160 FOV wide-angle lenses and placed either above the

enclosure (overhead view), perpendicular to the nest (nest view), or below the nest (bottom-up

view). Each camera was connected to a Raspberry Pi 4 single board computer that captured,

streamed, and saved uninterrupted video from the enclosures (30 frames per second) to a local

server using the Python library OpenCV. For some animals, videography was performed

continuously at all times up to weaning of the third litter; for other animals, we focused video

recording on 3-5 days before birth up until P5.

Behavioral Analysis. For analysis of video data, we used two different DeepLabCut (DLC)

networks: one trained to track the location of the dam and another one to track the pups. The

location of the nest was extracted with custom Matlab code using image segmentation, and we

processed frame-by-frame pose data obtained with DLC to identify episodes of dam nest entry or

exit. Nest entry time was defined as the moment when the two front paws entered the nest cavity.

We also used nests as a region of interest (ROI), to identify when a pup was out of the nest and

how long it took for the dam to it.

We used Datavyu to manually score other behaviors. Five independent coders identified

relevant individual and social behaviors and scored the videos blind to litter number and survival

rate. The results from each rater were compared, compiled, and cross-validated.

Thermal Imaging. For thermal imaging, we recorded continuous thermograms with a thermal

camera (FLIR C3) placed above the nest, streaming temperature data (15 frames per second) to a

Windows PC through a USB cable. Thermographic data were captured with the FLIR Tools software, timestamped, and saved as a csv file.

We synchronized the thermograms and the video recordings post-hoc using timestamp information. We isolated the episodes in which the dam was out of the nest, identified the video frame and thermogram corresponding to the onset of the episode, located the nest opening using image segmentation, and then extracted and averaged the corresponding pixel temperature information from that ROI. We saved these frame-by-frame average traces as csv files. For pup temperature analysis, we identified episodes in which pups were out of the nest and followed a similar procedure to the nest temperature analysis focusing on the out-of-nest pup as our ROI.

Litter Mortality. We monitored both the videos and the thermograms daily to establish how many pups were born alive, how many were stillbirths, and how many were cannibalized. We used color, temperature, movement, and physical integrity to determine if pups were alive. We calculated the litter survival rate as surviving pups on P5 divided by the total number of pups born alive.

Co-Housing. Dams with 0% litter survival during their first litter and less than 50% survival during their second litter were co-housed either with an experienced high-pup-survival dam with her pups (~P5 and older), a virgin, a paternal male, or left in isolation. Co-housing was performed for 16-20 days, before being mated a third and fourth time.

Data and Code Availability. The data that support the findings of this study are available on Databrary. Custom code are available on Github.

318 **Figures and Figure Legends**

Figure 1. Continuous long-term monitoring of mouse maternal care across litters. **A**. Schematic

of setup for 24/7 continuous monitoring of homecage behaviors. A standard mouse homecage was

placed within a custom enclosure containing cameras from top, side, and beneath the nest location.

B. System for long-term continuous imaging and environmental controls. **C**. Homecage behaviors

were monitored over four litters, scored by automated methods (in/out of nest) and manually (nest

building, nest manipulation, cannibalism). Top, timeline of monitoring of individual animals

(times are ages of the dams). Bottom, example ethograms showing several hours of spontaneous

homecage behaviors for the same animal during litters 1, 2, and 3. **D**. Litter survival at 24 hours

post-parturition. Left, fraction of pups surviving after the first postnatal day (N=15 dams). Each

gray circle represents one dam's litter, with the same animals monitored over litters 1-4. 7/15 dams

had 100% survival at litter 1 ('high-pup-survival dams'), and 6/7 of those animals had 100%

survival at litter 3 as well. 5/15 dams had 0% litter survival at litter 1, and 4/5 of those animals had

0% survival at litter 3. Right, same data with lines connecting litter survival from the same dam

across all 3-4 monitored litters. **E**. Correlation of litter survival over litters. Left, litter 1 survival

predicted litter 2 survival across dams (N=15, r(DF)=.80, p=.0003). Right, litter 1 survival also

predicted litter 3 survival (N=15, r(DF)=.96, p<0.0001).

Figure 2. Nests must be rebuilt to ensure pup thermoregulation. **A**. Visual scoring of nest quality;

higher nest integrity scores were correlated with higher litter survival rates (r(DF)=.69, p<.001).

B. Procedure for monitoring pup temperature. Video frames were used to compute nest mask,

registered with thermal imaging to isolate pup temperatures. **C**. Examples of heat loss when dam

leaves the nest; curves show average drop in pup temperature over many excursions of the dam

from the nest (8+ seconds in duration). Top, high-pup-survival dam; bottom, low-pup-survival

dam. Insets show representative nest structures from these animals. **D**. Pup temperature when dam

left the nest. Top: litter survival was not correlated with the initial temperature upon dam leaving,

345   r(DF)=.22, p=0.49. Bottom: litter survival was correlated with temperature drop after dam left the

346   nest, r(DF)=-.84, p=.007). **E.** Dam interaction with the nest. Top: litter survival was not correlated

347   with total time spent in the nest on the first full day after parturition, r(DF)=.006, p=.98. Bottom:

348   litter survival was correlated with amount of time dams spent rebuilding the nest, r(DF)=.65,

349   p=.04.

Figure 3. Co-housing with an experienced high-pup-survival dam improves maternal care. **A,**

Timeline for co-housing studies: virgin females were bred and those dams with 0% litter survival

for litter 1 and <50% survival for litter 2 were selected for these studies; dams were singly-housed

for all four litters. Between litters 2 and 3, dams were divided into four groups and placed in a

separate cage for 16-20 days before the third round of mating ('+Dam': housed with an

experienced high-pup-survival dam and her litter; 'Non': non-co-housed, i.e., singly housed

without pups; '+Virg': co-housed with a pup-naïve virgin female but no pups; '+Male': co-housed

with an experienced paternal male but no pups). **B**, Co-housing with an experienced high-pup-

survival dam led to improved litter survival of initially low-pup-survival dams. Litter survival of

all animals for litter 1: 0%, for litter 2: 10.2±3.7% (N=14). Nine of these animals were co-housed

prior to mating for litter 3; litter survival for litter 3: 88.5±4.8% (N=9), for litter 4: 100% (N=2).

For the five non-co-housed animals from the same cohort, litter survival remained low; for litter

3: 2.5±5.6% (N=5, $p<10^{-4}$ compared to litter 3 survival of co-housed animals, Student's unpaired

two-tailed t-test), for litter 4: 0% (N=4). Litter 2 survival for these two groups was not significantly

different (dams to be co-housed, litter 2 survival: 11.6±5.3%; non-co-housed dams, litter 2

survival: 7.7±4.8%; p=0.60, Student's unpaired two-tailed t-test). **C**, Comparison of litter survival

for litters 2 and 3 per group ('+Dam', litter 2 survival: 11.6±5.3%, litter 3 survival: 88.5±4.8%,

N=9, $p<10^{-4}$ significant increase, Student's paired two-tailed t-test; 'Non', litter 2 survival:

7.7±4.8%, litter 3 survival: 2.5±5.6%, N=5, p=0.45; '+Virg', litter 2 survival: 25.5±5.1%, litter 3

survival: 11.0±7.0%, N=5, p=0.02 significant decrease; '+Male', litter 2 survival: 14.1±7.3%, litter

3 survival: 16.2±8.2%, N=5, p=0.80). **, p<0.01; *, p<0.05.