Calcium signals tune AMPK activity and mitochondrial homeostasis in dendrites of developing neurons

Condensed title: Ca²⁺-AMPK tunes mitochondria in dendrites

Akane Hatsuda[1,2], Junko Kurisu[2], Kazuto Fujishima[2,†], Ayano Kawaguchi[3,4], Nobuhiko Ohno[5,6], Mineko Kengaku[1,2]

1. Graduate School of Biostudies, Kyoto University, Japan

2. Institute for Integrated Cell-Material Sciences (KUIAS-iCeMS), Kyoto University, Japan

3. Department of Anatomy and Cell Biology, Nagoya University Graduate School of Medicine, Japan

4. Department of Human Morphology, Okayama University Graduate School of Medicine, Density and Pharmaceutical Sciences, Japan.

5. Department of Anatomy, Division of Histology and Cell Biology, Jichi Medical University, Japan

6. Division of Ultrastructural Research, National Institute for Physiological Sciences, Japan

Corresponding authors. Email: kengaku@icems.kyoto-u.ac.jp

†Present address: Department of Anatomy and Cell Biology, Division of Life Sciences, Osaka Medical and Pharmaceutical University, Japan

Abstract

AMP-activated protein kinase (AMPK) governs multiple pathways regulating mitochondrial dynamics and homeostasis. AMPK overactivation is a hallmark of some neurodegenerative diseases, leading to excessive mitochondrial fission and mitophagy and subsequent degeneration of neuronal processes. Here we demonstrate that AMPK is involved in activity-dependent dendritic outgrowth of developing hippocampal neurons. AMPK deficiency phenocopies the inhibition of neuronal activity, inducing dendritic hypotrophy with abnormally elongated mitochondria. In growing dendrites, AMPK is activated by neuronal activity and dynamically oscillates in synchrony with calcium spikes, and this AMPK oscillation is inhibited by CaMKK2 knockdown. AMPK activation leads to phosphorylation of MFF and ULK1, which initiate mitochondrial fission and mitophagy, respectively. Dendritic mitochondria in AMPK-depleted neurons exhibit impaired fission and mitophagy and display multiple signs of dysfunction. Thus, AMPK activity is finely tuned by the calcium-CaMKK2 pathway and regulates mitochondrial homeostasis by facilitating removal of damaged components in rapidly growing neurons during normal brain development.

Introduction

Dendrite morphogenesis is a pivotal step for the establishment of neural circuit connectivity in the developing brain. Rapid extension and branching of dendrites in developing neurons is accompanied by extensive cytoskeletal and membrane remodeling and intracellular transport that increase the energetic burden particularly in growing dendritic branches. We have previously shown that mitochondria are actively transported to growing dendritic arbors to maintain local ATP levels necessary for the regulation of actin dynamics[1]. In addition to producing the majority of cellular ATP, mitochondria also participate in various processes indispensable for the development and function of dendrites, including Ca^{2+} homeostasis, lipid flux, and reactive oxygen species (ROS) generation[2,3]. The abnormal morphology of dendritic mitochondria is one of the pathogenic hallmarks of Alzheimer's disease (AD), supporting the notion that constant delivery of functional mitochondria throughout a large volume of dendrites is critical for maintenance as well as differentiation of neurons[4,5].

Mitochondrial homeostasis in postmitotic neurons is highly dependent on dynamic fission and fusion events. Mitochondrial fission is particularly important for the biogenesis of new mitochondria and for the removal of damaged or aged mitochondria through mitophagy[6]. Mitochondrial fission is triggered by the spiral assembly of an evolutionarily conserved GTPase dynamin-related protein 1 (Drp1) around the constriction site on the mitochondrial surface. During fission, Drp1 is recruited to the outer mitochondrial membrane by one of its receptors, such as Mitochondrial Fission Factor (MFF)[7–9]. Several groups, including us, have shown that the inhibition of mitochondrial fission leads to abnormal elongation and dysfunction of mitochondria, resulting in aberrant dendrite formation in vivo and in vitro[10–12]. On the other hand, excessive mitochondrial fission also leads to reduced respiratory activity and abnormal distribution of mitochondria, which impairs normal development of dendrites[13,14]. However, it remains unclear how the proper balance of mitochondrial fission is adaptively regulated in developing neurons whose shape, volume and activity states are dynamically changing.

The balance between fission and fusion should be finely tuned in response to changes in the cellular and subcellular metabolic state. Recent studies have highlighted the role of AMP-activated protein kinase (AMPK) as an energy sensor that sustains the metabolic state by regulating mitochondrial dynamics and functions[15]. AMPK regulates several aspects of mitochondrial homeostasis by direct phosphorylation of key factors, including MFF, peroxisome proliferator-activated receptor-coactivator 1 (PGC1α), Armadillo repeat-containing protein 10 (ARMC10), Unc-51 like autophagy activating kinases 1 and 2 (ULK1/2) and Mitochondrial

fission regulator 1-like protein (MTFR1L)[16–22]. AMPK is activated by direct allosteric binding of

AMP, facilitating phosphorylation at Thr172 by an upstream kinase LKB1 in response to cellular

ATP shortage[23]. Additionally, AMPK can be directly phosphorylated by the calcium (Ca^{2+})-

sensitive kinase CaMKK2 (also known as CaMKKβ) in response to an elevated intracellular Ca^{2+}

concentration[24,25]. In the central nervous system, overactivation of the CaMKK2-AMPK pathway

by excessive Ca^{2+} signals has been implicated in mitochondrial fragmentation and degradation

during the pathogenesis of Alzheimer's disease[26,27].

It is widely accepted that Ca^{2+} signals evoked by neuronal activity play important roles

in dendrite development. Ca^{2+} influx via voltage-gated calcium channels (VGCCs) and NMDA

receptors triggers activation of Ca^{2+}/calmodulin-dependent kinases (CaMKs) that positively and

negatively regulate dendrite growth through transcriptional control and local remodeling of the

cytoskeleton[28–30]. In mature neurons, activity-induced Ca^{2+} signals have been shown to induce

mitochondrial fission in dendrites[10,31]. However, the causal link between Ca^{2+} signals evoked by

neuronal activity and mitochondrial fission is less understood in the growing dendrites of

developing neurons. In this study, we investigated the impact of neuronal activity on the dynamics

of dendritic mitochondria in differentiating hippocampal neurons. We demonstrate that neuronal

activity finely tunes the CaMKK2-AMPK pathway in growing dendrites. Normal AMPK activity

induces mitochondrial fission and mitophagy via phosphorylation of MFF and ULK1, which are

critical to remove dysfunctional mitochondrial components and maintain functional mitochondria

in growing dendrites. Thus, there is indeed a causal link between activity-dependent dendritic

arbor development and AMPK-dependent mitochondrial quality control.

Results

Neuronal activity enhances dendrite formation and mitochondrial fission in developing

hippocampal neurons

We previously established long-term live-imaging of dissociated hippocampal neurons and

observed the gradual formation of dendritic arbors, with dynamic extension and retraction, until

the second week of culture[32]. To visualize neuronal activity in developing dendrites, we

transfected GCaMP6s in cultured hippocampal neurons and performed live imaging at 5 days in

vitro (DIV5). We observed intermittent Ca^{2+} transients in dendrites and cell soma, which were

abolished by the combined treatment with a blocker of voltage-gated sodium channels,

tetrodotoxin, and an NMDA receptor antagonist, D-(-)-2-Amino-5-phosphonopentanoic acid

(TTX+APV treatment) (Fig. 1A, Supplementary Video 1). There was no apparent difference in

the dynamic extension and retraction of dendrites in drug-treated and untreated neurons, but

growth retardation gradually became evident, with a significant decrease in total dendritic length

and the number of branches at 48 hr after TTX+APV treatment (Fig. 1B-D, Supplementary Video

2).

108   To examine if neuronal activity affects mitochondrial dynamics in growing dendrites,

we transfected Mito-DsRed into neurons that were then untreated or treated with TTX+APV.

Under both conditions, mitochondria were distributed throughout the dendritic arbors, and a

significant fraction of them moved along the dendritic shaft as previously reported by us and

others[1,33–35]. Inhibition of neuronal activity did not affect mitochondrial motility in dendrites:

neither the fraction of motile mitochondria nor their average speed was altered by TTX+APV

treatment (Extended Data Fig. 1A and B). There were no apparent changes in the density and

distribution of mitochondria in dendrites in the drug-treated neurons (Extended Data Fig. 1C).

However, we found a significant increase in the length of dendritic mitochondria in the drug-

treated neurons [Median (IQR)= 2.0 (1.2- 2.9) µm in control vs 2.5 (1.6-3.8) µm in TTX+APV

neurons; Fig. 1E, G, H, I]. In the axon, on the other hand, the length of mitochondria was much

shorter, as previously reported[36], and was unaffected by TTX+APV treatment (Fig. 1F). As

mitochondrial morphology is affected by the balance between fission and fusion, we analyzed the

frequency of fission and fusion in dendrites by high resolution time-lapse imaging (Extended Data

Fig. 1D and Supplementary Video 3). We found that inhibition of neuronal activity significantly

suppressed mitochondrial fission but had little or no effect on fusion (Fig. 1J). Conversely,

glutamate treatment led to a marked increase in smaller mitochondria in the cell soma and

dendrites, suggesting that enhanced neuronal activity facilitated mitochondrial fission (Extended

Data Fig. 1E). Taken together, these data suggest that neuronal activity is involved in the

regulation of the mitochondrial fission-fusion balance during dendritic outgrowth in developing

neurons.

AMPK regulates dendritic growth and mitochondrial fission

It has been shown that AMPK mediates mitochondrial fission in response to energy stress in

neurons and non-neuronal cells[17]. We therefore examined if AMPK is involved in activity-

dependent mitochondrial fission in developing hippocampal neurons. AMPK is a heterotrimeric

serine/threonine kinase composed of one catalytic (α1/α2), one regulatory (β1/β2) and one

AMP/ATP binding (γ1/γ2/γ3) subunit. Both AMPKα1 and AMPKα2 were expressed in cultured

hippocampal neurons, as assessed by western blotting using isoform-specific antibodies (Fig. 2A).

Specifically, AMPKα1 expression was high in earlier stages and declined by DIV10, whereas

AMPKα2 expression started at a level comparable to AMPKα1, and then gradually increased,

peaking at DIV10. The abundance of active AMPKα phosphorylated at Thr172 (detected by an

antibody that recognizes both phosphorylated isoforms, p-AMPKα) increased during dendritic

outgrowth in parallel with AMPKα2 expression (Fig. 2A).

To investigate the role of AMPK in mitochondrial dynamics and dendritic formation,

we knocked down AMPK subunits in cultured neurons by expressing short hairpin RNAs

(shRNA) against AMPKα1 and AMPKα2 (Extended Data Fig. 2A and B). At DIV3, we

transfected dissociated hippocampal neurons with AMPKα1 shRNA or AMPKα2 shRNA

plasmids together with a volume marker GFP construct, and observed dendritic morphology at

DIV5. AMPKα2 shRNA significantly reduced the size and complexity of dendritic branches,

reminiscent of the activity-deprived neurons (Fig. 2B-D and Extended Data Fig. 2C-E compared

to Fig. 1). Concomitant exogenous expression of an AMPKα2 mutant cDNA (AMPKα2 Rescue)

that was resistant to AMPKα2 shRNA largely prevented the dendritic defects, discounting any

major off-target effects of the shRNA (Fig. 2B-D and Extended Data Fig. 2C). In contrast,

AMPKα1 shRNA produced much weaker effects on dendritic morphology. Further, simultaneous

knockdown of AMPKα1 and α2 was not phenotypically different from AMPKα2 knockdown,

suggesting that AMPKα2 was the predominant isoform regulating dendritic morphogenesis in

our culture (Extended Data Fig. 2C-E). We further confirmed that repression of AMPKα2 using

CRISPR interference (CRISPRi) significantly delayed dendritic outgrowth at DIV5 (Extended

Data Fig. 2F-H). We thus focused on AMPKα2 in subsequent experiments.

To confirm the requirement of AMPKα2 for dendritic formation in vivo, we introduced

the AMPKα2 shRNA#1 together with GFP into CA1 pyramidal neurons at embryonic day (E)

14.5 by in utero electroporation. Compared to control neurons, AMPKα2-depleted neurons

showed significantly reduced length and complexity of the apical dendrite at P5, supporting the

notion that AMPKα2 functions in dendritic outgrowth of hippocampal pyramidal neurons (Fig.

2E-H). In contrast, the dendritic hypotrophy was less evident at P10, suggesting a compensatory

mechanism allows the recovery of dendritic growth retardation in the context of AMPKα2

deficiency in vivo.

Returning to cultured cells, we next observed mitochondria in AMPKα2-depleted

neurons by cotransfecting mito-DsRed and GFP. The AMPKα2 knockdown significantly

increased mitochondrial length in dendrites [Median (IQR)= 2.1 (1.3-3.1) μm in control; 2.6 (1.6-

4.1) μm in AMPKα2 knockdown; 2.0 (1.3-3.1) μm in rescue; Fig. 3A, C-E], but not in the axon

at DIV5 (Fig. 3B). There was no apparent change in the density and distribution of mitochondria

in the hypotrophic dendrites in AMPKα2-depleted neurons (Extended Data Fig. 3A). AMPKα2

CRISPRi also induced abnormal elongation of mitochondria in dendrites (Extended Data Fig. 3B-

E). These results prompted us to use time-lapse imaging to analyze the fission/fusion frequency

of dendritic mitochondria in the neurons transfected with AMPKα2 shRNA. As in the case for

neuronal activity inhibition, AMPKα2 knockdown specifically lowered the frequency of fission

but had little effect on fusion in dendrites (Fig. 3F).

We further documented the importance of AMPK activity in mitochondrial dynamics

by pharmacological activation and inhibition. We found that dendritic mitochondria were

significantly shortened after a 3-5 h treatment with an AMPK activator, AICAR (1 mM) (Fig. 3G-

J). In contrast, treatment with an AMPK inhibitor, Compound C (20 μM), elongated dendritic

mitochondria (Fig. 3K-N), underscoring the involvement of AMPK in the regulation of

mitochondrial fission in hippocampal neurons. Collectively, these data suggest that AMPKα2 has

critical roles in mitochondrial fission and dendritic formation in hippocampal neurons, and that

the two phenomena are mechanistically linked.

Neuronal activity induces dynamic oscillation of AMPK activity via Ca/CaMKK2

The striking phenotypic similarities between the modulation of neuronal activity and AMPK

activity prompted us to examine if AMPK mediates the activity-dependent control of

mitochondrial dynamics in developing dendrites. In order to analyze the spatiotemporal dynamics

of AMPK activity in developing hippocampal neurons, we transfected them with AMPKAR-EV,

a highly sensitive fluorescence resonance energy transfer (FRET)-based biosensor for AMPK[37].

We observed high basal FRET signals in the axon, which presumably reflected the activity of

AMPK-related kinases such as SAD-A/B[38–40]. Indeed, AMPKα2 shRNA specifically suppressed

FRET signals in the somatodendritic compartment but not in the axon (Extended Data Fig. 4A),

supporting the notion that the FRET signals in dendrites, but not in axons, reflect AMPK activity.

Consistent with previous reports, we observed a rapid increase in AMPK activity in

the cell body and dendrites upon glutamate treatment[41,42]; moreover, the AMPK activity was

abrogated by AMPKα2 knockdown (Extended Data Fig. 4B). Interestingly, we found an

unambiguous periodic fluctuation of AMPK activity with a frequency of 0 to 4 peaks in 3 minutes,

in the somatodendritic compartment of untreated neurons (Fig. 4A and C). The basal fluctuation

was suppressed by cotransfection of AMPKα2 shRNA or inhibition of neuronal activity by

TTX+APV treatment, suggesting that neuronal activity caused dynamic changes in AMPK

activity (Fig. 4A-D). To further prove the causal relationship of neuronal activity and AMPK

activation, we performed simultaneous imaging of Ca^{2+} and AMPK activity using a red

fluorescent Ca^{2+} indicator, jRGECO1a, together with AMPKAR-EV. We observed periodic Ca^{2+}

spikes which highly synchronized with AMPK activity fluctuation in dendrites (Fig. 4E, F and

Supplementary Video 4). The temporal cross-correlation analysis revealed the highest correlation

at 5 sec after Ca^{2+} influx, indicating that Ca^{2+} influx precedes AMPK activation (Fig. 4G).

Inhibition of neuronal activity by TTX+APV treatment suppressed both Ca^{2+} spikes and AMPK

fluctuation (Extended Data Fig. 4C). In contrast, AMPK knockdown suppressed only the AMPK

fluctuation, and did not alter the Ca^{2+} transient (Extended Data Fig. 4D), indicating that AMPK

fluctuation was induced by neuronal activity.

Two upstream kinases, CaMKK2 and LKB1, regulate AMPK activity by

phosphorylation[23–25]. As CaMKK2 has been implicated in activity-dependent formation of

dendrites[43,44], we examined whether CaMKK2 is required for AMPK fluctuation in response to

neuronal activity. Cotransfection of CaMKK2 shRNA or treatment with CaMKK2 inhibitor

(STO609, 10 μM) suppressed the periodic activation of AMPK (Fig. 4H-J). On the other hand,

the AMPK fluctuation was not affected by treatment with an LKB1 inhibitor (Pim1, 1μM)

Extended Data Fig. 4E).

Taken together, these data suggest that neuronal activity dynamically regulates AMPK

activity via Ca^{2+}/CaMKK2 in the somatodendritic region of developing hippocampal neurons.

AMPKα2 knockdown prevents asymmetric mitochondrial fission and mitophagy in

developing dendrites

Mitochondrial fission is involved in biogenesis and mitophagic degradation of mitochondria. It

has recently been reported that the downstream pathways could be distinguished by the position

of fission, either at the midzone or the periphery[45]. In normal developing hippocampal neurons,

we observed both midzone and peripheral fissions in growing dendrites, with the frequency of the

midzone fission, which was implicated in mitochondrial biogenesis in the previous study[45], being

approximately 2.1-fold higher than peripheral fission (Fig. 5A and B). TTX+APV treatment

hampered both midzone and peripheral fissions, but had a stronger inhibitory effect on peripheral

fission (Fig. 5B). AMPKα2 knockdown displayed strikingly similar effects, with stronger

suppression of the peripheral fission (Fig. 5C).

As the peripheral fission has been implicated in the mitophagy pathway, we asked

whether AMPK-induced mitochondrial fission is required for mitophagic degradation of dendritic

mitochondria. To this end, we used an antibody against the autophagy marker p62 to immunostain

neurons transfected with Mito-DsRed with or without AMPKα2 shRNA. The p62 signals formed

puncta, some of which were colocalized with mitochondria in the soma and dendrites in control

neurons. By contrast, the number of p62 puncta and those colocalized with mitochondria were

significantly decreased in AMPKα2 knockdown cells (Fig. 5D-G). These data suggest that

AMPK-mediated mitochondrial fission is important for the mitophagy initiation during dendrite

development.

We also examined the effect of AMPK activation on mitochondrial biogenesis. AMPK

has been shown to phosphorylate and activate PGC1α, the master regulator of mitochondrial

biogenesis. However, we detected no obvious difference in the phosphorylation level of PGC1α

by Phos-tag SDS-PAGE, nor the expression of PGC1α, in response to TTX+APV, Glutamate,

Compound C or AICAR (Extended Data Fig. 5A).

AMPK phosphorylates mitochondrial fission factor and ULK1 in response to neuronal

activity.

AMPK phosphorylates various downstream substrates involved in many aspects of mitochondrial

homeostasis. AMPK regulates mitochondrial fission by direct phosphorylation of MFF, which

promotes its association with Drp1 at the mitochondrial outer membrane[16,17]. AMPK is also

known to directly phosphorylate the autophagy initiating kinase ULK1[20]. We thus asked whether

AMPK phosphorylates these substrates in response to neuronal activity.

Consistent with our observation that AMPK activity was dynamically regulated in

synchrony with neuronal activity, AMPK phosphorylation rapidly increased within 5 min of

glutamate treatment (Fig. 6A). In contrast, TTX+APV treatment reduced the basal level of AMPK

phosphorylation. Concomitantly, MFF phosphorylation at Ser146 was increased by glutamate

treatment, while it was repressed by TTX+APV treatment. Likewise, the level of ULK1

phosphorylation at Ser555 paralleled that of AMPK phosphorylation, which was altered by

manipulation of the activity states in cultured neurons. We also confirmed that the

phosphorylation levels of MFF and ULK1 were upregulated by AICAR treatment, while they

were downregulated by Compound C treatment. Taken together, these results suggest that AMPK

activation by neuronal activity induces mitochondrial fission and mitophagy by direct

phosphorylation of MFF and ULK1.

AMPK knockdown compromises normal mitochondrial activity in developing dendrites

We next asked if AMPK-mediated mitochondrial fission and mitophagy is required for

mitochondrial quality control. Using the fluorescent dye tetramethylrhodamine methyl ester

(TMRM, 20 nM), we asked whether the mitochondrial membrane potential ($\Delta\Psi$m) was affected

in the elongated dendritic mitochondria in neurons that were defective in AMPK-mediated

mitochondrial fission. At first, we were unable to detect a significant difference in the average

intensity of basal TMRM fluorescence in individual mitochondria of control and AMPKα2

knockdown cells (Extended Data Fig. 5B and C). We then analyzed dynamic changes in $\Delta\Psi$m by

time-lapse imaging. We observed intermittent flickers of TMRM fluorescence in some dendritic

mitochondria in control neurons (Fig. 6B and Supplementary Video 5). TMRM flickering is

caused by the opening of the mitochondrial permeability transition pore (mPTP), an essential

process for maintaining mitochondrial homeostasis by preventing an overload of Ca^{2+} and

reactive oxygen species (ROS) in the mitochondrial matrix induced by respiration activity[46,47]. In

AMPKα2 knockdown neurons, the frequency of TMRM flickering was significantly reduced,

suggestive of lower respiratory activity of mitochondria in these cells (Fig. 6C).

We then analyzed mitochondrial ROS levels (mitoROS) using the fluorescent dye

MitoSOX (500 nM). MitoROS is a byproduct of electron transport chain activity, which is

increased by high respiration and triggers mPTP opening[48]. MitoSOX signals formed puncta in

the mitochondrial matrix, which were significantly decreased in AMPKα2 knockdown cells (Fig.

6D and E).

These results strongly suggest that AMPK-mediated mitochondrial fission and

mitophagy is required for the maintenance of functional mitochondria in growing dendrites of

hippocampal neurons.

Discussion

To meet the rapidly growing energy demands incurred by the extension of dendritic arbors, mitochondria are actively produced and transported throughout the expanding dendritic compartments. Here we demonstrate that neuronal activity regulates mitochondrial homeostasis through the activation of the Ca^{2+}-CaMKK2-AMPK axis to support dendritic outgrowth in developing neurons. AMPK deficiency prevented mitochondrial fission and mitophagy, leading to mitochondrial dysfunction with low respiratory activity and consequent growth retardation of dendrites. Our observations suggest that AMPK-dependent regulation of local mitochondrial homeostasis is an important mechanism of activity-induced dendritic growth besides the well-known transcription-dependent mechanism.

Mitochondrial fission is prerequisite for the biogenesis of mitochondria and for the elimination of damaged components through the mitophagy pathway[49,50]. Recent studies have demonstrated that symmetric fission generating two equal daughter mitochondria leads to doubling of mitochondrial biomass, while asymmetric fission generates a smaller mitochondrion that is subjected to the mitophagy pathway[45]. We observed both symmetric and asymmetric fissions in growing dendrites, but the asymmetric fission was more strongly affected by the inhibition of neuronal activity or AMPK. We showed that neuronal activity induces phosphorylation/activation of two downstream effectors of AMPK, MFF and ULK1, which are the key regulators of mitochondrial fission and mitophagy, respectively. Mitochondria with defective fission and mitophagy in AMPK-depleted neurons exhibit multiple signatures of mitochondrial dysfunction, such as reduced $\Delta\Psi$m flickering and ROS production, suggesting that AMPK-mediated mitochondrial fission and mitophagy is required for the maintenance of healthy mitochondria in growing dendrites.

On the other hand, we did not obtain evidence supporting the regulation of mitochondria biogenesis by activity-induced AMPK signaling, such as a change in PGC1α phosphorylation and expression. Nonetheless, the total biomass of dendritic mitochondria should decrease as a consequence of AMPK knockdown or neuronal activity inhibition, since the mitochondrial density remains the same but the dendritic volume is substantially reduced. We will need more sensitive assays to draw conclusions regarding the regulation of mitochondrial biogenesis by activity-induced AMPK signaling.

AMPK and AMPK-related kinases are expressed in both dendrites and axons of cortical neurons, but their activities seem differentially regulated in distinct subcellular compartments. Previous studies have shown that the activities of SAD-A/B and NUAK1 and their upstream

kinase LKB1 are strongly biased in the axon and regulate axon specification and branching[39,40].

The SAD kinase has been shown to regulate the fission/fusion balance of mitochondria through

phosphorylation of tau in the axon[51,52]. In contrast, we demonstrate that AMPK overactivation

and suppression alters mitochondrial fission, and those changes are seen only in dendrites but not

in the axon (Fig. 3). The activation of AMPK in dendrites and AMPK-related kinases in the axon

appears to be regulated by distinct molecular mechanisms: SAD-A/B and NUAK1 have been

shown to be constitutively activated in the axon by LKB1. In contrast, AMPK phosphorylation

dynamically oscillates in dendrites, depending on Ca^{2+} influx and CaMKK2 activity, while it is

independent of LKB1. AMPK activity was difficult to detect in the axon due to the high basal

FRET signals, presumably reflecting the high activity of SAD-A/B and/or NUAK1 in the axon.

Nonetheless, neither neural activity nor AMPK overactivation enhanced fission of axonal

mitochondria, suggesting that mitochondrial dynamics and transport are differentially regulated

in the axon and dendrites by distinct AMPK-related kinases.

One interesting finding in the present study is that AMPK activity dynamically oscillates in

synchrony with Ca^{2+} influx triggered by neuronal activity. Ca^{2+}-dependent AMPK

phosphorylation is rapidly downregulated in several tens of seconds, implying a regulatory loop

involving a negative regulator of AMPK activity which counteracts CaMKK2-mediated

phosphorylation. The molecular pathway of AMPK dephosphorylation is less understood, but a

Ca^{2+}-dependent phosphatase such as PP2A is a likely candidate, as PP2A has been implicated in

AMPK dephosphorylation and inactivation in non-neuronal cells[53]. The dynamic oscillatory

activation of AMPK would enable fine spatiotemporal tuning of mitochondrial dynamics in

response to the local activity states in dendritic compartments. The controlled fission and

mitophagy is critical for the maintenance of mitochondrial quality in developing dendrites, as the

inhibition of AMPK-dependent mitophagy caused aberrant elongation and dysfunction of

mitochondria.

Intriguingly, the CaMKK2-AMPK pathway regulating MFF and ULK2 activities has been

implicated in the excessive mitochondrial fission and mitophagy in degenerating dendrites in

Alzheimer's disease models[26,27]. It is suggested that disruption of Ca^{2+} homeostasis overactivates

the CaMKK2-AMPK pathway in these neurons. Thus, Ca^{2+}-sensitive activation of AMPK could

be a double-edged sword, promoting dendritic growth by regulating mitochondria turnover during

normal development, while contributing to neurodegeneration by excessive mitochondrial

degradation under pathological conditions.

It has been shown that genetic deletions of AMPK subunits exhibit little or no cell-

autonomous defects in neuronal development in mice[54–57]. Conditional deletion of AMPKα1/α2

(AMPKα1−/−; AMPKα2F/F; Emx1-Cre) exhibits no overt phenotypes in the polarization and

survival of cortical neurons, except for the thinning of the cortical layer at P3[55,58]. In contrast,

acute treatment with AICAR or metformin induces rapid AMPK phosphorylation and disrupts

axon formation in cultured cortical neurons[59]. Thus, AMPK is thought to be dispensable for

neuronal differentiation during normal development, but exerts its effects when overactivated

under stress conditions. In the present study, we show that defects in mitochondrial dynamics and

dendrite growth by acute AMPK inhibition are evident only in differentiating neurons in the first

postnatal week, and that they recovered in later stages in vivo. It is therefore possible that the

transient morphological changes might have been overlooked in previous studies. Mild and

transient phenotypes by chronic inhibition of AMPK might be due to the robust multi-step

regulation of mitochondrial dynamics. In mature neurons, long-term potentiation (LTP) of

synaptic activity causes mitochondrial fission via CaMKII and Drp1 phosphorylation[31]. The

intracellular Ca^{2+} rise also induces a distinct type of mitochondrial fission initiated from the inner

mitochondrial membrane[60]. AMPK might predominantly mediate spontaneous activity before

synaptic maturation, while other mechanisms may take over and control mitochondrial fission in

more mature neurons. Although no overt histological defects were found in mature neurons of

AMPK-deficient mice, whether these neurons form and operate normal neural circuits has not

been assessed. Whether AMPK has a distinct function in dendrites of mature neurons is one of

many important questions that entail further study.

The present study highlights a new role of AMPK in mitochondrial homeostasis in normal

development of neurons in addition to its major role in safeguarding neurons from energetic stress.

Dynamic activation and inactivation of AMPK by neuronal activity are clearly important for the

fine-tuning of mitochondrial homeostasis. Thus, future work should focus on the precise

dynamics and mechanism of AMPK activation, as well as inactivation, for a new perspective on

the physiology and pathology of AMPK signals.

Methods

Mice

All animals were treated in accordance with the guidelines of the Animal Experiment Committee of Kyoto University. ICR mice were kept in a 12 hr dark/light cycle at $23 \pm 3°C$/ 50% humidity, with standard food and water provided ad libitum, in group housing of up to three animals per cage.

Reagents

Reagents used in this study were as follows: D, L-APV (APV, 200 µM, Sigma Aldrich, A5282), Tetrodotoxin (TTX, 0.5 µM, Abcam, ab120054), AICAR (1-2 mM, AdipoGen. Inc, AG-CR1-0061), Compound C (CC, Dorsomorphin, 20 µM, Nacalai, 18768-04), L-Glutamic acid (Glutamate, 100 µM, Sigma-Aldrich, G1251), STO-609 (10 µM, MedChem Express, HY-19805), Pim1/AKK1-IN-1 (1 µM, MedChem Express, HY-10371), TMRM (20 nM, Fujifilm, 203-18041), Mito-SOX (500 nM, Invitrogen, M36008).

Plasmids

pCAG-EGFP, pAAV-CAG-Mito-DsRed and pAAV-CAG-Mito-EGFP were constructed as previously described[1,61]. For RNAi experiments, the target sequences for scramble control (5'-GACATTTCATCCGTTTAGTTA-3'), AMPKα1 sh#1 (5'-GGCACACCCTGGATGAATTAA-3'), AMPKα1 sh#2 (5'-GTTGTAAACCCCTATTATTTG-3'), AMPKα2 sh#1 (5'-GGTAGACAGTCGGAGCTATCT-3'), AMPKα2 sh#2 (5'-GACAATCGGAGAATAATGAAC-3'), and CaMKK2 (5'-CCCTTTCATGGATGAACGAAT-3') were cloned into pBAsi-hH1 vector (Takara, 3220). pCAG-AMPKα1(WT) and pCAG-AMPKα2 (WT) plasmids were created by PCR amplification of cDNAs for AMPKα1 (NCBI Reference Sequence: NM_001013367.3) and AMPKα2 (NCBI Reference Sequence: NM_178143.2) from a mouse brain cDNA library and inserted into the pCAGGS vector. Resistant mutants of AMPKα1 and AMPKα2 that contained three silent mutations within the respective shRNA target sequences were generated using the Primestar mutagenesis protocol. pPBbsr2-4031NES (AMPKAR-EV) was a gift from Michiyuki Matsuda. pCAG-jRGECO1a plasmid was generated by subcloning the jRGECO1a sequence from pGP-CMV-NES-jRGECO1a (a gift from Masayuki Sakamoto) into the pCAGGS vector. The pX458-CAG-dCas9KRAB-2A-GFP plasmid was generated by subcloning the CAG promoter from pCAGGS and the dCas9KRAB sequence from CRISPR-dCas9KRAB (a gift from Kuniya

Abe) into the pX458 vector (Addgene plasmid #48138). For CRISPR interference experiments,

the gRNA sequence for AMPKα2 (5'-TGCCGAAGGTGCCGACGCCCAGG-3') was inserted

into pX458-CAG-dCas9KRAB-2A-GFP.

Cell cultures and transfection

Primary cultures of hippocampal neurons were prepared as previously described with a few

modifications[32]. In brief, postnatal day 0 (P0) mouse hippocampi were dissected, dissociated

using Neuron Dissociation Solutions (Fujifilm, 291-78001), and plated on poly-D-lysine (Sigma-

Aldrich, P6407) coated coverslips at a density of $3.0 \sim 3.5 \times 10^5$ cells/cm^2 in MEM (Gibco,

11095080) supplemented with 10% horse serum (Gibco, 26050070), 0.6% D-glucose, 1 mM

sodium pyruvate (Sigma-Aldrich, S8636), and 1% penicillin-streptomycin. Three hours after

plating, media was replaced with Neurobasal medium (Gibco, 21103049) supplemented with 2%

(v/v) B-27 (Gibco, 17504044) and 0.25% (v/v) GlutaMax (Gibco, 35050061). All neurons were

maintained at 37°C in 5% CO_2. Five μM AraC (Sigma-Aldrich, C1768) was added at DIV3.

Neurons were transfected at DIV3 using Lipofectamine 2000 (Thermo Fisher Scientific,

11668019) according to the manufacturer's instructions. The CRISPRi plasmid was transfected

at DIV2.

HEK293T cells (Riken BRC Cell Bank, RCB2202) were cultured in DMEM (Gibco, 11965-092)

supplemented with 10% Fetal Bovine Serum (FBS, Sigma-Aldrich, F7524), 100 units/mL

penicillin and 100 ug/mL streptomycin (Gibco, 15140-122) at 37°C in 5% CO_2. Cells were

passaged every 2-3 days and maintained in plastic cell culture-treated dishes. shRNA constructs

were transfected using Lipofectamine 2000.

Western blotting

Cells were harvested in RIPA lysis buffer (50 mM Tris HCl (pH 7.4), 150 mM sodium chloride,

0.25% (w/v) Sodium Deoxycholate, 1% (v/v) NonidetP-40, 1 mM EDTA, and Protease and

phosphatase inhibitor cocktail (Thermo Fisher Scientific, 78442)). Lysates were subjected to

SDS-PAGE and immunoblotting.

Primary antibodies used were the following: mouse anti-GFP (1:5000, Thermo Fisher

Scientific, A11120); rabbit anti-AMPKα1 (1:3000, Abcam, ab32047); rabbit anti-AMPKα2

(1:3000, Abcam, ab3760); rabbit anti-AMPKα (1:1000, Cell Signaling Technology, 2532); rabbit

anti-phospho-AMPKα (Thr172) (1:1000, Cell Signaling Technology, 2535); rabbit anti-MFF

(1:3000, Proteintech, 17090-1-AP); rabbit anti-phospho-MFF (Ser172/146) (1:1000, Affinity

Biosciences, AF2365); rabbit anti-ULK1 (1:2000, Cell Signaling Technology, 8054); rabbit anti-

phospho-ULK1 (Ser555) (1:1000, Cell Signaling Technology, 5869); rabbit anti-PGC1α (1:5000,

Abcam, ab191838) and HRP-conjugated mouse anti-beta actin (1:10000, Santa Cruz, sc-47778

HRP). The secondary antibodies used were the following: HRP-conjugated goat anti-mouse IgG

(1:10000, Bio-Rad, 170-6516); HRP-conjugated goat anti-rabbit IgG (1:10000, Bio-Rad, 170-

6515).

In utero electroporation

Pregnant mice on day 14.5 of gestation were deeply anesthetized via the intra-abdominal injection

of a mixture of medetomidine, midazolam and butorphanol. Plasmid DNA diluted with saline

containing 0.01% Fast Green was injected into the lateral ventricle of mouse embryos with a glass

needle. Four current pulses (amplitude, 40 V; duration, 50 ms; intervals, 950 ms) were delivered

with a forceps-shaped electrode (CUY650P3, NepaGene) connected to an electroporator (CUY21,

NepaGene). Concentrations of the plasmids used are as follows: pCAG-EGFP (0.2 µg/µl);

shRNA-scramble control (3-6 µg/µl); shRNA- AMPKα2 (3-6 µg/µl); CRISPRi gRNA-empty (5

µg/µl); and CRISPRi gRNA- AMPKα2 (5 µg/µl).

Immunofluorescence and image acquisition

For immunocytochemistry, cultured cells were fixed with 4% PFA in PBS and permeabilized with

PBS containing 0.25% Triton X-100 (PBS-T). Cells were then blocked with blocking solution

(PBS with 2% BSA) and incubated with rabbit anti-AMPKα2 primary antibody (1:200,

Proteintech, 18167-1-AP) at 4°C overnight. After washing with PBS, cells were incubated with

goat anti-Rabbit Alexa Fluor 647 (1:400, Invitrogen, A21244) secondary antibody at 4°C

overnight.

For immunohistochemistry, the mice at P5 or P10 were perfused with PBS followed

by 4% paraformaldehyde (PFA) in phosphate buffer. Their brains were removed and postfixed for

2-3 hrs at 4°C. After washing with PBS, the brains were embedded in 3.5% low-melting agarose

in PBS. The brains were then sectioned into 200 µm thick sagittal slices with a vibratome (NLS-

AT, Dosaka EM). The sections were permeabilized in PBS with 0.25 % Triton (PBS-T) for 15

min and blocked with 2% skim milk in PBS-T for 30 min. The sections were incubated with a

chick anti-GFP (1:2000, Invitrogen, A10262) primary antibody in blocking solution at 4°C

overnight, followed by incubation with a goat anti-Chicken Alexa Fluor 488 (1:400, Invitrogen,

A11039) secondary antibodies at 4°C overnight.

Images of the fixed samples were acquired with a laser scanning confocal microscope

(FV1000, Olympus) equipped with UPlanSApo 20× (N.A. 0.75), UPLSAPO 40× (NA 0.95)

objectives, or a spinning disk confocal microscope (Dragonfly, Andor) equipped with Nikon

PLAN APO 40× (Silicone oil, NA 1.25), Apo TIRF 100× (oil-immersion, NA 1.49) objectives.

For morphometric analysis, dendrites were traced using the ImageJ plugin Simple

Neurite Tracer (SNT). Total dendrite length and the number of branches were calculated using

MATLAB software. For quantification of dendritic mitochondrial densities (ratio of

mitochondrial area to dendrite area), acquired images were binarized and the areas filled with

Mito-DsRed signals and volume marker GFP signals were quantified using ImageJ.

For p62 analysis, hippocampal neurons transfected with GFP and mito-DsRed were

fixed at DIV5 and immunostained with guinea pig anti-p62 primary antibody (1:100, Progen,

GP62-C,) and goat anti-Guinea Pig Alexa Fluor 633 secondary antibody (1:400, Invitrogen,

A21105). The number of p62 puncta in dendrites was counted using the ImageJ plugin Find

Maxima. For colocalization analysis of p62 puncta and mitochondria, Mander's overlap

coefficient was calculated from anti-p62 signals and mito-DsRed signals in dendrites using the

ImageJ plugin JaCoP. Surface rendering was done with Imaris software.

Time-lapse imaging and image analysis

For time-lapse imaging of mitochondrial dynamics, cultured hippocampal neurons were

transfected on DIV3 with pCAG-EGFP and pAAV-CAG-mitoDsRed plasmids. Time-lapse

images were acquired every 3 s for 20 min at DIV5 using a spinning-disc confocal microscope

(CV1000, Yokogawa) with a UPLSApo 100× objective (Oil immersion, N.A. 1.4, Olympus) at

37°C with 5% CO_2 flow. The mitochondrial movement was tracked using the ImageJ plugin

MTrackJ. The mitochondrial fission and fusion events were counted manually. Peripheral and

midzone mitochondrial fissions were defined as events in which the relative position of the

fission site along the length axis was 0~25% and 25~50%, respectively. The percentage of

motile mitochondria was calculated by dividing the number of mitochondria that were displaced

more than 4 μm from their original positions within 20 minutes by the total number of

mitochondria. A continuous movement of mitochondria (> 1.5 μm for > 6 sec) was defined as

one moving event, and their speed was calculated by using MATLAB software.

For AMPK FRET imaging, cultured hippocampal neurons transfected with AMPKAR-

EV were imaged every 3 sec for 3min at DIV5 with an inverted epifluorescence microscope

(IX83, Olympus) equipped with LUCPlanFLN 20× (NA 0.45), UplanSApo 40× (Silicone oil,

NA 1.25) objectives. For dual imaging of AMPK FRET and calcium imaging, cultured

hippocampal neurons were cotransfected with AMPKAR-EV and jRGECO1a, and imaged

every 5 sec for 5 min. Fluorescence images were acquired with the following filters and mirrors:

excitation filters FF02-438/24-25 (for CFP/ YFP); a dichroic mirror FF458-Di02-25x36 (for

CFP/ YFP); emission filters FF01-483/32-25 (for CFP); FF01-542/27-25 (for YFP); and filter

sets U-FMCHE (Ex 565-585 nm/ DM 595/ Em 600-690) (for jRGECO1a).

For FRET ratio analysis, after background subtraction, the YFP/CFP fluorescence

intensity ratio was calculated in the ROI of the proximal dendritic region. The YFP/CFP ratio

was plotted along the time axis, and the FRET peak frequency was quantified using the

MATLAB findpeaks function with a threshold for peak heights greater than 0.05. The Cross-

Correlation Function was calculated followed methods described previously [62] using MATLAB

software.

For TMRM imaging, cultured hippocampal neurons transfected with mito-EGFP

were incubated with 20 nM TMRM 1-2 h prior to imaging. Time-lapse images were acquired

every 5 sec for 10 min at DIV5 with a spinning-disc confocal microscope (CV1000, Yokogawa)

equipped with a UPLSApo 100× objective (Oil immersion, N.A. 1.4, Olympus).

For TMRM intensity analysis, after background subtraction, TMRM intensity in

individual mitochondria was divided by the mitochondrial area. TMRM flickering, defined as a

transient loss of TMRM intensity to the background level, was scored as the number of

flickering events that occurred within 10 minutes in the soma and proximal dendrites, divided

by the mitochondrial area.

For quantification of mitochondrial ROS, cultured hippocampal neurons transfected

with mito-EGFP at DIV3 were incubated with 500 nM Mito-SOX for 15 min at DIV5 and then

the Mito-Sox was washed out. Time-lapse imaging was initiated 18-20 h after treatment, using a

spinning-disc confocal microscope (CV1000, Yokogawa) equipped with a UPLSApo 100×

objective (Oil immersion, N.A. 1.4, Olympus). Mito-SOX intensity in individual mitochondria

$((F_{mitochondria}/ F_{background})/ (Mitochondrial\ Area))$ was quantified using ImageJ.

Acknowledgements

We thank Dr. Michiyuki Matsuda for the AMPKAR-EV plasmid, Dr. Yoshiaki Tagawa for the

GCaMP6s plasmid, Dr. Masayuki Sakamoto for the jRGECO1a plasmid, Drs. Kuniya Abe,

Yusuke Kishi, Shinnosuke Suzuki, Ryuta Kinoshita for the CRISPR-dCas9KRAB plasmid,

iCeMS Analysis Center for technical assistance with microscopy, Drs. Takaki Miyata, Mayumi

Okamoto, Daniel Packwood for advice, Dr. James Hejna for critical reading of the manuscript.

Author contributions

Conceptualization: A.H., K.F., M.K.; Methodology: A.H., K.F., A.K., N.O., M.K.; Investigation:

A.H., J.K., K.F.; Writing - original draft: M.K. and A.H., and all the authors read and

commented on the manuscript.

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) Kakenhi

(20H00483 and JP16H06280) and the Uehara Memorial Foundation to M.K., and a JST spring

grant (JPMJSP2110) to A.H.

Reference

1. Fukumitsu, K. *et al.* Synergistic Action of Dendritic Mitochondria and Creatine Kinase

Maintains ATP Homeostasis and Actin Dynamics in Growing Neuronal Dendrites. *J.*

Neurosci. **35**, 5707–5723 (2015).

2. Sheng, Z. H. & Cai, Q. Mitochondrial transport in neurons: Impact on synaptic

homeostasis and neurodegeneration. *Nat. Rev. Neurosci.* **13**, 77–93 (2012).

3. Rangaraju, V. *et al.* Pleiotropic Mitochondria: The Influence of Mitochondria on

Neuronal Development and Disease. *J. Neurosci.* **39**, 8200–8208 (2019).

4. Schon, E. A. & Przedborski, S. Mitochondria: The Next (Neurode) Generation. *Neuron*

70, 1033–1053 (2011).

5. Pekkurnaz Gulcin & Wang, X. Mitochondrial heterogeneity and homeostasis through the

lens of a neuron. *Nat. Metab.* (2022). doi:10.1038/s42255-022-00594-w

6. Youle, R. J. *et al.* Mitochondrial Fission, Fusion, and Stress. *Science* **337**, 1062–1065

(2012).

7. Gandre-Babbe, S. & van der Bliek, A. M. The Novel Tail-anchored Membrane Protein

Mff Controls Mitochondrial and Peroxisomal Fission in Mammalian Cells. *Mol. Biol.*

Cell **19**, 2402–2412 (2008).

8. Palmer, C. S. *et al.* MiD49 and MiD51, new components of the mitochondrial fission

machinery. *EMBO Rep.* **12**, 565–573 (2011).

9. Yoon, Y., Krueger, E. W., Oswald, B. J. & McNiven, M. A. The Mitochondrial Protein

hFis1 Regulates Mitochondrial Fission in Mammalian Cells through an Interaction with

the Dynamin-Like Protein DLP1. *Mol. Cell. Biol.* **23**, 5409–5420 (2003).

10. Li, Z., Okamoto, K. I., Hayashi, Y. & Sheng, M. The importance of dendritic

mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell* **119**, 873–

887 (2004).

11. Ishihara, N. *et al.* Mitochondrial fission factor Drp1 is essential for embryonic

development and synapse formation in mice. *Nat. Cell Biol.* **11**, 958–966 (2009).

12. Fukumitsu, K. *et al.* Mitochondrial fission protein Drp1 regulates mitochondrial

transport and dendritic arborization in cerebellar Purkinje cells. *Mol. Cell. Neurosci.* **71**,

56–65 (2016).

13. Chen, H., Mccaffery, J. M. & Chan, D. C. Mitochondrial Fusion Protects against

Neurodegeneration in the Cerebellum. *Cell* **130**, 548–562 (2007).

14. Wang, X. *et al.* Impaired balance of mitochondrial fission and fusion in Alzheimer's

disease. *J. Neurosci.* **29**, 9090–9103 (2009).

15.   Herzig, S. & Shaw, R. J. AMPK: Guardian of metabolism and mitochondrial

homeostasis. *Nat. Rev. Mol. Cell Biol.* **19**, 121–135 (2018).

16.   Ducommun, S. *et al.* Motif affinity and mass spectrometry proteomic approach for the

discovery of cellular AMPK targets : Identi fication of mitochondrial fission factor as a

new AMPK substrate. *Cell. Signal.* **27**, 978–988 (2015).

17.   Toyama, E. Q. *et al.* AMP-activated protein kinase mediates mitochondrial fission in

response to energy stress. *Science.* **351**, 275–281 (2016).

18.   Jäer, S., Handschin, C., St-Pierre, J. & Spiegelman, B. M. AMP-activated protein kinase

(AMPK) action in skeletal muscle via direct phosphorylation of PGC-1α. *Proc. Natl.*

Acad. Sci. U. S. A. **104**, 12017–12022 (2007).

19.   Chen, Z. *et al.* Global phosphoproteomic analysis reveals ARMC10 as an AMPK

substrate that regulates mitochondrial dynamics. *Nat. Commun.* **10**, (2019).

20.   Egan, D. F. *et al.* Phosphorylation of ULK1 (hATG1) by AMP-Activated Protein Kinase

Connects Energy Sensing to Mitophagy. *Science.* **331**, 456–461 (2011).

21.   Kim, J., Kundu, M., Viollet, B. & Guan, K. L. AMPK and mTOR regulate autophagy

through direct phosphorylation of Ulk1. *Nat. Cell Biol.* **13**, 132–141 (2011).

22.   Tilokani, L. *et al.* AMPK-dependent phosphorylation of MTFR1L regulates

mitochondrial morphology. *Sci. Adv.* **8**, 1–20 (2022).

23.   Hawley, S. A. *et al.* Complexes between the LKB1 tumor suppressor, STRADα/β and

MO25α/β are upstream kinases in the AMP-activated protein kinase cascade. *J. Biol.* **2**,

1–16 (2003).

24.   Hawley, S. A. *et al.* Calmodulin-dependent protein kinase kinase-β is an alternative

upstream kinase for AMP-activated protein kinase. *Cell Metab.* **2**, 9–19 (2005).

25.   Hurley, R. L. *et al.* The Ca2+/calmodulin-dependent protein kinase kinases are AMP-

activated protein kinase kinases. *J. Biol. Chem.* **280**, 29060–29066 (2005).

26.   Mairet-Coello, G. *et al.* The CAMKK2-AMPK Kinase Pathway Mediates the

Synaptotoxic Effects of Aβ Oligomers through Tau Phosphorylation. *Neuron* **78**, 94–108

(2013).

27.   Lee, A. *et al.* Aβ42 oligomers trigger synaptic loss through CAMKK2-AMPK-

dependent effectors coordinating mitochondrial fission and mitophagy. *Nat. Commun.*

13, 1–20 (2022).

28.   Redmond, L., Kashani, A. H. & Ghosh, A. Calcium regulation of dendritic growth via

628   CaM kinase IV and CREB-mediated transcription. *Neuron* **34**, 999–1010 (2002).

629 29. Wong, R. O. L. & Ghosh, A. Activity-dependent regulation of dendritic growth and

630   patterning. *Nat. Rev. Neurosci.* **3**, 803–812 (2002).

631 30. Konur, S. & Ghosh, A. Calcium signaling and the control of dendritic development.

632   *Neuron* **46**, 401–405 (2005).

633 31. Divakaruni, S. S. *et al.* Long-Term Potentiation Requires a Rapid Burst of Dendritic

634   Mitochondrial Fission during Induction. *Neuron* **100**, 860-875.e7 (2018).

635 32. Wu, Y. K., Fujishima, K. & Kengaku, M. Differentiation of apical and basal dendrites in

636   pyramidal cells and granule cells in dissociated hippocampal cultures. *PLoS One* **10**, 1–

637   16 (2015).

638 33. Spronsen, M. V. *et al.* TRAK/Milton Motor-Adaptor Proteins Steer Mitochondrial

639   Trafficking to Axons and Dendrites. *Neuron* **77**, 485–502 (2013).

640 34. Faits, M. C., Zhang, C., Soto, F. & Kerschensteiner, D. Dendritic mitochondria reach

641   stable positions during circuit development. *Elife* **5**, 1–17 (2016).

642 35. Loss, O. & Stephenson, F. A. Developmental changes in trak-mediated mitochondrial

643   transport in neurons. *Mol. Cell. Neurosci.* **80**, 134–147 (2017).

644 36. Lewis, T. L., Kwon, S. K., Lee, A., Shaw, R. & Polleux, F. MFF-dependent

645   mitochondrial fission regulates presynaptic release and axon branching by limiting

646   axonal mitochondria size. *Nat. Commun.* **9**, 1–15 (2018).

647 37. Konagaya, Y. *et al.* A Highly Sensitive FRET Biosensor for AMPK Exhibits

648   Heterogeneous AMPK Responses among Cells and Organs. *Cell Rep.* **21**, 2628–2638

649   (2017).

650 38. Sample, V., Ramamurthy, S., Gorshkov, K., Ronnett, G. V. & Zhang, J. Polarized

651   activities of AMPK and BRSK in primary hippocampal neurons. *Mol. Biol. Cell* **26**,

652   1935–1946 (2015).

653 39. Barnes, A. P. *et al.* LKB1 and SAD Kinases Define a Pathway Required for the

654   Polarization of Cortical Neurons. *Cell* **129**, 549–563 (2007).

655 40. Courchet, J. *et al.* Terminal Axon Branching Is Regulated by the LKB1-NUAK1 Kinase

656   Pathway via Presynaptic Mitochondrial Capture. *Cell* **153**, 1510–1525 (2013).

657 41. Weisová, P., Concannon, C. G., Devocelle, M., Prehn, J. H. M. & Ward, M. W.

658   Regulation of glucose transporter 3 surface expression by the AMP-activated protein

659   kinase mediates tolerance to glutamate excitation in neurons. *J. Neurosci.* **29**, 2997–3008

660   (2009).

42.   Connolly, N. M. C., Düssmann, H., Anilkumar, U., Huber, H. J. & Prehn, J. H. M.
Single-cell imaging of bioenergetic responses to neuronal excitotoxicity and oxygen and
glucose deprivation. *J. Neurosci.* **34**, 10192–10205 (2014).

43.   Wayman, G. A. *et al.* Regulation of Axonal Extension and Growth Cone Motility by
Calmodulin-Dependent Protein Kinase I. *J. Neurosci.* **24**, 3786–3794 (2004).

44.   Wayman, G. A. *et al.* Activity-Dependent Dendritic Arborization Mediated by CaM-
Kinase I Activation and Enhanced CREB-Dependent Transcription of Wnt-2. *Neuron*
50, 897–909 (2006).

45.   Kleele, T. *et al.* Distinct fission signatures predict mitochondrial degradation or
biogenesis. *Nature* **593**, 435–439 (2021).

46.   Buckman, J. F. & Reynolds, I. J. Spontaneous Changes in Mitochondrial Membrane
Potential in Cultured Neurons. *J. Neurosci.* **21**, 5054–5065 (2001).

47.   Bonora, M., Giorgi, C. & Pinton, P. Molecular mechanisms and consequences of
mitochondrial permeability transition. *Nat. Rev. Mol. Cell Biol.* **23**, 266–285 (2022).

48.   Zorov, D. B., Juhaszova, M. & Sollott, S. J. Mitochondrial reactive oxygen species
(ROS) and ROS-induced ROS release. *Physiol. Rev.* **94**, 909–950 (2014).

49.   Twig, G. *et al.* Fission and selective fusion govern mitochondrial segregation and
elimination by autophagy. *EMBO J.* **27**, 433–446 (2008).

50.   Lewis, S. C., Uchiyama, L. F. & Nunnari, J. ER-mitochondria contacts couple mtDNA
synthesis with Mitochondrial division in human cells. *Science.* **353**, (2016).

51.   DuBoff, B., Götz, J. & Feany, M. B. Tau Promotes Neurodegeneration via DRP1
Mislocalization In Vivo. *Neuron* **75**, 618–632 (2012).

52.   Di Meo, D., Ravindran, P., Sadhanasatish, T., Dhumale, P. & Püschel, A. W. The
balance of mitochondrial fission and fusion in cortical axons depends on the kinases
SadA and SadB. *Cell Rep.* **37**, 110141 (2021).

53.   Joseph, B. K. *et al.* Inhibition of AMP kinase by the protein phosphatase 2A
heterotrimer, PP2APpp2r2d. *J. Biol. Chem.* **290**, 10588–10598 (2015).

54.   Dzamko, N. *et al.* AMPK β1 deletion reduces appetite, preventing obesity and hepatic
insulin resistance. *J. Biol. Chem.* **285**, 115–122 (2010).

55.   Williams, T., Courchet, J., Viollet, B., Brenman, J. E. & Polleux, F. AMP-activated
protein kinase (AMPK) activity is not required for neuronal development but regulates
axogenesis during metabolic stress. *Proc. Natl. Acad. Sci.* **108**, 5849–5854 (2011).

56.   Ramamurthy, S., Chang, E., Cao, Y., Zhu, J. & Ronnett, G. V. AMPK activation

regulates neuronal structure in developing hippocampal neurons. *Neuroscience* **259**, 13–

24 (2014).

57. Muraleedharan, R. *et al.* AMPK-Regulated Astrocytic Lactate Shuttle Plays a Non-Cell-

Autonomous Role in Neuronal Survival. *Cell Rep.* **32**, (2020).

58. Muraleedharan, R. & Dasgupta, B. AMPK in the brain: its roles in glucose and neural

metabolism. *FEBS J.* 1–16 (2021). doi:10.1111/febs.16151

59. Amato, S. *et al.* AMP-activated protein kinase regulates neuronal polarization by

interfering with PI3-kinase localization. *Science.* **332**, 247–251 (2011).

60. Cho, B. *et al.* Constriction of the mitochondrial inner compartment is a priming event for

mitochondrial division. *Nat. Commun.* **8**, 15754 (2017).

61. Umeshima, H., Hirano, T. & Kengaku, M. Microtubule-based nuclear movement occurs

independently of centrosome positioning in migrating neurons. *Proc. Natl. Acad. Sci. U.*

S. A. **104**, 16182–7 (2007).

62. Kunida, K., Matsuda, M. & Aoki, K. FRET imaging and statistical signal processing

reveal positive and negative feedback loops regulating the morphology of randomly

migrating HT-1080 cells. *J. Cell Sci.* **125**, 2381–2392 (2012).

Figure 1. Neuronal activity enhances dendrite formation and mitochondrial fission in developing hippocampal neurons.
(A) GCAMP6s ΔF/F transients in cultured hippocampal neurons before and after TTX+APV treatment (see also Supplementary Video 1). **(B)** Morphology of hippocampal neurons treated with vehicle or TTX+APV from DIV3 to 5. Cells were labeled with GFP (see also Supplementary Video 2). **(C, D)** Quantification of total dendritic length (C) and number of branch points (D). N= 23 for control cells, 30 for TTX+APV treated cells. ***p= $4.8×10^{-5}$ (C); ***p= $1.8×10^{-4}$ (D); Wilcoxon rank-sum test. **(E, F)** Morphology of mitochondria in dendrites (E) and axons (F) of DIV5 hippocampal neurons treated with vehicle or TTX+APV for 6h. Cells were labeled with mito-DsRed and GFP. **(G, H)** Distribution (G) and cumulative frequency (H) of the mitochondrial length in dendrites. N= 576 mitochondria from 23 control cells, 613 mitochondria from 24 drug-treated cells. ***p= $1.3×10^{-9}$, Wilcoxon rank-sum test. **(I)** Average length of dendritic mitochondria in individual neurons. N= 23 control cells, 24 TTX+APV treated cells. ***p= $1.2×10^{-5}$, Wilcoxon rank-sum test. **(J)** Quantification of the mitochondrial fission and fusion frequencies in dendrites before and 1-3h after TTX+APV treatment. N= 14 cells, mean± SEM, ***p= $4.1×10^{-4}$ for fission, (n.s.) p= 0.57 for fusion, paired two-tailed t test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (C, D, G, I). Samples were collected from three independent experiments. Scale bars: 50 μm in (B), 5 μm in (E, F).

Figure 2. AMPK promotes dendrite formation of hippocampal neurons.

(A) Expression profiles of AMPKα subunits in developing hippocampal neurons in culture. Whole cell lysates were collected at indicated stages and subjected to western blotting with antibodies against AMPKα1 and/or α2, and AMPKα1/2 phosphorylated at threonine 172 (p-AMPKα). **(B)** Morphology of hippocampal neurons transfected with GFP and the following constructs: shRNA-scramble control (SCR); shRNA-AMPKα2 #1 (AMPKα2 KD); or shRNA-AMPKα2 #1 with shRNA-resistant AMPKα2 (Rescue). Cells were transfected at DIV3 and fixed at DIV5. **(C, D)** Quantification of total dendritic length (C) and number of branch points (D). N= 31 cells for SCR, 30 cells for AMPKα2 KD, and 26 cells for Rescue. ***p= 4.0×10^{-4} for SCR vs. AMPKα2 KD, **p= 0.0058 for AMPKα2 KD vs. Rescue (C); ***p= 2.0×10^{-4} for SCR vs. AMPKα2 KD, *p= 0.018 for AMPKα2 KD vs. Rescue (D), Kruskal–Wallis test with Bonferroni multiple comparisons test. **(E, F)** Morphology of CA1 pyramidal neurons transfected with GFP and shRNA-scramble control (SCR) or shRNA-AMPKα2 #1 (AMPKα2 KD) by in utero electroporation at E14.5, followed by fixation at P5 (E) and P10 (F). **(G, H)** Quantification of total dendritic length (G) and number of branch points (H) in CA1 pyramidal neurons at P5 and P10. N≥ 16 cells for SCR and AMPKα2 KD, **p= 0.0067 for P5; (n.s.) p= 0.20 for P10 (G); **p= 0.0021 for P5; (n.s.) p= 0.12 for P10 (H), Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (C, D, G, H). Samples were collected from three independent experiments. Scale bars: 50 μm in (B, E, F).

Figure 3. AMPK enhances mitochondrial fission in dendrites of developing hippocampal neurons.

(A, B) Morphology of mitochondria in dendrites (A) and axons (B) in hippocampal neurons transfected with GFP and mito-DsRed together with shRNA-scramble control (SCR), shRNA-AMPKα2#1 (AMPKα2 KD), or shRNA-AMPKα2#1 with shRNA-resistant AMPKα2 (Rescue). Cells were transfected at DIV3 and fixed at DIV5. **(C, D)** Distribution (C) and cumulative frequency (D) of the length of dendritic mitochondria in SCR, AMPKα2 KD and Rescue cells. N= 669 mitochondria from 26 cells (SCR), 548 mitochondria from 22 cells (AMPKα2 KD), and 543 mitochondria from 22 cells (Rescue), respectively. ***p< 0.001; (n.s.) p> 0.05; Kruskal–Wallis test with Bonferroni multiple comparisons test. **(E)** Average length of dendritic mitochondria in individual cells. N=26 cells (SCR), 22 cells (AMPKα2 KD), and 22 cells (Rescue), respectively. ***p< 0.001; (n.s.) p> 0.05; Kruskal–Wallis test with Bonferroni multiple comparisons test. **(F)** Quantification of the mitochondrial fission and fusion frequencies in dendrites in SCR and AMPKα2 KD cells. N= 12 cells for SCR, and 13 cells for AMPKα2 KD. Mean ± SEM, **p= 0.0042; (n.s.) p= 0.75; unpaired two-tailed t test. **(G)** Morphology of dendritic mitochondria in hippocampal neurons before and 3-5 h after treatment with 1 mM AICAR (AMPK activator). Cells were labeled with tdTomato and mito-EGFP. **(H, I)** Distribution (H) and cumulative frequency (I) of the length of dendritic mitochondria before and after AICAR treatment. N= 230 mitochondria from 12 control cells, 228 mitochondria from 12 drug-treated cells, ***p= 1.9×10^{-11}; Wilcoxon rank-sum test. **(J)** Average length of dendritic mitochondria in individual cells. N= 12 cells, *** p= 4.9×10^{-4}, Wilcoxon signed-rank test. **(K)** Morphology of dendritic mitochondria in hippocampal neurons treated with vehicle or 20 μM Compound C for 6h. Cells were labeled with GFP and mito-DsRed. **(L, M)** Distribution (L) and cumulative frequency (M) of the length of dendritic mitochondria treated with vehicle or Compound C. N= 575 mitochondria from 22 control cells, 483 mitochondria from 19 drug-treated cells, ***p= 8.5×10^{-8}, Wilcoxon rank-sum test. **(N)** Average length of the dendritic mitochondria in individual cells. N=22 control cells, 19 drug-treated cells, *** p= 2.9×10^{-4}, Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (C, E, H, J, L, N). Samples were collected from three independent experiments. Scale bars: 5 μm in (A, B, G, K).

Figure 4. Neuronal activity induces dynamic oscillation of AMPK activity via Ca2+/CaMKK2.

(A) Representative traces of AMPK activity (YFP/CFP ratio) in dendrites of hippocampal neurons expressing AMPKAR-EV with shRNA-scramble control (SCR) or shRNA-AMPKα2 #1 (AMPKα2 KD). Time-lapse images were acquired every 3 sec for 3 min at DIV5 using epifluorescence microscopy. Red arrowheads indicate peaks of AMPK activity. (B) Quantification of the frequency of AMPKAR-EV FRET peaks in hippocampal neurons described in (A). N= 46 cells for SCR, 40 cells for AMPKα2 KD; ***p= 1.2×10⁻⁸; Wilcoxon rank-sum test. (C) Representative traces of AMPK activity in hippocampal neurons expressing AMPKAR-EV before and after TTX+APV treatment. (D) Quantification of the frequency of AMPKAR-EV FRET peaks described in (C). N= 22 cells; ***p= 1.6×10⁻⁴; Wilcoxon signed-rank test. (E, F) Representative time-lapse images and traces of AMPKAR-EV FRET signals and Ca²⁺ influx in hippocampal neurons transfected with AMPKAR-EV and jRGECO1a. Scale bar: 10 μm (E). Blue and orange lines indicate AMPKAR-EV FRET and jRGECO1a signal intensity, respectively (F) (see also Supplementary Video 4). (G) The temporal cross-correlations between AMPK activity and Ca²⁺ influx. Gray lines show individual cells, and the red line indicates the average temporal cross-correlation function. The arrowhead shows the highest correlation at 5 sec after Ca²⁺ influx. N= 33 cells. (H) Representative traces of AMPK activity in hippocampal neurons expressing AMPKAR-EV with shRNA-scramble control (SCR) or shRNA-CaMKK2 (CaMKK2 KD). (I) Quantification of the frequency of AMPKAR-EV FRET peaks described in (H). N=32 cells for SCR, 45 cells for CaMKK2 KD; ***p= 6.8×10⁻⁴; Wilcoxon rank-sum test. (J) Quantification of the frequency of AMPKAR-EV FRET peaks in hippocampal neurons expressing AMPKAR-EV before and 2h after treatment with 10 μM STO-609 (CaMKK2 inhibitor). N= 17 cells from two independent experiments; *p= 0.011; Wilcoxon signed-rank test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (B, D, I, J). Samples were taken from three independent experiments in (B, D, G, I).

Figure 5. AMPK knockdown suppresses asymmetric mitochondrial fission and mitophagy in growing dendrites.

(A) Time-lapse images of mitochondria undergoing midzone or peripheral fission in dendrites of hippocampal neurons. Cells were labelled with GFP and mito-DsRed. The red arrows indicate fission sites. **(B)** Quantification of the frequencies of midzone and peripheral fissions of dendritic mitochondria in hippocampal neurons before and after treatment with TTX+APV. N= 14 cells; mean ± SEM; (n.s.) p= 0.060 for midzone fission; **p= 0.0025 for peripheral fission; paired two-tailed t test. **(C)** Quantification of the frequencies of midzone and peripheral fissions of dendritic mitochondria in hippocampal neurons transfected with shRNA-scramble control (SCR) or shRNA-AMPKα2 #1 (AMPKα2 KD). N= 12 cells for SCR, 13 cells for AMPKα2 KD; mean ± SEM; *p= 0.028 for midzone fission; **p= 0.0020 for peripheral fission; unpaired two-tailed t test. **(D, E)** Immunofluorescence of autophagy marker p62 in dendrites of hippocampal neurons expressing shRNA-scramble control (SCR) or shRNA-AMPKα2 (AMPKα2 KD). Cells were labeled with GFP and mito-DsRed. Yellow lines indicate the outlines of dendrites (D). Higher magnified views and surface rendering images of mitochondria (magenta) and p62 puncta (green) in control and AMPKα2 KD cells (E). **(F)** Quantification of the number of p62 puncta in SCR and AMPKα2 KD cells. The number of p62 puncta was divided by the dendritic area. N= 40 cells for SCR, 31 cells for AMPKα2; ***p= 2.9×10^{-7}; Wilcoxon rank-sum test. **(G)** Quantification of the co-localization (Manders overlap coefficient) of p62 puncta and mitochondrial area in SCR and AMPKα2 KD cells. N= 40 cells for SCR, 31 cells for AMPKα2; ***p= 2.1×10^{-7}; Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (F, G). Samples were taken from three independent experiments. Scale bars: 1 μm in (A, E), 5 μm in (D).

Figure 6. AMPK phosphorylates mitochondrial fission factor (MFF) and ULK1 in response to neuronal activity.

(A) Phosphorylation of AMPKα and its downstream effectors in neurons treated with either TTX+APV (6h), 100 μM glutamate (5 and 10 min), 20 μM Compound C (CC, 6h) or 2 mM AICAR (6h) at DIV5. Blots were probed with antibodies against total and p-Thr172 AMPKα, total and p-Ser146 MFF, total and p-Ser555 ULK1 and β-actin. Similar results were obtained from three independent cultures. **(B)** Time-lapse images of TMRM flickering in dendritic mitochondria in hippocampal neurons. Cells were transfected with mito-GFP and SCR or AMPKα2 shRNA and treated with 20 nM TMRM. Transient loss of TMRM fluorescence in a single mitochondrion is seen (yellow arrowheads) (see also Supplementary Video 5). **(C)** Quantification of the frequencies of TMRM flickering in control and AMPKα2 KD cells. N= 18 cells for SCR, 20 cells for AMPKα2 KD. Samples were taken from four independent experiments. *p= 0.041, Wilcoxon rank-sum test. **(D)** Mito-SOX signals in dendritic mitochondria in control and AMPKα2 KD cells. Cells were transfected with mito-GFP and SCR or AMPKα2 shRNA and treated with 500 nM Mito-SOX. **(E)** Quantification of the Mito-SOX intensity in individual mitochondria. N= 212 mitochondria from 13 cells for SCR, 225 mitochondria from 17 cells for AMPKα2 KD. Samples were taken from two independent experiments. ***p= 6.9×10^{-15}, Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (C, E). Scale bars: 1 μm in (B, D).

Extended Data Figure 1. Neuronal activity regulates mitochondrial dynamics.

(A) Motile fraction of mitochondria in dendrites before and after TTX+APV treatment. Cells were labeled with mito-DsRed and GFP. N= 14 cells, mean ± SEM, (n.s.) p= 0.45, unpaired two-tailed t test. **(B)** Mitochondrial speed before and after TTX+APV treatment. N= 294 events from 14 control cells, 376 events from 14 drug-treated cells, (n.s.) p= 0.52, Wilcoxon rank-sum test. **(C)** Mitochondrial distribution along the shaft of primary dendrites. The primary dendrites were compartmentalized by the relative length from the soma, with the tip of the dendrite as 100%. Hippocampal neurons expressing GFP and mito-DsRed were treated with vehicle or TTX+APV for 2 days from DIV3 and fixed at DIV5. Mitochondrial density was calculated by dividing the mitochondrial area (demarcated by mitoDsRed) by the dendritic area (demarcated by GFP). N= 23 control cells, 30 drug-treated cells, (n.s.) p> 0.05, unpaired two-tailed t test. **(D)** Time-lapse images of fission and fusion of dendritic mitochondria in hippocampal neurons labeled with GFP and mito-DsRed (see also Supplementary Video 3). **(E)** Mitochondrial morphology in soma and dendrites after 1-min treatment with vehicle or 100 μM Glutamate at DIV6. Scale bars: 1 μm in (D); 5 μm in (E). Samples were taken from three independent experiments.

Extended Data Figure 2. Validation of shRNA targeting AMPKα subunits.

(A) HEK293T cells were transfected with GFP and AMPKα1 or AMPKα2 cDNA plasmids with or without respective shRNA constructs. Cell lysates were prepared 2 days after transfection and analyzed by western blotting with antibodies against GFP, AMPKα1, AMPKα2 and β-actin. **(B)** DIV5 hippocampal neurons transfected with GFP and shRNA for the scrambled control (SCR) or AMPKα2 (α2 KD #1) were immunostained with an antibody against AMPKα2. Arrowheads indicate shRNA-transfected neurons. **(C)** Representative images of DIV5 hippocampal neurons transfected with GFP and shRNA for the scrambled control (SCR), AMPKα1 (α1 KD #1 or #2), AMPKα2 (α2 KD #1 or #2), AMPKα1 and α2 (α1KD #1 + α2KD #1), or AMPKα1 (α1 KD #1) or AMPKα2 (α2 KD #1) together with the respective shRNA-resistant mutants (Rescue). **(D, E)** Quantification of total dendritic length (D) and number of branch points (E) of hippocampal neurons described in (C). N= 29 cells for SCR, 29 cells for α1 KD #1, 22 cells for α2 KD #1 from three independent experiments, and 18 cells for α1KD #1 + α2KD #1 from two independent experiments; ***p<0.001; (n.s.) p> 0.05; Kruskal–Wallis test with Bonferroni multiple comparisons test. **(F)** Morphology of hippocampal neurons transfected with CRISPR interference plasmids for gRNA-empty control (Cont) or gRNA-AMPKα2 (AMPKα2 CRISPRi). Cells were transfected at DIV2 and fixed at DIV5. **(G, H)** Quantification of total dendritic length (G) and number of branch points (H) of neurons described in (F). N= 34 cells for control, 21 cells for AMPKα2 CRISPRi from three independent experiments. ***p= 9.7×10^{-4} (G), ***p= 6.4×10^{-4} (H), Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (D, E, G, H). Scale bars: 10μm in (B), 50 μm in (C, F).

Extended Data Figure 3. AMPK induces mitochondrial fission in dendrites of growing hippocampal neurons.

(A) Mitochondrial distribution along the shaft of primary dendrites. Hippocampal neurons were transfected with GFP, mito-DsRed and shRNA-scramble control (SCR) or shRNA-AMPKα2 (AMPKα2 KD) at DIV3 and fixed at DIV5. The primary dendrites were compartmentalized by the relative length from the soma, with the tip of dendrite defined as 100%. The mitochondrial density was calculated by dividing the mitochondrial area (demarcated by mitoDsRed) by the dendritic area (demarcated by GFP). N= 22 cells for SCR, 24 cells for AMPKα2 KD; (n.s.) p> 0.05; unpaired two-tailed t test. **(B)** Morphology of dendritic mitochondria in hippocampal neurons transfected at DIV2 with CRISPR interference plasmids for gRNA-empty control (Cont) or gRNA-AMPKα2 (AMPKα2 CRISPRi) and fixed at DIV5. Cells were labeled with mito-DsRed and GFP. Scale bars: 5 μm. **(C, D)** Distribution (C) and cumulative frequency (D) of the mitochondrial length in dendrites. N= 534 mitochondria from 22 control cells, 497 mitochondria from 21 CRISPRi cells; *** p= 8.0×10^{-12}; Wilcoxon rank-sum test. **(E)** Average mitochondrial length in individual cells. N=22 cells for control, 21 cells for CRISPRi; *** p= 2.3×10^{-6}; Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets in (C, E). Samples were taken from three independent experiments.

Extended Data Figure 4. AMPK activity in hippocampal neurons visualized by AMPKAR-EV.
(A) Subcellular distribution of AMPKAR-EV FRET signals in hippocampal neurons at DIV5. Cells were transfected with AMPKAR-EV together with shRNA-scramble control (SCR) or shRNA-AMPKα2 (AMPKα2 KD). Scale bar: 50 μm. **(B)** Activation of AMPK induced by neuronal activity. DIV5 hippocampal neurons expressing AMPKAR-EV with shRNA-scramble control (SCR) or shRNA-AMPKα2 #1 (AMPKα2 KD) were treated with 1 μM glutamate. The YFP/CFP ratio in dendrites was monitored every 2 min for 70 min before and after glutamate treatment. N= 18 cells for control and AMPKα2 from two independent experiments. **(C)** Representative traces of AMPK activity and Ca^{2+} influx in hippocampal neurons treated with TTX+APV. Cells were transfected with AMPKAR-EV and jRGECO1a. **(D)** Representative traces of AMPK activity and Ca^{2+} influx in hippocampal neurons transfected with AMPKAR-EV, jRGECO1a and shRNA-AMPKα2. **(E)** Quantification of the frequency of AMPKAR-EV FRET peaks in hippocampal neurons expressing AMPKAR-EV before and after treatment with 1 μM Pim1 (LKB1 inhibitor) at DIV5. N= 19 cells from two independent experiments, (n.s.) p= 0.30, Wilcoxon signed-rank test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets.

Extended Data Figure 5. Effects of AMPK deficiency on PGC1α phosphorylation and mitochondrial membrane potential.

(A) Phos-tag SDS-PAGE for a mitochondrial biogenesis marker, PGC1α in neurons treated with either TTX+APV (6h), 100 μM glutamate (5 or 10 min), 20 μM Compound C (CC, 6h), or 2 mM AICAR (6h). Cell lysates were subjected to Phos-tag SDS-PAGE (100 μM Phos-tag) and standard SDS-PAGE, followed by immunoblotting with antibodies for PGC1α and β-actin. **(B)** Representative images of dendritic mitochondria in neurons treated with 20 nM TMRM (magenta). Cells were transfected with mito-GFP (green) together with shRNA-scramble control (SCR) or shRNA-AMPKα2 (AMPKα2 KD). **(C)** Quantification of TMRM intensity in individual mitochondria. N= 227 mitochondria from 17 control cells, 216 mitochondria from 15 AMPKα2 KD cells from two independent experiments. (n.s.) p= 0.14, Wilcoxon rank-sum test. Box plots denote the median (50th percentile) and the 25th to 75th percentiles of datasets. Scale bars: 10 μm (B, upper panels), 1 μm (B, lower panels).

Supplementary information

Supplementary Video 1. Calcium transients in hippocampal neurons before and after TTX+APV treatment. Cells were transfected with GCAMP6s at DIV3. Images were acquired every 3 sec for 5 min at DIV5 using spinning disk confocal microscopy. Display rate: 10 frames/second. Scale bar: 10 μm. Related to Fig. 1A.

Supplementary Video2. Time-lapse movies of growing hippocampal neurons cultured in the presence or absence of TTX+APV. Cells were labeled with GFP. Images were acquired every 30 min for 52 h from DIV5 to 7 using spinning disk confocal microscopy. Display rate:10 frames/second. Scale bar: 50 μm. Related to Fig. 1B.

Supplementary Video3. Time-lapse movies of mitochondrial fission and fusion in dendrites Cells were labeled with mito-DsRed and GFP. Time-lapse images were acquired every 3 sec for 20 min at DIV5 using spinning disk confocal microscopy. Display rate: 8 frames/second. Scale bar: 1 μm. Related to Extended Data Fig. 1D.

Supplementary Video 4. Simultaneous imaging of AMPK activity and Ca^{2+} transients before and after TTX+APV treatment.
Cultured hippocampal neurons were transfected with AMPKAR-EV and jRGECO1a. The pseudo-color represents the YFP/CFP ratio (AMPK activity, left), and the Ca^{2+} influx (right), respectively. Images were acquired every 5 sec for 5 min at DIV5 using epifluorescence microscopy. Display rate: 5 frames/second. Scale bar: 10 μm. Related to Fig. 4E-G.

Supplementary Video 5. Time-lapse movie of mitochondrial TMRM flickering.
Cultured hippocampal neurons transfected with mito-GFP (green) were treated with 20 nM TMRM (magenta) at DIV5. Images were acquired every 5 sec for 10 min using spinning disk confocal microscopy. Display rate: 8 frames/ second. Scale bar: 1 μm. Related to Fig. 6B.