1

Spatial and functional arrangement of Ebola virus polymerase inside phase-separated viral factories

Jingru Fang[1,2], Guillaume Castillon[3,4], Sebastien Phan[4], Sara McArdle[1], Chitra Hariharan[1], Mark H. Ellisman[4], Ashok A. Deniz[2], Erica Ollmann Saphire[1,*]

[1]La Jolla Institute for Immunology, La Jolla, CA, USA, [2]Scripps Research, La Jolla, CA, USA, [3]CMM Electron Microscopy Core facility, University of California San Diego, La Jolla, CA, USA, [4]National Center for Microscopy and Imaging Research, Center for Research in Biological Systems, University of California San Diego, La Jolla, CA, USA.

*Corresponding author (erica@lji.org)

Abstract

Ebola virus (EBOV) infection induces formation of membrane-less, cytoplasmic compartments termed viral factories, in which multiple viral proteins gather and from which diverse viral biogenesis arises. Key to viral factory function is recruitment of the EBOV polymerase, a multifunctional machine that mediates replication and expression of the viral RNA genome. Here we show that intracellularly reconstituted EBOV viral factories are biomolecular condensates, with composition-dependent internal dynamics of exchange that likely facilitates viral replication. We report that EBOV viral factories display either droplet-like or network-like morphology, which could be influenced by multivalent intermolecular interactions between viral proteins. Within the viral factory, we found EBOV polymerase is not uniformly distributed, but instead clusters into foci. The distance between these foci increases when viral replication is enabled. This unique view of EBOV propagation suggests a form-to-function relationship that describes how physical properties and internal structures of biomolecular condensates influence and regulate viral biogenesis.

Introduction

Viruses are architects of cellular remodeling needed to fulfill life-cycle events inside host cells. Some viruses remodel host cells by inducing formation of nonequilibrium, membrane-less compartments, also termed inclusion bodies or viral factories (VFs), that separate essential viral replication and assembly events from other cellular processes[1-11]. As exemplified by many known cellular membrane-less organelles, such as the multiphasic nucleolus[12], different interdependent cellular processes can be spatially separated into distinct co-existing phases inside the membrane-less compartment. In contrast, how multiple interdependent viral biogenesis steps are coordinated inside viral factories remains unclear.

Among viruses that induce VF formation, many are non-segmented, negative-strand RNA viruses (nNSVs) [2,3,5-8,10,11]. The negative sense RNA genome of nNSVs is coated by oligomerized viral nucleoprotein to form a helical RNP structure. During viral biogenesis, the viral polymerase transcribes the negative-strand RNA genome (termed vRNA) into multiple mRNAs encoding individual viral genes that are translated by host cell ribosomes into viral proteins. Meanwhile, the same viral polymerase must also coordinate synthesis of a complementary, positive-strand of the viral genome, termed cRNA, which acts as a template to replicate progeny negative-strand vRNAs. As cRNA and vRNA are produced, both are immediately coated with the viral nucleoprotein to form helical cRNP and vRNP (ribonucleoprotein) assemblies, respectively. However, only vRNP, which is linked to viral polymerase, is assembled into progeny virions[13]. The occurrence of these different virus biogenesis reactions inside VFs and the presence of different species of viral RNAs with different functions and different fates likely requires some spatial regulation.

Recent studies indicated that several nNSV VFs are biomolecular condensates that have
viscoelastic material properties, including VFs derived from vesicular stomatitis, rabies, measles
viruses, and human metapneumovirus[11,14-17]. The importance and potential biological relevance
of these viscoelastic properties was highlighted by inhibition of RSV infection in an animal model
by a condensate-hardening drug[18].
Formation of biomolecular condensates follows basic principles of phase separation wherein
molecules separate from a well-mixed system into multiple phases with distinct constituents, at
different concentrations in each phase[19,20]. This phase separation in biological systems is driven
by multivalent intermolecular interactions, based on the intrinsic properties of the protein and RNA
constituents[21-24], and is further influenced by other features of the system's free energy
landscape[25,26], which include kinetic effects such as the macroscopic segregation of dynamically
asymmetric mixtures[27]. Under physiological conditions, the concentration of key constituents
essentially controls the state of intracellular phase separation, which can be approximated in a
phase diagram[19]. This theoretical framework can offer mechanistic and quantitative insights into
VF condensate dynamics from which biological functions arise or are enabled. Deciphering phase
behaviors of intracellular VF condensates, and the spatial localization of the steps of virus
biogenesis within them, will outline the form-to-function relationship of VFs and enhance our
understanding of nNSV replication.
Among nNSVs, Ebola virus is a zoonotic, human pathogen that causes near-annual outbreaks
of disease with up to 90% mortality (WHO[28], CDC[29]). The only FDA-approved vaccine for Ebola
virus protects against only Ebola virus Zaire (EBOV), but not against the other species, including
the Sudan ebolavirus linked to an outbreak of disease that began in Uganda in 2022. Further,
therapeutic antibodies approved thus far are also specific for EBOV and may poorly penetrate
immune-privileged sites where Ebola virus can lurk. Additional therapeutic strategies are needed,
and will be accelerated by a better understanding of essential and conserved intracellular events
in Ebola virus replication.
Here, we sought to define how EBOV VFs spatially accommodate and control viral RNA
synthesis with an emphasis on the organization of the EBOV polymerase complex within VFs. To
dissect the individual contribution of viral elements to VF organization and dynamics, we took a
bottom-up approach to reconstitute EBOV VFs within cells beginning with minimal viral protein
components and viral minigenome reporter RNA. We determined that intracellularly reconstituted
EBOV VFs exist as biomolecular condensates that have reduced internal dynamics upon
recruitment of EBOV polymerase. Although EBOV VFs typically have a droplet-shape, we
described an additional, network-like morphology of EBOV VFs. Using advanced confocal
microscopy, we found an unexpected punctate distribution pattern of EBOV polymerase inside
droplet-like VFs and that this distribution of EBOV polymerase links to polymerase-mediated viral
RNA synthesis. Leveraging thin-section electron microscopy (EM) and multi-tilt electron
tomography (ET), we further resolved several foci of EBOV polymerase at invaginated or
interconnected boundary of the network-like VF, surrounded by helical vRNP-like structures.
Collectively, these multi-scale imaging approaches provide an unprecedented view of the spatial
organization by which Ebola virus orchestrates multiple biogenesis steps and deploys viral
replication machinery inside viral factories.
93
Results
1. Engineering of a fluorescence protein-tagged EBOV VP35 for live cell imaging of
EBOV viral factories (VFs)
Intracellular EBOV VFs are found to undergo fusion during infection[6]. This observation served
as the first hint that EBOV VFs could be biomolecular condensates. Thus, we undertook a
quantitative approach to determine whether EBOV VFs are indeed fluid condensates rather than
solid aggregates and define the internal molecular dynamics within EBOV VFs. Since microscopy
of live, EBOV-infected cells under BSL4 containment is neither feasible nor amenable to

controlling VF composition, we first established a transfection-based, live-cell system to perform
fluorescence recovery after photobleaching (FRAP) analysis with intracellularly reconstituted
EBOV VFs.
To monitor EBOV VF with live-cell imaging, we adapted an approach used to image VFs in
live cells infected with Rabies (RABV)[15]. We fused an EBOV VF constituent, the polymerase
cofactor VP35, to an N-terminal HA epitope tag and a fluorescent protein, mNeonGreen (mNG)
to generate HA-mNG-VP35. We confirmed that HA-mNG-VP35 protein expresses at a similar
level as the wild-type (VP35-WT), and retains sufficient ability to act as the polymerase cofactor
that supports EBOV viral RNA synthesis (**Supplementary Figure 1a**).
VP35 interacts with EBOV nucleoprotein (NP) and chaperones the NP monomer (NP^0) prior
to NP oligomerization[30]. As a polymerase cofactor, VP35 bridges the EBOV large polymerase
protein (L) to NP-coated viral genomic material[31]. Since these multivalent intermolecular
interactions VP35 involved in might provide the foundation for biomolecule phase separation, we
first confirmed that the mNG tag did not affect these interactions. We performed HA-
immunoprecipitation (IP) using cells co-expressing NP and L with either mNG-HA-VP35, HA-
VP35 (positive control), or mNG-HA (negative control). In the presence of NP alone or with NP
and L, HA-VP35 and HA-mNG-VP35 both immunoprecipitated with NP and L (**Figure 1a**).
Immunofluorescence microscopy showed that the intracellular localization of HA-mNG-VP35
resembles that of VP35-WT, when co-expressed with NP and L (**Supplementary Figure 1b**).
Together, our results indicate that HA-mNG-VP35 preserves sufficient polymerase-cofactor
function and fully retains interactions with NP and L for use in live-cell imaging.

2. Reconstituted EBOV VFs display composition-dependent, viscoelastic properties in
live cells
To quantitatively measure the internal mobility of EBOV VF components in a cellular context,
we performed FRAP in HEK 293T cells co-transfected with EBOV NP and HA-mNG-VP35. Both
NP and VP35 proteins contain multiple disordered and low complexity regions (**Figure 1b**), bind
RNA[32,33] and self-oligomerize[34-37], which are common features associated with many known
cellular proteins to be involved in phase separation[19]. Here we used NP and VP35 as the minimal
components to reconstitute EBOV VFs.
Reconstituted EBOV VFs had diameters ranging from sub-micrometer to 10 μm
(**Supplementary Figure 1c**). Because the high mobility of sub-μm VF makes it challenging to
accurately measure fluorescence recovery within achievable microscopy frame-rate, we focused
on medium-sized VFs (4-5 μm diameter). We photobleached a 1 μm diameter center spot in the
VFs and monitored the fluorescence recovery of the bleached spot until the system returned to
equilibrium. To quantify the kinetics of fluorescence recovery from which we can extract
parameter values that describe the timescale of the fast vs. slow diffusion event and the fraction
of immobile molecules, we fitted double-normalized fluorescence intensity values from the
moment of photobleaching until equilibrium was reached to a two-phase association model
(**Methods**).
As a control, HA-mNG-VP35 expressed alone formed gel-like, dense phases in the cytoplasm
(**Figure 1c**). Most cells have large quantities of sub-μm VP35-condensates with high apparent
mobility relative to the dilute phase, but a small number of VP35-condensates had a larger size
suitable for FRAP. These large condensates have multiple dark internal regions that lacked
fluorescence. After photobleaching a center spot of VP35 condensates, we observed only partial
fluorescence recovery (**Figure 1f,** green curve). This slower recovery rate could be due to a non-
diffusion based binding reaction like that which occurs with VP35 self-oligomerization or RNA
binding, since diffusion of a similar-sized protein occurs within milliseconds for similar-sized
regions[38].
We next evaluated reconstituted EBOV VFs via co-expression of HA-mNG-VP35 and NP. The
combination of HA-mNG-VP35 and NP yielded homogenous binary condensates (**Figure 1d**) that

appeared to be fluid-like based on frequently observed fusions between condensates or fission
of one condensate into two parts (**Supplementary Figure 2a**), and on non-fluorescence objects
trafficking through VFs (**Supplementary Figure 2b**). Over half the transfected cells had easily
identifiable >5µm condensates. Fluorescence recovery inside VP35-NP condensates is highly
dynamic with faster molecular exchange than that measured in VP35-alone condensate (**Figure**
1f, red vs. green curves). This faster recovery likely reflects contributions of VP35-NP interactions.
In VP35-NP condensates, 84.6% of VP35 molecules (with a 95% CI [83.89%,85.90%]) are mobile,
while in VP35-alone condensates, <50% of VP35 are mobile. Moreover, the half-time of the fast
association phase ($t_{1/2FAST}$) for VP35-NP condensates was ~5.5 s, which is consistent with that
reported for VFs in RABV-infected cells[15] ($t_{1/2FAST}$ ~5.2 s), and the half time for the slow association
phase ($t_{1/2SLOW}$) was ~21.5 s (**Figure 1f**).
Co-expression of mNG-HA-VP35 with NP and EBOV large polymerase protein (L) resulted in
dense phases in the cytoplasm that contained all three proteins (**Supplementary Figure 1b**).
VP35 and NP had homogenous distribution in these dense ternary condensates, but whether the
distribution of L was also homogeneous was unclear (**Figure 1e**). In VP35-NP-L condensates,
78% of mNG-HA-VP35 (with a 95% CI [77.7%,78.4%]) are mobile (**Figure 1f**, blue curve), which
is on average 7% lower than VP35+NP, suggesting that VP35-L interactions may immobilize a
small population of VP35. In both binary and ternary condensates, bleach-pulses induced loss of
fluorescence signal was around 12% of the prebleach signal at equilibrium, which justifies the
comparison of VP35-mobile fraction in both condensates (**Supplementary Figure 2c).** Further,
the similar $t_{1/2FAST}$ and $t_{1/2SLOW}$ values for binary and ternary condensates likely indicates similar
molecular interactions mediate VP35 mobility, although the percentage of fast association events
fell from 80.3% to 64.3% (**Figure 1f**). This reduction suggests that L affects the stoichiometry of
different molecular interactions involving VP35.
Our FRAP results quantitatively showed that EBOV VFs reconstituted with NP and VP35
display composition-dependent, viscoelastic behaviors in live cells. The addition of L immobilizes
a small, but detectable, fraction of VP35 inside the reconstituted VFs.

3. Intracellular localization analysis of FLAG-tagged EBOV large polymerase protein L
with confocal fluorescence microscopy
We next assessed whether EBOV L is immobilized in a specific location within VFs using
immunofluorescence microscopy. To facilitate detection of L, we engineered recombinant L with
an N-terminal 2xFLAG tag (FLAG-L) as previously described[39] since the only currently available
L polyclonal antibody can non-specifically bind to proteins other than EBOV L (**Supplementary**
Figure 1d).
First, we co-expressed NP, VP35 and FLAG-L in HEK 293T cells and labeled EBOV VFs
using a monoclonal antibody targeting NP. Although NP was present throughout EBOV VFs, NP
molecules at the VF periphery were immunolabeled more efficiently than the NP inside VF,
leading to an "empty" droplet-like morphology of EBOV VFs, which was previously described[40,41].
Unexpectedly, FLAG-L was not homogenously distributed within reconstituted VFs and had a
different staining pattern than NP (**Figure 2a**). Instead, FLAG-L clustered into networks
comprising interconnected foci within the ternary condensate, suggesting that an intrinsic property
of L drives a different phase behavior than VP35 or NP.
To link this phase behavior of EBOV L to its biological function, we further analyzed L
localization in the presence of an active RNA substrate for L, the EBOV minigenome, which allows
reconstitution of L-mediated MG replication and transcription. The EBOV minigenome we used
here is bicistronic (2cis-MG), which contains two reporter genes encoding GFP (for imaging) and
Renilla luciferase (for quantification) in a tandem cassette carrying authentic EBOV gene start-
and end signals. EBOV regulatory sequences, the 3' leader and 5' trailer, required for replication,
transcription and encapsidation of viral RNAs[42], flank the bicistronic cassette. Using the
replication-competent version of the 2cis-MG (**Figure 2b**, Rep-comp. MG), we confirmed that the

L construct, FLAG-L, we engineered is competent to fulfill its biological function as it retained 74%
of the wildtype L (L-WT) activity (**Figure 2c**).
In GFP+ cells having successful reconstitution of EBOV MG replication and transcription,
FLAG-L still clustered into networks of interconnected foci inside VFs (**Figure 2d**). Besides, in
fixed cells, GFP reporter signals were preferentially trapped in both VFs and nucleoli, which both
have features of biomolecular condensates.
We next asked whether modulating viral RNA synthesis affects EBOV L organization inside
VFs. Since adding or removing MG system elements could affect the overall valency of protein-
protein/RNA interactions or the molecular composition inside EBOV VFs, we incorporated a
previously characterized mutation into the 5' trailer of the EBOV 2cis-MG that allows transcription
but disables vRNA replication[43]. Reporter activity measured in this replication-deficient MG (Rep-
def. MG) system reflects only viral transcription (**Figure 2b**, Rep-def. MG). With Rep-def. MG,
FLAG-L retained 33% of the L-WT activity, indicating that the 2xFLAG tag may have affected L-
mediated transcription (**Figure 2c**). Nevertheless, with FLAG-L, when replication was disabled,
the L foci inside EBOV VFs were more closely spaced compared to that seen for replication-
competent VFs (**Figure 2e, f**). This different spacing is unlikely to be associated with the
expression levels since FLAG-L was expressed at equivalent levels in the Rep-comp. and Rep-
def. MG systems (**Figure 2c**). Together, our results revealed a unique localization pattern of
EBOV polymerase L inside VF and established a potential link between the spatial distribution of
L within the VF and viral RNA synthesis events mediated by L.

4. Network-like VFs exist in EBOV-GFP-ΔVP30 infected cells
Among the EBOV VFs reconstituted with the transcription and replication of EBOV MG in
transfected HEK 293T cells, some display a granular, network-like morphology instead of the
typical droplet-like morphology (**Figure 3a**). We thus examined whether network-like VFs also
occur during virus infection.
To characterize VF morphology in EBOV-infected cells, we used the biologically contained
EBOV-GFP-ΔVP30 virus, which is morphologically indistinguishable from wild-type EBOV, but
approved for use in BSL2+ containment[44]. In this system, a GFP gene replaces the gene encoding
the viral transcription factor, VP30, so that EBOV-GFP-ΔVP30 virus can grow only in cell lines
stably expressing VP30. Here, Vero cells stably expressing EBOV VP30 (Vero-VP30) were
infected with EBOV-GFP-ΔVP30 in a synchronized manner and the cells were fixed and
inactivated at 18 hours post-infection, when the VF size is comparable to that in HEK 293T cells
transfected with the EBOV MG system[45]. We then used immunofluorescence-labeled NP as a
marker to detect VFs in GFP-positive (i.e., EBOV-GFP-ΔVP30 infected) Vero-VP30 cells.
EBOV-GFP-ΔVP30 infected cells had either droplet-like or network-like VFs (**Figure 3b**), with
the majority (77%) had droplet-like VFs and a stronger immunofluorescence staining of NP at the
VF periphery as previously reported[45]. The remaining 23% of infected cells harbored network-like
VFs, which, in contrast to droplet-like VFs located at discrete sites in the cytoplasm, occupied a
more extended region that included a group of small VFs either interconnected or separated by
only a small distance. Our results suggested that the both droplet-like and network-like
morphology of EBOV VFs exist during EBOV infection.

5. Engineering a split-APEX2 tag for electron microscopy analyses of EBOV polymerase
L-VP35 complexes
To dissect the spatial organization of EBOV polymerase inside VFs at nanometer resolution,
we used APEX2, a peroxidase tag engineered to indicate the location of tagged proteins in
electron microscopy (EM) imaging. Specifically, we used the split-APEX2 (sAPEX) system[46],
consisting of two inactive fragments, sAP and sEX. We genetically fused the small fragment sEX
to L (L-sEX) and the large fragment sAP, along with a V5 epitope tag, to VP35 (VP35-V5-sAP)

(**Figure 4a**). Interaction of L with VP35 during the formation of an active EBOV polymerase
complex joins the sAP and sEX fragments to reconstitute APEX2 peroxidase activity.
We confirmed that sAPEX-tagged EBOV polymerase is functionally active (**Figure 4b**), using
a Pol1-based, monocistronic EBOV minigenome (Pol1-MG) system[47] that works similarly to the
T7-based, bicistronic EBOV MG, but contains a single gene encoding the firefly luciferase reporter
(**Figure 4c**). VP35-V5-sAP expression levels were significantly higher than the V5-tagged VP35
control (VP35-V5). L-sEX was also expressed to higher levels than wild type L, which could
explain the correspondingly higher MG activity seen for sAPEX2-tagged EBOV polymerase
(**Figure 4d**).
In cells transfected with sAPEX2-tagged EBOV polymerase and the Pol1-MG system, we
confirmed a site-specific reconstitution of sAPEX2 activity, as evidenced by APEX2-mediated
conversion of a fluorogenic substrate at sites marked with VP35-V5-sAP immunofluorescence.
We noted that VP35-V5-sAP formed a network-like dense phase, whereas VP35-V5 formed
droplet-like dense phases (**Figure 4e,** upper panel). These two morphologies of VP35-dense
phase in transfected HEK293T cells are comparable to those we observed with EBOV VF in virus-
infected Vero-VP30 cells. Further, the two morphologies resemble possible outcomes of phase
separation occurring via nucleation-growth vs. spinodal decomposition[26,48]. In regions in the
phase diagram that correspond to a thermodynamically metastable state, a new phase as in a
small spherical "nucleus" can stochastically form and grow in size via intermolecular interactions.
Meanwhile, in regions that correspond to an unstable initial state, small fluctuations in composition
or density (i.e., increased concentration of one constituent) could translate into ripples in the
systems' free energy landscape, such that phase separation occurs spontaneously and features
in a network-like domain morphology that can be made persistent by kinetic arrest. The range of
intermolecular interactions underlining the dense vs. dilute phase controls the condensate
morphology[21,49].
The atypical network-like morphology of VP35-V5-sAP could be attributed to increased
valence of the inter-molecular interaction in the sAPEX2-tagged L-VP35 complex, since trans-
complementation of the sAPEX-tag creates an additional intermolecular interaction between L
and VP35 (**Figure 4e,** lower panel). To test this possibility independently of sAPEX2-tagging, we
increased the valence of inter-molecular interactions that involve VP35 by adding EBOV VP24,
an EBOV protein which also interacts with VP35[50]. Upon co-expression of VP24 with VP35-V5
and the Pol1-MG system components, VP35-V5 droplets were replaced by the network-like VP35-
V5 dense phase that colocalized with VP24 (**Figure 4f**, upper panel). This result could indicate
that increased valence of intermolecular interactions indeed alters VP35 phase behavior.
However, even in the presence of VP24, which is present during virus infection, the sAPEX2-
tagged EBOV L-VP35 complex remained in the network-like dense phase (**Figure 4f**, lower panel).
In summary, we successfully engineered an intracellularly active EBOV polymerase carrying
a split-APEX2 tag that will allow localization of EBOV polymerase in electron microscopy (EM)
analyses. Although the split-APEX2 tag did affect the intracellular localization pattern of VP35 in
cells reconstituted with the EBOV MG system, this altered VP35 pattern is likely not completely
artificial, since the same VP35 localization pattern occurred upon co-expression of EBOV VP24,
an EBOV protein present during natural EBOV infection. The altered VP35 pattern also coincides
with the network-like morphology of EBOV VF we also observed in EBOV-GFP-ΔVP30-infected
cells (**Figure 3b**). Therefore, we next carried out EM analysis on the localization of sAPEX2-
tagged EBOV polymerase in cells reconstituted with the Pol1-based EBOV MG system.

6. Nanoscale localization of split-APEX2-tagged EBOV polymerase complex with thin-
section transmission electron microscopy (thin-section TEM)
We used the sAPEX2 tag engineered into the L-VP35 complex, to locate the EBOV
polymerase within the compact cellular contents revealed by EM. Upon staining with 3,3′-
diaminobenzidine (DAB), trans-complementation of sAPEX2 catalyzes DAB polymerization with

minimal diffusion. The resulting DAB polymers alone at the site of active APEX2 are chromogenic,
which can be detected by light microscopy (LM). Further, DAB polymers are osmiophilic, and thus
capture osmium upon OsO_4 staining to increase electron density associated with DAB deposits
(**Figure 5a**). OsO4 also stains unsaturated lipids[51] and reacts with nucleic acids[52,53] to outline
cellular architecture in a specimen.
After optimization of transfection and staining conditions, we could detect sAPEX2-specific
DAB deposits in bright field light microscopy images of HEK 293T cells transfected with sAPEX2-
tagged EBOV polymerase together with other Pol1-MG components. DAB darkening was
intensified after osmification and DAB-positive (DAB+) cells were readily recognizable in resin-
embedded samples under light microscopy, which facilitates production of 70 nm thin-sections
containing a DAB+ cell and its detection with transmission electron microscopy (TEM) (**Figure
5b**). We observed within the same DAB+ cell network-like, electron-dense regions in cytoplasmic
areas that likely correspond to EBOV VFs (**Figure 5c, Supplementary Figure 3a**). These
electron-dense regions were present in cells transfected with the Pol1-MG system but not in
untransfected control cells (**Supplementary Figure 3b, c**). In the DAB+ cell, EBOV-specific
electron-dense regions were decorated with darker dots arranged in a pattern that correlated with
the shape of DAB darkening in light microscopy (**Figure 5c,** arrows), suggesting that the locations
of these darker dots in the electron micrograph corresponded to sites where sAPEX2-tagged
EBOV polymerases localize.
In the same thin-section, we also identified cells that had no DAB staining under light
microscopy (DAB-) (**Figure 5b**). These DAB- cells had the same transfection conditions as DAB+
cells but sAPEX2 activity was not reconstituted. DAB- cells exhibited network-like, electron-dense
VFs in the cytoplasm, similar to DAB+ cells (**Figure 5d**). By comparing the morphology of EBOV
VF in DAB+ and DAB- cells, the darker dots that associate with electron-dense VF in the DAB+
cell in fact indicate locations of sAPEX2-tagged EBOV polymerase (**Figure 5e**).
Our thin-section TEM observation revealed that sAPEX2-tagged EBOV polymerase non-
uniformly localizes to the periphery of network-like VFs. In several sites in the cytoplasm, sAPEX2-
tagged EBOV polymerase preferentially clusters at the junction of interconnected VFs.
Surrounding the electron-dense VFs were wire-like fragments that frequently associated with
neighboring VFs (**Figure 5c, d**).

7. 3D visualization of EBOV VFs and sAPEX2-tagged polymerase by electron tomography
To visualize the three-dimensional (3D) ultrastructure of EBOV VFs and elucidate the spatial
distribution of the sAPEX2-tagged polymerase complex, we applied a four-tilt electron
tomography (ET) collection scheme and reconstructed tomograms of subcellular volumes
containing EBOV VFs and sAPEX2-tagged polymerase. Similar to thin-section TEM specimens,
we transfected HEK 293T cells with the Pol1-based EBOV MG system containing sAPEX2-tagged
polymerase, except we generated 200-250 nm-thick sections from subcellular regions containing
DAB+ cells. Thin-section TEM allowed visualization of averaged 2D projection of all cellular
content across the thin-section specimen, but multi-tilt electron tomography allowed us to resolve
individual tomographic slices of the reconstructed volume with finer details of cellular matter
(**Figure 6a**). To understand the cellular surroundings of EBOV VF, we manually annotated a
representative tomogram at an organelle level (**Figure 6b**). Similar to previous EM analysis of
virus-infected cells[3,54], reconstituted EBOV VFs appeared as electron-dense clusters that lack
membrane boundaries themselves, but are in close proximity to membrane-bound cellular
organelles including mitochondria, the endoplasmic reticulum, the nuclear envelope and vesicles.
In addition to the membrane-bound organelles, the EBOV VFs and granularities in the cytosolic
surroundings were clearly resolved in the tomographic slices.
Across the tomographic slices, darker-stained patches near the edge of several electron-
dense VFs were apparent. By mapping these darker patches in the tomogram, we reconstructed
the 3D footprints of sAPEX2-tagged EBOV polymerase on a group of network-like VFs that were

in close proximity or even interconnected (**Figure 6b**). In the reconstructed tomograms the wire-
like fragments observed in thin-section TEM appeared to be continuous filaments having random
orientations. We then selected a region of interest from the whole tomogram, and performed a
finer manual segmentation focusing on smaller objects surrounding the VFs (**Figure 6c**).
sAPEX2-tagged EBOV polymerase appeared to dock at certain spots on the VF periphery.
Ribosomes were located outside, but adjacent to EBOV VFs. We observed several loosely coiled
structures either emerging from or passing through neighboring VFs, which likely gave rise to the
rough surface of the VFs and are the major component of VFs (**Figure 6d**). We propose that
these loosely coiled structures are the viral ribonucleoprotein complex (vRNP), which has at its
core helical viral genomic RNA bound to NP[55,56]. The helical structure is sufficiently relaxed to
allow the viral polymerase to slide along the genome during RNA synthesis. We saw no
condensed vRNP in our sample, likely because VP24, which has an inhibitory effect on vRNA
synthesis and can condense vRNP, was not included in the MG system[57,58].
In a selected region from the whole tomogram, two loose-coil structures can, in fact, be traced
back from their different branching sites to one parental loose coil (**Figure 6e**). One possible
explanation for these interconnected coils is that viral genomic RNA associates with active
transcription products. For several better-resolved loose coils in our tomogram, the coil terminus
often had a small loop linked to a solid globular structure. The identity of this terminal loop and
globular structure remains to be determined. In another selected region, several sAPEX2-tagged
EBOV spots were identified by manually tracing the darker stained patches on an invaginated
edge of electron-dense VFs. These darker-stained patches localized to the apical sides of
neighboring VFs where they formed a zipper-like structure (**Figure 6f**). In reconstructed
tomograms, the only obvious cellular structures that have direct contact with EBOV VFs are the
single-membrane vesicles (**Figure 6 b, f**).

Discussion
Ebola virus (EBOV) induces formation of viral factories (VFs) in host cells that allow physical
separation of viral biogenesis from other cellular processes. In this study we sought to understand
the mechanisms by which EBOV VFs spatially accommodate and control viral RNA synthesis,
with a particular focus on the localization of the EBOV polymerase complexes within VFs.
The internal dynamics we measured using FRAP for EBOV VFs produced in transfected cells
are strikingly comparable to those reported for VFs in rabies virus-infected cells[15]. Even though
the stoichiometry of transfected viral proteins cannot fully recapitulate that of virus infection, our
results demonstrate the validity of using transfection-based, intracellularly reconstituted VF to
study viscoelastic properties of VF in living cells.
The contribution of individual VF components to the internal molecular exchange within VFs
was poorly characterized in previous studies. Here, we used transfection-based VF reconstitution
and supplied individual EBOV components for stepwise assembly of intracellular VF. Our key
finding is a notable reduction in the molecular exchange rate within EBOV VFs upon recruitment
of L to the NP-VP35 binary condensate, which is likely due to interaction of L with a fraction of
mobile VP35. Reducing the internal molecular exchange rate within EBOV VF could functionally
impact viral RNA synthesis. Longer dwell times of EBOV polymerase and its cofactors inside VFs
may facilitate the initiation step of EBOV polymerase-mediated viral RNA synthesis. A slower
internal molecular exchange rate within EBOV VFs may also facilitate concurrent NP-
encapsidation of nascent v/cRNA. We additionally observed that VP35 expressed in the absence
of NP tends to form immobile aggregates, which indirectly supports a crucial role for NP as a
scaffold for VF condensates. Given that the EBOV NP protein is 3-times bigger than VP35, the
dynamic behavior of soluble NP[0]-VP35 complex[30] likely drives the mobility of VP35 in the NP-
bound state regardless of other molecular interactions VP35 is involved in. The same approach
we used here can be used to measure the contribution of other EBOV proteins, especially VP30,

to the internal dynamics within VFs and to characterize composition-dependent mobility of VFs
induced by other viruses.

Little was previously known about the functional localization of nNSV polymerases inside cells.
The lack of specific antibodies suitable for immunofluorescence labeling and the low protein
abundance of viral polymerase in infected cells has made imaging of nNSV polymerases
particularly challenging. An early study first identified that mCherry-tagged EBOV L polymerase
homogenously locates inside VFs during infection[6]. Despite the value of mCherry for live-cell
imaging, the activity of L-mCherry was only 10% that of unmodified L and the integrity of the L-
mCherry was unclear due to the lack of a detection antibody against L at the time of the study. In
contrast to insertion of a larger fluorescence protein, L protein can tolerate a small epitope tag
without substantial loss of RNA synthesis activity[59]. A recent study using super-resolution light
microscopy examined the intracellular localization of FLAG-tagged RSV L in RSV-infected cells[60].
In that study, RSV L concentrated non-uniformly at several sites inside RSV VFs. This result
aligns well with our confocal microscopy findings that FLAG-tagged EBOV L formed foci inside
VFs in cells reconstituted with the replication and expression of EBOV minigenome. A
recombinant EBOV expressing FLAG-tagged L should be generated to further validate our
findings with L localization in cells infected with authentic virus under BSL4 containment.

Beyond elucidating the functional localization of EBOV polymerase L, our work for the first
time links the localization pattern of EBOV L to particular types of L-mediated viral RNA synthesis.
We hypothesized that VFs have distinct sub-compartments that are dedicated to viral replication
or transcription. To dissect the EBOV L localization specific to replication or transcription, we
uncoupled L-mediated replication from transcription by generating a replication-deficient EBOV
minigenome (MG)[43]. We observed an intriguing difference in EBOV L localization in cells
reconstituted with replication-competent vs. replication-deficient MG systems. When viral
replication was active, the foci containing L were more widely spaced than when replication was
switched off. A wider spacing of L-foci could allow more NP to permeate and become available
for c/vRNA encapsidation, as reduced NP availability impairs viral genome replication[61].
Conversely, a narrower spacing of L-foci may reflect a switch from replication to transcription,
which produces flexible viral mRNAs that are less bulky, or may represent the waning activity of
overall viral RNA synthesis. Our result implies that, rather than having distinct sub-compartments
specialized for viral replication and transcription, the spatial organization of EBOV L expands and
contracts according to the type or activity of ongoing viral RNA synthesis.

In addition to the spatial-functional dynamics with EBOV polymerase, we revealed an atypical,
network-like morphology of EBOV VFs. In previous microscopy studies with virus-infected cells,
nNSV VFs, including those induced by EBOV, appear mostly as droplet-like structures[6-8,62]. We
did observe droplet-like VFs in EBOV-GFP-ΔVP30 infected cells, but a quarter of the cells
contained network-like VFs. Interestingly, some network-like EBOV VFs had similar shapes to
those observed in Marburg virus-infected cells[3]. The identification of both spherical and network-
like VFs in EBOV-infected cells suggests that the underlying mechanisms for EBOV VF phase
separation could include both nucleation-and-growth and spinodal decomposition[26,48]. Formation
of the network-like VFs could also be an outcome of viscoelastic phase separation of dynamically
asymmetric mixture[27], since we observed different viscoelastic behaviors in compositionally
distinct VF mixtures. Alternatively, the network-like VF could be driven by the formation of
extensive RNA-RNA interactions[63] that involve nascent viral RNAs or cellular RNAs. Future
experiments should identify the molecular basis of network-like EBOV VFs and elucidate the
biological role of such atypical VFs in virus infections.

Finally, we visualized the precise 3D localization of EBOV polymerase with nanometer
resolution in a subcellular volume using electron-tomography. Here EBOV polymerase with the
APEX2 EM-tag allowed localization of the polymerase among the heterogenous intracellular
content seen under electron microscopy[46]. Incorporating the split-APEX2 tag into L and VP35
preserved substantial levels of EBOV polymerase activity, although the valence of intermolecular

interaction was increased. This increased valence of intermolecular interactions in VF
components appeared to shift the EBOV VF morphology from the typical droplet-like to network-
like structures, which echoes the network-like VF we observed in EBOV-GFP-ΔVP30 infected
cells. Under such circumstances, we resolved multiple footprints of the EBOV polymerase
gathering at discrete sites in network-like EBOV VFs, where the VF phase boundary was
invaginated or interconnected. These sites may correlate to the EBOV L-foci under confocal light
microscopy.
Based on our confocal light and electron microscopy findings, we propose a model in which
EBOV polymerase molecules act in concert in a spatially restricted manner. Having a group of
EBOV polymerase molecules acting together on the same viral genome would compensate for
the low efficiency of viral RNA synthesis, as a productive viral RNA synthesis event can only
initiate at a single site at the 3' end of the viral genome[64]. Even so, during transcription, the EBOV
polymerase makes frequent stops at each intergenic region, and at each stop the polymerase
may detach from the genome template. How the same polymerase returns to the 3' end of the
viral genome and initiates a new round of RNA synthesis event is currently unclear. If a group of
EBOV polymerases act on the same viral genome, with some polymerases falling off the template
at certain points, other copies of the polymerase may eventually reach the distal end of the
genome where the L gene is located. By organizing the viral genome in tandem or in another
coordinated way, the same group of polymerases could process more copies of the viral genome
and maximize viral RNA production.
This study has several limitations. First, our FRAP analysis is based on the fluorescence
recovery of a sub-area within the VF condensate, which measures a combined rate of molecular
exchange that occur within the VF condensate and occur between the VF and the cytoplasm.
Because the fluorescence intensity of molecules in the VF condensate is ~200-folds higher than
that of the diffused molecules in the cytoplasm and the total VF condensate is significantly larger
than the photo-bleached spot, we reason that the combined rate we measured could approximate
the internal exchange rate within the VF condensate. Indeed, we observed a preferred internal
molecular exchange as in an immediate fluorescence-decay in internal regions within VF that
were not directly photobleached. Nevertheless, FRAP analysis by photobleaching of the full VF
condensate can measure the molecular exchange between the VF condensate and the rest of
the cytoplasm. This measurement is particularly relevant to understand several other stages of
the viral life cycle such as departure of condensed vRNP from VFs as part of viral egress. Second,
the intracellular localization of L we observed either by light or electron microscopy provides an
isolated "snapshot" of only the RNA synthesis stage of EBOV infection. However, EBOV infection
is a continuous process that involves expression of viral matrix proteins (VP24 and VP40) and
viral glycoprotein. The presence of VP24 will change the biological function as well as the physical
properties of EBOV VFs, as the loosely coiled vRNP will be condensed and inevitably reduce the
fluidity of EBOV VF. Meanwhile, viral polymerase will be immobilized in the condensed vRNP and
depart from VFs to the plasma membrane for budding. These virological events are not
recapitulated in the EBOV MG systems used here. Together, the results of our current study
increase our understanding of the mechanisms associated with EBOV VFs and the spatial
regulation of viral RNA synthesis within. Given the similar replication strategy shared among
nNSVs, our findings may be applicable to other critically important nNSV pathogens, such as
rabies, RSV and measles virus.
494
495   **Figures and figure legends**

Figure 1

496
497 **Figure 1. Reconstituted EBOV viral factories with minimal viral protein components display**
498 **viscoelastic behavior in live cells. a** Bioinformatic prediction of intrinsically disordered regions
499 and low complexity regions in the primary sequence of EBOV NP and V35 protein, using the

IUPred and PlaToLoCo webserver, respectively. **b** Coimmunoprecipitation (co-IP) of EBOV NP and L with HA-mNG-VP35. HA-VP35 and HA-mNG served as positive and negative control, respectively. Anti-HA antibody was used to detect HA-mNG-VP35, HA-VP35, and HA-mNG in the input cell lysate and in HA-IP fraction. Representative results from 3 biological replicates are shown. Confocal microscopy of **c** VP35 **d** VP35+NP **e** VP35+NP+L condensates inside live HEK 293T cells 1 d post-transfection. Representative cells (N>6 cells) for each condition from 4 biological replicates are shown. The cell body and nucleus (Nuc.) are marked by a dashed line. White arrow: individual condensate chosen for photobleaching. Image montage is composed of selected frames (including t = 0 s) with an interval of 2.88 s from each time-lapse of photobleached condensate displayed. The diameter of photobleached regions is 1 µm. Photobleaching occurred at t = 4.8 s. Scale bars: 5 µm. Schematic of EBOV VP35-involved molecular associations corresponding to each type of condensate. **f** Fluorescence recovery of mNG-HA-VP35 within the photobleached region inside intracellular condensates containing VP35. Normalized intensity corresponding to each time point before and after photobleaching of VP35 condensate is shown in green, VP35+NP condensate is shown in red, VP35+NP+L condensate is shown in blue. Each data point represents the mean with standard deviation (error bars) of N=9/9/6 (VP35/VP35+NP/VP35+NP+L condensates). Data points with t > 5s in red and in blue were used to fit a corresponding two-phase association curve, with the normalized intensity value in expressed as a percentage and the curve plateau marked on the side. Goodness of fit of each curve is indicated by an R^2 value. An extra sum-of-squares F test was performed to determine whether the best-fit value for unshared parameters differ between the blue and red curves. For each fitted curve, the best-fit value for each kinetic parameter is shown with a 95% confidence interval.

Figure 2

a NP+VP35+FLAG-L

b
- Rep-comp. MG
  EBOV 3' le — GFP — Rluc — EBOV 5' tr
- Rep-def. MG
  GFP — Rluc — 55nt deletion

○ T7 pol   ○ NP   ○ VP30
● T7-2cisMG   ○ VP35   ○ FLAG-L

T7-based EBOV 2cis-MG system

HEK293T

Reconstitution of L-mediated EBOV MG activities

	Rep-comp. MG	Rep-def. MG	
cRNA	↑ ↓	↑ ✗	Replication
vRNA	↑ ↓	↓	Transcription
mRNAs	↑ ↑	↑ ↑	

c Rep-comp. MG   Rep-def. MG
% EBOV minigenome activity
** p=0.0024   **** p<0.0001
no L, L-WT, FLAG-L (for each)

anti-FLAG (250 kD)
anti-L (250 kD)
β-actin (50 kD, 37 kD)

d Rep-comp. MG — FLAG-L | NP | MG reporter | Merge
Z-stacks
Scale bars: 5 μm

e Rep-def. MG — FLAG-L | NP | MG reporter | Merge
Z-stacks
Scale bars: 5 μm

f L-foci count vs Distance to the nearest neighbour (μm)
■ Rep-comp. MG   ■ Rep-def. MG

524
Figure 2. FLAG-tagged EBOV L cluster into foci within reconstituted viral factories. a
Confocal immunofluorescence microscopy of fixed HEK 293T cells co-expressing EBOV NP,
VP35 and FLAG-L at 1 d post-transfection. Represented Z-stacks in orthogonal view of 2
biological replicates each include > 4 fields of view. Scale bars: 5 μm. **b** Schematics of the T7
polymerase (T7 pol), bicistronic EBOV minigenome (2cis-MG) system with a replication
competent (Rep-comp.) or replication deficient (Rep-def.) MG. 3' le: 3' leader; 5' tr: 5' trailer; black
triangle: gene start; black bar: gene end. **c** Activity of FLAG-L relative to the L-WT control
measured in supporting the expression of *Renilla* luciferase (Rluc) in EBOV Rep-comp. vs. Rep-
def. MG. Background expression of MG assessed by excluding L in the MG system (no L). Results
from 3 biological replicates with technical triplicates are shown as individual data points with mean
± SD (error bars). **, *p*=0.0024, ****, *p*<0.0001 (N=9, two-tailed, unpaired t tests with Welch's
correction). Expression of FLAG-L compared to L-WT and detection of the FLAG tag analyzed by
western blot using a mouse monoclonal ani-FLAG and a rabbit polyclonal anti-EBOV L antibody,
respectively. Loading control: β-actin. Confocal microscopy of fixed HEK 293T cells transfected

with **d** Rep-comp. or **e** Rep-def. EBOV MG system at 2 d post-transfection. A representative
confocal image overview is shown, in which selected EBOV VFs are marked with a white box
(scale bars: 5 μm) and magnified in confocal z-stacks (scale bars: 2 μm). Fluorescence of the
GFP reporter in both Rep-comp. and Rep-def. EBOV MG is pseudo-colored in yellow for display
purposes. Nuclei are counterstained with Hoechst. Representative results from 4 biological
replicates with > 5 fields of view are shown. **f** A histogram showing the distribution of nearest
distances between EBOV L-foci within the same VF in the presence of Rep-comp. vs. Rep-def
MG, corresponding to z-stacks shown in (**d**) and (**e**). A total of 908 (Rep-comp.) or 683 (Rep-def.)
L-foci used in quantification.
548

Figure 3

a

b

Figure 3. Morphologically distinct EBOV viral factories in EBOV minigenome transfected cells and EBOV-GFP-ΔVP30 infected Vero-VP30 cells. a Confocal immunofluorescence microscopy of HEK 293T cells transfected with T7-pol based, Rep-comp. EBOV MG system, at 2 d post transfection. L-WT was used. A representative result from 2 biological replicates with 3 fields of view is shown. **b** Confocal immunofluorescence microscopy of representative Vero-VP30 cells infected with EBOV-GFP-ΔVP30 at MOI (multiplicity of infection) = 3, 18 h post infection. A total of N=39 cells from 2 biological replicates were analyzed. In both (**a**) and (**b**), fluorescence signals of the GFP reporter in Rep-comp. MG and in EBOV-GFP-ΔVP30 is pseudo-colored yellow for display purposes. EBOV NP was labeled with a human monoclonal anti-NP antibody paired

560   with Alexa-568 anti-human antibody. The morphologically distinct EBOV viral factory is magnified
561   in a single channel view. Nucleus counterstained with Hoechst. Scale bars: 5 µm.
562

Figure 4

Figure 4. Engineering and characterization of a split-APEX2 tagged EBOV polymerase
complex. a Protein constructs of the split-APEX2 (sAPEX2)-tagged EBOV L and VP35. sAPEX
contains two fragments: sAP and sEX. **b** Activity of L-sEX and VP35-V5-sEX (L-VP35-sAPEX2)
relative to wild-type EBOV L (L-WT) and V5-tagged VP35 (VP35-V5) control measured in
supporting expression of firefly luciferase (FLUC) in the EBOV MG. Background expression of
MG assessed by excluding L in the MG system (no L). Results from 2 biological replicates with
technical triplicates are shown as individual data points with mean ± SD (error bars). ****,
$p<0.0001$ (N=6, two-tailed, unpaired t tests with Welch's correction). **c** Schematic of the RNA
polymerase 1(Pol1)-based, monocistronic EBOV minigenome system (Pol1-EBOV MG). 3' le: 3'
leader; 5' tr: 5' trailer; black triangle: gene start; black bar: gene end. **d** Expression of L-sEX
compared to L-WT and VP35-V5-sAP compared to VP35-V5, each analyzed by western blot
using a rabbit polyclonal anti-EBOV L antibody and a mouse monoclonal anti-V5 antibody,
respectively. Loading control: β-actin. **e** Confocal immunofluorescence microscopy of HEK 293T
cells transfected with the Pol1-EBOV MG system containing either L-WT +VP35-V5 or L-VP35-
sAPEX2, at 2 d post transfection. **f** Confocal immunofluorescence microscopy of HEK 293T cells
transfected with the Pol1-EBOV MG system containing either L-WT +VP35-V5 or L-VP35-
sAPEX2, and in the presence of VP24, at 2 d post transfection. In (**e**) and (**f**), both VP35-V5-sAP
and VP35-V5 were labeled with a rabbit monoclonal anti-V5 antibody paired with Alexa-488 anti-

rabbit antibody. Functional reconstitution of APEX2 was indicated by fluorescence signals from
resorufin, the product of the Amplex UltraRed, a fluorogenic peroxidase substrate for APEX2. In
(**f**), EBOV VP24 was labeled with a mouse monoclonal anti-VP24 antibody paired with Alexa-488
anti-mouse antibody. Nuclei were counterstained with Hoechst. Fluorescence signals of VP35
and VP24 are pseudo-colored green and red for display purposes. Scale bars: 5 µm.
Representative results from 2 biological replicates with >5 fields of view are shown. The valence
of inter-molecular interaction is depicted as double-ended arrows.
590

591
592

Figure 5

593
Figure 5 Nanoscale localization of split-APEX2 tagged EBOV polymerase complex revealed
by thin-section electron microscopy (EM) a Schematic of the functional reconstitution of
APEX2 upon formation of EBOV polymerase (L-VP35) complex and the resulting EM contrast
upon DAB-OsO$_4$ staining. **b** HEK 293T cells were transfected with the Pol1-EBOV-MG system
containing sAPEX2-tagged EBOV polymerase (L-VP35) complex and were chemically fixed with
2.5% glutaraldehyde and stained with DAB at 2 d post-transfection. DAB+ and DAB- cells were
examined by transmitted light microscopy (left) and then embedded in resin and stained with
OsO$_4$ (middle). The samples were serially sliced into 70 nm-thick sections and imaged with
transmission electron microscopy (TEM) at low magnification (right). Scale bars: 20 µm. **c**
Electron micrographs of either the overview or a cytoplasmic region of the DAB+ cell indicated in

(**b**) at different magnifications. Arrows: sAPEX2 mediated deposition of electron-dense Osmium.

d Electron micrographs of either the overview or a cytoplasmic region of the DAB- cell indicated

in (**b**) at different magnifications. Scale bars: 5 µm/500 nm/200 nm in micrographs with a

low/medium/high magnification. **e** Cartoons annotating the whole cell overview, the electron-

dense viral factories (VF) and the sAPEX2-tagged EBOV polymerases (sAPEX2-pol) in both the

magnified DAB+ and DAB- cell, shown in (**c**) and (**d**), respectively. Representative results from 3

biological replicates with multiple fields of view are shown. Cyto.: cytoplasm. Nuc.: nucleus.

Figure 6

612
Figure 6. 3D visualization of EBOV viral factories (VFs) and sAPEX2-tagged polymerase by
613 **electron tomography. a** HEK 293 cell transfected with the Pol1-EBOV-MG system containing
614 sAPEX2-tagged EBOV polymerase (L-VP35) complex before fixation with 2.5% glutaraldehyde
615

at 2 d post-transfection. The samples were stained with DAB-OsO$_4$, embedded in resin and
sectioned into 250 nm-thick specimens. A reconstructed 3D electron tomogram of the specimen
originating from a DAB+ cell is shown in an orthogonal view with tomographic slices. Scale bar:
500 nm. **B** Segmented volume of the electron-dense EBOV VFs and surface representation of
numerous cellular organelles and sAPEX2-tagged EBOV polymerase, corresponding to the 3D
electron tomogram in (**a**). **c** Orthogonal view (with X-Y/Y-Z/X-Z plane) showing a magnified
volume of interest in the 3D electron tomogram in (a). Scale bar: 100 nm. **d** Segmented surface
representations of EBOV VFs (orange), sAPEX2-tagged EBOV polymerase (pink), viral RNP coils
(yellow), host ribosomes (purple), and single-membrane vesicle (blue), resulting from the
tomographic volume in (**c**). **e** Series of magnified tomographic slices on the X-Y plane from a
region containing multiple helical structures that correspond to viral RNPs. Yellow arrows pointing
toward one helical structure; white arrows pointing toward the branch site where multiple helical
structures meet. A cartoon annotation of the same region is shown on the side. **f** A series of
magnified tomographic slices on the X-Y plane from a region containing DAB-osmium deposits
that correspond to sAPEX-tagged EBOV polymerases (indicated by pink arrows) and a single-
membrane vesicle approaching the electron-dense VF. A cartoon annotation of the same region
is shown on the side. For (**e**) an (**f**), the scale bar in the tomographic overview slice and magnified
series represents 500 nm and 100 nm, respectively. Tomographic slice number is marked in the
lower left.
635

Methods:

Plasmids: If not specified, NEB HiFi assembly was used for incorporation of DNA fragments into
plasmid vector. To generate pCAGGS-mNG-HA-VP35, coding sequence of mNeonGreen (mNG)
was PCR amplified from pmNeonGreenHO-3xFLAG-G (addgene#127914), while an N-terminal
HA epitope tag was added to the mNG coding sequence. The resulting fragment was inserted
upstream of the VP35 coding sequence in pCAGGS-VP35. To generate pCAGGS-HA-VP35
control plasmid, a DNA fragment containing a HA epitope tag followed by a flexible linker was
synthesized and added to the N-terminus of VP35 coding frame. To generate pCMV-mNG-HA
control plasmid, a HA epitope tag was added to the C-terminus of mNG coding sequence. To
generate pCAGGS-VP35-V5-sAP, the fragment of sAP was PCR amplified from
FKBP_V5_AP_NEX_pLX304 (addgene#120912) and cloned into the intermediate vector
pcDNA5-VP35. The fragment of VP35-V5-sAP was subcloned to pCAGGS plasmid using
restriction sites Xho1 and Nhe1 and standard ligation method. To generate pCAGGS-VP35-V5
control plasmid, the fragment of VP35-V5 was PCR amplified with an addition of a stop codon
and Bgl2 restriction site, and was then incorporated into pCAGGS backbone using Kpn1 and Bgl2
restriction sites.
To generate pCAGGS-FLAG-L, a DNA fragment containing a 2xFLAG epitope tag followed
by a flexible linker was synthesized and added to the N-terminus of L coding frame. To generate
pCEZ-L-sEX, a fragment of L coding sequence was first subcloned into a pFastbacDual
intermediate plasmid using the natural restriction sites Pac1 and Hpa1. From there, coding
sequence of sEX was PCR amplified from HA-Halotag-FRB-EX-NES-pLX304 (addgene
#120913), flanked by two flexible linkers and was internally inserted to L fragment at the position
1705/1706 (TTIP/Q). The L-sEX fragment was shuffled back to the pCEZ-L backbone using the
same restriction sites Pac1 and Hpa1.
To generate pT7-2cis-MG-Replication competent (Rep-Comp.), the coding sequence of VP40,
GP and VP24, including non-coding sequences VP40/GP and VP30/VP24 in between were
removed from pT7-4cis-EBOV-vRNA-eGFP plasmid[65], while the coding sequence of *Renilla*
luciferase was inserted downstream of the non-coding sequence NP/VP35 and reassembled. To
generate pT7-2cis-MG-Replication deficient (Rep-def.), the last 55 nucleotide of the 5' trailer
sequence in pT7-2cis-MG-Rep-comp. plasmid was removed by PCR and re-assembled. Other
supporting plasmids used in T7-based EBOV minigenome system were obtained from Dr.
Thomas Hoenen (Friedrich-Loeffler-Institute). Plasmids used in Pol1-based EBOV minigenome
system were obtained from Dr. Yoshihiro Kawaoka (University of Wisconsin Madison).

Cell lines and virus: Human embryonic kidney cells HEK 293T (ATCC reference: CRL-3216)
were originally purchased from the ATCC organization. Vero cells stabling expressing Ebola virus
VP30 protein (Vero-VP30) were obtained from Dr. Yoshihiro Kawaoka (University of Wisconsin
Madison). Both cell lines were maintained in Dulbecco's modified Eagle medium (DMEM-
GlutaMAX) supplemented with 4.5 g/L D-Glucose, 10% fetal bovine serum (FBS), penicillin
(100 U/mL), streptomycin (100 µg/mL). In addition, puromycin (5 µg/mL) was included in the
culture media while maintaining Vero-VP30 cells but excluded for infection experiment. The
biologically contained Ebola virus, EBOV-ΔVP30-GFP was generated in HEK 293T cells and
grown in Vero-VP30 cells as previously described[44].

Fluorescence Recovery After Photobleaching (FRAP): IBIDI µ-slides (8 wells high-precision
glass bottom) were treated with human fibronectin (50 mg/mL) for 20 minutes at 37°C incubator,
prior to HEK 293T cells seeding (4×10^4 cells/well). Twenty-four hours later, the monolayer was
transfected with plasmids in three different combinations: 1) 125 ng of pCAGGS-mNG-HA-VP35
and 1 µg of pCAGGS-vector control; 2) 125 ng of pCAGGS-mNG-HA-VP35, 125 ng of pCAGGS-
NP, and 875 ng of pCAGGS-vector control; 3) 125 ng of pCAGGS-mNG-HA-VP35, 125 ng of

pCAGGS-NP, and 875 ng of pCAGGS-L (wild-type). Transfected cells were incubated with
complete FluoroBrite DMEM (+1x Glutamax, 10% FBS) for imaging. Time series were acquired
using ZEISS laser scanning microscope (LSM880) with an Airyscan detector while live cells
sample were kept in caged incubator maintaining 37˚C and 5% CO_2. mNG fluorescence was
excited with a 488 nm argon laser and detected with a 63x 1.4NA oil immersion objective (Plan-
Apochromat), with a combination of bandpass filters (420-480) and (495-550). Using a frame rate
of 240 millisecond per frame and 0.4% of 488 nm laser, 20 frames were recorded pre-bleach, and
then 330 frames were recorded post-bleach under the Airyscan fast mode. This duration of post-
bleach image acquisition was experimentally determined by the time when fluorescence recovery
reached a plateau. For photobleaching, 488 nm laser at 80% was used to pulse-bleach a spot
size of 1 micron in diameter in the center of large condensates, until the fluorescence intensity of
the bleach spot reduced to 30% of the prebleached value.
Time series in which the bleached object moved during post-bleach phase were manually
identified and removed from data analysis because the fluorescence intensity of the bleached
spot could not be accurately measured in ImageJ/FIJI[66]. For each bleach event, double
normalization of the FRAP curve was performed using ImageJ/FIJI as described[15]. To correct
unintentional photobleaching due to time-series acquisition, in every frame, fluorescence intensity
of the bleached spot (1 μm in diameter) was first normalized to the mean intensity of an
unbleached spot (1 μm in diameter) of another condensate in the same field of view. The resulting
values after correction for all frames were further normalized to the average intensity of the last
10 pre-bleach frames. The background intensity in each field of view was negligible as it was
lower than 0.5% of the average intensity of viral factory condensates. Due to biological mobility
of the intracellular condensate and the cellular specimens during imaging, we could not accurately
measure the bleach pulses-induced fluorescence loss in the whole condensate. Instead, we
assessed this parameter in the averaged fluorescence decay of internal reference-regions inside
the same condensate that was photobleached but are away from the bleached spot. Double
normalized data from 6-9 cells in a total of three independent experiments were pooled. Means
with standard deviation of double normalized intensity were plotted as a function of time, of which
post-bleach data-points were fitted to a two-phase association model in GraphPad prism,
according to previously described methods[67]. Best-fit values with 95% confidence interval and
goodness of fit indicated by R^2 for each data set were reported.

$$715 \quad A_{fast} = (Plateau - Y_0) \times \%Fast$$
$$716 \quad A_{slow} = (Plateau - Y_0) \times (1 - \%Fast)$$
$$717 \quad Y(t) = Y_0 + A_{fast}\left(1 - e^{-k_{fast}t}\right) + A_{slow}(1 - e^{-k_{slow}t})$$

Immunofluorescence staining and confocal light microscopy: Fixation and staining was
performed as described[68].
For samples shown in supplementary Figure S1b, human monoclonal anti-Ebola virus NP
(1:2000), mouse monoclonal anti-Ebola virus VP35 (1:500), and rabbit polyclonal anti-Ebola virus
L antibody (1:200) were used as the primary antibodies, goat anti-human Alexa-568 (1:1000),
goat anti-mouse Alexa-647 (1:500), and goat anti-rabbit Alexa-568 (1:500) were used as the
secondary antibodies.
For samples shown in Figure 2, human monoclonal anti-Ebola virus NP (1:2000) and mouse
monoclonal anti-FLAG (1:100) antibody were used as the primary antibodies, goat anti-human
Alexa-568 (1:1000) and goat anti-mouse Alexa-647 (1:500) were used as the secondary
antibodies.
For samples shown in Figure 3, human monoclonal anti-NP (1:2000) and goat anti-human
Alexa-568 (1:1000) were used.
For samples shown in Figure 4, rabbit monoclonal anti-V5 (1:500) and mouse monoclonal
anti-Ebola VP24 (1:1000, a kind gift from Dr. Yoshihiro Kawaoka) were used as the primary

antibodies, goat anti-human Alexa-568 (1:1000) and goat anti-mouse Alexa-647 (1:500) were
used as the secondary antibodies. In addition, trans-complemented APEX2 was stained with
Amplex UltraRed at 100 nM (with 0.02% H_2O_2 in DPBS) on ice for 20 minutes prior to fixation.
Unreacted Amplex UltraRed was washed off with DPBS.
Single-plane confocal images or confocal Z-stacks (interval= 0.224 μm) were acquired with
the ZEISS laser scanning microscope (LSM880)-Airyscan system under the superresolution
mode, using a plan-apochromat 20x/0.8 numerical aperture M27 objective or an alpha plan-
apochromat 63×/1.46 numerical aperture oil Korr M27 objective. All images or stacks that are
shown or quantified were Airyscan processed using the Zen black (ZEISS) build-in function.
For supplementary Figure 1C, size quantification and counting of reconstituted Ebola viral
factories (VFs) in single-plane confocal images using ImageJ/Fiji. HEK 293T cells were prepared
using identical transfection conditions described in FRAP experiment. VFs were identified on the
basis of mNG-fluorescence and analyzed for the count and size distribution using the analyze
particle function.
For Figure 2f, 3D segmentation of Ebola viral factories in confocal Z-stacks was performed in
Imaris 9.9.1 using the surface function and automatically thresholding the GFP intensity;
segmentation of FLAG-tagged Ebola virus L-foci inside each viral factory was performed using
the spot function and automatically thresholding the fluorescence intensity of FLAG-L with an
estimated diameter for each spot as 0.5 μm, which generates specific values of distance to
nearest neighbor for every random pair of segmented spot.

Virus infection with EBOV-GFP-ΔVP30: Vero-VP30 cells ($4X10^4$ cells/well) were seeded in
IBIDI μ-slides (8 wells high-precision glass bottom). Twenty-four hours later, the monolayer was
incubated with EBOV-ΔVP30-GFP (MOI=3 foci-forming unit/cell) on ice. After 1 hour, the
monolayer was washed three times with cold DPBS to remove unbound virions and was moved
to the 37°C incubator. Infection was terminated after 16 hours. Infected cells were inactivated with
4% PFA for 15 minutes.

Thin-section TEM with DAB staining: Sample preparation was adapted from[69]. Mattek 35 mm
dish with NO#1.5 gridded glass bottom (P35G-1.5-14-C-GRD) was treated with human fibronectin
(50 mg/mL) for 1 hour at 37°C incubator, prior to HEK 293T cells seeding ($1X10^5$ cells/well).
Twenty-four hours later, the monolayer was either untreated or transfected with plasmids in two
different combinations: 1) EBOV Pol1-MG system containing VP35-V5 and L-WT; 2) EBOV Pol1-
MG system containing sAPEX2 tagged L-VP35.
For the first combination, 250 ng of pCEZ-NP, 187.5 ng of pCEZ-VP30, 250 ng of pCEZ-
VP35-V5, 156.25 ng of pHH21-3E5E-fluc, and 1875 ng of pCEZ-L were used in the co-
transfection mix for each 35 mm dish. For the second combination, 250 ng of pCEZ-NP, 187.5 ng
of pCEZ-VP30, 250 ng of pCEZ-VP35-V5-sAP, 156.25 ng of pHH21-3E5E-fluc, and 1875 ng of
pCEZ-L-sEX were used in the co-transfection mix for each 35 mm dish.
Forty-eight hours post-transfection, cells were fixed first in 2.5% glutaraldehyde-0.1 M
cacodylate buffer pH 7.4 + 2 mM $CaCl_2$ at room temperature for 1 minute, then fixed in pre-chilled
2.5% glutaraldehyde-0.1 M cacodylate buffer pH 7.4 + 2 mM $CaCl_2$ on ice for 1 hour, and were
washed three times with cold 0.1 M cacodylate buffer pH 7.4 (cacodylate buffer). Unreacted
fixative in samples was quenched with cold 20 mM Glycine solution for 5 min on ice, followed by
three washes with cold cacodylate buffer. Samples were stained with 2.5 mM DAB (Sigma
#D8001)-0.1 M cacodylate solution in the presence of 1/1000V of 30% H_2O_2 for 45 minutes on
ice and washed three times with cold cacodylate buffer.
Brown DAB stain in samples was confirmed under light microscopy. Samples were then
stained in 1% osmium+1.5% potassium Ferrocyanide in 0.1 M cacodylate buffer on ice for 1 hour
and washed three times with cacodylate buffer. Samples were dehydrated with increasing
concentrations of ethanol (20%, 50%, 70%, 90%, 100%, 100%, 3 minutes each) and washed

once in room temperature anhydrous ethanol. Samples were then infiltrated with Durcupan ACM
resin (Electron Microscopy Sciences) using a mixture of anhydrous ethanol: resin (1V:1V) for 30
minutes, then with 100% resin overnight, followed by 48 hours polymerization step at 60 °C.
Ultrathin sections with a 70 mm thickness of embedded specimen were mounted on 200 mesh
hexagonal copper grids with no post-staining step. Electron micrographs were collected using a
FEI Tecnai Spirit transmission electron microscope operating at 120 kV.

Electron tomography and data processing: Electron tomography was performed using a
300kV Titan Halo equipped with an 8k x 8k direct detector (DE64, Direct Electron). During the
procedure, semi-thick (~250 nm) sections of the resin-embedded specimen were imaged at
different orientations using a 4-tilt acquisition scheme as described[70], for which the specimen was
tilted from -60 to +60 degrees every 0.25 degree at four different azimuthal orientation. For
aligning the micrographs, 5 nm colloidal gold particles were deposited on each side of the sections
to serve as fiducial markers. The TEM magnification was set at 11,000x, corresponding to a raw
pixel size of 0.36 nm. Tomograms were generated with an iterative reconstruction procedure[70],
and was binned by a factor of 4 for display and for volume segmentation using Amira 2020 3.1.
software. Membrane organelles were segmented manually combined with automatic thresholding.
Ribosomes were segmented manually. EBOV viral factories and sAPEX2-tagged EBOV
polymerase were segmented with a combination of automatic thresholding and the TopHat tool.

EBOV minigenome assays: Transfection and activity quantification with Pol1-based
monocistronic EBOV minigenome system was adapted from[47] and described previously[68].
Transfection with the T7 pol-based EBOV bicistronic minigenome system was adapted from[65].
Activity of bicistronic minigenome system was measured in *Renilla* luciferase assay. For each
transfection condition, the same cell lysates used in luciferase assay, were pooled from triplicated
assay wells and analyzed in western blot.

Coimmunoprecipitation (coIP): Co-IP reactions were performed as previously described[68]. HEK
293T cells ($1X10^6$ cells/well) were seeded in 6-wells plates. Twenty-four hours later, cells in each
well were transfected with 500 ng of pCMV-mNG-HA or pCAGGS-HA-VP35 or pCAGGS-mNG-
HA-VP35 plasmid combined with 500 ng of pCAGGS-NP, with and without 1 μg of pCAGGS-L,
using TransIT-LT1 transfection reagent. HA-affinity matrix was used to pull-down protein
complexes containing HA-tagged proteins.

Statistical analysis: Data shown in Figure 2c, 4a and supplementary Figure 1a were calculated
and plotted in mean with SD. Statistical analysis were analyzed by two-tailed unpaired t-test with
Welch's correction. P values were indicated in figure legends. Data shown in Figure 1f were
analyzed using an extra sum-of-squares F test to compare whether the best-fit values of *%Fast*,
k_{fast} and k_{slow} differ between two datasets. The resulting *p* value indicating the statistical
significance of the difference was reported in the figure.

Acknowledgments: We thank Dr. Yoshihiro Kawaoka (University of Wisconsin) for providing the
Pol1-based Ebola minigenome system, sharing the anti-VP30 and anti-VP24 antibodies and the
EBOV-GFP-ΔVP30 system. We thank Dr. Reika Watanabe (LJI) for providing the EBOV-GFP-
ΔVP30 virus stock. We thank Dr. Thomas Hoenen (Friedrich-Loeffler-Institute) for sharing the
Ebola tetracistronic trVLP system. We thank Diptiben Parekh (LJI) for plasmid preparations, Dr.
Sharon Schendel (LJI) for manuscript editing, NIH S10OD021831 for sponsoring the Zeiss LSM
880 microscope at the LJI microscopy core facility and an Imaging Scientist grant (2019‐198153)
from the Chan Zuckerberg Initiative to S.M. (LJI). We thank NIH R24GM137200 and
S10OD021784 to M.H.E. (UCSD) in support of the National Center for Microscopy and Imaging

Research. This research was supported by institutional funds of La Jolla Institute for Immunology
to E.O.S. (LJI). J.F. was supported by the Donald E. and Delia B. Baxter Foundation Fellowship.

Author contributions: J.F. conceived the study and wrote the paper under the supervision of
E.O.S.; J.F., G.C. and S.M. designed the experiments, J.F., G.C., S.P., S.M., and C.H. performed
the experiments; J.F., G.C., analyzed the data, M.H.E. supervised the data acquisition and
interpretation for electron microscopy analysis, A.A.D. contributed to data interpretation of
intracellular condensates. All authors edited and approved the paper.

References

References

1   Alenquer, M. *et al.* Influenza A virus ribonucleoproteins form liquid organelles at
endoplasmic reticulum exit sites. *Nat Commun* **10**, 1629, doi:10.1038/s41467-019-09549-
4 (2019).

2   Cifuentes-Munoz, N., Branttie, J., Slaughter, K. B. & Dutch, R. E. Human Metapneumovirus
Induces Formation of Inclusion Bodies for Efficient Genome Replication and Transcription.
J Virol **91**, doi:10.1128/JVI.01282-17 (2017).

3   Dolnik, O., Stevermann, L., Kolesnikova, L. & Becker, S. Marburg virus inclusions: A virus-
induced microcompartment and interface to multivesicular bodies and the late
endosomal compartment. *Eur J Cell Biol* **94**, 323-331, doi:10.1016/j.ejcb.2015.05.006
(2015).

4   Geiger, F. *et al.* Liquid-liquid phase separation underpins the formation of replication
factories in rotaviruses. *EMBO J* **40**, e107711, doi:10.15252/embj.2021107711 (2021).

5   Heinrich, B. S., Cureton, D. K., Rahmeh, A. A. & Whelan, S. P. Protein expression redirects
vesicular stomatitis virus RNA synthesis to cytoplasmic inclusions. *PLoS Pathog* **6**,
e1000958, doi:10.1371/journal.ppat.1000958 (2010).

6   Hoenen, T. *et al.* Inclusion bodies are a site of ebolavirus replication. *J Virol* **86**, 11779-
11788, doi:10.1128/JVI.01525-12 (2012).

7   Jobe, F., Simpson, J., Hawes, P., Guzman, E. & Bailey, D. Respiratory Syncytial Virus
Sequesters NF-kappaB Subunit p65 to Cytoplasmic Inclusion Bodies To Inhibit Innate
Immune Signaling. *J Virol* **94**, doi:10.1128/JVI.01380-20 (2020).

8   Lahaye, X. *et al.* Functional characterization of Negri bodies (NBs) in rabies virus-infected
cells: Evidence that NBs are sites of viral transcription and replication. *J Virol* **83**, 7948-
7958, doi:10.1128/JVI.00554-09 (2009).

9   Lee, C. H. *et al.* Reovirus Nonstructural Protein sigmaNS Recruits Viral RNA to Replication
Organelles. *mBio* **12**, e0140821, doi:10.1128/mBio.01408-21 (2021).

10   Ringel, M. *et al.* Nipah virus induces two inclusion body populations: Identification of
novel inclusions at the plasma membrane. *PLoS Pathog* **15**, e1007733,
doi:10.1371/journal.ppat.1007733 (2019).

11   Zhou, Y., Su, J. M., Samuel, C. E. & Ma, D. Measles Virus Forms Inclusion Bodies with
Properties of Liquid Organelles. *J Virol* **93**, doi:10.1128/JVI.00948-19 (2019).

12   Feric, M. *et al.* Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* **165**,
1686-1697, doi:10.1016/j.cell.2016.04.047 (2016).

13   Whelan, S. P., Barr, J. N. & Wertz, G. W. Transcription and replication of nonsegmented
negative-strand RNA viruses. *Curr Top Microbiol Immunol* **283**, 61-119, doi:10.1007/978-
3-662-06099-5_3 (2004).

14   Heinrich, B. S., Maliga, Z., Stein, D. A., Hyman, A. A. & Whelan, S. P. J. Phase Transitions
Drive the Formation of Vesicular Stomatitis Virus Replication Compartments. *mBio* **9**,
doi:10.1128/mBio.02290-17 (2018).

15   Nikolic, J. *et al.* Negri bodies are viral factories with properties of liquid organelles. *Nat*
Commun **8**, 58, doi:10.1038/s41467-017-00102-9 (2017).

16 Guseva, S. *et al.* Measles virus nucleo- and phosphoproteins form liquid-like phase-
separated compartments that promote nucleocapsid assembly. *Sci Adv* **6**, eaaz7095,
doi:10.1126/sciadv.aaz7095 (2020).
17 Boggs, K. B. *et al.* Human Metapneumovirus Phosphoprotein Independently Drives Phase
Separation and Recruits Nucleoprotein to Liquid-Like Bodies. *mBio* **13**, e0109922,
doi:10.1128/mbio.01099-22 (2022).
18 Risso-Ballester, J. *et al.* A condensate-hardening drug blocks RSV replication in vivo.
Nature **595**, 596-599, doi:10.1038/s41586-021-03703-z (2021).
19 Alberti, S., Gladfelter, A. & Mittag, T. Considerations and Challenges in Studying Liquid-
Liquid Phase Separation and Biomolecular Condensates. *Cell* **176**, 419-434,
doi:10.1016/j.cell.2018.12.035 (2019).
20 Banani, S. F., Lee, H. O., Hyman, A. A. & Rosen, M. K. Biomolecular condensates: organizers
of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298, doi:10.1038/nrm.2017.7
(2017).
21 Brangwynne, Clifford P., Tompa, P. & Pappu, Rohit V. Polymer physics of intracellular
phase transitions. *Nature Physics* **11**, 899-904, doi:10.1038/nphys3532 (2015).
22 Ruff, K. M. *et al.* Sequence grammar underlying the unfolding and phase separation of
globular proteins. *Mol Cell* **82**, 3193-3208 e3198, doi:10.1016/j.molcel.2022.06.024
(2022).
23 Boeynaems, S. *et al.* Spontaneous driving forces give rise to protein-RNA condensates
with coexisting phases and complex material properties. *Proc Natl Acad Sci U S A* **116**,
7889-7898, doi:10.1073/pnas.1821038116 (2019).
24 Harmon, T. S., Holehouse, A. S., Rosen, M. K. & Pappu, R. V. Intrinsically disordered linkers
determine the interplay between phase separation and gelation in multivalent proteins.
Elife **6**, doi:10.7554/eLife.30294 (2017).
25 Cahn, J. W. & Hilliard, J. E. Free Energy of a Nonuniform System. I. Interfacial Free Energy.
The Journal of Chemical Physics **28**, 258-267, doi:10.1063/1.1744102 (1958).
26 Cahn, J. W. & Hilliard, J. E. Free Energy of a Nonuniform System. III. Nucleation in a Two-
Component Incompressible Fluid. *The Journal of Chemical Physics* **31**, 688-699,
doi:10.1063/1.1730447 (1959).
27 Hajime, T. Viscoelastic phase separation. *Journal of Physics: Condensed Matter* **12**, R207,
doi:10.1088/0953-8984/12/15/201 (2000).
28 *Ebola virus disease*, <https://www.who.int/news-room/fact-sheets/detail/ebola-virus-
disease> (2022).
29 *History of Ebola Virus Disease (EVD) Outbreaks*,
<https://www.cdc.gov/vhf/ebola/history/chronology.html> (2022).
30 Kirchdoerfer, R. N., Abelson, D. M., Li, S., Wood, M. R. & Saphire, E. O. Assembly of the
Ebola Virus Nucleoprotein from a Chaperoned VP35 Complex. *Cell Rep* **12**, 140-149,
doi:10.1016/j.celrep.2015.06.003 (2015).
31 Becker, S., Rinne, C., Hofsass, U., Klenk, H. D. & Muhlberger, E. Interactions of Marburg
virus nucleocapsid proteins. *Virology* **249**, 406-417, doi:10.1006/viro.1998.9328 (1998).
32 Noda, T., Hagiwara, K., Sagara, H. & Kawaoka, Y. Characterization of the Ebola virus
nucleoprotein-RNA complex. *J Gen Virol* **91**, 1478-1483, doi:10.1099/vir.0.019794-0
(2010).

33   Shu, T. *et al.* Ebola virus VP35 has novel NTPase and helicase-like activities. *Nucleic Acids*
Res **47**, 5837-5851, doi:10.1093/nar/gkz340 (2019).
34   Watanabe, S., Noda, T. & Kawaoka, Y. Functional mapping of the nucleoprotein of Ebola
virus. *J Virol* **80**, 3743-3751, doi:10.1128/JVI.80.8.3743-3751.2006 (2006).
35   Bruhn, J. F. *et al.* Crystal Structure of the Marburg Virus VP35 Oligomerization Domain. *J*
Virol **91**, doi:10.1128/JVI.01085-16 (2017).
36   Chanthamontri, C. K. *et al.* The Ebola Viral Protein 35 N-Terminus Is a Parallel Tetramer.
Biochemistry **58**, 657-664, doi:10.1021/acs.biochem.8b01154 (2019).
37   Zinzula, L. *et al.* Structures of Ebola and Reston Virus VP35 Oligomerization Domains and
Comparative Biophysical Characterization in All Ebolavirus Species. *Structure* **27**, 39-54
e36, doi:10.1016/j.str.2018.09.009 (2019).
38   Sprague, B. L. & McNally, J. G. FRAP analysis of binding: proper and fitting. *Trends Cell Biol*
15, 84-91, doi:10.1016/j.tcb.2004.12.001 (2005).
39   Trunschke, M. *et al.* The L-VP35 and L-L interaction domains reside in the amino terminus
of the Ebola virus L protein and are potential targets for antivirals. *Virology* **441**, 135-145,
doi:10.1016/j.virol.2013.03.013 (2013).
40   Nelson, E. V. *et al.* Ebola Virus Does Not Induce Stress Granule Formation during Infection
and Sequesters Stress Granule Proteins within Viral Inclusions. *J Virol* **90**, 7268-7284,
doi:10.1128/JVI.00459-16 (2016).
41   Zhu, L. *et al.* Ebola virus VP35 hijacks the PKA-CREB1 pathway for replication and
pathogenesis by AKIP1 association. *Nat Commun* **13**, 2256, doi:10.1038/s41467-022-
29948-4 (2022).
42   Muhlberger, E., Weik, M., Volchkov, V. E., Klenk, H. D. & Becker, S. Comparison of the
transcription and replication strategies of marburg virus and Ebola virus by using artificial
replication systems. *J Virol* **73**, 2333-2342, doi:10.1128/JVI.73.3.2333-2342.1999 (1999).
43   Hoenen, T., Jung, S., Herwig, A., Groseth, A. & Becker, S. Both matrix proteins of Ebola
virus contribute to the regulation of viral genome replication and transcription. *Virology*
403, 56-66, doi:10.1016/j.virol.2010.04.002 (2010).
44   Halfmann, P. *et al.* Generation of biologically contained Ebola viruses. *Proc Natl Acad Sci*
U S A **105**, 1129-1133, doi:10.1073/pnas.0708057105 (2008).
45   Nanbo, A., Watanabe, S., Halfmann, P. & Kawaoka, Y. The spatio-temporal distribution
dynamics of Ebola virus proteins and RNA in infected cells. *Sci Rep* **3**, 1206,
doi:10.1038/srep01206 (2013).
46   Han, Y. *et al.* Directed Evolution of Split APEX2 Peroxidase. *ACS Chem Biol* **14**, 619-635,
doi:10.1021/acschembio.8b00919 (2019).
47   Jasenosky, L. D., Neumann, G. & Kawaoka, Y. Minigenome-based reporter system suitable
for high-throughput screening of compounds able to inhibit Ebolavirus replication and/or
transcription. *Antimicrob Agents Chemother* **54**, 3007-3010, doi:10.1128/AAC.00138-10
(2010).
48   Schmelzer, J. W., Abyzov, A. S. & Moller, J. Nucleation versus spinodal decomposition in
phase formation processes in multicomponent solutions. *J Chem Phys* **121**, 6900-6917,
doi:10.1063/1.1786914 (2004).
49   Tanaka, H. Viscoelastic phase separation in biological cells. *Communications Physics* **5**, 167,
doi:10.1038/s42005-022-00947-7 (2022).

50 Banadyga, L. *et al.* Ebola virus VP24 interacts with NP to facilitate nucleocapsid assembly
and genome packaging. *Sci Rep* **7**, 7698, doi:10.1038/s41598-017-08167-8 (2017).
51 Wigglesworth, V. B. The use of osmium in the fixation and staining of tissues. *Proceedings*
of the Royal Society of London. Series B - Biological Sciences **147**, 185-199,
doi:10.1098/rspb.1957.0043 (1957).
52 Neidle, S. & Stuart, D. I. The crystal and molecular structure of an osmium bispyridine
adduct of thymine. *Biochim Biophys Acta* **418**, 226-231, doi:10.1016/0005-
2787(76)90072-1 (1976).
53 Zhang, J., Li, D., Zhang, J., Chen, D. & Murchie, A. I. Osmium tetroxide as a probe of RNA
structure. *RNA* **23**, 483-492, doi:10.1261/rna.057539.116 (2017).
54 Geisbert, T. W. & Jahrling, P. B. Differentiation of filoviruses by electron microscopy. *Virus*
Res **39**, 129-150, doi:10.1016/0168-1702(95)00080-1 (1995).
55 Beniac, D. R. *et al.* The organisation of Ebola virus reveals a capacity for extensive,
modular polyploidy. *PLoS One* **7**, e29608, doi:10.1371/journal.pone.0029608 (2012).
56 Bharat, T. A. *et al.* Structural dissection of Ebola virus and its assembly determinants using
cryo-electron tomography. *Proc Natl Acad Sci U S A* **109**, 4275-4280,
doi:10.1073/pnas.1120453109 (2012).
57 Watanabe, S., Noda, T., Halfmann, P., Jasenosky, L. & Kawaoka, Y. Ebola virus (EBOV) VP24
inhibits transcription and replication of the EBOV genome. *J Infect Dis* **196 Suppl 2**, S284-
290, doi:10.1086/520582 (2007).
58 Watt, A. *et al.* A novel life cycle modeling system for Ebola virus shows a genome length-
dependent role of VP24 in virus infectivity. *J Virol* **88**, 10511-10524,
doi:10.1128/JVI.01272-14 (2014).
59 Bodmer, B. S. & Hoenen, T. Assessment of Life Cycle Modeling Systems as Prediction Tools
for a Possible Attenuation of Recombinant Ebola Viruses. *Viruses* **14**,
doi:10.3390/v14051044 (2022).
60 Blanchard, E. L. *et al.* Polymerase-tagged respiratory syncytial virus reveals a dynamic
rearrangement of the ribonucleocapsid complex during infection. *PLoS Pathog* **16**,
e1008987, doi:10.1371/journal.ppat.1008987 (2020).
61 Nilsson-Payant, B. E. *et al.* Reduced Nucleoprotein Availability Impairs Negative-Sense
RNA Virus Replication and Promotes Host Recognition. *J Virol* **95**, doi:10.1128/JVI.02274-
20 (2021).
62 Hu, Z. *et al.* Inclusion bodies of human parainfluenza virus type 3 inhibit antiviral stress
granule formation by shielding viral RNAs. *PLoS Pathog* **14**, e1006948,
doi:10.1371/journal.ppat.1006948 (2018).
63 Ma, W., Zheng, G., Xie, W. & Mayr, C. In vivo reconstitution finds multivalent RNA–RNA
interactions as drivers of mesh-like condensates. *eLife* **10**, e64252,
doi:10.7554/eLife.64252 (2021).
64 Deflube, L. R. *et al.* Ebolavirus polymerase uses an unconventional genome replication
mechanism. *Proc Natl Acad Sci U S A* **116**, 8535-8543, doi:10.1073/pnas.1815745116
(2019).
65 Schmidt, M. L., Tews, B. A., Groseth, A. & Hoenen, T. Generation and Optimization of a
Green Fluorescent Protein-Expressing Transcription and Replication-Competent Virus-

Like Particle System for Ebola Virus. *J Infect Dis* **218**, S360-S364, doi:10.1093/infdis/jiy405
(2018).

66   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682, doi:10.1038/nmeth.2019 (2012).

67   Taylor, N. O., Wei, M.-T., Stone, H. A. & Brangwynne, C. P. Quantifying Dynamics in Phase-
Separated Condensates Using Fluorescence Recovery after Photobleaching. *Biophysical*
Journal **117**, 1285-1300, doi:https://doi.org/10.1016/j.bpj.2019.08.030 (2019).

68   Fang, J. *et al.* Functional interactomes of the Ebola virus polymerase identified by
proximity proteomics in the context of viral replication. *Cell Rep* **38**, 110544,
doi:10.1016/j.celrep.2022.110544 (2022).

69   Martell, J. D., Deerinck, T. J., Lam, S. S., Ellisman, M. H. & Ting, A. Y. Electron microscopy
using the genetically encoded APEX2 tag in cultured mammalian cells. *Nat Protoc* **12**,
1792-1816, doi:10.1038/nprot.2017.065 (2017).

70   Phan, S. *et al.* 3D reconstruction of biological structures: automated procedures for
alignment and reconstruction of multiple tilt series in electron tomography. *Adv Struct*
Chem Imaging **2**, 8, doi:10.1186/s40679-016-0021-2 (2017).

1023