1 **Deucravacitinib, a tyrosine kinase 2 pseudokinase inhibitor, protects human beta cells**
2 **against proinflammatory insults**
3

4 **Reinaldo Sousa Dos Santos[1,2*], Daniel Guzman Llorens[1,2], Atenea Alexandra Perez-Serna[1,2],**
5 **Angel Nadal[1,2], Laura Marroqui[1,2*]**
6

7 *[1]Instituto de Investigación, Desarrollo e Innovación en Biotecnología Sanitaria de Elche (IDiBE),*
8 *Universidad Miguel Hernández de Elche, Alicante, Spain.*
9 *[2]CIBER de Diabetes y Enfermedades Metabólicas Asociadas, Instituto de Salud Carlos III.*
10

* Corresponding authors:
Reinaldo Sousa Dos Santos
Instituto de Investigación, Desarrollo e Innovación en Biotecnología Sanitaria de Elche
Universidad Miguel Hernández de Elche, Ed. Torregaitán, Av. de la Universidad s/n 03202 Elche,
Spain.
Email: r.sousa@umh.es
Laura Marroqui
Instituto de Investigación, Desarrollo e Innovación en Biotecnología Sanitaria de Elche
Universidad Miguel Hernández de Elche, Ed. Torregaitán, Av. de la Universidad s/n 03202 Elche,
Spain.
Email: lmarroqui@umh.es

Abstract

Aims/hypothesis: Type 1 diabetes is characterised by pancreatic islet inflammation and

autoimmune-driven pancreatic beta cell destruction. Type I interferons, such as IFNα, are key

players in early human type 1 diabetes pathogenesis, as the activation of the tyrosine kinase 2

(TYK2)-signal transducer and activator of transcription (STAT) pathway induces inflammation, a

long-lasting MHC class I overexpression, endoplasmic reticulum (ER) stress, and beta cell

apoptosis (in synergy with IL-1β). As TYK2 inhibition has been suggested as a potential

therapeutic target for the prevention or treatment of type 1 diabetes, we investigated whether the

selective TYK2 inhibitor deucravacitinib could protect beta cells against the damaging effects of

IFNα and other proinflammatory cytokines (i.e. IFNγ and IL-1β).

Methods: Inflammation, ER stress, and apoptosis were evaluated by real-time PCR, immunoblot,

immunofluorescence, and nuclear dyes. The promoter activity was assessed by luciferase assay

and insulin secretion and content by ELISA. All experiments were performed in the human EndoC-

βH1 cell line.

Results: Pre-treatment with deucravacitinib prevented IFNα effects, such as STAT1 and STAT2

phosphorylation and protein expression as well as MHC class I hyperexpression, in a dose-

dependent manner without affecting beta cell survival and function. Comparison between

deucravacitinib and two Janus kinase inhibitors, ruxolitinib and baricitinib, showed that

deucravacitinib blocked IFNα- but not IFNγ-induced signalling pathway. Pre-treatment with

deucravacitinib protected beta cells from the pro-apoptotic and proinflammatory effects of two

different combinations of cytokines: IFNα + IL-1β and IFNγ + IL-1β. Moreover, this TYK2

inhibitor could partially revert apoptosis and inflammation in cells previously treated with IFNα +

IL-1β or IFNγ + IL-1β.

Conclusions/interpretation: Our findings suggest that, by protecting beta cells against the

deleterious effects of proinflammatory cytokines without affecting beta cell function and survival,

deucravacitinib could be repurposed for the prevention or treatment of early type 1 diabetes.

Keywords: Apoptosis, Deucravacitinib, Inflammation, Pancreatic beta cells, TYK2, Type 1

diabetes, Type I interferons

Introduction

Type 1 diabetes is characterised by pancreatic islet inflammation and specific destruction of pancreatic beta cells by an autoimmune assault, which develops in the context of an inadequate "dialogue" between beta cells and the invading immune cells [1, 2].

A growing body of evidence places type I interferons (IFNs) as key players in the early stages of human type 1 diabetes pathogenesis [3]. IFNα was found in islets from type 1 diabetes patients [4–6], and laser-captured islets from living donors with recent-onset type 1 diabetes showed increased expression of IFN-stimulated genes (ISGs) [7]. In genetically susceptible children, an IFN signature was temporarily amplified preceding the development of autoantibodies and throughout the progress of type 1 diabetes [8, 9]. Recently, three type I IFN response markers, namely human MX Dynamin Like GTPase 1 (MX1), double-stranded RNA sensor protein kinase R, and HLA class I, were found to be expressed in a significantly higher percentage of insulin-containing islets from autoantibody-positive and/or recent-onset type 1 diabetes donors [10]. In human beta cells, IFNα induces inflammation, endoplasmic reticulum (ER) stress as well as a long-lasting overexpression of HLA class I via activation of the tyrosine kinase 2 (TYK2)-signal transducer and activator of transcription (STAT) pathway. Moreover, IFNα induces apoptosis in the presence of IL-1β [11–14].

Targeting the type I IFN signalling pathway has been proposed as a potential adjuvant therapy to treat at-risk individuals or patients still in the very early stages of the disease [3, 15]. Among some of the strategies that have been suggested, inhibitors of Janus kinase (JAK) proteins (JAK1-3 and TYK2) show great promise. Treatment with AZD1480 (a JAK1/JAK2 inhibitor) and ABT 317 (a JAK1-selective inhibitor) protected non-obese diabetic mice against autoimmune diabetes and reversed diabetes in newly diagnosed non-diabetic mice [16, 17]. In human beta cells, clinically used JAK inhibitors, namely ruxolitinib, cerdulatinib, and baricitinib, prevented MHC class I overexpression, ER stress, chemokine production, and apoptosis [13, 14].

Lately, attention has been focused on *TYK2*, a candidate gene for type 1 diabetes whose genetic variants that decrease TYK2 activity are associated with protection against the disease [18–20]. TYK2 is crucial for cell development and IFNα-mediated responses in human beta cells [11, 21, 22]. Partial TYK2 knockdown protected human beta cells against apoptosis and inflammation induced by polyinosinic-polycitidilic acid (poly(I:C)), a mimic of double-stranded RNA produced during viral infection [21]. In mature stem cell-islets, TYK2 knockout or pharmacological

inhibition decreased T-cell-mediated cytotoxicity by preventing IFNα-induced antigen processing

and presentation, including MHC class I expression [22]. As these findings place TYK2 as a

critical regulator of the type I IFN signalling pathway in beta cells, selective TYK2 inhibition has

emerged as a drug target to treat type 1 diabetes. Recently, two novel small molecule inhibitors

binding to the TYK2 pseudokinase domain protected human beta cells against the deleterious

effects of IFNα without compromising beta cell function and susceptibility to potentially

diabetogenic viruses [23].

Deucravacitinib (BMS-986165), a small molecule that selectively targets the TYK2 pseudokinase

domain, has shown great therapeutic potential for immune-mediated diseases, such as lupus

nephritis and systemic lupus erythematosus [24, 25]. In fact, deucravacitinib has been recently

approved for treatment of plaque psoriasis [26–28]. However, no preclinical studies have deeply

explored the possible use of deucravacitinib in the context of type 1 diabetes. Notably, Chandra et

al. recently used deucravacitinib to validate their CRISPR-Cas9-generated *TYK2* knockout in

human induced pluripotent stem cells, but did not provide further characterisation of its effects in

beta cells [22].

In the present study, we described the effects of deucravacitinib on the human EndoC-βH1 beta

cell line, including its ability to prevent IFNα-triggered signalling pathway and subsequent

damaging effects on beta cells. We report that deucravacitinib prevented IFNα effects in a dose-

dependent manner without affecting beta cell survival and function. Compared with ruxolitinib

and baricitinib, deucravacitinib inhibited the IFNα- but not IFNγ-stimulated signalling pathway.

Interestingly, this TYK2 inhibitor protected beta cells not only against the deleterious effects of

IFNα but also from other proinflammatory cytokines, namely IFNγ and IL-1β, which suggests that

deucravacitinib could be introduced as an adjuvant protective therapy at different stages of the

disease to avoid the progressive loss of beta cell mass.

Methods

Culture of EndoC-βH1 cells

The human EndoC-βH1 beta cell line [research resource identifier (RRID): CVCL_L909,

Univercell-Biosolutions, France] was cultured in Matrigel/fibronectin-coated plates as previously

described [29]. Cells were cultured in DMEM containing 5.6 mmol/l glucose, 10 mmol/l

nicotinamide, 5.5 μg/ml transferrin, 50 μmol/l 2-mercaptoethanol, 6.7 ng/ml selenite, 2% BSA

fatty acid free, 100 U/ml penicillin, and 100 μg/ml streptomycin. We confirmed that cells were

mycoplasma-free using the MycoAlert Mycoplasma Detection Kit (Lonza, Basel, Switzerland).

Cell treatments

Proinflammatory cytokine concentrations were selected according to previously established

experiments in human beta cells [11, 30]: recombinant human IFNα (PeproTech Inc., Rocky Hill,

NJ) at 1000 U/ml; recombinant human IFNγ (PeproTech Inc., Rocky Hill, NJ) at 1000 U/ml; and

recombinant human IL-1β (R&D Systems, Abingdon, UK) at 50 U/ml. Cells were transfected with

1 μg/ml poly(I:C) (InvivoGen, San Diego, CA) as indicated [31]. Ruxolitinib, baricitinib, or

deucravacitinib (Selleckchem, Planegg, Germany) were prepared by dissolution in DMSO (used

as vehicle) and cells were treated as indicated in the figures. Ruxolitinib and baricitinib

concentrations were selected based on previous dose-response experiments (unpublished data).

For treatments involving cytokines, 2% FBS was added to the culture medium.

Cell viability assessment

129 The percentage of apoptosis was measured by fluorescence microscopy upon staining with the

130 DNA-binding dyes Hoechst 33342 and propidium iodide (Sigma-Aldrich, Saint Louis, MO, USA)

131 as described [32]. At least 600 cells were counted for each experimental condition. Viability was

132 assessed by two independent observers, one of whom was unaware of sample identity, with an

133 agreement between results of >90%.

Caspase 3/7 activity

136 Caspase 3/7 activity was determined using the Caspase-Glo® 3/7 assay (Promega, Madison, WI,

137 USA) following the manufacturer's instructions. Briefly, upon treatment in 100 μl culture medium,

138 cells were incubated with 100 μl Caspase-Glo® 3/7 reagent at room temperature for 1 h before

139 recording luminescence with a POLASTAR plate reader (BMG Labtech, Ortenberg, Germany).

C-X-C motif chemokine ligand 10 (CXCL10) measurements

142 The release of CXCL10 to the culture medium was detected using Human ProcartaPlex

143 immunoassays (Invitrogen, Vienna, Austria) following the manufacturer's recommendations.

144 Reactions were read with a MagPix system (Luminex, Austin, TX, USA).

Luciferase reporter assays

EndoC-βH1 cells were transfected using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) with pRL-CMV encoding *Renilla* luciferase (Promega) and luciferase reporter constructs for either gamma-interferon activation site (GAS) (Panomics, Fremont, CA, USA) or IFN-stimulated regulatory element (ISRE) (kindly provided by Dr Izortze Santin, University of the Basque Country, Spain). After recovery, cells were treated with either IFNα for 2 h or IFNγ for 24 h [33]. Luciferase activity was measured in a POLASTAR plate reader (BMG Labtech) using the Dual-Luciferase Reporter Assay System (Promega) and corrected for the luciferase activity of the internal control plasmid, i.e. pRL-CMV.

RNA analysis

Poly(A)$^+$ mRNA was extracted using Dynabeads mRNA DIRECT kit (Invitrogen) and cDNA synthesis was performed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). Real-time PCR was performed on the CFX96 Real Time System (Bio-Rad) as described [34]. The housekeeping gene β-actin was used to correct expression values ($2^{-\Delta Ct}$) [35]. All primers used here are listed in ESM Table 1.

Immunoblotting and immunofluorescence analyses

Western blotting analysis was performed as described [21]. Briefly, cells were washed with cold PBS and lysed in Laemmli buffer. Immunoblot was performed using antibodies against STAT1 and STAT2 (phosphorylated and total forms; all at 1:1000 dilution), and α-tubulin (1:5000). Peroxidase-conjugated antibodies (1:5000) were used as secondary antibodies. SuperSignal West Femto chemiluminescent substrate (Thermo Scientific, Rockford, IL, USA) and ChemiDoc XRS+ (Bio-Rad Laboratories, Hercules, CA, USA) were used to detect bands.

Immunofluorescence was carried out as described [11, 21]. First, cells were washed with cold PBS and fixed with 4% paraformaldehyde. Afterwards, cells were permeabilised and incubated with the mouse anti-MHC Class I (W6/32) antibody (1:1000). The Alexa Fluor 568 polyclonal goat anti-mouse IgG (1:500) was used as secondary antibody. Upon staining with Hoechst 33342, coverslips were mounted with fluorescent mounting medium (Dako, Carpintera, CA, USA) and immunofluorescence was observed with an inverted fluorescence microscope Zeiss Confocal

LSM900 with Airyscan 2 microscope equipped with a camera (Zeiss-Vision, Munich, Germany),

and images were acquired at x40 magnification and analysed using ZEN software (version 3.3;

Zeiss-Vision, Munich, Germany) and open-source FIJI software (version 2.0; https://fiji.sc).

All antibodies used here are provided in ESM Table 2.

Glucose-stimulated insulin secretion

After preincubation in modified Krebs-Ringer for 1 h, cells were sequentially stimulated with low

(0 mmol/l) and high glucose (20 mmol/l) for 1 h (each stimulation) as previously described [36].

Insulin secreted and insulin content from lysed cells were measured using a human insulin ELISA

kit (Mercodia, Uppsala, Sweden) following the manufacturer's instructions. See ESM Methods for

further details.

Statistical analyses

The GraphPad Prism 7.0 software (GraphPad Software, La Jolla, CA, USA) was used for statistical

analyses. Data are shown as mean ± SEM of independent experiments (i.e. considering EndoC-

βH1 cells from different passages as $n = 1$, with individual data added to the column bars). The

statistical significance of differences between groups was evaluated using one-way ANOVA

followed by Dunnett's test or two-way ANOVA followed by Sidak's test or by Dunnett's test, as

appropriate.

Results

Deucravacitinib prevents IFNα effects in EndoC-βH1 cells

IFNα-mediated JAK/TYK2 activation leads to phosphorylation of STAT1 and STAT2, which will

eventually upregulate several ISGs, including STATs, *HLA-ABC*, *CXCL10*, and *MX1* (ESM Fig.

1a). Pre-treatment of EndoC-βH1 cells with deucravacitinib inhibited IFNα-induced STAT1 and

STAT2 phosphorylation in a dose-dependent manner, where deucravacitinib showed greater

potency towards IFNα-stimulated STAT1 inhibition (Fig. 1a,b). We then selected two doses, 10

and 1000 nmol/l, for the follow-up experiments. Next, we examined the effect of deucravacitinib

on the kinetics of IFNα-induced STAT activation. As expected, the phosphorylation of STAT1

and STAT2 was markedly amplified by IFNα at early time points (1–4 h) and returned to baseline

by 24 h (Fig. 1c,d and ESM Fig. 1b). STAT1 and STAT2 protein expression augmented in a time-

dependent manner. Although both proteins were already upregulated by 8 h, STAT2 reached an

expression peak at 16 h, while STAT1 expression was still increasing by 24 h (Fig. 1c-f). Exposure

to 1000 nmol/l deucravacitinib abrogated the IFNα-stimulated STAT1 and STAT2

phosphorylation and protein expression, whereas 10 nmol/l deucravacitinib had only a minor effect

(Fig. 1c-f and ESM Fig. 1b). These findings are better evidenced by analysing the area under the

curve of the phosphorylated and total forms of STAT1 and STAT2 (ESM Fig. 1c-f). Finally, MHC

class I protein expression stimulated by IFNα- was completely blocked by 1000 nmol/l

deucravacitinib (Fig. 1g,h).

Beta cell survival and function are not affected by deucravacitinib

We examined whether deucravacitinib would interfere with beta cell survival and function in the

absence or presence of IFNα. After 24 h exposure, 1000 nmol/l deucravacitinib did not affect beta

cell viability (ESM Fig. 2a) nor change glucose-stimulated insulin secretion and insulin content

(ESM Fig. 2b,c). IFNα did not affect beta cell viability and function as described [11, 23].

IFNα, but not IFNγ signalling pathway is blocked by deucravacitinib

We compared deucravacitinib with ruxolitinib and baricitinib, two JAK1/JAK2 inhibitors

previously tested in beta cells [13, 14, 37]. First, we measured STAT1 and STAT2 phosphorylation

upon stimulation with IFNα or IFNγ (Fig. 2a-d and ESM Fig. 3a,b). Ruxolitinib, baricitinib, and

deucravacitinib prevented IFNα-stimulated increase in P-STAT1 and P-STAT2 levels (Fig. 2a,c

and ESM Fig. 3b). Nevertheless, deucravacitinib did not change IFNγ-induced STAT1

phosphorylation, whereas ruxolitinib and baricitinib blocked it (Fig. 2b,d). Of note, IFNγ did not

induce STAT2 phosphorylation (Fig. 2b) [33]. Next, we assessed ISRE and GAS activities upon

stimulation with IFNα and IFNγ (Fig. 2e,f and ESM Fig. 3c,d). All three inhibitors abrogated

IFNα-stimulated ISRE reporter activity (Fig. 2e), whereas IFNγ-induced GAS activation was

barely affected by deucravacitinib (Fig. 2f). As expected, ISRE and GAS activities were not

stimulated by, respectively, IFNγ and IFNα (ESM Fig. 3c,d). These findings corroborate the

deucravacitinib mode of action, which specifically binds to and inhibits TYK2 without affecting

JAK1/JAK2 pathways. As TYK2 does not participate in the IFNγ pathway, the lack of

deucravacitinib effect in IFNγ-stimulated changes is expected.

Deucravacitinib blocks IFNα-induced upregulation of ISGs, but not ER stress markers

We assessed the effects of these three inhibitors on the expression of some ISGs and ER stress markers. All three inhibitors prevented IFNα-induced upregulation of *HLA-ABC*, *CXCL10*, and *MX1* in a dose-dependent manner (Fig. 3a-c). Although ruxolitinib and baricitinib inhibited the expression of the ER stress markers C/EBP homologous protein (*CHOP*, also known as *DDIT3*) and spliced isoform of XBP1 X-box binding protein 1 (*XBP1s*), only the lower dose of deucravacitinib reduced CHOP expression (Fig. 3d,e). None of the three inhibitors changed the mRNA expression of activating transcription factor 3 (*ATF3*) (Fig. 3f).

Deucravacitinib prevents IFNα + IL-1β-induced effects in beta cells

Previous studies showed that a combination of IFNα + IL-1β, two cytokines that might be present in the islet milieu at early stages of insulitis, induce beta cell apoptosis, inflammation, and ER stress [11, 14, 23] (Fig. 4a). Then, we investigated whether deucravacitinib protects beta cells after IFNα + IL-1β exposure for 24 h. We observed that deucravacitinib completely prevented IFNα + IL-1β-induced apoptosis, which was assessed by two approaches, namely DNA-binding dyes and caspase 3/7 activity assay (Fig. 4b,c). Moreover, cells treated with deucravacitinib showed reduced levels of P-STAT1 and STAT1 (Fig. 4d-f) as well as *HLA-ABC*, *MX1*, *CXCL10*, and *CHOP* mRNA expression (Fig. 4g-j). MHC class I protein expression and CXCL10 secretion to the medium were also decreased by TYK2 inhibition (Fig. 4k-m).

Deucravacitinib abrogates IFNγ + IL-1β-induced apoptosis in beta cells

We then evaluated whether deucravacitinib would protect against cytokines that, as compared with IFNα, probably appear later in the progression of islet inflammation, such as IFNγ and IL-1β [1, 38] (Fig. 5a). After treatment for 24 h, deucravacitinib inhibited IFNγ + IL-1β-induced apoptosis in a dose-dependent manner (60% and 92% protection at 10 and 1000 nmol/l, respectively) (Fig. 5b). These results were confirmed by the caspase 3/7 activity assay (Fig. 5c). Pre-treatment with deucravacitinib did not block IFNγ + IL-1β-induced STAT1 phosphorylation and protein expression (Fig. 5d-f) or *HLA-ABC* mRNA expression (Fig. 5g). Conversely, deucravacitinib diminished *MX1* and *CXCL10* mRNA expression, whereas *CHOP* was reduced at the lower dose of deucravacitinib (Fig. 5h-j).

Poly(I:C)-induced apoptosis is not changed by TYK2 inhibition

As intracellular exposure to poly(I:C), a mimic of viral infection, results in the production and secretion of type I IFNs as well as beta cell apoptosis [21, 39, 40], we tested deucravacitinib after treatment with poly(I:C) for 24 h. Contrary to what we observed for cytokine-triggered cell death (Figs 4 and 5), apoptosis induced by poly(I:C) was not altered by deucravacitinib (ESM Fig. 4a). Curiously, despite the absence of an effect on viability, deucravacitinib inhibited poly(I:C)-stimulated *HLA-ABC*, *MX1*, and *CHOP* upregulation (ESM Fig. 4b-d). While *CXCL10* mRNA expression was dampened by TYK2 inhibition, CXCL10 protein secretion was not significantly reduced (ESM Fig. 4e,f).

The harmful effects of cytokines are partially reverted by deucravacitinib

Up to this point, we investigated whether pre-treatment with deucravacitinib would prevent the effects of different cytokines in beta cells. Here, we assessed if deucravacitinib could reverse these damaging effects. EndoC-βH1 cells were pre-treated with either IFNα + IL-1β or IFNγ + IL-1β for 24 h. Afterwards, 1000 nmol/l deucravacitinib was added for an additional 24 h still in the presence of the different mixes of cytokines (Figs 6a and 7a). IFNα + IL-1β-induced apoptosis was partially reversed by deucravacitinib (60% decrease) (Fig. 6b). The expression of *HLA-ABC* mRNA stimulated by IFNα + IL-1β remained unchanged in the presence of deucravacitinib (Fig. 6c), which agrees with previous data showing a long-lasting expression of *HLA-ABC* [13]. STAT1 protein levels and CXCL10 secretion as well as *CHOP* mRNA expression were reduced by 26%-42%, while the expression of *MX1* and *CXCL10* was completely inhibited by deucravacitinib (Fig. 6 d-i). Of note, STAT1 phosphorylation was not detected, probably due to a process of desensitisation following the initial IFNα stimulus [33] (Fig. 6c).

Similarly to IFNα + IL-1β (Fig. 6), deucravacitinib diminished IFNγ + IL-1β-induced apoptosis (64% decrease) but did not modify *HLA-ABC* mRNA expression (Fig. 7b,c). Protein levels of STAT1 and CXCL10, however, were not altered by the TYK2 inhibitor, whereas a slight, non-significant 30% reduction was seen in *CHOP* expression (Fig. 7d,e,g,i). Expression of *MX1* and *CXCL10* was only partially blocked by deucravacitinib under IFNγ + IL-1β conditions (Fig. 7f,h).

Discussion

Due to its implication in type 1 diabetes pathogenesis, IFNα pathway has arisen as an interesting
therapeutic target. Reduction of IFNα extracellular levels, blockade of IFNα itself and/or its
receptor, and reduction of the activity of proteins mediating IFN effects have been proposed as
means to diminish IFNα deleterious effects [41]. In rodent models, some of these strategies have
successfully prevented diabetes development [16, 17, 42–44].

One of the most promising therapeutic approaches for type 1 diabetes prevention/early treatment
is the targeting of the JAK-STAT pathway with JAK inhibitors [3, 15]. This strategy has been
clinically approved for the treatment of some autoimmune diseases, including systemic lupus
erythematosus [45], rheumatoid arthritis [46], and psoriasis [26, 27]. Although there are no
approved JAK inhibitors being clinically used for type 1 diabetes, recent preclinical data suggest
that these inhibitors could be repurposed to treat this disease [13, 14, 16, 17, 22, 23]. In fact, a
clinical trial aiming to determine whether baricitinib could slow the progressive, immune-
mediated loss of beta cell function and mass that occurs in type 1 diabetes is currently ongoing
[47].

In the current study, we tested whether the TYK2 inhibitor deucravacitinib could protect human
beta cells against the deleterious effects of IFNα and other cytokines. We chose to focus on this
TYK2 inhibitor for two main reasons: first, due to the importance of TYK2 to type 1 diabetes
pathogenesis. For instance, TYK2 regulates pro-apoptotic and proinflammatory pathways via
regulation of the IFNα signalling, antigen processing and presentation, and modulation of cytokine
and chemokine production in beta cells [21, 22]. Second, exploring a drug that has been recently
approved by the US Food and Drug Administration to treat another autoimmune disease, namely
plaque psoriasis [28], increases its repositioning potential for type 1 diabetes and facilitates the
bench-to-bedside transition.

Deucravacitinib is a small-molecule ligand that binds to and stabilizes the TYK2 pseudokinase
domain, leading to highly potent and selective allosteric TYK2 inhibition [24, 48]. Inhibition of
IFNα-induced STAT phosphorylation by deucravacitinib has been shown in several cell types,
such as CD3[+] T cells, CD19[+] B cells, and CD14[+] monocytes [24]. Here we showed that
deucravacitinib also prevents IFNα-stimulated STAT1 and STAT2 phosphorylation in human
EndoC-βH1 cell line. Furthermore, in agreement with previous findings [24], deucravacitinib also
showed higher potency against TYK2-mediated phosphorylation of STAT1 compared with
STAT2 phosphorylation in our experimental model. Notably, at the concentrations used in our

study, deucravacitinib did not affect beta cell function and viability, which is a desired feature for a drug with therapeutic potential.

Compared with ruxolitinib and baricitinib, two clinically available JAK1/JAK2 inhibitors, deucravacitinib was more potent against IFNα-stimulated STAT phosphorylation, ISRE activity, and mRNA expression of *HLA-ABC*, *MX1*, and *CXCL10*. However, unlike ruxolitinib and baricitinib, deucravacitinib did not block the IFNα-mediated upregulation of the ER stress markers *CHOP* and *XBP1s*. Our results partially agree with the ones reported by Coomans de Brachène et al., where two TYK2 inhibitors failed to prevent IFNα-induced *CHOP* expression in EndoC-βH1 cells but inhibited *CHOP* and *ATF3* expression in dispersed human islets [23].

Prior studies have shown that other JAK/TYK2 inhibitors could prevent the detrimental effects of IFNα + IL-1β, such as apoptosis and inflammation [14, 23]. Therefore, we proceeded to investigate whether deucravacitinib could protect beta cells against the harmful effects of two different combinations of cytokines: IFNα + IL-1β (early insulitis) and IFNγ + IL-1β (late insulitis). In both scenarios, pre-treatment with deucravacitinib not only protected against cytokine-induced apoptosis but prevented the upregulation of *HLA-ABC*, *MX1*, *CXCL10*, and *CHOP*. Additionally, in cells treated with IFNα + IL-1β, deucravacitinib blocked the overexpression of MHC class I at the cell surface and CXCL10 secretion to the medium. Importantly, the addition of deucravacitinib when cytokine exposure was already ongoing could reverse, at least in part, the deleterious effects of these cytokines. Although it seems clear that deucravacitinib confers protection against IFNα + IL-1β by directly inhibiting the TYK2-mediated pathway, it remains to be answered how deucravacitinib protects against IFNγ + IL-1β-induced effects. Indeed, our present data suggest that deucravacitinib does not interfere with the IFNγ-mediated signalling pathway.

Due to TYK2 role in the IFNα-mediated antiviral response, as evidenced by the regulation of *MX1* mRNA expression and STAT1 protein levels [11, 22, 23], another desired characteristic for a TYK2 inhibitor would be to block IFNα pathway without sensitising beta cells to viral infections. To test this hypothesis, we mimicked a viral infection by exposing EndoC-βH1 cells to poly(I:C). In our model, deucravacitinib abrogated poly(I:C)-stimulated inflammation but did not change poly(I:C)-induced beta cell apoptosis. Even though our results were obtained with a synthetic viral double-stranded RNA analogue, they conform to a previous study showing that two novel TYK2 inhibitors did not sensitize human beta cells to potentially diabetogenic coxsackieviruses CVB1

and CVB5 [23]. Importantly, deucravacitinib and these two TYK2 inhibitors share the same
mechanism of action, i.e. binding to the TYK2 pseudokinase domain.

Based on our findings, it will be interesting to test whether novel small molecule TYK2
pseudokinase ligands [49] could also protect beta cells from IFNα deleterious effects.
Nevertheless, we must bear in mind that completely inhibiting TYK2 may be counterproductive,
as it might lead to susceptibility to microorganisms (e.g. mycobacteria and virus) and
immunodeficiency [50]. Thus, regardless of the TYK2 inhibitor chosen, we should focus on doses
that induce a partial inhibition, as seen in individuals with a protective single nucleotide
polymorphism in the *TYK2* gene [18], as it could offer maximal efficacy with reduced risk of
developing secondary infections.

One potential limitation of our study is its purely in vitro nature, which may limit our conclusions
regarding the use of deucravacitinib to treat a disease as complex as type 1 diabetes. Conversely,
our findings, along with those reported by Coomans de Brachène et al. [23] and Chandra et al.
[22], provide further preclinical evidence that TYK2 inhibitors could be considered a strategy for
an early therapy for type 1 diabetes. The next logical step would be to investigate whether our in
vitro findings could be translated to animal models of type 1 diabetes (e.g. NOD and RIP-B7.1
mice).

In conclusion, we have provided evidence that deucravacitinib protects beta cells against the
deleterious effects of proinflammatory cytokines, such as IFNα, IFNγ and IL-1β, without affecting
beta cell function and survival. Our present findings add to the growing body of evidence showing
that TYK2 inhibition may be an efficient strategy to treat type 1 diabetes. Moreover, we suggest
that deucravacitinib could be repurposed to treat pre-symptomatic type 1 diabetes subjects (i.e.
positive for 2–3 autoantibodies but still normoglycemic) or be introduced in the early stages of
type 1 diabetes onset.

Acknowledgements
The authors are grateful to Beatriz Bonmati Botella, Maria Luisa Navarro, and Salomé Ramon for
their excellent technical support. Once again, we thank Dr. Izortze Santin, University of the Basque
Country, Spain, for providing the luciferase reporter construct for ISRE. Figures 4a, 5a, 6a, 7a,
ESM Fig. 1a, and ESM Fig. 3a were created with BioRender.com.

Data availability

All data generated or analysed during this study are included in this published article (and its supplementary information files). Data are however available from the corresponding authors upon reasonable request.

Funding

LM is funded by the grant PID2020-117569RA-I00 by MCIN/AEI/10.13039/501100011033 and by the grant SEJI/2018/023 by Generalitat Valenciana. AN is supported by European Union's Horizon 2020 research and innovation programme under grant agreement GOLIATH No. 825489, by the grant PID2020-117294RB-I00 by MCIN/AEI/10.13039/501100011033, and by the grant PROMETEO II/2020/006 by Generalitat Valenciana. This research was supported by CIBER-Consorcio Centro de Investigación Biomédica en Red (CB07/08/0002), Instituto de Salud Carlos III, Ministerio de Ciencia e Innovación.

Authors' relationships and activities

The authors declare that there are no relationships or activities that might bias, or be perceived to bias, their work.

Contribution statement

RSDS and LM: Conceptualization, Supervision, Visualization, Investigation, Formal analysis, and Writing – original draft & editing; DGL and AAP-S: Investigation, Formal analysis, and Writing – review & editing; AN: Resources and Writing – review & editing; LM: Resources, Funding acquisition, and Project administration. All authors have read and given approval to the final version of the manuscript. RSDS and LM are guarantors of this work and, as such, had full access to all the data presented herein and take responsibility for the integrity of the data and the accuracy of data analyses.

References

1.   Eizirik DL, Pasquali L, Cnop M (2020) Pancreatic β-cells in type 1 and type 2 diabetes mellitus: different pathways to failure. Nat Rev Endocrinol 26(7):349–362. https://doi.org/10.1038/s41574-020-0355-7
2.   Mallone R, Eizirik DL (2020) Presumption of innocence for beta cells: why are they vulnerable autoimmune targets in type 1 diabetes? Diabetologia 63(10):1999–2006.

https://doi.org/10.1007/s00125-020-05176-7

3.   Marroqui L, Perez-Serna AA, Babiloni-Chust I, Dos Santos RS (2021) Type I interferons as key players in pancreatic β-cell dysfunction in type 1 diabetes. Int Rev Cell Mol Biol 359:1–80. https://doi.org/10.1016/bs.ircmb.2021.02.011

4.   Foulis A, Farquharson M, Meager A (1987) Immunoreactive alpha-interferon in insulin-secreting beta cells in type 1 diabetes mellitus. The Lancet 2(8573):1423–1427. https://doi.org/10.1016/S0140-6736(87)91128-7

5.   Somoza N, Vargas F, Roura-Mir C, et al (1994) Pancreas in recent onset insulin-dependent diabetes mellitus: Changes in HLA, adhesion molecules and autoantigens, restricted T cell receptor Vβ usage, and cytokine profile. J Immunol 153(3):1360–1377

6.   Huang X, Yuan J, Goddard A, et al (1995) Interferon expression in the pancreases of patients with type I diabetes. Diabetes 44(6):658–664. https://doi.org/10.2337/diab.44.6.658

7.   Lundberg M, Krogvold L, Kuric E, Dahl-Jørgensen K, Skog O (2016) Expression of interferon-stimulated genes in insulitic pancreatic islets of patients recently diagnosed with type 1 diabetes. Diabetes 65(10):3104–3110. https://doi.org/10.2337/db16-0616

8.   Ferreira RC, Guo H, Coulson RMR, et al (2014) A type I Interferon transcriptional signature precedes autoimmunity in children genetically at risk for type 1 diabetes. Diabetes 63(7):2538–2550. https://doi.org/10.2337/db13-1777

9.   Kallionpaa H, Elo LL, Laajala E, et al (2014) Innate Immune Activity Is Detected Prior to Seroconversion in Children With HLA-Conferred Type 1 Diabetes Susceptibility. Diabetes 63(7):2402–2414. https://doi.org/10.2337/db13-1775

10.   Apaolaza PS, Balcacean D, Zapardiel-Gonzalo J, et al (2021) Islet expression of type I interferon response sensors is associated with immune infiltration and viral infection in type 1 diabetes. Sci Adv 7(9):eabd6527. https://doi.org/10.1126/sciadv.abd6527

11.   Marroqui L, Dos Santos RS, Op de beeck A, et al (2017) Interferon-α mediates human beta cell HLA class I overexpression, endoplasmic reticulum stress and apoptosis, three hallmarks of early human type 1 diabetes. Diabetologia 60(4):656–667. https://doi.org/10.1007/s00125-016-4201-3

12.   Lombardi A, Tomer Y (2017) Interferon alpha impairs insulin production in human beta cells via endoplasmic reticulum stress. J Autoimmun 80:48–55. https://doi.org/10.1016/j.jaut.2017.02.002

13.   Coomans de Brachène A, Dos Santos RS, Marroqui L, et al (2018) IFN-α induces a preferential long-lasting expression of MHC class I in human pancreatic beta cells. Diabetologia 61(3):636–640. https://doi.org/10.1007/s00125-017-4536-4

14.   Colli ML, Ramos-Rodríguez M, Nakayasu ES, et al (2020) An integrated multi-omics approach identifies the landscape of interferon-α-mediated responses of human pancreatic beta cells. Nat Commun 11(1):2584. https://doi.org/10.1038/s41467-020-16327-0

15.   Eizirik DL, Szymczak F, Alvelos MI, Martin F (2021) From Pancreatic β-Cell Gene Networks to Novel Therapies for Type 1 Diabetes. Diabetes 70(9):1915–1925. https://doi.org/10.2337/dbi20-0046

16.   Trivedi PM, Graham KL, Scott NA, et al (2017) Repurposed JAK1/JAK2 inhibitor reverses established autoimmune insulitis in NOD mice. Diabetes 66(6):1650–1660. https://doi.org/10.2337/db16-1250

17.   Ge T, Jhala G, Fynch S, et al (2020) The JAK1 Selective Inhibitor ABT 317 Blocks Signaling Through Interferon-γ and Common γ Chain Cytokine Receptors to Reverse Autoimmune Diabetes in NOD Mice. Front Immunol 11:588543.

https://doi.org/10.3389/fimmu.2020.588543

18. Dendrou CA, Cortes A, Shipman L, et al (2016) Resolving TYK2 locus genotype-To-
phenotype differences in autoimmunity. Sci Transl Med 8(363):363ra149.
https://doi.org/10.1126/scitranslmed.aag1974

19. Tao JH, Zou YF, Feng XL, et al (2011) Meta-analysis of TYK2 gene polymorphisms
association with susceptibility to autoimmune and inflammatory diseases. Mol Biol Rep
38(7):4663–4672. https://doi.org/10.1007/s11033-010-0601-5

20. Wallace C, Smyth DJ, Maisuria-Armer M, Walker NM, Todd JA, Clayton DG (2010) The
imprinted DLK1-MEG3 gene region on chromosome 14q32.2 alters susceptibility to type 1
diabetes. Nat Genet 42(1):68–71. https://doi.org/10.1038/ng.493

21. Marroqui L, Dos Santos RS, Fløyel T, et al (2015) TYK2, a candidate gene for type 1
diabetes, modulates apoptosis and the innate immune response in human pancreatic β-cells.
Diabetes 64(11):3808–3817. https://doi.org/10.2337/db15-0362

22. Chandra V, Ibrahim H, Halliez C, et al (2022) The type 1 diabetes gene TYK2 regulates β-
cell development and its responses to interferon-α. Nat Commun 13(1):6363.
https://doi.org/10.1038/s41467-022-34069-z

23. Coomans de Brachène A, Castela A, Op de Beeck A, et al (2020) Pre-clinical evaluation
of TYK2 inhibitors for human beta cell protection in type 1 diabetes. Diabetes Obes Metab
22(10):1827–1836. https://doi.org/10.1111/dom.14104

24. Burke JR, Cheng L, Gillooly KM, et al (2019) Autoimmune pathways in mice and humans
are blocked by pharmacological stabilization of the TYK2 pseudokinase domain. Sci Transl Med
11(502):eaaw1736. https://doi.org/10.1126/scitranslmed.aaw1736

25. Morand E, Pike M, Merrill JT, et al Deucravacitinib, a Tyrosine Kinase 2 Inhibitor, in
Systemic Lupus Erythematosus: A Phase II, Randomized, Double-Blind, Placebo-Controlled
Trial. Arthritis Rheumatol n/a(n/a). https://doi.org/10.1002/art.42391

26. Armstrong AW, Gooderham M, Warren RB, et al (2022) Deucravacitinib versus placebo
and apremilast in moderate to severe plaque psoriasis: efficacy and safety results from the 52-
week, randomized, double-blinded, placebo-controlled phase 3 POETYK PSO-1 trial. J Am Acad
Dermatol 88(1):P29-39. https://doi.org/10.1016/j.jaad.2022.07.002

27. Strober B, Thaçi D, Sofen H, et al (2022) Deucravacitinib versus placebo and apremilast
in moderate to severe plaque psoriasis: efficacy and safety results from the 52-week, randomized,
double-blinded, phase 3 POETYK PSO-2 trial. J Am Acad Dermatol 88(1):P40-51.
https://doi.org/10.1016/j.jaad.2022.08.061

28. Bristol Myers Squibb. U.S. Food and Drug Administration Approves Sotyktu™
(deucravacitinib), Oral Treatment for Adults with Moderate-to-Severe Plaque Psoriasis.
https://news.bms.com/news/details/2022/U.S.-Food-and-Drug-Administration-Approves-
Sotyktu-deucravacitinib-Oral-Treatment-for-Adults-with-Moderate-to-Severe-Plaque-
Psoriasis/default.aspx. Accessed 27 Nov 2022

29. Ravassard P, Hazhouz Y, Pechberty S, et al (2011) A genetically engineered human
pancreatic β cell line exhibiting glucose-inducible insulin secretion. J Clin Invest 121(9):3589–97.
https://doi.org/10.1172/JCI58447

30. Brozzi F, Nardelli TR, Lopes M, et al (2015) Cytokines induce endoplasmic reticulum
stress in human, rat and mouse beta cells via different mechanisms. Diabetologia 58(10):2307–
2316. https://doi.org/10.1007/s00125-015-3669-6

31. Santin I, Moore F, Grieco FA, Marchetti P, Brancolini C, Eizirik DL (2012) USP18 is a
key regulator of the interferon-driven gene network modulating pancreatic beta cell inflammation

and apoptosis. Cell Death Dis 3(11):e49. https://doi.org/10.1038/cddis.2012.158

32.   Santin I, Dos Santos RS, Eizirik DL (2016) Pancreatic beta cell survival and signaling
pathways: Effects of type 1 diabetes-associated genetic variants. In: Methods in Molecular
Biology. Humana Press, New York, NY, USA, pp 21–54

33.   Dhayal S, Leslie KA, Baity M, et al (2022) Temporal regulation of interferon signalling in
human EndoC-βH1 cells. J Mol Endocrinol 69(2):299–313. https://doi.org/10.1530/JME-21-0224

34.   Villar-Pazos S, Martinez-Pinna J, Castellano-Muñoz M, et al (2017) Molecular
mechanisms involved in the non-monotonic effect of bisphenol-A on Ca2+ entry in mouse
pancreatic β-cells. Sci Rep 7(1):11770. https://doi.org/10.1038/s41598-017-11995-3

35.   Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-
Time   Quantitative   PCR   and   the   2−ΔΔCT   Method.   Methods   25(4):402–408.
https://doi.org/10.1006/meth.2001.1262

36.   Dos Santos RS, Medina-Gali RM, Babiloni-Chust I, Marroqui L, Nadal A (2022) In Vitro
Assays to Identify Metabolism-Disrupting Chemicals with Diabetogenic Activity in a Human
Pancreatic β-Cell Model. Int J Mol Sci 23(9):5040. https://doi.org/10.3390/ijms23095040

37.   Colli ML, Hill JLE, Marroquí L, et al (2018) PDL1 is expressed in the islets of people with
type 1 diabetes and is up-regulated by interferons-α and-γ via IRF1 induction. EBioMedicine
36:367–375. https://doi.org/10.1016/j.ebiom.2018.09.040

38.   Colli ML, Szymczak F, Eizirik DL (2020) Molecular Footprints of the Immune Assault on
Pancreatic   Beta   Cells   in   Type   1   Diabetes.   Front   Endocrinol   11:666.
https://doi.org/10.3389/fendo.2020.568446

39.   Dogusan Z, Garcia M, Flamez D, et al (2008) Double-stranded RNA induces pancreatic β-
cell apoptosis by activation of the toll-like receptor 3 and interferon regulatory factor 3 pathways.
Diabetes 57(5):1236–1245. https://doi.org/10.2337/db07-0844

40.   Dos Santos RS, Marroqui L, Velayos T, et al (2019) DEXI, a candidate gene for type 1
diabetes, modulates rat and human pancreatic beta cell inflammation via regulation of the type I
IFN/STAT signalling pathway. Diabetologia 62(3):459–472. https://doi.org/10.1007/s00125-018-
4782-0

41.   Stewart TA (2003) Neutralizing interferon alpha as a therapeutic approach to autoimmune
diseases.   Cytokine   Growth   Factor   Rev   14(2):139–154.   https://doi.org/10.1016/S1359-
6101(02)00088-6

42.   Stewart T, Hultgren B, Huang X, Pitts-Meek S, Hully J, MacLachlan N (1993) Induction
of type I diabetes by interferon-alpha in transgenic mice. Science 260(5116):1942–1946.
https://doi.org/10.1126/science.8100367

43.   Li Q, Xu B, Michie SA, Rubins KH, Schreriber RD, McDevitt HO (2008) Interferon-α
initiates type 1 diabetes in nonobese diabetic mice. Proc Natl Acad Sci U S A 105(34):12439–
12444. https://doi.org/10.1073/pnas.0806439105

44.   Marro BS, Ware BC, Zak J, De La Torre JC, Rosen H, Oldstone MBA (2017) Progression
of type 1 diabetes from the prediabetic stage is controlled by interferon-α signaling. Proc Natl
Acad Sci U S A 114(14):3708–3713. https://doi.org/10.1073/pnas.1700878114

45.   Wallace DJ, Furie RA, Tanaka Y, et al (2018) Baricitinib for systemic lupus erythematosus:
a double-blind, randomised, placebo-controlled, phase 2 trial. The Lancet 392(10143):222–231.
https://doi.org/10.1016/S0140-6736(18)31363-1

46.   Taylor PC, Keystone EC, Van Der Heijde D, et al (2017) Baricitinib versus Placebo or
Adalimumab   in   Rheumatoid   Arthritis.   N   Engl   J   Med   376(652–662).
https://doi.org/10.1056/NEJMoa1608345

47.   Waibel M, Thomas HE, Wentworth JM, et al (2022) Investigating the efficacy of baricitinib
in new onset type 1 diabetes mellitus (BANDIT)—study protocol for a phase 2, randomized,
placebo controlled trial. Trials 23(1):433. https://doi.org/10.1186/s13063-022-06356-z
48.   Wrobleski ST, Moslin R, Lin S, et al (2019) Highly Selective Inhibition of Tyrosine Kinase
2 (TYK2) for the Treatment of Autoimmune Diseases: Discovery of the Allosteric Inhibitor BMS-
986165. J Med Chem 62(20):8973–8995. https://doi.org/10.1021/acs.jmedchem.9b00444
49.   Zhou Y, Li X, Shen R, et al (2022) Novel Small Molecule Tyrosine Kinase 2 Pseudokinase
Ligands Block Cytokine-Induced TYK2-Mediated Signaling Pathways. Front Immunol 13
50.   Minegishi Y, Saito M, Morio T, et al (2006) Human Tyrosine Kinase 2 Deficiency Reveals
Its Requisite Roles in Multiple Cytokine Signals Involved in Innate and Acquired Immunity.
Immunity 25:745–755. https://doi.org/10.1016/j.immuni.2006.09.009
574
575

Figure 1. Deucravacitinib inhibits IFNα-mediated STAT phosphorylation and MHC class I overexpression. (**a,b**) EndoC-βH1 cells were treated with vehicle (V) or pre-treated with the

indicated deucravacitinib concentrations for 1 h. Afterwards, cells were left untreated or treated with IFNα (1000 U/ml) in the absence or presence of deucravacitinib for 1 h. (**a**, **b**) Protein expression was measured by western blot. (**a**) Images are representative of four to six independent experiments. (**b**) Densitometry results are shown for P-STAT1 (black circles) and P-STAT2 (white circles). Values were normalised to α-tubulin, and then to the value of IFNα alone of each experiment (considered as 100%). (**c-h**) EndoC-βH1 cells were treated with vehicle (V or Veh, black circles) or pre-treated with deucravacitinib (10 [D10, soft blue circles] and 1000 nmol/l [D1000, dark blue circles]) for 1 h. Afterwards, cells were left untreated or treated with IFNα (1000 U/ml) in the absence or presence of deucravacitinib for 1–24 h (**c-f**) or 24 h (**g**, **h**). (**c-f**) Protein expression was measured by western blot. (**c**) Images are representative of three to six independent experiments. (**d-f**) Densitometry results are shown for P-STAT1 (**d**), STAT1 (**e**), and STAT2 (**f**). Values were normalised to α-tubulin, and then to the highest value of each experiment (considered as 1). (**g**, **h**) Immunocytochemistry analysis of MHC class I (red) and Hoechst 33342 (blue) upon exposure to IFNα in the absence (white bars) or presence of 1000 nmol/l deucravacitinib (dark blue bars) for 24 h. Representative images of three independent experiments (13–30 images/coverslip) (**g**) and quantification (**h**) are shown. Data are means ± SEM of three to six independent experiments. (**d-f**) *$p \leq 0.05$, **$p \leq 0.01$ and ***$p \leq 0.001$ vs Vehicle + IFNα (two-way ANOVA plus Dunnett's test). (**h**) ***$p < 0.001$ vs the respective untreated (NT) (two-way ANOVA plus Sidak's test); ###$p \leq 0.001$, as indicated by bars (two-way ANOVA plus Dunnett's test).

Figure 2. Deucravacitinib blocks IFNα- but not IFNγ-induced pathway. (a-d) EndoC-βH1 cells were treated with vehicle (V, white bars) or pre-treated with ruxolitinib (500 and 5000 nmol/l; R500 and R5000), baricitinib (500 and 5000 nmol/l; B500 and B5000), or deucravacitinib (10 and 1000 nmol/l; D10 and D1000) for 1 h. Afterwards, cells were left untreated (NT, white circles) or treated with either IFNα (1000 U/ml) **(a, c)** or IFNγ (1000 U/ml) **(b, d)** in the absence or presence of each inhibitor for 1 h. **(a-d)** Protein expression was measured by western blot. **(a, b)** Images are representative of five to six independent experiments. **(c, d)** Densitometry results are shown for P-

STAT1. Values were normalised to α-tubulin, and then to the value of IFNα (**c**) or IFNγ alone (**d**)

of each experiment (considered as 1). (**e**, **f**) EndoC-βH1 cells were transfected with a pRL-CMV

plasmid (used as internal control) plus either ISRE (**e**) or GAS (**f**) promoter reporter constructs.

After 48 h of recovery, cells were treated with vehicle (white bars) or pre-treated with ruxolitinib

(5000 nmol/l; R5000), baricitinib (5000 nmol/l; B5000), or deucravacitinib (1000 nmol/l; D1000)

for 1 h. Afterwards, cells were left untreated (NT, white circles) or treated with either IFNα (1000

U/ml) for 2 h (**e**) or IFNγ (1000 U/ml) for 24 h (**f**) in the absence or presence of each inhibitor.

Relative luciferase units (RLU) were measured by a luminescent assay. Data are means ± SEM of

three to six independent experiments. $*p \leq 0.05$, $**p \leq 0.01$ and $***p \leq 0.001$ vs the respective

untreated (NT) (one-way ANOVA plus Dunnett's test). $^{\#}p \leq 0.05$, $^{\#\#}p \leq 0.01$ and $^{\#\#\#}p \leq 0.001$ vs IFNα

(**c**, **e**) or IFNγ (**d**, **f**) (one-way ANOVA plus Dunnett's test).

Figure 3. Deucravacitinib abrogates IFNα-induced expression of ISGs but not ER stress markers. (**a-f**) EndoC-βH1 cells were treated with vehicle (white bars) or pre-treated with ruxolitinib (500 and 5000 nmol/l; R500 and R5000), baricitinib (500 and 5000 nmol/l; B500 and B5000), or deucravacitinib (10 and 1000 nmol/l; D10 and D1000) for 1 h. Afterwards, cells were

left untreated (NT) or treated with IFNα (1000 U/ml) in the absence or presence of each inhibitor

for 24 h. mRNA expression of *HLA-ABC* (**a**), *MX1* (**b**), *CXCL10* (**c**), *CHOP* (**d**), *XBP1s* (**e**), and

ATF3 (**f**) was analysed by real-time PCR, normalised to β-actin and then to the value of IFNα alone

of each experiment (considered as 1). Data are means ± SEM of three independent experiments.

**p≤0.01 and ***p≤0.001 vs the respective untreated (NT) (one-way ANOVA plus Dunnett's

test). #*p*≤0.05, ##*p*≤0.01 and ###*p*≤0.001 vs IFNα (one-way ANOVA plus Dunnett's test).

Figure 4. Pre-treatment with deucravacitinib prevents IFNα + IL-1β effects. (a) Experimental design of the pre-treatment with deucravacitinib and subsequent exposure to IFNα + IL-1β for 24 h. EndoC-βH1 cells were treated with vehicle (V, white bars) or pre-treated with deucravacitinib (10 [D10, soft blue circles] and 1000 nmol/l [D1000, dark blue circles]) for 1 h. Afterwards, cells were left untreated (NT) or treated with IFNα + IL-1β (1000 U/ml + 50 U/ml, respectively) in the absence or presence of deucravacitinib for 24 h. **(b)** Apoptosis was evaluated using Hoechst

33342/propidium iodide staining. (**c**) Caspase 3/7 activity was measured by a luminescent assay.

Results are expressed as % vehicle-treated cells in the absence of cytokines (NT). (**d-f**) Protein

expression was measured by western blot. (**d**) Images are representative of four independent

experiments. Densitometry results are shown for P-STAT1 (**e**) and P-STAT2 (**f**). Values were

normalised to α-tubulin. (**g-j**) mRNA expression of *HLA-ABC* (**g**), *CXCL10* (**h**), *MX1* (**i**), and

CHOP (**j**) was analysed by real-time PCR, normalised to β-actin and then to the value of Vehicle

treated with IFNα + IL-1β (considered as 1). (**k**, **l**) Immunocytochemistry analysis of MHC class

I (red) and Hoechst 33342 (blue) upon exposure to IFNα + IL-1β in the absence (white bars) or

presence of deucravacitinib (dark blue bars) for 24 h. Representative images of three independent

experiments (12–23 images/coverslip) (**g**) and quantification (**h**) are shown. (**j**) CXCL10 secreted

to the medium was determined by ELISA. Data are means ± SEM of three to four independent

experiments. **$p{\leq}0.01$ and ***$p{\leq}0.001$ vs the respective untreated (NT) (two-way ANOVA plus

Sidak's test). #$p{\leq}0.05$, ##$p{\leq}0.01$ and ###$p{\leq}0.001$, as indicated by bars (two-way ANOVA plus

Dunnett's test).

Figure 5. Pre-treatment with deucravacitinib prevents IFNγ + IL-1β effects. (a) Experimental design of the pre-treatment with deucravacitinib and subsequent exposure to IFNγ + IL-1β for 24 h. EndoC-βH1 cells were treated with vehicle (V, white bars) or pre-treated with deucravacitinib (10 [D10, soft blue circles] and 1000 nmol/l [D1000, dark blue circles]) for 1 h. Afterwards, cells were left untreated (NT) or treated with IFNγ + IL-1β (1000 U/ml + 50 U/ml, respectively) in the absence or presence of deucravacitinib for 24 h. **(b)** Apoptosis was evaluated using Hoechst 33342/propidium iodide staining. **(c)** Caspase 3/7 activity was measured by a luminescent assay. Results are expressed as % vehicle-treated cells in the absence of cytokines (NT). **(d-f)** Protein expression was measured by western blot. Images are representative of four independent experiments **(d)**. Densitometry results are shown for P-STAT1 **(e)** and P-STAT2 **(f)**. Values were

normalised to α-tubulin. (**g-j**) mRNA expression of *HLA-ABC* (**g**)*, CXCL10* (**h**)*, MX1* (**i**), and

CHOP (**j**)was analysed by real-time PCR, normalised to β-actin and then to the value of Vehicle

treated with IFNγ + IL-1β (considered as 1). Data are means ± SEM of three to four independent

experiments. *$p \le 0.05$, **$p \le 0.01$ and ***$p \le 0.001$ vs the respective untreated (NT) (two-way

ANOVA plus Sidak's test). #$p \le 0.05$, ##$p \le 0.01$ and ###$p \le 0.001$, as indicated by bars (two-way

ANOVA plus Dunnett's test).

Figure 6. Treatment with deucravacitinib partially reverts IFNα + IL-1β-induced changes.
(**a**) Experimental design of the pre-treatment with IFNα + IL-1β and subsequent exposure to IFNα + IL-1β in the presence of deucravacitinib for 24 h. EndoC-βH1 cells were left untreated (NT) or pre-treated with IFNα + IL-1β (1000 U/ml + 50 U/ml, respectively) for 24 h. Afterwards, cells were treated with vehicle (V, white bars) or 1000 nmol/l deucravacitinib (D1000, dark blue bars) in the absence (NT) or presence of IFNα + IL-1β for 24 h. (**b**) Apoptosis was evaluated using Hoechst 33342/propidium iodide staining. (**c**, **f-h**) mRNA expression of *HLA-ABC* (**c**), *MX1* (**f**), *CHOP* (**g**), and *CXCL10* (**h**) was analysed by real-time PCR, normalised to β-actin and then to the value of Vehicle treated with IFNα + IL-1β (considered as 1). (**d**, **e**) Protein expression was measured by western blot. (**d**) Images are representative of four independent experiments. (**e**)

671   Densitometry results are shown for STAT1. Values were normalised to α-tubulin. (**i**) CXCL10

672   secreted to the medium was determined by ELISA. Data are means ± SEM of three to four

673   independent experiments. **$p \leq 0.01$ and ***$p \leq 0.001$ vs the respective untreated (NT) (two-way

674   ANOVA plus Sidak's test). #$p \leq 0.05$, ##$p \leq 0.01$ and ###$p \leq 0.001$, as indicated by bars (two-way

675   ANOVA plus Dunnett's test).

Figure 7. Treatment with deucravacitinib partially reverts IFNγ + IL-1β-induced changes.
(**a**) Experimental design of the pre-treatment with IFNγ + IL-1β and subsequent exposure to IFNγ + IL-1β in the presence of deucravacitinib for 24 h. EndoC-βH1 cells were left untreated (NT) or pre-treated with IFNγ + IL-1β (1000 U/ml + 50 U/ml, respectively) for 24 h. Afterwards, cells were treated with vehicle (V, white bars) or 1000 nmol/l deucravacitinib (D1000, dark blue bars) in the absence (NT) or presence of IFNγ + IL-1β for 24 h. (**b**) Apoptosis was evaluated using Hoechst 33342/propidium iodide staining. (**c**, **f-h**) mRNA expression of *HLA-ABC* (**c**), *MX1* (**f**), *CHOP* (**g**), and *CXCL10* (**h**) was analysed by real-time PCR, normalised to β-actin and then to the value of Vehicle treated with IFNγ + IL-1β (considered as 1). (**d**, **e**) Protein expression was measured by western blot. (**d**) Images are representative of four independent experiments. (**e**)

686   Densitometry results are shown for STAT1. Values were normalised to α-tubulin. (**i**) CXCL10

687   secreted to the medium was determined by ELISA. Data are means ± SEM of three to four

688   independent experiments. *$p \leq 0.05$, **$p \leq 0.01$ and ***$p \leq 0.001$ vs the respective untreated (NT)

689   (two-way ANOVA plus Sidak's test). #$p \leq 0.05$ and ##$p \leq 0.01$, as indicated by bars (two-way

690   ANOVA plus Dunnett's test).