Exercise rejuvenates the skeletal muscle methylome and transcriptome in humans

Sarah Voisin[1*], Kirsten Seale[1], Macsue Jacques[1], Shanie Landen[1], Nicholas R Harvey[2,3], Larisa M Haupt[3], Lyn R Griffiths[3], Kevin J Ashton[2], Vernon G Coffey[2], Jamie-Lee M Thompson[2], Thomas M Doering[4], Malene E Lindholm[5], Colum Walsh[6], Gareth Davison[7], Rachelle Irwin[6], Catherine McBride[7], Ola Hansson[8,9], Olof Asplund[8], Aino E Heikkinen[9], Päivi Piirilä[10], Kirsi Pietiläinen[11,12], Miina Ollikainen[9,13], Sara Blocquiaux[14], Martine Thomis[14], Adam P Sharples[15], Nir Eynon[1*].

[1]Institute for Health and Sport (iHeS), Victoria University, Footscray, VIC 3011, Australia.
[2]Faculty of Health Sciences & Medicine, Bond University, Gold Coast, QLD 4226, Australia.
[3]Genomics Research Centre, Centre for Genomics and Personalised Health, School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD 4000, Australia.
[4]School of Health, Medical and Applied Sciences, Central Queensland University, Rockhampton, Queensland, Australia
[5]Department of Medicine, School of Medicine, Stanford University, Stanford, CA 94305, USA.
[6]Genomic Medicine Research Group, School of Biomedical Sciences, Ulster University, Coleraine, BT52 1SA, UK
[7]Sport and Exercise Sciences Research Institute, Ulster University, Belfast, BT15 1ED, UK
[8]Department of Clinical Sciences, Genomics, Diabetes and Endocrinology Unit, Lund University Diabetes Center, Lund University, Malmö, Sweden
[9]Institute for Molecular Medicine Finland (FIMM), Helsinki University, Helsinki, Finland
[10]Unit of Clinical Physiology, Helsinki University Hospital and University of Helsinki, Helsinki, Finland
[11]Obesity Research Unit, Research Program for Clinical and Molecular Metabolism, Faculty of Medicine, University of Helsinki
[12]HealthyWeightHub, Abdominal Center, Helsinki University Hospital and University of Helsinki, Helsinki, Finland
[13]Minerva Foundation Institute for Medical Research, Helsinki, Finland
[14]Department of Movement Sciences, Physical Activity, Sports & Health Research Group, KU Leuven, Leuven, BE 3001, Belgium
[15]Institute of Physical Performance, Norwegian School of Sport Sciences, Oslo, Norway

*Corresponding Authors:

Dr Sarah Voisin
Institute for Health and Sport (iHeS),
Victoria University, Melbourne, Australia 8001
Phone: +61 466469673
E-mail: sarah.voisin@vu.edu.au

Professor Nir Eynon
Institute for Health and Sport (iHeS),
Victoria University, Melbourne, Australia 8001
Phone: +61 399195615
Fax: +61 399199185
E-Mail: Nir.Eynon@vu.edu.au

Abstract

Exercise training prevents age-related decline in muscle function. Targeting epigenetic aging is a promising actionable mechanism and late-life exercise mitigates epigenetic aging in rodent muscle. Whether exercise training can decelerate, or reverse epigenetic aging in humans is unknown. Here, we performed a powerful meta-analysis of the methylome and transcriptome of an unprecedented number of human skeletal muscle samples (n = 3,176). We show that: 1) individuals with higher baseline aerobic fitness have younger epigenetic and transcriptomic profiles, 2) exercise training leads to significant "rejuvenation" of molecular profiles, and 3) muscle disuse "ages" the transcriptome. Higher fitness levels were associated with attenuated differential methylation and transcription during aging. Furthermore, both epigenetic and transcriptomic profiles shifted towards a younger state after exercise training interventions, while the transcriptome shifted towards an older state after forced muscle disuse. We demonstrate that exercise training targets many of the age-related transcripts and DNA methylation loci to maintain younger methylome and transcriptome profiles, specifically in genes related to muscle structure, metabolism and mitochondrial function. Our comprehensive analysis will inform future studies aiming to identify the best combination of therapeutics and exercise regimes to optimize longevity.

Keywords

Aging, Epigenetics, Exercise, Physical activity, OMIC, Transcriptome, Proteome, Methylome, Skeletal Muscle

Introduction

The United Nations has declared 2021-2030 the "Decade of Healthy Ageing" to assist the aging population in living healthier for longer. Identifying reliable aging biomarkers that can be targeted by longevity-promoting interventions is a global priority, however, requires sizable human cohorts across a broad range of ages and relevant tissues, which is both costly and time-consuming. The last two decades have seen an open science revolution with the creation of free-access repositories overflowing with molecular data from all levels of gene regulation (e.g. epigenomics, transcriptomics). These rich repositories allow for the exploration of aging mechanisms and their susceptibility to environmental stressors, even in healthy and/or young individuals, since age-related changes gradually accumulate from early life and affect organ systems years before disease manifestation[1].

Aging is associated with a loss of muscle mass and function that leads to increased adverse outcomes including falling injury, functional decline, frailty, earlier morbidity and mortality[2]. Exercise training is one of the most affordable and effective ways to promote healthy aging[3], as being physically active reduces mortality from all causes, independent of levels and changes in several established risk factors (overall diet quality, body mass index, medical history, blood pressure, triglycerides, and cholesterol)[4]. Cardiorespiratory fitness, as estimated by maximal oxygen uptake (VO_{2max}) during an exercise test, shows a strong, graded, and inverse association with overall mortality[5]. However, we have an incomplete understanding of the fundamental mechanisms by which physical activity delays the age-related decline in skeletal muscle function.

At the molecular level, aging arises from a tip in the balance between cellular damage and compensatory mechanisms[6,7]. Cells undergo constant damage, such as genomic instability, telomere attrition, epigenetic alteration, and loss of proteostasis[6]. Compensatory mechanisms, such as impaired nutrient sensing, mitochondrial

dysfunction and cellular senescence, initially mitigate the damage, but eventually, they become deleterious themselves when chronically activated. This leads to stem cell exhaustion, and altered intercellular communication, which are ultimately responsible for the physiological decline associated with aging. The effect of aging on DNA methylation patterns are so profound that machine learning has spawned highly accurate predictors of both chronological and biological age (termed "epigenetic clocks")[8]. We developed an epigenetic clock for human skeletal muscle[9] and reported widespread changes in the muscle methylome at genes involved in muscle structure and function[10]. Downstream of epigenetic processes, changes in transcriptional patterns at genes involved in central metabolic pathways and mitochondrial function have also been reported in muscle during aging[11]. Regular exercise mitigates the age-related loss of proteostasis[12,13], mitochondrial dysfunction[3,14], and stem cell exhaustion[3,15] in muscle, but there is currently limited evidence for its effect on age-related epigenetic and transcriptomic changes.

Several cross-sectional analyses found only weak associations between self-reported physical activity levels and epigenetic age in blood[16–18] or skeletal muscle[18], after adjusting for confounders such as diet. However, self-reported measures of physical activity poorly reflect actual physical activity levels, particularly in individuals with higher body fat and females who typically overestimate energy expenditure[19]. Furthermore, these studies relied solely on DNA methylation clocks to quantify age-related changes in epigenetic patterns. Epigenetic age, as measured by epigenetic clocks, is a single value that encompasses a very small portion of the aging methylome (typically a few hundred age-related DNA methylation loci, also called CpGs). Therefore, it offers a very narrow and incomplete view of the aging methylome, and exercise training may affect aging regions that are not captured by epigenetic clocks. Interestingly, a recent study found evidence that late-life exercise mitigates age-related epigenetic changes in mouse gastrocnemius muscle[20]. In this study, the authors did not use clocks but investigated all DNA methylation loci that change with age in mouse muscle and applied a direct exercise training intervention. A couple of human studies are in line with these results: resistance training was shown to offset age-related changes both in the nuclear[21,22] and mitochondrial[23] epigenome. Unfortunately, these studies had small sample sizes and have not been replicated to date, so they should be regarded as preliminary[24]. They were also restricted to resistance training in males, which does not speak for the effect of exercise training in general in males or females across the lifespan. In another, large-scale meta-analysis, age-related changes in the transcriptome showed an inverse correlation with higher cardiorespiratory fitness (CRF)[11]. However, these associations were cross-sectional and could be confounded by other environmental and lifestyle factors that co-occur with higher CRF levels (e.g. a better diet). Therefore, there is a great need for a comprehensive, integrative and robust assessment of the effects of CRF, exercise training and inactivity on age-related molecular changes in human muscle.

To address these limitations, we identified relevant published datasets from online databases that we combined with our own original data to characterise the effect of cardiorespiratory (CRF), exercise training and inactivity on human skeletal muscle aging across the methylome and transcriptome. We first compiled a list of age-related changes in 1,251 samples across 16 cohorts (DNA methylation) and 1,925 samples across 21 cohorts (mRNA expression). We then tested cross-sectional associations between aging molecular profiles and CRF; we hypothesized that individuals with higher CRF would display 'younger' profiles than expected at age-related CpGs and mRNAs. Then, we investigated directly whether exercise training could 'rejuvenate' molecular profiles in human skeletal muscle, using high-resolution longitudinal data collected from exercise training studies of

various types and durations. Finally, we tested whether inactivity could 'age' transcriptomic profiles in human skeletal muscle, using longitudinal data from forced immobilization interventions in humans. This work provides a comprehensive and integrative map of the effect of physical activity on the age-related changes in fundamental processes controlling gene expression in human muscle.

Results

Methodology overview

We describe our study design in **Fig 1**.

Figure 1. *Study overview*. *First, we performed a large-scale data mining exercise to gather all existing DNA methylation (DNAm) and mRNA expression datasets from our own lab, our network of collaborators, and open-access repositories (GEO, dbGAP, ArrayExpress). See Methods for inclusion criteria of datasets. Step 1: we identified age-related changes in DNAm and mRNA expression in human skeletal muscle by meta-analysing 1,251 samples across 16 cohorts (DNAm) and 1,925 samples across 21 cohorts (mRNA). Step 2: first, we performed a cross-sectional association between DNAm levels at age-related CpGs or expression levels at age-related mRNAs, and VO_{2max}. Then, we determined whether DNAm levels at age-related CpGs, and expression levels at age-related mRNAs changed after exercise training, and we assessed whether expression levels at age-related mRNAs changed after muscle disuse. Finally, we performed a series of OMIC integrations and pathway analyses to identify the molecular pathways affected by age, VO_{2max}, exercise training, and/or muscle disuse across both OMIC layers. Note: we only had OMIC data available at the transcriptomic level following muscle disuse. Note 2: summary statistics for the exercise- and disuse-induced mRNA changes came from two recent meta-analyses[24,25].*

First, we identified the DNA methylation loci and transcripts that change with age in human skeletal muscle. We conducted a random-effects epigenome-wide association study (EWAS) meta-analysis of age across 1,251 samples from 16 independent cohorts (Supplementary Table 1), and a random-effects transcriptome-wide association study (TWAS) meta-analysis of age across 1,925 human muscle samples from 21 independent cohorts (Supplementary Table 2). To uncover the effect of aerobic fitness and exercise training on the aging methylome and transcriptome of skeletal muscle, we then restricted our analysis to the identified age-related Differentially Methylated Positions (DMPs) and Differentially Expressed Genes (DEGs).

We first assessed whether an objective gold-standard measure of aerobic fitness (VO_{2max}) was associated with younger methylome and transcriptome profiles in skeletal muscle (**Fig 1**). At the epigenetic level, we conducted a random-effects meta-analysis of VO_{2max} across 439 samples from 5 independent cohort; at the transcriptomic level, we conducted a random-effects meta-analysis of VO_{2max} across 354 samples from 5 independent cohorts (**Table 1**). There was a reasonably large range of VO_{2max} in each of these cohorts (SD > 5 mL/min/kg, **Table 1**), which is essential to detect differences in OMIC aging between individuals with variable fitness levels.

	Cohort	Number of samples	Number of unique individuals	Age (years)	Sex (% male)	VO₂max (mL/min/kg)	Data availability
DNA methylation	Gene SMART[9,10]	234	66	32 ± 8.1	80%	48 ± 9.0	GSE151407, GSE171140
	Finnish Twin Cohort[17]	73	73	43 ± 16	38%	36 ± 8.5	Per request
	E-MTAB-11282[25]	52	13	45 ± 11	38%	23 ± 5.0	E-MTAB-11282
	EXACT	48	16	33 ± 10	100%	42 ± 7.8	Per request
	CAUSE[22]	32	16	60 ± 5.3	0%	30 ± 6.1	GSE213029
mRNA expression	GSE18732[26]	117	117	54 ± 11	69%	28 ± 9.6	GSE18732
	HERITAGE[27,28]	82	42	34 ± 14	58%	36 ± 9.0	GSE117070
	GSE44818[29]	72	12	30 ± 7.2	100%	60 ± 6.0	GSE44818
	The Malmö Prevention Study[30]	43	43	66 ± 1.5	100%	28 ± 6.4	E-CBIL-30
	MSAT[31]	39	39	36 ± 8.3	100%	52 ± 8.1	Per request

Table 1. Characteristics of the cohorts used in the cross-sectional analysis (association between age-related CpGs or mRNAs and maximal oxygen uptake (VO₂max)).

As cross-sectional analyses can be confounded by unmeasured factors (e.g. lifelong dietary patterns, socioeconomic status), we tested directly whether an exercise training program could shift OMIC profiles towards a younger state, in an interventional, longitudinal setting. We conducted a random-effects meta-analysis of DNA methylation changes following aerobic, high-intensity interval, or resistance training across 401 samples from 6 independent exercise training interventions (**Fig 1, Table 2**); we also extracted summary statistics at age-related DEGs from a published meta-analysis of transcriptomic changes induced by exercise training[32]. Finally, and to further support the causal effect of exercise training in the rejuvenation of muscle OMIC profiles, we tested whether age-related transcriptomic profiles were altered following a *decrease* in physical activity; we extracted summary statistics at age-related DEGs from a published meta-analysis following forced immobilisation protocols[33].

	Cohort	Number of samples	Number of unique individuals	Age (years)	Sex (% male)	Intervention	Dataset access
DNA methylation	Gene SMART[9,10]	196	66	32 ± 8.1	80%	4 weeks of HIIT repeated twice after a > 1 year washout, and an additional 8 weeks of HIIT	GSE151407, GSE171140
	E-MTAB-11282[25]	52	13	35 ± 11	39%	8 weeks of aerobic training	E-MTAB-11282
	EPIK[21]	48	14	45 ± 22	100%	12 weeks of resistance training repeated twice after a 12-week washout (older individuals) or 2-week immobilization (younger individuals)	Per request
	GSE114763[34]	39	8	29 ± 6	100%	7 weeks of resistance training repeated twice after a 7-week washout	GSE114763

	EpiTrain[35]	34	17	27 ± 4	44%	3 months of aerobic training	GSE60655
	CAUSE[22]	32	16	60 ± 5.3	0%	5 months of aerobic training	GSE213029
mRNA expression	ExTraMeta[32]	952	476	Meta-analysis of 28 datasets		Exercise training of various types and duration	https://www.extrameta.org
	MetaMEx[33]	250	125	Meta-analysis of 7 datasets		Forced immobilisation of various durations	https://metamex.serve.scilifelab.se/

Table 2. Participant characteristics from the cohorts in the interventional analysis (changes in levels of age-related CpGs or mRNA after exercise training or muscle disuse).

We purposely refrained from using our established MEAT clock[9], optimised to predict epigenetic age in human skeletal muscle, as most of the cohorts analysed in this paper were used to build the clock. Hence, epigenetic age estimations in these cohorts using MEAT would be inherently biased. Furthermore, the MEAT clock was trained to predict chronological (instead of biological) age and would therefore be of limited use to understand the effect of aerobic fitness and exercise training on muscle biological aging.

Skeletal muscle aging alters DNA methylation and expression of genes involved in muscle structure and metabolism

Age-related changes to the methylome are widespread yet small (typically ~1% change in methylation per decade of age); out of the 595,541 tested CpG sites, we identified 3,168 differentially methylated positions (DMPs) associated with age at FDR < 0.005 (Supplementary Table 3), 73% of which were hypomethylated (Supplementary Fig 1A). This is in concordance with our previous findings[10], and in line with recent findings at the single-cell level[36]. While DMPs were not enriched in any particular canonical pathway or expression signatures of genetic and chemical perturbations, they were overrepresented in several gene ontology terms related to muscle structure (contractile fibre, I band, supramolecular polymer and supramolecular complex; Supplementary Fig 1B), and in two human phenotype ontologies entirely consistent with musculoskeletal aging ("difficulty climbing stairs", and "muscle weakness"; Supplementary Fig 1B).

Supplementary Figure 2. ***EWAS meta-analysis of age reveals small, widespread DNA methylation changes at genes encoding structural components of muscle***. *A) Volcano plot of the magnitude (% DNA methylation change per year of age) and significance of age effects for each CpG included in the meta-analysis (pooled results of 16 independent muscle datasets); colored dots are CpGs that are differentially methylated at a false discovery rate (FDR) < 0.005; we highlighted the top hypo- (left) and hyper- (right) methylated CpG in forest plots to show the estimates and confidence intervals in each dataset. B) Overrepresentation analysis of the age-related differentially methylated positions for Gene Ontology, and Human Phenotype Ontology (HPO) from MSigDB; pathways in red are significant at an FDR < 0.005. The circular plot displays genes from the two significant HPOs ("difficulty climbing stairs", and "muscle weakness") that showed age-related alterations at the DNA methylation level. C) Dotplot of the enrichment of DMPs in marker genes for 12 cell types identified in a single-cell sequencing study of human skeletal muscle[37].*

Out of the 16,657 tested transcripts, we identified 330 differentially expressed genes (DEGs) at FDR < 0.005 (Supplementary Table 4), 68% of which were downregulated with older age (Supplementary Fig 2A). While DEGs were not enriched in any particular canonical pathway, they were overrepresented in several gene ontology

terms related to mitochondrial function and energy production (ADP and ATP metabolic processes, generation of precursor metabolites and energy, energy derivation by oxidation of organic compounds, mitochondrial protein containing complex, organelle inner membrane; Supplementary Fig 2B). Furthermore, they were also overrepresented in expression signatures of genetic and chemical perturbations (incl. genes up-regulated in differentiating myoblasts upon expression of *PPARGC1A*[38], genes differentially regulated in myoblasts with *IGF2BP2* knockdown[39], and genes that comprise the mitochondria gene module[40]). DEGs were also enriched in two human phenotype ontologies related to muscle function ("exercise intolerance", and "myoglobinuria"; Supplementary Fig 1B).

Finally, we estimated whether the age-related DNAm and mRNA changes were possibly driven by changes in cell type composition. We tested whether the DMPs and DEGs were overrepresented in twelve gene sets containing curated cluster markers for cell types identified in a single-cell sequencing study of human skeletal muscle[37]. We found no significant enrichment of DMPs in any of the cell type marker genes (Supplementary Fig 1C), suggesting that the age-related DNAm changes are not confounded by changes in cell type proportions. In contrast, DEGs were enriched for genes whose expression differs between type I and type IIa fibres (Supplementary Fig 2C), with a change in mRNA expression suggestive of an increase in type I fibre % with older age; *TPM3, PDLIM1, MYOZ2* are all markers of type I fibres and increased in expression with age, while *PKM, ENO3 and PFKM* are all markers of type IIa fibres and decreased in expression with age. DEGs also showed a trend for enrichment in marker genes of smooth muscle cells that make up the walls of blood vessels (Supplementary Fig 2C), but it is unclear whether it reflected an increase or a decrease in the proportion of smooth muscle cells, as the signal was inconsistent: 17/28 of the marker genes increased in expression, and 11/28 decreased in expression.

Supplementary Figure 2. ***TWAS meta-analysis of age reveals mRNA expression changes at genes involved in mitochondrial function and energy production***. *A) Volcano plot of the magnitude (log2 fold-change (log2FC) per year of age) and significance of age effects for each gene included in the meta-analysis (pooled results of 21 independent muscle transcriptomic datasets); coloured dots are genes that are differentially expressed at a false discovery rate (FDR) < 0.005; we highlighted the top down- (left) and up- (right) regulated genes in forest plots to show the estimates and confidence intervals in each dataset. B) Overrepresentation analysis of the age-related differentially expressed genes (DEGs) for Gene Ontology (GO), and Human Phenotype Ontology (HPO) from MSigDB; pathways in red are significant at an FDR < 0.005. The network shows the membership of the DEGs in the significant GO terms. The circular plot displays DEGs from the two significant HPOs ("exercise intolerance", and "myoglobinuria"). C) Dotplot of the enrichment of DEGs in marker genes for 11 cell types identified in a single-cell sequencing study of human skeletal muscle[37]. The heatmap displays the log2FC of the DEGs that are marker genes for smooth muscle cells.*

As the DMPs and DEGs were associated with related yet distinct ontologies, we further examined the overlap between differentially methylated genes (DMGs) and DEGs. We identified 63 genes that were altered both at the epigenetic and transcriptional levels during aging (Supplementary Fig 3A, Supplementary Table 5). This overlap between age-related changes at the epigenetic and transcriptomic levels was greater than expected by chance

alone, as DMGs were more likely to also be DEGs than non-DMGs (p-value for over-representation = 0.00011, Supplementary Fig 3B); conversely, DEGs were more likely to be DMGs than non-DEGs (p-value for over-representation = 1.7×10^{-5}, Supplementary Fig 3B).

Supplementary Figure 3. ***Overlap between age-related differentially methylated genes (DMGs) and differentially expressed genes (DEGs)****. A) The 3,168 DMPs were located in 2,622 unique genes, 63 of which were also DEGs. On the bottom graph, each dot represents a DMP-DEG pair, with the change in DNAm level (x-axis) and mRNA level (y-axis) per year of age. Note that some genes appear multiple times on the plot as multiple DMPs could be annotated to the same gene (e.g. FEZ2, USP54). B) Top, proportion of genes within DEGs and non-DEGs that are also differentially methylated with age; bottom, proportion of genes within DMGs and non-DMGs that are also differentially expressed with age. *p < 0.005 in over-representation test (Fisher's exact test).*

Aerobic fitness is associated with younger epigenetic and transcriptomic profiles in human skeletal muscle

We identified 25 age-related DMPs one age-related DEGs associated with VO_{2max} at FDR < 0.005 (Supplementary Tables 3 & 4), but the inflation of p-values suggests that we were merely underpowered to detect more associations at a high level of confidence (Supplementary Fig 4). At the age-related DMPs and DEGs most significantly associated with VO_{2max} (the points on the far right of the Q-Q plots), VO_{2max} had a "rejuvenating" effect, particularly at the epigenetic level (Supplementary Fig 4). DMPs that were hypomethylated with age tended to be hypermethylated with higher VO_{2max}, and vice versa; DEGs that were downregulated with age tended to be upregulated with VO_{2max}, and vice versa.

Supplementary Figure 4*. **Quantile-quantile plot of p-values for the effect of VO$_{2max}$, aerobic training and muscle disuse on the aging methylome and transcriptome**. The observed p-values for each age-related DMP or DMG are sorted from largest to smallest and plotted against expected values from a theoretical χ2-distribution. If some observed p-values are clearly more significant than expected under the null hypothesis, points will move towards the y-axis; green points represent DMPs or DEGs for which the effect of age contrasts with the effect of VO$_{2max}$ or exercise or disuse ("rejuvenating" effect), while black points represent DMPs or DEGs for which the effect of age is in line with the effect of VO$_{2max}$ or exercise or disuse ("pro-aging" effect).*

The "rejuvenating" effect of VO$_{2max}$ was visible when all age-related DMPs were taken into account; a strong negative correlation was noted between the effects of age and VO$_{2max}$ across all age-related DMPs (**Fig 2A**, Spearman correlation ρ = -0.39, $p < 2.2 \times 10^{-16}$, **Fig 2B**). To put the effects of VO$_{2max}$ and age into perspective, we calculated the ratio between the effect size of VO$_{2max}$ and age at the 25 age-related DMPs that were also significantly associated with VO$_{2max}$: every additional unit of VO$_{2max}$ was accompanied by a change in methylation equivalent to 1.6 years of "rejuvenation" at those specific DMPs (**Fig 2C**). It should be noted that what we term "rejuvenation" should not be taken as a literal translation to an addition of life years, but a simple comparison of magnitude in effect size between age and VO$_{2max}$.

*Figure 2. **Aerobic fitness and exercise training have similar, rejuvenating effects on muscle epigenetic profiles.** A) Pairwise correlation (Spearman) between the effect sizes of age, aerobic fitness (VO$_{2max}$) and exercise training at age-related DMPs. Each dot corresponds to one of the 3,168 age-related DMPs, and the axes represent the magnitude of effect for age, VO$_{2max}$ and exercise training. B) Unsupervised hierarchical clustering of the effect sizes of age, VO$_{2max}$ and exercise training at age-related DMPs (ordered from the most hypomethylated to most hypermethylated with age). Note that the legend is arbitrary as effect sizes were scaled to an SD of 1 for age, VO$_{2max}$ and exercise training to be comparable. C) Volcano plots of the relative effects of age and VO$_{2max}$ (top), or age and exercise training (bottom) at age-related DMPs ("Rejuvenation effect of VO$_{2max}$" = $\frac{\% \, DNAm \, change \, per \, unit \, of \, VO2max}{\% \, DNAm \, change \, per \, year \, of \, age}$; "Rejuvenation effect of exercise training" = $\frac{\% \, DNAm \, change \, after \, exercise \, training}{\% \, DNAm \, change \, per \, year \, of \, age}$); red dots are age-related DMPs that are associated with VO$_{2max}$ at FDR < 0.005.*

We observed similar yet weaker associations at the transcriptomic level (**Fig 3**); the effects of age and VO$_{2max}$ were negatively correlated when all age-related DEGs were included (**Fig 2A**, Spearman correlation ρ = -0.18, $p = 0.0013$, **Fig 2B**).

Figure 3*. **Aerobic fitness and exercise training have similar, rejuvenating effects on muscle transcriptomic profiles that contrast with the "pro-aging" effect of muscle disuse.** A) Pairwise correlation (Spearman) between the effect sizes of age, aerobic fitness (VO$_{2max}$), exercise training and muscle disuse at age-related DEGs. Each dot corresponds to one of the 330 age-related DEGs, and the axes represent the magnitude of effect for age, VO$_{2max}$, exercise training and disuse. B) Unsupervised hierarchical clustering of the effect sizes of age, VO$_{2max}$, exercise training and disuse at age-related DEGs (ordered from the most downregulated to most upregulated with age). Note that the legend is arbitrary as effect sizes were scaled to an SD of 1 for age, VO$_{2max}$, exercise training and disuse to be comparable. C) Volcano plots of the relative effects of age and VO$_{2max}$ (top), age and exercise training (middle) or age and disuse (bottom) at age-related DEGs ("Rejuvenating effect of VO$_{2max}$" = $\frac{\% \, DNAm \, change \, per \, unit \, of \, VO2max}{\% \, DNAm \, change \, per \, year \, of \, age}$; "Rejuvenating effect of training" = $\frac{\% \, DNAm \, change \, after \, exercise \, training}{\% \, DNAm \, change \, per \, year \, of \, age}$; "Rejuvenating effect of disuse" = $\frac{\% \, DNAm \, change \, after \, disuse}{\% \, DNAm \, change \, per \, year \, of \, age}$); red dots are age-related DMPs that are associated with VO$_{2max}$ at FDR < 0.005 (top) or that significantly change after muscle disuse at FDR < 0.005 (bottom).*

To further highlight the association between VO$_{2max}$ and muscle OMIC aging, we performed principal component analysis (PCA) for DMPs in the Gene SMART cohort, and for DEGs in the GSE18732 cohort. These two cohorts have the largest sample size and VO$_{2max}$ range in our study (**Table 2**). In the Gene SMART cohort, individuals clustered by VO$_{2max}$ levels both on Dimension 1 and Dimension 3 (Pearson correlation $p = 0.0011$ for PC1 and

$p = 0.0047$ for PC3), indicating that individuals of similar fitness levels show similar patterns of DNA methylation at age-related DMPs (**Fig 4A**). Similarly, in the GSE18732 cohort, individuals tended to cluster by VO$_{2max}$ levels both on Dimension 1 and Dimension 3 (Pearson correlation $p = 0.05$ for PC1 and $p = 0.0019$ for PC3), indicating that individuals of similar fitness levels have similar mRNA levels at age-related DEGs (**Fig 4B**).

__Figure 4__. Principal component analysis of individuals from the Gene SMART cohort at age-related DMPs, and individuals from the GSE18732 cohort at age-related DEGs. We used principal component analysis (PCA) to reduce dimensionality and show each individual on a two-dimensional graph. Individuals from the Gene SMART cohort (A) and from the GSE18732 cohort (B) are colored according to their baseline VO$_{2max}$ levels. Lower levels of VO$_{2max}$ are indicated by smaller circles in lighter colors, while higher levels of are indicated by larger circles in darker reds. To objectively test the clustering of individuals according to VO$_{2max}$, we ran Pearson correlations between individual coordinates on Dimension 1, Dimension 2 or Dimension 3 and VO$_{2max}$.

Exercise training rejuvenates muscle OMIC profiles while inactivity accelerates transcriptomic aging

We did not identify age-related DMPs that were significantly altered following exercise training at FDR < 0.005, but the inflation of p-values suggests that we were merely underpowered to detect associations at a high level of confidence (Supplementary Fig 4). With substantially more samples (n = 952 from 28 datasets) and therefore statistical power at the transcriptional level, we found 40 age-related DEGs that were altered following exercise training at FDR < 0.005 (Supplementary Table 3). The age-related DMPs and DEGs that were most significantly changed after exercise training were overwhelmingly green (Supplementary Fig 4), suggesting a "rejuvenating" effect of exercise training on muscle OMIC aging. DMPs that were hypomethylated with age tended to increase in methylation following exercise training, and vice versa. Similarly, DEGs that were downregulated with age increased in expression following exercise training, and vice versa. The "rejuvenating" effect of exercise training was visible when all age-related DMPs and DEGs were taken into account; we found a strong negative correlation between the effects of age and exercise training over all age-related DMPs, and over all age-related DMPs (**Fig 2A**, Spearman correlation ρ = -0.37, $p < 2.2 \times 10^{-16}$), and over all age-related DEGs (**Fig 3A**, Spearman correlation ρ = -0.38, $p = 6.7 \times 10^{-12}$). To put the effects of exercise training and age into perspective, we calculated the ratio between the change in mRNA expression following exercise training and age: exercise training led to changes in mRNA expression equivalent to 12.6 years of "rejuvenation" at the 40 DEGs altered by training (**Fig 3C**).

We observed a very pronounced effect of muscle disuse on the aging transcriptome, despite a substantially lower sample size (n = 250 from 7 datasets, **Table 2**): 68 age-related DEGs were significantly altered following forced

immobilisation at FDR < 0.005 (Supplementary Table 4, Supplementary Fig 4, **Fig 3**). DEGs that were downregulated with age tended to decrease in expression following muscle disuse, and vice versa. The "pro-aging" effect of muscle disuse was visible at the scale of the whole aging transcriptome; we found a strong positive correlation between the effects of age and immobilisation over all age-related DEGs (**Fig 3A**, Spearman correlation $\rho = 0.43$, $p < 2.2 \times 10^{-16}$). To put the effects of muscle disuse and age into perspective, we calculated the ratio between the change in mRNA expression following disuse and age: forced immobilization led to changes in mRNA expression equivalent to 31 years of "aging" at the 68 DEGs altered by disuse (**Fig 3C**).

We compared the effects of age, aerobic fitness, exercise training, and muscle disuse in an integrated, multi-contrast enrichment analysis that uses a rank-MANOVA based statistical approach[41]. We could only perform this multi-contrast analysis at the transcriptional level, as this statistical approach has not been adapted for DNA methylation data (i.e. it has not been optimised to take into account the severe bias in gene-set analysis applied to genome-wide methylation data[42,43]). This multi-contrast analysis identified mitochondrial and metabolic pathways as simultaneously inhibited by age and muscle disuse, while enhanced by aerobic fitness and exercise training (**Fig 3C**). The effects of VO_{2max} and exercise training were highly consistent both at the epigenetic and transcriptomic levels (**Fig 2 & 3**). Conversely, the effect of muscle disuse contrasted with both VO_{2max} and exercise training (**Fig 2 & 3**).

Discussion

We showed, in a large sample (> 3,200) of human muscles, that higher aerobic fitness is associated with younger epigenetic and transcriptomic profiles. In line with this, exercise training "rejuvenates" the aging muscle epigenome and transcriptome, while muscle disuse significantly accelerates transcriptomic aging. The magnitude of "rejuvenation" or "aging" effects were highly site- and gene-dependent, suggesting that exercise training can reverse specific OMIC changes occurring during normal aging in human skeletal muscle. Transcriptomic integration revealed a pronounced deterioration of mitochondrial function and energy production during aging, which is accelerated by forced immobilisation but restored by exercise training.

Although there were few age-related DMPs and DMGs significantly associated with exercise training or forced immobilisation, a close inspection of all statistical tests performed suggested that this was more likely due to a lack of statistical power than an absence of effect. In support of this, we observed a striking global "rejuvenation" effect of aerobic fitness and training, as well as a "pro-aging" effect of forced immobilisation at the transcriptomic level. Every additional unit of VO_{2max} was accompanied by a change in methylation equivalent to ~1-2 years of "rejuvenation" at a subset of age-related DMPs. This is interesting considering that adults lose about ~2.5-3 mL/min/kg of VO_{2max} per decade of age[44], partly because of primary aging (i.e. the inevitable deterioration of cellular structure and biological function, independent of disease or harmful lifestyle or environmental factors[45]), but also due to a decline in physical activity levels as we age[46]. Therefore, the rejuvenating effect of exercise and the aging effect of disuse are perhaps unsurprising, given that some of the age-related signal captured in the EWAS and TWAS meta-analysis of age reflect a decline in physical activity levels, rather than primary aging per se. This exemplifies the complex nature of aging biology, and highlights the challenge of assessing which factors mutually affect each other when causality becomes circular[47] (e.g. aging leads to a decline in physical activity/fitness levels, and a decline in physical activity/fitness levels leads to aging[48]). In a small study of older men, master athletes

showed hypomethylation in the promoter of genes involved in energy metabolism and muscle structure, compared with men who reported lifelong sedentary behaviour[49]. However, without a young control group and adjusting for confounders such as diet, this study could not distinguish the age-related changes that can be counteracted by exercise, from those that remain unaffected by even the most extreme exercise regimes.

We used a large-scale data mining approach to achieve an unprecedented sample size (> 1,200 epigenetic profiles and >1,900 transcriptomic profiles across 37 cohorts). We applied random-effects meta-analyses allowing the effects of age, fitness, exercise and disuse to vary between cohorts while maintaining the specificity of each dataset (i.e. we did not force a normalisation across datasets). We therefore detected robust changes, applicable to a broad range of individuals (sex, age, health/training status) and exercise/immobilization regimes (training type/duration). However, it is possible that specific populations (e.g. diabetics) show blunted "rejuvenation" potential from exercise training, or that specific training regimes (e.g. endurance vs resistance training) lead to "rejuvenation" effects that are entirely specific to that training regime. Furthermore, our interventional data (i.e. exercise training & muscle disuse protocols) entirely supported the cross-sectional findings (i.e. associations with aerobic fitness), suggesting that the effect of aerobic fitness on OMIC aging is due to exercise training rather than any other unmeasured environmental factor.

While we observed the "rejuvenating" effect of exercise training at both the epigenetic and transcriptomic levels, and despite a significant overlap between DMPs and DEGs, the molecular pathways affected by age were distinct between the two OMIC layers. It may be due to differences in gene coverage between the DNA methylation and mRNA arrays, or it could reflect differences in aging mechanisms at the epigenetic and transcriptomic levels. Nevertheless, the integration of all effects at the transcriptomic level clearly showed a downregulation of mitochondrial and energy metabolism pathways during aging and following muscle disuse, which was restored by aerobic fitness and exercise training. This is in line with the known beneficial effect of exercise on mitochondrial function[50]. The integration was not feasible at the DNA methylation level because a given CpG can be annotated to multiple genes, and a given gene can harbour multiple CpGs, severely biasing the statistical test for enrichment[42,43].

A major challenge in OMIC analysis is determining whether changes are due to a modification of the intrinsic profiles of the cells, or to a change in the relative proportions of different cell types in the sample. We could not directly estimate the proportions of different cell types in our samples, as deconvolution algorithms have currently not been developed for human skeletal muscle, whether at the methylation[51] or transcriptional level. We therefore tested for an enrichment of DMGs and DEGs in genes whose expression differ between muscle cell types[37]. While we found no evidence of confounding by cell type in the epigenetic analysis, some of the age-related changes in mRNA were indicative of an increase in the proportion of type I fibres (vs type II fibres). This is surprising given that age does not affect the relative proportion of different fibre types, but rather the size and distribution of fibres within the muscle[52]. Furthermore, we observed a "rejuvenation" of OMIC patterns in individuals of higher aerobic fitness levels, yet fitter individuals typically harbour greater proportions of type I fibres[53]. Therefore, it is likely that the DNA methylation and mRNA expression changes were intrinsic to muscle cells, rather than reflecting a shift in the proportions of different cell types within the muscle. To answer this question, future studies should investigate the effects of age, aerobic fitness, exercise training and muscle disuse on OMIC profiles within individual cell types using cell sorting, or single-cell methods.

We avoided using epigenetic clocks in this analysis for multiple reasons. There are only two clocks currently available that could be applied to muscle DNA methylation data, namely the Horvath pan-tissue clock[54] and the MEAT clock we recently developed for human muscle[9] (all other clocks were developed for non-muscle tissue). First, both the pan-tissue and MEAT clocks were trained to predict *chronological* age, which is a poor proxy for clinically relevant measures of biological age (this has been highlighted by others[8,55,56] and is the reason for the development of second-generation clocks, such as PhenoAge[57], GrimAge[58] and DunedinPoAm[59]). There are currently too few DNA methylation datasets with corresponding measurements of muscle function (e.g. mitochondrial function, contractile properties, etc.) to develop an epigenetic clock that would capture muscle biological age. Second, the pan-tissue clock is poorly calibrated in skeletal muscle, and most of the muscle datasets from the present study were used to generate the MEAT clock[9]. This means that epigenetic age estimations using the MEAT clock would be severely biased and unsuitable to assess the effects of fitness and exercise on epigenetic aging. Finally, until recently[60], epigenetic clocks only selected a limited number of CpGs to maximise prediction accuracy, which means they would discard information at many potentially relevant CpGs associated with age. We adopted a broader perspective to look at the entire aging methylome and examine the potential "rejuvenating" effect of exercise training on this aging trend. We were unable to assess the functional effects of the age-related OMIC changes on muscle structure, function, and metabolism, so we cannot firmly conclude that the "rejuvenating" effect of exercise training led to gains in muscle function or quality. Future studies combining OMIC profiles with genetic data to implement Mendelian Randomization analyses[61] that would determine whether OMIC aging leads to a decline in muscle function, and whether the beneficial effects of fitness and exercise training on muscle function are mediated by a reversal of OMIC aging.

In conclusion, using an unprecedented number of epigenetic and transcriptomic human muscles profiles, meta-analyses and OMIC integration, we demonstrated the power of exercise training in shifting the epigenome and transcriptome towards a younger state. We hope that this work will inspire future studies to look deeper at the mechanisms underlying this epigenetic and transcriptomic rejuvenation in muscle.

Methods

This study was a large-scale investigation of the effect of exercise training on the aging muscle methylome and transcriptome in humans. We used a wide range of bioinformatics and computational techniques (data mining, epigenome-wide association studies, transcriptome-wide association studies, random effects meta-analysis, overrepresentation analysis and multi-contrast enrichment analysis) to analyse and interpret large amounts of OMIC data in human muscle. By exploiting the power of meta-analysis, we overcome many limitations of 'omics' research in humans. Specifically, large sample sizes are required to detect changes with small effect sizes, which is the case of age[10,11] and exercise[32,33,62,63]-related changes in muscle OMIC profiles. All bioinformatics and statistical analyses were performed using the R statistical software.

Data mining

Description of muscle DNA methylation and mRNA expression datasets

First, we gathered all existing DNA methylation and mRNA expression datasets from our laboratory and our collaborators', in conjunction with public repositories, to assemble an exhaustive database of DNA methylation and mRNA expression profiles in muscle (**Fig 1,** Supplementary Tables 1 & 2). We focused exclusively on *microarray* experiments, as they are widely used, scalable (so individual datasets have larger sample sizes), and

they measure the same CpGs or transcripts across datasets (so they are straightforward to meta-analyse). We collected the methylomes of 1,251 human samples from 16 datasets, profiled on the Illumina HumanMethylation platform (27K, 450K and EPIC) (Supplementary Table 1), as well as the transcriptomes of 1,926 samples from 21 datasets, profiled on Affimetrix, Illumina, and Agilent platforms (Supplementary Table 2). For robustness, we only included datasets with > 20 samples, with an age SD > 5 years (age-associated changes cannot be detected if age is invariant). Cohorts varied in their age range, health status, ethnicity and potential treatments, so were adjusted for relevant covariates in the statistical analysis to detect age-, CRF-, exercise- and disuse-related changes that are independent of undesirable confounders (see Supplementary Tables 1 & 2 for the list of confounders adjusted in each cohort).

Pre-processing

We downloaded the raw IDAT files and pre-processed all DNA methylation datasets, except for dataset GSE50498 for which we were missing batch information (we used the already pre-processed matrix for this dataset). Details on the pre-processing steps have been previously published[10], and the preprocessing code is available on Sarah Voisin's Github account.

We downloaded the already pre-processed mRNA expression datasets, resolved any gene ID ambiguity (e.g. outdated gene names) and averaged the expression of transcripts annotated to the same EntrezID gene.

Identifying age-related changes in the muscle methylome and transcriptome

EWAS and TWAS meta-analysis of age

To determine whether exercise training can slow down/reverse OMIC aging in human muscle, we first need to know which CpGs and mRNAs change during aging, in which direction, and to what extent. Therefore, we first conducted independent EWAS or TWAS of age in each methylation or transcription dataset. Details on the EWAS pipeline are available elsewhere[10], and TWAS were conducted in a similar manner. The code for each EWAS and each TWAS is available on Sarah Voisin's Github account. Briefly, we regressed the DNAm level for each CpG (or the mRNA expression level for each transcript) against age, and adjusted the models for dataset-specific covariates known to influence DNAm or mRNA expression levels (e.g. sex, ethnicity). We then conducted a random-effects meta-analysis to pool the summary statistics at each CpG and mRNA across datasets. Not all CpGs were present in all DNAm datasets, and not all mRNAs were present in all transcriptional datasets, and restricted the analysis to the 595,541 CpG sites present in at least 10 of the 16 DNAm cohorts, and we restricted the analysis to the 16,657 genes present in at least 15 of the 21 datasets. Meta-analysis was carried out using *metaphor*[64] using "EB" (Empirical Bayes) as the residual heterogeneity estimator, 0.5 as the step length, 10000 iterations, and an accuracy of 1^{-8} in the algorithm that estimates τ^2. CpGs and mRNAs that showed a meta-analysis false discovery rate (FDR) < 0.005 were considered age-related and selected for downstream analyses.

Over-representation analysis (ORA) of ontologies (molecular pathways, human phenotypes)

To gain insights into the cellular and physiological consequences of aging on the muscle methylome and transcriptome, we tested whether genes belonging to canonical pathways (CP gene set in MSigDB), expression signatures of genetic and chemical perturbations (CGP gene set in MsigDB), gene ontology terms (GO gene set in MsigDB), and human phenotype ontologies (HPO gene set in MsigDB) were over-represented among the age-related CpGs and mRNAs. We used the *missmethyl* package for DNA methylation[42,43], and the *clusterProfiler* package[65] for transcription. Gene sets showing an FDR < 0.005 were considered significantly overrepresented.

Confounding by changes in muscle cell type proportions

We used the same ORA technique to estimate whether age-related changes in DNAm signal were potentially confounded by changes in muscle cell type proportions. We created a gene set containing markers genes for muscle cell types identified in a recent single-cell transcriptional study of human muscle[37] (marker genes for muscle endothelial cells, smooth muscle cells, pericytes, FAP cells, PCV endothelial cells, satellite cells, FBN1 FAP cells, NK cells, myeloid cells, B cells and T cells were from the "Rubenstein_skeletal_muscle" gene set in MSigDB, while marker genes for type I and type II fibers were downloaded directly from the original paper's supplementary table). Cell types showing an FDR < 0.005 were considered significantly overrepresented.

Integration of aging methylome & transcriptome

We used the same ORA technique to estimate whether there was a significant overlap between age-related changes at DNA methylation and mRNA expression levels. Age-related differentially methylated genes (DMGs) were used as a gene set in the ORA for transcription, and age-related differentially expressed genes (DEGs) were used as a gene set in the ORA for DNA methylation.

Estimating the effects of CRF, exercise training and muscle disuse on the aging methylome and transcriptome

We restricted the analysis to the age-associated CpGs and mRNAs identified in Step 1.

Cardiorespiratory fitness

We focused on the muscle datasets for which information on baseline maximal oxygen uptake (VO_{2max}) was available (**Table 1**). We only included datasets with baseline VO_{2max} SD > 5 mL/min/kg (CRF-associated changes cannot be detected if there is no variability in baseline CRF between participant). VO_{2max}, measured during a graded exercise test, is considered the gold-standard measurement of CRF.

We applied the same EWAS and TWAS meta-analysis pipeline described previously but regressing DNAm or mRNA expression levels against VO_{2max}. We then performed meta-analysis for each aging CpG and transcript across datasets and adjusted for multiple testing. Aging CpGs or mRNAs that were associated with VO_{2max} at FDR < 0.005 were considered significant.

Exercise training

For DNA methylation, we focused on the muscle datasets that came from exercise training studies (**Table 2**). We included all types of exercise training interventions (aerobic training, high-intensity interval training, resistance training) as our aim was to test whether exercise in general could counteract the effect of age on aging OMIC profiles. We applied the same EWAS meta-analysis pipeline described previously but looking at changes in DNAm levels after exercise training (i.e. regressing DNAm levels against Timepoint (PRE/POST training). Then, we performed the meta-analysis for each aging CpG across datasets and adjusted for multiple testing. Aging CpGs whose DNAm levels changed following exercise training at FDR < 0.005 were considered significant.

For mRNA expression, we extracted summary statistics at age-related mRNAs from a recently published meta-analysis of exercise-induced transcriptional changes[32]. For each transcript, we used the summary statistics from the model selected by the authors (all data came from the meta_analysis_input.RData file uploaded by the authors on their Github page)

Muscle disuse

There were no available muscle immobilisation studies that profiled DNA methylation patterns in human muscle, so we could not estimate the effect of muscle disuse on age-related DNAm patterns.

For mRNA expression, we extracted summary statistics at age-related mRNAs from a published meta-analysis of disuse-induced transcriptional changes[33]. We used summary statistics sent by the authors upon correspondence with them.

Transcriptomic integration of age, CRF, exercise and disuse

To contrast the effects of age, CRF, exercise training and muscle disuse, we used multi-contrast enrichment analysis as implemented in the *mitch* package[41]. As recommended in the package, we first created a score to represent the importance of the differential gene expression for each transcript and each contrast:

$$Score = sign(log_2FC) \times -log_{10}(pvalue)$$

For example, the *FEZ2* gene decreased in expression with age at a magnitude of log_2FC = -0.12 per year of age at a p-value of 1.24×10^{-10}. The age score for *FEZ2* was therefore $-1 \times -log_{10}(1.24 \times 10^{-10}) = -9.9$.

We then performed multi-contrast enrichment analysis using the same gene sets as in the ORA analysis described above (canonical pathways, expression signatures of genetic and chemical, gene ontology terms, and human phenotype ontologies), using the default parameters in the mitch_calc() function. Gene sets showing an FDR < 0.005 were considered significant.

Visualisation tools

Pairwise correlation plots using Spearman correlations were graphed using the *GGally* package, heatmaps were graphed using the *pheatmap* package after scaling the effect sizes for age, VO_{2max}, exercise and/or disuse, and forest plots were graphed using the *metafor* package.

Acknowledgements

This work was supported by Sarah Voisin's National Health & Medical Research Council (NHMRC) Early Career Research Fellowship (APP11577321) and by Nir Eynon's NHMRC Career Development Fellowship (APP1140644). The Gene SMART and LITER studies were both supported by the Collaborative Research Network for Advancing Exercise and Sports Science (201202) scheme awarded to VGC and KJA from the Department of Education and Training, Australia. Mr Nicholas Harvey was supported by a PhD stipend also provided by Bond University CRN-AESS. This research was also supported by infrastructure purchased with Australian Government EIF Super Science Funds as part of the Therapeutic Innovation Australia - Queensland Node project (LRG). We also greatly acknowledge Erika Guzman at the ATGC/IHBI/QUT for performing the HMEPIC assays in the LITER study. The EPIK study was supported by the Research Foundation Flanders (FWO G.0898.15).

The authors declare that they have no conflict of interest.

References

1.   Belsky, D. W. *et al.* Quantification of biological aging in young adults. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E4104–E4110 (2015).

2.   Cruz-Jentoft, A. J. & Sayer, A. A. Sarcopenia. *Lancet* **393**, 2636–2646 (2019).

3.   Cartee, G. D., Hepple, R. T., Bamman, M. M. & Zierath, J. R. Exercise Promotes Healthy Aging of Skeletal Muscle. *Cell Metab.* **23**, 1034–1047 (2016).

4.   Mok, A., Khaw, K.-T., Luben, R., Wareham, N. & Brage, S. Physical activity trajectories and mortality: population based cohort study. *BMJ* **365**, l2323 (2019).

5.   Laukkanen, J. A. *et al.* Cardiovascular Fitness as a Predictor of Mortality in Men. *Arch. Intern. Med.* **161**, 825–831 (2001).

6.   López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The Hallmarks of Aging. *Cell* **153**, 1194–1217 (2017).

7.   Thuault, S. Reflections on aging research from within the National Institute on Aging. *Nat. Aging* **1**, 14–18 (2021).

8.   Bell, C. G. *et al.* DNA methylation aging clocks: challenges and recommendations. *Genome Biol.* **20**, 249 (2019).

9.   Voisin, S. *et al.* An epigenetic clock for human skeletal muscle. *J. Cachexia. Sarcopenia Muscle* **11**, 887–898 (2020).

10.   Voisin, S. *et al.* Meta-analysis of genome-wide DNA methylation and integrative omics of age in human skeletal muscle. *J. Cachexia. Sarcopenia Muscle* **12**, 1064–1078 (2021).

11.   Su, J. *et al.* A novel atlas of gene expression in human skeletal muscle reveals molecular changes associated with aging. *Skelet. Muscle* **5**, 35 (2015).

12.   Fernando, R., Drescher, C., Nowotny, K., Grune, T. & Castro, J. P. Impaired proteostasis during skeletal muscle aging. *Free Radic. Biol. Med.* **132**, 58–66 (2019).

13.   Ubaida-Mohien, C. *et al.* Physical Activity Associated Proteomics of Skeletal Muscle: Being Physically Active in Daily Life May Protect Skeletal Muscle From Aging . *Frontiers in Physiology* **10**, 312 (2019).

14.   Short, K. R. *et al.* Decline in skeletal muscle mitochondrial function with aging in humans. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 5618 LP – 5623 (2005).

15.   Sousa-Victor, P., García-Prat, L., Serrano, A. L., Perdiguero, E. & Muñoz-Cánoves, P. Muscle stem cell aging: regulation and rejuvenation. *Trends Endocrinol. Metab.* **26**, 287–296 (2015).

16.   Quach, A. *et al.* Epigenetic clock analysis of diet, exercise, education, and lifestyle factors. *Aging (Albany. NY).* **9**, 419–446 (2017).

17.   Sillanpää, E. *et al.* Blood and skeletal muscle ageing determined by epigenetic clocks and their associations with physical activity and functioning. *Clin. Epigenetics* **13**, 110 (2021).

18.   Sillanpää, E. *et al.* Leisure-time physical activity and DNA methylation age—a twin study. *Clin.*

Epigenetics **11**, 12 (2019).

19. Prince, S. A. *et al.* A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int. J. Behav. Nutr. Phys. Act.* **5**, 56 (2008).

20. Murach, K. A. *et al.* Late-life exercise mitigates skeletal muscle epigenetic aging. *Aging Cell* **21**, e13527 (2022).

21. Blocquiaux, S. *et al.* Recurrent training rejuvenates and enhances transcriptome and methylome responses in young and older human muscle. *JCSM Rapid Commun.* **5**, 10–32 (2022).

22. Gorski, P. P. *et al.* Aerobic exercise training resets the human skeletal muscle methylome 10 years after breast cancer treatment and survival. *FASEB J.* **37**, e22720 (2023).

23. Ruple, B. A. *et al.* Resistance training rejuvenates the mitochondrial methylome in aged human skeletal muscle. *FASEB J.* **35**, e21864 (2021).

24. Fanelli, D., Costas, R. & Ioannidis, J. P. A. Meta-assessment of bias in science. *Proc. Natl. Acad. Sci.* **114**, 3714 LP – 3719 (2017).

25. Garcia, L. A. *et al.* Can Exercise Training Alter Human Skeletal Muscle DNA Methylation? *Metabolites* **12**, (2022).

26. Gallagher, I. J. *et al.* Integration of microRNA changes in vivo identifies novel molecular features of muscle insulin resistance in type 2 diabetes. *Genome Med.* **2**, 9 (2010).

27. Clarke, K. *et al.* The Role of Eif6 in Skeletal Muscle Homeostasis Revealed by Endurance Training Co-expression Networks. *Cell Rep.* **21**, 1507–1520 (2017).

28. Takeshita, L. Y. *et al.* Genomics and transcriptomics landscapes associated to changes in insulin sensitivity in response to endurance exercise training. *Sci. Rep.* **11**, 23314 (2021).

29. Rowlands, D. S. *et al.* Protein-leucine ingestion activates a regenerative inflammo-myogenic transcriptome in skeletal muscle following intense endurance exercise. *Physiol. Genomics* **48**, 21–32 (2015).

30. De, S., Zhang, Y., Garner, J. R., Wang, S. A. & Becker, K. G. Disease and phenotype gene set analysis of disease-based gene expression in mouse and human. *Physiol. Genomics* **42A**, 162–167 (2010).

31. Oskolkov, N. *et al.* High-throughput muscle fiber typing from RNA sequencing data. *Skelet. Muscle* **12**, 16 (2022).

32. Amar, D. *et al.* Time trajectories in the transcriptomic response to exercise - a meta-analysis. *Nat. Commun.* **12**, 3471 (2021).

33. Pillon, N. J. *et al.* Transcriptomic profiling of skeletal muscle adaptations to exercise and inactivity. *Nat. Commun.* **11**, 470 (2020).

34. Seaborne, R. A. *et al.* Human Skeletal Muscle Possesses an Epigenetic Memory of Hypertrophy. *Sci.*

Rep. **8**, 1898 (2018).

35.   Lindholm, M. E. *et al.* An integrative analysis reveals coordinated reprogramming of the epigenome and the transcriptome in human skeletal muscle after training. *Epigenetics* **9**, 1557–1569 (2014).

36.   Tarkhov, A. E. *et al.* Nature of epigenetic aging from a single-cell perspective. *bioRxiv* 2022.09.26.509592 (2022). doi:10.1101/2022.09.26.509592

37.   Rubenstein, A. B. *et al.* Single-cell transcriptional profiles in human skeletal muscle. *Sci. Rep.* **10**, 229 (2020).

38.   Mootha, V. K. *et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat. Genet.* **34**, 267–273 (2003).

39.   Boudoukha, S., Cuvellier, S. & Polesskaya, A. Role of the RNA-binding protein IMP-2 in muscle cell motility. *Mol. Cell. Biol.* **30**, 5710–5725 (2010).

40.   Wong, D. J. *et al.* Revealing targeted therapy for human cancer by gene module maps. *Cancer Res.* **68**, 369–378 (2008).

41.   Kaspi, A. & Ziemann, M. mitch: multi-contrast pathway enrichment for multi-omics and single-cell profiling data. *BMC Genomics* **21**, 447 (2020).

42.   Phipson, B., Maksimovic, J. & Oshlack, A. missMethyl: an R package for analyzing data from Illumina's HumanMethylation450 platform. *Bioinformatics* **32**, 286–288 (2016).

43.   Maksimovic, J., Oshlack, A. & Phipson, B. Gene set enrichment analysis for genome-wide DNA methylation data. *Genome Biol.* **22**, 173 (2021).

44.   Rapp, D., Scharhag, J., Wagenpfeil, S. & Scholl, J. Reference values for peak oxygen uptake: cross-sectional analysis of cycle ergometry-based cardiopulmonary exercise tests of 10 090 adult German volunteers from the Prevention First Registry. *BMJ Open* **8**, e018697 (2018).

45.   Holloszy, J. O. The Biology of Aging. *Mayo Clin. Proc.* **75**, S3–S9 (2000).

46.   Hallal, P. C. *et al.* Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* **380**, 247–257 (2012).

47.   Cohen, A. A. *et al.* A complex systems approach to aging biology. *Nat. Aging* **2**, 580–591 (2022).

48.   Booth, F. W., Laye, M. J. & Roberts, M. D. Lifetime sedentary living accelerates some aspects of secondary aging. *HIGHLIGHTED Top. Physiol. Pathophysiol. Phys. Inact. J Appl Physiol* **111**, 1497–1504 (2011).

49.   Sailani, M. R. *et al.* Lifelong physical activity is associated with promoter hypomethylation of genes involved in metabolism, myogenesis, contractile properties and oxidative stress resistance in aged human skeletal muscle. *Sci. Rep.* **9**, 3272 (2019).

50.   Goh, J., Wong, E., Soh, J., Maier, A. & Kennedy, B. Targeting the Molecular & Cellular Pillars of

Human Aging with Exercise. *FEBS J.* (2021). doi:10.1111/febs.16337

51. Zhu, T. *et al.* A pan-tissue DNA methylation atlas enables in silico decomposition of human tissue methylomes at cell-type resolution. *Nat. Methods* **19**, 296–306 (2022).

52. Deschenes, M. R. Effects of aging on muscle fibre type and size. *Sports Med.* **34**, 809–824 (2004).

53. Stuart, C. A. *et al.* Slow-twitch fiber proportion in skeletal muscle correlates with insulin responsiveness. *J. Clin. Endocrinol. Metab.* **98**, 2027–2036 (2013).

54. Horvath, S. DNA methylation age of human tissues and cell types. *Genome Biol.* **14**, R115 (2013).

55. Zhang, Q. *et al.* Improved precision of epigenetic clock estimates across tissues and its implication for biological ageing. *Genome Med.* **11**, 54 (2019).

56. Field, A. E. *et al.* DNA Methylation Clocks in Aging: Categories, Causes, and Consequences. *Molecular Cell* **71**, 882–895 (2018).

57. Levine, M. E. *et al.* An epigenetic biomarker of aging for lifespan and healthspan. *Aging (Albany. NY).* **10**, 573–591 (2018).

58. Lu, A. T. *et al.* DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging (Albany. NY).* **11**, 303–327 (2019).

59. Belsky, D. W. *et al.* Quantification of the pace of biological aging in humans through a blood test, the DunedinPoAm DNA methylation algorithm. *Elife* **9**, (2020).

60. Higgins-Chen, A. T. *et al.* A computational solution for bolstering reliability of epigenetic clocks: implications for clinical trials and longitudinal tracking. *Nat. Aging* **2**, 644–661 (2022).

61. Relton, C. L. & Davey Smith, G. Two-step epigenetic Mendelian randomization: a strategy for establishing the causal role of epigenetic processes in pathways to disease. *Int. J. Epidemiol.* **41**, 161–176 (2012).

62. Voisin, S., Eynon, N., Yan, X. & Bishop, D. J. Exercise training and DNA methylation in humans. *Acta Physiol.* **213**, 39–59 (2015).

63. Jacques, M. *et al.* Epigenetic changes in healthy human skeletal muscle following exercise- a systematic review. *Epigenetics* 1–16 (2019). doi:10.1080/15592294.2019.1614416

64. Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Softw.* **36**, 1–48 (2010).

65. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innov.* **2**, 100141 (2021).

23