# The *shh* limb enhancer is activated in patterned limb regeneration but not in hypomorphic limb regeneration in *Xenopus laevis*

Reimi Tada[1,*], Takuya Higashidate[2,*], Takanori Amano[3], Shoma Ishikawa[1], Chifuyu Yokoyama[1], Saki Nara[1], Koshiro Ishida[1], Akane Kawaguchi[4,#], Haruki Ochi[5], Hajime Ogino[6], Nayuta Yakushiji-Kaminatsui[7], Joe Sakamoto[8], Yasuhiro Kamei[8,9], Koji Tamura[2], and Hitoshi Yokoyama[1‡].

Affiliation: [1]Department of Biochemistry and Molecular Biology
Faculty of Agriculture and Life Science, Hirosaki University
3 Bunkyo-cho, Hirosaki, Aomori 036-8561, Japan.
[2]Department of Ecological Developmental Adaptability Life Sciences,
Graduate School of Life Sciences, Tohoku University,
Aramaki-Aza-Aoba 6-3, Aoba-ku, Sendai 980-8578, Japan.
[3]Next Generation Human Disease Model Team
RIKEN BioResource Research Center
3-1-1 Koyadai, Tsukuba, Ibaraki, 305-0074, Japan
[4]Graduate School of Biological Sciences, Nara Institute of Science and Technology (NAIST), Ikoma, Nara, Japan.
[5]Institute for Promotion of Medical Science Research, Faculty of Medicine, Yamagata University, Yamagata 990-9585, Japan.
[6]Amphibian Research Center / Graduate School of Integrated Sciences for Life, Hiroshima University, 1-3-1 Kagami-yama, Higashi-Hiroshima, Hiroshima 739-8526, Japan.
[7]RIKEN Center for Integrative Medical Sciences (IMS)
1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama 230-0045, Japan
[8] Laboratory for Biothermology, National Institute for Basic Biology, Myodaiji, Okazaki, Aichi 444-8585, Japan
[9]Department of Basic Biology in the School of Life Science of the Graduate University for Advanced Studies (SOKENDAI), Okazaki,

Aichi 444-8585, Japan

#Present address: Molecular Life History Laboratory, Department of

Genomics and Evolutionary Biology, National Institute of Genetics, Japan

*These authors contributed equally to this work.

Running Title:  *shh* enhancer in limb regeneration

Key words: limb regeneration; *shh*; long-range enhancer; *Xenopus*; regenerative capacity

Summary statement: We visualized the activity of a limb-specific enhancer of *shh* in the

regenerating limb of an amphibian, *Xenopus*: it correlates with the capacity for

regeneration of a well-patterned limb.

‡Author for correspondence:

Hitoshi Yokoyama

Department of Biochemistry and Molecular Biology

Faculty of Agriculture and Life Science, Hirosaki University

3 Bunkyo-cho, Hirosaki, Aomori 036-8561, Japan.

E-mail:  yokoyoko@hirosaki-u.ac.jp

Phone:  +81-172-39-3773

FAX:  +81-172-39-3773

ORCID ID

Reimi Tada: 0000-0003-4474-1372

Takanori Amano: 0000-0002-0213-4745

Shoma Ishikawa: 0000-0002-6688-0288

Chifuyu Yokoyama: 0000-0002-7476-7496

Saki Nara: 0000-0003-2565-8944

Koshiro Ishida: 0000-0001-6443-4033

Haruki Ochi: 0000-0002-0088-581X

Hajime Ogino: 0000-0003-2707-2330

Nayuta Yakushiji-Kaminatsui: 0000-0001-7324-1161

Yasuhiro Kamei: 0000-0001-6382-1365

Koji Tamura: 0000-0002-1956-2826

Hitoshi Yokoyama: 0000-0003-4821-9506

**ABSTRACT**

*Xenopus* young tadpoles regenerate a limb with the anteroposterior (AP) pattern, but metamorphosed froglets regenerate a hypomorphic limb after amputation. The key gene for AP patterning, *shh*, is expressed in a regenerating limb of the tadpole but not in that of the froglet. Genomic DNA in the *shh* limb-specific enhancer, MFCS1 (ZRS), is hypermethylated in froglets but hypomethylated in tadpoles: *shh* expression may be controlled by epigenetic regulation of MFCS1. Is MFCS1 specifically activated for regenerating the AP-patterned limb? We generated transgenic *Xenopus laevis* lines that visualize the MFCS1 enhancer activity with a GFP reporter. The transgenic tadpoles showed GFP expression in *hoxd13*- and *shh*-expressing domains of developing and regenerating limbs, whereas the froglets showed no GFP expression in the regenerating limbs despite having *hoxd13* expression. Genome sequence analysis and co-transfection assays using cultured cells revealed that Hoxd13 can activate *Xenopus* MFCS1. These results suggest that MFCS1 activation correlates with regeneration of AP-patterned limbs and that re-activation of epigenetically inactivated MFCS1 would be crucial to confer the ability to non-regenerative animals for regenerating a properly patterned limb.

## INTRODUCTION

Among tetrapods, only urodele amphibians (newts and salamanders) can regenerate an amputated limb completely throughout their life cycles (Han et al., 2005; Joven et al., 2019; Otsuki and Tanaka, 2022; Agata and Inoue, 2012). In the anuran amphibian *Xenopus laevis*, a tadpole can completely regenerate its limbs prior to the onset of metamorphosis, but it loses regenerative capacity gradually as metamorphosis proceeds (Dent, 1962; Muneoka et al., 1986). The froglet, a young post-metamorphosis adult, regenerates only an unbranched cartilaginous spike-like structure after limb amputation (Dent, 1962; reviewed in Suzuki et al., 2006): the spike lacks the pattern formation along the anteroposterior (AP) axis. In the limb formation process, *sonic hedgehog* (*shh*) is expressed in the posterior part of the limb bud and serves as the key gene for the AP axis formation (Riddle et al., 1993; Harfe et al., 2004). *shh* is also expressed in the posterior part of a regenerating limb (blastema) in urodeles (Imokawa and Yoshizato, 1997; Torok et al., 1999). In *Xenopu*s, *shh* is expressed in the posterior part of the limb blastema of a tadpole (Endo et al., 1997) but is not expressed in the blastema of a froglet limb (Endo et al., 2000; Yakushiji et al., 2007). Treatment with a Hedgehog inhibitor results in hypomorphic limb regeneration in both a *Xenopus* tadpole and an axolotl (Roy and Gardiner, 2002; Satoh et al., 2006). These results suggest that *shh* re-expression correlates with the limb regenerative capacity: whether a limb regenerates with the AP pattern or not.

Genetic studies in mice have revealed that a long-range cis-acting enhancer called MFCS1 (mammal-fish-conserved-sequence 1), also called ZRS (ZPA regulating sequence) or Rr29, regulates limb-specific *Shh* expression (Lettice et al., 2003; Sagai et al., 2004). In transgenic (Tg) mouse embryos, MFCS1 with a minimal promoter drives *lacZ* expression in the posterior part of a limb bud (Lettice et al., 2003). Depletion of a large part of this sequence in a mouse results in limb-specific loss of *shh* expression and in loss of the AP polarity of a limb (Sagai et al., 2005). The activity of MFCS1 is positively regulated by Hoxd13 (5' Hoxd) and ETS transcription factors (ETS1 and GABPα), downstream factors of FGF signaling in a mouse limb bud that directly bind to MFCS1 (Capellini et al., 2006; Lettice et al., 2012). In amphibian limbs, the methylation level of genomic DNA in MFCS1 is relatively high in the limbs (both intact and

regenerating limbs) of a *Xenopus* froglet, while the methylation level is low to medium

in the limbs of a *Xenopus* tadpole, an axolotl and a Japanese newt (Yakushiji et al., 2007).

These findings strongly suggest that epigenetic inactivation of MFCS1 leads to the

shutdown of *shh* expression that results in hypomorhic limb regeneration without the AP

pattern in a froglet. The importance of MFCS1 for limb regeneration is further supported

by the results of partial deletion of the MFCS1 sequence by CRISPR/Cas9 in an Iberian

ribbed newt: the partial deletion resulted in reduction in *shh* expression and reduction in

the number of digits in limb regeneration (Suzuki et al., 2018). However, the spatio-

temporal activation pattern of MFCS1 has not been visualized in regenerating limbs of

amphibians. Thus, it is still unknown whether MFCS1 is activated in regenerating limbs

of amphibians.

Here, we succeeded in visualizing and monitoring the spatio-temporal pattern of

MFCS1 activation in regenerating limbs as well as in developing limbs by establishing

Tg *Xenopus laevis* lines carrying a GFP reporter linked to MFCS1 (MFCS1-GFP).

Expression patterns of MFCS1-GFP were overlapped with those of *shh* both in limb

development and limb regeneration. MFCS1-GFP expression was specifically localized

in the regenerating limb of a young tadpole. The pattern of MFCS1-GFP expression

coincided with the expression pattern of *hoxd13* in both developing and regenerating

tadpole limbs. Furthermore, genome sequence analysis and luciferase assays *in vitro*

suggested a positive regulation of *Xenopus* MFCS1 by Hoxd13. However, despite *hoxd13*

expression, MFCS1-GFP expression was not found in the regenerating froglet limb. We

revealed that the DNA methylation level of exogenous MFCS1 of the transgene as well

as that of endogenous MFCS1 in the regenerating limb were higher in the froglet than in

the tadpole. Our results suggest that MFCS1 is activated in limb regeneration with the AP

pattern but is inactivated in hypomorphic limb regeneration after metamorphosis when

MFCS1 has a higher DNA methylation status than that in tadpole stages.

**RESULTS AND DISCUSSION**

**MFCS1-GFP expression localizes in the posterior mesenchyme of a developing limb**

**bud**

To visualize the MFCS1 activity by the GFP reporter, we generated a transgene construct

containing GFP immediately downstream of a *Xenopus shh* promoter sequence

(Yakushiji et al., 2007) (Fig. 1A). At the same time, an approximately 3.2-kbp *Xenopus*

MFCS1 sequence was cloned and inserted upstream of the *shh* promoter in the transgene

construct (Fig. 1A). We generated transgenic *X. laevis* containing this transgene named

"*Xla.Tg(Xtr.shh:GFP)^{Tamura}*" (according to the Xenbase nomenclature guideline). For

simplicity and for highlighting the involvement of the limb-specific enhancer, MFCS1,

we hereafter refer to this Tg as MFCS1-GFP. We observed the GFP pattern in G0 founder

animals. Since *shh* is most prominently expressed in the posterior part of a limb bud at

stage 51-52 (Endo et al., 1997; Fig. 1B), we firstly focused on limb buds at stage 51-52

(Nieuwkoop and Faber, 1994). GFP was found in the posterior part of the limb bud in at

least eleven independent G0 individuals and the fluorescence gradually diminished after

stage 52. Thus, we considered that the pattern of MFCS1-GFP expression reflects the

activity of the MFCS1 enhancer. We established two independent lines by mating selected

GFP-expressing founder animals with wild-type frogs (Table S1). About 50% of the N1

individuals of line 2 as well as the N1 and N2 individuals of line 1 were GFP-positive in

limb buds, presumably being inherited in a Mendelian manner. Consistent with G0

animals, limb buds of both line 1 and line 2 at stage 51-52 showed GFP expression in the

posterior part (Fig. 1C, C' and Fig. S1A, A'). In the following experiments, we used GFP-

positive individuals of line 1 (N1 or N2) and line 2 (N1).

Confocal imaging revealed that GFP was localized specifically in the posterior

mesenchyme of the limb bud at stage 51-52 (Fig. 1D, E and Fig. S1B, C). In limb

development, GFP was firstly observed at stage 50 around the apical tip of a limb bud

(Fig. 1F, F' and Fig. S1D, D'). This observation is consistent with the initial weak *shh*

expression detected at stage 50 by whole mount *in situ* hybridization (Endo et al., 1997).

GFP expression became strongest at stages 51 and 52 and the fluorescence was more

posteriorly polarized by stage 52 (Fig. 1G, G', H, H' and Fig. S1E, E', F, F'). At stage 53,

GFP was found in the distal-posterior part of the limb bud (Fig. 1I, I' and Fig. S1G, G')

and the fluorescence was weaker than that at stage 52. Finally GFP expression was almost

undetectable at stage 54 (Fig. 1J, J' and Fig. S1H, H'). These observations were, in

principle, consistent with the spatio-temporal *shh* expression patterns (Endo et al., 1997),

while the domain of GFP expression appeared to be broader than that of *shh* expression.

We concluded that the MFCS1-GFP expression pattern in the Tg lines mostly visualizes

the activity of the endogenous MFCS1 enhancer in a developing limb bud.

**MFCS1-GFP expression predicts the AP duplication of a limb bud**

To further examine whether MFCS1-GFP expression correlates with limb morphogenesis

along the AP axis, we conducted a 180° rotation experiment on limb buds of Tg (line 1)

tadpoles (Fig. 2A). Cameron and Fallon (1977) found limb duplication along the AP axis

after rotation of the distal limb bud tips 180° on the proximo-distal axis at stage 51-53 in

*X. laevis*. Endo et al. (1997) repeated this experiment at stage 53 and found that ectopic

*shh* expression was induced in the posterior region of the graft and stump. To test the

correlation between MFCS1 enhancer activity and limb duplication along the AP axis,

we repeated this experiment and examined the GFP expression pattern (Fig. 2). Consistent

with the preceding results in control limb buds, in which the amputated distal tip of each

limb bud was returned to the stump in the original orientation, showing only faint ectopic

*shh* expression (Endo et al., 1997), the majority of the control limb buds (ten of 12

samples) without rotation did not show GFP expression and did not result in limb

duplication (Fig. 2B, C). In contrast, the majority of 180° rotated limb buds (eight of 11

samples) showed ectopic induction of GFP expression in the posterior part of the distal

limb bud tip (graft) and finally formed duplicated limbs along the AP axis (the number

of digits in a limb being more than 5) (Fig. 2D-G). These results (summarized in Table

S2) suggest that ectopic induction of MFCS1 activity is closely related to ectopic *shh*

expression and subsequent limb duplication along the AP axis, further suggesting that

MFCS1-GFP expression faithfully reflects the endogenous *shh* enhancer activity in a

limb.

**MFCS1 is activated in a blastema that regenerates a limb with the AP pattern.**

To examine whether MFCS1 is activated in limb regeneration with the AP pattern, we

amputated a stage 54 limb bud at the level of the presumptive ankle: this amputation

results in almost complete limb regeneration along the AP axis (Muneoka et al., 1986).

By this stage, *shh* expression and MFCS1-GFP expression in an intact limb bud have

become almost undetectable by *in situ* hybridization and observation under a fluorescence

stereomicroscope, respectively (Fig. 1 and Fig. S1) (Endo et al., 1997). At 4 days post

amputation (dpa), GFP was observed broadly in the regenerating blastema (Fig. 3R and

215 Fig. S2B-B''), indicating that MFCS1 is re-activated in the blastema that regenerates a

216 well-patterned limb.

217   MFCS1-GFP expression was distributed throughout the blastema and did not

218 appear to be localized posteriorly. It became stronger at 6 dpa (Fig. S2C-C'') and the

219 expression gradually diminished after the blastema showed a pallet shape and became

220 undetectable at 10 dpa (Fig. 3S, T and Fig. S2). *shh* expression was detected in the

221 posterior part of a limb bud at 4 dpa by *in situ* hybridization (Fig. 3E). Based on these

222 results, we judged that MFCS1-GFP expression is distributed broadly in regenerating

223 blastemas and that the temporal expression pattern of MFCS1-GFP coincidences with

224 that of *shh* expression, while the spatial MFCS1-GFP expression pattern is wider than the

225 *shh*-expressing domain. MFCS1-GFP signal in the regenerating blastema of a young

226 tadpole suggests that MFCS1 is re-activated in a regenerating limb with the AP pattern

227 formation when *shh* is expressed: *shh* expression in the regenerating limb as well as in

228 the developing limb seems to be regulated by the common enhancer, MFCS1.

230 **MFCS1-GFP expression pattern correlates with the pattern of *hoxd13* expression in**

231 **a tadpole**

232 Hoxd13 and FGF downstream factors, ETS transcription factors, positively regulate the

233 MFCS1 enhancer in mice (Capellini et al., 2006; Lettice et al., 2012). We examined *dusp6*

234 (*mkp3*) expression and *hoxd13* expression as indicators of positive upstream regulators

235 of MFCS1. *dusp6* expression is induced by FGF signaling (Eblaghie et al., 2003;

236 Kawakami et al., 2003) and it seems to reflect the activity of downstream factors of FGF

237 signaling, especially ETS transcription factors (Ekerot et al., 2008). We chose a stage 51

238 limb bud and a 4 dpa blastema of a tadpole in which the limb bud was amputated at the

239 presumptive ankle level at stage 54 as representatives of developing and regenerating

240 limbs, respectively: both initiate prominent *shh* expression in the posterior side (Fig. 3E,

241 I). In a developing limb bud, *dusp6* is expressed broadly in a limb bud and strong

242 expression is seen especially in the apical mesenchyme (Fig. 3K). *hoxd13* is expressed in

243 the posterior mesenchyme of a limb bud (Fig. 3L) and its expression domain is wider than

244 the *shh*-expressing domain (Fig. 3I). The posterior mesenchymal region of a limb bud is

245 positive for both *dusp6* and *hoxd13* and the double positive region seems to coincide with

246 the fluorescence of MFCS1-GFP (Fig. 1C', D, G') and *gfp* transcript (Fig. 3J) patterns.

The *gfp* transcript was not detected in line 2 limb bud samples (n=0/6) but was detected

in line 1 limb bud samples (n=10/10) by *in situ* hybridization, suggesting that line 1 has

stronger transgene expression than that of line 2. The coincidence between upstream

regulators and MFCS1-GFP expression seems to be clearer in a regenerating limb: *dusp6*

(Fig. 3G) and *hoxd13* (Fig. 3H) were strongly expressed in whole region of a blastema.

MFCS1-GFP expression (Fig. 3R) and *gfp* transcript expression (Fig. 3F) are robustly

distributed in the entire blastema, while *shh* expression is restricted to the posterior edge

of the blastema (Fig. 3E). Therefore, the MFCS1-GFP expression domain is larger than

the *shh* expression domain, especially in a regenerating limb. Transcriptional repression

by a silencer or defined regulation of chromatin conformational change resulting in the

proximity of MFCS1 and the *shh* promoter (Amano et al., 2009) may further restrict

endogenous *shh* expression to the posterior edge of the blastema. Even though *shh*

expression is spatially narrower than the GFP expression pattern, the expression pattern

of MFCS1-GFP covers the *shh*-expressing domain and coincides with the expression

domains of upstream regulators represented by *hoxd13* and *dusp6*.

**MFCS1 is inactive in a limb blastema of a metamorphosed froglet despite *hoxd13***

**expression**

In contrast to a young tadpole, a limb of a metamorphosed froglet does not express *shh*

after amputation, and amputation results in regeneration of a cartilaginous spike, while

an amputated limb of a froglet forms a regeneration blastema (Endo et al., 2000; Suzuki

et al., 2006; Yakushiji et al., 2007). Hypermethylated status of genomic DNA at the

MFCS1 enhancer strongly suggested that this enhancer is inactivated in a froglet limb:

the enhancer inactivation implies transcriptional inactivation of *shh* in the limb (Yakushiji

et al., 2007). To examine whether inactivation of the enhancer occurs after

metamorphosis, we examined MFCS1-GFP expression in limb regeneration of a froglet.

From the early stage (early bud) blastema to the late stage rod-shaped blastema, no

specific GFP was observed in the blastema, while non-specific fluorescence was

sometimes observed in both MFCS1-GFP Tg froglets and wild-type ones (controls) (Fig.

3M-P and Fig. S3). Therefore, we conclude that MFCS1 is inactivated completely or is

much more inactive in the blastema of a froglet limb than in that of a young tadpole.

We then examined expression of upstream regulators of MFCS1. By 10 dpa, an

amputated limb of a froglet forms a cone-shaped blastema and the blastema already starts

expressing *fgf-8*, *fgf-10*, *msx1*, and *hand2,* genes that are involved in maintenance/

induction of *shh* expression in a limb bud (Suzuki et al., 2005; Yakushiji et al., 2007). In

addition, Yakushiji et al. (2007) reported hypermethylation status of genomic DNA in a

limb blastema at 10 dpa in a froglet. Therefore, we chose blastema at 10 dpa and examined

gene expression at that time point. Similar to the results of a study by Satoh et al. (2006)

for the blastema at a later time point (14 dpa), *hoxd13* was broadly expressed in the

blastema at 10 dpa (Fig. 3D). *dusp6* was also expressed in the blastema (Fig. 3C), while

no *shh* (and *gfp*) expression was detected in the blastema as Yakushiji et al. (2007)

reported previously (Fig. 3A, B). The expression of *hoxd13* and *dusp6* in a froglet

blastema suggests that upstream regulators of MFCS1 (5' Hoxd and ETS transcription

factors) as well as *hand2* (Yakushiji et al., 2007) are not absent in the froglet blastema.

Thus, MFCS1 in a froglet limb appears to be unable to respond to positive regulators

represented by Hoxd13.

**Higher DNA methylation level of MFCS1 may inactivate the enhancer's**

**responsiveness to Hoxd13 in a froglet limb**

To confirm the potential responsiveness of MFCS1 to the upstream regulators, we

conducted luciferase assays *in vitro* using MFCS1 with the *shh* promoter. The *luc2*

reporter under control of the MFCS1 enhancer with either *shh* or a minimum promoter

(Fig. 4A, upper diagram) was prepared. A local alignment of the highly conserved

sequences from different species revealed that there are multiple putative Hox-binding

motifs in the conserved sequences (Fig. 4A, see also Fig. S4). The full lengths of *Hoxd13*,

*Fgf2*, *Fgf4*, and *Fgf8* were flanked with the CMV promoter as gene expression constructs.

MFCS1-*shh* promoter-*luc2* or MFCS1-minP-*luc2* reporter constructs were co-transfected

with various combinations of the gene expression constructs into NIH3T3 cells. The

reporter activities of cells transfected with MFCS1-shh promoter-*luc2* were significantly

increased when co-transfecting with *Fgf2* (1.4-fold vs. mock), *Fgf4* (1.5-fold vs. mock),

*Fgf8* (1.3-fold vs. mock), or *Hoxd13* (2.3-fold vs. mock) (Fig. 4B). The combination of

*Fgf4* and *Hoxd13* additively affected MFCS1-driven gene activation (1.1-fold vs. *Hoxd13*

alone). Meanwhile, transfection of the MFCS1-*luc2* construct containing minP had did

not result in significant activation of the luciferase reporter even in the presence of any

expression constructs. When the MFCS1-*luc2* was artificially methylated by CpG

methyltransferase (M.SssI) in advance, the responsiveness of the reporter to Hoxd13 was

strongly suppressed by ~260 fold (Fig. S5). Taken together with the highly methylated

status of MFCS1 in a *Xenopus* froglet limb (Yakushiji et al., 2007), this result raises the

possibility that DNA methylation of MFCS1 inactivates its responsiveness to Hoxd13.

Genomic DNA methylation leads to chromatin condensation, inhibits access of

transcription factors to the enhancer sequence and finally results in transcriptional

repression (Bird and Wolffe, 1999). To further examine whether a high methylation level

of MFCS1 exerts a suppressive effect on gene expression, we investigated the

methylation status of MFCS1 as the transgene in MFCS1-GFP Tg tadpoles and froglets.

Within an MFCS1 highly conserved sequence in amphibians, there are nine CpG sites

both in *X. tropicalis* and *X. laevis* (Fig. S6). Firstly, we examined DNA methylation status

of MFCS1 in wild-type tadpoles and froglets by bisulfite sequencing as reported by

Yakushiji et al. (2007) and obtained consistent results: While limb buds of young tadpoles

have a lower methylation level (26.3%), froglet limbs have a higher methylation level

(87.2%) as shown in Fig. 4C. We then examined the methylation level of the MFCS1

sequence as the transgene in MFCS1-GFP Tg (line 1). Again, froglet limbs have a higher

methylation level (17.2%) than that in limb buds of young tadpoles (2.6%) (Fig. 4C).

After limb amputation in a froglet and after completion of spike regeneration, a higher

methylation level than that in tadpoles was still found in the Tg regenerates (line 1,

14.4%) as well as in wild-type ones (Fig. S7). The nine CpG sites in MFCS1 (highly

conserved sequence in amphibians) as the transgene were susceptible to methylation to

some extent also in line 2 (Fig. S7). Although the basal methylation level differs between

endogenous MFCS1 and transgene MFCS1, the increase in the methylation level of the

transgene MFCS1 after metamorphosis also correlated with the ontogenetic inactivation

of gene expression under control of the enhancer.

**Conclusions**

339  In this study, we visualized MFCS1 activity in limb regeneration of an amphibian.

340  MFCS1-GFP Tg *Xenopus* revealed that MFCS1 enhancer is re-activated specifically in

341  the regenerating blastema for a patterned limb in a young tadpole but not in the blastema

342  for a hypomorphic limb in a froglet. MFCS1 seems to be activated by Hoxd13 in a tadpole

343 but seems to be unable to respond to Hoxd13 in a froglet. An ontogenetic increase in the

344 MFCS1 methylation level may block the function of Hoxd13 (and other positive

345 regulators) as a transcriptional activator(s) to the enhancer so that transcription of a gene

346 under control of the enhancer is inactivated (Fig. 4D). For improvement of regenerative

347 capacity in a froglet limb by re-activation of *shh* expression, removal of epigenetic

348 silencing of MFCS1 would be important. The combinational application of an inhibitor

349 of DNA methyltransferase and a histone deacetylase inhibitor stimulated *shh* expression

350 in cultured blastema cells of a froglet limb in a previous *in vitro* study (Yakushiji et al.,

351 2009), suggesting that application of such reagents affecting epigenetic modification

352 could be hopeful if the treatment condition is optimized *in vivo*. As an alternative

353 approach, CRISPR-based epigenome editing that removes DNA methylation (e.g., Nuñez

354 et al., 2021) may be effective. In these approaches, the MFCS1-GFP Tg line would be

355 useful for evaluating the effectiveness. The enhancer activation is easily detected by GFP

356 and it would thus be possible to screen chemicals that remove epigenetic silencing of the

357 enhancer in a large scale. If regeneration of a well-patterned limb is induced in a froglet

358 by re-activation of MFCS1, it would be a precious model case for recovery of organ-level

359 regeneration in non-regenerative higher vertebrates by manipulating epigenetics.

## MATERIALS AND METHODS

### Transgene construction and *in situ* hybridization

363 A 3,164-bp MFCS1 enhancer sequence was amplified from *Xenopus tropicalis* genomic

364 DNA prepared from a Nigerian line with primers (XtMFCS1-Fw and Xt-MFCS1-Rv;

365 Table S3). The PCR product was cloned into the pCRII-TOPO vector (Invitrogen) and

366 sequenced. The *Xenopus laevis shh* promoter (Yakushiji et al., 2007) was cloned into a

367 reporter construct that contains an EGFP expression cassette. Then the MFCS1-*shh*

368 promoter-*egfp* was generated by inserting the 3,164-bp MFCS1 enhancer sequence

369 upstream of the *shh* promoter (Fig. 1A). For synthesis of probes for *in situ* hybridization,

370 a *hoxd13* cDNA clone, BC080434, inserted in pCMV-SPORT6 was obtained from

371 Dharmacon. The *hoxd13*/ pCMV-SPORT6 plasmid was linearized with SalI and

372 transcribed with T7 RNA polymerase. DIG-labeled RNA probes of *hoxd13*, *shh* (Endo et

373 al., 1997), *gfp* (Hayashi et al., 2015) and *dusp6* (Gomez et al., 2005) were prepared

according to the protocol of the manufacturer (Roche). To prepare serial cryosections, the
specimens were fixed in MEMFA (0.1 M MOPS at pH 7.4, 2 mM EGTA, 1 mM MgSO$_4$,
3.7% formaldehyde) overnight at room temperature, embedded in OCT compound
(Sakura), and serially sectioned at 8-10 μm by a cryostat. *In situ* hybridization on frozen
sections was performed according to Yoshida et al. (1996) with slight modifications.
Sections were observed under an upright microscope (Olympus BX53) with an objective
lens (Olympus UPlan PLN 10X) and were photographed by a digital camera (Leica
DFC7000T).
382

**Animal husbandry and transgenesis in *Xenopus laevis***

Wild-type *Xenopus laevis* adult frogs were purchased from domestic animal venders,
Watanabe Zoushoku (http://www5d.biglobe.ne.jp/~zoushoku/top.htm) and Hamamatsu
Seibutsu Kyozai (http://www.h-seibutsu.co.jp/index.html). Founder (G0) MFCS1-GFP
Tg *Xenopus laevis* embryos were generated by a REMI technique (Kroll and Amaya,
1996) using oocyte extract instead of egg extract (Hirsch et al., 2002; reviewed in Ogino
and Ochi, 2009). MFCS1-GFP N1 Tg *Xenopus laevis* lines were obtained by crossing
sexually mature MFCS1-GFP G0 frogs with wild-type frogs. Tadpoles and froglets were
reared at 22–23°C in dechlorinated tap water. The rearing containers were cleaned daily,
and the tadpoles were fed powdered barley grass (Odani Kokufun, Japan or Yamamoto
Kanpoh, Japan). At stage 58 (Nieuwkoop and Faber, 1994), the feeding was ceased until
metamorphosis was completed. After metamorphosis, the froglets were fed dried tubifex
every other day. All surgery was performed under ethyl-3-aminobenzoate anesthesia, and
all efforts were made to minimize suffering. When tadpoles and froglets needed to be
euthanized, they were immersed in 0.05% ethyl-3-aminobenzoate (Tokyo Chemical
Industry, 886-86-2). When adult male frogs were sacrificed to excise their testes for
transgenesis, 2% ethyl-3-aminobenzoate dissolved in pure water was injected into the
intracardiac cavity (400 μg of ethyl-3-aminobenzoate/1 g of animal). After the male frog
was confirmed to be completely unconscious, its testes were excised with a blade, and
then the unconscious frog was put in a freezer for euthanization. All animal care and
experimentation procedures were conducted in accordance with the animal care and use
committee guidelines of Tohoku University (2015LsA-023) and Hirosaki University
(A15003, A15003-1).

**Observation of developing and regenerating limbs, limb amputation and 180°**

**rotation of a limb bud**

For observation of limb development and limb regeneration in tadpoles, we used hindlimb

buds since it is much easier to observe and to operate on hindlimb buds than forelimb

buds due to their size and accessibility. To observe the regenerating limb bud (blastema)

of a tadpole, tadpoles at stage 54 were anesthetized with 0.025% ethyl-3-aminobenzoate

and hindlimb buds were amputated using ophthalmologic scissors at the presumptive

ankle level (according to the outside view and a fate map by Tschumi, 1957). For

preparation of a 180° rotated limb bud, Nile Blue A particles were embedded into the

anterior side of the distal tip of a stage 53 hindlimb bud with a tungsten needle as the

reference point indicating the original AP axis of the graft according to Cameron and

Fallon (1977). Subsequently, the limb bud was amputated at the presumptive ankle level

(Tschumi, 1957) and then the distal limb bud tip was rotated 180° (not rotated for control

limbs) on the proximo-distal axis, returned to the stump, and fixed with tungsten pins

according to Cameron and Fallon (1977) and Endo et al. (1997). After metamorphosis,

the resulting limbs were fixed in 10% formalin in Holtfreter's solution and stained with

0.1% Alcian blue as previously reported (Yokoyama et al., 1998). Then digit identity was

determined by the number of phalanges in association with the presence/absence of a

claw (Cameron and Fallon, 1977). To observe the blastema of a froglet, froglets were

anesthetized with 0.05% ethyl-3-aminobenzoate and then forelimbs were amputated

through the distal zeugopodium with ophthalmologic scissors and the amputation surface

was trimmed to be flat. For observation of limb regeneration in froglets, we used

forelimbs since hindlimb amputation often results in drowning of the froglets.

**Imaging of whole-mount limb samples**

Tadpoles and froglets anesthetized with 0.025% or 0.05% ethyl-3-aminobenzoate were

observed under a fluorescence stereomicroscope (Nikon SMZ18) with a 0.5x objective

lens using a GFP longpass filter (if in a fluorescent view) and photographed by a digital

camera (Olympus DP74 or DP73). For confocal microscopy (Fig. 1D, E and Fig. S1B,

C), anesthetized tadpoles were each placed on a 35-mm glass-bottom dish (Iwaki, Japan)

so that the dorsal surface of the hindlimb bud was tightly attached to the glass-bottom.

The prepared samples were observed under an FV3000 confocal laser scanning
microscope (Olympus) equipped with an Olympus UPlanSApo 20× dry objective lens.
Fluorescence between 500 nm and 600 nm was detected by a GaAsP photo multiplier
tube.
442

**Genome sequence analysis**

The 449-b genomic sequences corresponding to a conserved core region of human
MFCS1 (hg38, chr7: 156,791,242－156,791,690) and its orthologous sequences in the
mouse (mm39, chr5: 29,520,033–29,520,474), chicken (galGal6, chr2: 8,553,351–
8,553,801), *X. tropicalis* (xenTro10, chr6: 17,346,249–17,346,700) and *X. laevis* (J-strain
10.1, Chr6L: 15,900,483–15,900,931) were downloaded from the UCSC Genome
Browser and Xenbase (Karolchik et al., 2003; Fortriede et al., 2020). These sequences
were aligned using MUSCLE (Edgar, 2004). Searches for potential transcription factor-
binding sites were performed with transcription factor-binding motifs collected from the
TRANSFAC and JASPAR databases (Wingender, 2008; Castro-Mondragon et al., 2022).
The *X. tropicalis* MFCS1 element used for our functional analyses contains a 12-
thymidine repeat in place of the 14-thymidine repeat underlined in the alignment (Fig.
4A, underlined). This difference may be because of a polymorphism between the strain
used for our study and that used for the genome sequencing project (Igawa et al., 2015).
457

**Reporter construction, cell culture and luciferase assay**

The highly conserved 998-bp core sequence of *Xenopus tropicalis* MFCS1 was amplified
by PCR with specific primers (Table S3) and inserted into the Acc65I and XhoI sites of
the pGL4.23 vector (Promega, E8411). Then the minimal promoter of the pGL4.23 vector
was replaced with a 486-bp sequence of *Xenopus laevis shh* promoter (Fig. 4A). It should
be noted that the 14-thymidine repeat in *X. tropicalis* MFCS1 sequence from the UCSC
Genome Browser (Fig. 4A, underlined) was shortened to a 12-thymidine repeat in the
transgene construct (Fig. S6) and the reporter construct for the luc assay. pGL4.74
containing the *Renilla* luciferase gene was used for normalization of reporter expression
(Promega, E6921). Coding sequences of mouse *Hoxd13*, *Fgf2*, *Fgf4* and *Fgf8* were cloned
into the pcDNA3.1 vector (ThermoFisher, V79020). NIH3T3 cell line (RCB2767) was

obtained from the RIKEN BioResource Research Center. NIH3T3 cells were maintained

in Dulbecco's modified Eagle's medium (ThermoFisher, 11960-044) containing 10%

fetal bovine serum, 1x GlutaMax and 100 μg/ml penicillin–streptomycin. Cells at 70–

90% confluency were transfected with equal amounts of luciferase reporter and gene

expression plasmids using Lipofectamine 3000 (ThermoFisher). Luciferase activity in the

cell lysates was measured using a Dual-Luciferase Reporter Assay System (Promega)

according to the manufacturer's protocol. For statistical analysis, Student's $t$-tests (two-

tailed) were applied when equal variance tests were passed.

**478    Bisulfite genomic sequencing**

Presumptive autopods of hindlimb buds (Tschumi, 1957) from thirty-five wild-type and

sixteen MFCS1-GFP Tg (line 1) tadpoles at stage 54 were collected. Forelimb autopods

from five wild-type and six MFCS1-GFP Tg (line 1) froglets were collected. Spike

regenerates from two MFCS1-GFP Tg (line 1) and two same Tg (line 2) frogs were

collected. Then genomic DNA was extracted from these tissue samples. The genomic

DNA was treated with sodium bisulfite using an EZ DNA Methylation-Gold Kit

following the protocol of the manufacturer (Zymo Research). DNA was amplified by 30

cycles of a three-step PCR (98℃ for 10 sec, 60℃ for 30 sec and 68℃ for 30 sec) with

MFCS1_BS_Fw and MFCS1_BS_Rv primers (Table S3) using KOD -Multi & Epi-

(Toyobo). Each product was cloned into the pANT vector (Nippon Gene) or pGEM-T

easy (Promega) and sequenced with M13F-pUC or M13R-pUC primers (Table S3).

**491    ACKNOWLEDGMENTS**

We thank Yoshiko Yoshizawa and Natsume Sagawa for excellent animal care. We thank

Drs. Makoto Asashima, Shuji Takahashi and Yoshio Yaoita for providing the Nigerian

line of *X. tropicalis*. We thank Dr. Ikuo Uchiyama for support in gene expression analysis.

This work was partially supported by the NIBB Collaborative Research Program to HY

(19-451, 20-436, 21-307, 21-401). This work was supported by the National Bio-

Resource Project (NBRP) of the MEXT, Japan. We acknowledge Shared Facility Center

for Science and Technology, Hirosaki University (SFCST) for confocal microscope

observation by FV3000 (Olympus). We thank Dr. Chisato Ushida, Dr. Michiko Sasabe,

Shunsuke Hosoi, Misako Saida and Hibiki Yokoyama for their help in using confocal

microscopy. We thank Dr. Daisuke Kurita for help in TA cloning. We thank Drs. Gembu

Abe and Kazuya Kobayashi for their help in preparation of tungsten needles. We thank

Dr. Jose F. de Celis for providing *Xenopus laevis dusp6* cDNA.

**Competing interests**

506  The authors declare no competing or financial interests.

**Author contributions**

Conceptualization: K.T., H.Y.; Methodology: T.A., N.Y.-K., J.S., Y.K., H.Y.; Validation:

T.A., N.Y.-K., H.Y.; Formal analysis: T.A., H.Ogino, H.Y.; Investigation: R.T., T.H., T.A.,

S.I., C.Y., S.N., K.I., A.K., H.Ochi, H.Ogino, N.Y.-K., Y.K., H.Y.; Resource: T.A., A.K.,

H.Ochi, H.Ogino, N.Y.-K., Y.K., K.T., H.Y.; Writing – original draft preparation: T.A.,

H.Ogino, J.S., Y.K., H.Y.; Writing – review and editing: A.K., H.Ochi, H.Ogino, N.Y.-K.,

K.T., H.Y.; Visualization: R.T., T.H., T.A., S.I., H.Ogino, N.Y.-K., H.Y.; Supervision:

K.T., H.Y.; Project administration: H.Ogino, K.T., H.Y.; Funding acquisition: K.T., H.Y.

**Funding**

518  H.Y is funded by MEXT and JSPS KAKENHI (grant number 22124005). H.Y. is funded

519  by JSPS KAKENHI (grant number 16H04790, 21K06195). H.Y. is funded by Grant for

520  Basic Science Research Projects from The Sumitomo Foundation. H.Y is funded by

521  Hirosaki University Institutional Research Grant for Young investigators.

**REFERENCES**

**Agata, K. and Inoue, T.** (2012). Survey of the differences between regenerative and
non-regenerative animals. *Dev Growth Differ* **54**, 143-152.
**Amano, T., Sagai, T., Tanabe, H., Mizushina, Y., Nakazawa, H. and Shiroishi, T.**
(2009). Chromosomal dynamics at the Shh locus: limb bud-specific differential
regulation of competence and active transcription. *Dev Cell* **16**, 47-57.
**Bird, A. P. and Wolffe, A. P.** (1999). Methylation-induced repression--belts, braces,
and chromatin. *Cell* **99**, 451-454.
**Cameron, J. A. and Fallon, J. F.** (1977). Evidence for polarizing zone in the limb buds
of Xenopus laevis. *Dev Biol* **55**, 320-330.
**Capellini, T. D., Di Giacomo, G., Salsi, V., Brendolan, A., Ferretti, E., Srivastava,
D., Zappavigna, V. and Selleri, L.** (2006). Pbx1/Pbx2 requirement for distal

limb patterning is mediated by the hierarchical control of Hox gene spatial
distribution and Shh expression. *Development* **133**, 2263-2273.

**Castro-Mondragon, J. A., Riudavets-Puig, R., Rauluseviciute, I., Lemma, R. B.,**
**Turchi, L., Blanc-Mathieu, R., Lucas, J., Boddie, P., Khan, A., Manosalva**
**Pérez, N., et al.** (2022). JASPAR 2022: the 9th release of the open-access
database of transcription factor binding profiles. *Nucleic Acids Res* **50**, D165-
D173.

**Dent, J. N.** (1962). Limb regeneration in larvae and metamorphosing individuals of the
South African clawed toad. *J Morphol* **110**, 61-77.

**Eblaghie, M. C., Lunn, J. S., Dickinson, R. J., Münsterberg, A. E., Sanz-Ezquerro,**
**J. J., Farrell, E. R., Mathers, J., Keyse, S. M., Storey, K. and Tickle, C.**
(2003). Negative feedback regulation of FGF signaling levels by Pyst1/MKP3 in
chick embryos. *Curr Biol* **13**, 1009-1018.

**Edgar, R. C.** (2004). MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Res* **32**, 1792-1797.

**Ekerot, M., Stavridis, M. P., Delavaine, L., Mitchell, M. P., Staples, C., Owens, D.**
**M., Keenan, I. D., Dickinson, R. J., Storey, K. G. and Keyse, S. M.** (2008).
Negative-feedback regulation of FGF signalling by DUSP6/MKP-3 is driven by
ERK1/2 and mediated by Ets factor binding to a conserved site within the
DUSP6/MKP-3 gene promoter. *Biochem J* **412**, 287-298.

**Endo, T., Tamura, K. and Ide, H.** (2000). Analysis of gene expressions during
Xenopus forelimb regeneration. *Dev Biol* **220**, 296-306.

**Endo, T., Yokoyama, H., Tamura, K. and Ide, H.** (1997). Shh expression in
developing and regenerating limb buds of Xenopus laevis. *Dev Dyn* **209**, 227-
232.

**Fortriede, J. D., Pells, T. J., Chu, S., Chaturvedi, P., Wang, D., Fisher, M. E.,**
**James-Zorn, C., Wang, Y., Nenni, M. J., Burns, K. A., et al.** (2020).
Xenbase: deep integration of GEO & SRA RNA-seq and ChIP-seq data in a
model organism database. *Nucleic Acids Res* **48**, D776-D782.

**Gómez, A. R., López-Varea, A., Molnar, C., de la Calle-Mustienes, E., Ruiz-**
**Gómez, M., Gómez-Skarmeta, J. L. and de Celis, J. F.** (2005). Conserved
cross-interactions in Drosophila and Xenopus between Ras/MAPK signaling and
the dual-specificity phosphatase MKP3. *Dev Dyn* **232**, 695-708.

**Han, M., Yang, X., Taylor, G., Burdsal, C. A., Anderson, R. A. and Muneoka, K.**
(2005). Limb regeneration in higher vertebrates: developing a roadmap. *Anat*
*Rec B New Anat* **287**, 14-24.

**Harfe, B. D., Scherz, P. J., Nissim, S., Tian, H., McMahon, A. P. and Tabin, C. J.**
(2004). Evidence for an expansion-based temporal Shh gradient in specifying
vertebrate digit identities. *Cell* **118**, 517-528.

**Hayashi, S., Kawaguchi, A., Uchiyama, I., Kawasumi-Kita, A., Kobayashi, T.,**
**Nishide, H., Tsutsumi, R., Tsuru, K., Inoue, T., Ogino, H., et al.** (2015).
Epigenetic modification maintains intrinsic limb-cell identity in Xenopus limb
bud regeneration. *Dev Biol* **406**, 271-282.

**Hirsch, N., Zimmerman, L. B., Gray, J., Chae, J., Curran, K. L., Fisher, M.,**
**Ogino, H. and Grainger, R. M.** (2002). Xenopus tropicalis transgenic lines and
their use in the study of embryonic induction. *Dev Dyn* **225**, 522-535.

**Igawa, T., Watanabe, A., Suzuki, A., Kashiwagi, A., Kashiwagi, K., Noble, A.,**

**Guille, M., Simpson, D. E., Horb, M. E., Fujii, T., et al.** (2015). Inbreeding Ratio and Genetic Relationships among Strains of the Western Clawed Frog, Xenopus tropicalis. *PLoS One* **10**, e0133963.

**Imokawa, Y. and Yoshizato, K.** (1997). Expression of Sonic hedgehog gene in regenerating newt limb blastemas recapitulates that in developing limb buds. *Proc Natl Acad Sci U S A* **94**, 9159-9164.

**Joven, A., Elewa, A. and Simon, A.** (2019). Model systems for regeneration: salamanders. *Development* **146**.

**Karolchik, D., Baertsch, R., Diekhans, M., Furey, T. S., Hinrichs, A., Lu, Y. T., Roskin, K. M., Schwartz, M., Sugnet, C. W., Thomas, D. J., et al.** (2003). The UCSC Genome Browser Database. *Nucleic Acids Res* **31**, 51-54.

**Kawakami, Y., Rodríguez-León, J., Koth, C. M., Büscher, D., Itoh, T., Raya, A., Ng, J. K., Esteban, C. R., Takahashi, S., Henrique, D., et al.** (2003). MKP3 mediates the cellular response to FGF8 signalling in the vertebrate limb. *Nat Cell Biol* **5**, 513-519.

**Kroll, K. L. and Amaya, E.** (1996). Transgenic Xenopus embryos from sperm nuclear transplantations reveal FGF signaling requirements during gastrulation. *Development* **122**, 3173-3183.

**Lettice, L. A., Heaney, S. J., Purdie, L. A., Li, L., de Beer, P., Oostra, B. A., Goode, D., Elgar, G., Hill, R. E. and de Graaff, E.** (2003). A long-range Shh enhancer regulates expression in the developing limb and fin and is associated with preaxial polydactyly. *Hum Mol Genet* **12**, 1725-1735.

**Lettice, L. A., Williamson, I., Wiltshire, J. H., Peluso, S., Devenney, P. S., Hill, A. E., Essafi, A., Hagman, J., Mort, R., Grimes, G., et al.** (2012). Opposing functions of the ETS factor family define Shh spatial expression in limb buds and underlie polydactyly. *Dev Cell* **22**, 459-467.

**Muneoka, K., Holler-Dinsmore, G. and Bryant, S. V.** (1986). Intrinsic control of regenerative loss in Xenopus laevis limbs. *J Exp Zool* **240**, 47-54.

**Nieuwkoop, P. D. and Faber, J.** (1994). *Normal Table of Xenopus laevis (Daudin)*. New York: Garland Publishing.

**Nuñez, J. K., Chen, J., Pommier, G. C., Cogan, J. Z., Replogle, J. M., Adriaens, C., Ramadoss, G. N., Shi, Q., Hung, K. L., Samelson, A. J., et al.** (2021). Genome-wide programmable transcriptional memory by CRISPR-based epigenome editing. *Cell* **184**, 2503-2519.e17.

**Ogino, H. and Ochi, H.** (2009). Resources and transgenesis techniques for functional genomics in Xenopus. *Dev Growth Differ* **51**, 387-401.

**Otsuki, L. and Tanaka, E. M.** (2022). Positional Memory in Vertebrate Regeneration: A Century's Insights from the Salamander Limb. *Cold Spring Harb Perspect Biol* **14**.

**Riddle, R. D., Johnson, R. L., Laufer, E. and Tabin, C.** (1993). Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell* **75**, 1401-1416.

**Roy, S. and Gardiner, D. M.** (2002). Cyclopamine induces digit loss in regenerating axolotl limbs. *J Exp Zool* **293**, 186-190.

**Sagai, T., Hosoya, M., Mizushina, Y., Tamura, M. and Shiroishi, T.** (2005). Elimination of a long-range cis-regulatory module causes complete loss of limb-specific Shh expression and truncation of the mouse limb. *Development* **132**, 797-803.

**Sagai, T., Masuya, H., Tamura, M., Shimizu, K., Yada, Y., Wakana, S., Gondo, Y., Noda, T. and Shiroishi, T.** (2004). Phylogenetic conservation of a limb-specific, cis-acting regulator of Sonic hedgehog ( Shh). *Mamm Genome* **15**, 23-34.

**Satoh, A., Endo, T., Abe, M., Yakushiji, N., Ohgo, S., Tamura, K. and Ide, H.** (2006). Characterization of Xenopus digits and regenerated limbs of the froglet. *Dev Dyn* **235**, 3316-3326.

**Suzuki, M., Hayashi, T., Inoue, T., Agata, K., Hirayama, M., Shigenobu, S., Takeuchi, T., Yamamoto, T. and Suzuki, K. T.** (2018). Cas9 ribonucleoprotein complex allows direct and rapid analysis of coding and noncoding regions of target genes in Pleurodeles waltl development and regeneration. *Dev Biol* **443**, 127-136.

**Suzuki, M., Satoh, A., Ide, H. and Tamura, K.** (2005). Nerve-dependent and -independent events in blastema formation during Xenopus froglet limb regeneration. *Dev Biol* **286**, 361-375.

**Suzuki, M., Yakushiji, N., Nakada, Y., Satoh, A., Ide, H. and Tamura, K.** (2006). Limb regeneration in Xenopus laevis froglet. *ScientificWorldJournal* **6 Suppl 1**, 26-37.

**Torok, M. A., Gardiner, D. M., Izpisúa-Belmonte, J. C. and Bryant, S. V.** (1999). Sonic hedgehog (shh) expression in developing and regenerating axolotl limbs. *J Exp Zool* **284**, 197-206.

**Tschumi, P. A.** (1957). The growth of the hindlimb bud of Xenopus laevis and its dependence upon the epidermis. *J Anat* **91**, 149-173.

**Wingender, E.** (2008). The TRANSFAC project as an example of framework technology that supports the analysis of genomic regulation. *Brief Bioinform* **9**, 326-332.

**Yakushiji, N., Suzuki, M., Satoh, A., Sagai, T., Shiroishi, T., Kobayashi, H., Sasaki, H., Ide, H. and Tamura, K.** (2007). Correlation between Shh expression and DNA methylation status of the limb-specific Shh enhancer region during limb regeneration in amphibians. *Dev Biol* **312**, 171-182.

**Yakushiji, N., Yokoyama, H. and Tamura, K.** (2009). Repatterning in amphibian limb regeneration: A model for study of genetic and epigenetic control of organ regeneration. *Semin Cell Dev Biol* **20**, 565-574.

**Yokoyama, H., Endo, T., Tamura, K., Yajima, H. and Ide, H.** (1998). Multiple digit formation in Xenopus limb bud recombinants. *Dev Biol* **196**, 1-10.

**Yoshida, K., Urase, K., Takahashi, J., Ishii, Y. and Yasugi, S.** (1996). Mucus-associated antigen in epithelial cells of the chicken digestive tract: developmental change in expression and implications for morphogenesis-function relationships. *Dev. Growth Differ.* **38**, 185-192.

**Fig. 1. MFCS1-GFP expression coincides with *shh* expression in a developing limb bud.** (A) Map of the MFCS1-GFP transgene. (B) *shh* expression in a limb bud at stage 52. (C, C') A limb bud of an MFCS1-GFP tadpole at stage 51. Bright-field (C) and fluorescent (C') images. (D, E) Optical-section images by confocal microscopy (approximately 60-70 μm in depth from the dorsal surface) of the right limb bud of an MFCS1-GFP tadpole (D) and that of a wild-type tadpole (E) at stage 51. Bright-field (F-J) and fluorescent (F'-J') images of the left limb bud of the same MFCS1-GFP tadpole. Limb buds are outlined by dotted lines in the fluorescent images. All of the MFCS1-GFP tadpoles shown are line 1. a, anterior: p, posterior. Arrowheads show *shh* expression (B) or GFP expression (D, E, G', H') at the posterior side. Brackets indicate the epidermal cell layer. Bars = 100 μm for (B, D ,E) and 200 μm for (C, C', F-J, F'-J').

**Fig. 2. MFCS1-GFP expression precedes limb duplication along the AP axis in 180°**
**rotated limb buds.** (A) 180° rotation of a limb bud at stage 53. A magenta arrowhead
indicates an embedded Nile blue A particle. Note the inverted position of the particle
before (lower side) and after (upper side) grafting. Black arrowheads indicate a tungsten
pin. (B, C) A 0° rotated limb bud at 6 days after grafting (B) and the resultant normal limb

formation (C). (D-G) 180° rotated limb buds. Rotated limb buds at 6 days after grafting
(D,F) resulted in supernumerary limb formation (E,G). Note that the sample with brighter
GFP (F) resulted in more supernumerary limbs (G). All of the operated MFCS1-GFP
tadpoles are line 1. A pair of lines indicates the host-graft boundary. An asterisk indicates
ectopic GFP. An Arabic number indicates the digit identify. A black arrow shows an
unidentifiable digit. Bars = 200 μm for (B, D, F) and 1 mm for (C, E, F).
695

**Fig. 3. Comparison of gene expression patterns and MFCS1-GFP expression pattern in a regenerating limb in a young tadpole and a froglet.** (A-D) Adjacent sections of a blastema of a froglet. n=5 of 5 blastemas from a total of five froglets of line 1. (E-H) Adjacent sections of a blastema of a tadpole. (I-L) Adjacent sections of a limb bud. To guarantee an accurate comparison of gene expression levels, limb bud and blastema sections were treated in exactly the same way. n=9 of 9 blastemas from a total of five tadpoles of line 1 and 6 of 6 blastemas from a total of three tadpoles of line 2, respectively. n=10 of 10 limb buds from a total of five tadpoles of line 1 and 6 of 6 from a total of three tadpoles of line 2, respectively. (M-P) A blastema of a froglet of line 1. See also Fig. S3 for details. (Q-T) A blastema of a tadpole of line 1. The blastema and the stump are outlined by a dotted line. See also Fig. S2 for details. a, anterior: p, posterior. A pair of lines indicates the amputation level. Arrows indicate *gfp* expression at the posterior side. All pictures shown in Fig. 3 are samples of line 1. Bar = 200 μm for (A-L, Q-T) and 500 μm for (M-P).

**Fig. 4. Transactivation and DNA methylation of *Xenopus* MFCS1 enhancer.** (A) Upper panel: Map of MFCS1-luc2 used for assays in (B). Lower panel: Alignment of MFCS1 highly conserved sequences. Nucleotides identical in more than two sequences are shaded in grey. Conserved Hox-binding motifs are boxed. A light bule line underlies the 14-thymidine repeat in *X. tropicalis* (See Materials and Methods for details.). (B) Firefly luciferase activity of MFCS1-luc2 using a minimum promoter (minP) or *shh* promoter (shh pro) was normalized to *Renilla* luciferase control activity. The mean value of mock samples by the control construct with the minimum promoter was adjusted to 1.0. Error bars: standard deviation from the mean (n=3). *Significant increase from mock (p<0.01). **Significant increase in Fgf2+Hoxd13, Fgf4+Hoxd13 and Fgf8+Hoxd13 from sole Fgf2, Fgf4 and Fgf8, respectively (p<0.01). (C) Methylation pattern of endogenous and transgene MFCS1 (line 1). Samples were obtained from the autopod (pink). Each

circle corresponds to a CpG site (see Fig. S6). Lower panel: Increases in average

methylation rate of the MFCS1 sequences after metamorphosis. (D) Model for epigenetic

silencing of gene expression through DNA methylation of MFCS1. Hoxd13 activates

hypomethylated MFCS1 and results in transcription of a gene under control of MFCS1

in the tadpole limb (left). However, Hoxd13 is unable to activate highly methylated

MFCS1 in a froglet limb and results in inactive transcription of the gene (right).

**Fig. S1. MFCS1-GFP expression pattern was recapitulated in line 2.** (A) A developing limb bud of an MFCS1-GFP Tg tadpole of line 2 at stage 52. Bright-field (A) and fluorescent (A') images of the same limb bud. (B, C) Optical-section images by confocal microscopy (approximately 80 μm in depth from the dorsal surface) of the right limb bud of an MFCS1-GFP Tg tadpole of line 2 (B) and a wild-type tadpole (C) at stage 52. GFP expression was detected in the posterior side mesenchyme specifically in the MFCS1-GFP Tg tadpole. Bright field (D-H) and fluorescent (D'-H') images of the limb bud of the same Tg tadpole of line 2. Limb buds are outlined by dotted lines. GFP expression was detected in the posterior side of limb buds at stage 51-52 and then gradually diminished. a, anterior: p, posterior. Arrowheads show GFP expression (A, A', B, E', F') at the posterior side of the limb buds. Brackets indicate the epidermal cell layer. A pair of lines indicates the amputation level. Bars = 100 μm for (A, B, C), 200 μm for (D-H, D'-H').

**Fig. S2. Spatio-temporal expression pattern of MFCS1-GFP in a regenerating limb**
**of a young tadpole.** All of the pictures are focused on a limb blastema of the same tadpole
(identical to Fig. 3Q-T) of line 1. Left and right limb buds were amputated at the
presumptive ankle level of a stage 54 tadpole. GFP fluorescence was distributed

throughout the regenerating limb blastema at 4 dpa and 6 dpa. (A'', B'', C'', D'', E''')

Limb blastemas and stumps are outlined by dotted lines. n=16 of 16 blastemas from a

total of ten MFCS1-GFP tadpoles of line 1 and 6 of 6 blastemas from a total of three

MFCS1-GFP tadpoles of line 2, respectively. A pair of lines indicates the amputation

level. Arrows show GFP fluorescence of the contralateral right limb blastemas. a,

anterior: p, posterior. Bar = 200 μm.

**Fig. S3. MFCS1-GFP expression is inactive in a limb blastema of a froglet.** A blastema of the same froglet of line 1 (right two columns; identical to Fig. 3M-N) and a blastema of the same wild-type froglet (left two columns) from 2 dpa to 14 dpa. No specific GFP fluorescence was observed in the blastema of the Tg froglet at any timepoint. n=4 of 4 blastemas from a total of four froglets of line 1, 6 of 6 blastemas from a total of four

765     froglets of line 2 and 8 of 8 blastemas from a total of six wild-type froglets as controls,

766     respectively. A pair of lines indicates the amputation level. An arrow indicates non-

767     specific fluorescence commonly observed both in the Tg froglet and wild-type one. a,

768     anterior: p, posterior. Bar = 500 μm.

**Fig. S4. Alignment of MFCS1 highly conserved sequences (identical sequences shown in Fig. 4A) from the human, mouse, chick, *X. tropicalis* and *X. laevis*.** Nucleotides identical in more than two sequences are shaded in grey. Conserved transcription factor-binding motifs (Hox and other transcription factors) are boxed.

**Fig. S5. DNA methylation suppresses the response of MFCS1 to Hoxd13 *in vitro*.**

(A) Methylation-sensitive restriction digestion visualized by agarose gel

electrophoresis. Plasmid DNA of the MFCS1-*shh* promoter-*luc2* reporter was treated

with CpG methyltransferase (M. SssI, NEB) according to the manufacturer's protocol.

Methylation efficiency of the plasmid DNA was evaluated by digestion with

methylation-sensitive (HpaII) and -insensitive (MspI) restriction enzymes. The CpG

methylation prohibited the digestion of the plasmid DNA specifically with HpaII. M,

l/HindIII marker. (B) Luciferase activity in NIH3T3 cells transfected with the

methylated or non-treated MFCS1-luc2 reporter constructs and the *Hoxd13* expression

construct. The mean value of luciferase activity in cells transfected with the non-treated

reporter and mock expression constructs was set to 1.0. Error bars: standard deviation

from the mean (n=3). While a significant increase of *Hoxd13*-induced luciferase activity

was observed in cells transfected with the unmethylated reporter construct (*$p<0.01$),

being consistent with Fig. 4B, there was no significant increase in cells transfected with

the methylated reporter construct despite *Hoxd13* expression. N. S., not significant.

**Fig. S6. *X. tropicalis* and *X. laevis* have 9 CpG sites in the highly conserved sequence**

**of MFCS1.** Alignment of MFCS1 highly conserved sequences between *X. tropicalis* and

*X. laevis*. Green letters correspond to MFCS1_BS_Fw and MFCS1_BS_Rv primers for

bisulfite sequencing (see also Table S3). Open and black circles represent unmethylated

and methylated CpG sites, respectively.

**Fig. S7. Methylation pattern of MFCS1 in a spike regenerate after limb regeneration**

**of a froglet.** After limb blastemas of froglets completed spike regeneration, regenerate

samples (pink) were obtained from line 1 and line 2 Tg frogs.

**Table S1. Rate of germline transmission and activity of MFCS1-GFP**

| G0 | N1 | N2 |
|---|---|---|
| | GFP+/ total | GFP+/ total |
| line 1 | 195/368 (53%) | 70/172 (41%) |
| line 2 | 79/180 (44%) | not applicable |

## Table S2. Supernumerary limb formation and ectopic GFP in rotated limb buds

180° rotated limb buds

| Sample | No. of original digits | No. of supernumerary digits | No. of digits (total) | Is there ectopic GFP? | Figure |
|--------|------------------------|------------------------------|------------------------|------------------------|--------|
| 1 | 5 | 2 | 7 | Yes | - |
| 2 | 5 | 2 | 7 | Yes | - |
| 3 | 5 | 2 | 7 | Yes | - |
| 4 | 5 | 2 | 7 | Yes | - |
| 5 | 5 | 4 | 9 | Yes | Fig. 2F, G |
| 6 | 5 | 2 | 7 | Yes | - |
| 7 | 5 | 1 | 6 | Yes | Fig. 2D, E |
| 8 | 5 | 0 | 5 | No | - |
| 9 | 5 | 0 | 5 | No | - |
| 10 | 5 | 0 | 5 | No | - |
| 11 | 5 | 2 | 7 | Yes | - |

0° rotated limb buds (controls)

| Sample | No. of original digits | No. of supernumerary digits | No. of digits (total) | Is there ectopic GFP? | Figure |
|--------|------------------------|------------------------------|------------------------|------------------------|--------|
| 1 | 5 | 0 | 5 | No | Fig. 2B, C |
| 2 | 5 | 0 | 5 | No | - |
| 3 | 5 | 0 | 5 | No | - |
| 4 | 5 | 0 | 5 | No | - |
| 5 | 5 | 0 | 5 | No | - |
| 6 | 5 | 0 | 5 | No | - |
| 7 | 5 | 0 | 5 | No | - |
| 8 | 5 | 0 | 5 | No | - |
| 9 | 5 | 0 | 5 | No | - |
| 10 | 5 | 0 | 5 | No | - |
| 11 | 5 | 5 | 10 | Yes | - |
| 12 | 5 | 3 | 8 | Yes | - |

# Table S3. Primers for genomic- and RT-PCR

| name | Sequence (5' -> 3') | note |
|------|---------------------|------|
| xtMFCS1-Fw | ACTGTAACCAGTGAAAATCATCAGTCATGAATATC | transgene |
| xtMFCS1-Rv | TGGCCTAGTTCATAGACATTAATTATTTTCTATGGG | transgene |
| xtM1-Acc65I-Fw | ttggtaccTTCAAATGATGGATTTTCATATTATAA | Luc reporter |
| xtM1-XhoI-Rv | ggctcgaGACCATGAAGGCACCCTAATACT | Luc reporter |
| xlpS-XhoI-Fw | ggctcgAGTGACACCGTGGAAGCCTTTATTATC | Luc reporter |
| xlpS-NcoI-Rv | ggccatggtGTCCGAGCGAAGCCAATTACATCA | Luc reporter |
| Fgf2_cds_F1 | GCCATGGCTGCCAGCGGCATCACC | Gene Expression |
| Fgf2_cds_R1 | TCAGCTCTTAGCAGACATTGGAAG | Gene Expression |
| Fgf4_cds_F1 | CCGGGATGGCGAAACGCGGGCCGA | Gene Expression |
| Fgf4_cds_R1 | TCACAGTCTAGGAAGGAAGTGGGT | Gene Expression |
| Fgf8_cds_F1 | GACCCCGCTCGGGCTCTCAGTGCT | Gene Expression |
| Fgf8_cds_R1 | CTATCGGGGCTCCGGGGCCCAAGT | Gene Expression |
| Hoxd13-F | ATGAGCCGCTCGGGGACTTG | Gene Expression |
| Hoxd13-R | TCAGGAGACAGTGTCTTTGA | Gene Expression |
| MFCS1_BS_Fw | GTGTAATTAATTAGGTGGGyyAGGTGGAAA | Bisulfite sequencing |
| MFCS1_BS_Rv | TTAArCArCACTCCCTTAATCACTCATTTC | Bisulfite sequencing |
| M13F-pUC | GTTTTCCCAGTCACGAC | Bisulfite sequencing |
| M13R-pUC | CAGGAAACAGCTATGAC | Bisulfite sequencing |