Displacement of extracellular chloride by sulfated glycosaminoglycans of the brain's extracellular matrix[*]

Kieran P Normoyle,[1,2] Kyle P Lillis,[1,2] Kiyoshi Egawa,[3] Melanie A McNally,[1,2] Mousumi Paulchakrabarti,[4] Biswa P Coudhury,[4] Lauren Lau,[1,2] Thomas Shiu,[1,2] and Kevin J Staley[1,2]

1. Department of Neurology, Division of Child Neurology, Massachusetts General Hospital, Boston, MA, United States
2. Department of Neurology, Harvard Medical School, Boston, MA, United States
3. Department of Medicine, Hokaiddo University, Sapporo, Hokaiddo, Japan
4. GlycoAnalytics Core, University of California San Diego, La Jolla, CA, United States

Abstract

GABA is the primary inhibitory neurotransmitter. Membrane currents evoked by $GABA_A$ receptor activation have uniquely small driving forces: their reversal potential (E_{GABA}) is very close to the resting membrane potential. As a consequence, $GABA_A$ currents can flow in either direction, depending on both the membrane potential and the local intra and extracellular concentrations of the primary permeant ion, chloride (Cl). Local cytoplasmic Cl concentrations vary widely due to displacement of mobile Cl ions by relatively immobile anions. Here we use new reporters of extracellular chloride (Cl^-_o) to demonstrate that Cl is displaced in the extracellular space by high and spatially heterogenous concentrations of sulfated glycosaminoglycans (sGAGs). Cl^-_o varies widely, and the mean Cl^-_o is only half the canonical concentration, i.e. the Cl concentration in the cerebrospinal fluid. These unexpectedly low and heterogenous Cl^-_o domains provide a mechanism to link the varied but highly stable distribution of sGAGs in the brain's extracellular space to neuronal signal processing via the effects on the amplitude and direction of $GABA_A$ transmembrane Cl currents.

Introduction

$GABA_A$ synapses comprise the primary output for inhibitory interneurons in the brain (Hu *et al.*, 2014; Tremblay *et al.*, 2016). A unique feature of $GABA_A$ synapses are the very small driving forces for ion flux through open $GABA_A$ receptor-operated membrane channels. The reversal potential for the primary permeant ion, chloride, is usually within a few mV of the neuronal resting membrane potential (RMP) (Thompson *et al.*, 1988) and may even be positive to RMP (Staley & Mody, 1992; Tyzio *et al.*, 2008). This low driving force makes possible a novel form of synaptic plasticity at $GABA_A$ synapses: the direction of current flow, i.e. the effect of $GABA_A$ receptor activation on the resting membrane potential, can be reversed by low millimolar alterations in the local chloride concentration (Coombs *et al.*, 1955). Such physiological variance is routinely observed in individual studies (Ebihara *et al.*, 1995; Huberfeld *et al.*, 2007) and is likely to underlie the 25 mV range of reported group mean reversal potentials in 20 recent studies of evoked $GABA_A$ currents recorded with gramicidin perforated patch electrodes in area CA1 of the hippocampus (tabulated in (Rahmati *et al.*, 2021). This variance has substantial effects on synaptic signaling: the reversal potential of $GABA_A$ currents evoked in a single

pyramidal cell by stimulation of 8 different presynaptic interneurons also varied by over 20 mV
(Rahmati *et al.*, 2021).

We have proposed that the intracellular chloride concentration (Cl^-_i) is not monotonic across the
neuronal cytoplasm, but rather varies inversely with the local concentration of macromolecular
anions that comprise the vast majority of intracellular anions (Glykys *et al.*, 2014a; Rahmati *et*
al., 2021). This phenomenon is known as Donnan exclusion (Marinsky, 1985a; Helfferich, 1995;
Fatin-Rouge *et al.*, 2003; Epsztein *et al.*, 2018; Gao *et al.*, 2022). Mobile ions, in this case Cl,
are repelled by relatively immobile ions such as proteins with negative surface charges. Many
proteins such as gephyrin that are associated with $GABA_A$ synapses bear large negative
surface charges (Sola *et al.*, 2001), suggesting that variance in the $GABA_A$ reversal potential
(E_{GABA}) is determined by local fixed cytoplasmic charge densities. However, E_{GABA} is not
determined solely by Cl^-_i but rather by the ratio of intra- to extracellular permeant ions. This
raises the possibility that spatial variance in the extracellular Cl concentration (Cl^-_o) could also
be an important determinant of E_{GABA}.

The displacement of Cl^- by fixed negative charges should also occur in the extracellular space,
where the high negative charge density of the polysulfated glycosaminoglycans (GAGs) that
comprise the bulk of the extracellular matrix (Karamanos *et al.*, 2021) can be as high as the Cl^-
concentration in the CSF (Urban *et al.*, 1979; Chahine *et al.*, 2005; Morawski *et al.*, 2015). Thus
some to all of the local Cl^-_o could be displaced, depending on GAG density and the degree of
GAG sulfation. Such displacement has been most clearly demonstrated in the matrix of
cartilaginous tissue, where the negative charge density of the extracellular matrix can reach 400
meq / liter (Lesperance *et al.*, 1992), but dozens of mM of Cl^-_o are also displaced by sulfated
GAGs in tissues such as skin (Titze *et al.*, 2004). Mucous, which comprises the extracellular
space for airway epithelial cells, is also rich in polysulfated GAGs that displace about half of the
Cl^-_o (Reuter *et al.*, 1998).

Many findings suggest that the brain Cl^-_o is lower than the canonical value. Xray emission
spectroscopy of the extracellular space has demonstrated sulfate concentrations as high as 100
mM (Morawski *et al.*, 2015). Astrocytic buffering of Cl^-_o (Egawa *et al.*, 2013) would not be
necessary if Cl^-_o were indeed > 100 mM. The gating of Group II and III metabotropic glutamate
receptors, kainate receptors, acid-sensing membrane ion channels, and the transport rate of the
GlyT1 glycine transporter are all strongly modulated by Cl^-_o over the entire range of 1 - 100 mM
(Kuang & Hampson, 2006; Plested & Mayer, 2007; Kusama *et al.*, 2010; DiRaddo *et al.*, 2015;
Tora *et al.*, 2018; Zhang *et al.*, 2021). Such strong modulation by low Cl^-_o would not make
sense if Cl^-_o was uniformly > 100 mM.

Studies of Cl^-_o are few, particularly in live tissue, and have only been carried out with relatively
large multibarreled ion-sensitive electrodes with tip diameters of 3-5 microns (Jiang *et al.*, 1992).
This technique may alter the reported Cl^-_o via leakage of Cl from the high-Cl barrel of the
electrode or distortion of the extracellular milieu through which the electrode passes. Modern
Cl-selective microelectrodes are less invasive, yet recent studies also demonstrate modest
variance in Cl^-_o of only a few mM (Kroeger *et al.*, 2010).

A fluorometric study of Cl^-_o would provide a noninvasive means to establish the distribution of
chloride in the extracellular space. However, no suitable Cl-sensitive organic fluorophores exist,
and transgenic fluorophores such as SuperClomeleon (Grimley *et al.*, 2013a) and CloPhensor
(Sato *et al.*, 2017) are not distributed in the extracellular space. We therefore developed a

chloride-sensitive fluorophore (Normoyle *et al.*, 2024) from previously described organic
backbones (Biwersi *et al.*, 1992b; Biwersi *et al.*, 1994) that satisfied the following criteria. First,
the fluorophore should be sensitive to Cl in the range of the extracellular chloride concentration
i.e., up to 100 mM; available fluorophores have been optimized for the low millimolar chloride
concentrations of the cytoplasm. Second, the fluorophore should report Cl independently of the
concentration of fluorophore. Ratiometric fluorophores suitable for the extracellular space do
not exist, so we used $\underline{F}$luorescence $\underline{L}$ifetime $\underline{Im}$aging (FLIM) (Kaneko *et al.*, 2004) to remove the
dependence of the fluorescence emission on fluorophore concentration. Third, the fluorophore
should be constrained to the extracellular space, for which dextran conjugation is optimal (Xiao
et al., 2008; Tønnesen *et al.*, 2018). Fourth, the fluorophore should be excited by wavelengths
that are not damaging to brain tissue, i.e. below the ultraviolet range and in the visible or
infrared range.   Fifth, many Cl sensors are also sensitive to pH (Kuner & Augustine, 2000;
Grimley *et al.*, 2013b), so a pH-insensitive fluorophore would be ideal. Finally, the fluorophore
needs to be sufficiently bright so that its emission can be reliably distinguished from
autofluorescence.

Here, we utilize a newly synthesized Cl-sensitive fluorophore that meets these 6 criteria to
demonstrate: stable Donnan exclusion of chloride by sulfates fixed to GAG-like carbohydrate
polymers; directly measured Cl^-_o of *in vitro* and *in vivo* brain tissue that is much lower than
cerebrospinal or plasma chloride; a remarkable spatial variance in Cl^-_o; and, as evidence that Cl^-_o
is determined by Donnan exclusion of Cl by sulfated GAGs, Cl^-_o increases after release of
these sulfates by exogenous chondroitinase.

Methods

<u>General</u>: Chemicals were purchased from Sigma-Aldrich (St Louis, MO) unless otherwise noted.
Solutions are brought up in distilled water unless otherwise noted. Standard aCSF was brought
up in 18MΩ water and consisted of (in mM): NaCl 126; KCl 3.5; $CaCl_2$ 2; $MgCl_2$ 1.3; NaH_2PO4
1.2; Glucose 11; $NaHCO_3$ 15. Low chloride aCSF variant was identical except that sodium
gluconate was substituted for NaCl, resulting in 10mM Cl^-. Calibration solutions were made by
mixing appropriate proportions of standard aCSF with low chloride aCSF such that a chloride
and gluconate sum total of 136mM was maintained while Cl^- ranged from 10 to 136mM.
Colorimetric assays were read using a Wallac Victor-2 1420 spectrophotometer with a halogen
continuous wave light source and spectral line filters at listed wavelength +/- 5 to 10nm.
Centrifugation steps were accomplished in a tabletop Eppendorf 5417R microcentrifuge. Reflux
apparatus for synthesis consisted of a recirculating cooling bath (Fisher Scientific) filled with
ethylene glycol cooling a 24/40 double-lumen coiled reflux tube (Ace Glassware, Vineland, NJ)
with a 500mL round bottom flask (Corning) heated with a heating mantle regulated by a timed
power controller (Glas-Col #O406 and #104A PL312, respectively).

<u>Synthesis of ABP-dextran</u>: The process of optimizing a bright, redshifted, pH-insensitive
fluorophore that is responsive to chloride concentration over a dynamic range inclusive of
expected extracellular chloride concentrations is beyond the scope of this paper and has been
published separately (Normoyle *et al.*, 2024). Synthesis techniques closely followed those of the
Verkman group when synthesizing related compounds (Verkman, 1990; Biwersi *et al.*, 1992a)
and conjugating to dextran (Biwersi *et al.*, 1992a). Briefly, N(4-***A***mino***B***utyl)***P***henantridinium
(ABP) was synthesized by a reflux of equimolar amounts of phenanthridine and N(4-

bromobutyl)phthalimide in acetonitrile followed by a second reflux in 6N HCl to hydrolyze
phthalate. Product was twice recrystallized from 95% ethanol before conjugation to cyanogen
bromide-activated 10,000g/moL (10KDa) dextran, producing 10KDa ABP-dextran with an
approximate molar labeling ratio of 4.4.

<u>Sulfation of agarose</u>: We modified the method of Fuse and Suzuki (Fuse & Toshiya, 1975) to
add sulfate groups to the repeating disaccharide units of agarose. The goal of maximizing
sulfation must be balanced with the consequent erosion of gel properties such as gelation and
water holding capacity. Briefly, Agarose was added to a well-dried Erlenmeyer flask and
dissolved in DMSO at a concentration of 4%(w/v) at room temperature before being cooled to
4°C with stirring. Addition of 4:1 mixture of acetic anhydride / glacial acetic acid, also pre-chilled
to 4°C, was added to agarose/DMSO mixture (0.4 volumes) in 4°C coldroom. Vigorous mixing
by handheld rotation was necessary to thin the gelatinous mixture to a slurry and allow stirring
at 4°C for 15 minutes; gentle heat transfer from experimenter's hands helped loosen the initial
gel but care is taken to return the mixture to 4°C. Sulfuric acid (15.6M) is then added slowly at a
6:1 molar ratio to agarose repeating disaccharide units, taking care to keep temperature
between 4-10°C. Mixture is then moved to ambient temperature and allowed to stir for 15
minutes before reaction is stopped by neutralization with 10M NaOH after again cooling to 4°C
and maintaining temperature between 4-10°C in coldroom. Sulfated agarose is then dialyzed at
ambient temperature in 12-14KDa dialysis tubing (SpectraPor S432703) against 15 volumes of
50mM Tris pH 7.4 once, then against distilled water changed twice daily until dialysate clarifies.
Agarose is then precipitated in 95% ethanol cooled to 4°C >4hours; over-dialysis may hinder
ethanol precipitation due to a lack of sufficient counter-ion presence. Precipitate is then vacuum
filtered and oven-dried at 50°C. Material is loosened with polished glass pestle and stored at
ambient temperature in sealed glass vial until use. Note that careful temperature and pH control
is necessary throughout exothermic processes to maintain gelation characteristics of the
product.

<u>Measurement of sulfate concentration in polysaccharide gels</u>: $BaCl_2$-gelatin turbidimetric method
(Torres *et al.*, 2021) was used to quantify the sulfate content of gel materials. Sulfated agarose
product or commercially available agarose or agar powder was dissolved in 0.5M HCl at a
concentration of 5% (w/v) in 2mL screw-top microtubes with gasket (Fisher Scientific). Tubes
are then placed in a heat block at 100-110°C for 3hours with periodic vortex mixing, hydrolyzing
the sugar backbone and releasing sulfate. All samples had varying amounts of insoluble dark
brown precipitated sugar, which was clarified at 20,000 x g for 10 minutes. Standards are
diluted from 1M Na_2SO_4 stock in 0.5M HCl. $BaCl_2$-gelatin reagent is freshly prepared by
dissolving 30mg gelatin in 10mL distilled water at 80°C for 10minutes, then promptly adding
100mg $BaCl_2$ and vortex mixing. $BaCl_2$-gelatin is then allowed to cool passively before use.
Samples are prepared by combining 1 part sample (diluted as desired), 1 part $BaCl_2$-gelatin,
and 3 parts 0.5M HCl. Assay is performed in a glass-bottom 96-well plate reading absorbance
at 405nm. Multiple sample dilutions were prepared and concentrations were calculated only
from dilutions whose readings were well within the dynamic range of the standards.

<u>Measurement of Cl^- in agar or agarose gels (chemical method)</u>: Chloride was measured
colorimetrically using mercury(II) thiocyanate $(Hg(SCN)_2)$ / iron(III) nitrate $(Fe(HNO_3)_3)$
absorbance method of Florence and Farrar (Florence & Farrar, 1971) to infer Cl^- from $Fe(SCN)_3$
production through the two-step reaction,

$$[Hg(SCN)_2 + 2Cl^- \rightleftharpoons HgCl_2 + 2SCN^-]x3$$

$$[3SCN^- + Fe(NO_3)_3 \rightleftharpoons Fe(SCN)_3 + 3(NO_3)^-]x2$$

reading $Fe(SCN)_3$ absorbance at 490nm. Briefly, agar or agarose gels cast at 5% (w/v) in 3.5cm
Petri dishes were drained by gravity and then Kimwipes were used to gently wick away
remaining excess fluid at sides of dish. Similar amounts of each sample are transferred to 2mL
screw-top microtubes with gasket (Fisher Scientific) and 1 gel volume, estimated by weight, of
7.8M nitric acid (HNO_3) is added (final $[HNO_3]$ = 3.9M). Samples are then placed in a heat block
at 100-110°C for 3hours with periodic vortex mixing, hydrolyzing the sugar backbone to break
down the gel lattice. All samples had varying amounts of insoluble dark brown precipitated
sugar, which was clarified at 20,000 x g for 10 minutes. A portion of supernatant or KCl
standard, diluted with 0.3M HNO_3, was combined with one-tenth volumes of 0.4M $Fe(NO_3)_3$ (in
4M nitric acid) and saturated $Hg(SCN)_2$ (in ethanol), triturated, and allowed 20minutes for
reaction to reach endpoint. Stock solutions are used within 1week. Multiple sample dilutions
were prepared and concentrations were calculated only from dilutions whose readings were well
within the dynamic range of the standards. Final concentrations are adjusted to account for
dilution from original gel sample volume.

<u>Fluorescence Lifetime Imaging (FLIM)</u>: Time-correlated single-photon counting (TCSPC) FLIM
measurements were obtained using a custom laser-scanning two-photon rig driven by custom
ScanImage software (MBF Bioscience, Williston, VT) equipped with a MaiTai Ti:Sapph laser
(SpectraPhysics) and high-sensitivity PMT detectors (Hamamatsu C6438-01) as described
previously (Normoyle *et al.*, 2024). Briefly, ABP-dextran was excited at 760nm and emitted
photons were subjected to a 445/58 bandpass filter (Chroma) prior to detection using a high-
sensitivity PMT. A fluorescence lifetime is calculated for each pixel by measuring the time it
takes for an emitted photon to be detected, binning these photons according to arrival time, and
to these data a first-order exponential equation is fit to derive the rate constant representing the
fluorescence lifetime (Gehlen, 2020). The presence of a quencher will shorten fluorescence
lifetime to an extent which is linearly related to quencher concentration through the Stern-
Volmer equation,

$$\frac{\tau_o}{\tau_i} = 1 + [Q]k_{SV}$$

where τ_o / τ_i represents the unquenched lifetime divided by the shortened lifetime at a given
quencher concentration ($[Q]$) (Gehlen, 2020). The Stern-Volmer constant (k_{SV}) of the resultant
line relates the concentration of quencher to the shortening of fluorescence lifetime. Photons
were collected over an acquisition period of 60 seconds (*in vivo* and agarose gels) to 90
seconds (*in vitro* slice cultures) and the lag between incident laser pulse and resulting photon
detection recorded. The population of photons in each individual pixel is fit to an exponential to
yield a lifetime value and summed to yield an intensity value using the FLIMJ plug-in for ImageJ.
Further image processing, calculations, and statistics were accomplished using built-in functions
and custom routines developed in Matlab (v2018b and v2022a; Mathworks, Natick, MA). For
composite image generation, see *Image processing in Matlab* section below.

<u>Calibration of ABP-dextran:</u> ABP-dextran is synthesized, conjugated, and calibrated as
described previously (Normoyle *et al.*, 2024). Briefly, we calibrated ABP-dextran against aCSF
with varying chloride concentrations by preparing standard aCSF (in mM: NaCl 126; KCl 3.5;

CaCl$_2$ 2; MgCl$_2$ 1.3; NaH$_2$PO4 1.2; Glucose 11; NaHCO$_3$ 15) and a low-chloride aCSF in which
NaCl was replaced with sodium gluconate. We mixed appropriate amounts of each to vary
chloride concentrations (10-136mM) while maintaining the sum of sodium chloride and sodium
gluconate at 136mM. Although k_{SV} values did not differ appreciably between ABP-dextran
batches nor when measured using different FLIM-capable microscopes, out of an abundance of
caution we used a single batch of ABP-dextran and obtained FLIM calibration parameters
specific for this batch of fluorophore and the custom microscope used for all experiments
reported in this manuscript. Because calibrations done in aCSF have a minimum Cl$^-$ of 10mM as
described, the unquenched lifetime (τ_o) used for Stern-Volmer calculations is approximated as
the y-intercept (Cl$^-$ = 0mM) of a best-fit curve of the raw calibration data; empirically, a
biexponential fit was used. Each concentration of Cl$^-$ in the calibration curve has a normally
distributed set of ABP-dextran lifetime values with a near-constant relative standard deviation
(RSD = standard deviation divided by mean) (Normoyle *et al.*, 2024). All calculations are
otherwise done as described under *Fluorescence Lifetime Imaging (FLIM)* section above.

<u>Animals</u>: All animal protocols were approved by the Massachusetts General Hospital
Institutional Animal Care and Use Committee. Wild type mice (C57bl/6; Jackson Labs 000664)
of either sex were used for this study. Mouse pups remained in the home cage with the dam
under standard husbandry conditions until postnatal day 6 to 8 (P6–8) when organotypic slice
cultures were prepared or until a cortical window was surgically placed at P26-30.

<u>In vitro organotypic slice cultures and imaging:</u> Organotypic hippocampal slice cultures were
prepared either as glass-mounted (Berdichevsky *et al.*, 2016) or membrane insert-mounted
(Stoppini *et al.*, 1991) cultures. Briefly, in either case hippocampi are obtained from P6-8 mice
and cut to 400μm thick slices. These are then gently transferred to a 6-well dish containing
either a membrane insert (Millipore) or a poly-L-lysine coated coverslip (Electron Microscopy
Sciences), are fed twice weekly with 1mL neurobasal-A media supplemented with 500μM
Glutamax, 2% B-27, and 0.03mg/mL gentamycin (all from Invitrogen), and are incubated at 35C
in a 5% CO$_2$ incubator. Cultures are typically imaged between DIV 12-21 in aCSF warmed to
33°C and bubbled with 95%O$_2$ / 5% CO$_2$ perfused at a rate of approximately 100mL/hour unless
otherwise noted. Slices were pretreated with 500μg/mL ABP-dextran for 1-2 hours prior to
perfusing with aCSF containing 136mM chloride and the same concentration of ABP-dextran.
Slices were allowed to equilibrate in perfusate for 20-30 minutes before imaging, equaling at
least 2 hours of total exposure to ABP-dextran prior to initiation of imaging. Two photon images
were captured on a perfusion-enabled stage with a 20x water-immersion objective, NA 0.95,
and analyzed as described (see *Image processing in Matlab* section below).

In vivo murine cortical window imaging: Using a modified method based on that of Che et
al.,(Che & De Marco García, 2021) we placed cortical windows in young adult mice (P26-30) for
acute (non-survival) imaging in accordance with Massachusetts General Hospital Institutional
Animal Care and Use Committee policies and procedures (protocol 2018N000221). Briefly,
anaesthetized mice were immobilized with standard ear bars and nosepiece until custom
headbar is placed. A section of scalp is removed to expose a roughly 1cm section of skull.
Acrylic dental cement powder (Lang Dental, Wheeling IL) mixed with cyanoacrylate adhesive is
used to fix the headbar to the exposed skull at 4 o'clock to Bregma, creating a 5mm diameter
working area just lateral to midline. A 2.5mm round section of skull and underlying dura is
removed and a 3mm No. 1 coverslip is placed over the exposed cortex. Coverslip has 10μL
concentrated ABP-dextran / agarose mixture (1%w/v low gelation temperature agarose, Sigma-

Aldrich, with 10mg/mL ABP-dextran in aCSF) pipetted from 42°C heat block immediately before
inversion and placement over cortex. One hour is allowed before image acquisition for ABP-
dextran diffusion into cortex from overlying agarose (between cortex and coverslip) during which
time the warmed breadboard with immobilization apparatus and anesthetized mouse is
transferred to the microscope table for *in vivo* imaging. This custom-made gantry-type
microscope accommodates the breadboard as a stage for *in vivo* imaging. Two photon images
were captured with a 16x water-immersion long focal distance objective, NA 0.80, and were
analyzed as described (see *Image processing in Matlab* section below).

Agarose or agar gel Cl⁻ determination using ABP-dextran FLIM: Cl⁻ within an agar or agarose
gel was measured using the two-photon rig described above in the *in vivo* section, except with a
4x (NA 0.28) objective. Using an 8-well microchamber with silicone dividing gasket (Ibidi,
Grafelfing, Germany), gels were cast in aCSF containing ABP-dextran and subsequently
transferred to a single Petri dish where they were equilibrated with aCSF containing ABP-
dextran for at least 1 hour prior to imaging. FLIM measurements were taken as described,
generally one field per quadrant at a depth of between 50-100microns. Importantly, the gasket
separating gels of differing compositions was removed such that all gels in an experiment were
bathed in the same aCSF with the same Cl⁻. Where gels were kept for more than 1 day, they
were prepared in aCSF without glucose and kept refrigerated between readings to prevent
fouling.

Chondroitinase treatment of organotypic hippocampal slice cultures: Chondroitinase ABC
(ChABC; AMSBIO, #E1028-02) was brought up in aCSF containing 0.1% bovine serum albumin
(BSA) to a concentration of 1U/mL. Slices grown on membrane inserts were treated for 1hour
either with aCSF containing ChABC or sham aCSF containing only the 0.1% BSA vehicle. Both
experiment and control had 0.5mg/mL ABP-dextran (50μM), and FLIM imaging was done using
the two-photon rig described in the *in vivo* section above. At the conclusion of imaging, the
aCSF was collected for detection of liberated chondroitin sulfate.

Measurement of sulfated glycosamioglycans (sGAGs) liberated in aCSF: Blyscan 1,9-
dimethylmethylene blue-based kit (Biocolor, #B1000) was utilized for measurement of sGAGs
including chondroitin sulfate proteoglycans (CSPGs). After treatment with aCSF containing
ChABC or BSA vehicle as detailed above, aCSF was collected and measured as directed
against standards of bovine trachea chondroitin (Blyscan kit standard) with appropriate amounts
of 10x stocks of aCSF and 0.1% BSA to obtain concentrations identical to samples. 1,9-
Dimethylmethylene blue reagent absorbance is read at 650nm.

Measurement of chondroitin sulfate by liquid chromatography / mass spectroscopy (LC-MS):
aCSF containing either chondroitinase or vehicle was recovered after incubation with slice (see
above), volume recovered carefully noted before freezing at -20°C, and shipped over dry ice
with blinded labels for analysis. The samples were filtered using 3K MWCO spin filter units (Pall
Corporation Nanosep with 3K Omega Ref: OD003C34) at 14000 RPM at 10°C for 20minutes.
The flow through containing CS disaccharides was dried in speed vac. The dried flow through
were reconstituted in 200µl ultrapure distilled water (Invitrogen, Cat # 10977-015) and 100µl
was dried taken for $^{12}C_6$ aniline tagging by reductive amination method (Lawrence *et al.*, 2008).
Briefly, the CS disaccharides were dissolved in 17µl of $^{12}C_6$ aniline followed by addition of equal
volume of freshly prepared 1 M NaCNBH$_3$ (Sigma-Aldrich) in dimethyl sulfoxide: acetic acid
mixture (65:35, v/v). The reaction mixture was kept at 70°C for 45min and then transferred to

dry oven at 37°C for overnight reaction (16 h). After aniline tagging samples were dried in
speed-vac for 48h to remove excess reagents. The samples were then dissolved in 17µL of LC-
MS grade water (Honeywell), centrifuged at 7000g for 5min. 5µL of the sample was taken and
mixed with 5µL reconstitution buffer containing known amount of internal standard (5µl of the
reconstitution buffer contains, 1 µl of 10X LC-MS buffer-A (80 mM acetic acid and 50 mM
dibutylamine; Sigma-Aldrich), 2 µl of $^{13}C_6$ aniline-tagged CS disaccharide internal standard
mixture (20pmole each) and 2 µl LC-MS grade water. CS disaccharides were analyzed by LTQ
Orbitrap Discovery (Thermo Scientific) mass spectrometry attached with liquid chromatography
system (Ultimate3000, Thermo-Dionex). All disaccharides were separated on a reverse phase
C18 column (1mm x 150mm, 5µm, Targa, Higgins Analytical) using buffer containing dibutyl
amine as ion-pairing reagent, followed by detection of disaccharides on mass spectrometer in
negative ionization mode. The chromatographic steps were: 100% buffer-A (8mM acetic acid,
5mM DBA) for 10 min followed by stepwise increase of buffer-B amount; 17% buffer-B (70%
methanol, 8 mM acetic acid, 5 mM DBA) for 10 min; 32% buffer B for 15 min; 40% buffer B for
15 min; 50% buffer B for 15min; 60% buffer B for 15 min; and 100% buffer A for 10 min. Flow
rate was kept at 50µL per min.

<u>Calculation of ECM sulfate lost after chondroitinase digestion</u>: In order to contextualize the Cl^-_o
increase we observe after chondroitinase treatment of HOSCs, we used our HPLC-derived data
and approximated HOSC volume to derive a reasonable expected Cl^-_o value for comparison.
HPLC data was obtained from collected chondroitinase (or sham) digestion aCSF of a known
volume. Chondroitin disaccharide concentration and sulfate content were found analytically, and
a total chondroitin and total sulfate amount were calculated from mean values of chondroitin
sulfate content (0.2mg/mL), sample volume (0.725mL), sulfation ratio (0.95 sulfate per
disaccharide), and a monosulfated disaccharide molar mass of 481.4g/moL. Given mean HOSC
thickness of 266µm and typical HOSC dimensions of 3x1.5mm, we calculate the best
approximation of extracellular volume (20% of total slice volume) to be 0.293mL. The organic
sulfate content lost from total ECS volume comes out to 1.65mM when assuming full-depth
uniform chondroitinase digestion, or 8.8mM assuming depth-dependent digestion affecting only
50µm nearest the chondroitinase reservoir (i.e., nearest the perfusate). For comparison to
measured Cl^-_o, we assume the molar relationship between sulfate lost and chloride gained to be
1:1.

<u>Image processing in Matlab</u>: ABP-dextran fluorescence lifetime is converted to Cl^-_o as described
above. Composite Cl^-_o images, depicting Cl^-_o values via colormap while incorporating structural
data through brightness adjustment, were produced using custom Matlab routines that apply
normalized intensity data to Cl^-_o images in a manner similar to commercial FLIM software. Note
that intensity images were generated by summation of photons for each pixel across 32 time
bins after excitation, resulting in a higher intensity background than is typical in standard two-
photon images.

Pixel exclusion: We sought to perform as little processing of either intensity or lifetime data as
possible, using statistical methods to objectively exclude pixels greater than three standard
deviations from the mean of either property. Any remaining pixels reflecting a fluorescence
lifetime outside the calibrated range (0-150mM Cl^-), as happens in pixels with poor data quality
typically due to low signal, were also excluded. More specifically, pixels were excluded if ABP-
dextran lifetimes were in excess of the mean at 0mM chloride, or if lifetimes were shorter than
twice the standard deviation less than the mean at 150mM (calculated from calibration data).

This yielded total objective exclusion of 0.1-0.5% of pixels in a given image, which were set to
black. Subjective exclusions were done either by manually drawn regions of interest (ROI) to
exclude areas of perfusate at the margins of HOSCs or large blood vessels *in vivo*, or else by
image-specific thresholding to exclude low-intensity areas coinciding with apparent somatic
silhouettes. All exclusions were performed prior to analysis of extracellular Cl^-_o. To ensure that
somatic silhouette-defined pixels, in particular, were not a subpopulation of pixels possibly
reflecting ABP-dextran in an atypical compartment (e.g., the intracellular space), multiple
analyses of excluded versus included pixels were performed to confirm faithful extracellular
ABP-dextran signal and appropriateness of pixel exclusion (see Results).

Neuronal tracing analysis: Where somatic borders were traced for analysis, these were not
dependent upon thresholding but were traced manually where neurons within the focal plane
could be confidently identified based upon morphology. Traces followed the exterior aspect of
the visually apparent border between bright and dim signal (see Results). Data was linearized
and is presented as the pixelwise change in Cl^-_o versus linearized distance, the latter calculated
from pixel centerpoints. Comparison control data is derived by applying the same traces over
calibration solution images, in which stochastic pixel to pixel differences are isolated (e.g.,
machine noise, Brownian motion artifact, etc).

Regions of interest (ROIs) selection: ROIs are defined using intensity images for HOSCs either
as a differentiation between tissue and surrounding bath, or this plus a regular division of the
field (i.e., quarters or eighths) as indicated. ROI fields containing a majority of excluded pixels
(e.g., perfusate) are excluded from analysis. ROIs are defined *in vivo* as cortical regions
between anatomical landmarks, such as large vessels, which are easily identified through their
exclusion of ABP-dextran. ROIs are then converted to masks and applied to fluorescence
lifetime or Cl^-_o images. ROIs in fields in which somatic silhouettes are evident are distinguished
from ROIs that are too superficial to contain discernable silhouettes such as cellular (Layer 2/3)
or acellular (Layer 1) cortical fields, respectively.

<u>Statistics</u>: Statistical analysis was done with Matlab 2022a, with access to statistical toolbox
functions. Individual tests used and *p*-values are detailed in figure legends. Corresponding *p*-
value levels are indicated within figures by asterisks and listed in figure legends. *In vivo*
statistical analysis of cortical Cl^-_o was accomplished with ROI mean values, but because ROI
size and number varies between groups we report mean+/-standard deviation of all analyzed
pixels in addition to the ROI mean+/-standard error associated with ANOVA analysis. In the
interest of transparency we elected to display violin plots of all pixels analyzed overlaid with ROI
mean+/-standard error, using ANOVA analysis of the latter to determine significance of findings.

<u>Data availability statement</u>: All data files used to generate the figures in the paper can be found
on GitHub under KieranNormoyle/NeuronalChloride2023.

Results

<u>Non-invasive measurement of extracellular chloride using ABP-dextran</u>: ABP is a novel Cl-
sensitive fluorophore compatible with <u>F</u>luorescence <u>L</u>ifetime <u>IM</u>aging (FLIM) that we have
optimized to be bright, red-shifted, and sensitive to chloride concentrations through the range
that would be encountered in in the extracellular space. We conjugated ABP to a 10,000 Dalton
(10KDa) dextran resulting in an amine-bonded moiety (Figure 1A) that is resistant to pH
changes in the physiological range (Normoyle *et al.*, 2024) and is restricted to the extracellular
space (see below). We validated the ability of ABP-dextran to reliably report chloride

concentration by correlating the shortening of ABP-dextran fluorescence lifetime to chloride
concentration (Figure 1B) through the Stern-Volmer relationship (Figure 1C), generating a k_{SV} of
15.9M^{-1} for ABP-dextran used in these experiments. The fluorescence lifetime of ABP-dextran is
not dependent upon the concentration of ABP-dextran, only the concentration of chloride.
Chloride concentration also affects ABP-dextran intensity, such that high chloride
concentrations not only shorten the emission lifetime but also reduce the intensity of emission.
However, the concentration of chloride can be calculated (independently of the local ABP-
dextran concentration) as long as sufficient photons are collected to calculate fluorescence
lifetime in a given pixel.

Sulfated agarose gels as an *in vitro* model of perineuronal space: We recently observed that
differing concentrations of chloride juxtaposed to one another were stable on time scales of at
least 100 minutes due to the presence of immobile anionic charges present in the neuronal
cytoplasm (Rahmati *et al.*, 2021), an effect known as Donnan exclusion (Marinsky, 1985b;
Helfferich, 1995; Fatin-Rouge *et al.*, 2003; Epsztein *et al.*, 2018; Gao *et al.*, 2022). We tested
whether Donnan exclusion was also operative in the extracellular space as a consequence of
the fixed negative charges on the sulfate moieties of the glycosaminoglycans that comprise the
bulk of the extracellular matrix (Karamanos *et al.*, 2021). This matrix is most frequently
visualized as the perineuronal lattice-like structures found around all neurons and first described
by Golgi in 1898 (Golgi, 1898; Celio *et al.*, 1998). The process of Donnan exclusion does not
require an ongoing energy input, although this issue has been contested (Glykys *et al.*, 2014b;
Voipio *et al.*, 2014). We therefore modeled the brain's extracellular matrix using a reduced
system to remove the possibility of active buffering of Cl^-_o by astrocytes (Egawa *et al.*, 2013) or
other cellular or vascular elements. We used gels of sulfated agarose to mimic the sulfated
glycosaminoglycans such as the chondroitin sulfate-decorated aggrecan family members that
are prominent in the perineuronal matrix. Agarose is a nominally sulfate-free linear glycopolymer
that is a major component of agar (Figure 2B; bars 1 and 2) that was used as a chemically
homogenous starting material for covalent sulfate linkage. We optimized the method of Fuse
and Suzuki (Fuse & Toshiya, 1975) to balance the gelatinizing properties of agarose with the
dispersive, hydrophilic Donnan effects arising from incorporation of the charged sulfate groups
(Gregor, 1951; Helfferich, 1995). We obtained a sulfated agarose that could be used to cast
gels with 50 mM effective sulfate concentration (Figure 2B, bar 3; 49.8+/-0.9mM sulfate across
three batches [mean+/-SD]) covalently linked to the polymerized agarose lattice. Gels cast from
sulfated agarose serve as a simple *in vitro* model of the perineuronal extracellular space,
allowing us to investigate the ability of covalently linked anions attached to a three-dimensional
matrix to displace mobile chloride ions.

We measured the chloride concentrations within non-sulfated agarose gels vs. sulfated agarose
gels, each equilibrated in aCSF, and compared these to the chloride concentration in the
surrounding aCSF using either a standard colorimetric iron(III) thiocyanate assay (Figure 2C,
dark gray bars) or ABP-dextran FLIM imaging (Figure 2C, light gray bars). Agarose gels with
fixed anions in the form of covalently bound sulfate had significantly decreased chloride
concentrations versus both control aCSF and non-sulfated agarose gels (Figure 2C, right pair vs
left and center pairs, respectively) measured either with the standard chemical assay (dark gray
bars; $p = 0.00264$ and $p = 0.00354$ by Student's paired *t*-test, respectively) or ABP-dextran FLIM
(light gray bars; $p < 0.001$ by Student's paired *t*-test for either comparison). We measured a
small chloride increase within non-sulfated agarose gels vs control aCSF in either assay,
representing an insignificant increase when using the standard chemical assay (8.05+/-6.14mM,

mean+/-standard deviation; p = 0.190 by Student's paired t-test) and a smaller yet statistically
significant increase when measured using ABP-dextran FLIM (4.95+/-0.51mM, mean+/-
standard deviation; p = 0.00328 by Student's paired t-test). Later ABP-dextran FLIM
experiments comparing aCSF perfusate to non-sulfated gels show no such difference (see
below and Figure 2E). This Cl^- increase is curious but is quantitatively small and qualitatively
distinct from the Cl^- decrease we see in sulfated agarose gels (3.5% increase from non-sulfated
gel vs. aCSF control compared to a 40.7% decrease from sulfated agarose gel vs. aCSF
control, respectively), meaning that any effect would be a small overestimation of Cl^- within
sulfated agarose gels. The overall agreement of results derived from both assays lends
confidence to the reliable performance of ABP-dextran FLIM versus standard chemical
colorimetric methods of Cl^- measurement.

Next we placed separately cast non-sulfated and sulfated gels adjacent to each other and
equilibrated in the same aCSF bath with ABP-dextran (Figure 2D; schematic in Figure 2A).
ABP-dextran FLIM measurements of chloride concentration within the non-sulfated and sulfated
gels demonstrated significant chloride concentration differences between gels that had no fixed
anionic charge to displace mobile chloride (non-sulfated agarose; Figure 2D light blue bars) vs.
gels that had covalently bound sulfate groups (sulfated agarose; Figure 2D dark blue bars). The
chloride concentrations were inversely proportional to the sulfate concentration in the gel, were
significantly lower for sulfated gels at each time point (each p < 0.001), and were stable for over
three weeks (agarose gel comparisons: each p = 1.0000; sulfate gel comparisons: 2-4hours vs
24hours p = 1.0000, 3weeks vs 2-4 hours and 24 hours p = 0.211 and p = 0.113, respectively;
ANOVA with Bonferroni correction). These findings support the displacement of mobile chloride
ions by the sulfates fixed to the gel (Figure 2D) by Donnan exclusion. We next imaged both non-
sulfated and sulfated gels immediately juxtaposed to one another, along with aCSF perfusate, in
the same field (Figure 2E) and demonstrated that the presence of sulfate groups covalently
bound within the polymerized gel results in a significantly decreased chloride concentration
relative to both non-sulfated gels and aCSF perfusate (Figure 2E, bar 3 vs bar 2 and bar 1,
respectively; each p < 0.001 by Student's paired t-test). Non-sulfated agarose gels
demonstrated no difference versus aCSF perfusate (p = 0.0647 by Student's paired t-test).

<u>Demonstration of chloride displacement in living systems:</u> To test whether Donnan exclusion is
also operative in the extracellular matrix of the brain, we studied mouse hippocampal
organotypic slice cultures. We tested whether extracellular perineuronal fixed anions, such as
the chondroitin sulfate moieties of aggrecan family proteoglycans (Frantz et $al.$, 2010;
Karamanos et $al.$, 2021), displace chloride in this living system. After incubation with ABP-
dextran and equilibration with aCSF containing ABP-dextran and 136mM chloride, we measured
extracellular chloride concentration using ABP-dextran FLIM and found that Cl^-_o was
unexpectedly low and heterogenous (46.7 +/- 18.8 mM, calculated from 25.8×10^6 pixels across
32 fields of 7 slices). Representative fields are shown in Figure 3A (ABP-dextran intensity) and
Figure 3B (Cl^-_o composite image, quantified in Figure 3C). Paired analysis of slices pre- and
post-ABP-dextran incubation confirmed that autofluorescence signals in pixels in pre-ABP
images did not contribute to the unexpectedly low values of Cl^-_o obtained from post-ABP
images. This is because pre-ABP pixels that had both sufficient photon count to allow lifetime
calculation and lifetimes within the calibrated range of ABP-dextran had short lifetimes whose
effect would be to increase the reported Cl^-_o, not decrease it.

Perineuronal traces along the outer edge of neuronal silhouettes (Figure 3B, white dashes)
were used to analyze the spatial change in Cl_o in the extracellular space immediately adjacent
to the neuronal cytoplasmic membrane. Pyramidal neuron silhouettes were identified
morphologically and pixels underlying the extracellular margin (i.e., perineuronal) traces were
linearized for analysis. We compared the size of the effects of fixed anions on local Cl_o to
Brownian and machine noise arising from FLIM of ABP-dextran in bulk solutions with similar
mean Cl_o. The pixel-to-pixel change in Cl_o (ΔCl_o) versus distance in the presence of fixed
anions in the perineuronal space (Figure 3D, blue) was much larger than the ΔCl_o versus
distance in the absence of fixed anions in bulk solution containing ABP-dextran and a single
concentration of Cl^- (Figure 3D, red). Figure 3E shows similar data for a larger cohort of
perineuronal tracings and their corresponding aCSF solution counterparts (N = 72). We found a
consistently increased probability of greater ΔCl_o per micron in the extracellular space where
fixed extracellular anions were present compared to the perfusate aCSF (Figure 3E, blue vs
red).

We further analyzed dim pixels typical of the neuronal silhouettes, which had not already been
statistically excluded and displayed as black. These dim pixels retain their Cl_o colormap values
but are often so dim they are difficult to appreciate visually (e.g., Figure 3B). They are included
in presented images in the interest of transparency but were excluded from Cl_o analysis after
applying an empirically derived threshold value that best approximates the clearest silhouettes
in each image. We reasoned that these pixels represented photons originating from extracellular
ABP-dextran above or below the focal plane, likely present due to the higher-background FLIM-
derived intensity images (Figure 4A) which represent a summation of all photons in each pixel.
Because Cl_o is derived from the intrinsic, intensity-independent property of fluorescence
lifetime, structural context is lost in Cl_o images (Figure 4B) and must be reestablished by
incorporation of intensity information into a composite image (Figure 4C). This is not unique to
Cl imaging; commercial FLIM software creates intensity-lifetime composite images in a similar
manner. In our images we first converted from fluorescence lifetime to Cl_o to allow the use of a
linear colormap for readers' ease of interpretation (see Methods).

The dim, excluded pixels occurring in neuronal silhouettes offered an opportunity to assess
whether they were in fact out-of-focus, extracellular ABP-dextran. If true, this would produce a
similar, if wider (Normoyle *et al.*, 2024), distribution of Cl_o values. Conversely, if the dim,
excluded pixels were autofluorescent in origin or were reflective of low Cl_i reported by ABP-
dextran leaked into neurons, then we would find a right-shifted, high-Cl_o distribution or a left-
shifted, low-Cl_o distribution, respectively.

<u>Verification of ABP-dextran extracellular compartmentalization:</u> We used two approaches to
assessing the Cl values reported in the dimmest pixels. We first compared all pixels below the
exclusion threshold to all pixels above the threshold. Only pixels outside the actual slice (i.e.,
perfusate pixels) and those pixels with lifetimes outside the calibrated range of ABP-dextran
were not considered. We found that included and excluded pixels had histograms with similar
characteristics, such as mean and mode, although excluded pixels were fewer in number and
had a wider distribution (Figure 4D, blue vs gray). There is neither a high-Cl_o peak indicative of
autofluorescence nor is there a low-Cl_o peak indicative of intracellular signal in either included
pixels or excluded pixels. Notably, there are virtually no included pixels in the typically
intracellular range (i.e., $\approx$15mM Cl^-), and such pixels are far more likely to be excluded from
analysis (Figure 4D, gray arrow). We conclude that intensity-excluded pixels are not a low- or

high-Cl_o subset, which would seem to preclude errant Cl_i signal. However, we wanted to rule
out any possible non-extracellular signal more specifically as most pixels at these resolutions
are expected to contain a portion of both neuropil and extracellular space.

As a second means to evaluate the Cl values reported in the dimmest pixels, we assessed
pixels within individual somatic silhouettes. Visually apparent somatic edges are ideal to
compare extracellular ABP-dextran signal in the bright perineuronal exterior of silhouettes to
ostensibly intracellular pixels inside the silhouette margins. The fact that the brightest of these
pixels are located nearest the outer silhouette margin and the least intense pixels are located
toward silhouette centers supports our hypothesis that this signal is extracellular in origin from
outside the focal plane. Alternatively, if ABP-dextran leaked inside cells, despite its conjugation
to 10KDa dextran, we would expect that such an intracellular signal would be brightest in the
silhouettes formed by cells firmly within the focal plane (i.e., with the most ABP-containing
cytoplasm in the focal plane). After the objective pixel exclusion step (intensity greater than
three standard deviations from the mean, or lifetime outside the calibrated range), we compared
Cl_o in the perineuronal margin within one micron external to the traced border to those pixels
inside the traced border. In this way our analysis was restricted to only those cells
morphologically identified as pyramidal neurons adequately centered in the focal plane. Pixels
from inside the silhouette are greater in number than perineuronal extracellular pixels in this
analysis, yet we saw no subpopulation of intracellular chloride values (i.e., in the range of 15mM
Cl) that would be expected if this signal were originating from 'leaked' intracellular ABP-dextran
(Figure 4E). Signal from the one micron external to the perineuronal perimeter is five times as
bright on average as pixels from within the silhouette (Figure 4F), which helps explain why the
excluded intracellular signal has a wider distribution of Cl_o values (Normoyle *et al.*, 2024). If no
intracellular leakage of ABP-dextran can be detected even within in-focus neuronal somatic
silhouettes, where such signal is expected to be strongest if it were present, one can infer that
neuropil will not contribute any such signal in pixels containing both neuropil and extracellular
space. While we look forward to future experiments with greater spatial resolution, we conclude
that neither objectively excluded pixels nor those excluded by intensity thresholding are
indicative of low- or high-Cl_o subsets. Further, we provide ample evidence supporting the fidelity
of ABP-dextran in its reporting of chloride exclusively in the extracellular space.

<u>Demonstration of chloride displacement by chondroitin sulfate:</u> We then confirmed that fixed
anions, including chondroitin sulfate glycoproteins, are a significant contributor to this difference
by measuring Cl_o before and after treatment with chondroitinase. Analogous to matrix
metalloproteases, chondroitinase is a bacterial enzyme that cleaves the sulfated glycopolymer
chondroitin. Chondroitinase treatment significantly increased Cl_o in organotypic hippocampal
slice cultures (Figure 5A&B; pre- and post-digestion, respectively). Cl_o increased on average
8.3+/-3.3mM (mean+/-standard deviation) in freely diffusible extracellular chloride within the
slice culture (Figure 5A vs 5B; quantified in 5D-F) and was significantly greater in chondroitinase
digested slices than in sham digested slices (Figure 5B vs 5C; quantified in 5F). This increase
in Cl_o coincides with a significant decrease in the extracellular fixed anions within the slice
culture. The actual decrease in fixed anion concentration in the extracellular space was
estimated from the concentration of polysulfated chondroitin released into the perfusate after
chondroitinase treatment (Figure 5G) and confirmed by HPLC analysis (Figure 5H). We
approximated an expected Cl_o increase using HPLC-derived measurements of sulfated
chondroitin disaccharide and HOSC volume (see Methods). Our own empirical observations, in
agreement with others' observations (e.g. Orlando and colleagues, especially Figure 5C therein)

(Orlando *et al.*, 2012) indicate that chondroitinase activity decreases with distance from its
diffusive reservoir and full activity is restricted to an estimated 50μm; as such we only measure
Cl_o^- to a depth of 50μm. If all digested chondroitin sulfate indeed came from only the most
superficial 50μm of our HOSCs, the Cl_o^- increase expected in this same superficial 50μm is
calculated to be 8.8mM. This agrees well with experimental mean Cl_o^- values calculated from
regular subfields of pre- and post-chondroitinase digestion, 8.8mM versus 8.3+/-3.3mM (Figure
5E), or when using the full field to calculate the minimum and maximum range of Cl_o^- increases
from 0.6-16.1mM (Figure 5D; pre-digestion mean- standard deviation to post-digestion mean+
standard deviation). Further, we find that even if we assume chondroitin sulfate is lost uniformly
from the full thickness of the HOSC and recalculate a smaller expected Cl_o^- increase of 1.65mM,
this still lies within the empirically measured range of 0.6-16.1mM. These observations are
consistent with the expected inverse relationship between fixed extracellular anions such as
sulfate and mobile Cl arising from Donnan exclusion. It would be advantageous to similarly
estimate a predicted Cl_o^- based upon local sulfate concentrations and compare this to empirical
Cl_o^- measurements similar to our sulfated agarose experiments (Figure 2E), but without direct
and independent measurement of the extracellular sulfate content of HOSCs this is not currently
feasible. In summary, the immobile ions associated with the extracellular matrix displace mobile
anions such as chloride consistent with the well-established effects of Donnan equilibrium
forces, such that Cl_o^- is much lower and less uniform than would otherwise be expected.

<u>Chloride displacement in the perineuronal space *in vivo*</u>: To test whether Donnan exclusion of
Cl_o^- by sulfated GAGs was significant *in vivo*, we performed live 2 photon imaging of P26-30
mice via a cortical window (Figure 6A). Direct application of high concentrations of ABP-dextran
into the non-sulfated agarose overlying the neocortex (Figure 6B) resulted in a robust ABP
fluorescence signal from the extracellular space of the brain parenchyma. Control
measurements were obtained from the agarose above the cortex, which had been brought up in
aCSF (136mM Cl^-) and allowed to equilibrate with *in vivo* CSF. Notable exclusion of ABP-
dextran fluorescence emission from blood vessels (Figure 6C,D&F) and cell bodies (Figure 6C)
was observed. Areas of locally elevated Cl_o^- and overall bright signal seen in HOSCs (Figures 3-
5) were absent *in vivo* (Figure 6), suggesting these features are specific to *in vitro* preparations.
At several microns in size, we reason the areas seen in HOSCs are most likely damaged areas
where intracellular space makes up less than the typical 80% of volume allowing ABP-dextran
access to a greater proportion of volume, so that the signal is bright despite quenching by the
high Cl_o^- in these spaces. ECM damage would also be likely in such circumstances and would
be expected to result in a loss of extracellular immobile anions, limiting the Donnan exclusionary
force and leading to relatively high concentrations of Cl_o^-. This combination would be expected
to result in a paradoxically bright, short-lifetime ABP-dextran signal yielding an unusually intense
high Cl_o^- area. Given the absence of these features *in vivo*, we speculate that these areas are
the result of slicing trauma but longitudinal studies will be needed to adequately categorize
them. Regions of interest (ROIs) were defined using intensity images before rendering
composite images so that only structural information was considered; an example intensity and
composite image pair can be seen in Figure 6D&F. Cortical Cl_o^- was less than half the chloride
concentration measured in the overlying agarose (48.1 +/- 17.0mM, calculated from 68 ROIs
and 4 animals, vs 135.4 +/- 5.0mM in agarose between cortex and coverglass). Cl_o^- in cellular
and acellular fields displayed in Figure 6CD&F were not significantly different from each other
(Figure 6E entries 2&3, respectively), nor were either significantly different from all ROIs
collected from this animal (entry 4) or from all animals (entry 5; each p = 1.0000 by ANOVA with

Bonferroni correction), while all were significantly different when compared to aCSF measured
above the cortex (each $p < 0.001$ by ANOVA with Bonferroni correction). Regional mean Cl^-_o
values were approximately 45-55mM (Figure 6E) but there was notable spatial variation
between regions and even within regions, similar to *in vitro* HOSC results and consistent with
local Donnan exclusion of Cl^-_o by variably sulfated GAGs. Future experiments will be optimized
for higher resolution imaging, but our results suggest that the well-established heterogeneity of
sulfation patterns around neurons represent a means by which neurons and supporting cells
can indirectly modulate Cl^-_o through Donnan forces in a manner similar to how cytoskeletal
components, in particular actin, have been recently shown to influence local intracellular Cl^-_i
(Rahmati *et al.*, 2021).

Discussion

We demonstrate stable Donnan exclusion of chloride by fixed anions in a transport-free system
of sulfated agar using FLIM of a novel fluorophore, ABP-dextran, and validate these
measurements with direct chemical assays of sulfate and chloride (Figure 2). FLIM of ABP-
dextran demonstrated that Cl^-_o in the brain is less than half the Cl concentration in the
cerebrospinal fluid and plasma (Figures 3-6). This low Cl^-_o arises from Donnan exclusion of
chloride by fixed sulfated glycosaminoglycans (sGAGs), because release of sGAGs by
chondroitinase results in an increase in Cl^-_o that varies with the amount of sulfates released
(Figure 5). Major findings are confirmed *in vivo* (Figure 6).

Limitations of this study

The finding that Cl^-_o is much lower than expected in the extracellular space of living brain tissue
is based on FLIM of a new Cl-sensitive fluorophore, ABP-dextran. These results are in contrast
to the higher values measured using ion sensitive electrodes (Jiang *et al.*, 1992; Kroeger *et al.*,
2010). The ABP-dextran fluorophore is based on a validated structure (Biwersi *et al.*, 1992a;
Biwersi *et al.*, 1992b) and was calibrated against Cl, other anions, and direct chemical analysis
of chloride concentrations in situ. ABP-dextran FLIM accurately reported Cl values in sulfated
carbohydrate polymers (Figure 2), so it is unlikely that the sGAGs or other anions in the
extracellular environment (Figure 1C) affected the accuracy of the ABP-dextran FLIM. ABP-
dextran overcomes several barriers to accurate measurement of extracellular chloride, including
sufficient sensitivity in the relevant range of chloride concentrations, competition with
autofluorescence, and permeation into the cytoplasmic space, where lower chloride values
would be encountered. ABP-dextran was clearly excluded from the intracellular space (Figure
3A&B; Figure 4A&C; Figure 6C) and blood vessels (Figure 6C,D&E). We speculate that the
difference between our novel fluorophore-based measures and the ion-sensitive electrode
measures of Cl^-_o could arise from distortion of the extracellular milieu created by the stab wound
of the microelectrode, although our imaging techniques do not provide sufficient spatial
resolution to test this possibility. Alternatively, microelectrode-based chloride measurements
may be increased by electrode solutions leaking from the high-chloride barrel of the ion-
sensitive micropipette into the extracellular space (Figure S1). A study comparing fluorometric
and microelectrode techniques would benefit greatly from an ABP-based ratiometric chloride
sensor, currently in development (Normoyle *et al.*, 2024), and higher-resolution imaging.

Stable Donnan exclusion

An important finding in this study is that Cl microdomains established by Donnan exclusion are
stable for weeks, and do not require active transport to be maintained (Figure 2). This

phenomenon is well-established in biophysics (Epsztein *et al.*, 2018; Gao *et al.*, 2022), but is
more confusing in neuronal physiology where membranes with time-varying ionic permeabilities
and transporters create a more complex environment (Voipio *et al.*, 2014; Doyon *et al.*, 2016;
Düsterwald *et al.*, 2018; Rahmati *et al.*, 2021). A point of particular confusion is the role of
active Cl transport vs. Donnan exclusion in maintaining a particular E_{GABA} in the face of GABA-
gated Cl flux that alters the local Cl concentration. After a GABA-gated Cl influx, and re-
establishment of local charge balance by cationic flux via voltage- and ligand-gated cation
channels, the local cytoplasm will contain excess chloride salts. This Cl has two fates: either
excess salt diffuses through the cytoplasm to re-establish a new equilibrium with fixed charges,
or the excess salt is transported back across the membrane by cation-chloride cotransporters
(Brumback & Staley, 2008; Lewin *et al.*, 2012; Doyon *et al.*, 2016; Beckstein & Naughton, 2022).
The transporters are required to re-establish the original steady state volume, but in the
absence of transport, diffusion maintains local Cl concentrations remarkably well (Brumback &
Staley, 2008; Rahmati *et al.*, 2021). In the absence of membrane Cl salt transport, this diffusion
of excess Cl salts will be associated with either a cytoplasmic volume increase from associated
water influx or an increase in the mean Cl concentration in the cytoplasm. At pathologically high
rates of Cl influx, membrane chloride salt transport becomes rate-limiting (Staley & Proctor,
1999; Jedlicka *et al.*, 2011; Lewin *et al.*, 2012; Doyon *et al.*, 2016), and chloride concentrations
become sufficiently labile to depolarize E_{GABA} (Barker & Ransom, 1978; Alger & Nicoll, 1982;
Huguenard & Alger, 1986; Staley *et al.*, 1995). Thus Donnan exclusion *establishes* the local
value of E_{GABA}, and active transport *maintains* that value (and neuronal volume) in the face of
synaptic Cl flux (Glykys *et al.*, 2014a).   The same principals should apply to the local
extracellular chloride concentrations described here, with astrocytic chloride buffering (Egawa *et*
al., 2013) and diffusion between the extracellular fluid, the CSF, and vascular spaces
(Rasmussen *et al.*, 2022) taking the place of neuronal membrane cation-chloride cotransport.

<u>Low extracellular Cl</u>

The Cl^-_o measured by ABP FLIM is less than half of the canonical Cl^-_o. Such a low value of Cl^-_o
should affect not only E_{GABA} but also the inhibitory $GABA_A$ conductance, because there are only
half as many ions to carry the outward membrane currents (i.e. inward Cl- flux) (Coombs *et al.*,
1955). Shouldn't such a low value of Cl^-_o have been obvious from electrophysiological studies?
E_{GABA} is determined by the ratio of Cl^-_i to Cl^-_o (Coombs *et al.*, 1955; Staley, 2023). If either
concentration is known, the other can be determined from E_{GABA}. However if both Cl^-_i and Cl^-_o
are determined by the concentrations of local immobile anions, neither can be set
experimentally and thus E_{GABA} cannot be used to determine either Cl^-_i or Cl^-_o; only the ratio can
be inferred from the Nernst equation. Similarly, although the conductance of inward Cl flux
(outward membrane current) is determined in part by Cl^-_o, it is also determined by the $GABA_A$
permeability, i.e. the number of open membrane channels and their capacity for ion flux. Thus
membrane currents are not sufficient to determine Cl^-_o (Coombs *et al.*, 1955). Control of the
ionic milieu is greater in single-channel recordings, but whether the complement of immobile
anions is physiological under these conditions is difficult to determine, and even in these
conditions there is substantial uncertainty regarding local ionic concentrations (Li *et al.*, 2015).

<u>Cl^-_o microdomains</u>

The extracellular matrix of the brain is often depicted as perineural nets surrounding
parvalbumin neurons that are detected by specific agglutinins and antibodies (Celio & Blümcke,
1994). However, these assays detect very specific motifs of glycosaminoglycan sulfation; using

more permissive ionic methods, a highly anionic extracellular matrix can be seen to surround
virtually all neurons (Golgi, 1898; Glykys *et al.*, 2014a; Morawski *et al.*, 2015). This extracellular
matrix fills the extracellular space, which comprises about 20% of the brain's volume (Syková &
Nicholson, 2008; Tønnesen *et al.*, 2018). The sGAG moiety of proteoglycans are the primary
constituent of extracellular matrix (Tønnesen *et al.*, 2018; Karamanos *et al.*, 2021). GAGs are
polymers of widely varied length comprised of sugars and substituted sugars (N-
acetylglucosamine, glucuronic acid, and galactose) that are synthesized without templates. Up
to 3 sulfates are added to the sugar moieties before export to the extracellular space. This
enables the creation of a stable but highly variable spatial distribution of dense, fixed anionic
charge in the extracellular matrix. Figure 3 demonstrates that this variance is substantial at the
level of resolution of light microscopy. However, the distribution of fixed anions at the
extracellular face of $GABA_A$ receptors, and the corresponding Cl^-_o, is currently unknown.

Physiological implications

Cl^-_o microdomains in the region of the extracellular openings of $GABA_A$ receptor-operated
channels are a critical determinant of both E_{GABA} and the $GABA_A$ conductance. The wide range
of Cl^-_o values established via Donnan exclusion by sulfated GAGs (Figures 2, 3, 4, 5 and 6)
makes possible a substantial variety of effects of $GABA_A$ receptor activation based on the
direction of chloride flux. These effects range from hyperpolarizing inhibition through shunting
inhibition through excitation by activation of low threshold calcium currents and removal of the
magnesium block of the NMDA receptor. In addition, the effects of Cl^-_o on the quantity of charge
that is carried by $GABA_A$ membrane currents (the conductance) (Coombs *et al.*, 1955) makes
possible a broad range of strengths of $GABA_A$ synapses that is independent of the number or
subunit composition of receptors or their posttranslational modifications.

The turnover of the macromolecules comprising the extracellular matrix is very slow (Tsien,
2013), on the order of months to years based on rodent experiments (Margolis & Margolis,
1973) as well as the rate of deterioration of patients with genetic defects of GAG catabolism
(Héron *et al.*, 2011; Hampe *et al.*, 2020; Seo *et al.*, 2020). This slow turnover provides a
mechanism for long-lasting influences on synaptic GABA signaling. Some elements of GAG
synthesis are activity dependent (Sidharthan *et al.*, 2013), but currently nothing is known about
the insertion of GAGs into the extracellular matrix, in situ modification, and the capacity for
directed alterations of Cl^-_o microdomains in the regions of $GABA_A$ receptors and synapses. Thus
the role of extracellular chloride microdomains in synaptic signal processing and memory
remains to be discovered.

Pathological implications

Cytotoxic cerebral edema: If Cl^-_o is normally only half of the value of Cl in CSF and plasma, then
the capacity for Cl^-_o to increase acutely may be an important and hitherto unappreciated
element of the pathophysiology of cerebral edema (Glykys *et al.*, 2017). Matrix
metalloproteinases (MMPs) are released and activated after neuronal injury and death (Zhang
et al., 2016). Release of sulfates in sGAGs from the extracellular matrix would reduce the
Donnan exclusion of Cl by fixed sulfates. This would result in an acute increase of Cl^-_o as in
Figure 4. The action of equilibrative cation-Cl cotransporters would then lead to a
corresponding increase in chloride salts in the cytoplasm of neighboring healthy neurons, with
attendant volume increases. This novel mechanism of cytotoxic edema could potentially be
ameliorated by pharmacological MMP inhibition. Alternatively, inhibition of the influx of chloride

salts that cross the BBB from the vascular space (Jha *et al.*, 2019) would limit the influx of Cl
into the extra and intracellular spaces.

Post Traumatic Epilepsy: Richard Miles and colleagues demonstrated that $GABA_A$ receptor
activation could be excitatory in human brain tissue resected for control of medically intractable
epilepsy (Cohen *et al.*, 2002). However, the pathomechanism has never been elucidated
(Huberfeld *et al.*, 2007; González, 2016; Karlócai *et al.*, 2016). Gliosis is the most common
finding in medically intractable epilepsy (Thom, 2014). In glial scars, reactive astrocytes
synthesize new matrix elements (Roll & Faissner, 2014; Song & Dityatev, 2018). If the
replacement matrix was more heavily sulfated than the original, as would be expected from the
firm texture that is responsible for the name "scar" (Lesperance *et al.*, 1992), then in areas of
gliosis Cl_o^- would be low. Low Cl_o^- could contribute to degradation of $GABA_A$ receptor mediated
inhibition, frank excitation, and failure of anticonvulsants whose action are predicated on intact
synaptic inhibition. Thus replacement of the extracellular matrix and distortion of extracellular
chloride microdomains could comprise an important mechanism of medically intractable
epilepsy after brain injury.

Future studies

The findings that Cl_o^- is both lower than expected and spatially heterogenous opens the door for
many investigations into the mechanisms by which Cl_o^- is set, such as the synthesis, export, and
in situ modification of sulfated GAGs. The activity dependence of these mechanisms and the
effect of Cl_o^- microdomains on synaptic signaling are additional areas of investigation.
Pathologically, the impact of changes in Cl_o^- in both acute edema and long-term complications
such as epilepsy may provide much-needed therapeutic insights into these frequently-
intractable conditions.

Figure 1 Characterization and Calibration of Chloride-Sensitive Fluorophore ABP. *A)* N-
(**a**mmonium)-**b**utyl **p**henanthridine (ABP) is synthesized in-house and conjugated to activated
dextran, a glucose polymer, at a ratio of 10-14 fluorescent molecules per 10KDa Dextran. *B)*
Representative calibration of ABP-dextran fluorescence lifetime versus Cl⁻ demonstrating ABP-
dextran sensitivity to chloride-induced quenching. Calibration of ABP-dextran fluorescence
lifetime vs Cl⁻$_o$ is displayed as a violin plot with mean +/- standard deviation overlaid. ABP-
dextran lifetime at 0mM chloride (t_o) is extrapolated using a best-fit approach (see Methods). *C)*
Stern-Volmer plot demonstrates a linear relationship between ABP-dextran fluorescence lifetime
and concentration of chloride and was used to convert lifetimes to chloride concentrations. The
ratio of unquenched ABP-dextran fluorescence lifetime (t_o) to ABP-dextran lifetime at a given
concentration of quencher (t) is plotted versus concentration of quencher (t_o/t vs [Q⁻]). The slope
of the linear regression is the Stern-Volmer constant (k_{SV}) which allows conversion of ABP-
dextran lifetime to concentration of quencher, specifically Cl⁻ (blue circles / black line;
k_{SV}=15.9M⁻¹). Four biologically relevant anions were found to have no experimentally relevant
effect on ABP-dextran fluorescence lifetime: bicarbonate (k_{SV}=1.21M⁻¹), gluconate (k_{SV}=0.70M⁻
¹), methyl sulfonate (k_{SV}=0.18M⁻¹), and phosphate (k_{SV}=0.11M⁻¹). Gluconate was subsequently
used as a chloride substitute in non-standard aCSF formulations used for calibrations.

Figure 2 Effects of immobile sulfate on Cl⁻ in a volume of agarose gel. *A)* Experimental
schematic demonstrating how polysaccharide gels, cast in aCSF separately, were equilibrated
in aCSF after transfer to a common vessel. Gels were either equilibrated with aCSF containing
ABP-dextran, processed for chemical analysis, or subjected to multiple investigations in tandem
(see Methods). ***B)*** We covalently bound sulfate groups to algal biopolymer agarose and cast
this as a gel in aCSF to model perineuronal interstitial matrix that is rich in chondroitin sulfate-
decorated glycosaminoglycans. We compared sulfated agarose gel to native agar and agarose
gels, achieving a 250-fold increase in sulfate content compared to commercially available agar
(bar 3 vs bar 1, respectively; *p* < 0.001, Student's paired *t*-test; note discontinuous y-axis).
Commercially available agarose, which is intended as a sulfate-free product, had no
measurable sulfate content (bar 2) and subsequently was used as a control gel. ***C)*** Chloride
concentration was reduced in sulfated agarose gels. Cl⁻ was measured using both a standard
colorimetric assay (dark gray bars) and ABP-dextran FLIM (light gray bars) in aCSF control
(leftmost pair), within non-sulfated agarose gel (center pair) and within sulfated agarose gel
containing ≈50mM (49.2+/-0.289mM) covalently bound sulfate (rightmost pair). Measured
chemically or using ABP-dextran FLIM, Cl⁻ in sulfated agarose gels differed significantly from
aCSF (*p* = 0.00264 and *p* < 0.001, respectively) and non-sulfated agarose gels (*p* = 0.00354
and *p* < 0.001, respectively), while Cl⁻ measured in non-sulfated agarose gels was not
significantly different from aCSF when measured by chemical assay (*p* = 0.190) but registered a
small but significant difference (3.5% increased Cl- relative to aCSF, *p* = 0.00328) when
measured with ABP-dextran FLIM, each compared using Student's paired *t*-test. ***D)*** Significant

chloride concentration differences between agarose gels with 50mM covalently bound sulfate
(dark blue) and non-sulfated agarose gels (light blue) do not equilibrate with time but instead are
stable for as long as experimentally observed (> 3 weeks). ANOVA testing with Bonferroni
correction demonstrates that Cl^- significantly differs between native agarose and sulfated
agarose at all time points (overall and each pair $p < 0.001$) while neither agarose (each $p =$
1.0000) nor sulfated agarose gels (from left to right, $p = 1.0000$; $p = 0.2119$; $p = 0.1129$) change
significantly with time. *E)* Demonstration of stable juxtaposition of significantly different chloride
concentrations at equilibrium. Separately cast sulfated and non-sulfated gel pieces are placed in
a single vessel as described in *A* and equilibrated in common aCSF with ABP-dextran. Sulfated
agarose gels (dark blue) again contain significantly lower Cl^- than do non-sulfated agarose gels
(light blue; $p < 0.001$) or perfusate aCSF (gray; $p < 0.001$). Cl^- within non-sulfated agarose gels
is again not significantly different than perfusate aCSF ($p = 0.0647$), all analyzed using
Student's paired *t*-test. *Statistical significance indicated on graph for p-values <0.05 (*); <0.01*
(**); <0.001 (***); "ns" indicates "not significant". Error bars reflect standard deviation from the
mean.

Figure 3 Cl⁻ in the extracellular space is not uniform and is considerably lower than aCSF perfusate. *A)* High-magnification HOSC intensity image of ABP-dextran excluded from cell bodies in the pyramidal cell layer (CA1; scale bar 20μm). *B)* Chloride-composite image again demonstrating the exclusion of ABP-dextran from intracellular spaces as well as the non-uniformity of Cl_o in the pyramidal cell layer of a representative HOSC. Representative pyramidal cell visible just above and left of image center, identified by ABP-dextran exclusion and morphology of resulting silhouette, is one of 6-10 neurons per field deemed to have a regular and nearly-continuous border analyzed as described in *D&E* (scale bar 20μm). *C)* Histogram of full field Cl_o in HOSC pictured in *B*, demonstrating non-uniformity visually apparent in *B*. Notably, Cl_o = 44.5+/-15.9mM despite perfusate $[Cl^-]$ = 136mM (across all depths of this slice Cl_o = 49.7+/-18.5mM; mean +/- SD). *D)* Change in chloride ($\Delta[Cl^-]_o$) plotted against distance in the perineuronal space (blue line) or the same trace overlaid onto a calibration aCSF FLIM image (red line) corresponding to the perineuronal trace in *B*. $\Delta[Cl^-]_o$ of calibration aCSF (red) encapsulates machine and experimental noise, and $\Delta[Cl^-]_o$ in the perineuronal space (blue) exceeds that of control in this representative neuron. *E)* Perineuronal $\Delta[Cl^-]_o$ per micron calculated from many neurons' circumferential traces demonstrates that the variability of perineuronal Cl_o in living *in vitro* system exceeds the variability of control aCSF (72 neurons versus matched calibration aCSF controls from depths of 20-100μm in this representative HOSC). *Cl_o image rendering and presentation details can be found in Methods and in subsequent figures. Scale bars A&B: 20μm.*

Figure 4 Generation of composite chloride images and cell body masking for pixel quantification. ***A)*** ABP-dextran intensity data of an HOSC, derived by taking the total number of photons detected in each pixel throughout the acquisition period, is used to inform on tissue structure. Image series ***A-C*** corresponds to Figure 3B and its traced neuron example. Neuron silhouette traced in ***B&C*** is indicated with a white asterisk. ***B)*** Cl_o calculated from ABP-dextran lifetime. The loss of discernible structure is due to the fact that the intensity of the ABP-dextran signal has no effect on the Cl_o calculated from fluorescence lifetime (see Methods). Areas corresponding to cell body silhouettes in ***A*** have notable pixelation with relatively high apparent Cl_o values interspersed with black pixels, the latter indicating statistical exclusion due to low signal intensity (see Methods). ***C)*** In order to observe Cl_o with spatial context restored, we use intensity data to inform the brightness of the raw chloride data in ***B***, generating a composite image in which the color corresponds to Cl_o and brightness is attenuated based on the intensity of the signal in each pixel (see Methods). The most popular proprietary FLIM software packages use similar methods of composite data display. Note that prior to Cl_o analysis a further intensity-based thresholding step excludes pixels from inside somatic silhouettes in the focal plane, and pixels not corresponding to the slice (i.e., perfusate, top left) are also excluded by manually drawn ROI (masks not shown). ***D)*** Histogram of Cl_o in all analyzed pixels (blue) versus pixels excluded through thresholding step, principally marking somatic silhouettes (gray) in this representative HOSC, demonstrating no secondary peak at typical Cl_i values (i.e., =< 15mM). Further, this range of low Cl_o values is more likely excluded (gray) than included (blue) in our analysis (indicated by gray arrow). ***E&F)*** To further verify the cellular exclusion of ABP-dextran,

we looked specifically at the margins of well-defined somatic silhouettes, such as the example
marked across **A-C**, to search for signs of intracellular ABP-dextran leakage. We compared
perineuronal pixels up to 1mm outside neurons to those pixels which lie inside the somatic
margins of 72 neurons but were not excluded prior to thresholding (statistically excluded pixels
are not considered; see Methods). Consistent with out-of-focus extracellular ABP-dextran, we
found similar calculated Cl^-_o values between the two groups (**E**) with the expected wider
variance in the less intense intracellular signal (**F**; see text). *Scale bar: **A-C** 50μm.*

Figure 5 Chondroitinase treatment removes fixed perineuronal anions and attenuates Cl⁻ₒ displacement. Cl⁻ₒ composite images of representative mouse hippocampus slice cultures imaged in culture media (Cl⁻ ≈ 64mM) with 1mg/mL ABP-dextran (100mM ABP-Dextran) before (**A**) and after (**B**) treatment with bacterial MMP analog chondroitinase, or after sham digestion (**C**); shared colorbar for **A-C** at right. **D)** Histograms of [Cl⁻]ₒ corresponding to **A&B** demonstrate pre-chondroitinase digestion (dark blue) and post-chondroitinase digestion (light blue) [Cl⁻]ₒ populations of pixels are significantly different ($p < 0.001$ by Kolmogorov-Smirnov paired-sample test). **E)** When images from **A&B** are broken into regularly shaped regions of interest and analyzed as paired groups, pre-chondroitinase and post-chondroitinase are again significantly different (46.7+/-2.7mM vs 55.0+/- 5.0mM, respectively, mean+/-SD; $p = 0.00523$ by Student's paired *t*-test). **F)** Violin plot of data from **A&B** (dark and light blue, respectively) vs grouped sham-digested slice data (gray) demonstrating that Cl⁻ₒ increases significantly after chondroitin sulfate is removed by digestion, whether compared to pre-digestion of the same slice or a sham digestion (each $p < 0.001$ by Kolmogorov-Smirnov paired-sample test). **G)** Measurement of glycosaminoglycans, including chondroitin sulfate, released into the perfusate after HOSCs are treated with chondroitinase (vs sham digestion) using a commercial colorimetric assay. A significantly higher concentration of polyanionic glycosaminoglycans are found in the extracellular fluid after chondroitinase digestion ($p = 0.00632$ by Student's paired *t*-test), consistent with the release of chondroitin sulfate from extracellular matrix within slice cultures and a resultant increase in Cl⁻ₒ. **H)** HPLC analysis confirms the presence of chondroitin sulfate released into digestion buffer after chondroitinase treatment (mean = 0.20μg/mL); there was no detectable chondroitin sulfate in sham digests, negating formal significance testing. **A-C** *scale bars 50μm; all slice cultures were DIV14-21 at time of experiment; cultures were grown directly on glass with the exception of **G** where cultures were grown on membrane inserts (see Methods); significance values noted graphically for p-values <0.05 (*); <0.01 (**); <0.001 (***);*

907   *"ns" indicates "not significant". Note that data contained in **D-F** labeled 'pre' and 'post' are*
908   *derived from HOSCs in **A** and **B**, respectively.*

Figure 6 *In vivo* Cl$_o$ in mouse neocortex is consistent with *ex vivo* findings. *A&B)*
Schematic of *in vivo* imaging set-up, using a surgically implanted headbar to immobilize P26-
P30 mouse pup for imaging through 3mm coverglass window overlaying 1% agar brought up in
aCSF containing ABP-Dextran and 136mM chloride laid directly on exposed cortex. *C)* Intensity
image of mouse L2/3 cortex, demonstrating ABP-dextran exclusion from cells and vessels; large
vessels appear blurred because their widest diameter exists superficial to this field (depth
192µm). *D)* Chloride composite image corresponding to L1 cortex 138µm superficial to field *C*
(depth 54µm). *E)* Cl$_o$ in mouse neocortex is significantly lower than the CSF Cl$^-$ as shown in this
violin plot of all ROI pixels with population mean (black dashed lines) and overlaid mean+/-
standard error of ROIs analyzed. Compared to agarose brought up in 136mM aCSF overlying
the cortex (leftmost violin plot: 135.4+/-5.0mM; mean+/-standard deviation of all analyzed
pixels), *in vivo* mouse cortex had significantly lower Cl$_o$ whether measured in cellular fields such
as in *C* or in acellular fields such as in *D* (plots 2 and 3: 48.9+/-6.7mM and 55.0+/-17.9mM,
respectively). This was true when considering all ROIs in all fields from the animal from which
C&D were taken (plot 4: 47.2+/-14.9mM), and when considering all fields from all animals used
in this study (plot 5: 48.1+/-17.0mM). Statistically comparing discreet ROI means by ANOVA
analysis with Bonferroni correction (red overlay indicating mean+/-standard error of ROIs), all
groups demonstrated a statistically significant difference vs control (p < 0.001) but were not

significantly different vs each other (p = 1.0000). Note that ROIs are too numerous to display for
all groups, and we have instead overlaid the mean+/-standard error (red lines) associated with
ANOVA analysis of ROIs. Violin plot means (dashed black lines), reflecting aggregated pixels
from all ROIs, appear on the graph; associated standard deviations are listed above but are not
explicitly labeled on graph. *F)* Region of interest (ROI) selection example with ROIs shown
numbered in Cl^-_o composite image, corresponding to intensity image in ***D***, demonstrates
exclusion of large vessels but is otherwise permissive. *Scale bars **C,D&F***: 50μm; significance
values noted graphically for p-values <0.05 (*); <0.01 (**); <0.001 (***); "ns" indicates "not
significant".

.

<u>Supplemental Figure 1</u>: Evidence for pipette effects on local extracellular environment.
CA1 pyramidal cells in HOSCs can be seen to depolarize due to leakage of internal solution
from patch pipette containing 150mM KCl. Whether an area of pyramidal cells become bright
due to depolarization, as is the case in neurons expressing GcAMP *(A-C)*, or dim due to
depolarization, as is the case in neurons expressing JEDI-series voltage-sensitive system (Liu
et al., 2022) *(D-F)*, depolarization in response to high K^+ occurs in the vicinity of the pipette tip
(white dashed line). Neurons generally recover soon after a seal is achieved, precluding further
leakage and/or diffusion of internal solution from the pipette electrode. *Scale bars: 50μm*.

949   Bibliography

950   Alger BE & Nicoll RA. (1982). Pharmacological evidence for two kinds of GABA receptor on rat
951   hippocampal pyramidal cells studied in vitro. *J Physiol* **328,** 125-141.

952
953   Barker JL & Ransom BR. (1978). Amino acid pharmacology of mammalian central neurones grown in
954   tissue culture. *J Physiol* **280,** 331-354.

955
956   Beckstein O & Naughton F. (2022). General principles of secondary active transporter function. *Biophys*
957   *Rev (Melville)* **3,** 011307.

958
959   Berdichevsky Y, Saponjian Y, Park KI, Roach B, Pouliot W, Lu K, Swiercz W, Dudek FE & Staley KJ. (2016).
960   Staged anticonvulsant screening for chronic epilepsy. *Ann Clin Transl Neurol* **3,** 908-923.

961
962   Biwersi J, Farah N, Wang Y-X, Ketcham R & Verkman A. (1992a). Synthesis of cell-impermeable Cl-
963   sensitive fluorescent indicators with improved sensitivity and optical properties. *American*
964   *Journal of Physiology-Cell Physiology* **262,** C243-C250.

965
966   Biwersi J, Farah N, Wang YX, Ketcham R & Verkman AS. (1992b). Synthesis of cell-impermeable Cl-
967   sensitive fluorescent indicators with improved sensitivity and optical properties. *Am J Physiol*
968   **262,** C242-250.

969
970   Biwersi J, Tulk B & Verkman AS. (1994). Long-wavelength chloride-sensitive fluorescent indicators. *Anal*
971   *Biochem* **219,** 139-143.

972
973   Brumback AC & Staley KJ. (2008). Thermodynamic regulation of NKCC1-mediated Cl- cotransport
974   underlies plasticity of GABA(A) signaling in neonatal neurons. *J Neurosci* **28,** 1301-1312.

975
976   Celio MR & Blümcke I. (1994). Perineuronal nets--a specialized form of extracellular matrix in the adult
977   nervous system. *Brain Res Brain Res Rev* **19,** 128-145.

978
979   Celio MR, Spreafico R, De Biasi S & Vitellaro-Zuccarello L. (1998). Perineuronal nets: past and present.
980   *Trends Neurosci* **21,** 510-515.

981
982   Chahine NO, Chen FH, Hung CT & Ateshian GA. (2005). Direct measurement of osmotic pressure of
983   glycosaminoglycan solutions by membrane osmometry at room temperature. *Biophys J* **89,**
984   1543-1550.

985
986   Che A & De Marco García NV. (2021). An in vivo Calcium Imaging Approach for the Identification of Cell-
987   Type Specific Patterns in the Developing Cortex. *Front Neural Circuits* **15,** 747724.

989   Cohen I, Navarro V, Clemenceau S, Baulac M & Miles R. (2002). On the origin of interictal activity in
990   human temporal lobe epilepsy in vitro. *Science* **298,** 1418-1421.

991
992   Coombs JS, Eccles JC & Fatt P. (1955). The specific ionic conductances and the ionic movements across
993   the motoneuronal membrane that produce the inhibitory post-synaptic potential. *J Physiol* **130,**
994   326-374.

995
996   DiRaddo JO, Miller EJ, Bowman-Dalley C, Wroblewska B, Javidnia M, Grajkowska E, Wolfe BB, Liotta DC &
997   Wroblewski JT. (2015). Chloride is an Agonist of Group II and III Metabotropic Glutamate
998   Receptors. *Mol Pharmacol* **88,** 450-459.

999
1000   Doyon N, Vinay L, Prescott SA & De Koninck Y. (2016). Chloride regulation: a dynamic equilibrium crucial
1001   for synaptic inhibition. *Neuron* **89,** 1157-1172.

1002
1003   Düsterwald KM, Currin CB, Burman RJ, Akerman CJ, Kay AR & Raimondo JV. (2018). Biophysical models
1004   reveal the relative importance of transporter proteins and impermeant anions in chloride
1005   homeostasis. *Elife* **7**.

1006
1007   Ebihara S, Shirato K, Harata N & Akaike N. (1995). Gramicidin-perforated patch recording: GABA
1008   response in mammalian neurones with intact intracellular chloride. *J Physiol* **484 (Pt 1),** 77-86.

1009
1010   Egawa K, Yamada J, Furukawa T, Yanagawa Y & Fukuda A. (2013). Cl⁻ homeodynamics in gap junction-
1011   coupled astrocytic networks on activation of GABAergic synapses. *J Physiol* **591,** 3901-3917.

1012
1013   Epsztein R, Shaulsky E, Dizge N, Warsinger DM & Elimelech M. (2018). Role of Ionic Charge Density in
1014   Donnan Exclusion of Monovalent Anions by Nanofiltration. *Environ Sci Technol* **52,** 4108-4116.

1015
1016   Fatin-Rouge N, Milon A, Buffle J, Goulet RR & Tessier A. (2003). Diffusion and partitioning of solutes in
1017   agarose hydrogels: the relative influence of electrostatic and specific interactions. *The Journal of
1018   Physical Chemistry B* **107,** 12126-12137.

1019
1020   Florence TM & Farrar YJ. (1971). Spectrophotometric determination of chloride at the parts-per-billion
1021   level by the mercury(II) thiocyanate method. *Analytica chimica acta* **54,** 373--377.

1022
1023   Frantz C, Stewart KM & Weaver VM. (2010). The extracellular matrix at a glance. *J Cell Sci* **123,** 4195-
1024   4200.

1025
1026   Fuse T & Toshiya S. (1975). Preparation and properties of agar sulfates. *Agricultural and biological
1027   chemistry* **39,** 119--126.

1028

1029   Gao KW, Yu X, Darling RM, Newman J & Balsara NP. (2022). Increased Donnan exclusion in charged
1030   polymer networks at high salt concentrations. *Soft Matter* **18,** 282-292.

1031
1032   Gehlen MH. (2020). The centenary of the Stern-Volmer equation of fluorescence quenching: From the
1033   single line plot to the SV quenching map. *Journal of photochemistry and photobiology C,*
1034   *Photochemistry reviews* **42,** 100338.

1035
1036   Glykys J, Dzhala V, Egawa K, Balena T, Saponjian Y, Kuchibhotla K, Bacskai B, Kahle K, Zeuthen T & Staley
1037   K. (2014a). Local impermeant anions establish the neuronal chloride concentration. *Science* **343,**
1038   670-675.

1039
1040   Glykys J, Dzhala V, Egawa K, Balena T, Saponjian Y, Kuchibhotla KV, Bacskai BJ, Kahle KT, Zeuthen T &
1041   Staley KJ. (2014b). Response to comments on "Local impermeant anions establish the neuronal
1042   chloride concentration". *Science* **345,** 1130.

1043
1044   Glykys J, Dzhala V, Egawa K, Kahle KT, Delpire E & Staley K. (2017). Chloride dysregulation, seizures, and
1045   cerebral edema: a relationship with therapeutic potential. *Trends in neurosciences* **40,** 276-294.

1046
1047   Golgi C. (1898). Intomo alla struttura delle cellule nervose. *Boll Soc Med-Chir Pavia* **1,** 1-14.

1048
1049   González MI. (2016). Regulation of the cell surface expression of chloride transporters during
1050   epileptogenesis. *Neurosci Lett* **628,** 213-218.

1051
1052   Gregor HP. (1951). Gibbs-Donnan equilibria in ion exchange resin systems. *Journal of the American*
1053   *Chemical Society* **73,** 642-650.

1054
1055   Grimley JS, Li L, Wang W, Wen L, Beese LS, Hellinga HW & Augustine GJ. (2013a). Visualization of
1056   synaptic inhibition with an optogenetic sensor developed by cell-free protein engineering
1057   automation. *J Neurosci* **33,** 16297-16309.

1058
1059   Grimley JS, Li L, Wang W, Wen L, Beese LS, Hellinga HW & Augustine GJ. (2013b). Visualization of
1060   synaptic inhibition with an optogenetic sensor developed by cell-free protein engineering
1061   automation. *Journal of Neuroscience* **33,** 16297-16309.

1062
1063   Hampe CS, Eisengart JB, Lund TC, Orchard PJ, Swietlicka M, Wesley J & McIvor RS. (2020).
1064   Mucopolysaccharidosis Type I: A Review of the Natural History and Molecular Pathology. *Cells* **9**.

1065
1066   Helfferich F. (1995). *Ion Exchange*. Dover, New York, NY.

1067
1068   Hu H, Gan J & Jonas P. (2014). Interneurons. Fast-spiking, parvalbumin[+] GABAergic interneurons: from
1069   cellular design to microcircuit function. *Science* **345,** 1255263.

1070

1071 Huberfeld G, Wittner L, Clemenceau S, Baulac M, Kaila K, Miles R & Rivera C. (2007). Perturbed chloride
1072   homeostasis and GABAergic signaling in human temporal lobe epilepsy. *J Neurosci* **27,** 9866-
1073   9873.

1074

1075 Huguenard JR & Alger BE. (1986). Whole-cell voltage-clamp study of the fading of GABA-activated
1076   currents in acutely dissociated hippocampal neurons. *J Neurophysiol* **56,** 1-18.

1077

1078 Héron B, Mikaeloff Y, Froissart R, Caridade G, Maire I, Caillaud C, Levade T, Chabrol B, Feillet F, Ogier H,
1079   Valayannopoulos V, Michelakakis H, Zafeiriou D, Lavery L, Wraith E, Danos O, Heard JM &
1080   Tardieu M. (2011). Incidence and natural history of mucopolysaccharidosis type III in France and
1081   comparison with United Kingdom and Greece. *Am J Med Genet A* **155a,** 58-68.

1082

1083 Jedlicka P, Deller T, Gutkin BS & Backus KH. (2011). Activity-dependent intracellular chloride
1084   accumulation and diffusion controls GABA(A) receptor-mediated synaptic transmission.
1085   *Hippocampus* **21,** 885-898.

1086

1087 Jha RM, Kochanek PM & Simard JM. (2019). Pathophysiology and treatment of cerebral edema in
1088   traumatic brain injury. *Neuropharmacology* **145,** 230-246.

1089

1090 Jiang C, Agulian S & Haddad GG. (1992). Cl- and Na+ homeostasis during anoxia in rat hypoglossal
1091   neurons: intracellular and extracellular in vitro studies. *J Physiol* **448,** 697-708.

1092

1093 Kaneko H, Putzier I, Frings S, Kaupp UB & Gensch T. (2004). Chloride accumulation in mammalian
1094   olfactory sensory neurons. *J Neurosci* **24,** 7931-7938.

1095

1096 Karamanos NK, Theocharis AD, Piperigkou Z, Manou D, Passi A, Skandalis SS, Vynios DH, Orian-Rousseau
1097   V, Ricard-Blum S, Schmelzer CEH, Duca L, Durbeej M, Afratis NA, Troeberg L, Franchi M, Masola
1098   V & Onisto M. (2021). A guide to the composition and functions of the extracellular matrix. *Febs
1099   j* **288,** 6850-6912.

1100

1101 Karlócai MR, Wittner L, Tóth K, Maglóczky Z, Katarova Z, Rásonyi G, Erőss L, Czirják S, Halász P, Szabó G,
1102   Payne JA, Kaila K & Freund TF. (2016). Enhanced expression of potassium-chloride cotransporter
1103   KCC2 in human temporal lobe epilepsy. *Brain Struct Funct* **221,** 3601-3615.

1104

1105 Kroeger D, Tamburri A, Amzica F & Sík A. (2010). Activity-dependent layer-specific changes in the
1106   extracellular chloride concentration and chloride driving force in the rat hippocampus. *J
1107   Neurophysiol* **103,** 1905-1914.

1108

1109 Kuang D & Hampson DR. (2006). Ion dependence of ligand binding to metabotropic glutamate receptors.
1110   *Biochem Biophys Res Commun* **345,** 1-6.

1111

1112 Kuner T & Augustine GJ. (2000). A genetically encoded ratiometric indicator for chloride: capturing
1113   chloride transients in cultured hippocampal neurons. *Neuron* **27,** 447-459.

1114

1115 Kusama N, Harding AM & Benson CJ. (2010). Extracellular chloride modulates the desensitization
1116   kinetics of acid-sensing ion channel 1a (ASIC1a). *J Biol Chem* **285,** 17425-17431.

1117

1118 Lawrence R, Olson SK, Steele RE, Wang L, Warrior R, Cummings RD & Esko JD. (2008). Evolutionary
1119   differences in glycosaminoglycan fine structure detected by quantitative glycan reductive
1120   isotope labeling. *J Biol Chem* **283,** 33674-33684.

1121

1122 Lesperance LM, Gray ML & Burstein D. (1992). Determination of fixed charge density in cartilage using
1123   nuclear magnetic resonance. *J Orthop Res* **10,** 1-13.

1124

1125 Lewin N, Aksay E & Clancy CE. (2012). Computational modeling reveals dendritic origins of GABA(A)-
1126   mediated excitation in CA1 pyramidal neurons. *PLoS One* **7,** e47250.

1127

1128 Li M, Toombes GE, Silberberg SD & Swartz KJ. (2015). Physical basis of apparent pore dilation of ATP-
1129   activated P2X receptor channels. *Nat Neurosci* **18,** 1577-1583.

1130

1131 Liu Z, Lu X, Villette V, Gou Y, Colbert KL, Lai S, Guan S, Land MA, Lee J, Assefa T, Zollinger DR, Korympidou
1132   MM, Vlasits AL, Pang MM, Su S, Cai C, Froudarakis E, Zhou N, Patel SS, Smith CL, Ayon A,
1133   Bizouard P, Bradley J, Franke K, Clandinin TR, Giovannucci A, Tolias AS, Reimer J, Dieudonné S &
1134   St-Pierre F. (2022). Sustained deep-tissue voltage recording using a fast indicator evolved for
1135   two-photon microscopy. *Cell* **185,** 3408-3425.e3429.

1136

1137 Margolis RK & Margolis RU. (1973). The turnover of hexosamine and sialic acid in glycoproteins and
1138   mucopolysaccharides of brain. *Biochim Biophys Acta* **304,** 413-420.

1139

1140 Marinsky JA. (1985a). An interpretation of the sensitivity of weakly acidic (basic) polyelectrolyte (cross-
1141   linked and linear) equilibria to excess neutral salt. *Journal of physical chemistry (1952)* **89,** 5294--
1142   5302.

1143

1144 Marinsky JA. (1985b). An interpretation of the sensitivity of weakly acidic (basic) polyelectrolyte (cross-
1145   linked and linear) equilibria to excess neutral salt. *The Journal of Physical Chemistry* **89,** 5294-
1146   5302.

1147

1148 Morawski M, Reinert T, Meyer-Klaucke W, Wagner FE, Tröger W, Reinert A, Jäger C, Brückner G &
1149   Arendt T. (2015). Ion exchanger in the brain: quantitative analysis of perineuronally fixed anionic
1150   binding sites suggests diffusion barriers with ion sorting properties. *Scientific reports* **5,** 16471.

1151

1152 Normoyle KP, Lillis KP & Staley KJ. (2024). Synthesis and Characterization of a Novel Concentration-
1153   Independent Fluorescent Chloride Indicator, ABP-Dextran, Optimized for Extracellular Chloride
1154   Measurement. *Biomolecules* **14**.

1155
1156 Orlando C, Ster J, Gerber U, Fawcett JW & Raineteau O. (2012). Perisynaptic chondroitin sulfate
1157   proteoglycans restrict structural plasticity in an integrin-dependent manner. *J Neurosci* **32,**
1158   18009-18017, 18017a.

1159
1160 Plested AJ & Mayer ML. (2007). Structure and mechanism of kainate receptor modulation by anions.
1161   *Neuron* **53,** 829-841.

1162
1163 Rahmati N, Normoyle KP, Glykys J, Dzhala VI, Lillis KP, Kahle KT, Raiyyani R, Jacob T & Staley KJ. (2021).
1164   Unique Actions of GABA Arising from Cytoplasmic Chloride Microdomains. *J Neurosci* **41,** 4957-
1165   4975.

1166
1167 Rasmussen MK, Mestre H & Nedergaard M. (2022). Fluid transport in the brain. *Physiol Rev* **102,** 1025-
1168   1151.

1169
1170 Reuter D, Zierold K, Schröder WH & Frings S. (1998). A depolarizing chloride current contributes to
1171   chemoelectrical transduction in olfactory sensory neurons in situ. *J Neurosci* **18,** 6623-6630.

1172
1173 Roll L & Faissner A. (2014). Influence of the extracellular matrix on endogenous and transplanted stem
1174   cells after brain damage. *Front Cell Neurosci* **8,** 219.

1175
1176 Sato SS, Artoni P, Landi S, Cozzolino O, Parra R, Pracucci E, Trovato F, Szczurkowska J, Luin S & Arosio D.
1177   (2017). Simultaneous two-photon imaging of intracellular chloride concentration and pH in
1178   mouse pyramidal neurons in vivo. *Proceedings of the National Academy of Sciences* **114,** E8770-
1179   E8779.

1180
1181 Seo JH, Okuyama T, Shapiro E, Fukuhara Y & Kosuga M. (2020). Natural history of cognitive development
1182   in neuronopathic mucopolysaccharidosis type II (Hunter syndrome): Contribution of genotype to
1183   cognitive developmental course. *Mol Genet Metab Rep* **24,** 100630.

1184
1185 Sidharthan NP, Minchin RF & Butcher NJ. (2013). Cytosolic sulfotransferase 1A3 is induced by dopamine
1186   and protects neuronal cells from dopamine toxicity: role of D1 receptor-N-methyl-D-aspartate
1187   receptor coupling. *J Biol Chem* **288,** 34364-34374.

1188
1189 Sola M, Kneussel M, Heck IS, Betz H & Weissenhorn W. (2001). X-ray crystal structure of the trimeric N-
1190   terminal domain of gephyrin. *J Biol Chem* **276,** 25294-25301.

1191

1192   Song I & Dityatev A. (2018). Crosstalk between glia, extracellular matrix and neurons. *Brain Res Bull* **136,**
1193   101-108.

1194
1195   Staley K. (2023). Ionic mechanisms of ictogenic disinhibition. In *Jasper's Basic Mechanisms of the*
1196   *Epilepsies*. In Press.

1197
1198   Staley KJ & Mody I. (1992). Shunting of excitatory input to dentate gyrus granule cells by a depolarizing
1199   GABAA receptor-mediated postsynaptic conductance. *J Neurophysiol* **68,** 197-212.

1200
1201   Staley KJ & Proctor WR. (1999). Modulation of mammalian dendritic GABA(A) receptor function by the
1202   kinetics of Cl- and HCO3- transport. *J Physiol* **519 Pt 3,** 693-712.

1203
1204   Staley KJ, Soldo BL & Proctor WR. (1995). Ionic mechanisms of neuronal excitation by inhibitory GABAA
1205   receptors. *Science* **269,** 977-981.

1206
1207   Stoppini L, Buchs PA & Muller D. (1991). A simple method for organotypic cultures of nervous tissue. *J*
1208   *Neurosci Methods* **37,** 173-182.

1209
1210   Syková E & Nicholson C. (2008). Diffusion in brain extracellular space. *Physiol Rev* **88,** 1277-1340.

1211
1212   Thom M. (2014). Review: Hippocampal sclerosis in epilepsy: a neuropathology review. *Neuropathol Appl*
1213   *Neurobiol* **40,** 520-543.

1214
1215   Thompson SM, Deisz RA & Prince DA. (1988). Relative contributions of passive equilibrium and active
1216   transport to the distribution of chloride in mammalian cortical neurons. *J Neurophysiol* **60,** 105-
1217   124.

1218
1219   Titze J, Shakibaei M, Schafflhuber M, Schulze-Tanzil G, Porst M, Schwind KH, Dietsch P & Hilgers KF.
1220   (2004). Glycosaminoglycan polymerization may enable osmotically inactive Na+ storage in the
1221   skin. *Am J Physiol Heart Circ Physiol* **287,** H203-208.

1222
1223   Tora AS, Rovira X, Cao AM, Cabayé A, Olofsson L, Malhaire F, Scholler P, Baik H, Van Eeckhaut A,
1224   Smolders I, Rondard P, Margeat E, Acher F, Pin JP & Goudet C. (2018). Chloride ions stabilize the
1225   glutamate-induced active state of the metabotropic glutamate receptor 3. *Neuropharmacology*
1226   **140,** 275-286.

1227
1228   Torres PB, Nagai A, Jara CEP, Santos JP, Chow F & dos Santos DYAC. (2021). Determination of sulfate in
1229   algal polysaccharide samples: a step-by-step protocol using microplate reader. *Ocean and*
1230   *Coastal Research* **69**.

1231

1232 Tremblay R, Lee S & Rudy B. (2016). GABAergic Interneurons in the Neocortex: From Cellular Properties
1233   to Circuits. *Neuron* **91,** 260-292.

1234
1235 Tsien RY. (2013). Very long-term memories may be stored in the pattern of holes in the perineuronal
1236   net. *Proc Natl Acad Sci U S A* **110,** 12456-12461.

1237
1238 Tyzio R, Minlebaev M, Rheims S, Ivanov A, Jorquera I, Holmes GL, Zilberter Y, Ben-Ari Y & Khazipov R.
1239   (2008). Postnatal changes in somatic γ-aminobutyric acid signalling in the rat hippocampus.
1240   *European Journal of Neuroscience* **27,** 2515-2528.

1241
1242 Tønnesen J, Inavalli V & Nägerl UV. (2018). Super-Resolution Imaging of the Extracellular Space in Living
1243   Brain Tissue. *Cell* **172,** 1108-1121.e1115.

1244
1245 Urban JP, Maroudas A, Bayliss MT & Dillon J. (1979). Swelling pressures of proteoglycans at the
1246   concentrations found in cartilaginous tissues. *Biorheology* **16,** 447-464.

1247
1248 Verkman AS. (1990). Development and biological applications of chloride-sensitive fluorescent
1249   indicators. *Am J Physiol* **259,** C375-388.

1250
1251 Voipio J, Boron WF, Jones SW, Hopfer U, Payne JA & Kaila K. (2014). Comment on "Local impermeant
1252   anions establish the neuronal chloride concentration". *Science* **345,** 1130.

1253
1254 Xiao F, Nicholson C, Hrabe J & Hrabetová S. (2008). Diffusion of flexible random-coil dextran polymers
1255   measured in anisotropic brain extracellular space by integrative optical imaging. *Biophys J* **95,**
1256   1382-1392.

1257
1258 Zhang S, Kojic L, Tsang M, Grewal P, Liu J, Namjoshi D, Wellington CL, Tetzlaff W, Cynader MS & Jia W.
1259   (2016). Distinct roles for metalloproteinases during traumatic brain injury. *Neurochem Int* **96,**
1260   46-55.

1261
1262 Zhang YW, Uchendu S, Leone V, Bradshaw RT, Sangwa N, Forrest LR & Rudnick G. (2021). Chloride-
1263   dependent conformational changes in the GlyT1 glycine transporter. *Proc Natl Acad Sci U S A*
1264   **118**.

1265

1266

1267

Additional Information:

Competing Interests: None of the authors have any competing interests to declare, as indicated during the submission process for each author individually.

Funding: This work was funded by two grants to Kevin J. Staley from the National Institute for Neurological Disorders and Stroke (NINDS): R35 NS116852 and P01 NS127769.

Data availability statement: All data files used to generate the figures in the paper can be found on GitHub under KieranNormoyle/NeuronalChloride2023. Please note that the authors are still working to compile and upload data files as of article submission, but these will be ready prior to publication.

Authors' Translational Perspective: We hypothesized that if measured non-invasively, the extracellular chloride concentration (Cl^-_o) would be less than CSF chloride values ($\approx$110mM) due to Donnan exclusion by heterogeneously distributed extracellular anions so that Cl^-_o would be lower than expected and heterogenous. We found that Cl^-_o has a mean value much lower than previously appreciated ($\approx$50mM) and is indeed heterogenous. Our data further demonstrates that the loss of heterogenous sulfated glycosaminoglycans (sGAGs) in extracellular matrix results in an increased Cl^-_o as the Donnan equilibrium rebalances. Together with our previous work demonstrating heterogenous intracellular chloride (Cl^-_i), our data suggests the variance of GABA reversal potential (E_{GABA}) we and others have observed is due to the heterogeneity of the concentrations of the primary permeant anion, Cl^-, on both the intra- and extracellular side of the channel. Stable variations of perisynaptic Cl^-_o or Cl^-_i potentially allows for the tuning of E_{GABA} at individual $GABA_A$ synapses and suggests a mechanism for information storage. Pathophysiological alteration of Cl^-_o would be expected to occur when sGAG-containing extracellular matrix is damaged, such as when matrix metalloproteases (MMPs) are released following neuronal injury or death. The expected increase in extracellular chloride salts, in turn leading to equilibrative intracellular increases, represents a new mechanism of cytotoxic edema.