Title:

Human neurons lacking amyloid precursor protein exhibit cholesterol-associated developmental and presynaptic deficits

Running Title:

APP regulates Chol in neurodevelopment and synapses

Authors:

Haylee Mesa[1†], Elaine Y. Zhang[1,2†], Yingcai Wang[3], Qi Zhang[1,4]*

Affiliations:

[1] Stiles-Nicholson Brain Institute, Florida Atlantic University, Jupiter, Florida

[2] Brentwood High School, Brentwood, Tennessee

[3] Department of Biomedical Sciences, Florida Atlantic University, Boca Raton, Florida

[4] Department of Chemistry and Biochemistry, Florida Atlantic University, Boca Raton, Florida

[†] These authors contributed equally.

* Corresponding author, zhangq@fau.edu

Keywords:

Amyloid precursor protein, cholesterol, presynaptic, neuron, iPSC

Abstract:

Amyloid precursor protein (APP) produces aggregable β-amyloid peptides and its mutations are associated with familial Alzheimer's disease, which makes it one of the most studied proteins. However, APP's role in the human brain remains unclear despite years of investigation. One problem is that most studies on APP have been carried out in cell lines or model organisms, which are physiologically different from human neurons in the brain. Recently, human induced neurons (hiNs) derived from induced pluripotent stem cells provide a practical platform for studying the human brain *in vitro*. Here, we generated APP-null iPSCs using CRISPR/Cas9 genome editing technology and differentiate them to matured human neurons with functional synapses using a two-step procedure. During hiN differentiation and maturation, APP-null cells exhibited less neurite growth and reduced synaptogenesis in serum-free but not serum-containing media. We have found that cholesterol (Chol) remedies those developmental defects in APP-null cells, consistent with Chol's role in neurodevelopment and synaptogenesis. Phenotypic rescue was also achieved by co-culturing those cells with wildtype mouse astrocytes, suggesting that APP's developmental role is likely astrocytic. Next, we examined matured hiNs using patch-clamp recording and detected reduced synaptic transmission in APP-null cells. This change was largely due to decreased synaptic vesicle (SV) release and retrieval, which was confirmed by live-cell imaging using two SV-specific fluorescent reporters. Adding Chol shortly before stimulation mitigated the SV deficits in APP-null iNs, indicating that APP facilitates presynaptic membrane Chol turnover during SV exo-/endocytosis cycle. Taken together, our study in hiNs supports the notion that APP contributes to neurodevelopment, synaptogenesis and neurotransmission via maintaining brain Chol homeostasis. Given the vital of Chol in the central nervous system, the

functional connection between APP and Chol bears important implication in the pathogenesis of Alzheimer's disease.

Introduction:

Amyloid precursor protein (APP) is a type I membrane protein with single transmembrane domain which connects a long N-terminal extracellular fragment and a short C-terminal intracellular fragment (Müller et al., 2017). It has drawn tremendous attention because its proteolytic product (β-amyloid peptides, Aβs) makes amyloid plaques, a pathological hallmark for Alzheimer's disease (AD) (Selkoe and Hardy, 2016). Studies since early 1980s had elucidated how Aβs is carved from APP by β- and γ-secretases (β/γS) sequentially (de Strooper et al., 2010; Selkoe, 1998). However, the majority of APP undergoes nonamyloidogenic cleavage by α- and γ-secretases (α/γS) without producing Aβs. So far, genetic studies about the patients of rare and inheritable forms of AD with early onset (i.e., familial AD, fAD) have identified hundreds of mutations in APP and Presenilin 1/2 (i.e., catalytic subunits of γS), supporting the notion that APP proteolysis and/or its proteolytic products (namely Aβs) are likely causal factors for fAD (Selkoe and Hardy, 2016). Reportedly, most of those mutations either increase Aβ production or promotes Aβ aggregation (D'argenio and Sarnataro, 2020) and Aβ aggregates are believed to be disruptive to neuronal functionality and survival (Palop and Mucke, 2010). Together, these findings lead to the idea that soluble polymers or insoluble aggregates of Aβs causes Alzheimer's disease, so called amyloid cascade hypothesis (Selkoe and Hardy, 2016).

A powerful tool to study protein function is to remove it from its natural environment, i.e., gene knockout. In early 1990s, three strains of APP-deficient mice were generated (Li et al., 1996; Müller et al., 1994; Zheng et al., 1995). Although viable and fertile, mice completely lacking APP are 15-20% smaller and exhibit various neurological defects including reduced locomotor activity, reactive gliosis, and agenesis of corpus callosum (Li et al., 1996; Müller et al., 1994; Zheng et al., 1995), which suggests developmental and physiological roles for APP. Nevertheless, the underlying mechanism remains unsettled, especially in the aging brain (Zheng and Koo, 2011). For example, study on somatosensory cortex (layers III&V) of adult APP-null mice found an increase in dendritic spine density (Bittner et al., 2009) whereas examining cortical layers II/III and hippocampal CA1 pyramidal neurons of aging APP-null mice found a significant decrease instead (Lee et al., 2010). In addition to the variation between preparations, difference between organisms matters more, especially for complex neurological disorders like AD (Kokjohn and Roher, 2009). Recent development of induced pluripotent stem cells (iPSCs) circumvents ethical concern and leads to a variety of *in vitro* human cell preparations like 2D cultures and 3D organoids for studying aging as well as aging-related disorders (Mertens et al., 2018) like AD (Arber et al., 2017; Israel et al., 2012). One approach is to generate iPSCs from AD patients bearing disease-related genetic mutations or polymorphisms (Amponsah et al., 2021). Alternatively, one can modify AD-associated genes like APP and Presenilin in iPSCs originated from health subjects (Essayan-Perez et al., 2019). Fong *et al* showed that human astrocytes differentiated from APP-null iPSCs had reduced lipoprotein uptake, less cholesterol (Chol), and increased expression of genes for Chol synthesis and uptake (Fong et al., 2018). On the other hand, Kwart *et al* found that human neurons derived from APP-knockout iPSCs had significantly smaller endosomes and likely cellular defects in endocytosis pathways (Kwart et al., 2019). However, it remains unclear how the

deletion of APP and consequent disruption of Chol homeostasis and endocytosis can impair synaptic function and endanger neural health. Notably, AD is believed to be initiated at neuronal synapses because synaptic dysfunction proceeds and correlates to neurodegeneration and cognitive decline (Selkoe, 2002).

Different from fAD, the most common form of AD is sporadic or late-onset (sAD or LOAD) (Blennow et al., 2006). Although sAD bears similar histopathology and cognitive symptom, it has no association with APP or Presenilin polymorphism (Querfurth and LaFerla, 2010). Many Genome-wide association studies have been conducted in cohorts with thousands subjects across populations, nations, continents, and races. And dozens of genetic risk factors have been identified but not APP or Presenilins (Bellenguez et al., 2020; Tanzi, 2012). Much larger scale meta-analyses not only confirmed the relevance of those genes to sAD but also revealed the major cellular pathways they are involved (Jansen et al., 2019; Kunkle et al., 2019), which are Chol metabolism, cell membrane trafficking, and immune response (Karch and Goate, 2015). Notably, Chol is a great influencer for the latter two (Lingwood and Simons, 2010; Platt et al., 2016). While fAD and sAD seemingly have little commonality in genetics, APP has several functional ties to Chol. On one hand, APP distribution and cleavage are affected by Chol in the surface and intracellular membranes (Vetrivel and Thinakaran, 2010). On the other hand, APP and its proteolytic fragments modulate Chol uptake and synthesis (Fong et al., 2018; Kwart et al., 2019). More importantly, Barrett et al identified a Chol-interacting motif in APP (Barrett et al., 2012), providing a structural basis to connect APP and Chol together. However, how APP and Chol collaboratively contribute to AD pathogenesis remains unclear. Here, we use CRISPR/Cas9 genomic editing technology to selectively eliminate APP in human iPSCs. Using a two-step protocol, we differentiate those APP-

null iPSCs and its isogenic control to mature and synaptically connected neurons. Using morphological, optical, electrophysiological and pharmacological tools, we examine the developmental and physiological changes in human APP-null neurons. Moreover, we investigate whether and how Chol mediates those phenotypic changes caused by APP knockout.

Results

Generation of APP-null iPSCs

To obtain APP-null human neurons (hiNs) form iPSCs, we started with a wild type iPSC line (WT thereafter) that was provided by the NIH Regenerative Medicine Program as wildtype control. Within APP's exon 3, we selected a twenty-nucleotide sequence for designing sgRNA (**Figure 1a & S1**). Synthesized 20-base pair DNA sequence corresponding to this sgRNA was cloned into a CRISPR/Cas9 vector, pSpCas9(BB)-2A-Puro, which co-expresses Cas9 and Puromycin-resistant gene using a viral T2A sequence. Transfected cells were selected against Puromycin and separated to single-cell colonies. Clone 31 was selected using polymerase chain reaction (PCR) with two pairs of primers (i.e., p02/p04 for control and p05/p06 for CRISPR/Cas9 deletion (**Figure 1b & S1**)). DNA sequencing of Clone 31 revealed four-nucleotide deletion within APP's exon 3 (**Figure 1c**) which causes a reading frame shift and hence a premature termination of translation. Using Western blot, we confirmed the absence of APP in clone 31 (**Figure 1d**). To ensure that the genomic editing is limited to APP, we checked the other two members of APP gene family, amyloid precursor-like protein 1 and 2 (APLP1&2), which highly resemble APP in gene sequence and partially redundant in functionality. Western blot showed no significant difference at protein

level for both APLP1 and 2 between KO and wild-type iPSCs (**Figure 1d**). Therefore, we conclude that APP is selectively knocked out in the iPSCs clone 31 (KO thereafter).

Neurodevelopment of APP-null cells

Since APP reportedly plays a neurotrophic role during development (Müller et al., 2017), we examined the differentiation and maturation of hiNs from KO iPSCs. Based on our previous study (Gu et al., 2015), we utilized a two-step differentiation protocol, i.e., from iPSCs to neuronal precursor cells (NPCs) and then to hiNs (**Figure 2a**). First, we applied neuron induction medium (NIM) to differentiate iPSCs to NPCs. We assessed the differentiation by checking the expression of nestin, an NPC-specific marker (**Figure 2b**). The fraction of nestin-positive cells substantially increased after the application of NIM and peaked between day 4 and 8 (**Figure 2c**), suggesting the completion of the first phase. Notably, there was no significant difference in nestin-positive fractions between KO and WT cells (**Figure 2c**), indicating that APP has little effect on NPC differentiation. Next, we replated both WT and KO NPCs on Matrigel-coated coverslips and fed them with serum-free neurobasal plus medium every three days. We assessed the differentiation of hiNs by monitoring the appearance of Tuj1 (a neuron-specific marker) and glial fibrillary acidic protein (GFAP, an astrocyte-specific marker) (**Figure 2b**). Most NPCs differentiated to Tuj1- or GFAP-positive cells, whose fractions stabilized between day 20 to 30 (**Figure 2d**), suggesting a completion of differentiation. In order to achieve full synaptic connectivity and to simulate the aging brain, we maintained hiNs for at least 90 days. We examined synaptogenesis by reverse transcription quantitative polymerase chain reaction (RT-qPCR) and fluorescent immunostaining for a synapse-specific protein, Synaptophysin (i.e., Syp). RT-qPCR showed that Syp expression rose slowly and plateaued around day 50 and that there was no significant difference between WT

and KO cells (**Figure S2**). We also measured the extension and complexity of Tuj1-positive processes (likely neurites) at day 50 using Sholl analysis (Gensel et al., 2010) and found a moderate but significant decrease in APP-knockout cells (**Figure 2e&f**), in good agreement with APP's moderate contribution to neurodevelopment. Moreover, we found a significant difference between WT and KO when we measured the density of Syp (i.e., the number of Syp puncta per 20 µm Tuj1-positive processes) at day 50. It was significantly lower in KO cells (**Figure 2g**), suggesting that the lack of APP caused a reduction in synaptogenesis.

To improve neuronal differentiation and synaptogenesis in the second step, we tested serum-containing media widely used for primary neuronal culture (Liu and Tsien, 1995). Although adding serum made no difference for the resulting fractions of neurons (Tuj1+) and astrocytes (GFAP+) (**Figure 3a**), we did find that neurite growth and synaptogenesis in KO cells became similar to those of WT control at day 50 (**Figure 3b & c**). Alternatively, we experimented purified wild-type murine astrocytes as feeder cells based on our previous work with embryonic stem cells (Gu et al., 2015). We plated iPSC-derived NPCs on mouse cortical astrocytes and observed similar improvements in KO cells (**Figure 3a – c**). It is clear that factor(s) in serum or from astrocytes rescued the neurodevelopmental phenotypes of KO cells. The most likely candidate is Chol in the form of high-density lipoprotein (HDL) in serum or from astrocytes. Chol is known to be crucial for axon growth and synaptogenesis (Pfrieger, 2003a). To test that, we first applied 10 µg/mL HDL on to KO cells right after plating the iPSC-derived NPCs. We measured neurites growth and Syp puncta density at day 50. There was a significant improvement by both counts (**Figure 3d & e**). Next, we used 0.5 mM Chol-saturated methyl-β-cyclodextrin (Chol:MβCD), often used to supplement Chol to the plasma membrane (Zidovetzki and Levitan, 2007). Again, it significantly

increased neurite growth and synaptogenesis (**Figure 3d & e**). Together, our data suggest that the moderate developmental deficits in KO cells are mediated by Chol, although we cannot rule out other factors that is redundant or complimentary to Chol. Since serum-containing media has to be used with mitosis inhibitor to avoid glia overgrowth (Liu and Tsien, 1995), we used serum-free media and wildtype murine astrocytes as feeder cells for the rest of this study.

Presynaptic changes in APP-null neurons

To study the electrophysiological character of KO cells, we performed whole-cell voltage-clamp recordings after 90 days in differentiation. 17 out of 19 randomly selected cells in the KO group possessed a voltage-dependent and tetrodotoxin (TTX)-sensitive inward current (**Figure 4a**), suggesting the presence of voltage-gated Na^+-channels (VGSCs). This is comparable to the WT group (12/13 cells recorded). Since neurotransmitter release is one of the hallmarks for functional neurons, we measured the activity-evoked and Ca^{2+}-dependent release of synaptic vesicles (SVs) in both groups using FM1-43, a green styryl dye commonly used to label SVs (Betz et al., 1992). For loading, hiNs were exposed to Tyrode's solution containing 90 mM K^+ and 10 μM FM1-43 for 2 minutes, and then washed with dye-free normal Tyrode's solution (4 mM K^+) for 5 minutes. FM1-43 puncta were evident along the processes of hiNs (**Figure 4b1**). When challenged with 10-Hz, 2-min electric field stimulation, there were stimulation-induced FM1-43 fluorescence loss in both (**Figure 4b2**), which are similar to that of cultured mouse hippocampal neurons. We also recorded spontaneous excitatory postsynaptic currents (sEPSCs) in both WT and KO groups (**Figure 4c**), demonstrating that those cells were synaptically connected. Together with previous immunocytochemistry results, we conclude that both groups of cells were indeed functional neurons with functional synapses.

Notably, KO hiNs show moderate but significantly reduction as well as slow down of FM1-43 destaining, along with smaller sEPSC amplitude than the WT (**Figure 4b & c**). To understand such change, we measured the evoked EPSCs (eEPSCs) by applying five episodes of 0.5-Hz 8-sec (i.e., 4 pulses) field stimulation with a 10-second interval between consecutive episodes. We used 10-second interval because it was long enough for fast vesicle recycling but not enough for conventional clathrin-mediated endocytosis (Alabi and Tsien, 2013; Wu et al., 2014). Within the 4-pulse stimuli, we observed short-term facilitation of eEPSC amplitudes in both WT and KO groups. However, there was significant less facilitation in the KO (**Figure 4d1**). When comparing the first eEPSCs across the five episodes, we found a depression in both groups but significantly more in KO iNs (**Figure 4d2**). These results indicate that KO cells exhibit defective synaptic transmission and short-term plasticity, most likely at presynaptic terminals.

Such short-term eEPSCs facilitation (within every episode) and depression (across consecutive episodes) could result from changes in intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$), SV pool size, and SV release probability. First, we checked evoked Ca^{2+}-influx using a low-affinity and cell-permeant Ca^{2+}-indicator, Fluo-4·AM, ideal for measuring fast $[Ca^{2+}]_i$ entry (Gee et al., 2000; Grienberger and Konnerth, 2012). To identify synaptic boutons, we retrospectively loaded cells with FM4-64, a red styryl dye selectively labeling SVs (Rouze and Schwartz, 1998; Vida and Emr, 1995). In order to eliminate spontaneous neuronal firing and evoked postsynaptic Ca^{2+}-influx, we applied NBQX and D-AP5 to block AMPA and NMDA receptors respectively. We continuously imaged Fluo-4 fluorescence before, during and after a 5-s 10-Hz electric field stimulation (**Figure 5a1**). We did not detect any difference in the peak values or half rise times ($t_{1/2}$) of Fluo-4

fluorescence at synaptic boutons upon stimulation (**Figure 5a2**). After stimulation, the decay

constant (τ) of the Fluo-4 fluorescence was the same (**Figure 5a2**). Those results suggest that APP

knockout did not affect presynaptic Ca^{2+}-influx or post-stimulation $[Ca^{2+}]_i$ recovery. Can APP

deletion change presynaptic resting $[Ca^{2+}]_i$? To test that, we used Fura-2·AM, a high-affinity

ratiometric Ca^{2+} indicator suitable for measuring low $[Ca^{2+}]_i$ (Grynkiewicz et al., 1985; Miyawaki

et al., 1997). Again, FM4-64 staining was used to mark synaptic boutons. We applied ionomycin

to permeabilize hiNs and equalized their $[Ca^{2+}]_i$ using Tyrode's solutions with defined Ca^{2+}

concentrations. As such, we generated a working curve for Fura-2 ratio ($R_{340/380}$) *vs.* $[Ca^{2+}]_i$ at

synaptic boutons. We found no significant difference in the presynaptic resting $[Ca^{2+}]_i$ between

WT and KO cells (67.5 ± 4.8 nM and 70.9 ± 4.3 nM respectively) (**Figure 5b**).

To further evaluate synaptic change in KO cells, we measured miniature EPSCs (mEPSCs), whose

amplitudes and frequencies are often used to evaluate post- and presynaptic changes respectively

(Zucker and Regehr, 2002). Between WT and KO cells, the most significant difference was the

mEPSC frequency (i.e., inter-event interval, sample records shown in **Figure 5c1**). We detected

significantly longer inter-event intervals in the KO when counting mEPSC events (50 consecutive

events from every recording) or the average of average frequency for every recorded cell (**Figure

5c2** and insert respectively). As lower mEPSC frequency (i.e. longer inter-event interval) generally

indicates reduced $P_{r,v}$. This result indicates presynaptic deficiency in SV release for KO cells. In

case of mEPSC amplitudes, there was no significant difference between the two groups (**Figure

5c3** and inset showing average of averages), arguing against a significant postsynaptic change in

the KO group.

APP KO has less releasable synaptic vesicles and slower recycling

To directly examine the effect of APP KO on presynaptic terminals especially SVs, we used Synaptophysin-pHTomato (SypHTm) to label SVs. Synaptophysin is SV-specific and pHTomato is a red fluorescence protein sensitive to pH. The latter is inserted in the luminal domain of the former, so that pHTm can report deacidification and reacidification following SV exo-/endocytosis respectively (Li and Tsien, 2012). Notably, pHTm weakly fluoresces at low pH, which makes synaptic boutons visible even at rest. To mobilize all releasable SVs, we again applied 10-Hz 2-min electric stimulation; then, we sequentially applied 50mM NH_4Cl that neutralized all SVs and pH5.5 Tyrode's solution that quench cell surface SypHTm (**Figure 6a**). Expectedly, we detected (1) a rapid increase of SypHTm fluorescence representing evoked SV exocytosis, (2) a subsequent decrease representing compensatory endocytosis and reacidification, (3) maximal SypHTm fluorescence by 50mM NH_4Cl, and (4) minimal SypHTm fluorescence by pH5.5 Tyrode's solution. We normalized the fluorescence changes at individual boutons to the maximal (as 1.0) and the minimal (as 0.0) which were set by NH_4Cl and pH5.5 treatments respectively. Hence, the fractions of SypHTm at the presynaptic surface (*i*), in the releasable pool (*ii*) and in the total pool (*iii*) (**Figure 6a**) are all measurable. Among the three fractions, only (*ii*) was significantly lower (~15 % less than WT) in the KO iNs (**Figure 6b**), suggesting a reduction of the releasable pool. Although there was no significant difference in the rapid increase of pHTm fluorescence, there was a difference in the decay phase (i.e., $\tau_{WT} = 41.8 \pm 2.3$ s, $\tau_{KO} = 49.2 \pm 3.7$ s; **Figure 6c**). This result suggests a slower retrieval and/or recycling of released SVs.

Since APP directly interacts with Chol (Barrett et al., 2012), since Chol is important for exo- and endocytosis (Linetti et al., 2010; Yue and Xu, 2015), and since co-cultured wild type astrocytes

amended chronic Chol deficits, we hypothesize that the presynaptic changes in KO iNs is also related to Chol. But it is likely about acute Chol change instead of Chronic alternation. To test that, we applied MβCD or Chol:MβCD right before stimulation to acutely decrease or increase Chol in the cell membrane. Using FM1-43 imaging, we found that Chol:MβCD brought the FM1-43 destaining kinetic in KO hiNs back to the WT level whereas MβCD had little effect if not causing further reduction (**Figure 6d**). Similarly, SypHTm imaging also showed that Chol:MβCD but not MβCD amend the difference between KO and WT hiNs (**Figure 6e-g**). Together, these results suggested that the absence of APP unbalanced the membrane Chol and consequently impairs SV release and recycle. Since the plasma membrane at axon terminal is the most likely membrane section affected by the acute application of Chol:MβCD, we can further conclude that the reduction of Chol in the presynaptic surface membrane in KO cells is the major cause of reduced SV turnover.

DISCUSSION

APP is one of the most studied protein due to its proteolytic product (i.e., Aβs) and its close tie to AD. While the genetics, biochemistry, and cleavage of APP are well understood, the intrinsic functionality of APP, especially in the adult brain, is far from clear (Zheng and Koo, 2006). Studies of APP-knockout mice revealed its moderate impact on neurodevelopment and synaptic transmission, but the underlying mechanisms remain elusive (Müller and Zheng, 2012). To better understand APP's role in the human brain and its implication in AD pathogenesis, recent knockout studies utilizing human iPSCs and CRISPR/Cas9-genome editing suggest APP's relevance to Chol regulation in astrocytes (Fong et al., 2018) and endosome size in neurons (Kwart et al., 2019).

However, it remains unanswered if and how APP's relationship with Chol and endocytosis is relevant to neurodevelopment and synaptic function, especially in the context of AD and neurodegeneration in general. Using genome editing, we produced human iPSCs lacking APP expression. Using a refined two-step procedure, we differentiated APP-knockout iPSCs and the isogenic controls to NPCs and then to matured and synaptically connected hiNs. During the differentiation and maturation, we observed moderate but significant neurodevelopmental deficiency and presynaptic defects. By manipulating Chol, we have demonstrated that those chronic and acute phenotypes can be mostly mitigated by increasing Chol supply and cell membrane Chol content respectively. Based on our morphological and physiological data, and based on the fact that Chol is vital to neurons and synapses, we conclude that Chol dysregulation is at least one of the explanations for the phenotypes of APP-null cells. Furthermore, we propose that disruption of neuronal Chol homeostasis, especially at axon terminals, causes synaptic dysfunction and triggers neuronal loss. In other words, neuronal Chol is a likely converging point of various AD risk factors associated with Chol metabolism and endocytosis.

At the molecular level, APP and Chol cross with each other quite a lot. APP isoforms containing the KPI domain at its N-terminal can bind to lipoprotein receptor-related protein 1 (Knauer et al., 1996; Kounnas et al., 1995), which involves in Chol uptake. The cytoplasmic part of APP is linked to multiple LDL receptors by an adaptor protein, FE65 (Pietrzik et al., 2004; Trommsdorff et al., 1998). Additionally, APP has diverse intracellular and extracellular binding partners including SORLA (Schmidt et al., 2007) and ApoE (Haß et al., 1998), again indicating its potential to influence Chol uptake. More importantly, APP possesses a Chol interactive motif, which is within the Aβ sequence and partially overlaps with APP's transmembrane domain (Barrett et al., 2012).

This structural feature enables APP to interact with Chol embedded in the cell membranes (Beel et al., 2010; DelBove et al., 2019; Song et al., 2014, 2013). On the other hand, membrane Chol significantly influences APP cleavage, i.e., suppressing αS but facilitating β/γSs (Bodovitz and Klein, 1996; Hata et al., 2020; Kim et al., 2016; Kojro et al., 2001; Wahrle et al., 2002). This means that reducing membrane Chol will reduce the production of Aβs. However, reducing Chol is adversary to many membrane proteins, signaling pathways, and cellular functions, which is testified by the cognitive side effect of blood-brain barrier-permeable statins (Eckert et al., 2005; Shepardson et al., 2011a, 2011b). Furthermore, γ-secretase cleavage of APP is termed regulated intramembrane proteolysis (RIP), the same as the proteolytic cleavage of SREBP2 (the master Chol regulator in the brain). Although SREBP2 itself does not interact with membrane Chol, its constitutive binding partner, SCAP, serves as a Chol sensor and shuttles SREBP2 to different membrane compartments depending on membrane Chol content (Goldstein and Brown, 2015). In summary, APP has the ability to detect Chol in cell membranes and membrane Chol affects APP cleavage and distribution, a two-way interplaying which positions APP to be a membrane Chol regulator.

While possessing about a quarter of total body Chol, brain metabolizes and regulates Chol independently from other parts of the body due to the separation by blood-brain brain (Dietschy and Turley, 2004). Chol is crucial for the development, functionality and survival of neurons in the brain (Segatto et al., 2014; Zhang and Liu, 2015). Almost all brain Chol is synthesized in astrocytes and supplied to neurons in the form of HDL (Dietschy and Turley, 2004). Intriguingly, ApoE is the major brain lipoprotein and ApoE4 (the highest genetic risk factor for sAD) less intends to form HDL than E2 and E3, indicating a higher burden for maintaining neuronal Chol

homeostasis for ApoE4 carriers (Jeong et al., 2019; Leduc et al., 2010; Mahley, 2016). In neurons and almost all mammalian cells, the plasma membrane possesses the majority of cellular Chol whereas intracellular membrane has much less Chol content (Yeagle, 1985). The maintenance of such membrane Chol gradient involves active membrane trafficking and many membrane proteins (Enrich et al., 2015; Schroeder et al., 1996; Yeagle, 1985). On the other hand, disturbance of membrane Chol can affect a lot of membrane proteins (including ion channels, receptors, and transporters in neurons) and cellular signaling (including neurotransmitter secretion) (Jeremic et al., 2006; Koudinov and Koudinova, 2001; Pfrieger, 2003b; Pfrieger and Ungerer, 2011). Clearly, it is extremely important for neurons to maintain Chol gradient between the surface and intracellular membranes. This is particularly challenging at axons and axon terminals where vesicle trafficking and membrane exchanging are the most active (Bloom and Morgan, 2011). Our results agree well with the idea that APP is a local Chol sensor/regulator for membrane Chol homeostasis. Thus, APP deletion or mutation may disrupt neuronal membrane Chol and consequently synaptic function as well as neuronal survival. While guided by the popular amyloid hypothesis, previous studies on fAD mutations or sAD risk factors mostly focused on Aβ production and aggregation. With new findings linking APP to Chol, it is time to ask if and how those genetic changes alter Chol metabolism and homeostatic regulation in neurons and other brain cells. And the advances in iPSC and genome-editing technologies have paved way to tackle these questions in a more clinically relevant way.

FIGURES AND LEGENDS

Figure 1. Generation of APP-null human iPSCs. (**a**) sgDNA and targeting sequences (red) are within APP's exon 3. The nucleotide number is in reference to the beginning of exon 3. Deletion of 4 bp by CRISPR/Cas9 introduces two STOP codons, prematurely terminated APP translation. (**b**) Single clones of resulting iPSCs are assessed by PCR using two pairs of primers P02/P04 and P05/P06. Clone #31 has amplicons with 02/04 but not 05/06, indicating the absence of correct APP transcripts. (**c**) Sequencing of genomic DNA extracted from clone #31 confirmed 4-bp deletion. (**d**) Western blots of APP, APLP1 and APLP2 along with Actin from WT and KO (#31) iPSCs demonstrate the absence of APP but not APLP1 and 2.

(a)

Days: 0   6-10   80-90

Medium: Neuron Induction | Neuron Basal Plus

Status: Precursor cells | Neurons & Astrocytes

(b)

NPC (Day 6)

Nestin   DAPI

hiN (Day 80)

GFAP   Tuj1   DAPI

(c)

Nestin-Positive Fraction vs Time (day)

n.s. *n.s.* *n.s.* *n.s.*

—— WT
—— KO

(d)

Tuj1+   GFAP+   D.N.
WT
KO

Fraction (%) vs Time (day): 13, 20, 30, 50

(e)

Syp   Tuj1   DAPI

(f)

Intersections vs Diameter (μm)

***

—●— WT
—■— KO

(g)

Syp bouton per 20 μm vs Neurite Length (μm)

**

—●— WT
—■— KO

Figure 2. Differentiation of human iPSCs to hiNs and detection of neurodevelopmental defects in KO cells. (**a**) Two-step differentiation procedure. (**b**) Sample images of immunofluorescence labeling of WT cells for Nestin (red) at the end of the first step (left); and labeling for GFAP (red) and Tuj1 (green) at the end of the second step (right). Nuclear staining by DAPI (blue) is used to count all cells. Scale bar, 100 μm. (**c**) Violin plot with medians (solid lines) and quartiles (dash lines) shows a progressive appearance of Nestin-positive cells until they plateaued by day 4-8 in the first step. There are no significant differences (*n.s.*) between WT (gray) and KO (black) cells at all four dates. All n = 15 FOVs (i.e., 5 FOVs per batch and 3 batches per group) for both WT and KO at four dates. Two-tailed Student's *t*-test for comparing WT and KO at every date, all $p > 0.05$. (**d**) Bar plot for the cell fractions of Tuj1+ (green), GFAP+ (red), and double negative (D.N.) cells at four dates during the second step. There are no significant differences (*n.s.*) between WT (open) and KO (filled) groups for all three staining types at all dates. All n = 15 FOVs (i.e., 5 FOVs per batch and 3 batches per group) for both WT and KO at four dates. Two-tailed Student's *t*-test, all $p > 0.05$. (**e**) Sample image of immunofluorescence labeling of WT cells (at 50 days into the second step) for Tuj1 (green, to illustrate neurites) and Synaptophysin (Syp, red), with DAPI marking all cell nuclei. The concentric dash-line cycles with a 20-μm interval are used for Sholl analysis. The semi-transparent solid white line illustrates one of the neurites (by Tuj1 staining). Scale bar, 50 μm. (**f**) Mean and error plot of intersection counts between neurites and concentric cycles at 50 days into the second step. There is a significant difference in the total intersection counts between WT and KO cells. Both n = 18 iNs (i.e., 6 iNs per batch and 3 batches per group) for WT and KO. Two-tailed Student's *t*-test, $p < 0.001$ (***). (**g**) Mean and error plot of Syp bouton counts (per 20 μm along neurites) in the same samples as (**f**). There is a significant difference in the average counts at all lengths between WT and KO cells. Both sample sizes are the same as those in (**f**). Two-tailed Student's *t*-test, $p < 0.01$ (**).

Figure 3. Chol rescues KO's neurodevelopmental defects. (**a**) Bar plot shows that adding serum or astrocytes during the second step makes no significant difference (*n.s.*) on the fractions of differentiated cell-types. All n = 15 FOVs (i.e., 5 FOVs per batch and 3 batches per group) for the four groups. ANOVA and post-hoc Tukey-Kramer test between WT and each KO groups, all *p* > 0.05. (**b**) Mean and error plot shows that adding serum or astrocytes to KO group rescues neurite growth deficits. All n = 18 cells (i.e., 6 cells per batch and 3 batches per group) for the four groups.

Two-tailed Student's *t*-test, $p < 0.01$ for KO *vs.* WT but both $p > 0.05$ for KO + serum or KO + astrocytes *vs.* WT. (**c**) Mean and error plot shows that adding serum or astrocytes to KO group improves synaptogenesis. All sample sizes are the same as those in (**b**). Two-tailed Student's *t*-test, $p < 0.005$ for KO *vs.* WT and $p > 0.05$ for KO + serum or KO + astrocytes *vs.* WT. (**d**) Mean and error plot shows that adding HDL or Chol:MβCD to KO group improves neurite growth. All $n = 18$ cells (i.e., 6 cells per batch and 3 batches per group) for the four groups. By two-tailed Student's *t*-test, both $p < 0.005$ in comparison to KO. (**e**) Mean and error plot shows that adding HDL or Chol:MβCD to KO group increases synaptogenesis. All sample sizes are the same as those in (**d**). Two-tailed Student's *t*-test, both $p < 0.001$ in comparison to KO.

Figure 4. Mature KO hiNs exhibits reduced excitatory neurotransmission. (**a**) Sample image (left) and records (right) of whole-cell patch clamp recording in a WT hiN. Scale bar, 100 μm. Voltage-dependent inward current is detected (upper right), and it is blocked by TTX (lower right), suggesting the presence of voltage-gated sodium channels in hiNs. (**b1**) Sample image of hiNs loaded with FM1-43 whose fluorescent puncta indicates presynaptic terminals. Scale bar, 20 μm. (**b2**) Mean and error plot shows KO hiNs (iN$_{KO}$, red cycles) have slower and less FM1-43 destaining than WT hiNs (iN$_{WT}$, black triangles) or wild-type mouse hippocampal neurons

(Neuron$_{mouse}$, dotted line) during a 10-Hz, 2-min electric field stimulation (indicated by black bar). Inset is a 2D mean and error plot for average maximal FM1-43 florescence decrease ($\Delta F_{max}/F_0$) and average decay time constant (τ) between KO and WT hiNs, showing significant less florescence decrease and slower decay in KO hiNs. Both n = 180 FM1-43 puncta (i.e., 20 randomly selected FM1-43 puncta per FOV, 3 FOVs per batch, and 3 batches per group). Two-tailed Student's t-test, $p < 0.01$ (**) for $\Delta F_{max}/F_0$ and $p < 0.05$ for τ (*). (**c1**) Sample recordings of sEPSCs in WT (black) and KO (red) hiNs. (**c2**) 2D mean and error plot shows sEPSCs in KO hiNs (red cycle) exhibit decreased amplitude but similar frequency (i.e., longer inter-event interval) in comparison to WT hiNs (iN$_{WT}$, black triangle). There is a significant difference in sPESC amplitudes between WT and KO but not in inter-event interval. Data from wild-type mouse hippocampal neurons is used as reference. All n = 450 events (i.e., 30 consecutive events per cell, 5 cells per batch, and 3 batches per group). Two-tailed Student's t-test for KO $vs.$ WT, $p < 0.05$ (*) for the amplitudes and $p > 0.05$ ($n.s.$) for the inter-event interval. (**d1**) Mean and error plot shows the average of four pluses in all five stimulation episodes between KO (red cycle) and WT (black triangle) hiNs. The amplitude increases of the 2nd, 3rd, and 4th pluses are significantly less in KO. Both n = 18 iNs (i.e., 6 iNs per batch and 3 batches per group) for WT and KO. Two-tailed Student's t-test, all three $p < 0.05$ (*). (**d2**) Mean and error plot shows the average of the first pluses across all five stimulation episodes between KO and WT hiNs. The amplitude decreases of the first pluses in episodes 2, 3, 4 and 5 are significantly more in KO. Both sample sizes are the same as those in (**d1**). Two-tailed Student's t-test, $p < 0.05$ (*) for the first pulses in 2nd and 3rd episodes, and $p < 0.01$ (**) for the first pulses in 4th and 5th episodes.

Figure 5. APP knockout affects neurotransmitter release but not intracellular Ca²⁺ at presynaptic terminals. (**a1**) Mean and error plot of presynaptic (determined by FM4-64 staining) Ca²⁺ imaging, i.e., relative fluorescence change of Fluo-4 ($\Delta F_t/\Delta F_{max}$), before, during, and after 5-s 10-Hz electrical stimulation (11th - 15th seconds, black bar). In term of total fluorescence increase during the stimulation, no significant difference is detected between KO (red cycle) and WT (black triangle). Both n = 180 FM1-43 puncta (i.e., 20 randomly selected FM1-43 puncta per FOV, 3 FOVs per batch, and 3 batches per group). Two-tailed Student's *t*-test, $p > 0.05$ (*n.s.*). (**a2**) Bar plots show no significant differences in peak response (at 14th - 15th seconds), in response speed (i.e., half rising time during stimulation, $t_{1/2}$), and in recovery speed (i.e., fluorescence decay time

constant after stimulation, τ) are detected between KO (red) and WT (black). Both n = 180 FM1-43 puncta (i.e., 20 randomly selected FM1-43 puncta per FOV, 3 FOVs per batch, and 3 batches per group). Two-tailed Student's t-test, all $p > 0.05$ (*n.s.*). (**b**) Mean and error plot of ratiometric Ca^{2+} imaging using Fura-2 ($R_{340/380}$) shows intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) calibration curve (black open square) and presynaptic (marked by FM4-64 staining) $[Ca^{2+}]_i$. The presynaptic resting $[Ca^{2+}]_i$ of WT and KO are estimated to be 66.8 ± 5.1 nM and 71.1 ± 3.9 nM respectively. No significant difference is detected between KO and WT. For calibration, n = 90 hiNs (i.e., 10 randomly selected hiNs per FOV, 3 FOVs per batch, and 3 batches in total). For KO and WT, both n = 180 FM1-43 puncta (i.e., 20 randomly selected FM1-43 puncta per FOV, 3 FOVs per batch, and 3 batches per group). Two-tailed Student's t-test, $p > 0.05$ (*n.s.*). (**c1**) Sample mEPSC recordings of WT (black) and KO (red). Cumulative distribution plot of inter-event interval (i.e., inverse of frequency) (**c2**) and cumulative distributions of mEPSC amplitude (**c3**) show significant difference in mEPSC frequency but not amplitudes between WT (black) and KO (red). Both n = 750 events (i.e., 50 consecutive events per cell, 5 cells per batch, and 3 batches per group) for KO and WT. *Kolmogorov-Smirnov* test, $p < 0.01$ for inter-event interval and $p < 0.05$ for amplitude. Insets in (**c2**) and (**c3**) are box plots with all data points (i.e., average values of individual cells) of KO (red cycles) and WT (black triangles). According to the average values of individual cells, there is a significant difference in frequencies but not amplitudes between KO and WT.

Figure 6. APP knockout reduces SV turnover via disrupting Chol. (**a**) Mean and error plot shows SypHTm fluorescence intensity change during 10-Hz 2-min electrical field stimulation, which is normalized to maximal (by 50 mM NH$_4$Cl) and minimal (by pH5.5 Tyrode's solution) fluorescence. KO (red cycles) showed less increase and slower recovery than WT (black triangles). Both n = 180 SypHTm puncta (i.e., 20 randomly selected puncta per FOV, 3 FOVs per batch, and 3 batches per group) for KO and WT. (**b**) Bar plots of average values from three periods (i.e., shaded areas, i, ii, and iii, in **a**) demonstrates that KO (red) exhibits significantly less SypHTm fluorescence responses than WT (black) only during stimulation (period ii). Same sample sizes as

(**a**). Two-tailed Student's *t*-test, $p > 0.05$ (*n.s.*) for i and iii and $p < 0.01$ (******) for ii. (**c**) The left is single-exponential decay fittings for average SypHTm fluorescence changes between 61^{th} and 150^{th} second of the imaging (i.e., during the last 90 seconds of the stimulation). KO (red) is slower than WT (black). The right is bar plot of the average decay time constants (τ) shows that KO is significantly longer than WT, suggesting slower SV recycling. Same sample sizes as (**a**). Two-tailed Student's *t*-test, $p < 0.05$ (*). (**d**) Mean and error plot shows FM1-43 destaining by 10-Hz 2-min electrical field stimulation, which is normalized to pre-stimulation baseline (F_0). KO acutely treated with MβCD (open red cycles) showed little difference from untreated KO cells (semi-transparent solid red line) whereas adding Chol: MβCD (filled red cycles) improves KO cells to be like WT (semi-transparent solid black line). n = 180 FM1-43 puncta (i.e., 20 randomly selected puncta per FOV, 3 FOVs per batch, and 3 batches per group) for every group. Inset is mean and error plot of average relative FM1-43 detaining and time constant from individual FOVs (i.e., n = 9). Same symbols for MβCD and Chol: MβCD treated KO hiNs; semi-transparent red and black ovals represent the ranges (i.e., mean $\pm$ SEM) of KO and WT respectively. Two-tailed Student's *t*-test, $p < 0.05$ (*) for $\Delta Fmax/F_0$ and τ between MβCD and Chol: MβCD treated KO hiNs. (**e**) Mean and error plot shows SypHTm fluorescence changes by 10-Hz 2-min electrical field stimulation, which is normalized in the same way as (**a**). Again, acute MβCD application (open red cycles) had little effect in comparison to KO alone (semi-transparent solid red line) whereas adding Chol: MβCD (filled red cycles) makes KO cells behave like WT (semi-transparent solid black line). n = 180 FM1-43 puncta (i.e., 20 randomly selected puncta per FOV, 3 FOVs per batch, and 3 batches per group) for every group. (**f**) Bar plot of maximal SypHTm increase during the stimulation shows that Chol: MβCD but not MβCD recovers the response back to WT level. Same n as (**e**). ANOVA and post-hoc Tukey-Kramer test between WT (1) and KO with MβCD (3) or with Chol: MβCD (4), $p < 0.05$ (*) for (1) *vs.* (3) and $p > 0.05$ (*n.s.*) for (1) vs. (4). (**g**) Bar plot of SypHTm fluorescence recovery time constant shows that Chol: MβCD but not MβCD recovers the time constant back to WT level. Same n as (**e**). ANOVA and post-hoc Tukey-Kramer test between WT (1) and KO with MβCD (3) or with Chol: MβCD (4), $p < 0.05$ (*) for (1) *vs.* (3) and $p > 0.05$ (*n.s.*) for (1) vs. (4).

Author Contributions

Q.Z. designed and supervised all experiments. H.M. and E.Y.Z. carried out imaging experiments and analyze the data. H.M. made cell cultures, differentiated hiNs, collected samples, performed tests. Y.W. generated APP knockout iPSCs and validated them. Q.Z. also conducted imaging experiments, electrophysiology recording, and data analysis. Q.Z. wrote the paper with the help and inputs from all authors.

Funding Sources

This work is funded by the Florida Department of Health Ed and Ethel Moore Alzheimer's Disease Research Award (21A04) and the Stiles-Nicholson Brain Institute at Florida Atlantic University to Q.Z.

Acknowledgment

We thank R. Blakely for advice and comments and O. Pelletier for technical support. We thank all members of the Zhang Lab for their input and support.

Abbreviations

Aβ, β-amyloid peptide; AD, Alzheimer's disease; α/β/γS, α/β/γ-secretases; fAD, familial Alzheimer's disease; sAD, sporadic Alzheimer's disease; APLP, amyloid precursor-like protein; APP, amyloid precursor protein; $[Ca^{2+}]_i$, intracellular calcium concentration; Chol, cholesterol; Chol:MβCD, Chol-saturated methyl-β-cyclodextrin; D.N., double negative; eEPSCs, evoked excitatory postsynaptic currents; FOV, field of view; GFAP, glial fibrillary acidic protein; HDL, high-density lipoprotein; hiN, human induced neuron derived from induced pluripotent stem cell;

iPSC, induced pluripotent stem cell; KO, knock-out; LOAD, late-onset Alzheimer's disease; MβCD, methyl-β-cyclodextrin; mEPSCs, miniature excitatory postsynaptic currents; NPC, neuronal precursor cells; RT-qPCR, reverse transcription quantitative polymerase chain reaction; sEPSCs, spontaneous excitatory postsynaptic currents; SV, synaptic vesicle; Syp, Synaptophysin; SypHTm, Synaptophysin-pHTomato; TTX, tetrodotoxin; VGSC, voltage-gated sodium channel; WT, wild-type;

Materials and Methods

Genomic editing and clone selection

The wild-type human iPSC line ND50022 was obtained from NINDS Human Cell and Data

Repository. To disrupt APP gene, we designed sgRNA (5' – ATCCAGAACT GGUGCAAGCG

– 3') and cloned it into CRISPR/Cas9 vector, pSpCas9(BB)–2A–Puro (62988, Addgene). The

resulting plasmid was transfected into the wild-type human iPSCs using Lipofectamine

(CMAX00015, ThermoFisher Scientific). Transfected iPSC clones were selected against 5 µg/ml

Puromycin (A1113802, ThermoFisher Scientific). Puromycin-resistant iPSCs were dissociated

by Accutase (07922, STEMCELL Technologies) and plated in very low density for single

clones. After one passage, fractions of viable clones were collected and their genomic DNAs

were extracted. To identify APP–null iPSC clones, we used two pairs of PCR primers: p02 (5' –

CACTCCACCT GTACCTTACA GT – 3') & p04 (5' – GCATAGCGTA TCTACTAAGA

GTAC – 3') and p05 (5' – ATCCAGAACT GGTGCAAGCG – 3') & p06 (5' – TACTGCTCCT

ATAGGGTCAG TGCA – 3'). Their relative positions in the APP cDNA sequences are shown in

Figure S1. PCR amplification with both pairs were done with 55°C annealing and 45s extension.

Clone(s) positive for p02/p04 but negative for p05/p06 were selected for further characterization.

Animals and astrocyte culture

All animal handling and procedures were carried out in accordance with the Care and Use of

Laboratory Animals of the National Institutes of Health and approved by the Florida Atlantic

University Institutional Animal Care and Use Committee. Pregnant (E14) C57BL/6J (wild-type)

female mice were purchased from Jackson Laboratory. After arrival, those mice were acclimated

at FAU vivarium until giving birth. Postnatal cortical cultures were prepared as previously

described (Liu and Tsien, 1995) Briefly, mouse cortices were collected from postnatal day 0-2 new born pups and dissociated into a single-cell suspension with a 10-min incubation in 0.01% Trypsin-EDTA (Life Technologies) followed by gentle trituration using three glass pipettes of different diameters (~ 1, 0.5, and 0.2 mm), sequentially. Dissociated cells were recovered by centrifugation (x 200 g, 5 minutes) at 4 °C and re-suspended in media composed of Minimal Essential Medium (MEM, Life Technologies) with 10%/vol fetal bovine serum (Omega). 5 mL of cell suspension was plated in T75 culture flasks precoated with Matrigel (Corning). Cells were allowed to settle down for 2 hours before the addition of 10 mL media. After 7-10 days in vitro, cells (mostly astrocytes) reached confluency whereas neurons die out. Cells in the flasks were washed three times with ice-cold PBS and shaken repeatedly during every wash. To ensure the depletion of neurons, oligodendrocytes, and microglia, flasks were filled with cold PBS and placed in a culture shaker (180rpm) for at least 2 hours. Then, cells in the flasks were washed with ice-cold PBS for another three times with stronger tapping in between. The absence of neurons and other glial cells were confirmed by RT-qPCR. Purified astrocytes in the flasks were detached by 0.01% Trypsin-EDTA and collected as cell pellets after (x 200 g, 5 minutes) at 4 °C. After wash and spin down, purified astrocytes were resuspended in serum-free Neurobasal Plus medium (ThermoFisher Scientific) and plated on 12mm-∅ round glass coverslips (~200,000 cells/mL) precoated with Matrigel (1 coverslip per well in 24- well plates) for more than 1 hour. After two hours, we added 1mL media for every coverslip and maintained them until the plating of NPCs.

iPSCs culture and their differentiation to neurons and astrocytes

Human iPSCs with or without CRISPR/Cas9-based genome editing were grown in 35mm culture dish until ~70% confluency. After three times wash with pre-warmed PBS, cells were detached

by Accutase (ThermoFisher Scientific), and replated in new 35mm dishes or 6-well plates precoated with Matrigel (~1,000 cells/mL). For storage, iPSC pellets were resuspended in mFreSR™ Cryopreservation media (STEMCELL Technologies) and placed in -20°C for 2 hours, -80°C for overnight and finally in liquid nitrogen for long-term storage. For the 1st step of iPSC differentiate, Neural Induction Medium (STEMCELL Technologies) was applied in replated iPSCs and exchanged every other day for 6-10 days. In the second step of differentiation, cells were washed and detached as previous described. The detached cells were plated on Matrigel-coated 12mm-∅ round glass coverslips (~200,000 cells/mL) or mouse cortical astrocytes monolayers and fed with Neurobasal Plus medium every three days until reach 80-90 days.

RT-qPCR

Every step of the procedure after sample preparation, including RNA extraction, cDNA synthesis, qPCR and analysis, was performed blind. Cells were collected at the defined time points and stored in Ambion RNAlater (Thermo Fisher Scientific, Waltham, Massachusetts) at 4 °C for no more than two weeks. RNA was extracted using the Aurum Total RNA minikit (skipping the DNase step) (Bio-Rad, Hercules, California) and immediately processed or stored at 80 °C. QuantiTect Reverse Transcription Kit (Qiagen) was used for genomic DNA removal and cDNA synthesis according to the provided instructions. cDNA was stored at -20 °C. Quantitative PCR was performed using Maxima SYBR Green qPCR Master Mix (Thermo Scientific), 96-well clear plates (Thermo Scientific), and optical flat 8-cap strips (Bio-Rad) on an Opticon 2 from MJ Research with Opticon Monitor 3 software (BioRad). Reactions were set up manually using an 8-channel pipette (30-300μL) for the master mix and a single channel P2

pipette for the cDNA samples. Appropriate controls, including a reverse transcriptase-negative sample, a no template control, and a pooled cDNA sample (taken from cultured hippocampal neurons), were used on every plate to ensure consistency. Information about the primers, including sequences, sequence accession number, and annealing temperature used, can be found in **Table S1**. Optimal annealing temperature under the conditions described was experimentally determined by running a temperature gradient. The program for qPCR is as follows: (1) 95 °C for 15 minutes, (2) 95 °C for 10 seconds, (3) in designated annealing temperature (**Table S1**) for 30 seconds, (4) 72 °C for 30 seconds, (5) plate read, (6) repeat steps 2 - 5 44 times, (7) 72 °C for 10 minutes, (8) melting curve from 50 °C to 95 °C and read every 0.2 °C, (9) 72 °C for 10 minutes, (10) 4°C for storage. C(t) values were exported analyzed using Microsoft Excel and SigmaPlot. The $2^{-\Delta C(t)}$ method (Schmittgen and Livak, 2008) was used to quantify the expression of the target genes relative to the expression of three reference genes, β-actin, CypA, and Ppp1ca, which were selected based on their consistent expression in neurons and astrocytes (Bonefeld et al., 2008). The $2^{-C(t)}$ values were first normalized to the average $2^{-C(t)}$ of all samples of the gene, or intra-normalized, before further quantification.

Immunocytochemistry, Fluorescence image and Analysis

For immunostaining, coverslips were fixed in PBS containing 4% paraformaldehyde, washed, blocked for 1 h with PBS containing 1% BSA, and incubated overnight at 4 ºC with primary antibodies. Secondary antibodies with distinct fluorophores were then incubated at room temperature for 2 hours. Fluorescence imaging was performed on a Nikon Ti-E inverted microscope equipped with 20X Apo and 100X (N.A. = 0.75 and 1.40, respectively) objective and a Flash4.0 sCMOS camera (Hamamatsu). The optical filter sets used are from Chroma or Semrock.

For each fluorescence channel, images were taken with the same acquisition settings (i.e., excitation intensity and exposure time). For image analysis, Fluorescence images were thresholded in ImageJ and used as a mask with the same thresholding value applied across different image sets. The masks were subjected to minimum particle size restrictions and ROI sets were generated for every image sets. These ROI sets were applied to the original image and mean intensities were measured. Background subtraction was performed on intensity values measured from every FOV by subtraction of the average of at least 4 different ROIs manually selected from cell free areas. Data were pooled for analysis and total number of ROIs analyzed for every test are stated in the figure legends.

Electrophysiology

For every test, we used three different batches of cultures, randomly chose at least five coverslips per batch, and recorded at least one neuron per coverslip. Whole-cell voltage clamp recordings were performed using a Multi-Clamp 700B amplifier, digitized through a Digidata 1440A, and interfaced via pCLAMP 10 software (all from Molecular Devices). All recordings were performed at room temperature. Cells were voltage clamped at -70 mV for all experiments. Patch pipettes were pulled from borosilicate glass capillaries with resistances ranging from 3 - 6 MΩ when filled with pipette solution. The bath solution (Tyrode's solution) contained (in mM): 150 NaCl, 4 KCl, 2 MgCl$_2$, 2 CaCl$_2$, 10 N-2 hydroxyethyl piperazine-n-2 ethanesulphonic acid (HEPES), 10 glucose, pH 7.35. The pipette solution contained (in mM): 120 Cesium Methanesulfonate, 8 CsCl, 1 MgCl$_2$, 10 HEPES, 0.4 ethylene glycol-bis-(aminoethyl ethane)-N,N,N',N'-tetraacetic acid (EGTA), 2 MgATP, 0.3 GTP-Tris, 10 phosphocreatine, QX-314 (50 μM), pH 7.2. For the recordings of eEPSCs, a pair of platinum wires were positioned at the edge of the recording chamber and

connected to a Grass SD9 stimulator set at 70mV and 1ms duration for single pulses. For the recording of mEPSCs, bath solution was supplied with 1 μM tetrodotoxin (TTX, Abcam). The last 100 seconds of the recording were collected. For event detection of sEPSCs and mEPSCs, we used a template-based method, in which the templates were the average of thousands of sEPSC and mEPSC events randomly selected from hundreds of records collected from different batches of controls. We did not use the threshold-based event detection because the increase of sEPSC or mEPSC amplitudes might lead to more events being larger than an arbitrary threshold and thus an artificial increase of event frequency. The template was generated from our own representative data. All signals were digitized at 20 kHz, filtered at 2 kHz, and analyzed offline with Clampfit software (Molecular Devices).

Live Cell Fluorescence Imaging and Image Analysis

Live cell imaging was performed using the same Nikon Ti-E microscope setup previously described. For Ca^{2+}-imaging, cells were pre-incubated with Fura-2·AM or Fluo-4·AM (30 minutes, 1 μM, Biotium) at 37° C with 5% CO_2. After dye loading, coverslips were washed and mounted in an RC-26G imaging chamber (Warner Instruments) bottom-sealed with a 24 × 40 mm size 0 cover glass (Fisher Scientific). The chamber was fixed in a PH-1 platform (Warner Instruments) fixed on the MP-78 stage and bath solutions containing 10 μM NBQX and 20 μM D-AP5 were applied via gravity perfusion with a constant rate of ~50 μL/sec. All perfusion lines were merged into an SHM-6 in-line solution heater (Warner Instruments). The temperatures of both the imaging chamber and the perfusion solution were maintained at 34°C by a temperature controller (TC344B, Warner Instruments). For FM dye loading of the evoked pool of synaptic vesicles, mounted coverslips were incubated with 10 μM FM1-43 or FM4-64 (Life Technologies) for 2 minutes in

high K^+ bath solution containing (in mM): 64 NaCl, 90 KCl, 2 $MgCl_2$, 2 $CaCl_2$, 10 N-2 hydroxyethyl piperazine-n-2 ethanesulphonic acid (HEPES), 10 glucose, 1 µM TTX, pH 7.35. After FM loading, cells were washed with normal Tyrode's solution containing 10 µM NBQX and 20 µM D-AP5 for 5 or 10 minutes respectively. Electrical field stimulation (10 Hz, 70V) was triggered by a 5-V 2-ms TTL pulse generated by Clampex software 30 seconds after imaging began, and delivered via a pair of platinum wires attached to both sides of the imaging chamber by a Grass SD9 stimulator. Synchronization of perfusion with image acquisition was via a VC-6 valve system (Warner Instruments) and controlled in Clampex. For SypHTm imaging, the exposure time was 100 ms for all images. For Fluo-4 and FM imaging, the exposure time was 50 ms and the acquisition rate was 1 Hz. For Fura-2, two excitation filters, 340/20 and 380/20 were combined with a Prior L200 light source to excite Fura-2. For calcium concentration calibration, cells were permeabilized with ionomycin and incubated with Tyrode solutions containing designated concentration of calcium. The exposure time was 50 ms with an EM gain of 900 for all images. All images were taken with the same acquisition settings within the same measurements (laser intensity, exposure time, and EM gain).

Image analyses were performed in ImageJ. Four rectangular ROIs were drawn in cell-free regions in every FOV and their intensities averaged. For every type of fluorescence imaging including Fura-2, Fluo-4, FM1-43, and SypHTm, we pooled all background ROIs regardless of transgene differences in order to calculate the mean and standard deviation of the background intensity. Again, a masked threshold approach was applied in ImageJ, and the mean intensity plus two standard deviations was used as the common threshold for all images or image stacks. For every FOV, ROIs were generated by particle analysis based on the binary threshold mask image. FM and SypHTm, watershed segmentation and particle size limits (0.3 – 3 µm) were applied in ImageJ

to isolate ROIs representing synaptic boutons (~1 μm). For Fura-2, Fluo-4, and FM time-lapse imaging, the average intensity from 4 background ROIs was subtracted from the average intensity of each individual ROI in the same FOV. Normalization was performed using the average intensity of the first 5 frames.

Statistical analysis

All experiments were carried out in a double-blind fashion and repeated with at least three different batches of cultures (N ≥ 3). Every imaging test was repeated in at least three randomly picked coverslips per batch with randomly chosen FOVs per coverslip. No statistical methods were used to predetermine sample size. All values presented are mean ± s.e.m.. All fluorescence intensity values are background subtracted. Unpaired two-tailed Student *t*-tests or *Kolmogorov-Smirnov* test were used for two-group comparison. For more than three groups in one measure, statistics were calculated using Two-Way ANOVA with repeated measures followed by the unpaired two-tailed Student *t*-tests. *Kolmogorov-Smirnov* tests were used to compare data of cumulative distributions.

References

Alabi, A.A., Tsien, R.W., 2013. Perspectives on kiss-and-run: role in exocytosis, endocytosis, and neurotransmission. Annu Rev Physiol 75, 393–422. https://doi.org/10.1146/annurev-physiol-020911-153305

Amponsah, A.E., Guo, R., Kong, D., Feng, B., He, J., Zhang, W., Liu, X., Du, X., Ma, Z., Liu, B., Ma, J., Cui, H., 2021. Patient-derived iPSCs, a reliable in vitro model for the investigation of Alzheimer's disease. Rev Neurosci 32. https://doi.org/10.1515/revneuro-2020-0065

Arber, C., Lovejoy, C., Wray, S., 2017. Stem cell models of Alzheimer's disease: Progress and challenges. Alzheimers Res Ther. https://doi.org/10.1186/s13195-017-0268-4

Barrett, P.J., Song, Y., van Horn, W.D., Hustedt, E.J., Schafer, J.M., Hadziselimovic, A., Beel, A.J., Sanders, C.R., 2012. The amyloid precursor protein has a flexible transmembrane domain and binds cholesterol. Science (1979) 336, 1168–1171. https://doi.org/10.1126/science.1219988

Beel, A.J., Sakakura, M., Barrett, P.J., Sanders, C.R., 2010. Direct binding of cholesterol to the amyloid precursor protein: An important interaction in lipid-Alzheimer's disease relationships? Biochim Biophys Acta 1801, 975–982. https://doi.org/10.1016/j.bbalip.2010.03.008

Bellenguez, C., Grenier-Boley, B., Lambert, J.C., 2020. Genetics of Alzheimer's disease: where we are, and where we are going. Curr Opin Neurobiol 61, 40–48. https://doi.org/10.1016/j.conb.2019.11.024

Betz, W.J., Mao, F., Bewick, G.S., 1992. Activity-dependent fluorescent staining and destaining of living vertebrate motor nerve terminals. J. Neurosci. 12, 363–375.

Bittner, T., Fuhrmann, M., Burgold, S., Jung, C.K.E., Volbracht, C., Steiner, H., Mitteregger, G., Kretzschmar, H.A., Haass, C., Herms, J., 2009. γ-secretase inhibition reduces spine density in vivo via an amyloid precursor protein-dependent pathway. Journal of Neuroscience 29. https://doi.org/10.1523/JNEUROSCI.2288-09.2009

Blennow, K., de Leon, M.J., Zetterberg, H., 2006. Alzheimer's disease. Lancet 368, 387–403. https://doi.org/10.1016/S0140-6736(06)69113-7

Bloom, O.E., Morgan, J.R., 2011. Membrane trafficking events underlying axon repair, growth, and regeneration. Molecular and Cellular Neuroscience. https://doi.org/10.1016/j.mcn.2011.04.003

Bodovitz, S., Klein, W.L., 1996. Cholesterol modulates alpha-secretase cleavage of amyloid precursor protein. J Biol Chem 271, 4436–4440.

Bonefeld, B.E., Elfving, B., Wegener, G., 2008. Reference genes for normalization: A study of rat brain tissue. Synapse 62, 302–309. https://doi.org/10.1002/syn.20496

D'argenio, V., Sarnataro, D., 2020. New insights into the molecular bases of familial alzheimer's disease. J Pers Med. https://doi.org/10.3390/jpm10020026

de Strooper, B., Vassar, R., Golde, T., 2010. The secretases: enzymes with therapeutic potential in Alzheimer disease. Nat Rev Neurol 6, 99–107. https://doi.org/10.1038/nrneurol.2009.218

DelBove, C.E., Strothman, C.E., Lazarenko, R.M., Huang, H., Sanders, C.R., Zhang, Q., 2019. Reciprocal modulation between amyloid precursor protein and synaptic membrane cholesterol revealed by live cell imaging. Neurobiol Dis 127, 449–461. https://doi.org/10.1016/j.nbd.2019.03.009

Dietschy, J.M., Turley, S.D., 2004. Thematic review series: brain Lipids. Cholesterol metabolism in the central nervous system during early development and in the mature animal. J Lipid Res 45, 1375–1397. https://doi.org/10.1194/jlr.R400004-JLR200

Eckert, G.P., Wood, W.G., Muller, W.E., 2005. Statins: drugs for Alzheimer's disease? J Neural Transm (Vienna) 112, 1057–1071. https://doi.org/10.1007/s00702-004-0273-1

Enrich, C., Rentero, C., Hierro, A., Grewal, T., 2015. Role of cholesterol in SNARE-mediated trafficking on intracellular membranes. J Cell Sci 128, 1071–1081. https://doi.org/10.1242/jcs.164459

Essayan-Perez, S., Zhou, B., Nabet, A.M., Wernig, M., Huang, Y.W.A., 2019. Modeling Alzheimer's disease with human iPS cells: advancements, lessons, and applications. Neurobiol Dis. https://doi.org/10.1016/j.nbd.2019.104503

Fong, L.K., Yang, M.M., dos Santos Chaves, R., Reyna, S.M., Langness, V.F., Woodruff, G., Roberts, E.A., Young, J.E., Goldstein, L.S.B., 2018. Full-length amyloid precursor protein regulates lipoprotein metabolism and amyloid-beta clearance in human astrocytes. J Biol Chem 293, 11341–11357. https://doi.org/10.1074/jbc.RA117.000441

Gee, K.R., Brown, K.A., Chen, W.N.U., Bishop-Stewart, J., Gray, D., Johnson, I., 2000. Chemical and physiological characterization of fluo-4 Ca2+-indicator dyes. Cell Calcium 27, 97–106. https://doi.org/10.1054/ceca.1999.0095

Gensel, J.C., Schonberg, D.L., Alexander, J.K., McTigue, D.M., Popovich, P.G., 2010. SEMI-AUTOMATED SHOLL ANALYSIS FOR QUANTIFYING CHANGES IN GROWTH AND DIFFERENTIATION OF NEURONS AND GLIA. J Neurosci Methods 190, 71–79. https://doi.org/10.1016/j.jneumeth.2010.04.026

Goldstein, J.L., Brown, M.S., 2015. A century of cholesterol and coronaries: from plaques to genes to statins. Cell 161, 161–172. https://doi.org/10.1016/j.cell.2015.01.036

Grienberger, C., Konnerth, A., 2012. Imaging calcium in neurons. Neuron 73, 862–885. https://doi.org/10.1016/j.neuron.2012.02.011

Grynkiewicz, G., Poenie, M., Tsien, R.Y., 1985. A new generation of Ca2+ indicators with greatly improved fluorescence properties. J Biol Chem 260, 3440–3450.

Gu, H., Lazarenko, R.M., Koktysh, D., Iacovitti, L., Zhang, Q., 2015. A Stem Cell-Derived Platform for Studying Single Synaptic Vesicles in Dopaminergic Synapses. Stem Cells Transl Med 4, 887–893. https://doi.org/10.5966/sctm.2015-0005

Haß, S., Fresser, F., Köchl, S., Beyreuther, K., Utermann, G., Baier, G., 1998. Physical interaction of ApoE with amyloid precursor protein independent of the amyloid Aβ region in vitro. Journal of Biological Chemistry 273. https://doi.org/10.1016/s0021-9258(19)57838-3

Hata, S., Hu, A., Piao, Y., Nakaya, T., Taru, H., Morishima-Kawashima, M., Murayama, S., Nishimura, M., Suzuki, T., 2020. Enhanced amyloid-β generation by γ-secretase complex in DRM microdomains with reduced cholesterol levels. Hum Mol Genet 29, 382–393. https://doi.org/10.1093/hmg/ddz297

Israel, M.A., Yuan, S.H., Bardy, C., Reyna, S.M., Mu, Y., Herrera, C., Hefferan, M.P., van Gorp, S., Nazor, K.L., Boscolo, F.S., Carson, C.T., Laurent, L.C., Marsala, M., Gage, F.H., Remes, A.M., Koo, E.H., Goldstein, L.S.B., 2012. Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. Nature 482. https://doi.org/10.1038/nature10821

Jansen, I.E., Savage, J.E., Watanabe, K., Bryois, J., Williams, D.M., Steinberg, S., Sealock, J., Karlsson, I.K., Hägg, S., Athanasiu, L., Voyle, N., Proitsi, P., Witoelar, A., Stringer, S.,

Aarsland, D., Almdahl, I.S., Andersen, F., Bergh, S., Bettella, F., Bjornsson, S., Bræhkus, A., Bråthen, G., de Leeuw, C., Desikan, R.S., Djurovic, S., Dumitrescu, L., Fladby, T., Hohman, T.J., Jonsson, P. v, Kiddle, S.J., Rongve, A., Saltvedt, I., Sando, S.B., Selbæk, G., Shoai, M., Skene, N.G., Snaedal, J., Stordal, E., Ulstein, I.D., Wang, Y., White, L.R., Hardy, J., Hjerling-Leffler, J., Sullivan, P.F., van der Flier, W.M., Dobson, R., Davis, L.K., Stefansson, H., Stefansson, K., Pedersen, N.L., Ripke, S., Andreassen, O.A., Posthuma, D., 2019. Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. Nat Genet 51, 404–413. https://doi.org/10.1038/s41588-018-0311-9

Jeong, W., Lee, H., Cho, S., Seo, J., 2019. ApoE4-Induced Cholesterol Dysregulation and Its Brain Cell Type-Specific Implications in the Pathogenesis of Alzheimer's Disease. Mol Cells 42, 739–746. https://doi.org/10.14348/molcells.2019.0200

Jeremic, A., Jin Cho, W., Jena, B.P., 2006. Cholesterol is critical to the integrity of neuronal porosome/fusion pore. Ultramicroscopy 106, 674–677. https://doi.org/10.1016/j.ultramic.2006.01.012

Karch, C.M., Goate, A.M., 2015. Alzheimer's disease risk genes and mechanisms of disease pathogenesis. Biol Psychiatry 77, 43–51. https://doi.org/10.1016/j.biopsych.2014.05.006

Kim, Y., Kim, C., Jang, H.Y., Mook-Jung, I., 2016. Inhibition of Cholesterol Biosynthesis Reduces gamma-Secretase Activity and Amyloid-beta Generation. J Alzheimers Dis 51, 1057–1068. https://doi.org/10.3233/jad-150982

Knauer, M.F., Orlando, R.A., Glabe, C.G., 1996. Cell surface APP751 forms complexes with protease nexin 2 ligands is internalized via the low density lipoprotein receptor-related protein (LRP). Brain Res 740. https://doi.org/10.1016/S0006-8993(96)00711-1

Kojro, E., Gimpl, G., Lammich, S., Marz, W., Fahrenholz, F., 2001. Low cholesterol stimulates the nonamyloidogenic pathway by its effect on the alpha -secretase ADAM 10. Proc Natl Acad Sci U S A 98, 5815–5820. https://doi.org/10.1073/pnas.081612998

Kokjohn, T.A., Roher, A.E., 2009. Amyloid precursor protein transgenic mouse models and Alzheimer's disease: Understanding the paradigms, limitations, and contributions. Alzheimer's and Dementia 5, 340–347.

Koudinov, A.R., Koudinova, N. v, 2001. Essential role for cholesterol in synaptic plasticity and neuronal degeneration. FASEB J 15, 1858–1860.

Kounnas, M.Z., Moir, R.D., Rebeck, G.W., Bush, A.I., Argraves, W.S., Tanzi, R.E., Hyman, B.T., Strickland, D.K., 1995. LDL receptor-related protein, a multifunctional ApoE receptor, binds secreted β-amyloid precursor protein and mediates its degradation. Cell 82. https://doi.org/10.1016/0092-8674(95)90320-8

Kunkle, B.W., Grenier-Boley, B., Sims, R., Bis, J.C., Damotte, V., Naj, A.C., Boland, A., Vronskaya, M., van der Lee, S.J., Amlie-Wolf, A., Bellenguez, C., Frizatti, A., Chouraki, V., Martin, E.R., Sleegers, K., Badarinarayan, N., Jakobsdottir, J., Hamilton-Nelson, K.L., Moreno-Grau, S., Olaso, R., Raybould, R., Chen, Y., Kuzma, A.B., Hiltunen, M., Morgan, T., Ahmad, S., Vardarajan, B.N., Epelbaum, J., Hoffmann, P., Boada, M., Beecham, G.W., Garnier, J.-G., Harold, D., Fitzpatrick, A.L., Valladares, O., Moutet, M.-L., Gerrish, A., Smith, A. v, Qu, L., Bacq, D., Denning, N., Jian, X., Zhao, Y., del Zompo, M., Fox, N.C., Choi, S.-H., Mateo, I., Hughes, J.T., Adams, H.H., Malamon, J., Sanchez-Garcia, F., Patel, Y., Brody, J.A., Dombroski, B.A., Naranjo, M.C.D., Daniilidou, M., Eiriksdottir, G., Mukherjee, S., Wallon, D., Uphill, J., Aspelund, T., Cantwell, L.B., Garzia, F., Galimberti, D., Hofer, E., Butkiewicz, M., Fin, B., Scarpini, E., Sarnowski, C., Bush, W.S., Meslage, S.,

Kornhuber, J., White, C.C., Song, Y., Barber, R.C., Engelborghs, S., Sordon, S., Voijnovic, D., Adams, P.M., Vandenberghe, R., Mayhaus, M., Cupples, L.A., Albert, M.S., de Deyn, P.P., Gu, W., Himali, J.J., Beekly, D., Squassina, A., Hartmann, A.M., Orellana, A., Blacker, D., Rodriguez-Rodriguez, E., Lovestone, S., Garcia, M.E., Doody, R.S., Munoz-Fernadez, C., Sussams, R., Lin, H., Fairchild, T.J., Benito, Y.A., Holmes, C., Karamujić-Čomić, H., Frosch, M.P., Thonberg, H., Maier, W., Roschupkin, G., Ghetti, B., Giedraitis, V., Kawalia, A., Li, S., Huebinger, R.M., Kilander, L., Moebus, S., Hernández, I., Kamboh, M.I., Brundin, R., Turton, J., Yang, Q., Katz, M.J., Concari, L., Lord, J., Beiser, A.S., Keene, C.D., Helisalmi, S., Kloszewska, I., Kukull, W.A., Koivisto, A.M., Lynch, A., Tarraga, L., Larson, E.B., Haapasalo, A., Lawlor, B., Mosley, T.H., Lipton, R.B., Solfrizzi, V., Gill, M., Longstreth, W.T., Montine, T.J., Frisardi, V., Diez-Fairen, M., Rivadeneira, F., Petersen, R.C., Deramecourt, V., Alvarez, I., Salani, F., Ciaramella, A., Boerwinkle, E., Reiman, E.M., Fievet, N., Rotter, J.I., Reisch, J.S., Hanon, O., Cupidi, C., Andre Uitterlinden, A.G., Royall, D.R., Dufouil, C., Maletta, R.G., de Rojas, I., Sano, M., Brice, A., Cecchetti, R., George-Hyslop, P.S., Ritchie, K., Tsolaki, M., Tsuang, D.W., Dubois, B., Craig, D., Wu, C.-K., Soininen, H., Avramidou, D., Albin, R.L., Fratiglioni, L., Germanou, A., Apostolova, L.G., Keller, L., Koutroumani, M., Arnold, S.E., Panza, F., Gkatzima, O., Asthana, S., Hannequin, D., Whitehead, P., Atwood, C.S., Caffarra, P., Hampel, H., Quintela, I., Carracedo, Á., Lannfelt, L., Rubinsztein, D.C., Barnes, L.L., Pasquier, F., Frölich, L., Barral, S., McGuinness, B., Beach, T.G., Johnston, J.A., Becker, J.T., Passmore, P., Bigio, E.H., Schott, J.M., Bird, T.D., Warren, J.D., Boeve, B.F., Lupton, M.K., Bowen, J.D., Proitsi, P., Boxer, A., Powell, J.F., Burke, J.R., Kauwe, J.S.K., Burns, J.M., Mancuso, M., Buxbaum, J.D., Bonuccelli, U., Cairns, N.J., McQuillin, A., Cao, C., Livingston, G., Carlson, C.S., Bass, N.J., Carlsson, C.M., Hardy, J., Carney, R.M., Bras, J., Carrasquillo, M.M., Guerreiro, R., Allen, M., Chui, H.C., Fisher, E., Masullo, C., Crocco, E.A., DeCarli, C., Bisceglio, G., Dick, M., Ma, L., Duara, R., Graff-Radford, N.R., Evans, D.A., Hodges, A., Faber, K.M., Scherer, M., Fallon, K.B., Riemenschneider, M., Fardo, D.W., Heun, R., Farlow, M.R., Kölsch, H., Ferris, S., Leber, M., Foroud, T.M., Heuser, I., Galasko, D.R., Giegling, I., Gearing, M., Hüll, M., Geschwind, D.H., Gilbert, J.R., Morris, J., Green, R.C., Mayo, K., Growdon, J.H., Feulner, T., Hamilton, R.L., Harrell, L.E., Drichel, D., Honig, L.S., Cushion, T.D., Huentelman, M.J., Hollingworth, P., Hulette, C.M., Hyman, B.T., Marshall, R., Jarvik, G.P., Meggy, A., Abner, E., Menzies, G.E., Jin, L.-W., Leonenko, G., Real, L.M., Jun, G.R., Baldwin, C.T., Grozeva, D., Karydas, A., Russo, G., Kaye, J.A., Kim, R., Jessen, F., Kowall, N.W., Vellas, B., Kramer, J.H., Vardy, E., LaFerla, F.M., Jöckel, K.-H., Lah, J.J., Dichgans, M., Leverenz, J.B., Mann, D., Levey, A.I., Pickering-Brown, S., Lieberman, A.P., 2019. Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. Nat Genet 51, 414–430. https://doi.org/10.1038/s41588-019-0358-2

Kwart, D., Gregg, A., Scheckel, C., Murphy, E., Paquet, D., Duffield, M., Fak, J., Olsen, O., Darnell, R., Tessier-Lavigne, M., 2019. A Large Panel of Isogenic APP and PSEN1 Mutant Human iPSC Neurons Reveals Shared Endosomal Abnormalities Mediated by APP β-CTFs, Not Aβ. Neuron. https://doi.org/https://doi.org/10.1016/j.neuron.2019.07.010

Leduc, V., Jasmin-Bélanger, S., Poirier, J., 2010. APOE and cholesterol homeostasis in Alzheimer's disease. Trends Mol Med 16, 469–477. https://doi.org/http://dx.doi.org/10.1016/j.molmed.2010.07.008

Lee, K.J., Moussa, C.E.H., Lee, Y., Sung, Y., Howell, B.W., Turner, R.S., Pak, D.T.S., Hoe, H.S., 2010. Beta amyloid-independent role of amyloid precursor protein in generation and maintenance of dendritic spines. Neuroscience 169, 344–356. https://doi.org/10.1016/J.NEUROSCIENCE.2010.04.078

Li, Y., Tsien, R.W., 2012. pHTomato, a red, genetically encoded indicator that enables multiplex interrogation of synaptic activity. Nat Neurosci 15, 1047–1053. https://doi.org/10.1038/nn.3126

Li, Z.W., Stark, G., Gotz, J., Rulicke, T., Gschwind, M., Huber, G., Muller, U., Weissmann, C., 1996. Generation of mice with a 200-kb amyloid precursor protein gene deletion by Cre recombinase-mediated site-specific recombination in embryonic stem cells. Proc Natl Acad Sci U S A 93, 6158–6162. https://doi.org/10.1073/pnas.93.12.6158

Linetti, A., Fratangeli, A., Taverna, E., Valnegri, P., Francolini, M., Cappello, V., Matteoli, M., Passafaro, M., Rosa, P., 2010. Cholesterol reduction impairs exocytosis of synaptic vesicles. J Cell Sci 123, 595–605. https://doi.org/10.1242/jcs.060681

Lingwood, D., Simons, K., 2010. Lipid rafts as a membrane-organizing principle. Science (1979) 327, 46–50. https://doi.org/10.1126/science.1174621

Liu, G., Tsien, R.W., 1995. Properties of synaptic transmission at single hippocampal synaptic boutons. Nature 375, 404–408.

Mahley, R.W., 2016. Central Nervous System Lipoproteins: ApoE and Regulation of Cholesterol Metabolism. Arterioscler Thromb Vasc Biol 36, 1305–1315. https://doi.org/10.1161/ATVBAHA.116.307023

Mertens, J., Reid, D., Lau, S., Kim, Y., Gage, F.H., 2018. Aging in a dish: IPSC-derived and directly induced neurons for studying brain aging and age-related neurodegenerative diseases. Annu Rev Genet. https://doi.org/10.1146/annurev-genet-120417-031534

Miyawaki, A., Llopis, J., Heim, R., McCaffery, J.M., Adams, J.A., Ikura, M., Tsien, R.Y., 1997. Fluorescent indicators for Ca2+ based on green fluorescent proteins and calmodulin. Nature 388, 882–887. https://doi.org/10.1038/42264

Müller, U., Cristina, N., Li, Z.W., Wolfer, D.P., Lipp, H.P., Rülicke, T., Brandner, S., Aguzzi, A., Weissmann, C., 1994. Behavioral and anatomical deficits in mice homozygous for a modified β-amyloid precursor protein gene. Cell 79. https://doi.org/10.1016/0092-8674(94)90066-3

Müller, U.C., Deller, T., Korte, M., 2017. Not just amyloid: Physiological functions of the amyloid precursor protein family. Nat Rev Neurosci. https://doi.org/10.1038/nrn.2017.29

Müller, U.C., Zheng, H., 2012. Physiological functions of APP family proteins. Cold Spring Harb Perspect Med 2, 1–18. https://doi.org/10.1101/cshperspect.a006288

Palop, J.J., Mucke, L., 2010. Amyloid-beta-induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. Nat Neurosci 13, 812–818. https://doi.org/nn.2583 [pii] 10.1038/nn.2583

Pfrieger, F.W., 2003a. Role of cholesterol in synapse formation and function. Biochim Biophys Acta 1610, 271–280.

Pfrieger, F.W., 2003b. Cholesterol homeostasis and function in neurons of the central nervous system. Cell Mol Life Sci 60, 1158–1171. https://doi.org/10.1007/s00018-003-3018-7

Pfrieger, F.W., Ungerer, N., 2011. Cholesterol metabolism in neurons and astrocytes. Prog Lipid Res 50, 357–371. https://doi.org/10.1016/j.plipres.2011.06.002

Pietrzik, C.U., Yoon, I.S., Jaeger, S., Busse, T., Weggen, S., Koo, E.H., 2004. FE65 Constitutes the Functional Link between the Low-Density Lipoprotein Receptor-Related Protein and the

Amyloid Precursor Protein. Journal of Neuroscience 24. https://doi.org/10.1523/JNEUROSCI.5451-03.2004

Platt, N., Speak, A.O., Colaco, A., Gray, J., Smith, D.A., Williams, I.M., Wallom, K.L., Platt, F.M., 2016. Immune dysfunction in Niemann-Pick disease type C. J Neurochem 136 Suppl, 74–80. https://doi.org/10.1111/jnc.13138

Querfurth, H.W., LaFerla, F.M., 2010. Alzheimer's disease. N Engl J Med 362, 329–344. https://doi.org/10.1056/NEJMra0909142

Rouze, N.C., Schwartz, E.A., 1998. Continuous and Transient Vesicle Cycling at a Ribbon Synapse. The Journal of Neuroscience 18, 8614.

Schmidt, V., Sporbert, A., Rohe, M., Reimer, T., Rehm, A., Andersen, O.M., Willnow, T.E., 2007. SorLA/LR11 regulates processing of amyloid precursor protein via interaction with adaptors GGA and PACS-1. Journal of Biological Chemistry 282. https://doi.org/10.1074/jbc.M705073200

Schmittgen, T.D., Livak, K.J., 2008. Analyzing real-time PCR data by the comparative CT method. Nat. Protocols 3, 1101–1108.

Schroeder, F., Frolov, A.A., Murphy, E.J., Atshaves, B.P., Jefferson, J.R., Pu, L., Wood, W.G., Foxworth, W.B., Kier, A.B., 1996. Recent advances in membrane cholesterol domain dynamics and intracellular cholesterol trafficking. Proc Soc Exp Biol Med 213, 150–177.

Segatto, M., Leboffe, L., Trapani, L., Pallottini, V., 2014. Cholesterol homeostasis failure in the brain: implications for synaptic dysfunction and cognitive decline. Curr Med Chem 21, 2788–2802.

Selkoe, D.J., 2002. Alzheimer's disease is a synaptic failure. Science (1979) 298, 789–791. https://doi.org/10.1126/science.1074069

Selkoe, D.J., 1998. The cell biology of beta-amyloid precursor protein and presenilin in Alzheimer's disease. Trends Cell Biol 8, 447–453.

Selkoe, D.J., Hardy, J., 2016. The amyloid hypothesis of Alzheimer's disease at 25 years. EMBO Mol Med 8, 595–608. https://doi.org/10.15252/emmm.201606210

Shepardson, N.E., Shankar, G.M., Selkoe, D.J., 2011a. Cholesterol level and statin use in Alzheimer disease: II. Review of human trials and recommendations. Arch Neurol 68, 1385–1392. https://doi.org/10.1001/archneurol.2011.242

Shepardson, N.E., Shankar, G.M., Selkoe, D.J., 2011b. Cholesterol level and statin use in Alzheimer disease: I. Review of epidemiological and preclinical studies. Arch Neurol 68, 1239–1244. https://doi.org/10.1001/archneurol.2011.203

Song, Y., Hustedt, E.J., Brandon, S., Sanders, C.R., 2013. Competition between homodimerization and cholesterol binding to the C99 domain of the amyloid precursor protein. Biochemistry 52, 5051–5064. https://doi.org/10.1021/bi400735x

Song, Y., Mittendorf, K.F., Lu, Z., Sanders, C.R., 2014. Impact of bilayer lipid composition on the structure and topology of the transmembrane amyloid precursor C99 protein. J Am Chem Soc 136, 4093–4096. https://doi.org/10.1021/ja4114374

Tanzi, R.E., 2012. The Genetics of Alzheimer Disease. Cold Spring Harb Perspect Med 2. https://doi.org/10.1101/cshperspect.a006296

Trommsdorff, M., Borg, J.P., Margolis, B., Herz, J., 1998. Interaction of cytosolic adaptor proteins with neuronal apolipoprotein E receptors and the amyloid precursor protein. Journal of Biological Chemistry 273. https://doi.org/10.1074/jbc.273.50.33556

Vetrivel, K.S., Thinakaran, G., 2010. Membrane rafts in Alzheimer's disease beta-amyloid production. Biochim Biophys Acta Mol Cell Biol Lipids. https://doi.org/10.1016/j.bbalip.2010.03.007

Vida, T.A., Emr, S.D., 1995. A new vital stain for visualizing vacuolar membrane dynamics and endocytosis in yeast. J Cell Biol 128, 779–792. https://doi.org/10.1083/jcb.128.5.779

Wahrle, S., Das, P., Nyborg, A.C., McLendon, C., Shoji, M., Kawarabayashi, T., Younkin, L.H., Younkin, S.G., Golde, T.E., 2002. Cholesterol-dependent gamma-secretase activity in buoyant cholesterol-rich membrane microdomains. Neurobiol Dis 9, 11–23. https://doi.org/10.1006/nbdi.2001.0470

Wu, L.G., Hamid, E., Shin, W., Chiang, H.C., 2014. Exocytosis and endocytosis: modes, functions, and coupling mechanisms. Annu Rev Physiol 76, 301–331. https://doi.org/10.1146/annurev-physiol-021113-170305

Yeagle, P.L., 1985. Cholesterol and the cell membrane. Biochimica et Biophysica Acta (BBA) - Reviews on Biomembranes 822, 267–287. https://doi.org/http://dx.doi.org/10.1016/0304-4157(85)90011-5

Yue, H.Y., Xu, J., 2015. Cholesterol regulates multiple forms of vesicle endocytosis at a mammalian central synapse. J Neurochem 134, 247–260. https://doi.org/10.1111/jnc.13129

Zhang, J., Liu, Q., 2015. Cholesterol metabolism and homeostasis in the brain. Protein Cell 6, 254–264. https://doi.org/10.1007/s13238-014-0131-3

Zheng, H., Jiang, M., Trumbauer, M.E., Sirinathsinghji, D.J., Hopkins, R., Smith, D.W., Heavens, R.P., Dawson, G.R., Boyce, S., Conner, M.W., Stevens, K.A., Slunt, H.H., Sisoda, S.S., Chen, H.Y., van der Ploeg, L.H., 1995. beta-Amyloid precursor protein-deficient mice show reactive gliosis and decreased locomotor activity. Cell 81, 525–531.

Zheng, H., Koo, E.H., 2011. Biology and pathophysiology of the amyloid precursor protein. Mol Neurodegener 6, 27. https://doi.org/10.1186/1750-1326-6-27

Zheng, H., Koo, E.H., 2006. The amyloid precursor protein: beyond amyloid. Mol Neurodegener 1, 5. https://doi.org/10.1186/1750-1326-1-5

Zidovetzki, R., Levitan, I., 2007. Use of cyclodextrins to manipulate plasma membrane cholesterol content: evidence, misconceptions and control strategies. Biochim Biophys Acta 1768, 1311–1324. https://doi.org/10.1016/j.bbamem.2007.03.026

Zucker, R.S., Regehr, W.G., 2002. Short-term synaptic plasticity. Annu Rev Physiol 64, 355–405.