Exact distribution of the quantal content in synaptic transmission

Krishna Rijal,[1] Nicolas I.C. Müller,[2] Eckhard Friauf,[2] Abhyudai Singh,[3] Ashok Prasad,[4, *] and Dibyendu Das[1, †]

[1]*Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India*
[2]*Animal Physiology Group, Department of Biology,*
University of Kaiserslautern, Kaiserslautern, Germany
[3]*Departments of Electrical and Computer Engineering,*
Biomedical Engineering and Mathematical Sciences,
University of Delaware, Newark, DE 19716, USA
[4]*Department of Chemical and Biological Engineering,*
Colorado State University, Fort Collins, Colorado, USA
(Dated: July 12, 2023)

During electro-chemical signal transmission through synapses, triggered by an action potential (AP), a stochastic number of synaptic vesicles (SV), called the *quantal content*, release neurotransmitters in the synaptic cleft. It is widely accepted that the quantal content probability distribution is a binomial based on the number of ready-release SVs in the pre-synaptic terminal. But the latter number itself fluctuates due to its stochastic replenishment, hence the actual distribution of quantal content is unknown. We show that exact distribution of quantal content can be derived for general stochastic AP inputs in the steady-state. For fixed interval AP train, we prove that the distribution is a binomial, and corroborate our predictions by comparison with electrophysiological recordings from MNTB-LSO synapses of juvenile mice. For a Poisson train, we show that the distribution is non-binomial. Moreover, we find exact moments of the quantal content in the Poisson and other general cases, which may be used to obtain the model parameters from experiments.

The human brain is arguably the most complex natural object in the known universe. Its complexity, as well as its capabilities, are a product of its roughly 86 billion [1] neurons that make approximately 10^{15} connections [2] with each other. The majority of these connections are chemical synapses, a schematic diagram of which is shown in Fig 1a, with the vesicle-rich pre-synaptic terminal on the left and the post-synaptic terminal with neurotransmitter receptors on the right. In the pre-synaptic terminal, ready-release synaptic vesicles (SVs) packed with neurotransmitters attach to docking sites. The arrival of an AP controls the calcium dynamics which triggers the stochastic fusion of a few SVs (the *quantal content*) with the pre-synaptic membrane, thereby releasing their neurotransmitter cargo in the synaptic cleft as shown. A stochastic train of APs is shown in Fig 1b (top) and the corresponding time dependent quantal content is shown schematically (below). Experimentally the quantal content is obtained as the ratio of the measured peak amplitude of evoked (excitatory or inhibitory) postsynaptic currents [3] to average quantal size. Thus, SV release from pre-synaptic terminals is a fundamental feature of all information processing in the nervous system.

Addressing how neurons communicated with another, Castillo and Katz [4] first proposed what they described as a quantal theory of release of certain binomially distributed 'units'. Over the years micro-structural and electrophysiological studies revealed the relevant biological details, and a picture emerged of spatially distributed pools of neurotransmitter containing SVs in the presynaptic neuron. These are the resting, recycling

FIG. 1: (a) Schematic picture of a chemical synapse. The SVs (circular balls) filled with neurotransmitters (black dots) get docked to sites (in green) and remain ready to release at the presynaptic terminal. Each AP triggers release of some SVs to the synaptic cleft, which reach the receptors in post-synaptic terminal to produce evoked post-synaptic current. (b) A schematic input AP train with random intervals t_s (top), and (bottom) the output quantal content varying with time, shown to reach a steady-state at long times.

and ready-release (RRP) pools. SVs from the RRP may release their neurotransmitter cargo on stimulation by an AP through membrane fusion at the pre-synaptic terminal [5–8]. Sustained APs require RRP replenishment, and early theoretical analysis assumed deterministic replenishment, and quantal release [13–15]. Later works developed probabilistic descriptions and stochastic models [16–22]. Empirical observations had shown that the binomial distribution was a good fit to the statistics of quantal release, and this has been the basis of the majority of fluctuation analysis [12, 16, 20, 22–

27]. The basic model thus assumes that at the instant of AP stimulation, if there are n_- docked SVs in RRP and p_{rt} is the instantaneous release probability, then quantal content b (fused and released SVs) is distributed as $B(n_-, p_{rt}, b) = \binom{n_-}{b} p_{rt}^b (1 - p_{rt})^{n_- - b}$ [12, 20–22, 28–30].

As n_- is itself stochastic, it is generally expected that the actual distribution of quantal content is not a binomial, as would result from summing over the likelihood of random values of n_-. This fact has largely been overlooked in the literature and some works even used binomial with n_- approximated by its mean value [20–22, 28, 29, 31, 32]. Systematic formalism treating the docked vesicle number stochastically appeared in a set of works [33–36] – although the focus of the authors was the estimation of the model parameters. While exact relationships between probability of postsynaptic currents after two successive APs were formally written down, the problem could not be solved analytically further because of the presence of multiple nested summations due to the explicit history of the AP train [34].

In this work, we show that in the *steady state* limit after sustained AP stimulation, for uncorrelated stochastic inter-spike intervals (ISI), the full distribution of quantal content b as well as its moments becomes *analytically tractable*. The commonly used fixed ISI stimulation follows as a special case of our general result. Although the transient plastic regime (Fig. 1b) is often studied in synapses, the steady state is also of interest [7, 9, 12, 20] and is attained in experimentally reasonable times [37, 38]. Poisson AP stimulation has been experimentally studied [39], and such ISI distributions have been observed in visual cortical neurons [40, 41]. Thus stochastic stimulations are important *in vivo* scenarios. These facts provide motivation for our focus here on steady state quantal content under stochastic stimulations.

FIG. 2: The time variation of docked SV number $n(t)$ is shown as a function of time over two successive ISIs between the $m - 2$ & $m - 1$ & m^{th} APs. The respective numbers before an AP (n_-) after an AP (n_+) and quantal content (b) are shown.

In Fig 2, the time dependent number $n(t)$ of docked SVs in the pre-synaptic neuron is shown schematically. Starting with $n = n_{+,m-1}$ after the $(m-1)^{th}$ AP, the number grows stochastically to $n = n_{-,m}$ up to a time t_{sm}, at which the appearance of the m^{th} AP leads to the

fusion and release of b_m vesicles, resulting in a sudden decrease of SVs from $n_{-,m} = n_{+,m} + b_m$ to $n_{+,m}$.

We assume that docked vesicles may be released probabilistically after an AP but do not otherwise detach. Following the empirical literature [12, 20, 21, 28, 30], as mentioned above the distribution of b_m is $B(n_{-,m}, p_{rt}, b_m) = \binom{n_{-,m}}{b_m} p_{rt}^{b_m} (1 - p_{rt})^{n_{-,m} - b_m}$, where the release probability p_{rt} is time dependent but saturates at long time. However, $n_{-,m}$ itself is stochastic and is replenished as follows. We assume M is the maximum number of available docking sites, and that docking sites are refilled from the vesicle pool at a rate proportional to the number of empty sites, i.e., $k_t(M - n)$ [20, 22], where k_t is the time dependent docking rate per empty site. The stochastic replenishment of docked SVs from an initial $n_{+,m-1}$ to some n in time t is described by the conditional distribution $p(n, t | n_{+,m-1})$.

Exact recursive relations may be written between the probabilities $P_+^m(n_{+,m}, t_{sm})$ and $P_+^{m-1}(n_{+,m-1}, t_{sm-1})$ for the SV numbers $n_{+,m}$ and $n_{+,m-1}$ after m^{th} and $(m-1)^{th}$ AP respectively – similar to ideas in [33, 34]:

$$P_+^m(n_{+,m}, t_{sm}) = \sum_{n_{+,m-1}, b_m} B(n_{+,m} + b_m, p_{rt}, b_m) \times$$
$$p(n_{+,m} + b_m, t_{sm} | n_{+,m-1}) P_+^{m-1}(n_{+,m-1}, t_{sm-1}) \tag{1}$$

Similarly, the probability of SV number $n_{-,m}$ just prior to the m^{th} AP namely $P_-^m(n_{-,m}, t_{sm})$ is given by:

$$P_-^m(n_{-,m}, t_{sm}) = \sum_{n_{+,m-1}} p(n_{-,m}, t_{sm} | n_{+,m-1}) \times$$
$$P_+^{m-1}(n_{+,m-1}, t_{sm-1}) \tag{2}$$

The nested nature of the Eqs. (1) and (2) depending on the full history of successive ISIs make problems of these type analytically very challenging and various computational methods have been developed in this context [33, 34, 36]. In this work we focus on the long time limit after sustained AP stimulation, when interestingly we find that the problem becomes analytically tractable.

Whether the input stimulation is a deterministic AP train [38] or a stochastic one [39], on sustained stimulation (i.e. the index m of AP and time is large), a *steady state* [7, 12, 20, 37, 38] is expected to be reached. In this limit, the mean quantal content $\langle b \rangle$ becomes a constant, and its distribution $Q^{ss}(b)$ becomes time independent. Analytically obtaining $Q^{ss}(b)$ is the main aim of this work. Note that in this $t \to \infty$ limit, the time-varying calcium concentration-dependent replenishment rate k_t and release probability p_{rt} saturates to constant values $k = k_\infty$ and $p_r = p_{r\infty}$ (see representative curves in Section 8 of supplementary information (SI) following a model of Ref. [15]). As we solely focus on the steady state, the only two neuronal model parameters we assume in our calculations are k and p_r. We will show that these can be estimated by comparing experimen-

tal data with exact moments of quantal content that we derive.

We study stochastic AP stimulations but assume that the ISIs are uncorrelated, i.e. the joint distribution of two intervals t_{sm} and t_{sn} namely $g_2(t_{sm}, t_{sn}) = g(t_{sm})g(t_{sn})$. Under this assumption, for any general stochastic ISI distribution $g(t_s)$, in steady state, it may be shown (Section 2 of SI) that the Eqs. (1) and (2) become

$$P_+^{ss}(n_+) = \sum_{n'_+,b} \int_0^\infty dt_s g(t_s) B(n_+ + b, p_r, b) \times$$
$$p(n_+ + b, t_s|n'_+)P_+^{ss}(n'_+) \quad (3)$$

$$P_-^{ss}(n_-) = \sum_{n'_+} \int_0^\infty p\left(n_-, t_s|n'_+\right) P_+^{ss}(n'_+)g(t_s)dt_s \quad (4)$$

At long times, m is large. Therefore, we have used $n_{+,m-1} = n'_+$, $n_{+,m} = n_+$, $n_{-,m} = n_-$ and $b_m = b$, as they become history independent random variables. For the case of common experimental interest of fixed ISIs we need to set $g(t_s) = \delta(t_s - T)$. The conditional distribution $p(n, t|n'_+)$ for n'_+ to become n in time $t = t_s$ used in Eqs. (3) and (4), satisfy the Master equation [45]:

$$\frac{\partial p(n, t|n'_+)}{\partial t} = k\left[M - (n-1)\right]\, p(n-1, t|n'_+)$$
$$-k[M - n]\, p(n, t|n'_+). \quad (5)$$

This equation can be solved using the generating function $F(q, t|n'_+) = \sum_n q^n p(n, t|n'_+)$ (see SI Section 1 for details), and the solution is:

$$p(n, t|n'_+) = \binom{M - n'_+}{n - n'_+}\left(1 - e^{-kt}\right)^{n - n'_+} e^{-[k(M-n)]t}. \quad (6)$$

The true steady state distribution of quantal content $Q^{ss}(b)$, is obtained by summing its Binomial distribution at every release step over all possible docked random SV numbers n_-:

$$Q^{ss}(b) = \sum_{n_-=0}^{M} B(n_-, p_r, b)P_-^{ss}(n_-). \quad (7)$$

Thus to calculate $Q^{ss}(b)$, we need the steady state distribution $P_-^{ss}(n_-)$, and in turn $P_+^{ss}(n_+)$ because of the relationships in Eqs. (3) and (4). The Eq. (3) has $P_+^{ss}(n_+)$ on both sides. Through a lengthy calculation (see details in SI Section 3), we show that the corresponding generation function, namely $F_+(q) = \sum_{n_+=0}^\infty q^{n_+} P_+^{ss}(n_+)$ can be obtained using the knowledge of $F(q, t|n'_+)$, the generating function for $p(n, t|n'_+)$ obtained above (Eq. (6)). This generating function leads us to two other relevant generating functions $F_-(q) = \sum_{n_-=0}^\infty q^{n_-} P_-^{ss}(n_-)$ and

$F_b(q) = \sum_{b=0}^\infty q^b Q^{ss}(b)$ corresponding to the distributions $P_-^{ss}(n_-)$ and $Q^{ss}(b)$. Since the three stochastic variables, $n_-, n_+,$ and b are related to each other through the relationship $n_+ = n_- - b$, the different generating functions are also related (see details in SI Section 4):

$$F_-(q) = F_+\left(\frac{q - p_r}{1 - p_r}\right) \,\&\, F_b(q) = F_+\left(\frac{p_r(q-1)}{1 - p_r} + 1\right). \quad (8)$$

Once we know $F_b(q)$, we may obtain the desired distribution $Q^{ss}(b)$ (see SI Section 4) using the standard relationship $Q^{ss}(b) = \frac{1}{b!}\frac{\partial^b F_b(q)}{\partial q^b}\Big|_{q=0}$ [45]:

$$Q^{ss}(b) = \sum_{n=0}^\infty \binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1 - (1 - p_r)^n L_n}\left[\phi_{n,0} \right.$$
$$\left. + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i}\phi_{n,m_z}\phi_{m_z,m_{z-1}}\cdots\phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})}\right], \quad (9)$$

where, $\phi_{n,m} = \binom{n}{m}\int_0^\infty dt_s g(t_s)e^{-nkt_s}(e^{kt_s} - 1)^{n-m}$ and $L_n = \int_0^\infty dt_s g(t_s)e^{-nkt_s}$. In Eq. (9), $\{S_{(n-1)}\}$ denotes the set of all the subsets $S_{(n-1)} = \{m_i\} = (m_z, m_{z-1}, \cdots, m_1)$ with $m_i \in (1, 2, \cdots, n-1)$ such that $m_z > m_{z-1} > \cdots > m_1$. Although the upper limit of the sum is shown up to ∞, due to the combinatorial factor $\binom{M}{n}$, n never exceeds M. Note that for an arbitrary $g(t_s)$ there is no reason to expect that $Q^{ss}(b)$ in Eq. (9) would reduce to a binomial distribution.

The closed form expressions of the mean and $CV^2 = \frac{\langle b^2\rangle}{\langle b\rangle^2} - 1$ of the distribution in Eq. (9) (see Section 4 in SI) are:

$$\langle b\rangle = \frac{M p_r \phi_{1,0}}{1 - (1 - p_r)L_1}\,, \text{ and}$$

$$CV^2 = \frac{\phi_{1,0} + p_r L_1}{M\phi_{1,0}}\left[\frac{(M-1)}{1 - (1 - p_r)^2 L_2}\left[\phi_{2,0}\left(1 + \frac{p_r L_1}{\phi_{1,0}}\right)\right.\right.$$
$$\left.\left. + (1 - p_r)\phi_{2,1}\right] + \frac{1}{p_r}\right] - 1 \quad (10)$$

As fixed frequency trains are commonly used in experiments, and Poisson-like statistics have been observed in the ISI distribution from neurons in the visual cortex [40, 41], explicit expressions of $Q^{ss}(b)$ in these two special cases are desirable.

For constant ISI $t_s = T$, also called a fixed frequency $f = 1/T$ AP train, we have $g(t_s) = \delta(t_s - T)$. In SI Section 5(a), we shown that $Q^{ss}(b)$ in this case reduces to a binomial distribution:

$$Q^{ss}(b) = B(M, p_{rb}, b) \quad (11)$$

$$\text{where, } p_{rb} = \frac{(1 - e^{-kT})p_r}{1 - (1 - p_r)e^{-kT}} \quad (12)$$

The exact binomial distribution has the parameters M and a modified release probability p_{rb} given by Eq. (12), which depends on the interval T and growth rate k apart from p_r. We do Kinetic Monte Carlo (KMC) simulations to verify our analytical predictions. In Fig 3a, we see that the exact distribution $Q^{ss}(b)$ (Eq. (11)) matches perfectly the data from KMC simulations (code in SI) of this process (see the red symbols). Note that Eq. (11) is not equivalent to an approximate binomial often used in the literature $B(\langle n_- \rangle, p_r, b)$. In fact we calculate exactly $\langle n_- \rangle = M p_{rb}/p_r$ (see SI Section 5(a)), and use this to plot $B(\langle n_- \rangle, p_r, b)$ (dashed line) in Fig 3a which deviates significantly. In the large T limit, from Eq. (12), $p_{rb} \to p_r$ (the release probability in the steady state), as one would expect. The mean and CV^2 of b from Eq. (11) are:

$$\langle b \rangle = M p_{rb} \quad \text{and,} \quad CV^2 = \frac{(1 - p_{rb})}{M p_{rb}}, \qquad (13)$$

(a)   (b)

FIG. 3: Steady state quantal content distributions for (a) fixed AP arrival times ($f = 1/T = 20$ Hz), and (b) exponentially distributed ISIs (rate $f_0 = 20$ Hz). The common parameters are $M = 50$, $p_r = 0.5$, $k = 2$ sec^{-1}. The exact theoretical curves (thick lines) match the KMC data (red symbols). In (a) an approximate (see text) binomial distribution (dashed line) is shown alongside the exact binomial. In (b) the binomial distribution of Eq. (11) with $T \to 1/f_0$ (dashed line) deviates from the exact non-binomial distribution.

For Poisson AP train, the ISIs are exponentially distributed with $g(t_s) = f_0 e^{-f_0 t_s}$. Obtaining $\phi_{n,m}$ and L_n in this case, Eq. (9) simplifies (see details in SI Section 5(b)) to yield the following quantal content distribution:

$$Q^{ss}(b) = \sum_{n=0}^{\infty} (-1)^{n-b} \binom{n}{b} \binom{M}{n} \frac{n!(kp_r)^n}{\prod_{i=1}^{n}(f_0 + ki - (1 - p_r)^i f_0)}.$$

(14)

The corresponding mean and CV^2 for the Poisson case

(see Section 5(b) in SI) are:

$$\langle b \rangle = \frac{M k p_r}{k + f_0 p_r} \quad \text{and}$$

$$CV^2 = \frac{1}{M} \left(\frac{2(M - 1)(k + f_0 p_r)}{2k + f_0 p_r (2 - p_r)} + \frac{f_0}{k} - M + \frac{1}{p_r} \right).$$

(15)

Note that this distribution is no longer a binomial. In Fig 3b we see that the exact formula (solid line) for $Q^{ss}(b)$ matches very well with the data obtained from KMC simulation (in symbols). Just to demonstrate that a Binomial approximation would be completely inadequate in this Poisson case, we plot Eq. (11)) with $T = 1/f_0$ where f_0 is the Poisson rate – the deviation is stark.

Experimental verification and estimation of model parameters: To validate the predictions of our model, we performed electrophysiological experiments (see Section 9 of SI). The comparison between theory and experiment is done in Fig 4. In Figs 4a and 4b, for synapse-1 (left) and synapse-2 (right) respectively, we plot the time-varying fluctuating quantal contents for a single tetanic burst (in blue), as well as the mean quantal content over 20 repeats (in red). In both neurons the mean quantal content reaches a steady-state value beyond 0.8 s (see the yellow box). Considering quantal contents between times $0.8 - 1.0$ s over all the 20 histories, we obtained the normalised histogram of the experimental quantal contents – these are shown (in green) in Figs 4c and 4d.

After the first AP (at 0 sec), the quantal content b_0 should follow a binomial distribution $B(M, p_{r0}, b_0)$, since docking sites should have maximal occupancy. From the 20 experimental histories we obtained the mean and variance of b_0 and using the theoretical formulas of mean = $M p_{r0}$ and variance = $M p_{r0}(1 - p_{r0})$, we obtained M by eliminating p_{r0}. Next we equate the experimental steady-state $\langle b \rangle$ (Figs 4c and 4d) to the theoretical mean $\langle b \rangle = M p_{rb}$ (Eq. (13)), and determine the parameter p_{rb}. With these parameters, we plot the exact distribution Eq. (11) (black curves) for the two synapses and they match quite well to the experimental data in Figs 4c and 4d (in green). The CV^2 automatically turn out quite close: for synapse-1, it is 0.039 (theory) vs 0.032 (experiment), while for synapse-2, the values are 0.028 (theory) vs 0.022 (experiment). Thus we have shown how the full experimental distribution, not just few moments, may be matched to the analytical distribution function. Such comparative study of full experimental and theoretical distributions maybe done for stochastic inputs with Poisson APs (using Eq. 14) or any other $g(t_s)$ (using Eq. 9) in future.

Note that the moments in the fixed ISI case depend on the composite parameter p_{rb} (Eq. (13)) and not separately on the model parameters p_r and k, and hence we cannot estimate them using the fixed ISI data. But

FIG. 4: The experimental mean quantal content $\langle b \rangle$ is plotted (in red) against time, for (a) synapse-1 and (b) synapse-2. The blue curves represent one of the 20 experimental realizations of the quantal content. The yellow box shows the part of data we considered to be in the steady state for plotting the experimental histograms of quantal content in (c) and (d) (in green). In (c) and (d), the parameter values used for the exact theoretical distributions (black curves) for synapse-1 are $M = 50$ and $p_{rb} = 0.345$ and for the synapse-2 are $M = 98$ and $p_{rb} = 0.266$.

that would not be the case for Poisson APs – see Eq. (15)) where the explicit dependence on p_r and k appear. Thus future experimental data on Possion APs may be used to estimate the model parameters. In fact this may be done for stochastic inputs with any other $g(t_s)$ – Eq. (10) may be used in such cases.

Exact results are rare and desirable in biological contexts. Here, we analyze the basic model of synaptic transmission between neurons applicable to all chemical synapses. Although we look at the restricted regime of steady state and uncorrelated stochastic AP inputs, we provide full exact general distributions including moments. We also show how the results may be used to compare with experimental data and indicate how model parameters may be estimated in future.

Acknowledgements: DD acknowledges IIT Bombay for funding. KR thanks IIT Bombay for Institute Ph.D. fellowship. AS acknowledges support by NIH/NIDCD grant R01DC019268. EF acknowledges support by Bundesministerium für Bildung und Forschung grant 01GQ2001.

* ashok.prasad@colostate.edu
† dibyendu@phy.iitb.ac.in

[1] S. Herculano-Houzel, Front Hum Neurosci **3**, 31 (2009).

[2] Y. Tang, J. R. Nyengaard, D. M. De Groot, and H. J. Gundersen, Synapse **41**, 258 (2001).

[3] E. R. Kandel, J. Koester, S. Mack, and S. Siegelbaum, *Principles of neural science*, sixth edition. ed. (McGraw Hill, New York, 2021) pp. l, 1646 pages.

[4] J. Del Castillo and B. Katz, The Journal of Physiology **124**, 560 (1954).

[5] V. N. Murthy and P. De Camilli, Annu Rev Neurosci **26**, 701 (2003).

[6] S. O. Rizzoli and W. J. Betz, Nat Rev Neurosci **6**, 57 (2005).

[7] A. A. Alabi and R. W. Tsien, Cold Spring Harb Perspect Biol **4**, a013680 (2012).

[8] E. Neher, Neuron **87**, 1131 (2015).

[9] S. Weis, R. Schneggenburger, and E. Neher, Biophysical Journal **77**, 2418 (1999).

[10] D. A. Rusakov, L. P. Savtchenko, and P. E. Latham, Trends in Neurosciences **43**, 363 (2020).

[11] M. S. Goldman, Neural Computation **16**, 1137 (2004).

[12] C. Zhang and C. S. Peskin, Proceedings of the National Academy of Sciences **112**, 14954 (2015).

[13] M. V. Tsodyks and H. Markram, Proceedings of the National Academy of Sciences **94**, 719 (1997).

[14] J. S. Dittman and W. G. Regehr, Journal of Neuroscience **18**, 6147 (1998).

[15] J. S. Dittman, A. C. Kreitzer, and W. G. Regehr, Journal of Neuroscience **20**, 1374 (2000).

[16] E. M. McLachlan, Int Rev Physiol **17**, 49 (1978).

[17] S. Redman, Physiological Reviews **70**, 165 (1990).

[18] A. Arleo, T. Nieus, M. Bezzi, A. d'Errico, E. d'Angelo, and O. J.-M. Coenen, Neural Computation **22**, 2031 (2010).

[19] R. Rosenbaum, J. Rubin, and B. Doiron, PLoS Computational Biology **8**, e1002557 (2012).

[20] C. Pulido, F. F. Trigo, I. Llano, and A. Marty, Neuron **85**, 159 (2015).

[21] G. Malagon, T. Miki, I. Llano, E. Neher, and A. Marty, Journal of Neuroscience **36**, 4010 (2016).

[22] Z. Vahdat, Z. Xu, and A. Singh, in *2019 IEEE 58th Conference on Decision and Control (CDC)* (IEEE, 2019) pp. 4729–4734.

[23] M. D. Miyamoto, J Physiol **250**, 121 (1975).

[24] R. Malinow and R. W. Tsien, Nature **346**, 177 (1990).

[25] S. S. Kelly and N. Robbins, The Journal of Physiology **385**, 507 (1987).

[26] R. A. Silver, A. Momiyama, and S. G. Cull-Candy, The Journal of Physiology **510**, 881 (1998).

[27] H. Taschenberger, V. Scheuss, and E. Neher, The Journal of Physiology **568**, 513 (2005).

[28] E. Hanse and B. Gustafsson, Journal of Neuroscience **21**, 8362 (2001).

[29] C. Pulido and A. Marty, Physiological Reviews **97**, 1403 (2017).

[30] G. Malagon, T. Miki, V. Tran, L. C. Gomez, and A. Marty, eLife **9**, e52137 (2020).

[31] A. Brémaud, D. C. West, and A. M. Thomson, Proceedings of the National Academy of Sciences **104**, 14134 (2007).

[32] H. Wen, M. J. McGinley, G. Mandel, and P. Brehm, Proceedings of the National Academy of Sciences **113**, E378 (2016).

[33] A. Barri, Y. Wang, D. Hansel, and G. Mongillo, ENeuro **3** (2016).

[34] A. D. Bird, M. J. Wall, and M. J. Richardson, Frontiers in Computational Neuroscience **10**, 116 (2016).

[35] O. Bykowska, C. Gontier, A.-L. Sax, D. W. Jia, M. L. Montero, A. D. Bird, C. Houghton, J.-P. Pfister, and R. P. Costa, Frontiers in Synaptic Neuroscience **11**, 21 (2019).

[36] C. Gontier, S. C. Surace, and J.-P. Pfister, arXiv preprint arXiv:2201.07539 (2022).

[37] K. Mahfooz, M. Singh, R. Renden, and J. F. Wesseling, PLoS Computational Biology **12**, e1004855 (2016).

[38] E. G. Krächan, A. U. Fischer, J. Franke, and E. Friauf, The Journal of Physiology **595**, 839 (2017).

[39] R. P. Costa, P. J. Sjöström, and M. C. Van Rossum, Frontiers in Computational Neuroscience **7**, 75 (2013).

[40] M. N. Shadlen and W. T. Newsome, Journal of Neuroscience **18**, 3870 (1998).

[41] G. Maimon and J. A. Assad, Neuron **62**, 426 (2009).

[42] J. S. Lee, W.-K. Ho, E. Neher, and S.-H. Lee, Proceedings of the National Academy of Sciences **110**, 15079 (2013).

[43] K. Ikeda and J. M. Bekkers, Proceedings of the National Academy of Sciences USA **106**, 2945 (2009).

[44] B. G. Wilhelm, S. Mandad, S. Truckenbrodt, K. Kröhnert, C. Schäfer, B. Rammner, S. J. Koo, G. A. Claßen, M. Krauss, V. Haucke, H. Urlaub, and S. O. Rizzoli, Science **344**, 1023 (2014), https://www.science.org/doi/pdf/10.1126/science.1252884.

[45] C. W. Gardiner, *Handbook of stochastic methods*, Vol. 3 (springer Berlin, 1985).

[46] G. W. Davis, Annual Review of Neuroscience **29**, 307 (2006), pMID: 16776588, https://doi.org/10.1146/annurev.neuro.28.061604.135751.

[47] H. Taschenberger, A. Woehler, and E. Neher, Proceedings of the National Academy of Sciences **113**, E4548 (2016), https://www.pnas.org/doi/pdf/10.1073/pnas.1606383113.

[48] J. M. Ortega, O. Genc, and G. W. Davis, eLife **7**, e40385 (2018).

[49] B. O. Orr, R. D. Fetter, and G. W. Davis, Neuron (2022), 10.1016/j.neuron.2022.08.016.

[50] D. Keller, N. Babai, O. Kochubey, Y. Han, H. Markram, F. Schürmann, and R. Schneggenburger, PLoS Computational Biology **11**, e1004253 (2015).

[51] Y. Nakamura, H. Harada, N. Kamasawa, K. Matsui, J. S. Rothman, R. Shigemoto, R. A. Silver, D. A. DiGregorio, and T. Takahashi, Neuron **85**, 145 (2015).

Exact distribution of the quantal content in synaptic transmission Supplementary Information File

Krishna Rijal[a], Nicolas I.C. Müller[b], Eckhard Friauf[b], Abhyudai Singh[c], Ashok Prasad[d,1], and Dibyendu Das[a,1]

[a]Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India

[b]Animal Physiology Group, Department of Biology, University of Kaiserslautern, Kaiserslautern, Germany

[c]Departments of Electrical and Computer Engineering, Biomedical Engineering and Mathematical Sciences, University of Delaware, Newark, DE 19716, USA

[d]Department of Chemical and Biological Engineering, Colorado State University, Fort Collins, Colorado, USA

[1]To whom correspondence should be addressed: dibyendu@phy.iitb.ac.in; ashok.prasad@colostate.edu

July 12, 2023

1 Probability distribution of docked vesicle number n between two consecutive action potentials

Let $p(n, t|n'_+)$ denote the probability distribution of the number n of docked (ready to release) vesicles at time t, starting from $n = n'_+$ at $t = 0$ (i.e. from the instant of the last action potential (AP)). Thus t denotes any time up to the next AP, i.e. $t < t_s$ where t_s is a time when the next AP arrives. In the interval t to $t + dt$, the number of docked vesicles may increase from $n - 1$ to n with the probability $k(M - (n-1))dt$. Here k is the growth rate per empty docking sites, and M is maximum number of docking sites. Consequently, the master equation for the time evolution of $p(n, t|n'_+)$ between two successive APs is:

$$\frac{\partial p(n, t|n'_+)}{\partial t} = k\left(M - (n-1)\right) p(n-1, t|n'_+) - k(M - n)p(n, t|n'_+) \qquad (1)$$

We define a generating function for the above probability distribution:

$$F(q, t|n'_+) = \sum_n q^n p(n, t|n'_+) \qquad (2)$$

Here the limits of the above sum are $n'_+ \leq n \leq M$. Otherwise, we assume $p(n, t|n'_+) = 0$ if n'_+ falls outside this range. Multiplying Eq. (1) by $\sum_n q^n$ on both sides, we get a PDE for the generating function F:

$$\frac{\partial F}{\partial t} = + kM \sum_n q^n p(n-1, t|n'_+) - k \sum_n q^n (n-1)p(n-1, t|n'_+)$$

$$- kM \sum_n q^n p(n, t|n'_+) + k \sum_n q^n n p(n, t|n'_+) = kMqF - kq^2\frac{\partial F}{\partial q} - kMF + kq\frac{\partial F}{\partial q}$$

Rearranging the terms, we have,

$$\frac{\partial F}{\partial t} + kq(q-1)\frac{\partial F}{\partial q} = kM(q-1)F \qquad (3)$$

We use the method of Lagrange characteristics to solve the Eq. (3). The result is:

$$F(q, t|n'_+) = q^M \left[1 - \frac{(q-1)}{q} e^{-kt} \right]^{M-n'_+} \tag{4}$$

We do binomial expansion of Eq. (4) to get

$$F = q^{n'_+} \left[q(1 - e^{-kt}) + e^{-kt} \right]^{M-n'_+}$$

$$= \sum_{n=0}^{M-n'_+} \binom{M - n'_+}{n} \left[q^{n+n'_+} \left(1 - e^{-kt} \right)^n \right] \left(e^{-kt} \right)^{M-n'_+-n}$$

and relabelling $n + n'_+ \to n$ we get

$$F = \sum_{n=n'_+}^{M} \binom{M - n'_+}{n - n'_+} q^n \left(1 - e^{-kt} \right)^{n-n'_+} \left(e^{-kt} \right)^{M-n} \tag{5}$$

Next we can compare Eq. (5) with Eq. (2) to get the expression of $p(n, t|n'_+)$:

$$\boxed{p(n, t|n'_+) = \binom{M - n'_+}{n - n'_+} \left(1 - e^{-kt} \right)^{n-n'_+} e^{-[k(M-n)]t}} \tag{6}$$

for $n'_+ \leq n < \infty$. Note that for $n > M$, the probability goes to zero automatically in the above expression.

2   Self-consistent equation for the quantal content distribution $P_+^{ss}(n_+)$ in the steady state

Let us denote the time intervals between $(m - 1)^{th}$ AP and m^{th} AP by t_{sm} and between the $(m - 2)^{th}$ AP and $(m - 1)^{th}$ AP by t_{sm-1}. We first relate the probability $P_+^m(n_{+,m}, t_{sm})$ of having $n_{+,m}$ number of docked vesicles after the m^{th} AP, to the probability $P_+^{m-1}(n_{+,m-1}, t_{sm-1})$ of having $n_{+,m-1}$ number of vesicles just after the $(m-1)^{th}$ AP. To do so, we need to sum over all possible paths beginning at $n = n_{+,m-1}$ after the $(m - 1)^{th}$ action potential and ending at $n = n_{+,m}$ after the m^{th} AP.

The sum involves the product of three probabilities due to a sequence of steps: (i) initial $P_+^{m-1}(n_{+,m-1}, t_{sm-1})$, then (ii) the probability $p(n_{+,m} + b, t_{sm}|n_{+,m-1})$ that $n_{+,m-1}$ rises to $n_{+,m} + b$ in time t_{sm}, and (iii) finally the probability that at t_{sm}, the number $n_{+,m} + b$ of docked vesicles suddenly reduce to $n_{+,m}$ due to the AP triggered vesicle release, with the associated quantal content b being binomially distribution $B(n_{+,m}+b, p_r, b)$. After summing over all possible initial number $n_{+,m-1}$ and quantal content b we have,

$$
P_+^m(n_{+,m}, t_{sm}) = \sum_{n_{+,m-1},b} B(n_{+,m}+b, p_r, b) p(n_{+,m}+b, t_{sm}|n_{+,m-1}) P_+^{m-1}(n_{+,m-1}, t_{sm-1})
$$

(7)

We may follow the same procedure for $P_+^{m-1}(n_{+,m-1}, t_{sm-1})$ and so on, and build up a hierarchy of relations for any interval between two APs. The problem remains hard to solve as result for every interval depends on the history of the previous interval. We are particularly interested in the steady state attained after several APs appear, i.e. $m \gg 1$ and the output quantal contents statistically stabilise. We integrate Eq. (7) over the joint probability distribution $g_2(t_{sm}, t_{sm-1})$ of t_{sm} and t_{sm-1}. We get

$$
\int_0^\infty \int_0^\infty dt_{sm} dt_{sm-1} g_2(t_{sm}, t_{sm-1}) P_+^m(n_{+,m}, t_{sm})
$$
$$
= \int_0^\infty \int_0^\infty dt_{sm} dt_{sm-1} g_2(t_{sm}, t_{sm-1})
$$
$$
\times \sum_{n_{+,m-1},b} p(n_{+,m} + b, t_{sm}|n_{+,m-1}) B(n_{+,m} + b, p_r, b) P_+^{m-1}(n_{+,m-1}, t_{sm-1}) \quad (8)
$$

If the interspike intervals (ISIs) are not correlated, we would have $g_2(t_{sm}, t_{sm-1}) = g(t_{sm}) g(t_{sm-1})$, where $g(t_{sm})$ and $g(t_{sm-1})$ are the normalised distributions of t_{sm} and t_{sm-1} respectively. Then,

$$
\int_0^\infty \int_0^\infty dt_{sm} dt_{sm-1} g(t_{sm}) g(t_{sm-1}) P_+^m(n_{+,m}, t_{sm})
$$
$$
= \int_0^\infty \int_0^\infty dt_{sm} dt_{sm-1} g(t_{sm}) g(t_{sm-1})
$$

$$\times \sum_{n_{+,m-1},b} p(n_{+,m}+b, t_{sm}|n_{+,m-1})B(n_{+,m}+b, p_r, b)P_+^{m-1}(n_{+,m-1}, t_{sm-1})$$

$$\Rightarrow \int_0^\infty dt_{sm}g(t_{sm})P_+^m(n_{+,m}, t_{sm})$$

$$= \sum_{n_{+,m-1},b} \int_0^\infty dt_{sm}g(t_{sm})p(n_{+,m}+b, t_{sm}|n_{+,m-1})B(n_{+,m}+b, p_r, b)$$

$$\times \int_0^\infty dt_{sm-1}g(t_{sm-1})P_+^{m-1}(n_{+,m-1}, t_{sm-1}) \tag{9}$$

⁴⁹ In the long time limit, the steady state probability distribution of vesicle number
⁵⁰ after any AP is $P_+^{ss}(n_{+,m}) = \int_0^\infty dt_{sm-1}g(t_{sm-1})P_+^{m-1}(n_{+,m-1}, t_{sm-1})$, which does not
⁵¹ depend any more on the exact history of the spike train. Dropping the subscripts m
⁵² (in the steady state), and setting $t_{sm} = t_s$ and $n_{+,m-1} = n'_+$, the Eq. (9) becomes

$$\boxed{P_+^{ss}(n_+) = \sum_{n'_+,b} \int_0^\infty dt_s g(t_s)p(n_++b, t_s|n'_+)B(n_++b, p_r, b)P_+^{ss}(n'_+)} \tag{10}$$

⁵³ Note that this is a self-consistent equation for the unknown $P_+^{ss}(n_+)$, when we
⁵⁴ know $g(t_s)$, $B(n_++b, p_r, b)$ and $p(n_++b, t_s|n'_+)$ (Eq. (6)). This equation appears in
⁵⁵ the main manuscript. We analytically treat it in the next section.

3 Generating function of the distribution $P_+^{ss}(n_+)$ in the steady state

⁵⁸ We define a generating function of $P_+^{ss}(n_+)$ as follows and solve it in this section:

$$F_+(q) = \sum_{n_+} q^{n_+} P_+^{ss}(n_+) \tag{11}$$

⁵⁹ In Eq. (10), the limits of the summation are $0 \le b \le M - n_+$ and $0 \le n'_+ \le n_+ + b$.
⁶⁰ Since anyways, $p(n_++b, t_s|n'_+) = 0$ outside the limits (see Eq. (6)), we may as well
⁶¹ use the range $0 \le b, n'_+ \le \infty$. Multiplying Eq. (10) by q^{n_+} and summing over n_+

 (with $0 \leq n_+ \leq \infty$), we have:

$$F_+(q) = \sum_{n'_+,b} \int_0^\infty dt_s g(t_s) \sum_{n_+} q^{n_+} p(n_+ + b, t_s | n'_+) B(n_+ + b, p_r, b) P_+^{ss}(n'_+) \qquad (12)$$

 We introduce a new variable $\tilde{n_+} = n_+ + b$ with $b \leq \tilde{n_+} \leq \infty$. Note that $B(\tilde{n_+}, p_r, b) = 0$
 for $\tilde{n_+} < b$, so we may keep the range $0 \leq \tilde{n_+} \leq \infty$.

$$F_+(q) = \int_0^\infty dt_s g(t_s) \sum_{n'_+,b} \sum_{\tilde{n_+}} q^{\tilde{n_+}-b} p(\tilde{n_+}, t_s | n'_+) B(\tilde{n_+}, p_r, b) P_+^{ss}(n'_+) \qquad (13)$$

Summing over b after substituting $B(\tilde{n_+}, p_r, b) = \binom{\tilde{n_+}}{b} p_r^b (1 - p_r)^{\tilde{n_+}-b}$, we have

$$F_+(q) = \int_0^\infty dt_s g(t_s) \sum_{n'_+} \sum_{\tilde{n_+}} \left(1 - p_r + \frac{p_r}{q}\right)^{\tilde{n_+}} q^{\tilde{n_+}} p(\tilde{n_+}, t_s | n'_+) P_+^{ss}(n'_+)$$

$$= \int_0^\infty dt_s g(t_s) \sum_{n'_+} \sum_{\tilde{n_+}} (q - qp_r + p_r)^{\tilde{n_+}} p(\tilde{n_+}, t_s | n'_+) P_+^{ss}(n'_+)$$

 Doing summation over $\tilde{n_+}$ and using the definition of the generating function F (from
 Eq. (2)), we get

$$F_+(q) = \int_0^\infty dt_s g(t_s) \sum_{n'_+} F\left((q - qp_r + p_r), t_s | n'_+\right) P_+^{ss}(n'_+)$$

Defining $h(q) = (q - qp_r + p_r)$ and using Eq. (4) for the explicit form of function F:

$$F_+(q) = \int_0^\infty dt_s g(t_s) \sum_{n'_+} h^M \left[1 - \frac{h-1}{h} e^{-kt_s}\right]^{M-n'_+} P_+^{ss}(n'_+)$$

$$= \int_0^\infty dt_s g(t_s) \left[h - (h-1)e^{-kt_s}\right]^M \sum_{n'_+} \left(\frac{h}{h - (h-1)e^{-kt_s}}\right)^{n'_+} P_+^{ss}(n'_+)$$

$$= \int_0^\infty dt_s g(t_s) \left[h - (h-1)e^{-kt_s}\right]^M F_+ \left[\frac{h}{h - (h-1)e^{-kt_s}}\right]$$

$$= \int_0^\infty dt_s g(t_s) \left[h(1 - e^{-kt_s}) + e^{-kt_s}\right]^M F_+ \left[\frac{h}{h(1 - e^{-kt_s}) + e^{-kt_s}}\right] \qquad (14)$$

 In the above we have used the definition of F_+ from Eq(11). In Eq. (14) we have

a self-consistent integral equation for the function F_+ with two different arguments, and involving an integration over the inter-spike interval t_s. Clearly, it is difficult to solve the equation directly for $F_+(q)$. But as we show below, $F_+(q)$ may be obtained as a series expansion around $q = 1$, that is

$$F_+(q) = \sum_{n=0}^{\infty} \frac{(q-1)^n}{n!} F_+^{(n)}(1) \tag{15}$$

where the series coefficients $F_+^{(n)}(1) = \left. \frac{\partial^n F_+(q)}{\partial q^n} \right|_{q=1}$. The procedure of obtaining these coefficients are as follows.

Taking derivative with respect to q on both side of Eq. (14), we have

$$\begin{aligned}
\frac{\partial F_+(q)}{\partial q} =& \frac{\partial h}{\partial q} \int_0^\infty dt_s g(t_s) \left(M \left[h(1 - e^{-kt_s}) + e^{-kt_s} \right]^{M-1} (1 - e^{-kt_s}) F_+ \left[\frac{h}{h(1 - e^{-kt_s}) + e^{-kt_s}} \right] \right. \\
&\left. + \left[h(1 - e^{-kt_s}) + e^{-kt_s} \right]^M \frac{\partial F_+[\,.\,]}{\partial [\,.\,]} \frac{e^{-kt_s}}{(h(1 - e^{-kt_s}) + e^{-kt_s})^2} \right)
\end{aligned} \tag{16}$$

where $[\,.\,]$ is a short hand for $\left[\frac{h}{h(1-e^{-kt_s})+e^{-kt_s}} \right]$. Since $\frac{\partial h}{\partial q} = (1 - p_r)$, $h(q) \big|_{q=1} = 1$, $[\,.\,] \big|_{q=1} = 1$, $F_+(1) = 1$, and $\left. \frac{\partial F_+[\,.\,]}{\partial [\,.\,]} \right|_{q=1} = \left. \frac{\partial F_+(q)}{\partial q} \right|_{q=1}$, we have

$$\left. \frac{\partial F_+(q)}{\partial q} \right|_{q=1} = (1 - p_r) \int_0^\infty dt_s g(t_s) \left(M(1 - e^{-kt_s}) + \left. \frac{\partial F_+(q)}{\partial q} \right|_{q=1} e^{-kt_s} \right) \tag{17}$$

Defining the quantities $\psi_{n,m}$ and L_n as follows,

$$\begin{aligned}
\psi_{n,m} &= \int_0^\infty dt_s g(t_s) e^{-nkt_s} (e^{kt_s} - 1)^{n-m} \\
L_n &= \psi_{n,n} = \int_0^\infty dt_s g(t_s) e^{-nkt_s}
\end{aligned} \tag{18}$$

we have

$$F_+^{(1)}(1) = (1 - p_r) \left(M \psi_{1,0} + L_1 F_+^{(1)}(1) \right) \tag{19}$$

where we have set $\left. \frac{\partial F_+(q)}{\partial q} \right|_{q=1} = F_+^{(1)}(1)$ according to the general definition of the n-th series coefficient above.

80 Next we may take derivative of Eq. (16) with respect to q and set $q = 1$ and obtain

81 an equation for $F_+^{(2)}(1)$. We find that it is linearly coupled to $F_+^{(1)}(1)$ as follows:

$$F_+^{(2)}(1) = (1 - p_r)^2 \left(M(M-1)\psi_{2,0} + (M-1)\psi_{2,1}F_+^{(1)}(1) + L_2 F_+^{(2)}(1) \right). \quad (20)$$

82 We may follow the same procedure to obtain equations for the higher derivatives.

83 It is clumsy to do it manually, so we use Mathematica for this. We use symbolic

84 derivatives of Eq. (16) and automatic replacement of terms by the ψ functions using

85 the *Replace* syntax. The result are the equations for the next two series coefficients.

$$F_+^{(3)}(1) = (1 - p_r)^3 \left(M(M-1)(M-2)\psi_{3,0} + (M-1)(M-2)\psi_{3,1}F_+^{(1)}(1) \right.$$
$$\left. + (M-2)\psi_{3,2}F_+^{(2)}(1) + L_3 F_+^{(3)}(1) \right) \quad (21)$$

$$F_+^{(4)}(1) = (1 - p_r)^4 \left(M(M-1)(M-2)(M-3)\psi_{4,0} + (M-1)(M-2)(M-3)\psi_{4,1}F_+^{(1)}(1) \right.$$
$$\left. + (M-2)(M-3)\psi_{4,2}F_+^{(2)}(1) + (M-3)\psi_{4,3}F_+^{(3)}(1) + L_4 F_+^{(4)}(1) \right) \quad (22)$$

86 Eqs. (19) - (22) relate the four derivates $F_+^{(1)}(1), F_+^{(2)}(1), F_+^{(3)}(1)$ and $F_+^{(4)}(1)$ linearly.

87 So, it is easy to obtain solutions by inverting a 4×4 matrix, using Mathematica.

88 The explicit expressions of the four coefficients are

$$F_+^{(1)}(1) = \frac{(1-p_r)M\psi_{1,0}}{1 - (1-p_r)L_1} \quad (23)$$

$$F_+^{(2)}(1) = \frac{(1-p_r)^2 M(M-1)}{1 - (1-p_r)^2 L_2} \left[\psi_{2,0} + \frac{2(1-p_r)\psi_{2,1}\psi_{1,0}}{1 - (1-p_r)L_1} \right] \quad (24)$$

$$F_+^{(3)}(1) = \frac{(1-p_r)^3 M(M-1)(M-2)}{1 - (1-p_r)^3 L_3} \left[\psi_{3,0} + \frac{3(1-p_r)\psi_{3,1}\psi_{1,0}}{1 - (1-p_r)L_1} \right.$$
$$\left. + \frac{3(1-p_r)^2 \psi_{3,2}\psi_{2,0}}{1 - (1-p_r)^2 L_2} + \frac{6(1-p_r)^3 \psi_{3,2}\psi_{2,1}\psi_{1,0}}{(1 - (1-p_r)L_1)(1 - (1-p_r)^2 L_2)} \right] \quad (25)$$

$$F_+^{(4)}(1) = \frac{(1-p_r)^4 M(M-1)(M-2)(M-3)}{1 - (1-p_r)^4 L_4} \left[\psi_{4,0} + \frac{4(1-p_r)\psi_{4,1}\psi_{1,0}}{1 - (1-p_r)L_1} \right.$$
$$\left. + \frac{4(1-p_r)^3 \psi_{4,3}\psi_{3,0}}{1 - (1-p_r)^3 L_3} + \frac{6(1-p_r)^2 \psi_{4,2}\psi_{2,0}}{1 - (1-p_r)^2 L_2} + \frac{12(1-p_r)^3 \psi_{4,2}\psi_{2,1}\psi_{1,0}}{(1 - (1-p_r)L_1)(1 - (1-p_r)^2 L_2)} \right.$$

$$+\frac{12(1-p_r)^4\psi_{4,3}\psi_{3,1}\psi_{1,0}}{(1-(1-p_r)L_1)(1-(1-p_r)^3L_3)}+\frac{12(1-p_r)^5\psi_{4,3}\psi_{3,2}\psi_{2,0}}{(1-(1-p_r)^2L_2)(1-(1-p_r)^3L_3)}$$

$$+\frac{24(1-p_r)^6\psi_{4,3}\psi_{3,2}\psi_{2,1}\psi_{1,0}}{(1-(1-p_r)L_1)(1-(1-p_r)^2L_2)(1-(1-p_r)^3L_3)}\Bigg] \tag{26}$$

In the above expressions, Eqs. (23) - (26), we notice the following patterns. The terms outside the square brackets are of the form $\binom{M}{n}n!\frac{(1-p_r)^n}{1-(1-p_r)^nL_n}$ for the n-th coefficient. Except for the first term, the denominators of the terms inside the square brackets for the n-th coefficient involve factors $(1-(1-p_r)^jL_j)$ with $j\neq n$ or 0. The most crucial fact is that each such term may be associated with a set of integers which are subsets of the set of integers $\{1,2,\cdots,n-1\}$. For example, a subset of $z\leq(n-1)$ positive integer elements is $S_{(n-1)}=(m_z,m_{z-1},\cdots,m_1)=\{m_i\}$ with $i=z,z-1,\cdots,1$, and every $m_i\in\{1,2,\cdots,n-1\}$ and $m_z>m_{z-1}>\cdots m_1$. The power of $(1-p_r)$ in the numerator of each term is equal to $\sum_{i=1}^z m_i$. The subscript of ψ factors are formed by pairs of successive integers drawn from the ordered array of numbers: $n,m_z,m_{z-1},\cdots,m_1,0$. The numeric factors in any numerator can be absorbed into the ψ factors by introducing the quantities $\phi_{n,m}=\binom{n}{m}\psi_{n,m}$. Putting together all these observations of the patterns, we may write down a general form of $F_+^{(n)}(1)$ as follows:

$$F_+^{(n)}(1)=\binom{M}{n}n!\frac{(1-p_r)^n}{1-(1-p_r)^nL_n}\left[\phi_{n,0}+\sum_{\{S_{(n-1)}\}}\frac{(1-p_r)^{\sum_i m_i}\phi_{n,m_z}\phi_{m_z,m_{z-1}}\cdots\phi_{m_1,0}}{\prod_i(1-(1-p_r)^{m_i}L_{m_i})}\right] \tag{27}$$

where the sum over $\{S_{(n-1)}\}$ runs over all such subsets as $S_{(n-1)}$ mentioned above.

Using Eq. (27) in Eq. (15), we can finally write down the expression of $F_+(q)$ as a series expansion around $q=1$,

$$F_+(q)=\sum_{n=0}^{\infty}(q-1)^n\binom{M}{n}\frac{(1-p_r)^n}{1-(1-p_r)^nL_n}\left[\phi_{n,0}+\sum_{\{S_{(n-1)}\}}\frac{(1-p_r)^{\sum_i m_i}\phi_{n,m_z}\phi_{m_z,m_{z-1}}\cdots\phi_{m_1,0}}{\prod_i(1-(1-p_r)^{m_i}L_{m_i})}\right] \tag{28}$$

4 Relations between the generating functions, and the quantal content distribution $Q^{ss}(b)$

Let us denote the probability distribution of having n_- number of docked vesicles before an AP in the steady state by $P_-^{ss}(n_-)$. It is easy to see from similar arguments as in Section 2, that $P_-^{ss}(n_-)$ will be related to the three distributions: $P_+^{ss}(n_+)$ (to have an initial number), $p(n_-|t_s, n_+)$ (for docked vesicles to build up), and $g(t_s)$ (for AP to appear). Thus

$$P_-^{ss}(n_-) = \sum_{n_+} \int_0^\infty p(n_-, t_s|n_+) P_+^{ss}(n_+) g(t_s) dt_s \tag{29}$$

We define its generating function as

$$F_-(q) = \sum_{n_-=0}^\infty q^{n_-} P_-^{ss}(n_-) \tag{30}$$

The ultimate object of our interest, namely the distribution of quantal content b in the steady state, is related to the distribution $P_-^{ss}(n_-)$ as:

$$Q^{ss}(b) = \sum_{n_-} P_-^{ss}(n_-) B(n_-, p_r, b) \tag{31}$$

Note that the the crucial fact is that we are averaging the well-known binomial distribution now, over the stochastically generated docked vesicle number n_- just prior to an action potential. As we would see, the resultant averaged distribution would be different from the original binomial. The generating function corresponding to $Q^{ss}(b)$ is defined as

$$F_b(q) = \sum_{b=0}^\infty q^b Q^{ss}(b) \tag{32}$$

We have four distributions $p(n_-, t_s|n_+)$, $P_+^{ss}(n_+)$, $P_-^{ss}(n_-)$, and $Q^{ss}(b)$ and four corresponding generating functions F, F_+, F_- and F_b, respectively. Of these, we have already calculated F (Eq. (4)) and F_+ (Eq. (28)). We would now proceed to relate the generating functions F_- and F_b to F_+. Once F_b is known explicitly using the

expression of F_+, we would then obtain $Q^{ss}(b)$ by taking suitable derivatives of $F_b(q)$ with respect to q.

Using Eq. (29) in Eq. (30) we get,

$$
\begin{aligned}
F_-(q) &= \sum_{n_-} q^{n_-} \left(\sum_{n_+} \int_0^\infty p\left(n_-, t_s | n_+\right) P_+^{ss}\left(n_+\right) g(t_s) dt_s \right) \\
&= \int_0^\infty g(t_s) dt_s \left\{ \sum_{n_+} P_+^{ss}\left(n_+\right) \sum_{n_-} p\left(n_-, t_s | n_+\right) q^{n_-} \right\} \\
&= \int_0^\infty g(t_s) dt_s \left\{ \sum_{n_+} P_+^{ss}\left(n_+\right) F\left(q, t_s | n_+\right) \right\}
\end{aligned}
$$

In the above we have used the definition of the function F. Substituting the explicit form of $F(q, t_s | n_+)$ from Eq. (4), we have

$$
\begin{aligned}
F_-(q) &= \int_0^\infty g(t_s) dt_s \left\{ \sum_{n_+} P_+^{ss}\left(n_+\right) q^M \left[1 + \frac{1-q}{q} e^{-kt_s}\right]^{M-n_+} \right\} \\
&= \int_0^\infty g(t_s) dt_s \left\{ \left[q + (1-q)e^{-kt_s}\right]^M F_+ \left(\frac{1}{1 + \frac{1-q}{q} e^{-kt_s}} \right) \right\}
\end{aligned} \tag{33}
$$

In the above we have used the definition of F_+.

Substituting $\frac{q-p_r}{1-p_r}$ in place of q in Eq. (14), and noticing $h\left(\frac{q-p_r}{1-p_r}\right) = q$, we get

$$
F_+ \left(\frac{q - p_r}{1 - p_r} \right) = \int_0^\infty g(t_s) dt_s \left\{ \left[q + (1-q)e^{-kt_s}\right]^M F_+ \left(\frac{1}{1 + \frac{1-q}{q} e^{-kt_s}} \right) \right\} \tag{34}
$$

Comparing Eqs. (33) and (34), we immediately get,

$$
F_-(q) = F_+ \left(\frac{q - p_r}{1 - p_r} \right) \tag{35}
$$

Substituting Eq. (31) in Eq. (32) with the binomial distribution $B(n_-, p_r, b)$ written explicitly, we have

$$
F_b(q) = \sum_b q^b \left(\sum_{n_-} P_-^{ss}(n_-) \binom{n_-}{b} p_r^b (1 - p_r)^{n_- - b} \right)
$$

$$= \sum_{n_-} P_-^{ss}(n_-) \left(qp_r + (1 - p_r)\right)^{n_-}$$

$$= F_- \left(qp_r + 1 - p_r\right)$$

$$= F_+ \left(\frac{p_r(q - 1)}{1 - p_r} + 1\right) \tag{36}$$

Here we have used the definition of the function F_-, and the relation of F_+ to F_- from Eq. (35). Using the explicit form of F_+ from Eq. (28), the Eq. (36) gives,

$$F_b(q) = \sum_{n=0}^{\infty} (q - 1)^n \binom{M}{n} \frac{p_r^n}{1 - (1 - p_r)^n L_n} \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})}\right] \tag{37}$$

We may invert Eq. (32), to obtain the quantal content distribution $Q^{ss}(b)$ from successive derivatives of $F_b(q)$ as follows:

$$Q^{ss}(b) = \frac{1}{b!} \left.\frac{\partial^b F_b(q)}{\partial q^b}\right|_{q=0} \tag{38}$$

Note that q is present in $F_b(q)$ only through the factor $(q - 1)^n$ in Eq. (37). Using the identity $\left.\frac{\partial^b (q-1)^n}{\partial q^b}\right|_{q=0} = \frac{n!}{(n-b)!}(-1)^{n-b}$, we obtain our final desired result

$$Q^{ss}(b) = \sum_{n=0}^{\infty} \binom{n}{b}(-1)^{n-b}\binom{M}{n} \frac{p_r^n}{1 - (1 - p_r)^n L_n} \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})}\right] \tag{39}$$

Moments:

(i) For the general distribution of ISIs, following Eq. (39), the mean quantal content $\langle b \rangle$ is given by

$$\langle b \rangle = \sum_{b=0}^{M} b Q^{ss}(b)$$

$$= \sum_{n=0}^{M} \left[\sum_{b=0}^{n} (-1)^b b \binom{n}{b} \right] (-1)^n \binom{M}{n} \frac{p_r^n}{1 - (1 - p_r)^n L_n}$$

$$\times \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})} \right] \tag{40}$$

Note that

$$\sum_{b=0}^{n} (-1)^b b \binom{n}{b} = x \frac{\partial}{\partial x} \left[\sum_{b=0}^{n} \binom{n}{b} x^b \right] \Bigg|_{x=-1}$$

$$= x \frac{\partial}{\partial x} (1 + x)^n \Bigg|_{x=-1} = xn(1 + x)^{n-1} \Bigg|_{x=-1} = -\delta_{n,1} \tag{41}$$

Substituting Eq. (41) in Eq. (40), we have

$$\langle b \rangle = - \sum_{n=0}^{M} \delta_{n,1} (-1)^n \binom{M}{n} \frac{p_r^n}{1 - (1 - p_r)^n L_n} \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})} \right]$$

$$= \frac{M p_r \phi_{1,0}}{1 - (1 - p_r) L_1} \tag{42}$$

(ii) Similarly, the second moment $\langle b^2 \rangle$ is given as:

$$\langle b^2 \rangle = \sum_{b=0}^{M} b^2 Q^{ss}(b)$$

$$= \sum_{n=0}^{M} \left[\sum_{b=0}^{n} (-1)^b b^2 \binom{n}{b} \right] (-1)^n \binom{M}{n} \frac{p_r^n}{1 - (1 - p_r)^n L_n}$$

$$\times \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1 - p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1 - p_r)^{m_i} L_{m_i})} \right] \tag{43}$$

Where,

$$\sum_{b=0}^{n} (-1)^b b^2 \binom{n}{b} = x^2 \frac{\partial^2}{\partial x^2} \left[\sum_{b=0}^{n} \binom{n}{b} x^b \right] \Bigg|_{x=-1} + x \frac{\partial}{\partial x} \left[\sum_{b=0}^{n} \binom{n}{b} x^b \right] \Bigg|_{x=-1}$$

$$=x^2 n(n-1)(1+x)^{n-2}\Big|_{x=-1} - \delta_{n,1}$$

$$=2\delta_{n,2} - \delta_{n,1} \qquad (44)$$

Substituting Eq. (44) in Eq. (43), we have

$$\langle b^2 \rangle = \sum_{n=0}^{M} [2\delta_{n,2} - \delta_{n,1}] \, (-1)^n \binom{M}{n} \frac{p_r^n}{1 - (1-p_r)^n L_n}$$

$$\times \left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}} \frac{(1-p_r)^{\sum_i m_i} \phi_{n,m_z} \phi_{m_z,m_{z-1}} \cdots \phi_{m_1,0}}{\prod_i (1 - (1-p_r)^{m_i} L_{m_i})} \right]$$

$$= \frac{M(M-1)p_r^2}{1 - (1-p_r)^2 L_2} \left[\phi_{2,0} + \frac{(1-p_r)\phi_{2,1}\phi_{1,0}}{1 - (1-p_r)L_1} \right] + \frac{Mp_r\phi_{1,0}}{1 - (1-p_r)L_1} \qquad (45)$$

Using Eqs. (42) and (45), CV^2 can be easily obtained as

$$CV^2 = \frac{\langle b^2 \rangle}{\langle b \rangle^2} - 1$$

$$= \left(\frac{1 - (1-p_r)L_1}{Mp_r\phi_{1,0}} \right)^2 \frac{M(M-1)p_r^2}{1 - (1-p_r)^2 L_2} \left[\phi_{2,0} + \frac{(1-p_r)\phi_{2,1}\phi_{1,0}}{1 - (1-p_r)L_1} \right]$$

$$+ \frac{1 - (1-p_r)L_1}{Mp_r\phi_{1,0}} - 1$$

Note that $1 - L_1 = \phi_{1,0}$. So, we have

$$CV^2 = \left(\frac{\phi_{1,0} + p_r L_1}{Mp_r\phi_{1,0}} \right)^2 \frac{M(M-1)p_r^2}{1 - (1-p_r)^2 L_2} \left[\phi_{2,0} + \frac{(1-p_r)\phi_{2,1}\phi_{1,0}}{\phi_{1,0} + p_r L_1} \right]$$

$$+ \frac{\phi_{1,0} + p_r L_1}{Mp_r\phi_{1,0}} - 1$$

$$= \frac{\phi_{1,0} + p_r L_1}{M\phi_{1,0}} \left[\frac{(M-1)}{1 - (1-p_r)^2 L_2} \left[\phi_{2,0}\left(1 + \frac{p_r L_1}{\phi_{1,0}}\right) + (1-p_r)\phi_{2,1} \right] + \frac{1}{p_r} \right] - 1$$

$$(46)$$

5 The distribution $Q^{ss}(b)$ for two special cases of inter AP interval distribution $g(t_s)$

(a) The deterministic case – APs at a fixed frequency $f = 1/T$

Assuming the APs arrive after fixed time duration T, the corresponding distribution $g(t_s) = \delta(t_s - T)$. This is most often the pattern of input spike train used by experimentalists to excite neuronal response at a fixed frequency $f = 1/T$. We may calculate using Eq. (18),

$$\psi_{n,m} = \int_0^\infty dt_s \delta(t_s - T) e^{-nkt_s}(e^{kt_s} - 1)^{n-m} = e^{-nkT}(e^{kT} - 1)^{n-m},$$

$$L_n = \psi_{n,n} = e^{-nkT},$$

$$\psi_{n,m_z}\psi_{m_z,m_{z-1}} \cdots \psi_{m_1,0} = e^{-nkT}(e^{kT} - 1)^n e^{-kT\sum_{i=1}^z m_i} = (1 - e^{-kT})^n e^{-kT\sum_{i=1}^z m_i} \tag{47}$$

Substituting these expressions in the Eq. (39), and recalling $\phi_{n,m} = \binom{n}{m}\psi_{n,m}$, we have

$$Q^{ss}(b) = \sum_{n=0}^\infty \binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1 - (1-p_r)^n L_n}\left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}}\frac{(1-p_r)^{\sum_i m_i}\phi_{n,m_z}\phi_{m_z,m_{z-1}}\cdots\phi_{m_1,0}}{\prod_i(1 - (1-p_r)^{m_i}L_{m_i})}\right]$$

$$= \sum_{n=0}^\infty \binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1 - (e^{-kT}(1-p_r))^n}(1 - e^{-kT})^n$$

$$\left[1 + \sum_{\{S_{(n-1)}\}}\binom{n}{m_z}\binom{m_z}{m_{z-1}}\cdots\binom{m_2}{m_1}\prod_i\frac{(e^{-kT}(1-p_r))^{m_i}}{(1 - (e^{-kT}(1-p_r))^{m_i})}\right] \tag{48}$$

It is important to remind that $\{S_{(n-1)}\}$ is the set of all the subsets of the set $\{1, 2, \cdots, n-1\}$. To proceed further, we need to use the following identity (for

143 $|x| < 1$) – for proof see Section 6:

$$1 + \sum_{\{S_{(n-1)}\}} \binom{n}{m_z}\binom{m_z}{m_{z-1}}\cdots\binom{m_2}{m_1}\prod_i \frac{x^{m_i}}{(1-x^{m_i})} = \frac{1-x^n}{(1-x)^n} \tag{49}$$

Using the identity (Eq. (49)) in Eq. (48), the expression simplifies to give,

$$\begin{aligned}
Q^{ss}(b) &= \sum_{n=0}^{\infty}\binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1-(e^{-kT}(1-p_r))^n}(1-e^{-kT})^n\frac{1-(e^{-kT}(1-p_r))^n}{(1-e^{-kT}(1-p_r))^n} \\
&= \sum_{n=0}^{\infty}\binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{(1-e^{-kT}(1-p_r))^n}(1-e^{-kT})^n \\
&= \sum_{n=0}^{\infty}\binom{n}{b}(-1)^b\binom{M}{n}\left(\frac{-p_r(1-e^{-kT})}{1-e^{-kT}(1-p_r)}\right)^n
\end{aligned}$$

144 Further using the identity, $\sum_{n=b}^{M} x^n \binom{n}{b}\binom{M}{n} = \binom{M}{b}x^b(x+1)^{M-b}$ we finally get

$$\boxed{Q^{ss}(b) = \binom{M}{b}\left(\frac{(1-e^{-kT})p_r}{1-(1-p_r)e^{-kT}}\right)^b\left(1-\frac{(1-e^{-kT})p_r}{1-(1-p_r)e^{-kT}}\right)^{M-b} = B(M, p_{rb}, b)} \tag{50}$$

145 Hence, the steady state quantal content distribution for the deterministic $g(t_s)$
146 is still a binomial distribution, but with the parameters M (maximal docked vesicle
147 number) and an effective release probability p_{rb} (different from p_r) given by:

$$p_{rb} = \frac{(1-e^{-kT})p_r}{1-(1-p_r)e^{-kT}} \tag{51}$$

148 In order to compare the exact distribution (Eq. (50)) with the approximate
149 binomial distribution $B(\langle n_-\rangle, p_r, b)$ where $\langle n_-\rangle$ is the mean number of docked vesicles
150 just before the AP spikes, we need to calculate an expression for $\langle n_-\rangle$. It is easy to
151 calculate $\langle n_-\rangle$ from the generating function $F_-(q)$ as $\langle n_-\rangle = \frac{\partial F_-(q)}{\partial q}\Big|_{q=1}$.
152 Taking derivative of Eq. (35) and setting $q = 1$, we get

$$\frac{\partial F_-(q)}{\partial q}\Big|_{q=1} = \frac{\partial F_+(q)}{\partial q}\Big|_{q=1}\frac{1}{(1-p_r)} \tag{52}$$

If we take the derivative of Eq. (28) with respect to q at $q = 1$, only $n = 1$ term in the summation survives. Eq. (52) will become

$$\left.\frac{\partial F_-(q)}{\partial q}\right|_{q=1} = \frac{M(1-p_r)}{1-(1-p_r)L_1}\phi_{1,0}\frac{1}{(1-p_r)}$$

Substituting the forms of L_1 and $\phi_{1,0}$ from Eq. (47), we get

$$\left.\frac{\partial F_-(q)}{\partial q}\right|_{q=1} = \frac{M(1-p_r)}{1-(1-p_r)e^{-kT}}(1-e^{-kT})\frac{1}{(1-p_r)}$$

Hence,

$$\boxed{\langle n_-\rangle = \frac{M(1-e^{-kT})}{1-(1-p_r)e^{-kT}} = M\frac{p_{rb}}{p_r}} \tag{53}$$

In the above we have used the expression of p_{rb} from Eq. (51). The Eq. (53) implies that the mean $\langle b \rangle$ of the exact binomial and the approximate binomial are equal, i.e. $Mp_{rb} = \langle n_-\rangle p_r$. But the variance of the exact distribution is $Mp_{rb}(1-p_{rb})$, different from that of the approximate binomial $\langle n_-\rangle p_r(1-p_r)$.

(b) Poisson distributed AP arrival times

APs may arrive at intervals which may be noisy. A simple way to incorporate stochasticity in the input spike train is to assume that they arrive at time intervals which are exponentially distributed with $g(t_s) = f_0 e^{-f_0 t_s}$. Note that the mean as well as characteristic spike interval is $1/f_0$. In this case using Eq. (18),

$$\begin{aligned}
\psi_{n,m} &= \int_0^\infty dt_s g(t_s)e^{-mkt_s}(1-e^{-kt_s})^{n-m} \\
&= \int_0^\infty dt_s f_0 e^{-(mk+f_0)t_s}(1-e^{-kt_s})^{n-m} \\
&= \frac{f_0}{k}\int_0^1 dz\, z^{\frac{mk+f_0}{k}-1}(1-z)^{n-m} \\
&= \frac{f_0}{k}B\left(m+\frac{f_0}{k}, n-m+1\right) = \frac{f_0}{k}\frac{\Gamma(-m+n+1)\Gamma\left(\frac{f_0}{k}+m\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)},
\end{aligned} \tag{54}$$

 where we have used Beta function $B(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$, and then

$$L_n = \psi_{n,n} = \frac{f_0}{k} \frac{\Gamma(1)\Gamma\left(\frac{f_0}{k} + n\right)}{\Gamma\left(\frac{f_0}{k} + n + 1\right)} = \frac{f_0}{f_0 + nk}. \tag{55}$$

Moreover using Eq. (54), we have

$$\phi_{n,m} = \binom{n}{m}\psi_{n,m} = \binom{n}{m}\frac{f_0}{k}\frac{\Gamma(-m+n+1)\Gamma\left(\frac{f_0}{k}+m\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)} = \frac{f_0}{k}\frac{\Gamma(n+1)\Gamma\left(\frac{f_0}{k}+m\right)}{\Gamma(m+1)\Gamma\left(\frac{f_0}{k}+n+1\right)} \tag{56}$$

$$\Rightarrow \phi_{n,m}\phi_{m,l} = \frac{f_0^2}{k^2}\frac{\Gamma(n+1)\Gamma\left(\frac{f_0}{k}+m\right)}{\Gamma(m+1)\Gamma\left(\frac{f_0}{k}+n+1\right)}\frac{\Gamma(m+1)\Gamma\left(\frac{f_0}{k}+l\right)}{\Gamma(l+1)\Gamma\left(\frac{f_0}{k}+m+1\right)}$$

$$= \frac{f_0^2}{k^2}\frac{k\Gamma(n+1)\Gamma\left(\frac{f_0}{k}+l\right)}{\Gamma(l+1)(f_0+km)\Gamma\left(\frac{f_0}{k}+n+1\right)} \tag{57}$$

 The above Eq. (57) may be extended to get the following product:

$$\phi_{n,m_z}\phi_{m_z,m_{z-1}}\cdots\phi_{m_1,0} = \frac{f_0^{z+1}}{k^{z+1}}\frac{n!\Gamma\left(\frac{f_0}{k}\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)}\prod_i\frac{k}{f_0+km_i} \tag{58}$$

Substituting Eq. (58) into Eq. (39), we have

$$Q^{ss}(b) = \sum_{n=0}^{\infty}\binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1-(1-p_r)^nL_n}\left[\phi_{n,0} + \sum_{\{S_{(n-1)}\}}\frac{(1-p_r)^{\sum_i m_i}\frac{f_0^{z+1}}{k^{z+1}}\frac{n!\Gamma\left(\frac{f_0}{k}\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)}\prod_i\frac{k}{f_0+km_i}}{\prod_i(1-(1-p_r)^{m_i}L_{m_i})}\right]$$

$$= \sum_{n=0}^{\infty}\binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1-(1-p_r)^nL_n}\frac{n!\Gamma\left(\frac{f_0}{k}\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)}\frac{f_0}{k}\left[1 + \sum_{\{S_{(n-1)}\}}\frac{(1-p_r)^{\sum_i m_i}\prod_i\frac{f_0}{f_0+km_i}}{\prod_i(1-(1-p_r)^{m_i}L_{m_i})}\right]$$

 From Eq. (55), substituting $\frac{f_0}{f_0+km_i} = L_{m_i}$, we have

$$Q^{ss}(b) = \sum_{n=0}^{\infty}\binom{n}{b}(-1)^{n-b}\binom{M}{n}\frac{p_r^n}{1-(1-p_r)^nL_n}\frac{n!\Gamma\left(\frac{f_0}{k}+1\right)}{\Gamma\left(\frac{f_0}{k}+n+1\right)}\left[1 + \sum_{\{S_{(n-1)}\}}\prod_i\frac{(1-p_r)^{m_i}L_{m_i}}{(1-(1-p_r)^{m_i}L_{m_i})}\right] \tag{59}$$

 We may simplify the expression above by using the following identity valid for any

167 arbitrary function f – for proof see Section 7:

$$1 + \sum_{\{S_{(n-1)}\}} \prod_i \frac{f(m_i)}{1 - f(m_i)} = \frac{1}{\prod_{i=1}^{n-1}(1 - f(i))} \tag{60}$$

168 Using Eq. (60) in Eq. (59), we get

$$Q^{ss} = \sum_{n=0}^{\infty} \binom{n}{b}(-1)^{n-b}\binom{M}{n} \frac{p_r^n n! \Gamma\left(\frac{f_0}{k} + 1\right)}{(1 - (1-p_r)^n L_n)\Gamma\left(\frac{f_0}{k} + n + 1\right)} \frac{1}{\prod_{i=1}^{n-1}(1 - (1-p_r)^i L_i)}$$

Substituting back $L_i = \frac{f_0}{f_0 + ki}$, we have

$$Q^{ss} = \sum_{n=0}^{\infty} \binom{n}{b}(-1)^{n-b}\binom{M}{n} \frac{p_r^n n! \Gamma\left(\frac{f_0}{k} + 1\right)}{\Gamma\left(\frac{f_0}{k} + n + 1\right)} \frac{1}{\prod_{i=1}^{n} \frac{(f_0 + ki - (1-p_r)^i f_0)}{f_0 + ki}}$$

$$\boxed{Q^{ss}(b) = \sum_{n=0}^{\infty}(-1)^{n-b}\binom{n}{b}\binom{M}{n} \frac{n!(kp_r)^n}{\prod_{i=1}^{n}(f_0 + ki - (1-p_r)^i f_0)}} \tag{61}$$

169 Thus the steady state quantal content distribution for the spike intervals drawn
170 from a Poisson processes is not a Binomial, but instead the above distribution in Eq.
171 (61).

172 **_Moments:_**

(i) For the exponentially distributed ISIs, the mean quantal content $\langle b \rangle$ can be ob-
tained by substituting $\phi_{1,0} = \frac{k}{f_0 + k}$ and $L_1 = \frac{f_0}{f_0 + k}$ from Eqs. (56) and (55) respec-
tively, into the general $\langle b \rangle$ expression (Eq. (42)) as follows

$$\langle b \rangle = \frac{M p_r \frac{k}{f_0 + k}}{1 - (1 - p_r)\frac{f_0}{f_0 + k}}$$

$$= \frac{M k p_r}{k + f_0 p_r} \tag{62}$$

(ii) Similarly, the CV^2 can be obtained by substituting the expressions: $\phi_{1,0} = \frac{k}{f_0+k}$, $\phi_{2,0} = \frac{2k^2}{(f_0+k)(f_0+2k)}$, $\phi_{2,1} = \frac{2f_0k}{(f_0+k)(f_0+2k)}$, $L_1 = \frac{f_0}{f_0+k}$, and $L_2 = \frac{f_0}{f_0+2k}$ from Eqs. (56) and (55), into the general CV^2 expression (Eq. (46)) as follows

$$
\begin{aligned}
CV^2 =& \frac{k+f_0p_r}{Mk}\left[\frac{(M-1)(f_0+2k)}{f_0+2k-(1-p_r)^2f_0}\left(\frac{(k+f_0p_r)2k^2+2(1-p_r)f_0k^2}{k\,(f_0+k)\,(f_0+2k)}\right)+\frac{1}{p_r}\right]-1 \\
=& \frac{k+f_0p_r}{Mk}\left[\frac{(M-1)(f_0+2k)}{f_0+2k-(1-p_r)^2f_0}\left(\frac{2k^2(f_0+k)}{k\,(f_0+k)\,(f_0+2k)}\right)+\frac{1}{p_r}\right]-1 \\
=& \frac{1}{M}\left(\frac{2(M-1)\,(k+f_0p_r)}{2k+f_0p_r\,(2-p_r)}+\frac{f_0}{k}-M+\frac{1}{p_r}\right)
\end{aligned}
\tag{63}
$$

6 Proof of the identity in Eq. (49):

The identity can be proved using the principle of strong mathematical induction as follows.

We first verify the identity for $n=1$ and 2 explicitly. For $n=1$, LHS of Eq. (49) will be simply equal to 1 as the summation term does not contribute anything. The RHS is also equal to 1, so the identity is true for $n=1$. For $n=2$, $\{S_{(1)}\} = \{\{1\}\}$. So, the LHS of Eq. (49) will be equal to $1+\binom{2}{1}\frac{x}{1-x} = \frac{1+x}{1-x} = \frac{1-x^2}{(1-x)^2} = RHS$. Hence, we have verified the identity for $n=2$ as well.

Let us assume that the identity is true for all integer values of $n \in [1,k]$. We would then show below that it is true for $n=k+1$. For $n=k+1$ we have:

$$
\begin{aligned}
LHS =& 1 + \sum_{\{S_{(k)}\}}\binom{k+1}{m_z}\binom{m_z}{m_{z-1}}\cdots\binom{m_2}{m_1}\prod_i\frac{x^{m_i}}{(1-x^{m_i})} \\
=& 1 + \binom{k+1}{1}\frac{x}{1-x} + \sum_{\substack{\{S_{(k)}\}\\ m_z\neq 1}}\binom{k+1}{m_z}\binom{m_z}{m_{z-1}}\cdots\binom{m_2}{m_1}\prod_i\frac{x^{m_i}}{(1-x^{m_i})}
\end{aligned}
\tag{64}
$$

where in the last step, we have separated out the contribution of $m_z = 1$ in the summation. The crucial step in our proof comes from the idea that the subsets in the set $\{S_{(k)}\}$ (with $m_z \neq 1$) can be clubbed into different groups, such that the largest element in all the subsets in a particular group are equal. There will be $k-1$ such groups as the maximum element m_z can range from 2 to k. With the summing

over all such groups explicitly shown, Eq. (64) becomes

$$LHS = 1 + \binom{k+1}{1}\frac{x}{1-x} + \sum_{m_z=2}^{k}\binom{k+1}{m_z}\frac{x^{m_z}}{1-x^{m_z}}\left[\sum_{\{S_{(m_z-1)}\}}\binom{m_z}{m_{z-1}}\cdots\binom{m_2}{m_1}\prod_i\frac{x^{m_i}}{(1-x^{m_i})}+1\right]$$

(65)

Note that 1 is added manually inside the square bracket to take care of the subsets with only one element m_z, as these subsets are not taken into account by the summation term over the subsets $\{S_{(m_z-1)}\}$. The expression inside the square bracket is equal to the LHS of Eq. (49) with $n = m_z$. Note that as positive integer $m_z \leq k$, by assumption the identity Eq. (49) is true. So, substituting the form of RHS from Eq. (49) into the square bracket of Eq. (65), we have

$$
\begin{aligned}
LHS =& 1 + \binom{k+1}{1}\frac{x}{1-x} + \sum_{m_z=2}^{k}\binom{k+1}{m_z}\frac{x^{m_z}}{1-x^{m_z}}\left[\frac{1-x^{m_z}}{(1-x)^{m_z}}\right]\\
=& 1 + \binom{k+1}{1}\frac{x}{1-x} + \sum_{m_z=2}^{k}\binom{k+1}{m_z}\left(\frac{x}{1-x}\right)^{m_z}\\
=& \sum_{m_z=0}^{k}\binom{k+1}{m_z}\left(\frac{x}{1-x}\right)^{m_z}\\
=& \sum_{m_z=0}^{k+1}\binom{k+1}{m_z}\left(\frac{x}{1-x}\right)^{m_z} - \left(\frac{x}{1-x}\right)^{k+1}\\
=& \left(\frac{1}{1-x}\right)^{k+1} - \left(\frac{x}{1-x}\right)^{k+1}\\
=& \frac{1-x^{k+1}}{(1-x)^{k+1}} = RHS
\end{aligned}
$$

(66)

Hence, we have shown above that the identity is true for $n = k+1$. Since integer $k \geq 2$ is arbitrary, we have proved the identity in Eq. (49) using the method of strong induction.

7 Proof of the identity in Eq. (60):

Here is the proof of the identity using the method of mathematical induction:

For $n = 2$, $\{S\}$, the set of all the subsets of $\{1\}$, is simply $\{\{1\}\}$. Hence, the LHS

of Eq. (60) will be equal to

$$LHS = 1 + \prod_{\{1\}} \frac{f(m_i)}{1 - f(m_i)} = 1 + \frac{f(1)}{1 - f(1)} = \frac{1}{1 - f(1)} = RHS \qquad (67)$$

Let us assume that the identity is true for an arbitrary value of $n = k$. For, $n = k+1$, the set $\{S\}$ of all the subsets of $\{1, 2, 3, \cdots k\}$ can be divided into three disjoint subsets:

(a) the set of all the subsets of $\{1, 2, 3, \cdots k - 1\}$.

(b) the set of all the subsets in (a) with an extra element k added to each of the subsets.

(c) $\{\{k\}\}$

Clearly, the contribution of (a) in the summation term in the LHS of Eq. (60) will be equal to that for $n = k$, that is equal to $1/\prod_{i=1}^{k-1}(1 - f(i)) - 1$. On the other hand, the contribution of (b) will be equal to that of (a) multiplied by an additional factor $\frac{f(k)}{1-f(k)}$. The contribution from (c) will be simply $\frac{f(k)}{1-f(k)}$. Adding all the three contributions,

$$
\begin{aligned}
LHS =& 1 + \left(\frac{1}{\prod_{i=1}^{k-1}(1 - f(i))} - 1 \right) + \left(\frac{1}{\prod_{i=1}^{k-1}(1 - f(i))} - 1 \right) \frac{f(k)}{1 - f(k)} + \frac{f(k)}{1 - f(k)} \\
=& \frac{1}{\prod_{i=1}^{k-1}(1 - f(i))} \left(1 + \frac{f(k)}{1 - f(k)} \right) \\
=& \frac{1}{\prod_{i=1}^{k}(1 - f(i))} = RHS
\end{aligned}
$$

Hence, the identity is true for $n = k+1$. Thus from the principle of mathematical induction, the identity is proved for any integer $n > 1$.

8   Saturation of time dependent p_r and k, following calcium level saturation, in steady state

(a)   (b)

Figure 1: The time-dependence of p_{rt} and k_t are shown for the parameter values given in Table 2. of Ref. [1], for parallel fiber. The initial and steady state values are marked with red lines, and the corresponding values are mentioned.

The vesicle release probability p_{rt} and the rate of docking of vesicles k_t are dependent on the time t, through their dependence on time-varying Ca^{2+} level inside the presynaptic terminal. There are different models available in the literature to describe this period of plasticity. For the purpose of demonstrating the approach to steady state, we use one such model studied in the paper by Dittman and colleagues [1]. Two different types of calcium binding targets are considered, namely F and D. The p_{rt} is related to the number CaX_F of Ca^{2+}-bound-F molecules as $p_{rt} = p_{r0} + \frac{1-p_{r0}}{\frac{K_F}{CaX_F(t)}+1}$. On the other hand, k_t is related to the number CaX_D of Ca^{2+}-bound-D sites as $k_t = k_0 + \frac{k_{\max}-k_0}{\frac{K_D}{ca(t)}+1}$. Given an AP train, with m^{th} AP arriving at $t = t_m$, the dynamics of CaX_F and CaX_D are given by

$$\frac{dCaX_F}{dt} = -\frac{CaX_F}{\tau_F} + \Delta_F \sum_m \delta(t - t_m) \tag{68}$$

and,

$$\frac{dCaX_D}{dt} = -\frac{CaX_D}{\tau_D} + \Delta_D \sum_m \delta(t - t_m) \tag{69}$$

²²¹ respectively. CaX_F (or CaX_D) jumps by Δ_F (or Δ_D) after each AP, and falls

²²² exponentially between two APs, with a decay constant τ_F (or τ_D). In Fig. 1, for the

²²³ parameter values given in Ref. [1] for parallel fiber, we have plotted the p_{r_t} and k_t as

²²⁴ a function of t. Note that the initial values at $t \to 0$ are p_{r_0} and k_0 respectively. We

²²⁵ see that $p_{r_t} \to p_r$ and $k_t \to k$ in the long time limit. This steady state limit is the

²²⁶ focus of the entire analysis described in this paper, in which the release probability

²²⁷ and replenishment rate has attained the steady values p_r and k respectively.

9   Details of the experiments performed

²²⁹ We performed electrophysiological experiments in acute brainstem slices of juvenile

²³⁰ mice (postnatal day 11) at MNTB-LSO synapses. These fast auditory synapses

²³¹ transfer inhibitory glycinergic information and can maintain reliable neurotransmis-

²³² sion even during prolonged high-frequency stimulation [2, 3]. We refer the reader

²³³ to the Methods section of Ref. [3] for details on the slice preparation and details of

²³⁴ whole-cell patch-clamp recordings. The stimulation protocol consisted of repetitive

²³⁵ tetanic AP bursts (100 Hz, 1 s) that were given 20 times. A 30-s rest period was

²³⁶ introduced between repeats to allow for recovery from short-term depression. The

²³⁷ number of fused SVs was determined as the ratio of the peak amplitude of evoked in-

²³⁸ hibitory postsynaptic currents (eIPSC) to the current produced by a single quantum

²³⁹ (q) [pA/(pA/SV)]. Detailed information is available in Refs. [3, 4].

10   Kinetic Monte Carlo (KMC) code

241 The code used to simulate the distribution of quantal content for constant ISIs as
242 well as Poisson train is as follows.

```python
import math as mp
import numpy as np
```

```python
burst=[]

for col in range (0,100,1):
    burstt=[]
    time=[]
    M=50
    f=100
    pr =0.5
    k = 1.0
    g=1
    while g<2:
        t=0
        n=M
        nstate=[M]
        j=[]
        x=1
        #####################################################################
        #Use this part for constant ISIs

        dlist=[0]
        for i in range (100000):
            dlist.append(max(dlist)+(1/f))

        #####################################################################
        #Use this part for exponentially distributed ISIs

        dlist=[0]
        for i in range (100000):
            dlist.append(max(dlist)+(1/f)*mp.log(1/np.random.uniform(0,1)))

        #####################################################################
        while x<5000:
            p=np.random.uniform(0,1)
            for i in range (0,100,1):
                if dlist[i]<=t<=dlist[i+1] and \
                t+(1/(k*(M-n+0.000001)))*mp.log(1/p)<=dlist[i+1]:
                    n=n+1
                    t=t+(1/(k*(M-n+1+0.000001)))*mp.log(1/p)
                    time.append(t)
                    break
                elif dlist[i]<=t<=dlist[i+1] and \
                t+(1/(k*(M-n+0.000001)))*mp.log(1/p)>=dlist[i+1]:
                    time.append(dlist[i+1])
                    zz=np.random.binomial(n, pr, size=None)
                    burstt.append(zz)
                    n=n-zz
                    t=dlist[i+1]+0.0001
                    time.append(t)
                    break
            x=x+1
        g=g+1
    burst.append(burstt)
```

References

[1] J. S. Dittman, A. C. Kreitzer, and W. G. Regehr, "Interplay between facilitation, depression, and residual calcium at three presynaptic terminals," *Journal of Neuroscience*, vol. 20, no. 4, pp. 1374–1385, 2000.

[2] S. E. Brill, A. Maraslioglu, C. Kurz, F. Kramer, M. F. Fuhr, A. Singh, and E. Friauf, "Glycinergic transmission in the presence and absence of functional glyt2: Lessons from the auditory brainstem," *Frontiers in Synaptic Neuroscience*, vol. 12, p. 560008, 2020.

[3] N. I. C. Müller, I. Paulussen, L. N. Hofmann, J. O. Fisch, A. Singh, and E. Friauf, "Development of synaptic fidelity and action potential robustness at an inhibitory sound localization circuit: effects of otoferlin-related deafness," *The Journal of Physioloy*, vol. 600, no. 10, pp. 2461–2497, 2022.

[4] N. I. C. Müller, M. Sonntag, A. Maraslioglu, J. J. Hirtz, and E. Friauf, "Topographic map refinement and synaptic strengthening of a sound localization circuit require spontaneous peripheral activity," *The Journal of Physiology*, vol. 597, no. 22, pp. 5469–5493, 2019.