Neuronal TIMP2 regulates hippocampus-dependent plasticity and extracellular matrix complexity

Ana Catarina Ferreira[1,2,3,4], Brittany M. Hemmer[1,2,3,4,5], Sarah M. Philippi[1,2,3,4,5], Hanxiao Liu[1,2,3,4], Jeffrey D. Zhu[1,2,3,4], Tatyana Kareva[1,2,3,4], Tim Ahfeldt[1,2,3,4], Merina Varghese[1], Patrick R. Hof[1,2], Joseph M. Castellano[1,2,3,4,†]

[1]Nash Family Department of Neuroscience, Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY, United States

[2]Ronald M. Loeb Center for Alzheimer's Disease, Icahn School of Medicine at Mount Sinai, New York, NY, United States

[3]Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, United States

[4]Black Family Stem Cell Institute, Icahn School of Medicine at Mount Sinai, New York, NY, United States

[5]Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, United States

[†]To whom all correspondence should be addressed

Email: joseph.castellano@mssm.edu

Joseph M. Castellano, Ph.D.

Icahn School of Medicine at Mount Sinai

Nash Family Department of Neuroscience, Department of Neurology

Friedman Brain Institute and Black Family Stem Cell Institute

Ronald M. Loeb Center for Alzheimer's disease

1425 Madison Ave., Box 1065

New York, NY 10029

Keywords: TIMP2, neurogenesis, synaptic plasticity, memory, extracellular matrix

Abstract

The functional output of the hippocampus, a brain region subserving memory processes, depends

on highly orchestrated cellular and molecular processes that regulate synaptic plasticity

throughout life. However, little is known regarding the structural requirements of such plasticity

and molecular processes involved in this regulation. Specific molecules, including tissue inhibitor

of metalloproteinases-2 (TIMP2) have been implicated in serving a pro-plasticity role within the

hippocampus, a role that decreases with brain aging. Here, we report that TIMP2 is expressed at

high levels in the hippocampus by neurons and its loss in TIMP2 KO mice drives changes in

cellular programs related to adult neurogenesis and dendritic spine turnover with corresponding

impairments in hippocampus-dependent memory performance. We find that TIMP2 regulates the

accumulation of extracellular matrix (ECM) around synapses in the hippocampus with

concomitant hindrance in the migration of newborn neurons through a denser ECM network.

Deleting the neuronal pool of TIMP2 using a conditional TIMP2 KO mouse reveals that neuronal

TIMP2 regulates adult neurogenesis and hippocampus-dependent memory, as well as

accumulation of ECM. Our results define a mechanism whereby hippocampus-dependent

function is regulated by TIMP2 and its interactions with the ECM to regulate diverse processes

associated with synaptic plasticity.

Introduction

A variety of cellular, molecular, and functional changes have now been documented in the brain

as aging progresses, leaving it susceptible to diseases like Alzheimer's disease. Given that aging

serves as the major risk factor for many neurological disorders, aging processes have increasingly

been considered to be valid therapeutic targets, especially as the aged population continues to

increase. Growing evidence in animal models indicates that the aged brain can be rejuvenated

through manipulation of the systemic environment, including through young plasma transfer or

sharing of young blood through heterochronic parabiosis (Castellano et al., 2017; Katsimpardi et

al., 2014; Villeda et al., 2011; Villeda et al., 2014). Aged mice exposed to young blood exhibit

improved synaptic plasticity and cognitive performance (Villeda et al., 2014). We and others have

begun to identify factors present in young blood that confer brain rejuvenation phenotypes in aged

mice (Castellano et al., 2017; Katsimpardi et al., 2014). Among those, we showed that tissue

inhibitor of metalloproteinases-2 (TIMP2), a protein elevated early in life, revitalizes function of

the aged mouse hippocampus (Castellano et al., 2017). Despite demonstrating that this factor is

critical for young plasma-mediated memory improvements in aged mice (Castellano et al., 2017),

little is known of the cellular and molecular details of how TIMP2 regulates function within the

hippocampus.

TIMP2 is a secreted protein of approximately 21-kDa in size primarily known for its role in

regulating peripheral extracellular matrix (ECM) remodeling (Brew and Nagase, 2010) through its

unique dual roles in regulating the activation of latent matrix metalloproteinases (MMPs), including

MMP2, to proteolytically active forms, and the inhibition of MMP activity (Caterina et al., 2000;

Wang et al., 2000). Through these roles, TIMP2 has been shown to play a key role in the

maintenance of ECM network structure in many tissues with myriad disease implications,

including heart failure (Kandalam et al., 2011), cancer (Kaczorowska et al., 2020), ischemic stroke

(Guo et al., 2018), among others. While early reports described expression of TIMP2 from

development into adulthood (Fager and Jaworski, 2000), and its role in regulating cortical

neuronal differentiation (Perez-Martinez and Jaworski, 2005), and subtle motor phenotypes

(Jaworski et al., 2006), few studies have examined the biological role of TIMP2 in the adult brain

and how it modulates function of the hippocampus. While we recently demonstrated that TIMP2

is critical in revitalizing overall hippocampus-dependent cognitive function in aged mice

(Castellano et al., 2017), the biological role of TIMP2 expressed within the hippocampus, where

it is highly expressed, has not been studied. TIMP2 is part of a small family of tissue

metalloproteinase inhibitors that include TIMP1, TIMP3, and TIMP4, all of which perform broad

MMP targeting functions but in diverse contexts in the brain, including in myelin repair (Nicaise et

al., 2019), neurovascular regulation (Dewing et al., 2019), or in regulation of the tumor

microenvironment (Solga et al., 2019). Interestingly, TIMP2 shares only ~40-50% sequence

homology with other TIMP family members and is particularly expressed in the brain and enriched

specifically in hippocampus relative to other TIMPs. Since levels of TIMP2 decline in the

hippocampus with age (Castellano et al., 2017), understanding the function of this local source

will be important to understand its normal function and position it for putative therapies age-related

brain disorders. Moreover, we sought to understand the extent to which ECM regulation is related

to disparate processes of plasticity to regulate complex behavioral outputs.

Here we define a mechanism by which TIMP2 regulates ECM components in the

hippocampus in a manner that regulates processes linked to plasticity and memory. We find that

TIMP2 is predominantly expressed by neurons in the hippocampus and is present at high levels

in the brain extracellular space adjacent to these cells. Deletion of TIMP2 induces transcriptomic

changes in the hippocampus that are consistent with processes related to synaptic plasticity and

adult neurogenesis. We find that loss of TIMP2 results in fewer dendritic spines in dentate gyrus

(DG) granule cells, decreased adult DG neurogenesis along several stages, and impaired

memory in various hippocampus-dependent tasks. Loss of TIMP2 also causes an accumulation

of ECM proteins, particularly around synapses within the DG, and impaired migration of

neuroblasts within the neurogenic niche, reflecting dysregulated ECM turnover. Finally, we

developed a mouse model for conditional deletion of TIMP2 from neurons that exhibits the

functional phenotypes observed in TIMP2 KO mice, further arguing for a role of the neuronal pool

of TIMP2 in shaping plasticity-related function of the hippocampus. Our findings indicate

molecular mechanisms through which neuronal TIMP2 remodeling of the ECM promotes synapse

plasticity and is required for memory, with functional implications for neurogenic rejuvenation.

Materials and Methods

Animals

All animal care and procedures were performed in accordance with the National Institutes of

Health Guide for Care and Use of Laboratory Animals and the Icahn School of Medicine at Mount

Sinai Institutional Animal Care and Use Committee. All efforts were made to reduce the number

of animals used and to minimize their suffering and pain. TIMP2 knockout (KO) (Jackson) and

WT female mice or $\text{Syn}^{Cre/+}$; $\text{TIMP2}^{fl/fl}$ and $\text{TIMP2}^{fl/fl}$ mice were used at 2-3 months or 5-6 months

of age unless indicated otherwise. Littermate controls were used for all experiments. Mice were

maintained on a 12h light/dark cycle at constant temperature (23°C) with *ad libitum* access to food

and water.

Generation of TIMP2-floxed mice

To achieve conditional deletion of *Timp2*, a loxP-flanked *Timp2* allele ($\text{TIMP2}^{fl/fl}$) was generated

by the two-cell homologous recombination (2C-HR)-CRISPR/Cas9-based genome editing

method, as previously described (Gu et al., 2018). Briefly, two loxP sites were inserted into the

Timp2 gene flanking exon 2, with homologous arms (HA) on each side. Two point-mutations

(C928->A near the end of the left arm and G1795->T at the 5' junction of the right arm) were

introduced to disrupt the PAM sequence to avoid re-cutting of the recombined allele. Three

nucleotides adjacent to each loxP site were altered to create restriction sites for diagnostic

purposes: XbaI near the upstream loxP, and HindIII near the downstream loxP site. The two-cell

stage embryo microinjection procedure was performed as in (Gu et al. 2018), using Cas9-mSA

mRNA, biotinylated double-stranded PCR product (1016 nts), and pre-assembled ctRNA

prepared accordingly (Gu et al., 2018). F_0 mice were screened by sequencing from TOPO cloned

PCR product, using tail genomic DNA, amplified from the TIMP2 target region, and using the

following primers: FW-AGCGACCGATAAGCAGGAAG, Rev-CACTAGCAGACACCACCACA.

Identified founders were crossed with wild-type (WT) C57BL/6J mice, purchased from The

Jackson Laboratory, to produce heterozygous mice.

To achieve neuron-specific *Timp2* deletion, neuron-specific heterozygous Synapsin (Syn)-

Cre ($Syn^{Cre/+}$) transgenic mice (Zhu et al., 2001) (JAX stock #003966) were crossbred with

$TIMP2^{fl/fl}$ mice to generate $Syn^{Cre/+}$; $TIMP2^{fl/+}$ progeny. The F_1 mice were then crossed to generate

$Syn^{Cre/+}$; $TIMP2^{fl/fl}$ and $TIMP2^{fl/fl}$ littermate controls. Cre expression was maintained in the female

mice for breeding (Zhu et al., 2001). All animals were maintained on a C57Bl/6J background.

Genotyping

Genotyping of $TIMP2^{fl/fl}$ mice was performed using the following primers: *Primer 1*: 5'-

TGACCCCTCTGCTCTAGTCC, *Primer 2:* 5'-GCTCCGTCTCTTCGTCCATC (**Figure 6A**).

Genomic DNA was extracted from the tail using the QIAamp Fast DNA Tissue Kit (Qiagen). PCR

was performed in a total volume of 50 µl at 94°C for 2 min, 94°C for 30 s, 61°C for 30 s, 68°C for

2 min and 15 sec for 10 cycles, then 94°C for 30 s, 56°C for 30 s, 68°C for 2 min and 15 s for 25

cycles, and ending at 72°C for 5 min using Prime STAR (#R050A, TaKaRa). For loxP site

verification based on restriction digest, the PCR product (2.1 kb) was digested with XbaI and

HindIII restriction enzymes (New England Biolabs, Ipswich, MA) at 37°C for 1 hour, followed by

10 min at 65°C for heat inactivation. Digested PCR products were run on an agarose gel and

imaged for the identification of founders: HindIII (Wt: 2133 bp; upstream loxP positive: 829 bp and

1,304 bp); Xba1 (Wt: 2133 bp; downstream loxP positive: 566 bp and 1,567 bp).

In vivo microdialysis

For guide cannula implantation into the hippocampus, mice were anesthetized with 2% isoflurane and hair over the skull was shaved prior to being fixed in a stereotaxic apparatus (Stoelting Co., Wood Dale, IL, USA). After preparing skin with 70% ethanol and povidone-iodine, an anterior-to-posterior incision was made along the midline of the head to expose the skull. Skull position was leveled to within 0.1 mm along the bregma-lambda axis, and at 2.2 mm to the left and right of midline on the lateral axis. Holes were drilled to target the left hippocampus using the stereotaxic coordinates: bregma -3.1 mm, 2.5 mm lateral. A second hole was made diagonally to the first, to serve as an anchoring position with a bone screw. To target caudal hippocampus, AtomosLM Guide Cannula (PEG-12, Eicom, San Diego, CA, USA) was inserted at a 12° angle, 1.2 mm below the dura mater. Dental cement was applied to secure the cannula, and the skin was secured using a surgical adhesive glue. An AtmosLM Dummy Cannula (PED-12, Eicom, San Diego, CA, USA) was inserted into the guide cannula and secured with a plastic cap nut. Animals were then placed in a clean cage on a heating pad and allowed to recover from surgery.

Approximately 12 hours following surgery, mice underwent 1000-kDa *in vivo* microdialysis. The inlet port of AtmosLM Microdialysis Probe (PEP-12-02, Emicon) was connected to a syringe pump (kdScientific) perfusing artificial CSF (aCSF; in mM: 1.3 $CaCl_2$, 1.2 $MgSO_4$, 3 KCl, 0.4 KH_2PO_4, 25 $NaHCO_3$, 122 NaCl, pH 7.35) at a rate of 1.2 µl/min (Ulrich et al., 2013). The outlet of the probe was connected to the peristaltic pump (MAB 20, SciPro), which was calibrated to obtain a pull rate between 1-1.1 µl/min. After the probe was connected to the push-pull mechanism, the mice were briefly anesthetized with isoflurane to allow for the removal of the dummy cannula and insertion of the probe into the hippocampus. The probe was secured with a cap nut, and a plastic collar was loosely placed around the mouse neck. Mice were then moved to a Raturn (Stand-Alone Raturn System, BASi, West Lafayette, IN, USA) and tethered by the collar to avoid tangling microdialysis tubing (FEP tubing 0.65 mm OD x 0.12 mm ID, BASi), while

allowing for free movement of the mouse during the sample collection period. Samples were

collected hourly in a refrigerated fraction collector (MAB 85 Fraction Collector, SciPro) and frozen

after collection at −80°C. The procedure was performed under constant light conditions, and food

and water were provided *ad libitum*.

Bulk RNA-sequencing

RNA-sequencing was performed on hippocampi dissected from WT and TIMP2 KO male and

female mice. Dissected hippocampi were preserved in RNAlater (Invitrogen, Carlsbad, CA)

overnight at 4°C before storing at −80°C until use. RNA was extracted using the RNeasy Mini Kit

(Qiagen), according to the manufacturer's instructions. RNA quality was measured using the

Agilent TapeStation Bioanalyzer (Agilent Technologies, USA), and all samples exhibited RNA

Integrity Number (RIN) > 8. cDNA libraries were prepared with poly(A) selection and sequenced

using Illumina Hiseq (2x150bp paired-end) (Genewiz). At least 25 M clean reads were generated

from each sample. The reads were mapped to the *Mus musculus* GRCm38 reference genome

available on ENSEMBL, using STAR aligner (v.2.5.2b). After extraction of gene hit counts, the

gene hit counts table was used for downstream differential expression analysis, using DESeq2

software. The Wald test was used to generate *P*-values and log2 fold changes. Differentially

expressed genes (DEGs; *P*<0.05*)* were used for Gene Set Enrichment Analysis (GSEA;

https://www.gsea-msigdb.org/gsea/index.jsp). Volcano plot was generated with R v4.1.2.

BrdU injections

For cell proliferation assessment, mice were injected intraperitoneally (i.p.) once with 5-Bromo-2′-

deoxyuridine (BrdU; Sigma) at a concentration of 150 mg/kg dissolved in 0.9% sodium chloride

solution and sacrificed 24 hours later. For cell fate ("survival") analysis, an additional group of

mice was i.p.-injected with BrdU once a day for 5 consecutive days, followed by a chase period

of 28 days before perfusion.

Immunohistochemistry and microscopy

For all immunohistochemistry experiments, mice were anesthetized with a cocktail of ketamine (90 mg/kg) and xylazine (10 mg/kg) and transcardially perfused with ice-cold 0.9% saline. Brains were dissected, and hemibrains isolated for immunohistochemistry were postfixed in 4% paraformaldehyde (PFA) for 48 hours at 4°C before preservation in 30% sucrose (Sigma) solution in phosphate-buffered saline (PBS). Hemibrains were sectioned coronally at 40-µm thickness on a freezing-sliding microtome (SM2010R, Leica), and sections were stored in cryoprotective medium at −20°C.

For immunohistochemistry, free-floating brain sections were permeabilized with Tris-Buffer Saline (TBS) plus 0.05% Tween (TBST) and incubated in a blocking solution consisting of 10% normal goat (#S-1000, Vector Laboratories) or donkey serum (#017-000-121, Jackson ImmunoResearch) in TBST. Sections were then incubated with primary antibodies in 2% goat or donkey serum in TBST overnight at 4°C. Sections were incubated with primary antibodies, including the following: TIMP2 (1:100; #AF971, R&D Systems), NeuN (1:200; #MAB377, Millipore), doublecortin (DCX; 1:200; #4604S, Cell Signaling), Sox2 (1:200; #sc365823, Santa Cruz), Ki67 (1:200; #14-5698-82, Invitrogen), aggrecan (1:500; #AB1031, Millipore), Homer1 (1:500; #160006, Synaptic Systems). The next day, sections were washed in TBST three times for 5 min each and incubated with Alexa Fluor-conjugated secondary antibodies at 1:200 in TBST for 1 hour at RT. Secondary antibodies used were: Alexa Fluor 594-labeled donkey anti-rabbit (#A21206), Alexa Fluor 594-labeled donkey anti-mouse (#A21203), Alexa Fluor 488-donkey anti-rat (#A21208), Alexa Fluor 488-donkey anti-goat (#A11055), Alexa Fluor 647-donkey anti-mouse (#A31571), Alexa Fluor 594-goat anti-rabbit (#A11012), Alexa Fluor 5488-goat anti-chicken (#A11039), all from Life Technologies. Brain sections were then washed, stained with 4′,6-diamidino-2-phenylindole (DAPI, Sigma-Aldrich) for 15 min to visualize nuclei, mounted and coverslipped using Prolong Gold (Life Technologies) and dried overnight before imaging. For

BrdU staining, brain sections were pre-treated with 3M HCl (Thermo Fisher) for 30 min at 37°C

before overnight incubation with primary antibody anti-BrdU (1:500; #ab6326, Abcam) at 4°C.

Secondary staining and mounting were performed as described above.

Image processing was performed using an LSM 780 confocal microscope (Zeiss,

Oberkochen, DE) using 40x/1.4 Oil DIC objective. Four equally-spaced sections per mouse

(rostral through the caudal extent of the DG) were used to count the total number of positive cells

within the DG (subgranular zone and hilar subregions) of the hippocampus according to

stereological principles. All counts were performed using FIJI in a blinded fashion.

Aggrecan and Homer1 puncta quantification

Quantification of the number of puncta by super-resolution microscopy proceeded similar to the

method described in (Nguyen et al., 2020). Briefly, images were acquired by confocal microscopy

imaging using an LSM 880 with AiryScan in super-resolution mode (Zeiss) set with a 63X/1.4 Oil

DIC objective with 5X optical zoom. Using consistent laser strength, gain, and digitial offset

settings across all sections/experiments, Z stacks were acquired with steps less than the optical

slices acquired. Using Zen software (Zeiss), a setting of 6 ("optimal") was used during AiryScan

processing. Aggrecan puncta were quantified using the Puncta Analyzer plugin (Ippolito and

Eroglu, 2010) in ImageJ, and thresholding was applied equally across images. For colocalization

of co-stained Aggrecan and Homer1 puncta, images were analyzed with the Puncta Analyzer

plugin with a minimum pixel specification of 4. Three images in the molecular layer of the DG

were averaged per mouse for analysis.

Dendritic spine analysis by iontophoretic dye injections

Tissue processing. Mice were anesthetized using 30% chloral hydrate and transcardially perfused

with 1% PFA in phosphate buffer (PB, pH 7.4) for 2 min, followed by 4% PFA with 0.125%

glutaraldehyde in PB for 10 min at a rate of 5 ml/min. Brains were dissected and postfixed

overnight at 4°C in the same fixative, and then transferred to PBS with 0.1% sodium azide at 4°C

until sectioned. Brains were hemisected, and the right hemisphere was cut into 250 µm-thick

sections using a vibratome (VT1000S, Leica Microsystems, Buffalo Grove, IL).

Intracellular dye injection and confocal imaging. Coronal sections were incubated in 250 ng/ml

DAPI for 5 min to enable identification of the DG. Sections were mounted on nitrocellulose

membrane filters, immersed in ice-cold PB, and DG granule cells were iontophoretically injected

with 5% Lucifer Yellow (Invitrogen, Carlsbad, CA) in distilled water under a direct current of 3-8

nA until the dye filled the distal ends of the dendrites. Six to eleven neurons were injected per

section and neurons selected for injection were spaced to avoid overlapping of dendrites.

Sections were mounted with Fluoromount-G (#0100-01, Southern Biotech, Birmingham, AL) on

gelatin-coated glass slides (#22-214-320, Fisher).

To select dendritic segments for imaging on a Zeiss LSM 780 confocal microscope (Zeiss),

whole cells were first imaged at 20x/0.8 M27 Plan-Apochromat objective, using an Ar/Kr laser at

an excitation wavelength of 488 nm. Confocal stacks were acquired at 512 x 512-pixel resolution

with a Z-step of 1 µm, a pinhole setting of 1 Airy Unit, and optimal settings for gain and offset.

Basal dendritic segments at 50 µm from the soma were selected for high-resolution imaging

according to the following criteria: not a primary dendrite, no overlap with other dendrites or

branching that would obscure spines, not too deep in the section, and parallel or at acute angles

to the coronal plane. Dendritic segments were imaged using a 100x/1.46 Oil DIC M27 Plan-

Apochromat objective, and stacks were acquired at 512 x 512-pixel resolution with a Z-step of 0.1

µm, optical zoom of 3.3x, a pinhole setting of 1 Airy Unit, and optimal settings for gain and offset.

The stacks were imaged with approximately 1 µm above and below the segment to fully include

all spines. Three z-stacks were imaged from each neuron. Confocal stacks were deconvolved

using an iterative blind deconvolution algorithm (AutoQuant X, vX3.0.1, MediaCybernetics,

Bethesda, MD).

Spine reconstruction and analysis. Deconvolved stacks were analyzed using Neurolucida 360

(v2019.2.1; MBF Bioscience) for semi-automated reconstruction of dendrites and spines to

determine spine density and morphology. Spines were classified as stubby, thin, mushroom, and

filopodia morphologies. 4-5 mice per genotype, 6 neurons per mice, and 3 dendrites per neuron

were analyzed, and violin plots were used to better visualize the distribution of individual data

points.

Scanning electron microscopy (SEM)

Following ketamine/xylazine anesthesia and transcardial perfusion with PBS, 2-mm thick brain

slices containing the hippocampus were decellularized, following previously described methods

(De Waele et al., 2015). SEM methods were adapted from a previous study (Tajerian et al., 2018).

Decellularized brain slices were fixed in 4% PFA with 2% glutaraldehyde in 0.1M sodium

cacodylate buffer (pH 7.4) for 24 hours at 4°C. Samples were then briefly rinsed in the same

buffer before post-staining with 1% OsO4 for 1 hour. OsO_4-treated samples were rinsed in water

and gradually dehydrated in increasing concentrations of ethanol (50, 70, 90 100, 100%, 10 min

each). Samples were then stacked horizontally onto wire mesh dividers to keep them flattened

and critical point-dried with liquid CO_2. Dried samples were mounted onto Aluminum SEM stubs

using conductive copper tape and sputter-coated before imaging with a Zeiss Supra 55 VP

FESEM using InLens SE detection at 5 kV operating voltage (Carl Zeiss Microscopy Inc, USA).

Fiber diameters were analyzed using Image J (Bethesda, USA).

Behavior

All behavioral analyses were performed during the 0700–1900 light cycle. For all behavioral

experiments, age- (2-3 months old) and sex-matched mice with corresponding littermate control

animals were used.

Novel location recognition. Hippocampus-dependent memory was assessed using the novel location recognition as previously described (Castellano et al., 2017). On day 1, the training day, mice were habituated to the open-field arena for 6 min, an arena that contained wall-mounted visual cues. Mice were exposed to three consecutive trials of 6 min each, during which they explored two different objects in fixed positions. On day 2, the testing day, mice explored the same arena as on the training day but with one object displaced to a novel position. Time spent exploring each object was manually scored in a blinded fashion to assess the discrimination index for the novel location as the displaced object interaction time/total interaction time of both objects.

Contextual fear-conditioning. Fear-conditioning experiments were performed as previously described (Castellano et al., 2017) to assess hippocampus-dependent memory. Briefly, mice were trained to associate the cage context with an aversive stimulus (light foot-shock; Ugo Basile). On the first day, training parameters included two periods of 30 s consisting of a paired cue light and a tone of 1,000-Hz, followed by a light foot-shock (2 s, 0.5 mA) separated by a 180-s interval. Chambers were cleaned with 70% ethanol between experiments by the experimenter. Twenty-four hours later, mice were re-exposed to the same context for 3 min, and freezing levels (contextual) were measured. Two hours later, for the cued task, mice were placed in a novel context and exposed to the same tone and cue light from day 1 (training) after 120 s of exploration. Freezing levels were analyzed for pre- and post-cue phases. Freezing levels were measured using EthoVision XT system software (v14.0.1319, Noldus).

Barnes maze. Barnes maze testing was performed similarly to that previously described (Castellano et al., 2017). A large circular maze containing 40 holes was centered over a pedestal and elevated approximately 40 cm above the floor with a video recording device mounted directly above. The escape hole consisted of a PVC box similar in texture and color to the surface of the

maze. Distinct visual cues were placed at four equally spaced points around the maze to serve

as spatial navigation cues as the mice navigate the task. The task proceeded over 4 days, with

four trials on each day for each mouse. With overhead illumination and a persistent 2 kHz tone,

mice were given 90 s for each trial to identify the escape hole by jumping in or identifying the hole

with extended/overhead pokes. If mice failed to find the escape hole within 90 s, mice were gently

guided by light tapping/directing towards the escape hole and scored with 90 s. The escape hole

position was fixed within a day but changed for each successive day of testing. For each trial

within a day, the starting location for the mouse was rotated relative to the escape hole position.

Data collection and analyses were performed using EthoVision video-tracking system.

Search strategy classification was manually performed and based on methods adapted

from previous work (Mateus-Pinheiro et al., 2017), with data collected by EthoVision video-

tracking system software. Search strategies were coded according to the following: (i) localized -

minimal movement from the starting position; (ii) serial - sequential nose points within the outer

ring area; (iii) random - unorganized searching; (iv) scanning - arc-like trajectory, without nose

points; (v) focal - direct search within a maze quadrant; (vi) focal missense - direct search within

a maze quadrant from previous trials; (vii) targeted - direct movement to hole with deviations in

the trajectory direction; (viii) direct - no shifts in the trajectory direction with single nose point to

target hole. For strategy analysis, each related strategy was categorized into non-hippocampus-

dependent strategies comprising localized, serial, random, and scanning strategies, and

hippocampus-dependent strategies that included focal, focal missense, targeted and direct

strategies. Cognitive performance of each trial on day 3 was scored such that cognitive strategies

received higher scores according to the following scale, accounting for high cognitive complexity,

with a separation of 1 point between non-hippocampus-dependent and hippocampus-dependent

strategies: localized = 0, serial = 1, random = 2, scanning = 3, focal = 5, focal missense = 5,

targeted = 6, direct = 6. Focal and focal missense strategies are assigned a score of 5 since both

are goal-directed strategies used without achieving the final target. Target and direct strategies

were assigned score of 6 since both are goal-directed strategies with successful achievement of

the final target.

Open field. The open field test was used to evaluate locomotion and exploratory behavior, and

anxiety-like behavior in the center of the arena. The apparatus consisted of a brightly illuminated

square arena of 50 x 50 cm closed by walls 45 cm high. Mice were placed individually in the

center of the open field arena, and their movement was traced for 6 min. The resulting data was

analyzed using EthoVision software, considering two previously defined areas: a central and an

outer area. Due to the thigmotaxic exploratory activity of rodents, the ratio between the time spent

in the center and in the periphery of the open field arena can reflect anxiety-like behaviors.

Distance traveled, average velocity, and time spent in each of the zones were recorded and

analyzed.

Rotarod. Balance and motor function was measured using the fixed and standard accelerated

rotarod test (Med Associates, St. Albans, VT). The protocol consisted of a first day of training at

a constant speed (4 rpm) for a maximum of 2 min in three trials, with a 2-min interval between

each trial. On day 2, animals were tested for each of 3 different fixed speeds over a range of 4 to

40 rpm for a maximum of 60 sec each in three trials, with a 2-min interval between each trial. For

the acceleration test, animals were placed on the rod rotating at a constant speed (4 rpm), then

the rod started to accelerate continuously from 4 to 40 rpm over 5 min, in three trials with a 5-min

interval between each trial. The latency to fall off the rotarod was recorded for each trial, and data

from three trials were averaged for each mouse.

Wire grip test. Motor coordination and strength was measured by the wire grip task. Each mouse

was placed on a wire cage top, which was slowly inverted and suspended at approximately 30

cm above a mouse cage. The latency of time until the mouse fell off the wire was recorded in

three trials. Mice that did not fall within the 60-s trial period, were removed and assigned a

maximal time of 60 s. Data from three trials were averaged for each mouse.

Clasping. Mice were suspended by their tail, and the extent of hindlimb clasping was observed

for 15 s. Clasping scoring was based on a previously described method (Zhu et al., 2016). Briefly,

if both hindlimbs were splayed outward away from the abdomen with splayed toes, a score of 0

was given. If one hindlimb was retracted or both hindlimbs were partially retracted toward the

abdomen without touching it, and the toes were splayed, a score of 1 was assigned. If both

hindlimbs were partially retracted toward the abdomen and were touching the abdomen without

touching each other, a score of 2 was given. If both hindlimbs were fully clasped and touching the

abdomen, a score of 3 was assigned.

Grip strength. For the grip strength test, forelimb, hindlimb, and 4-limbs grip strength was

evaluated using a grip strength meter (Bioseb). Animals were held by the tail and allowed to place

forelimbs on the bar, or they were restrained to place hindlimbs on the bar connected to the force

sensor. Mice were slowly pulled away until they released the bar, and the maximal grip force

measured during the pulling session was recorded. For the 4-limb measurement, using a mesh

pull bar attachment, animals were handled by the tail so that all the paws latching onto the mesh

pull the bar together. The mice were slowly and steadily pulled away from the apparatus exactly

parallel from the countertop until paws were released from the mesh. A modified version of the

forelimb grip strength test was performed as previously described (Takeshita et al., 2017). Briefly,

the meter was rotated vertically and the measurement procedure was identical to the conventional

test, except for the direction in which the mouse's tail was pulled. Each animal's grip strength (in

g) was recorded, and data from five trials were collected.

Pole test. Motor coordination and balance with striatal involvement was assessed with the vertical pole test, which consisted of a plastic pole (50 cm high, diameter 1 cm) placed vertically on a mouse cage. Mice were individually placed at the top of the pole facing upward by their front paws on the pole, and the time taken to turn around and climb down the pole was recorded. A maximum time of 120 s was given to complete the task. If the mouse fell from the top of the pole, a time of 120 s was recorded. Each mouse underwent three trials, and the pole was cleaned with 70% ethanol between animals. Data from three trials were averaged for each mouse.

qPCR

After rapid dissection of the hippocampus, tissue was preserved in RNAlater (Invitrogen) overnight at 4°C before storing at −80°C until use. Hippocampal RNA was extracted according to the manufacturer's instructions using a RNeasy Mini Kit (Qiagen). Quality and concentration of total RNA were measured on the Nanodrop 8000 (Thermo Scientific). cDNA was subsequently synthesized with SuperScript III (Invitrogen), following the manufacturer's recommendations. Samples were mixed with SYBR Green master mix (Thermo Fisher) and primers for Mmp2 (FW: 5'-CAAGTTCCCCGGCGATGTC-3'; REV: 5'-TTCTGGTCAAGGTCACCTGTC-3') before loading as technical replicates for qPCR on a QuantStudio 7 Flex Real-Time PCR System (Thermo Scientific). Cycle counts for mRNA quantification were normalized to *Rplp0* (FW: 5'-AGATTCGGGATATGCTGTTGGC-3'; REV: 5'-CCAGTTGGTAACAATGCCATGT-3'). The ΔΔCT method was used to provide gene expression values, and expression levels were normalized to *Rplp0*.

Western blotting

For Western blotting of MMP2 or TIMP2 within hippocampus, dissected hippocampi were manually homogenized with 60 strokes in RIPA lysis buffer (Thermo Fisher), with protease inhibitors (Roche), before centrifugation at 20,000g for 25 min at 4°C. Protein concentration of the

supernatant was determined using the BCA kit (Thermo Fisher). Samples with equal amounts of

protein were separated on 4-12% NuPAGE Bis-Tris precast denaturing gels (Invitrogen) and

transferred onto nitrocellulose membranes (BioRad) at 125 V for 90 min. Membranes were

blocked with 3% milk-TBST (0.125% Tween) for 1 hour at RT and then probed with primary

antibodies diluted in 3% milk-TBST solution overnight at 4°C: anti-TIMP2 (1:500; #D18B7, Cell

Signaling), anti-MMP2 (1:1000; #ab86607, Abcam), and anti-actin (1:10000; #A5060, Sigma).

Membranes were washed and probed with horseradish-peroxidase-conjugated anti-mouse

(1:15,000; #1705047, BioRad) or anti-rabbit (1:15,000; #1705046, BioRad) for 1 hour at RT.

Membranes were developed using ECL Clarity reagent (BioRad) and imaged using ChemiDoc

MP Imaging System and Image Lab software (BioRad). Band intensities were quantified using

ImageJ software as previously described (Castellano et al., 2011).

For TIMP2 detection in ISF samples, 4 consecutive baseline eISF samples from

microdialysis were pooled and concentrated ~6 fold using Pierce Concentrator (PES, 3K MWCO,

Thermo Scientific). Concentrated samples were then processed for western blotting as described

above. Estimation of eISF TIMP2 protein levels was determined by extrapolation from a standard

curve of recombinant mouse TIMP2 (R&D/Biotechne) ranging from 0.31 ng to 20 ng run alongside

ISF samples in SDS-PAGE/Western that had been concentrated ~9.2-fold.

Data analysis

Statistical analyses were performed using GraphPad Prism version 9.0 software (GraphPad

Software, San Diego, California USA). Statistical tests are as described in text and figure legends,

including the use of student's t test for 2-group comparisons, nested t tests with multi-level

variables for 2-group comparisons, or chi-square analysis for 2-group comparisons of proportions.

All experiments were performed in a randomized and blinded fashion.

Results

TIMP2 is expressed in hippocampal neurons, and its deletion induces transcriptomic changes

TIMP2 exhibits high expression in the hippocampus relative to other brain regions (Castellano et al., 2017). To identify the cellular sources of TIMP2 in the hippocampus and to begin to probe its function in this region, we examined TIMP2 protein levels by confocal microscopy in various subfields of the hippocampus of 2-month-old wildtype (WT) male and female mice. We found that TIMP2 is expressed in the hilus of dentate gyrus (DG), as well as CA3 and CA1 subfields of the hippocampus, and we found that the majority of TIMP2-expressing cells co-localized with staining for pan-neuronal marker, NeuN, arguing that the major cellular source of TIMP2 expression in these areas was neuronal (**Figure 1A, 1B**). We did not detect differential TIMP2 neuronal expression in the hilus/DG, CA3, and CA1 subfields between males and females (**Figures S1A-C**).

Given the high level of TIMP2 expression seen in neurons in the hippocampus, we asked whether levels were elevated in the extracellular space adjacent to this cellular source. To measure extracellular levels of TIMP2 in this region, we implanted high molecular-weight cut-off (1-MDa) probes into hippocampi of freely moving, 2-3-month-old WT mice to measure interstitial fluid (ISF) levels of TIMP2 by *in vivo* microdialysis. TIMP2 was readily detectable in the ISF in hippocampus (**Figure 1D**), and extrapolated values of exchangeable ISF TIMP2 were ~10.1 ± 0.16 ng/mL (N=3), as measured at a flow rate of 1 μl/min. Importantly, the 21- to 22-kDa TIMP2 band was absent in the ISF of hippocampus from mice in which TIMP2 has been deleted (KO) (**Figure 1D**).

The high levels of TIMP2 within neurons and the extracellular space of the hippocampus argues for unappreciated roles for its function locally in this region, prompting us to examine the cellular and molecular processes regulated by TIMP2. In order to identify putative pathways involved in TIMP2 function in the hippocampus, we isolated hippocampi from WT and TIMP2 KO male and female mice and processed the tissue for RNA-sequencing (**Figure 1E**). A total of 905

differentially expressed genes (DEGs; *P<0.05*) were identified in TIMP2 KO compared to WT

(**Figure 1F**), with TIMP2 being the top differentially expressed gene, as expected. Of these genes,

449 DEGs were downregulated in TIMP2 KO hippocampi (**Figure 1F**). To gain insights into how

the altered transcriptome may reflect altered biological processes, we performed gene set

enrichment analysis (GSEA). Downregulated DEGs in TIMP2 KO were enriched for a number of

Gene Ontology (GO) biological processes related to plasticity, including "cell morphogenesis",

"neurogenesis", "cell projection organization", and "synapse organization" (**Figure 1G**).

Upregulated DEGs in TIMP2 KO were enriched in functions that included "regulation of cell death

and cell cycle", among others (**Figure S1E**). Together, our data suggest that TIMP2 in the normal

hippocampus may be involved in processes of plasticity.

TIMP2 is necessary for adult neurogenesis in the normal dentate gyrus

To evaluate the extent to which our discovery-based pathway approach from RNAseq highlighted

processes that may indeed be regulated by TIMP2, we examined the key cellular processes

implicated. As one of the top downregulated pathways in hippocampus of TIMP2 KO vs. WT mice

was "neurogenesis", we used confocal microscopy to begin to address whether processes of adult

neurogenesis within the DG are altered by absence of TIMP2. To examine whether TIMP2

regulates overall cell proliferation, we injected WT and TIMP2 KO mice with the proliferation

marker bromodeoxyuridine (BrdU), which labels the pool of proliferating cells in the subgranular

zone (SGZ) of DG (**Figure 2A**). We found that the DG of TIMP2 KO mice had significantly fewer

proliferating cells than in WT mice (**Figure 2C**), as quantified by the total number of BrdU$^+$ cells

(**Figure 2D**). We observed a similar result when analyzing the pool of Ki67$^+$ cells in the SGZ

(**Figure 2E-F**), which marks a larger pool of proliferating cells in other phases. We next analyzed

the pool of neural progenitor cells and the immature neuroblasts they ultimately give rise to, using

SRY-Box Transcription Factor 2 (Sox2) and doublecortin (DCX) markers, respectively. Deletion

of TIMP2 resulted in a significant decrease in the number of Sox2$^+$ neural progenitor cells (**Figure**

2G-H), and of DCX$^+$ immature neurons (**Figure 2I-J**).

To examine whether this disrupted pool of neuroblasts corresponds to a differing number

of surviving immature neurons, we injected TIMP2 KO and WT mice by pulse-chase labeling with

BrdU for 5 days and quantified the number of label-retaining, surviving BrdU$^+$ NeuN$^+$ neurons in

the SGZ of DG 4 weeks later (**Figure 2B**). Using this paradigm, we found that the DG of TIMP2

KO mice had significantly fewer newborn neurons compared with the DG of WT mice (**Figure 2K-**

L). Together, these results are consistent with TIMP2-mediated roles in adult neurogenesis and

related pathways uncovered by our transcriptomic analysis.

TIMP2 regulates dendritic spine plasticity in the dentate gyrus

Our transcriptomic data in TIMP2 KO hippocampi suggest a role for TIMP2 acting within the

hippocampus to modulate plasticity processes, including "synapse organization", possibly at the

structural level. To evaluate the impact of TIMP2 deletion on dendritic spines, the cellular

substrates of memory storage (Yang et al., 2009), we turned to a rigorous method to sensitively

quantify dendritic spines and their classification. We performed Lucifer Yellow iontophoretic dye-

filling of mature DG granule cells in isolated slices from TIMP2 KO and WT mice for high-

resolution imaging and reconstruction (**Figure 3A**) (Dickstein et al., 2016; Price et al., 2014). We

found that neurons in the DG of TIMP2 KO mice exhibited significantly reduced dendritic spine

density compared to those in WT DG (**Figure 3B, E**). We then used automated classification

algorithms (Dickstein et al., 2016; Rodriguez et al., 2008) to classify and quantify spines based

on morphology, as a readout of spine plasticity. Intriguingly, we found a significantly higher

proportion of immature thin spines in labeled DG neurons from TIMP2 KO mice compared to

those in WT mice (**Figure 3C, E**) and a significantly decreased proportion of mature mushroom

spines (**Figure 3D, E**) exhibited by TIMP2 KO relative WT DG neurons. Mushroom spines are a

particularly stable class of spines with lower turnover (Bourne and Harris, 2007) and are proposed

to serve as structural correlates of memory (Zuo et al., 2005). Of note, these described spine

observations in DG neurons of TIMP2 KO mice are in line with previous descriptions of altered

synaptic function upon TIMP2 modulation (Castellano et al., 2017), including long-term

potentiation, which was shown to be affected by modulating levels of TIMP2 (Castellano et al.,

2017). Collectively, these results suggest that TIMP2 may act to promote plasticity through

regulation of adult neurogenesis and dendritic spine formation.

TIMP2 is required for hippocampus-dependent memory function

The hippocampus is critical for spatial memory and learning, likely through regulation of adult

neurogenesis in the DG [reviewed in (Anacker and Hen, 2017)], as well as through changes in

synaptic structure (Jedlicka et al., 2008; Moser et al., 1994). The changes we observed in both

adult neurogenesis and dendritic spine plasticity in TIMP2 KO hippocampi prompted us to

investigate whether TIMP2 is involved in hippocampus-dependent memory and learning. We first

utilized the novel location recognition task (**Figure 4A**), in which mice use spatial cues in the

arenas to learn and associate the positions of objects in space. 24 hours after training, we

observed that TIMP2 KO mice exhibited significantly decreased preference for the training object

when displaced, as indicated by the reduced discrimination index (**Figure 4B**), thus suggesting a

critical role of TIMP2 in spatial memory in healthy mice. We next employed contextual and cued

fear-conditioning to evaluate whether the mice differ in the ability to discriminate the previously

aversive training context or a new cue-drive context (**Figure 4C**). Assessment of context

discrimination revealed significantly reduced freezing behavior by TIMP2 KO mice, suggesting

impaired hippocampus-associated contextual discrimination in the absence of TIMP2 (**Figure 4D-**

E), while amygdala-associated cued freezing levels were unchanged (**Figure S2A**). We next

examined whether more complex hippocampus-dependent spatial memory was altered in a

Barnes maze assay in which mice use a variety of strategies while navigating an illuminated maze

surface to an escape hole using spatial cues. Strategies were divided into non-hippocampus-

dependent and hippocampus-dependent with increasing levels of cognitive demand (**Figure 4F**)

(Mateus-Pinheiro et al., 2017). Analysis of the strategies used to reach the escape hole revealed

that TIMP2 KO mice delayed the switch from non-hippocampal-dependent to hippocampal-

dependent strategies while performing the task (**Figure 4G**). On day 3 of the Barnes maze, a

significant lower proportion of TIMP2 KO mice used hippocampus-dependent strategies

compared to WT mice (**Figure 4H**). Further categorization of cognitive performance by scoring

the different strategies according to level of complexity related to spatial learning (Illouz et al.,

2016) showed that TIMP2 KO mice achieved significantly lower cognitive scores relative to WT

mice (**Figure 4I**), consistent with impaired hippocampus-dependent function in TIMP2 KO mice

while performing in the Barnes maze.

Importantly, we did not detect differences in motor function and overall activity in TIMP2

KO mice (**Figure S2B-J**). Motor coordination perturbations have been reported in TIMP2 mutant

mice (Jaworski et al., 2006), so we subjected the mice to a wide battery of motor function,

coordination, and activity assays, including rotarod, open field, wire testing, pole testing, grip

strength, none of which revealed changes in corresponding function between TIMP2 KO and WT

mice (**Figure S2B-I**). We detected impaired clasping behavior in TIMP2 KO relative to WT mice

(**Figure S2J**), as previously reported (Jaworski et al., 2006), but this form of motor coordination

does not impact performance observed for TIMP2 KO mice in the wide range of hippocampus-

dependent tasks utilized. No differences in anxiety-related behaviors in the open field were

observed between WT and TIMP2 KO mice (**Figure S2F**). Collectively, these results indicate a

critical role for TIMP2 in regulation of hippocampus-dependent memory.

TIMP2 ablation leads to accumulation of ECM in the hippocampus

We next investigated potential mechanisms that may account for how spine formation and adult

DG neurogenesis are regulated by TIMP2. Outside the brain, TIMP2 has been characterized to

be involved in ECM degradation and remodeling through its canonical binding partner MMP2

(Caterina et al., 2000; Wang et al., 2000). Brain ECM is dynamic, and accumulating evidence

suggests a role for ECM and its components in the regulation of synaptic plasticity, including in

spine remodeling (Magnowska et al., 2016; Nguyen et al., 2020), adult neurogenesis (Yamada et

al., 2018), and cognition (Yamada et al., 2018). We first assessed the levels of pro-MMP2 in

lysates of hippocampi from TIMP2 KO and WT mice. Interestingly, we found significantly elevated

levels of MMP2, both at the gene (**Figure 5A**) and protein (**Figure 5B-C**) levels in the absence of

TIMP2, reflecting an accumulation in response to loss in TIMP2 signaling and pointing to a

potential disruption in ECM remodeling and turnover in hippocampus.

In the brain, chondroitin sulfate proteoglycans (CSPGs) are the major class of ECM

proteins (Miyata et al., 2018; Zimmermann and Dours-Zimmermann, 2008). Particularly, the

proteoglycan aggrecan is a core CSPG expressed in perineuronal nets in the brain and highly

enriched in the ECM of the DG (Rowlands et al., 2018) that has been linked to synaptic plasticity

(Fawcett et al., 2019) and memory (Gogolla et al., 2009; Romberg et al., 2013). We therefore

analyzed aggrecan distribution in the hippocampus of WT and TIMP2 KO mice. Using super-

resolution confocal imaging in brain sections isolated from TIMP2 KO and WT mice, we found

that TIMP2 KO mice exhibited a significant increase in the density of aggrecan puncta in the DG

molecular layer (**Figure 5D-E**), the same area where we observed impaired spine remodeling in

TIMP2 KO relative to WT mice (**Figure 3**). To further determine if TIMP2 remodeling of the ECM

could affect accumulation of ECM near synapses given the synaptic deficits we observed in these

neurons, we analyzed aggrecan colocalization with the postsynaptic protein Homer1. We

observed that the DG of TIMP2 KO mice exhibited significantly increased colocalization of

Aggrecan with Homer1 relative to that observed in WT DG (**Figure 5F-G**). These data indicate

that TIMP2 normally promotes ECM remodeling and limits its deposition near synapses.

We next further characterized the relationship between TIMP2's effect on ECM and adult

neurogenesis, considering the known roles of ECM and its related components in the regulation

of proliferation and migration of newborn neurons [reviewed in (Cope and Gould, 2019)]. To

explore this possibility, we evaluated the migration of newborn neurons toward the molecular layer

by examining the distribution of DCX$^+$ immature neuroblasts in the SGZ relative to that in the GCL

of the DG. This analysis revealed significantly decreased distribution of DCX$^+$ immature neurons

in the GCL of TIMP2 KO mice (**Figure 5H-I**), reflecting a process whereby migration of newborn

neurons within the DG is apparently impaired in the absence of TIMP2. These data suggest that

the absence of TIMP2 leads to a tighter ECM structural environment that acts to spatially restrict

the migration of newborn neurons and plasticity of synapses, thus affecting overall adult

neurogenesis and spine dynamics. In support of this concept, we measured structural changes in

the ECM of TIMP2 KO mice by performing surface scanning electron microscopy (SEM) on

decellularized brains isolated from TIMP2 KO and WT mice. Interestingly, we detected altered

ECM microarchitecture in the form of thicker ECM fibers (**Figure 5J-K**) in hippocampi from TIMP2

KO compared to WT mice. Together, our results suggest that TIMP2 modulates the flexibility of

the ECM within the hippocampus to regulate plasticity processes, including adult neurogenesis

and spine plasticity.

Creation of a conditional model to target neuronal pools of TIMP2

To gain further mechanistic insight into TIMP2's role in the function of the hippocampus with

increased cellular resolution, we generated *Timp2*-floxed (TIMP2$^{fl/fl}$) mice using two-cell

homologous recombination (2C-HR)-CRISPR/Cas9-based genome editing. We inserted two

loxP sites flanking exon 2 of the *Timp2* gene transcript (**Figure 6A-B**) and then crossbred *Timp2*$^{fl/fl}$

mice with Syn$^{Cre/+}$ mice (**Figure 6C**) to specifically excise *Timp2* in neurons. Immunoblotting

against TIMP2 from isolated hippocampus lysates of Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ or TIMP2$^{fl/fl}$ control mice

confirmed that conditional targeting of TIMP2 resulted in significantly reduced TIMP2 levels

compared to TIMP2$^{fl/fl}$ controls in hippocampus (**Figure 6D-E**). We performed further validation

by performing confocal microscopy in hippocampus of Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ or TIMP2$^{fl/fl}$ control mice

to examine the cells in hippocampus expressing TIMP2. Quantification of TIMP2$^+$ NeuN$^+$ cells in

the hilus/DG, CA3 and CA1 regions revealed significant abrogation of TIMP2 in neurons upon its

conditional deletion (**Figure 6F-G**). These data validate our model as a tool for addressing the

role of neuronal TIMP2 in regulating the key cellular and behavioral phenotypes observed in the

context of global TIMP2 deletion.

Neuronal TIMP2 regulation of the extracellular matrix regulates hippocampus-dependent

cognitive function and adult neurogenesis

Given that we found that TIMP2 was expressed by neurons in DG and present at significant levels

in the extracellular space adjacent to these cells, we sought to determine whether the pool of

neuronal TIMP2 is specifically involved in hippocampus-dependent memory. Using the Syn$^{Cre/+}$;

TIMP2$^{fl/fl}$ mice we generated and validated (**Figure 6**), we created cohorts of Syn$^{Cre/+}$; TIMP2$^{fl/fl}$

mice and TIMP2$^{fl/fl}$ control littermates to evaluate performance in spatial memory tasks. We found

that Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice exhibited significantly decreased preference for the displaced object

24 hours after training in the testing phase compared to TIMP2$^{fl/fl}$ controls, as indicated by a

reduced discrimination index in these mice (**Figure 7A**). Further assessment in the contextual

portion of fear-conditioning revealed a significant reduction in freezing levels by Syn$^{Cre/+}$; TIMP2$^{fl/fl}$

mice compared to TIMP2$^{fl/fl}$ controls, suggesting impaired contextual memory in the absence of

neuronal TIMP2 (**Figure 7B-C**). Cued freezing levels were indistinguishable between the groups

of mice (**Figure S3A**). We then subjected the mice to Barnes maze testing to examine

performance in search strategies to find the escape hole. Our search strategy analysis showed

that a smaller proportion of Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice used hippocampus-dependent strategies

compared to TIMP2$^{fl/fl}$ mice (**Figure 7D-E**). Scoring their search strategy utilization by complexity

shows that Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice score lower than their TIMP2$^{fl/fl}$ counterparts (**Figure 7F**),

suggesting TIMP2 from neurons is required for normal hippocampus-related memory and

learning. Importantly, we did not detect differences in overall motor function in terms of

coordination (**Figure S3B-C**), overall activity in the open field (**Figure S3D-E**), grip strength or

motor learning (**Figure S3G-I**) and clasping behavior (**Figure S3J**) in Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice

relative to TIMP2$^{fl/fl}$ mice. No differences in anxiety-related behaviors in the open field were

observed between TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (**Figure S3F**).

To test whether the local pool of neuronal TIMP2 affects the survival of newly generated

neurons, we analyzed the pool of BrdU^{+} NeuN^{+} cells after pulse-chase labeling with BrdU, as

described earlier (**Figure 2B**). Indeed, we observed that the deletion of TIMP2 within neurons

resulted in a significant decrease in the number of BrdU^{+} NeuN^{+} newborn neurons compared to

TIMP2$^{fl/fl}$ controls (**Figure 7G-H**).

To probe potential mechanisms by which TIMP2 expressed locally from neurons affects

hippocampus-dependent memory and adult neurogenesis, we next addressed ECM-related

remodeling in the DG of Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice. TIMP2 deletion in neurons in Syn$^{Cre/+}$; TIMP2$^{fl/fl}$

mice resulted in a significant increase in the density of aggrecan-positive puncta compared to

TIMP2$^{fl/fl}$ controls (**Figure 7I-J**), as well as increased colocalization of Aggrecan with Homer1

(**Figure 7I, 7K**), consistent with an accumulation of ECM components adjacent to synapses, as

observed earlier. Together, our results show that the pool of neuronal TIMP2 is necessary to

promote ECM remodeling to enable synaptic plasticity and adult neurogenesis to facilitate normal

hippocampus-dependent memory.

Discussion

Our study provides an overview of how alterations in the ECM within the hippocampus by TIMP2

can directly regulate features of plasticity, including dendritic spine plasticity, adult neurogenesis

and hippocampus-dependent memory. We find that TIMP2 is highly expressed in neurons within

hippocampus and expressed at high levels in the brain extracellular space. Its removal induces

structural synaptic changes and deficits in adult neurogenesis, as predicted from changes

observed at the transcriptomic level, both of which result in impairments in hippocampus-

dependent memory function. We demonstrate that TIMP2's role in ECM remodeling is associated

with changes in the flexibility of synaptic plasticity, impinging on processes like neuronal migration

and differentiation. We generate and characterize a novel mouse model for the conditional

deletion of TIMP2 within neurons in order to understand which phenotypes associated with TIMP2

can be attributed to the neuronal pool of TIMP2. We report that mice in which neuronal pools of

TIMP2 are targeted exhibited the major functional phenotypes observed in mice in which TIMP2

is deleted globally, highlighting molecular mechanisms through which neuronal TIMP2 promotes

synaptic plasticity through regulation in the ECM to facilitate normal memory.

TIMP2 regulation of synaptic remodeling and adult neurogenesis

The development and maintenance of synapses is critical in the establishment of functional

neuronal circuits, and emerging evidence indicates a role for the ECM and its components in

regulating synapse structure and morphology (Dansie and Ethell, 2011; Singhal and Martin,

2011). Few reports to date have shown that regulating the organization of the ECM is critical to

ensuring proper synaptic morphology (Kurshan et al., 2014). Herein, we show that the ECM

regulator TIMP2 is important for normal spine morphology in the hippocampus. These

observations are in line with previous *in vitro* reports that neurites of TIMP2 KO neurons are

significantly shorter than WT neurons (Perez-Martinez and Jaworski, 2005). We used high-

resolution imaging of labeled granule cells in the intact hippocampus of TIMP2 KO and WT mice

to show that TIMP2 regulates dendritic spine density, with a loss in KO mice in the class of spines

involved in memory, mushroom spines (Bourne and Harris, 2007). This finding is consistent with

reports that neutralization of TIMP2 leads to a dramatic loss in LTP (Castellano et al., 2017).

Moreover, when TIMP2 is targeted globally or specifically within neurons, ECM components like

aggrecan accumulate around synapses in the hippocampus and the ECM is more dense, based

on SEM data, suggesting that TIMP2 acts directly within the hippocampus to modulate plasticity

on a structural level. In agreement with its role in modulating plasticity, we also find that TIMP2

regulates the process of adult neurogenesis. While immature neuroblasts were unaltered in mice

with systemic modulation of TIMP2 in young mice (Castellano et al., 2017), presumably due to

the window over which TIMP2 was modulated being too short to reflect changes in DCX$^+$ cells,

we now show that removal of TIMP2, both globally and from neurons in the hippocampus, results

in disruption in adult neurogenesis, including in neural progenitor proliferation and maturation.

This is in line with sparse evidence *in vitro* indicating that TIMP2 expression was significantly

upregulated upon neuronal differentiation (Jaworski and Perez-Martinez, 2006), and that it may

be involved in differentiation (Perez-Martinez and Jaworski, 2005). Intriguingly, we observed that

TIMP2 deletion affected the migration of immature DCX$^+$ neurons towards the molecular layer,

arguing for a role of ECM in regulating neuronal migration (Wang et al., 2018). It has been

documented that ECM molecules participate in neurogenesis (Cope and Gould, 2019), though

the mechanisms by which they regulate neuronal differentiation are still limited. CSPGs are known

to surround new neurons in the hippocampus, and have recently been shown to be important in

the production of new neurons (Yamada et al., 2018). Moreover, ECM-related proteins, such as

reelin, have been shown to play critical roles in neuronal migration (Wang et al., 2018).

Roles of TIMP2 remodeling of the ECM within hippocampus

The molecular details regarding how molecules like TIMP2 may interact with the ECM to alter

functional plasticity remain unclear. TIMP2 is traditionally recognized for its unique dual role in

regulating both the activation of pro-MMP2 and the inhibition of MMP2 activity necessary for ECM

remodeling (Caterina et al., 2000; Wang et al., 2000). We observed that the deletion of TIMP2

leads to increased levels of *Mmp2* and pro-MMP2 levels within hippocampus, perhaps reflecting

compensatory responses to reduced TIMP2 and impaired MMP2 activation, as described outside

of the brain (Caterina et al., 2000; Wang et al., 2000). The accumulated pro-MMP2 appears to

translate to less remodeling of the ECM and its resulting accumulation around cells in the DG

neurogenic cells that we observe. A proposed mechanism of spine stabilization and remodeling

by ECM includes a structural spatial restriction mechanism in which ECM components, including

CSPGs, form a matrix around dendritic spines to provide extracellular rigidity and physically

restrict spine motion (Levy et al., 2014). MMPs themselves can modulate structural plasticity by

loosening the ECM structure, allowing for a more permissive environment (Cope and Gould,

2019). Moreover, stiffness of the surrounding environment determined by ECM complexity has

been shown to regulate cell migration (Choi et al., 2012; Vincent et al., 2013), including for neural

stem cell proliferation, differentiation, and migration *in vitro* (Leipzig and Shoichet, 2009;

Rammensee et al., 2017; Saha et al., 2008). This modulation has been proposed to be among

fundamental cell fate-determining factors in neural development and neurological disorders

(Barnes et al., 2017; Hall et al., 2021; Javier-Torrent et al., 2021). Our results argue that, in the

absence of TIMP2, ECM accumulation leads to a denser network that creates physical restrictions

that impact the turnover of synapses from thin into mature mushroom-shaped spines, while

impeding the proliferation and migration or cells related to the process of adult neurogenesis.

TIMP2 regulates functional effects exerted by the hippocampus

In line with reports implicating disrupted synaptic plasticity and neurogenesis in cognitive

dysfunction (Cope and Gould, 2019), we show how TIMP2 deletion affects hippocampus

plasticity, ultimately causing striking memory impairment. Both global and neuron-specific

targeting of TIMP2 levels leads to deficits in hippocampus-dependent memory tasks, mimicking

previously described phenotypes observed upon systemic targeting of TIMP2 in aging (Castellano

et al., 2017). Our results strongly reinforce a surprising link among TIMP2, synaptic plasticity, and

memory in the healthy hippocampus. Our data support the concept that TIMP2 regulates spine

dynamics and neurogenesis, which are important for memory. Importantly, the cognitive

phenotypes described here cannot be attributed to TIMP2's role in other brain regions related to

motor function, though earlier work in younger mice described subtle deficits in coordination upon

TIMP2 deletion (Jaworski et al., 2006), primarily related to cerebellar alterations deletion

(Jaworski et al., 2006). We show that TIMP2-mediated roles in ECM to regulate cellular

phenotypes are associated with changes in MMP2. TIMP2 likely interacts with many MMPs and

other binding partner to mediate functional changes under different conditions. Others have

shown that TIMP2 exerts diverse biological functions in addition to MMP2. Through its binding of

alpha3beta1 integrin, TIMP2 has been shown to regulate cell cycle progression in PC12 cells,

independent of MMP inhibitory activity (Perez-Martinez and Jaworski, 2005). It is possible that

TIMP2 exerts its diverse biological functions through interactions with different binding partners,

or that it exerts its activities differently *in vivo* versus *in vitro*, all of which will need to be further

explored in future studies. Moreover, while we mostly find TIMP2 expressed in neurons in our

study, TIMP2 expression from other cell types may be important in certain pathological contexts

(Keren-Shaul et al., 2017) that could be important to explore further, especially as a role for

microglia in remodeling the ECM to promote synapse plasticity in the hippocampus has been

recently described (Nguyen et al., 2020).

TIMP2 function in physiology, aging and pathology

Dysregulation of the ECM is implicated in cognitive dysfunction that occurs in aging (Birch, 2018)

and in disease (Sun et al., 2021). The progressive decline in TIMP2 expression with age

(Castellano et al., 2017) occurs in parallel with deficits in memory (Bettio et al., 2017) and ECM

accumulation (Vegh et al., 2014b), while systemic treatment with TIMP2 revitalizes hippocampus-

related function in aged mice (Castellano et al., 2017). Moreover, ECM clearance is beneficial in

mouse models of Alzheimer's disease pathology (Vegh et al., 2014a). It has been increasingly

appreciated that adult neurogenesis in the human DG is dysregulated in Alzheimer's disease

brain (Moreno-Jimenez et al., 2019), in addition to the loss observed in dendritic spine integrity.

Therefore, defining the precise molecular mechanisms accounting for how TIMP2 regulates

synaptic plasticity within the hippocampus in the healthy brain may lead to the development of

novel approaches for cognitive repair in contexts of aging or neurodegeneration.

Acknowledgments

We gratefully acknowledge the contributions of the Mouse Genetics and Gene Targeting (MGGT) CoRE facility (Kevin Kelley, ISMMS), which is supported by ISMMS and the National Cancer Institute of the NIH under award P30CA196521, for two-cell stage embryo microinjections for generating TIMP2$^{fl/fl}$ mice. We also thank Chingwen Yang of the CRISPR and Genome Editing Center (Rockefeller University) for assistance generating constructs to develop the TIMP2$^{fl/fl}$ line. We thank Yihang Wang (ISMMS) for RNA isolation support, Nikos Tzavaras and Deanna Benson of the Microscopy CoRE and Advanced Bioimaging Center (ISMMS) for advice and equipment for Airyscan microscopy and image analysis, Lee Cohen-Gould of the Electron Microscopy & Histology Services of Weill Cornell Medicine Microscopy & Image Analysis Core for SEM sample preparation, Jorge Morales of the Microscopy Facility (CCNY) for SEM acquisition support, and Ron Gordon and Andy Leonard of the EM Core (ISMMS) for advice and sample preparation for SEM. This work was funded by the National Institute on Aging (R01AG061382 (J.M.C), RF1AG072300 (J.M.C.), T32AG049688 (B.M.H.), and R01AG061382-02S1 (J.M.C., S.M.P)).

Author contributions

A.C.F. and J.M.C. conceived and designed the experiments, performed data analysis and interpretation, and wrote the manuscript. A.C.F. conducted the experiments. B.M.H. performed and analyzed microdialysis experiments. S.M.P analyzed data for RNAseq experiments. H. L. and J.D.Z performed or analyzed motor or Barnes maze experiments. M.V. performed intracellular dye injections for spine labeling, and analyzed/interpreted data with P.R.H. T.K. and T.A. assisted with design and analysis of TIMP2$^{fl/fl}$ validation data. All authors approved the final manuscript.

Declaration of interests

J.M.C. is listed as a co-inventor on patents for treating aging-associated conditions, including the

use of young plasma administration (US10688130B2) or youth-associated protein TIMP2

(US10617744B2), the latter of which is licensed to Alkahest, Inc.

References

Anacker, C., and Hen, R. (2017). Adult hippocampal neurogenesis and cognitive flexibility

- linking memory and mood. Nature reviews Neuroscience *18*, 335-346.

Barnes, J.M., Przybyla, L., and Weaver, V.M. (2017). Tissue mechanics regulate brain

development, homeostasis and disease. J Cell Sci *130*, 71-82.

Bettio, L.E.B., Rajendran, L., and Gil-Mohapel, J. (2017). The effects of aging in the

hippocampus and cognitive decline. Neurosci Biobehav Rev *79*, 66-86.

Birch, H.L. (2018). Extracellular Matrix and Ageing. Subcell Biochem *90*, 169-190.

Bourne, J., and Harris, K.M. (2007). Do thin spines learn to be mushroom spines that

remember? Curr Opin Neurobiol *17*, 381-386.

Brew, K., and Nagase, H. (2010). The tissue inhibitors of metalloproteinases (TIMPs): an

ancient family with structural and functional diversity. Biochimica et biophysica acta *1803*,

55-71.

Castellano, J.M., Kim, J., Stewart, F.R., Jiang, H., DeMattos, R.B., Patterson, B.W.,

Fagan, A.M., Morris, J.C., Mawuenyega, K.G., Cruchaga, C.*, et al.* (2011). Human apoE

isoforms differentially regulate brain amyloid-beta peptide clearance. Sci Transl Med *3*,

89ra57.

Castellano, J.M., Mosher, K.I., Abbey, R.J., McBride, A.A., James, M.L., Berdnik, D.,

Shen, J.C., Zou, B., Xie, X.S., Tingle, M.*, et al.* (2017). Human umbilical cord plasma

proteins revitalize hippocampal function in aged mice. Nature *544*, 488-492.

Caterina, J.J., Yamada, S., Caterina, N.C., Longenecker, G., Holmback, K., Shi, J.,

Yermovsky, A.E., Engler, J.A., and Birkedal-Hansen, H. (2000). Inactivating mutation of

the mouse tissue inhibitor of metalloproteinases-2(Timp-2) gene alters proMMP-2

activation. The Journal of biological chemistry *275*, 26416-26422.

Choi, Y.S., Vincent, L.G., Lee, A.R., Kretchmer, K.C., Chirasatitsin, S., Dobke, M.K., and

Engler, A.J. (2012). The alignment and fusion assembly of adipose-derived stem cells on

mechanically patterned matrices. Biomaterials *33*, 6943-6951.

Cope, E.C., and Gould, E. (2019). Adult Neurogenesis, Glia, and the Extracellular Matrix.

Cell Stem Cell *24*, 690-705.

Dansie, L.E., and Ethell, I.M. (2011). Casting a net on dendritic spines: the extracellular

matrix and its receptors. Dev Neurobiol *71*, 956-981.

De Waele, J., Reekmans, K., Daans, J., Goossens, H., Berneman, Z., and Ponsaerts, P.

(2015). 3D culture of murine neural stem cells on decellularized mouse brain sections.

Biomaterials *41*, 122-131.

Dewing, J.M., Carare, R.O., Lotery, A.J., and Ratnayaka, J.A. (2019). The Diverse Roles

of TIMP-3: Insights into Degenerative Diseases of the Senescent Retina and Brain. Cells

9.

Dickstein, D.L., Dickstein, D.R., Janssen, W.G.M., Hof, P.R., Glaser, J.R., Rodriguez, A.,

O'Connor, N., Angstman, P., and Tappan, S.J. (2016). Automatic Dendritic Spine

Quantification from Confocal Data with Neurolucida 360. Curr Protoc Neurosci *77*, 1 27

21-21 27 21.

Fager, N., and Jaworski, D.M. (2000). Differential spatial distribution and temporal

regulation of tissue inhibitor of metalloproteinase mRNA expression during rat central

nervous system development. Mech Dev *98*, 105-109.

Fawcett, J.W., Oohashi, T., and Pizzorusso, T. (2019). The roles of perineuronal nets and

the perinodal extracellular matrix in neuronal function. Nature reviews Neuroscience *20*,

451-465.

Gogolla, N., Caroni, P., Luthi, A., and Herry, C. (2009). Perineuronal nets protect fear

memories from erasure. Science *325*, 1258-1261.

Gu, B., Posfai, E., and Rossant, J. (2018). Efficient generation of targeted large insertions

by microinjection into two-cell-stage mouse embryos. Nat Biotechnol *36*, 632-637.

Guo, T., Hao, H., Zhou, L., Zhou, F., and Yu, D. (2018). Association of SNPs in the TIMP-

2 gene and large artery atherosclerotic stroke in southern Chinese Han population.

Oncotarget *9*, 4698-4706.

Hall, C.M., Moeendarbary, E., and Sheridan, G.K. (2021). Mechanobiology of the brain in

ageing and Alzheimer's disease. Eur J Neurosci *53*, 3851-3878.

Illouz, T., Madar, R., Louzoun, Y., Griffioen, K.J., and Okun, E. (2016). Unraveling

cognitive traits using the Morris water maze unbiased strategy classification (MUST-C)

algorithm. Brain Behav Immun *52*, 132-144.

Ippolito, D.M., and Eroglu, C. (2010). Quantifying synapses: an immunocytochemistry-

based assay to quantify synapse number. J Vis Exp.

Javier-Torrent, M., Zimmer-Bensch, G., and Nguyen, L. (2021). Mechanical Forces

Orchestrate Brain Development. Trends Neurosci *44*, 110-121.

Jaworski, D.M., and Perez-Martinez, L. (2006). Tissue inhibitor of metalloproteinase-2

(TIMP-2) expression is regulated by multiple neural differentiation signals. J Neurochem

98, 234-247.

Jaworski, D.M., Soloway, P., Caterina, J., and Falls, W.A. (2006). Tissue inhibitor of

metalloproteinase-2(TIMP-2)-deficient mice display motor deficits. J Neurobiol *66*, 82-94.

Jedlicka, P., Vlachos, A., Schwarzacher, S.W., and Deller, T. (2008). A role for the spine

apparatus in LTP and spatial learning. Behav Brain Res *192*, 12-19.

Kaczorowska, A., Miekus, N., Stefanowicz, J., and Adamkiewicz-Drozynska, E. (2020).

Selected Matrix Metalloproteinases (MMP-2, MMP-7) and Their Inhibitor (TIMP-2) in Adult

and Pediatric Cancer. Diagnostics (Basel) *10*.

Kandalam, V., Basu, R., Moore, L., Fan, D., Wang, X., Jaworski, D.M., Oudit, G.Y., and

Kassiri, Z. (2011). Lack of tissue inhibitor of metalloproteinases 2 leads to exacerbated

left ventricular dysfunction and adverse extracellular matrix remodeling in response to

biomechanical stress. Circulation *124*, 2094-2105.

Katsimpardi, L., Litterman, N.K., Schein, P.A., Miller, C.M., Loffredo, F.S., Wojtkiewicz,

G.R., Chen, J.W., Lee, R.T., Wagers, A.J., and Rubin, L.L. (2014). Vascular and

neurogenic rejuvenation of the aging mouse brain by young systemic factors. Science

344, 630-634.

Keren-Shaul, H., Spinrad, A., Weiner, A., Matcovitch-Natan, O., Dvir-Szternfeld, R.,

Ulland, T.K., David, E., Baruch, K., Lara-Astaiso, D., Toth, B.*, et al.* (2017). A Unique

Microglia Type Associated with Restricting Development of Alzheimer's Disease. Cell

169, 1276-1290 e1217.

Kurshan, P.T., Phan, A.Q., Wang, G.J., Crane, M.M., Lu, H., and Shen, K. (2014).

Regulation of synaptic extracellular matrix composition is critical for proper synapse

morphology. The Journal of neuroscience : the official journal of the Society for

Neuroscience *34*, 12678-12689.

Leipzig, N.D., and Shoichet, M.S. (2009). The effect of substrate stiffness on adult neural

stem cell behavior. Biomaterials *30*, 6867-6878.

Levy, A.D., Omar, M.H., and Koleske, A.J. (2014). Extracellular matrix control of dendritic

spine and synapse structure and plasticity in adulthood. Front Neuroanat *8*, 116.

Magnowska, M., Gorkiewicz, T., Suska, A., Wawrzyniak, M., Rutkowska-Wlodarczyk, I.,

Kaczmarek, L., and Wlodarczyk, J. (2016). Transient ECM protease activity promotes

synaptic plasticity. Sci Rep *6*, 27757.

Mateus-Pinheiro, A., Alves, N.D., Patricio, P., Machado-Santos, A.R., Loureiro-Campos,

E., Silva, J.M., Sardinha, V.M., Reis, J., Schorle, H., Oliveira, J.F.*, et al.* (2017).

AP2gamma controls adult hippocampal neurogenesis and modulates cognitive, but not

anxiety or depressive-like behavior. Mol Psychiatry *22*, 1725-1734.

Miyata, S., Nadanaka, S., Igarashi, M., and Kitagawa, H. (2018). Structural Variation of

Chondroitin Sulfate Chains Contributes to the Molecular Heterogeneity of Perineuronal

Nets. Front Integr Neurosci *12*, 3.

Moreno-Jimenez, E.P., Flor-Garcia, M., Terreros-Roncal, J., Rabano, A., Cafini, F.,

Pallas-Bazarra, N., Avila, J., and Llorens-Martin, M. (2019). Adult hippocampal

neurogenesis is abundant in neurologically healthy subjects and drops sharply in patients

with Alzheimer's disease. Nat Med *25*, 554-560.

Moser, M.B., Trommald, M., and Andersen, P. (1994). An increase in dendritic spine

density on hippocampal CA1 pyramidal cells following spatial learning in adult rats

suggests the formation of new synapses. Proceedings of the National Academy of

Sciences of the United States of America *91*, 12673-12675.

Nguyen, P.T., Dorman, L.C., Pan, S., Vainchtein, I.D., Han, R.T., Nakao-Inoue, H.,

Taloma, S.E., Barron, J.J., Molofsky, A.B., Kheirbek, M.A.*, et al.* (2020). Microglial

Remodeling of the Extracellular Matrix Promotes Synapse Plasticity. Cell *182*, 388-403

e315.

Nicaise, A.M., Johnson, K.M., Willis, C.M., Guzzo, R.M., and Crocker, S.J. (2019). TIMP-

1 Promotes Oligodendrocyte Differentiation Through Receptor-Mediated Signaling. Mol

Neurobiol *56*, 3380-3392.

Perez-Martinez, L., and Jaworski, D.M. (2005). Tissue inhibitor of metalloproteinase-2

promotes neuronal differentiation by acting as an anti-mitogenic signal. The Journal of

neuroscience : the official journal of the Society for Neuroscience *25*, 4917-4929.

Price, K.A., Varghese, M., Sowa, A., Yuk, F., Brautigam, H., Ehrlich, M.E., and Dickstein,

D.L. (2014). Altered synaptic structure in the hippocampus in a mouse model of

Alzheimer's disease with soluble amyloid-beta oligomers and no plaque pathology. Mol

Neurodegener *9*, 41.

Rammensee, S., Kang, M.S., Georgiou, K., Kumar, S., and Schaffer, D.V. (2017).

Dynamics of Mechanosensitive Neural Stem Cell Differentiation. Stem cells *35*, 497-506.

Rodriguez, A., Ehlenberger, D.B., Dickstein, D.L., Hof, P.R., and Wearne, S.L. (2008).

Automated three-dimensional detection and shape classification of dendritic spines from

fluorescence microscopy images. PloS one *3*, e1997.

Romberg, C., Yang, S., Melani, R., Andrews, M.R., Horner, A.E., Spillantini, M.G.,

Bussey, T.J., Fawcett, J.W., Pizzorusso, T., and Saksida, L.M. (2013). Depletion of

perineuronal nets enhances recognition memory and long-term depression in the

perirhinal cortex. The Journal of neuroscience : the official journal of the Society for

Neuroscience *33*, 7057-7065.

Rowlands, D., Lensjo, K.K., Dinh, T., Yang, S., Andrews, M.R., Hafting, T., Fyhn, M.,

Fawcett, J.W., and Dick, G. (2018). Aggrecan Directs Extracellular Matrix-Mediated

Neuronal Plasticity. The Journal of neuroscience : the official journal of the Society for

Neuroscience *38*, 10102-10113.

Saha, K., Keung, A.J., Irwin, E.F., Li, Y., Little, L., Schaffer, D.V., and Healy, K.E. (2008).

Substrate modulus directs neural stem cell behavior. Biophys J *95*, 4426-4438.

Singhal, N., and Martin, P.T. (2011). Role of extracellular matrix proteins and their

receptors in the development of the vertebrate neuromuscular junction. Dev Neurobiol

71, 982-1005.

Solga, R., Behrens, J., Ziemann, A., Riou, A., Berwanger, C., Becker, L., Garrett, L., de

Angelis, M.H., Fischer, L., Coras, R.*, et al.* (2019). CRN2 binds to TIMP4 and MMP14

and promotes perivascular invasion of glioblastoma cells. Eur J Cell Biol *98*, 151046.

Sun, Y., Xu, S., Jiang, M., Liu, X., Yang, L., Bai, Z., and Yang, Q. (2021). Role of the

Extracellular Matrix in Alzheimer's Disease. Front Aging Neurosci *13*, 707466.

Tajerian, M., Hung, V., Nguyen, H., Lee, G., Joubert, L.M., Malkovskiy, A.V., Zou, B., Xie,

S., Huang, T.T., and Clark, J.D. (2018). The hippocampal extracellular matrix regulates

pain and memory after injury. Mol Psychiatry *23*, 2302-2313.

Takeshita, H., Yamamoto, K., Nozato, S., Inagaki, T., Tsuchimochi, H., Shirai, M.,

Yamamoto, R., Imaizumi, Y., Hongyo, K., Yokoyama, S.*, et al.* (2017). Modified forelimb

grip strength test detects aging-associated physiological decline in skeletal muscle

function in male mice. Sci Rep *7*, 42323.

Ulrich, J.D., Burchett, J.M., Restivo, J.L., Schuler, D.R., Verghese, P.B., Mahan, T.E.,

Landreth, G.E., Castellano, J.M., Jiang, H., Cirrito, J.R.*, et al.* (2013). In vivo

measurement of apolipoprotein E from the brain interstitial fluid using microdialysis. Mol

Neurodegener *8*, 13.

Vegh, M.J., Heldring, C.M., Kamphuis, W., Hijazi, S., Timmerman, A.J., Li, K.W., van

Nierop, P., Mansvelder, H.D., Hol, E.M., Smit, A.B.*, et al.* (2014a). Reducing hippocampal

extracellular matrix reverses early memory deficits in a mouse model of Alzheimer's

disease. Acta Neuropathol Commun *2*, 76.

Vegh, M.J., Rausell, A., Loos, M., Heldring, C.M., Jurkowski, W., van Nierop, P.,

Paliukhovich, I., Li, K.W., del Sol, A., Smit, A.B*., et al.* (2014b). Hippocampal extracellular

matrix levels and stochasticity in synaptic protein expression increase with age and are

associated with age-dependent cognitive decline. Mol Cell Proteomics *13*, 2975-2985.

Villeda, S.A., Luo, J., Mosher, K.I., Zou, B., Britschgi, M., Bieri, G., Stan, T.M., Fainberg,

N., Ding, Z., Eggel, A*., et al.* (2011). The ageing systemic milieu negatively regulates

neurogenesis and cognitive function. Nature *477*, 90-94.

Villeda, S.A., Plambeck, K.E., Middeldorp, J., Castellano, J.M., Mosher, K.I., Luo, J.,

Smith, L.K., Bieri, G., Lin, K., Berdnik, D*., et al.* (2014). Young blood reverses age-related

impairments in cognitive function and synaptic plasticity in mice. Nat Med *20*, 659-663.

Vincent, L.G., Choi, Y.S., Alonso-Latorre, B., del Alamo, J.C., and Engler, A.J. (2013).

Mesenchymal stem cell durotaxis depends on substrate stiffness gradient strength.

Biotechnol J *8*, 472-484.

Wang, S., Brunne, B., Zhao, S., Chai, X., Li, J., Lau, J., Failla, A.V., Zobiak, B., Sibbe, M.,

Westbrook, G.L*., et al.* (2018). Trajectory Analysis Unveils Reelin's Role in the Directed

Migration of Granule Cells in the Dentate Gyrus. The Journal of neuroscience : the official

journal of the Society for Neuroscience *38*, 137-148.

Wang, Z., Juttermann, R., and Soloway, P.D. (2000). TIMP-2 is required for efficient

activation of proMMP-2 in vivo. The Journal of biological chemistry *275*, 26411-26415.

Yamada, J., Nadanaka, S., Kitagawa, H., Takeuchi, K., and Jinno, S. (2018). Increased

Synthesis of Chondroitin Sulfate Proteoglycan Promotes Adult Hippocampal

Neurogenesis in Response to Enriched Environment. The Journal of neuroscience : the

official journal of the Society for Neuroscience *38*, 8496-8513.

Yang, G., Pan, F., and Gan, W.B. (2009). Stably maintained dendritic spines are

associated with lifelong memories. Nature *462*, 920-924.

Zhu, J.W., Li, Y.F., Wang, Z.T., Jia, W.Q., and Xu, R.X. (2016). Toll-Like Receptor 4

Deficiency Impairs Motor Coordination. Front Neurosci *10*, 33.

Zhu, Y., Romero, M.I., Ghosh, P., Ye, Z., Charnay, P., Rushing, E.J., Marth, J.D., and

Parada, L.F. (2001). Ablation of NF1 function in neurons induces abnormal development

of cerebral cortex and reactive gliosis in the brain. Genes Dev *15*, 859-876.

Zimmermann, D.R., and Dours-Zimmermann, M.T. (2008). Extracellular matrix of the

central nervous system: from neglect to challenge. Histochem Cell Biol *130*, 635-653.

Zuo, Y., Lin, A., Chang, P., and Gan, W.B. (2005). Development of long-term dendritic

spine stability in diverse regions of cerebral cortex. Neuron *46*, 181-189.

1028

1029

 Figure 1

Figure 1. TIMP2 is expressed in neurons of the hippocampus, and its deletion induces

transcriptomic changes. **(A)** Low-magnification view (upper image) of mouse hippocampus and

high-magnification view (lower images) of hilus/DG, CA3, and CA1 subregions showing TIMP2$^+$

and TIMP2$^+$ NeuN$^+$ cells. Scale bars, 200 μm and 20 μm (inset). **(B)** Quantification of the total

number of TIMP2$^+$ cells and TIMP2$^+$ NeuN$^+$ cells across hippocampal subregions in WT mice (2

months of age; N = 8, males and females). **(C)** Schematic representation of high molecular-weight

cut-off (1-MDa) *in vivo* microdialysis to assay TIMP2 levels in mouse hippocampal ISF. **(D)** TIMP2

immunoblotting of hippocampal ISF dialyzed from 2-month-old WT and TIMP2 KO mice, with

corresponding Ponceau S stain. **(E)** Schematic representation of bulk RNA-seq workflow

performed in isolated WT and TIMP2 KO hippocampi (N=13-17/group, sex-matched) for

transcriptomic analysis. **(F)** Volcano plot showing the fold-change of genes (log$_2$ scale)

differentially expressed in hippocampus of TIMP2 KO vs. WT mice. Downregulated DEGs at *P* <

0.05 are highlighted in red (upregulated in black). **(G)** Top 10 significant pathways for

downregulated DEGs from Gene Set Enrichment Analysis. Data are represented as mean ± SEM.

DG, dentate gyrus; eISF, exchangeable interstitial fluid; aCSF, artificial cerebrospinal fluid.

1046

Figure 2.

Figure 2. TIMP2 is necessary for adult neurogenesis in the normal dentate gyrus

(A) Schematic timeline of BrdU intraperitoneal injection protocol to label proliferating cells in the

DG in isolated brain sections. **(B)** Schematic timeline of BrdU intraperitoneal injection protocol

used for cell fate "survival" labeling in DG of isolated brain sections. **(C)** Representative confocal

microscopy images of BrdU$^+$ cells in the DG of WT and TIMP2 KO mice (2-3 months of age, N =

11-12 mice per group; arrowheads indicate BrdU$^+$ cells; scale bar, 50 µm) with corresponding **(D)**

quantification of number of BrdU$^+$ cells in the dentate gyrus per unit area. **(E)** Representative

confocal microscopy images of proliferating Ki67$^+$ cells in the DG of WT and TIMP2 KO mice (2-

3 months of age, N = 11-12 mice per group; arrowheads indicate Ki67$^+$ cells; scale bar, 50 µm)

with corresponding **(F)** quantification of the number of Ki67$^+$ proliferating cells per DG in WT and

TIMP2 KO mice. **(G)** Representative confocal microscopy images of Sox2$^+$ in the DG of WT and

TIMP2 KO mice (2-3 months of age, N = 11-12 mice per group; arrowheads indicate Sox2$^+$ cells;

scale bar, 50 µm) with corresponding **(H)** quantification of Sox2$^+$ neural progenitor cells in the DG

of WT and TIMP2 KO mice. **(I)** Representative confocal microscopy images of DCX$^+$ cells in the

DG of WT and TIMP2 KO mice (2-3 months of age, N = 11-12 mice per group; scale bar, 50 µm)

with corresponding **(J)** quantification of DCX$^+$ immature neuroblasts in DG of WT and TIMP2 KO

mice. **(K)** Representative confocal microscopy images of BrdU$^+$ NeuN$^+$ cells in DG of WT and

TIMP2 KO mice (2-3 months of age. N = 8-11 mice per group; scale bar, 50 µm and 100 µm

(inset)) with corresponding **(L)** quantification of newborn neurons (BrdU$^+$ NeuN$^+$ cells) in the DG

of WT and TIMP2 KO mice. Data are represented as mean ± SEM. Student's *t*-test for two-group

comparisons. *$P<0.05$; **$P<0.01$; ***$P<0.001$. Data points represent individual mice. IHC,

immunohistochemistry; DG, dentate gyrus; DCX, doublecortin.

1071

1072   **Figure 3.**

1073

Figure 3. TIMP2 regulates dendritic spine plasticity within the dentate gyrus.

(A) Schematic representation of the workflow for dendritic spine quantification using Lucifer

Yellow-filled DG granule cells. **(B)** Overall dendritic spine density in DG granule cells

iontophoretically labeled with Lucifer Yellow using sections isolated from six-month-old WT and

TIMP2 KO mice (N=6 neurons per mouse from N=4-5 mice per group). **(C)** Quantification of the

percentage of spines categorized according to "thin" spine heads or **(D)** "mushroom" spine heads.

(E) Representative deconvolved confocal image of a dendritic segment from WT and TIMP2 KO

Lucifer Yellow-labeled brain sections, and the downstream 3D reconstruction, with dendritic

segment shown in pink, thin spines in green, stubby in blue, mushroom in red, and filopodia in

yellow. Scale bar, 2 µm. Data are represented as mean ± SEM. Nested t-test for comparisons

with neuron and mouse as levels. *$P<0.05$; **$P<0.01$. Data points represent neurons (left) and

mice (right) for each group. LY, Lucifer Yellow.

1086

1087 **Figure 4.**

Figure 4. TIMP2 is required for hippocampus-dependent memory function.

(A) Schematic of the novel location recognition assay. **(B)** Discrimination index for novel location recognition on day 2 for WT and TIMP2 KO mice (2-3 months of age, N = 11-12 mice per group). **(C)** Schematic of the contextual and cued fear-conditioning assay. **(D, E)** Freezing levels by interval and overall measured in the conditioned-fear context A in WT and TIMP2 KO mice (2-3 months of age, N = 11-12 mice per group). **(F)** Schematic diagram of modified Barnes maze with color coding for overall strategy classification. **(G)** Proportion of WT and TIMP2 KO mice using non-hippocampus-dependent (gray) and hippocampus-dependent (yellow) strategies during the testing trials (2-3 months of age, n = 9 mice per group). **(H)** Strategy utilization by WT and TIMP2 KO mice on day 3 of the Barnes Maze and corresponding **(I)** cognitive scores, ranked by strategy complexity for WT and TIMP2 KO mice on day 3 in the Barnes Maze. Data are represented as mean ± SEM. Student's t-test for two-group comparisons in **(B, D, E)**, chi-square test in **(H)**, and nested t-test **(I)** for trial and mouse levels. *$P<0.05$, **$P<0.01$. Data points represent individual mice.

1105 **Figure 5.**

1106

Figure 5. TIMP2 ablation leads to accumulation of ECM within the hippocampus.

(A) *MMP2* mRNA levels by qPCR from isolated hippocampus of WT and TIMP2 KO mice (2-3

months of age, N = 6 mice per group). **(B)** Immunoblotting of MMP2 from hippocampal lysates of

WT and TIMP2 KO mice (2-3 months of age, N = 9 mice per group) with corresponding **(C)**

quantification of MMP2 protein levels. **(D)** Representative confocal microscopy images showing

aggrecan puncta in the molecular layer of DG from WT and TIMP2 KO mice (2-3 months of age,

N = 8-13 mice per group; scale bar, 5 µm) with corresponding **(E)** quantification of puncta. **(F)**

Representative confocal microscopy images and corresponding **(G)** quantification of aggrecan

and homer1 co-localization in the molecular layer of DG from WT and TIMP2 KO mice (2-3 months

of age, N = 8-13 mice per group; scale bar, 5 µm). Circles indicate colocalized puncta. **(H)**

Representative images showing the migration of DCX$^+$ cells in the SGZ and GCL of WT and

TIMP2 KO mice (2-3 months of age, N = 11-12 mice per group, scale bar, 20 µm with

corresponding **(I)** quantification of DCX$^+$ cells distributed in the subgranular zone (SGZ) and

granule cell layer (GCL) of DG. **(J)** Representative surface scanning electron microscopy (SEM)

from DG of WT and TIMP2 KO mice (2-3 months of age, N = 3 mice per group, scale bar, 2 µm,

with corresponding **(K)** quantification of ECM microarchitecture in terms of fiber diameter

measurements. Data are represented as mean ± SEM. Student's t-test for two-group

comparisons. *$P<0.05$, ** $P<0.01$, ***$P<0.001$. Data points represent individual mice.

1125

1126   **Figure 6.**

1127

Figure 6. Creation of a conditional model to target neuronal pools of TIMP2.

(A) Schematic illustration of the targeting strategy of to insert loxP sites flanking exon 2 to

generate a model for conditional deletion of *TIMP2*. **(B)** Representative genotyping results

revealing the PCR products for the mutant LoxP and wild-type alleles. Wild-type (+/+),

heterozygous (fl/+), and homozygous (fl/fl) mice were identified according to this strategy. **(C)**

Schematic diagram of cross-breeding strategy to establish neuron-specific TIMP2 deletion: male

TIMP2$^{fl/fl}$ mice were mated with Syn$^{Cre/+}$ females to obtain Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ and their respective

TIMP2$^{fl/fl}$ littermate controls. **(D-E)** Representative TIMP2 immunoblot and corresponding

quantification of TIMP2 protein levels from hippocampal lysate of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$

(2-3 months of age, N = 5-6 mice per group). **(F)** High-magnification view of hilus/DG, CA3, and

CA1 sub-regions of TIMP2+ cells co-expressing NeuN in TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (2-

3 months of age, N = 5 mice per group, scale bar, 20 µm) with corresponding **(G)** quantification

of the total number of TIMP2+ cells with NeuN+ nuclei across hippocampal subregions. Data are

represented as mean ± SEM. Student's t-test for two-group comparisons. *P<0.05, **P<0.01,

***P<0.001, ****P<0.0001. Data points represent individual mice.

1143

Figure 7.

Figure 7. Neuronal TIMP2 regulation of the extracellular matrix regulates hippocampus-dependent cognitive function and adult neurogenesis.

(A) Discrimination index for novel location recognition on day 2 and **(B,C)** contextual fear-conditioning freezing levels in TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (2-3 months of age, N = 13-15 mice per group). **(D)** Proportion of mice of each genotype using non-hippocampus-dependent (gray) and hippocampus-dependent (yellow) search strategies during the testing trials of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice in Barnes maze (2-3 months of age, N = 12-14 mice per group) with **(E)** quantification of the percentage of mice using these strategies by on day 3. **(F)** Cognitive complexity scores for TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice based on strategies used on day 3 of Barnes maze. **(G)** Representative confocal microscopy images of BrdU$^+$ and NeuN$^+$ cells in DG of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (2-3 months of age, N = 7–8 mice per group, scale bar, 100 μm) with corresponding **(H)** quantification of BrdU$^+$ NeuN$^+$ newborn neurons in the DG. **(I)** Representative confocal microscopy images of aggrecan and homer1 puncta in molecular layer of DG of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (N = 7-8 mice per group, scale bar, 5 μm) with corresponding **(J)** quantification of aggrecan puncta and **(K)** co-localized puncta of aggrecan and homer1, as indicated by overlaid circles. Data are represented as mean ± SEM. Student's *t*-test for two-group comparisons in **(A-C)**, **(H)**, and **(J, K)**, chi-square test in **(E)**, and nested t-test **(F)** with trial and mice as levels. *$P<0.05$, ** $P<0.01$, ***$P<0.001$, ****$P<0.0001$. Data points represent individual mice.

1166   **Figure S1.**
1167

Figure S1. TIMP2 protein is found in hippocampal neurons in male and female mice, and

its loss affects hippocampal transcriptome.

(A) Quantification of the total number of TIMP2$^+$ and TIMP2$^+$ NeuN$^+$ cells at the hilus/DG, **(B)** CA3

and **(C)** CA1 subregions of the hippocampus from WT males and females at 2 months of age (N

= 4 mice per sex). **(D)** Cresyl violet-stained section depicting the microdialysis probe tract through

brain surface extending into hippocampus, with microdialysis probe position seen at Bregma -

3.08mm based on Paxinos and Watson atlas. Scale bar, 500 µm. **(E)** Top 10 significant pathways

for upregulated DEGs in hippocampi of TIMP2 KO mice relative to WT mice (N = 13-17 mice per

group). Data are represented as mean ± SEM.

1177

1178   **Figure S2.**

Figure S2. Loss of TIMP2 does not affect amygdala-associated behavior or overall motor function

(A) Percentage of freezing detected in the cued task of the fear-conditioning assay in WT and TIMP2 KO mice (2-3 months of age, N = 12 mice per group). **(B)** Latency of TIMP2 KO and WT mice to fall in the rotarod in the fixed and **(C)** acceleration protocol at 4, 20 and 40 rpm (N = 12 mice per group). **(D)** Total distance traveled by TIMP2 KO and WT mice in the open field, as well as **(E)** velocity, and **(F)** percentage of time spent in the center of the arena (N = 11-12 mice per group). **(G)** Latency of TIMP2 KO and WT mice to fall in the wire test (N = 12 mice per group). **(H)** Time for TIMP2 KO and WT mice to descend the pole in the pole test (N = 11-12 mice per group). **(I)** Grip strength of the fore-, hind- and four limbs in TIMP2 KO and WT mice (N = 12 mice per group). **(J)** Hindlimb extension by clasping score in TIMP2 KO and WT mice (N = 10-11 mice per group). Data are represented as mean ± SEM. Student's *t*-test for two-group comparisons. n.s., not significant.

1195 **Figure S3.**

Figure S3. Neuronal TIMP2 deletion does not impair amygdala-associated behavior or overall motor function

(A) Percentage of freezing detected in the cued task of the fear-conditioning assay in TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (2-3 months of age, N = 13-15 mice per group). **(B)** Latency of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice to fall in the rotarod in the fixed and **(C)** acceleration protocol at 4, 10 and 20 rpm (N = 13-15 mice per group). **(D)** Total distance traveled by TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice in the open field, as well as **(E)** velocity, and **(F)** percentage of time spent in the center of the arena (N = 13-15 mice per group). **(G)** Latency of TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice to fall in the wire test (N = 12-14 mice per group). **(H)** Time for TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice to descend the pole in the pole test (N = 12-14 mice per group). **(I)** Grip strength of the fore-, hind- and four limbs in TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (N = 13-15 mice per group). **(J)** Hindlimb extension by clasping score in TIMP2$^{fl/fl}$ and Syn$^{Cre/+}$; TIMP2$^{fl/fl}$ mice (N = 12-14 mice per group). Data are represented as mean ± SEM. Student's *t*-test for two-group comparisons. n.s., not significant.