Clinical Research Article

DI/cle, a Measure Consisting of Insulin Sensitivity, Secretion, and Clearance,

Captures Diabetic States

Hikaru Sugimoto,[1] Ken-ichi Hironaka,[2] Tomoko Yamada,[3] Kazuhiko Sakaguchi,[3] Wataru

Ogawa,[3]* and Shinya Kuroda[1,2]*

[1]Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The

University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[2]Department of Biological Sciences, Graduate School of Science, The University of Tokyo,

7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[3]Division of Diabetes and Endocrinology, Department of Internal Medicine, Kobe University

Graduate School of Medicine, Kusunoki-cho 7-5-1, Chuo-ku, Kobe, Hyogo 650-0017, Japan.

*Corresponding authors: Wataru Ogawa, ogawa@med.kobe-u.ac.jp; or Shinya Kuroda,

skuroda@bs.s.u-tokyo.ac.jp

Keywords. Insulin clearance, Insulin secretion, Insulin sensitivity, Disposition index

Funding and Assistance. This study was supported by the Japan Society for the Promotion of

Science (JSPS) KAKENHI (JP21H04759), CREST, the Japan Science and Technology

Agency (JST) (JPMJCR2123), and The Uehara Memorial Foundation.

Conflict of Interest. The authors declare no competing interests.

ABSTRACT

Context

Insulin clearance is implicated in regulation of glucose homeostasis independently of insulin

sensitivity and insulin secretion.

Objective

To understand the relation between blood glucose and insulin sensitivity, secretion, and

clearance.

Methods

We performed a hyperglycemic clamp, a hyperinsulinemic-euglycemic clamp, and an oral

glucose tolerance test (OGTT) in 47, 16, and 49 subjects with normal glucose tolerance

(NGT), impaired glucose tolerance (IGT), and type 2 diabetes mellitus (T2DM), respectively.

Mathematical analyses were retrospectively performed on this dataset.

Results

The disposition index (DI), defined as the product of insulin sensitivity and secretion, showed

a weak correlation with blood glucose levels, especially in IGT ($r =$

$0.04;\ 95\%\ \text{CI}, -0.63 - 0.44$). However, an equation relating DI, insulin clearance, and blood

glucose levels was well conserved regardless of the extent of glucose intolerance. As a

measure of the effect of insulin, we developed an index, designated "DI$_{\text{cle}}$," that is based on

this equation and corresponds to DI divided by the square of insulin clearance. DI$_{\text{cle}}$ was not

impaired in IGT compared with NGT, possibly as a result of a decrease in insulin clearance

in response to a reduction in DI, whereas it was impaired in T2DM relative to IGT.

Moreover, DI$_{\text{cle}}$ estimated from a hyperinsulinemic-euglycemic clamp, OGTT, or a fasting

blood test were significantly correlated with that estimated from two clamp tests ($r =$

$0.52;\ 95\%\ \text{CI}, 0.37 - 0.64, r = 0.43;\ 95\%\ \text{CI}, 0.24 - 0.58, \text{and } r =$

$0.54;\ 95\%\ \text{CI}, 0.38 - 0.68, \text{respectively}$).

50 **Conclusions**

51 $DI_{/cle}$ can serve as a new indicator for the trajectory of changes in glucose tolerance.

INTRODUCTION

The blood glucose concentration is maintained within a narrow range by feedback-compensatory responses in the body. Insulin sensitivity and insulin secretion are linked by such a relation, with insulin secretion increasing to maintain normal glucose homeostasis as insulin sensitivity declines (1). This compensatory response has been regarded as "moving up" the hyperbolic curve of the product of insulin sensitivity and insulin secretion, and the failure of this response, regarded as "falling off" the hyperbolic curve, is associated with the transition from normal glucose tolerance (NGT) to impaired glucose tolerance (IGT) and type 2 diabetes mellitus (T2DM) (1). The disposition index (DI), generally defined as the product of insulin sensitivity and insulin secretion, has therefore been adopted as a measure that correlates well with blood glucose levels (1,2).

Several models that explain this moving up–falling off phenomenon as an interplay among insulin sensitivity, insulin secretion, and pancreatic β-cell mass have been described (3–6). Such models account for the physiological features and pathological trajectory of glucose tolerance—including NGT, IGT, and T2DM—to a certain extent, but their conformity to clinical or experimental data is not high (6). Moreover, the moving up and falling off the curve of the product of insulin sensitivity and insulin secretion appear not to adequately represent the relation between blood glucose and insulin levels. For example, comparison of individuals with NGT at extremes of insulin sensitivity revealed that an approximately twofold increase in blood insulin levels was associated with an increase in insulin secretion of <50% (7,8), indicating that the increase in insulin secretion may not fully explain the increase in circulating insulin levels in individuals with insulin resistance (8).

Insulin clearance is implicated as a determinant of circulating insulin levels that is independent of insulin secretion (8–16). Insulin clearance likely contributes to the compensatory responses of the body, with a reduction in insulin clearance in the insulin-

resistant state resulting in an increase in circulating insulin levels and a lessening of the

demand for insulin secretion by β-cells (11). In some instances, changes in the circulating

insulin levels do not reflect changes in insulin secretion, instead being primarily influenced

by insulin clearance (17). Furthermore, changes in insulin clearance have been implicated in

the pathogenesis of glucose intolerance (18–21). Whereas these various findings suggest the

importance of insulin clearance in the physiology and pathology of glucose homeostasis, it is

difficult to directly evaluate this parameter from clinical data (22).

Mathematical models provide insight into phenomena in which various factors

mutually interact and for which effects of such factors cannot be measured directly (22–25).

We previously developed a mathematical model based on temporal changes in circulating

glucose and insulin levels during consecutive hyperglycemic and hyperinsulinemic-

euglycemic clamps (23). Using this model, we estimated insulin clearance and characterized

its role in the regulation of glucose homeostasis, and we identified a relation between insulin

clearance and the product of insulin sensitivity and insulin secretion (23). However, this

study did not examine the goodness-of-fit of the relation in individuals with each category of

glucose tolerance.

The objectives of this study were (1) to test the hypothesis that the product of insulin

sensitivity and insulin secretion cannot fully represent blood glucose levels and insulin

clearance, (2) to derive a better equation relating these parameters from an empirical method

that necessitates numerical calculations and simple mathematical models that can be solved

analytically, and (3) to develop a simple index to assess the relation. We here propose a new

index, "DI_{cle}," that consists of DI corrected by insulin clearance. We have characterized DI_{cle}

in individuals with various levels of glucose tolerance, and we provide evidence that this new

index can serve as an indicator of the effect of insulin on glucose homeostasis and can

capture the transition from IGT to T2DM.

MATERIALS AND METHODS

Subjects and Measurements

We analyzed a previously reported data set (23) in the present study. This study was

approved by the ethics committee of Kobe University Graduate School of Medicine

(UMIN000002359). Participants were recruited between October 2008 and December 2011.

In brief, a hyperglycemic clamp was performed by intravenous infusion first of a bolus of

glucose over 15 min and then of a variable amount of glucose to maintain the plasma glucose

level at 200 mg/dL. Ten minutes after the end of the hyperglycemic clamp, a

hyperinsulinemic-euglycemic clamp was performed by intravenous infusion of insulin with a

target plasma glucose level of 90 mg/dL or of the fasting level in the case of NGT and IGT

subjects whose fasting plasma glucose level was <90 mg/dL. Blood was obtained from

arterialized veins that were confirmed by oxygen saturation. These clamp analyses and a 75-g

oral glucose tolerance test (OGTT) were performed within a period of 10 days.

Fifty NGT, 18 IGT, and 53 T2DM subjects were initially used for parameter

estimation. As we previously described (23), three NGT, two IGT, and three T2DM

individuals were subsequently eliminated as outliers. In addition, one T2DM individual was

eliminated because of missing values, with the remaining 47 NGT, 16 IGT, and 49 T2DM

subjects being analyzed. Detailed characteristics of each subject are described in the previous

study (23).

Mathematical models

To investigate the relation among insulin sensitivity, insulin secretion, insulin clearance, and

blood glucose levels, we examined a previously developed mathematical model (23), which

is written as follows:

$$\frac{dG}{dt} = k_1 Y - k_2 G + \frac{k_3}{1+I} - k_4 GI,$$

$$\frac{dI}{dt} = k_6 X - k_7 I,$$

$$\frac{dY}{dt} = -k_1 Y + k_2 G + \text{influx } G,$$

$$\frac{dX}{dt} = k_5 Y - k_6 X + \text{influx } I,$$

where G and I denote blood glucose and insulin concentrations, respectively. The variable Y

corresponds to the effective glucose (the portion of glucose that affects insulin secretion) on

induction of variable X, which can be regarded as insulin secreted from β-cells. The fluxes

influx G and influx I denote glucose and insulin infusions, respectively. The indices k_4, k_5,

and k_7 correspond to insulin sensitivity, insulin secretion, and insulin clearance, respectively.

The parameters were estimated to minimize residual sum of the square (RSS) between the

time course during clamp tests and the model trajectories, as previously described (23).

To validate the relation among the parameters, we also calculated insulin secretion,

insulin sensitivity, and insulin clearance by using other methods. Insulin secretion was

calculated from the deconvolution of the fasting serum C-peptide concentration (26). We

investigated the first-phase insulin secretion, as follows:

$$\dot{C} = k_{Sb} G + k_{Sf} \dot{G}_+ - k_{Cc} C + k_{CQ} Q - k_{QC} C,$$

$$\dot{Q} = k_{QC} C - k_{CQ} Q,$$

where k_{Sb}, k_{Sf}, and k_{Cc} correspond to basal insulin secretion, first-phase insulin secretion,

and C-peptide clearance, respectively. The variable C and Q denote the concentrations of C-

peptide in the accessible and non-accessible compartments, respectively (26). k_{CQ} and k_{QC}

are rate parameters describing C-peptide exchange kinetics (26). By approximating the

second-order derivative as zero and $k_{Sb} G = k_{Cc} C$ at 0 min, we calculated k_{Sf} as follows:

$$k_{Sf} \sim \frac{\dot{C} + \dot{Q}}{\dot{G}_+} \sim \frac{C_5 - C_0}{G_5 - G_0} \left(1 + \frac{k_{QC}}{k_{CQ}} \right).$$

The subscripts of C and G denote the time (min) during a hyperglycemic clamp. Five

individuals with $k_{Sf} \leq 0$ were removed in this analysis.

As an index of insulin sensitivity, the insulin sensitivity index (ISI) was calculated,

as described previously (23,27,28). As an index of insulin clearance, we calculated metabolic

clearance rate of insulin (MCR) (23). To further examine the relation between blood insulin

and glucose, we calculated basal insulin effect (29).

To further validate the relation among the parameters, and to analytically estimate

the relation among the parameters, we also examined a simple and stable model, which can

be written as follows (30):

160
$$\frac{dG}{dt} = -k_{glu}G - k_{sen}IG + f,$$

161
$$\frac{dI}{dt} = \frac{k_{sec}}{k_{tim}} \int_{t-k_{tim}}^{t} G(s)ds - k_{cle}I,$$

where k_{glu}, k_{sen}, k_{sec}, k_{tim}, and k_{cle} are glucose effectiveness, insulin sensitivity, insulin

secretion, the duration of the preceding interval whose plasma glucose influence the current

insulin secretion, and insulin clearance, respectively. f is the sum of the constant baseline

hepatic glucose release and the amount of infused glucose.

To investigate how the relation among the parameters change with the progression of

glucose intolerance, we also examined βGI model (5,19), which has been used to explore the

characteristics of glucose homeostasis and is written as follows:

169
$$\frac{dG}{dt} = u_0 + u(t) - (k_{glu} + k_{sen}I)G,$$

170
$$\frac{dI}{dt} = p\beta \cdot \rho(G) - k_{cle}I,$$

171
$$\frac{d\beta}{dt} = \beta \cdot h(G),$$

where u_0, $u(t)$, p, and β are endogenous glucose production and infused glucose, insulin

secretion and beta-cell functional mass, respectively. $\rho(G)$ is a monotonically increasing

function of G. $h(G)$ is the beta-cell growth rate, and at the glucose set point, $G = G_0$,

$h(G_0) = 0$.

Statistical Analysis

We first evaluated the equations relating insulin sensitivity, insulin secretion, insulin

clearance, and blood glucose levels, which were given by the following equations:

$$\log k_4 k_5 = 1.98 \log k_7 + \log 0.0945, \qquad \text{Eq (1)}$$

$$\log k_4 k_5 = a \log \text{PG120} + b. \qquad \text{Eq (2)}$$

Eq (1) is a previously reported relation between insulin clearance and the product of insulin

sensitivity and secretion (23). Eq (2) is a relation between the plasma glucose concentration

at 120 min during the OGTT (PG120) and the product of insulin sensitivity and secretion. For

estimating the parameters a and b, RSS between variables was minimized with the use of a

nonlinear least squares technique, as described previously (23). After evaluating the

goodness-of-fit of these equations, we examined the goodness-of-fit of various other

equations to determine if a better equation existed. The goodness-of-fit was evaluated with

Pearson's correlation test (r). These values were reported with 95% confidence intervals

(CIs) by using bootstrap confidence intervals. The number of resamples performed to form

the bootstrap distribution was set at 10000. We also tested the relations among the variables

by standardized major axis regression (31).

193   Comparisons among NGT, IGT, and T2DM individuals were performed by the

194   Steel-Dwass test, a nonparametric test for multiple comparisons. A P value of <0.05 was

195   considered statistically significant. The association between indices was evaluated with

196   Spearman's correlation test. The correlation coefficients were reported with 95% CIs by

197   using bootstrap confidence intervals.

RESULTS

Derivation of a Conserved Equation for NGT, IGT, and T2DM from the Relations

Among Insulin Sensitivity, Insulin Secretion, Insulin Clearance, and Blood Glucose

Levels

We first investigated relations among insulin sensitivity, insulin secretion, insulin clearance, and blood glucose levels. We previously analyzed the relations using serum insulin and plasma glucose data obtained during consecutive hyperglycemic and hyperinsulinemic-euglycemic clamps in 47 NGT, 16 IGT, and 49 T2DM individuals (23), and we identified a relation regarding the three insulin-related parameters (23). This relation between insulin clearance and the product of insulin sensitivity and insulin secretion, was given by the following equation (23):

$$\log k_4 k_5 = 1.98 \log k_7 + \log 0.0945 \qquad \text{Eq (1)}$$

The indices k_4, k_5, and k_7, which correspond to insulin sensitivity, insulin secretion, and insulin clearance, respectively, were derived from the previous study (23).

We reevaluated the goodness-of-fit of the relation in individuals with each category of glucose tolerance. The coefficient of determination (R^2) of Eq (1) was estimated to be 0.26 for all subjects (Fig. S1A) and to be 0.28, 0.29, and –2.3 for NGT, IGT, and T2DM groups, respectively (Fig. S1B), indicating that the goodness-of-fit of Eq (1) was poor, especially for the T2DM group.

DI has been generally considered to reflect glucose disposal ability (2,32). The relation between DI and the blood glucose level can be represented as follows:

$$\log k_4 k_5 = a \log \text{PG120} + b \qquad \text{Eq (2)}$$

where the parameters a and b were estimated to minimize RSS between $\log k_4 k_5$ and $a \log \text{PG120} + b$, and PG120 indicates the plasma glucose concentration at 120 min during the OGTT, an important parameter for evaluation of glucose tolerance. Since the purpose of

this study was to investigate the relation among insulin secretion, insulin sensitivity, insulin

clearance, and blood glucose, DI was based on insulin secretion estimated by mathematical

models rather than the insulin concentration, which is affected by both insulin secretion and

clearance. Of note, the index defined in this way is also called adaptation index (33). The r

value of Eq (2) estimated for all subjects was 0.81 (Fig. 1A). However, r of Eq (2) for the

NGT, IGT, and T2DM groups was 0.41, 0.04, and 0.46, respectively (Fig. 1B), indicating that

the goodness-of-fit was particularly poor for IGT. The parameters a_n, a_i, and a_t estimated

from Eq (2) were –2.5, –0.24, and –1.8, respectively, and the parameters b_n, b_i, and b_t were

3.2, –1.7, and 1.2, respectively (Fig. 1B), suggesting that the relation between DI and blood

glucose levels changes with the progression of glucose intolerance. Collectively, these results

indicate that factors other than insulin secretion and insulin sensitivity may be involved in

glucose dynamics.

Given that the goodness-of-fit of Eq (1) and Eq (2) was poor, we attemted to show a

better equation. Empirically, we hypothesized that the following equation might have a better

fit:

$$\log k_4 k_5 = 2\log k_7 + a\log \text{PG120} + b \qquad \text{Eq (3)}$$

The parameters a and b were estimated to minimize RSS between $\log k_4 k_5$ and $2\log k_7 +$

$a\log \text{PG120} + b$. The r value of Eq (3) estimated for all subjects was 0.87 (Fig. 1C), which is

better than that of Eq (2). The r value of Eq (3) for the NGT, IGT, and T2DM groups was

0.60, 0.63, and 0.65, respectively (Fig. 1D). The parameters a_n, a_i, and a_t estimated from Eq

(3) were –1.4, –1.2, and –1.5, respectively, and the parameters b_n, b_i, and b_t were 1.7, 1.5,

and 1.7, respectively (Fig. 1D). Differences in r and in the parameters of Eq (3) among NGT,

IGT, and T2DM groups were smaller compared with those for Eq (2). These results indicated

that Eq (3) in general provides a better representation of blood glucose levels than does Eq

(2), especially for IGT. Of note, we here only indicated Eq (3) was one of the equations that

have a better fit than Eqs (1) and (2), which have been considered to have good fit, and did

not indicate that Eq (3) is the best. We further examined the characteristics of Eq (3) by

different mathematical models and statistical methods, with the results confirming the

validity of this equation (Appendix 1, Fig. S2, Fig. S3, and Fig. S4).

"DI/cle", a mearsure well correlated with blood glucose levels

Equation (3), which well describes the relation between the three insulin-related indices and

blood glucose levels among NGT, IGT, and T2DM individuals, can be transformed as

follows:

258
$$\mathrm{PG120} = 10^{-\frac{b}{a}} \left(\frac{k_4 k_5}{k_7^{\,2}}\right)^{\frac{1}{a}}$$
Eq (4)

This transformation separates blood glucose concentration and the insulin-related indices

onto the left and right sides. We designated this combination of the insulin-related indices on

the right side (insulin sensitivity multiplied by insulin secretion and divided by the square of

insulin clearance) as "DI/cle." The product of k_4 and k_5, which corresponds to DI, was

significantly lower in individuals with IGT or T2DM than in those with NGT (23), whereas

DI/cle was significantly lower in individuals with T2DM than in those with NGT or IGT but

was similar in the latter two groups (Fig. 2A). These results were validated by the values of

insulin sensitivity, insulin secretion, and insulin clearance estimated from the serum C-

peptide concentration (Appendix 1 and Fig. S5). Moreover, DI/cle can be approximated as the

extension of the basal insulin effect (29) to the insulin effect after glucose administration

(Appendix 2). Of note, the dimension of DI/cle is dL/mg, which is the same as the dimension

of the inverse of blood glucose levels. In addition, the sum of DI/cle and glucose effectiveness

determines blood glucose levels after glucose administration (Appendix 3). Although the

derivation from Eqs (1), (2), and (3) to Eq (4) is empirical and numerical, these analyses

analytically yielded similar results regarding the relation between $DI_{/cle}$ and blood glucose

levels, suggesting that $DI_{/cle}$ can literally be considered as the overall effect of insulin on

blood glucose levels.

Given that DI is the product of insulin secretion and insulin sensitivity, it is generally

represented as a hyperbolic curve with these two insulin-related parameters as the axes of

coordinates. The hyperbolic curves corresponding to the mean values of DI in NGT, IGT, and

T2DM subjects are shown in Figure 2B. On the other hand, $DI_{/cle}$ can be represented as a

hyperbolic curve with DI and 1/insulin clearance^2 as the axes of coordinates, given that $DI_{/cle}$

is the product of these two parameters. The hyperbolic curves corresponding to the mean

values of $DI_{/cle}$ in NGT, IGT, and T2DM subjects are shown in Figure 2C. In general, as

insulin sensitivity decreases, insulin secretion increases to maintain normal glucose

homeostasis (the phenomenon of moving up the hyperbolic curve) (2). The relation between

DI and 1/insulin clearance^2 can also be interpreted as moving up the curve, with 1/insulin

clearance^2 increasing as DI decreases. Of note, we previously found that k_7 (parameter for

insulin clearance) was significantly smaller in individuals with IGT or T2DM than in those

with NGT—that is, 1/insulin clearance^2 increased with the progression of glucose intolerance

(23).

We next investigated the relation between DI and $DI_{/cle}$. We therefore represented

$DI_{/cle}$ as a hyperbolic surface of insulin sensitivity, insulin secretion, and 1/insulin clearance^2

(Fig. 2D and E). The mean values of $DI_{/cle}$ in NGT, IGT, and T2DM subjects that are shown

as hyperbolic curves in Figure 2C are thus represented as hyperbolic surfaces in Figure 2D.

Figure 2E shows the relation between Figure 2B and Figure 2D (that is, the relation between

DI and $DI_{/cle}$), with DI corresponding to the intersection of the surface representing $DI_{/cle}$ and

the plane representing the mean value of 1/insulin clearance^2. The DI for IGT subjects was

significantly lower than that for NGT subjects (23), which can be interpreted as falling off the

hyperbolic curve (Fig. 2B). $DI_{/cle}$ was similar for NGT and IGT (Fig. 2A), however, with the

result that the corresponding hyperbolic curves (Fig. 2C) and hyperbolic surfaces (Fig. 2D)

were almost identical. The difference in DI between NGT and IGT is reflected as a change in

the intersection of the surface corresponding to $DI_{/cle}$ and the plane representing the mean

value of 1/insulin clearance^2 (Fig. 2E).

Of note, the phenomenon of moving up the curve of $DI_{/cle}$ for IGT represents almost

the same phenomenon as predicted in a previous modeling study using the βGI model (19),

indicating a pathway to failure of glucose homeostasis in which a change in insulin clearance

may alter glucose dynamics even when the normal glucose set point is maintained (Appendix

4).

Modified Mathematical Model of the Feedback Loop Linking Glucose and Insulin

We determined $DI_{/cle}$ by performing a hyperglycemic clamp and a hyperinsulinemic-

euglycemic clamp to measure insulin secretion and insulin sensitivity together with insulin

clearance (22,23,34), respectively. The performance of two clamp tests is laborious, however,

and so we attempted to estimate $DI_{/cle}$ from a single clamp.

Given that we were not able to estimate $DI_{/cle}$ with a single clamp analysis using our

previous model (23) (Appendix 5, Fig. S6, and Fig. S7A), we modified the model by

changing the positions of influx I and influx G as shown in Figure 3A (Appendix 6). We

simulated the time courses of blood glucose and insulin concentrations in the

hyperinsulinemic-euglycemic clamp analysis, with one example for an NGT subject being

shown in Figure 3B. To confirm that the modified model appropriately captured the

characteristics of the time-series data, we evaluated the consistency between the ISI

calculated from the actual measurements and that calculated from the simulation data. The

coefficient of determination (R^2) between measured ISI and simulated ISI was 0.96 (95% CI,

0.94 to 0.98) (Fig. 3C), indicating that the model was reasonably able to capture the essential

characteristics of the time-series data.

Prediction of DI/cle from only a hyperinsulinemic-euglycemic clamp

Some of the estimated parameters—including k_{m7}_ins, corresponding to insulin clearance

estimated from the modified model for the hyperinsulinemic-euglycemic clamp alone—

showed a tendency to diverge (Fig. 4A). However, k_{m5}/k_{m7}^2_ins was significantly

correlated with k_5/k_7^2_full ($r = 0.36$; 95% CI, 0.16 to 0.51), and $k_{m4}k_{m5}/k_{m7}^2$_ins was

significantly correlated with k_4k_5/k_7^2_full ($r = 0.52$; 95% CI, 0.37 to 0.64) (Fig. 4B and

Fig. S7B). Of note, insulin sensitivity, insulin secretion, and insulin clearance estimated from

the modified model for both the hyperglycemic clamp and the hyperinsulinemic-euglycemic

clamp (k_{m4}_full, k_{m5}_full, and k_{m7}_full, respectively) were significantly correlated with

those estimated from the original model (k_4_full, k_5_full, and k_7_full, respectively) (Fig. S8),

indicating that both models can be used to estimate the insulin-related parameters when both

the hyperglycemic and hyperinsulinemic-euglycemic clamps are performed. Thus, even with

this modified model, Eq (3) showed a better fit than Eq (2): the r value of Eq (2) for the

NGT, IGT, and T2DM groups was 0.03 (95% CI, –0.26 to 0.47), 0.25 (95% CI, –0.38 to

0.61), and 0.49 (95% CI, 0.24 to 0.66), respectively, and the r value of Eq (3) for the NGT,

IGT, and T2DM groups was 0.97 (95% CI, 0.83 to 0.99), 0.81 (95% CI, 0.34 to 0.94), and

0.66 (95% CI, 0.46 to 0.79), respectively.

We also attempted to estimate DI/cle using a simple analytically computable method.

We hypothesized that DI/cle can be analytically estimated as follows (Appendix 7):

345
$$\text{DI}_{/\text{cle}} = \frac{k_{\text{sen}}k_{\text{sec}}}{k_{\text{cle}}^2} \sim \frac{f_g I_0 (I_{SS} - I_0)}{f_i G_{SS} G_0 I_{SS}}$$

where k_{sen}, k_{sec}, and k_{cle} represent insulin sensitivity, insulin secretion, and insulin

clearance, respectively, and G_{SS}, G_0, I_{SS}, I_0, f_{g}, and f_{i} represent blood glucose level at the

steady state during insulin administration, blood glucose level in the fasting state, blood

insulin level at the steady state during insulin administration, blood insulin level in the fasting

state, glucose infusion rate at the steady state, and insulin infusion rate at the steady state,

respectively (Fig. 4C). This analytically calculated DI/cle ($k_{\text{sen}}k_{\text{sec}}/k_{\text{cle}}^2$) was significantly

correlated with $k_4 k_5/k_7^2$_full ($r = 0.67$; 95% CI, 0.52 to 0.77) (Fig. 4D). Of note,

$k_4 k_5/k_7^2$_full was also significantly correlated with DI/cle analytically estimated from a

hyperinsulinemic-euglycemic clamp alone by using the serum C-peptide concentration ($r =$

0.71; 95% CI, 0.58 to 0.80), OGTT ($r = 0.43$; 95% CI, 0.24 to 0.58), or a fasting blood test

($r = 0.54$; 95% CI, 0.38 to 0.68) (Appendix 1, Appendix 8, Fig. S9 and Fig. S10).

DISCUSSION

We have here derived an equation relating insulin sensitivity, insulin secretion, insulin

clearance, and blood glucose levels [Eq (3)] that was well conserved among individuals with

various levels of glucose tolerance, and we showed that this equation provides a better fit,

particularly for individuals with IGT, than does an equation relating only insulin sensitivity,

insulin secretion, and blood glucose levels [Eq (2)]. Insulin clearance is implicated in the

regulation of blood glucose concentration (8), and our present data also suggest that, together

with insulin sensitivity and insulin secretion, insulin clearance is important for such

regulation, especially in individuals with IGT. Our present results also suggest that DI/cle, an

index that corrects DI for insulin clearance, reflects glucose disposal ability, possibly by

representing the effect of insulin on blood glucose concentration.

DI/cle is represented as a hyperbolic curve of DI and 1/insulin clearance^2 or as a

hyperbolic surface of insulin sensitivity, insulin secretion, and 1/insulin clearance^2. The

deterioration of glucose tolerance is sometimes explained by the failure of insulin secretion to

increase sufficiently in compensation for a decline in insulin sensitivity, with this failure

being referred to as falling off the hyperbolic curve of DI (2,32). Our data now suggest that,

even when DI decreases, $DI_{/cle}$ is preserved if insulin clearance is also reduced. In general,

moving up the hyperbolic curve of DI eventually positions individuals at unstable extreme

points of the curve where the risk of disease onset increases (32), a phenomenon that has

been referred to as canalization and decanalization (2,32) or as glucose allostasis (35). Given

that individuals with IGT are at high risk of developing T2DM, the progression from NGT to

IGT and T2DM can be referred to as canalization and decanalization of the hyperbolic

surface of $DI_{/cle}$.

Our data also indicate that IGT encompasses the condition of decreased insulin

clearance with a relatively well preserved $DI_{/cle}$ and glucose set point (Appendix 4), and that

T2DM encompasses that of decreased $DI_{/cle}$ with a disrupted glucose set point. A modeling

study with simulated data indicated that changes in insulin clearance may affect glucose

dynamics even when the normal glucose set point is maintained, and can constitute a pathway

for progression of glucose intolerance (19). Our study based on mathematical models and

observational data now indicates that the phenomenon predicted by this previous model can

occur in IGT. Our findings are also consistent with those of a previous study indicating that

an increase in insulin secretion is followed by a decrease in first-pass hepatic insulin

clearance when insulin resistance is induced (15).

Despite the apparent analogy of the moving up–falling off phenomenon, whether the

body possesses the ability to reduce insulin clearance in response to a decline in DI and the

potential physiological mechanism underlying such a response remain unknown. However,

evidence suggests that changes in insulin clearance can be a determinant of glucose

intolerance (21,36). Whereas the direction of causation of decreased insulin clearance and

glucose intolerance remains unclear, a classification based on $DI_{/cle}$ may provide new insight

into the trajectories of glucose intolerance and T2DM.

Although these results were obtained from a limited number of individuals, the

results were validated by several different mathematical models. Moreover, we attempted to

estimate $DI_{/cle}$ from a hyperinsulinemic-euglycemic clamp alone, OGTT, or a fasting blood

test so that the characteristics of $DI_{/cle}$ can be validated more easily in the future. Whereas we

showed that $DI_{/cle}$ estimated from a hyperinsulinemic-euglycemic clamp test alone was

significantly correlated with that estimated from the two clamps, performance of even a

single clamp is labor-intensive. Moreover, the correlation between $DI_{/cle}$ estimated from the

two clamps and that estimated from the single clamp was only 0.52, 0.67 or 0.71. It will

therefore be important to develop more accurate and convenient methods for estimation of

$DI_{/cle}$ in order to validate its physiological relevance and clinical usefulness.

The current study has several limitations. For simplicity, we considered only insulin

sensitivity, insulin secretion, and insulin clearance. However, insulin secretion has basal,

first-phase, and second-phase components (37), and both insulin sensitivity and insulin

clearance have hepatic and peripheral components (3,22). Saturation of or time-dependent

changes in insulin clearance (6,22) were not considered. Although the models used in this

study possesses sufficient expressivity to fit the data from clamp tests (23) and IVGTT

(26,30), and to conclude the significance of incorporating insulin clearance in addition to

insulin secretion and insulin sensitivity, as well as a novel perspective on the progression of

glucose intolerance with the βGI model, it is important to devise a more comprehensive

measure that encompasses these more intricate effects in the future. Moreover, several better

approaches to evaluation of the hyperbolic relation have also been presented, and a power

function describing the relation between insulin sensitivity and insulin secretion gives a better

representation (38–40). Given that insulin sensitivity and insulin clearance are correlated

(41), the plane representing 1/insulin clearance^2 is not necessarily horizontal (Fig. 2E), and

the curve corresponding to DI also departs from a hyperbola when the plane departs from the

horizontal. Therefore, this study indicates that insulin clearance should be considered when

assessing the coefficient of disposition index, and future research on the coefficients of

insulin sensitivity, insulin secretion, and insulin clearance in the formula defining the effect

of insulin on blood glucose levels is warranted. In addition, the capacity for glucose disposal

is affected not only by insulin, but also by other hormones including glucagon as well as by

glucose (42,43). It is possible that differences in such factors may explain the glucose

dynamics of glucose tolerance. Further studies are also warranted to develop extended

equations that include glucose and glucagon to represent mechanisms of blood glucose

regulation.

In conclusion, we have here shown that DI$_{/cle}$, a new index that corrects DI by insulin

clearance, better reflects changes in glycemia compared with DI. By analogy with falling off

the hyperbolic curve of DI, the progression of glucose intolerance may be considered as

falling off the hyperbolic surface of DI$_{/cle}$. In theory, DI$_{/cle}$ represents the effect of insulin on

blood glucose levels, and it may serve as an indicator for the trajectory of changes in glucose

tolerance. Further study is warranted to understand the physiological and pathological

relevance of this new index.

Acknowledgments

We thank our laboratory members for critically reading this manuscript

Contribution statement. H.S. and K.H. analyzed the data. H.S., K.H., T.Y., K.S., W.O., and

S.K. wrote the manuscript. W.O. and S.K. supervised the study. S.K. is the guarantor of this

work and, as such, had full access to all the data in the study and takes responsibility for the

integrity of the data and the accuracy of the data analysis.

Conflict of Interest. The authors declare no competing interests.

Funding and Assistance. This study was supported by the Japan Society for the Promotion

of Science (JSPS) KAKENHI (JP21H04759), CREST, the Japan Science and Technology

Agency (JST) (JPMJCR2123), and The Uehara Memorial Foundation.

Data availability

The data set analyzed in the present study was described in our previous study (23).

Prior Presentation. A non–peer-reviewed version of this article was submitted to the

bioRxiv preprint server (https://doi.org/10.1101/2022.12.28.522156) on 30 December 2022.

References

1. **Bergman RN.** Origins and History of the Minimal Model of Glucose Regulation. *Front. Endocrinol.* 2020;11:583016.

2. **Kodama K, Tojjar D, Yamada S, Toda K, Patel CJ, Butte AJ.** Ethnic differences in the relationship between insulin sensitivity and insulin response: a systematic review and meta-analysis. *Diabetes Care* 2013;36(6):1789–1796.

3. **Ha J, Sherman A.** Type 2 diabetes: one disease, many pathways. *Am. J. Physiol. Endocrinol. Metab.* 2020;319(2):E410–E426.

4. **De Gaetano A, Hardy TA.** A novel fast-slow model of diabetes progression: Insights into mechanisms of response to the interventions in the Diabetes Prevention Program. *PLoS One* 2019;14(10):e0222833.

5. **Topp B, Promislow K, Devries G, Miura RM, Finegood DT.** A Model of β -Cell Mass, Insulin, and Glucose Kinetics: Pathways to Diabetes. *J. Theor. Biol.* 2000;206(4):605–619.

6. **Mari A, Tura A, Grespan E, Bizzotto R.** Mathematical Modeling for the Physiological and Clinical Investigation of Glucose Homeostasis and Diabetes. *Front. Physiol.* 2020;11:575789.

7. **Jones CN, Pei D, Staris P, Polonsky KS, Chen YD, Reaven GM.** Alterations in the glucose-stimulated insulin secretory dose-response curve and in insulin clearance in nondiabetic insulin-resistant individuals. *J. Clin. Endocrinol. Metab.* 1997;82(6):1834–1838.

8. **Kim SH, Reaven GM.** Insulin clearance: an underappreciated modulator of plasma insulin concentration. *J. Investig. Med.* 2016;64(7):1162–1165.

9. **Goodarzi MO, Cui J, Chen YDI.** Fasting insulin reflects heterogeneous physiological processes: role of insulin clearance. *Am. J. Physiol. Endocrinol. Metab.* 2011;301(2):E402–E408.

10. **de Oliveira DM, Tura A, Vasques ACJ, Camilo DF, Lima MM, de Lemos-Marini SHV, Goncalves EM, Guerra-Junior G, Geloneze B.** Insulin Resistance in Congenital Adrenal Hyperplasia is Compensated for by Reduced Insulin Clearance. *J. Clin. Endocrinol. Metab.* 2021;106(4):e1574–e1585.

11. **Mittelman SD, Van Citters GW, Kim SP, Davis DA, Dea MK, Hamilton-Wessler M, Bergman RN.** Longitudinal compensation for fat-induced insulin resistance includes reduced insulin clearance and enhanced beta-cell response. *Diabetes* 2000;49(12):2116–2125.

12. **Gastaldelli A, Abdul Ghani M, DeFronzo RA.** Adaptation of Insulin Clearance to Metabolic Demand Is a Key Determinant of Glucose Tolerance. *Diabetes* 2021;70(2):377–385.

13. **Haffner SM, Stern MP, Watanabe RM, Bergman RN.** Relationship of insulin
clearance and secretion to insulin sensitivity in non-diabetic Mexican Americans. *Eur. J.*
Clin. Invest. 1992;22(3):147–153.

14. **Galderisi A, Polidori D, Weiss R, Giannini C, Pierpont B, Tricò D, Caprio S.** Lower
Insulin Clearance Parallels a Reduced Insulin Sensitivity in Obese Youths and Is
Associated With a Decline in β-Cell Function Over Time. *Diabetes* 2019;68(11):2074–
2084.

15. **Kim SP, Ellmerer M, Kirkman EL, Bergman RN.** β-Cell "rest" accompanies reduced
first-pass hepatic insulin extraction in the insulin-resistant, fat-fed canine model.
American Journal of Physiology-Endocrinology and Metabolism 2007;292(6):E1581–
E1589.

16. **Bizzotto R, Tricò D, Natali A, Gastaldelli A, Muscelli E, De Fronzo RA, Arslanian**
S, Ferrannini E, Mari A. New Insights on the Interactions Between Insulin Clearance
and the Main Glucose Homeostasis Mechanisms. *Diabetes Care* 2021;44(9):2115–2123.

17. **DeFronzo RA, Tripathy D, Abdul-Ghani M, Musi N, Gastaldelli A.** The disposition
index does not reflect β-cell function in IGT subjects treated with pioglitazone. *J. Clin.*
Endocrinol. Metab. 2014;99(10):3774–3781.

18. **Ladwa M, Bello O, Hakim O, Shojaee-Moradie F, Boselli L, Charles-Edwards G,**
Stadler M, Peacock JL, Umpleby AM, Amiel SA, Bonadonna RC, Goff LM. Insulin
clearance as the major player in the hyperinsulinaemia of black African men without
diabetes. *Diabetes Obes. Metab.* 2020;22(10):1808–1817.

19. **Karin O, Swisa A, Glaser B, Dor Y, Alon U.** Dynamical compensation in
physiological circuits. *Mol. Syst. Biol.* 2016;12(11):886.

20. **Galderisi A, Giannini C, Weiss R, Kim G, Shabanova V, Santoro N, Pierpont B,**
Savoye M, Caprio S. Trajectories of changes in glucose tolerance in a multiethnic
cohort of obese youths: an observational prospective analysis. *Lancet Child Adolesc*
Health 2018;2(10):726–735.

21. **Bergman RN, Piccinini F, Kabir M, Kolka CM, Ader M.** Hypothesis: Role of
Reduced Hepatic Insulin Clearance in the Pathogenesis of Type 2 Diabetes. *Diabetes*
2019;68(9):1709–1716.

22. **Piccinini F, Bergman RN.** The Measurement of Insulin Clearance. *Diabetes Care*
2020;43(9):2296–2302.

23. **Ohashi K, Komada H, Uda S, Kubota H, Iwaki T, Fukuzawa H, Komori Y, Fujii**
M, Toyoshima Y, Sakaguchi K, Ogawa W, Kuroda S. Glucose Homeostatic Law:
Insulin Clearance Predicts the Progression of Glucose Intolerance in Humans. *PLoS One*
2015;10(12):e0143880.

24. **Ohashi K, Fujii M, Uda S, Kubota H, Komada H, Sakaguchi K, Ogawa W, Kuroda**
S. Increase in hepatic and decrease in peripheral insulin clearance characterize abnormal

temporal patterns of serum insulin in diabetic subjects. *npj Systems Biology and*
Applications 2018;4(1):1–12.

25. **Cobelli C, Dalla Man C.** Minimal and Maximal Models to Quantitate Glucose
Metabolism: Tools to Measure, to Simulate and to Run in Silico Clinical Trials. *J.*
Diabetes Sci. Technol. 2021:19322968211015268.

26. **Van Cauter E, Mestrez F, Sturis J, Polonsky KS.** Estimation of insulin secretion rates
from C-peptide levels. Comparison of individual and standard kinetic parameters for C-
peptide clearance. *Diabetes* 1992;41(3):368–377.

27. **DeFronzo RA, Tobin JD, Andres R.** Glucose clamp technique: a method for
quantifying insulin secretion and resistance. *Am. J. Physiol.* 1979;237(3):E214-23.

28. **Okuno Y, Komada H, Sakaguchi K, Nakamura T, Hashimoto N, Hirota Y, Ogawa**
W, Seino S. Postprandial serum C-peptide to plasma glucose concentration ratio
correlates with oral glucose tolerance test- and glucose clamp-based disposition indexes.
Metabolism 2013;62(10):1470–1476.

29. **Kahn SE, Bergman RN, Schwartz MW, Taborsky GJ Jr, Porte D Jr.** Short-term
hyperglycemia and hyperinsulinemia improve insulin action but do not alter glucose
action in normal humans. *Am. J. Physiol.* 1992;262(4 Pt 1):E518-E523.

30. **De Gaetano A, Arino O.** Mathematical modelling of the intravenous glucose tolerance
test. *J. Math. Biol.* 2000;40(2):136–168.

31. **Warton DI, Duursma RA, Falster DS, Taskinen S.** smatr 3 - an R package for
estimation and inference about allometric lines. *Methods Ecol. Evol.* 2012;3(2):257–259.

32. **Gibson G.** Decanalization and the origin of complex disease. *Nat. Rev. Genet.*
2009;10(2):134–140.

33. **Ahrén B, Pacini G.** Impaired adaptation of first-phase insulin secretion in
postmenopausal women with glucose intolerance. *Am. J. Physiol.* 1997;273(4):E701-
E707.

34. **Sjaarda L, Lee S, Tfayli H, Bacha F, Bertolet M, Arslanian S.** Measuring β-cell
function relative to insulin sensitivity in youth: does the hyperglycemic clamp suffice?
Diabetes Care 2013;36(6):1607–1612.

35. **Stumvoll M, Tataranni PA, Stefan N, Vozarova B, Bogardus C.** Glucose allostasis.
Diabetes 2003;52(4):903–909.

36. **Sugiyama S, Jinnouchi H, Hieshima K, Kurinami N, Jinnouchi K, Yoshida A,**
Suzuki T, Kajiwara K, Miyamoto F, Jinnouchi T. Potential identification of type 2
diabetes with elevated insulin clearance. *NEJM Evidence* 2022;1(4):EVIDoa2100052.

37. **Cobelli C, Toffolo GM, Man CD, Campioni M, Denti P, Caumo A, Butler P, Rizza**
R. Assessment of β-cell function in humans, simultaneously with insulin sensitivity and
hepatic extraction, from intravenous and oral glucose tests. *American Journal of*
Physiology-Endocrinology and Metabolism 2007;293(1):E1–E15.

38. **Denti P, Toffolo GM, Cobelli C.** The disposition index: from individual to population
approach. *Am. J. Physiol. Endocrinol. Metab.* 2012;303(5):E576-E586.
39. **Ferrannini E, Mari A.** Beta cell function and its relation to insulin action in humans: a
critical appraisal. *Diabetologia* 2004;47(5):943–956.
40. **Cobelli C, Dalla Man C, Toffolo G, Basu R, Vella A, Rizza R.** The oral minimal
model method. *Diabetes* 2014;63(4):1203–1213.
41. **Lorenzo C, Hanley AJG, Wagenknecht LE, Rewers MJ, Stefanovski D, Goodarzi
MO, Haffner SM.** Relationship of insulin sensitivity, insulin secretion, and adiposity
with insulin clearance in a multiethnic population: the insulin Resistance Atherosclerosis
study. *Diabetes Care* 2013;36(1):101–103.
42. **Dube S, Errazuriz-Cruzat I, Basu A, Basu R.** The forgotten role of glucose
effectiveness in the regulation of glucose tolerance. *Curr. Diab. Rep.* 2015;15(6):605.
43. **Lee YH, Wang M-Y, Yu X-X, Unger RH.** Glucagon is the key factor in the
development of diabetes. *Diabetologia* 2016;59(7):1372–1375.
581

Figure Legends

Figure 1. Relation between insulin-related parameters and blood glucose levels. (A) Scatter

plot for $a_a \log PG120 + b_a$ versus $\log k_4 k_5$ (Eq (2)). Each point corresponds to the values for

a single subject. Green circles, red crosses, and blue squares indicate NGT, IGT, and T2DM

subjects, respectively. r is Pearson's correlation coefficient, and the value in the parenthesis

is 95% confidence interval. The parameters a_a and b_a were estimated to minimize RSS

between $a_a \log PG120 + b_a$ and $\log k_4 k_5$ in all subjects. (B) Scatter plots for $a \log PG120 +$

b versus $\log k_4 k_5$ in NGT, IGT, and T2DM subjects (Eq (2)). The parameters a_n and b_n, a_i

and b_i, and a_t and b_t were estimated to minimize RSS between $a \log PG120 + b$ and

$\log k_4 k_5$ in NGT, IGT, and T2DM subjects, respectively. (C) Scatter plot for $2 \log k_7 +$

$a_a \log PG120 + b_a$ versus $\log k_4 k_5$ (Eq (3)). The parameters a_a and b_a were estimated to

minimize RSS between $2 \log k_7 + a_a \log PG120 + b_a$ and $\log k_4 k_5$ in all subjects. (D)

Scatter plots for $2 \log k_7 + a \log PG120 + b$ versus $\log k_4 k_5$ in NGT, IGT, and T2DM

subjects (Eq (3)). The parameters a_n and b_n, a_i and b_i, and a_t and b_t were estimated to

minimize RSS between $2 \log k_7 + a \log PG120 + b$ and $\log k_4 k_5$ in NGT, IGT, and T2DM

subjects, respectively. The indices k_4, k_5, and k_7, which correspond to insulin sensitivity,

insulin secretion, and insulin clearance, respectively, were calculated by a previously

described method (23), and PG120 is the plasma glucose concentration at 120 min during an

OGTT.

Figure 2. Relation between DI and DI$_{/cle}$. (A) Box plots of log DI$_{/cle}$, corresponding to the

logarithm of $k_4 k_5 / k_7{}^2$, in NGT, IGT, and T2DM subjects. Each point corresponds to the

value for a single subject. The boxes denote the median and upper and lower quartiles. *P*

values were determined by the Steel-Dwass test. (B) Scatter plots for k_4 and k_5 fitted with

hyperbolic curves (based on the function $xy = $ constant), and kernel density plots. The

hyperbolic curves correspond to the mean value of DI for NGT, IGT, or T2DM subjects.

Each point corresponds to the values for a single subject, with green, red, and blue indicating

NGT, IGT, and T2DM, respectively. The bandwidth of each kernel is based on the rule-of-

thumb bandwidth. (C) Scatter plots for $k_4 k_5$ and $1/k_7{}^2$ fitted with hyperbolic curves (based

on the function $xy = $ constant), and kernel density plots. The hyperbolic curves correspond

to the mean value of DI/cle in NGT, IGT, or T2DM subjects. Each point corresponds to the

values for a single subject, with green, red, and blue indicating NGT, IGT, and T2DM,

respectively. The bandwidth of each kernel is based on the rule-of-thumb bandwidth. (D)

Hyperbolic surfaces (based on the function $xyz = $ constant) corresponding to the mean

value of DI/cle in NGT, IGT, or T2DM subjects. (E) Hyperbolic surfaces corresponding to the

mean value of DI/cle as well as planes corresponding to the mean value of 1/insulin clearance^2

in NGT, IGT, or T2DM subjects. The axes correspond to k_4, k_5, and $1/k_7{}^2$ as in (D), but the

label for k_5 was omitted for clarity. The intersections (solid lines) of the surfaces representing

DI/cle and the planes representing the mean value of 1/insulin clearance^2 correspond to the

hyperbolic curves in (B).

Figure 3. Modified mathematical model of the feedback loop that links glucose and insulin.

(A) Model diagram. The letters within circles indicate the variables of the model, the arrows

denote the fluxes, and the lines ending in circles or bars represent activation or inactivation,

 respectively. ϕ indicates a fixed value. The variables G and I represent blood glucose and

 insulin concentrations, respectively. The variable Y corresponds to the effective glucose

 concentration for induction of variable X, which represents insulin secreted from pancreatic

 β-cells. (B) Time courses of blood glucose, blood insulin, infused glucose, and infused

 insulin during hyperinsulinemic-euglycemic clamp analysis for an NGT subject. The blue

 lines indicate the simulated patterns, and the red circles indicate the measured time course

 data. The simulated values of each variable were rescaled to absolute concentration and

 plotted. (C) Scatter plot for simulated ISI versus measured ISI. Each point corresponds to the

 values for a single subject, with green circles, red crosses, and blue squares indicating NGT,

 IGT, and T2DM subjects, respectively. The measured and simulated ISI values were

 calculated from the normalized time course and plotted. R^2 is the coefficient of

 determination, and the value in the parenthesis is 95% confidence interval.

Figure 4. Relation between parameters estimated from a hyperinsulinemic-euglycemic clamp

alone and those estimated from both hyperglycemic and hyperinsulinemic-euglycemic

clamps. (A) Estimated parameters of the modified model based on a hyperinsulinemic-

euglycemic clamp alone. (B) Spearman's correlation coefficients. Bars indicate 95%

confidence intervals. The parameters k_4_full, k_5_full, k_7_full, $k_5/k_7{}^2$_full, and $k_4k_5/k_7{}^2$_full

correspond to insulin sensitivity, insulin secretion, insulin clearance, the ratio of insulin

secretion to the square of insulin clearance, and DI$_{\text{cle}}$ estimated from both hyperglycemic and

hyperinsulinemic-euglycemic clamps by the previously described mathematical model (23)

(Fig. S6A), respectively. The parameters k_{m4}_ins, k_{m5}_ins, k_{m7}_ins, $k_{\text{m5}}/k_{\text{m7}}{}^2$_ins, and

$k_{\text{m4}}k_{\text{m5}}/k_{\text{m7}}{}^2$_ins correspond to insulin sensitivity, insulin secretion, insulin clearance, the

ratio of insulin secretion to the square of insulin clearance, and DI$_{\text{cle}}$ estimated from the

hyperinsulinemic-euglycemic clamp alone by the modified mathematical model (Fig. 3A),

respectively. (C) Simulated time courses of blood glucose, blood insulin, infused glucose, and

infused insulin during a hyperinsulinemic-euglycemic clamp. G_{SS}, blood glucose level at the

steady state during insulin administration; G_0, blood glucose level in the fasting state; I_{SS},

blood insulin level at the steady state during insulin administration; I_0, blood insulin level in

the fasting state; f_{g}, glucose infusion rate at the steady state; and f_{i}, insulin infusion rate at the

steady state. (D) Scatter plot for $\log k_{\text{sen}}k_{\text{sec}}/k_{\text{cle}}{}^2$ versus $\log k_4k_5/k_7{}^2$_full. The parameters

$k_{\text{sen}}k_{\text{sec}}/k_{\text{cle}}{}^2$ and $k_4k_5/k_7{}^2$_full correspond to DI$_{\text{cle}}$ analytically estimated from the

hyperinsulinemic-euglycemic clamp alone (Appendix 7) and that estimated from the

hyperglycemic and hyperinsulinemic-euglycemic clamps by the previously described

mathematical model (23) (Fig. S6A), respectively. Each point corresponds to the values for a

single subject, with green circles, red crosses, and blue squares indicating NGT, IGT, and

T2DM subjects, respectively. r is Spearman's correlation coefficient, and the value in the

parenthesis is 95% confidence interval.