Insulin Effectiveness, a Measure Consisting of Insulin Sensitivity, Secretion, and Clearance, Captures Differences Between Diabetic and Nondiabetic States

Hikaru Sugimoto,[1] Ken-ichi Hironaka,[2] Tomoko Yamada,[3] Kazuhiko Sakaguchi,[3] Wataru Ogawa,[3]* and Shinya Kuroda[1,2]*

[1]Department of Biochemistry and Molecular Biology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[2]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

[3]Division of Diabetes and Endocrinology, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kusunoki-cho 7-5-1, Chuo-ku, Kobe, Hyogo 650-0017, Japan.

*Corresponding authors: Wataru Ogawa, ogawa@med.kobe-u.ac.jp; or Shinya Kuroda, skuroda@bs.s.u-tokyo.ac.jp

OBJECTIVE

Insulin clearance is implicated in regulation of glucose homeostasis independently of insulin sensitivity and insulin secretion. To understand the relation between blood glucose levels and insulin sensitivity, insulin secretion, and insulin clearance, we performed a mathematical analysis.

RESEARCH DESIGN AND METHODS

We performed a hyperglycemic clamp, a hyperinsulinemic-euglycemic clamp, and an oral glucose tolerance test in 47, 16, and 49 subjects with normal glucose tolerance (NGT), impaired glucose tolerance (IGT), and type 2 diabetes mellitus (T2DM), respectively.

RESULTS

The disposition index (DI), defined as the product of insulin sensitivity and insulin secretion and regarded as a measure of the capacity for glucose handling, did not well explain blood glucose levels at the early stage of glucose intolerance. However, an equation relating DI, insulin clearance, and blood glucose levels was well conserved regardless of the extent of glucose intolerance. As a measure of the effect of insulin, we developed an index, designated "insulin effectiveness," that is based on this equation and corresponds to DI divided by the square of insulin clearance. Insulin effectiveness was not impaired in IGT compared with NGT, possibly as a result of a decrease in insulin clearance in response to a reduction in DI, whereas it was impaired in T2DM relative to IGT.

CONCLUSIONS

Insulin effectiveness, which likely reflects insulin action in the body, well represents changes in glycemia and is impaired during the transition from IGT to T2DM. It may serve as a new indicator for the trajectory of changes in glucose tolerance.

The blood glucose concentration is maintained within a narrow range by feedback-compensatory responses in the body. Insulin sensitivity and insulin secretion are linked by such a relation, with insulin secretion increasing to maintain normal glucose homeostasis as insulin sensitivity declines (1). This compensatory response has been regarded as "moving up" the hyperbolic curve of the product of insulin sensitivity and insulin secretion, and the failure of this response, regarded as "falling off" the hyperbolic curve, is associated with the transition from normal glucose tolerance (NGT) to impaired glucose tolerance (IGT) and type 2 diabetes mellitus (T2DM) (1). The disposition index (DI), generally defined as the product of insulin sensitivity and insulin secretion, has therefore been adopted as a measure of glucose handling in the body (1,2).

Several models that explain this moving up–falling off phenomenon as an interplay among insulin sensitivity, insulin secretion, and pancreatic β-cell mass have been described (3–6). Such models account for the physiological features and pathological trajectory of glucose tolerance—including NGT, IGT, and T2DM—to a certain extent, but their conformity to clinical or experimental data is not high (6). Moreover, the moving up and falling off the curve of the product of insulin sensitivity and insulin secretion appear not to adequately represent the relation between blood glucose and insulin levels. For example, comparison of individuals with NGT at extremes of insulin sensitivity revealed that an approximately twofold increase in blood insulin levels was associated with an increase in insulin secretion of <50% (7,8), indicating that the increase in insulin secretion may not fully explain the increase in circulating insulin levels in individuals with insulin resistance (8).

Insulin clearance is implicated as a determinant of circulating insulin levels that is independent of insulin secretion (8–16). Insulin clearance likely contributes to the compensatory responses of the body, with a reduction in insulin clearance in the insulin-resistant state resulting in an increase in circulating insulin levels and a lessening of the

demand for insulin secretion by β-cells (11). In some instances, changes in the circulating

insulin levels do not reflect changes in insulin secretion, instead being primarily influenced

by insulin clearance (17). Furthermore, changes in insulin clearance have been implicated in

the pathogenesis of glucose intolerance (18–21). Whereas these various findings suggest the

importance of insulin clearance in the physiology and pathology of glucose homeostasis, it is

difficult to directly evaluate this parameter from clinical data (22).

Mathematical models provide insight into phenomena in which various factors

mutually interact and for which effects of such factors cannot be measured directly (22–25).

We previously developed a mathematical model based on temporal changes in circulating

glucose and insulin levels in human subjects during consecutive hyperglycemic and

hyperinsulinemic-euglycemic clamps (23). Using this model, we estimated insulin clearance

and characterized its role in the regulation of glucose homeostasis, and we identified a

square-law relation between insulin clearance and the product of insulin sensitivity and

insulin secretion that was conserved among individuals with NGT, IGT, or T2DM (23).

We have now further examined the relation among insulin sensitivity, insulin

secretion, and insulin clearance, and we here propose a new index, "insulin effectiveness,"

that consists of DI corrected by insulin clearance. We have characterized insulin effectiveness

in subjects with various levels of glucose tolerance, and we provide evidence that this new

index can serve as an indicator of the effect of insulin on glucose homeostasis and can

capture the transition from IGT to T2DM.

RESEARCH DESIGN AND METHODS

Subjects and Measurements

We analyzed a previously reported data set (23) in the present study. In brief, a

hyperglycemic clamp was performed by intravenous infusion first of a bolus of glucose over

15 min and then of a variable amount of glucose to maintain the plasma glucose level at 200

mg/dL. Ten minutes after the end of the hyperglycemic clamp, a hyperinsulinemic-

euglycemic clamp was performed by intravenous infusion of insulin with a target plasma

glucose level of 90 mg/dL or of the fasting level in the case of NGT and IGT subjects whose

fasting plasma glucose level was <90 mg/dL. These clamp analyses and a 75-g oral glucose

tolerance test (OGTT) were performed within a period of 10 days.

The indices k_4, k_5, and k_7, which correspond to insulin sensitivity, secretion, and

clearance, respectively, were calculated with a mathematical model described in our previous

study (23). The insulin sensitivity index (ISI) and incremental area under the serum insulin

concentration curve from 0 to 10 min during the hyperglycemic clamp (AUC_IRI) were also

calculated as described previously (23,26,27).

Fifty NGT, 18 IGT, and 53 T2DM subjects were initially used for parameter

estimation. As we previously described (23), three NGT, two IGT, and three T2DM

individuals were subsequently eliminated as outliers. In addition, one T2DM individual was

eliminated because of missing values, with the remaining 47 NGT, 16 IGT, and 49 T2DM

subjects being analyzed.

Statistical Analysis

Comparisons among NGT, IGT, and T2DM subjects were performed by ANOVA followed

by Tukey's honestly significant difference test. A *P* value of <0.05 was considered

statistically significant. For examination of the relations among insulin sensitivity, insulin

secretion, insulin clearance, and plasma glucose levels, the residual sum of the square (RSS)

between variables was minimized with the use of a nonlinear least squares technique, as

described previously (23). We also tested the relations among the variables by standardized

major axis regression (28). The association between indices was evaluated with Spearman's

correlation test.

Data and Resource Availability

The data set analyzed in the present study was described in our previous study (23).

RESULTS

Derivation of a Conserved Equation for NGT, IGT, and T2DM from the Relations

Among Insulin Sensitivity, Insulin Secretion, Insulin Clearance, and Blood Glucose

Levels

We first attempted to derive a relation for the regulation of blood glucose levels that is

conserved across various levels of glucose tolerance. We previously analyzed the relations

among insulin sensitivity, insulin secretion, insulin clearance, and blood glucose levels using

serum insulin and plasma glucose data obtained during consecutive hyperglycemic and

hyperinsulinemic-euglycemic clamps in NGT, IGT, and T2DM individuals (23), and we

identified a law regarding the three insulin-related parameters (23). This law, a square-law

relation between insulin clearance and the product of insulin sensitivity and insulin secretion,

was given by the following equation (23):

$$\log k_4 k_5 = 1.98\log k_7 + \log 0.0945 \qquad \text{Eq (1)}$$

The indices k_4, k_5, and k_7, which correspond to insulin sensitivity, secretion, and clearance,

respectively, were derived from the previous study (23).

We reevaluated the fitness of the law in individuals with each category of glucose

tolerance. The coefficient of determination (R^2) of Eq (1) was estimated to be 0.26 for all

subjects (Fig. S1A) and to be 0.28, 0.29, and –2.3 for NGT, IGT, and T2DM groups,

respectively (Fig. S1B), indicating that the fitness of Eq (1) was poor, especially for the

T2DM group. Given that a high concentration of blood glucose is a characteristic of this latter

group, it is possible that Eq (1) needs to be modified by blood glucose level.

The DI (product of insulin secretion and insulin sensitivity) has been generally

considered to reflect glucose disposal ability (1,2). The relation between DI and the blood

glucose level can be represented as follows (29):

$$\log k_4 k_5 = a \log \text{PG120} + b \qquad\qquad \text{Eq (2)}$$

where the parameters a and b were estimated to minimize RSS between $\log k_4 k_5$ and

$a \log \text{PG120} + b$, and PG120 indicates the plasma glucose concentration at 120 min during

the OGTT, an important parameter for evaluation of glucose tolerance. The parameters a and

b estimated for all subjects are represented as a_a and b_a, respectively. The R^2 value of Eq (2)

estimated for all subjects was 0.66 (Fig. 1A), which is better than that of Eq (1) (Fig. S1A).

However, R^2 of Eq (2) for the NGT, IGT, and T2DM groups was 0.17, 0.00, and 0.22,

respectively (Fig. 1B), indicating that the goodness-of-fit was particularly poor for IGT. The

parameters a_n and b_n, a_i and b_i, and a_t and b_t were estimated to minimize RSS between

$\log k_4 k_5$ and $a \log \text{PG120} + b$ in NGT, IGT, and T2DM individuals, respectively. The

parameters a_n, a_i, and a_t estimated from Eq (2) were –2.5, –0.24, and –1.8, respectively, and

the parameters b_n, b_i, and b_t were 3.2, –1.7, and 1.2, respectively (Fig. 1B), suggesting that

the relation between DI and blood glucose levels changes with the progression of glucose

intolerance. Given that insulin clearance is likely a determinant of circulating insulin levels

that is independent of insulin secretion (8–11), we hypothesized that the addition of insulin

clearance might improve the representation of blood glucose, in particular for individuals

with IGT.

Given that Eq (1) likely needs to be corrected for blood glucose level and that Eq (2)

might be improved by adding insulin clearance, we modified these equations by integrating

them as follows:

$$167 \qquad \log k_4 k_5 = 2\log k_7 + a\log \mathrm{PG120} + b \qquad\qquad \text{Eq (3)}$$

The parameters a and b were estimated to minimize RSS between $\log k_4 k_5$ and $2\log k_7 +$

$a\log \mathrm{PG120} + b$. The R^2 value of Eq (3) estimated for all subjects was 0.76 (Fig. 1C), which

is better than that of Eq (1) or Eq (2) (Fig. S1A and Fig. 1A). It is also better than that of

another equation that connects insulin clearance and the product of insulin sensitivity and

insulin secretion (Fig. S1C). The R^2 value of Eq (3) for the NGT, IGT, and T2DM groups

was 0.35, 0.36, and 0.43, respectively (Fig. 1D). The parameters a_n, a_i, and a_t estimated

from Eq (3) were –1.4, –1.2, and –1.5, respectively, and the parameters b_n, b_i, and b_t were

1.7, 1.5, and 1.7, respectively (Fig. 1D). Differences in R^2 and in the parameters of Eq (3)

among NGT, IGT, and T2DM groups were smaller compared with those for Eq (1) and Eq

(2). These results indicated that Eq (3) in general provides a better representation of blood

glucose levels than does Eq (2), especially for IGT. We further examined the characteristics

of Eq (3) by different methods, with the results confirming the validity of this equation

(Appendix 1, Fig. S2, Fig. S3, and Fig. S4).

"Insulin Effectiveness" as a Glucose Homeostatic Law–Derived Index

Equation (3), which well describes the relation between the three insulin-related indices and

blood glucose levels among NGT, IGT, and T2DM individuals, can be transformed as

follows:

$$186 \qquad \mathrm{PG120} = 10^{-\frac{b}{a}} \left(\frac{k_4 k_5}{k_7{}^2} \right)^{\frac{1}{a}} \qquad\qquad \text{Eq (4)}$$

This transformation separates blood glucose concentration and the insulin-related indices

onto the left and right sides. We designated this combination of the insulin-related indices on

the right side (insulin sensitivity multiplied by insulin secretion and divided by the square of

insulin clearance) as "insulin effectiveness." The product of k_4 and k_5, which corresponds to

DI, was significantly lower in individuals with IGT or T2DM than in those with NGT (23),

whereas insulin effectiveness was significantly lower in individuals with T2DM than in those

with NGT or IGT but was similar in the latter two groups (Fig. 2A). These results were

validated by the values of insulin sensitivity, secretion, and clearance estimated from the

serum C-peptide concentration (Appendix 1 and Fig. S5). Moreover, insulin effectiveness can

be approximated as the extension of the basal insulin effect (30) to the insulin effect after

glucose administration (Appendix 2). In addition, the sum of insulin effectiveness and

glucose effectiveness determines blood glucose levels after glucose administration (Appendix

3), suggesting that insulin effectiveness can literally be considered as the overall effect of

insulin on blood glucose levels.

Given that DI is the product of insulin secretion and insulin sensitivity, it is generally

represented as a hyperbolic curve with these two insulin-related parameters as the axes of

coordinates. The hyperbolic curves corresponding to the mean values of DI in NGT, IGT, and

T2DM subjects are shown in Figure 2B. In general, as insulin sensitivity decreases, insulin

secretion increases to maintain normal glucose homeostasis (the phenomenon of moving up

the hyperbolic curve) (2). On the other hand, insulin effectiveness can be represented as a

hyperbolic curve with DI and 1/insulin clearance^2 as the axes of coordinates, given that

insulin effectiveness is the product of these two parameters. The hyperbolic curves

corresponding to the mean values of insulin effectiveness in NGT, IGT, and T2DM subjects

are shown in Figure 2C. The relation between DI and 1/insulin clearance^2 can also be

interpreted as moving up the curve, with 1/insulin clearance^2 increasing as DI decreases. Of

note, we previously found that k_7 (parameter for insulin clearance) was significantly smaller

in individuals with IGT or T2DM than in those with NGT—that is, 1/insulin clearance^2

increased with the progression of glucose intolerance (23).

We next investigated the relation between DI and insulin effectiveness. We therefore

represented insulin effectiveness as a hyperbolic surface of insulin sensitivity, insulin

secretion, and 1/insulin clearance[2] (Fig. 2D and E). The mean values of insulin effectiveness

in NGT, IGT, and T2DM subjects that are shown as hyperbolic curves in Figure 2C are thus

represented as hyperbolic surfaces in Figure 2D. Figure 2E shows the relation between Figure

2B and Figure 2D (that is, the relation between DI and insulin effectiveness), with DI

corresponding to the intersection of the surface representing insulin effectiveness and the

plane representing the mean value of 1/insulin clearance[2]. The DI for IGT subjects was

significantly lower than that for NGT subjects (23), which can be interpreted as falling off the

hyperbolic curve (Fig. 2B). Insulin effectiveness was similar for NGT and IGT (Fig. 2A),

however, with the result that the corresponding hyperbolic curves (Fig. 2C) and hyperbolic

surfaces (Fig. 2D) were almost identical. The difference in DI between NGT and IGT is

reflected as a change in the intersection of the surface corresponding to insulin effectiveness

and the plane representing the mean value of 1/insulin clearance[2] (Fig. 2E).

Of note, the phenomenon of moving up the curve of insulin effectiveness for IGT

represents almost the same phenomenon as predicted in a previous modeling study (19),

indicating a pathway to failure of glucose homeostasis in which a change in insulin clearance

may alter glucose dynamics even when the normal glucose set point is maintained (Appendix

4).

Modified Mathematical Model of the Feedback Loop Linking Glucose and Insulin

We determined insulin effectiveness by performing a hyperglycemic clamp and a

hyperinsulinemic-euglycemic clamp to measure insulin secretion and sensitivity together

with clearance (22,23,31), respectively. The performance of two clamp tests is laborious,

however, and so we attempted to estimate insulin effectiveness from a single clamp.

Given that we were not able to estimate insulin effectiveness with a single clamp

analysis using our previous model (23) (Appendix 5, Fig. S6, and Fig. S7A), we modified the

model by changing the positions of influx I and influx G as shown in Figure 3A (Appendix

6). We simulated the time courses of blood glucose and insulin concentrations in the

hyperinsulinemic-euglycemic clamp analysis, with one example for an NGT subject being

shown in Figure 3B. To confirm that the modified model appropriately captured the

characteristics of the time-series data, we evaluated the consistency between the ISI

calculated from the actual measurements and that calculated from the simulation data. The

coefficient of determination (R^2) between measured ISI and simulated ISI was 0.96 (Fig. 3C),

indicating that the model was reasonably able to capture the essential characteristics of the

time-series data.

Prediction of insulin effectiveness from only a hyperinsulinemic-euglycemic clamp

Some of the estimated parameters—including k_{m7}_ins, corresponding to insulin clearance

estimated from the modified model for the hyperinsulinemic-euglycemic clamp alone—

showed a tendency to diverge (Fig. 4A). However, $k_{m5}/k_{m7}{}^2$_ins was significantly

correlated with $k_5/k_7{}^2$_full ($r = 0.36, P = 1.2 \times 10^{-4}$), and $k_{m4}k_{m5}/k_{m7}{}^2$_ins was

significantly correlated with $k_4k_5/k_7{}^2$_full ($r = 0.52, P = 5.9 \times 10^{-9}$) (Fig. 4B and Fig.

S7B). Of note, insulin sensitivity, insulin secretion, and insulin clearance estimated from the

modified model for both the hyperglycemic clamp and the hyperinsulinemic-euglycemic

clamp (k_{m4}_full, k_{m5}_full, and k_{m7}_full, respectively) were significantly correlated with

those estimated from the original model (k_4_full, k_5_full, and k_7_full, respectively) (Fig. S8),

indicating that both models can be used to estimate the insulin-related parameters when both

the hyperglycemic and hyperinsulinemic-euglycemic clamps are performed.

We also attempted to estimate insulin effectiveness using a simple analytically

computable method. We hypothesized that insulin effectiveness can be analytically estimated

as follows (Appendix 7):

$$\text{Insulin effectiveness} = \frac{k_{\text{sen}}k_{\text{sec}}}{k_{\text{cle}}^2} \sim \frac{f_{\text{g}}I_0(I_{\text{SS}} - I_0)}{f_{\text{i}}G_{\text{SS}}G_0 I_{\text{SS}}}$$

where k_{sen}, k_{sec}, and k_{cle} represent insulin sensitivity, secretion, and clearance, respectively,

and G_{SS}, G_0, I_{SS}, I_0, f_{g}, and f_{i} represent blood glucose level at the steady state during insulin

administration, blood glucose level in the fasting state, blood insulin level at the steady state

during insulin administration, blood insulin level in the fasting state, glucose infusion rate at

the steady state, and insulin infusion rate at the steady state, respectively (Fig. 4C). This

analytically calculated insulin effectiveness ($k_{\text{sen}}k_{\text{sec}}/k_{\text{cle}}^2$) was significantly correlated

with k_4k_5/k_7^2_full ($r = 0.67, P = 5.8 \times 10^{-16}$) (Fig. 4D), indicating that insulin

effectiveness can be analytically estimated from a hyperinsulinemic-euglycemic clamp alone.

Of note, insulin effectiveness can also be analytically estimated from a hyperinsulinemic-

euglycemic clamp alone by using the serum C-peptide concentration ($r = 0.71, P =$

3.9×10^{-18}), OGTT ($r = 0.43, P = 4.2 \times 10^{-6}$), or a fasting blood test ($r = 0.54, P =$

1.1×10^{-9}) (Appendix 1, Appendix 8, Fig. S9 and Fig. S10).

CONCLUSIONS

We have here derived an equation relating insulin sensitivity, secretion, clearance, and blood

glucose levels [Eq (3)] that was well conserved among individuals with various levels of

glucose tolerance, and we showed that this equation provides a better fit, particularly for

individuals with IGT, than does an equation relating only insulin sensitivity, insulin secretion,

and blood glucose levels [Eq (2)]. Insulin clearance is implicated in the regulation of blood

glucose concentration (8), and our present data also suggest that, together with insulin

sensitivity and secretion, insulin clearance is important for such regulation, especially in

individuals with IGT. Our present results also suggest that insulin effectiveness, an index that

corrects DI for insulin clearance, reflects glucose disposal ability, possibly by representing

the effect of insulin on blood glucose concentration.

Insulin effectiveness is represented as a hyperbolic curve of DI and 1/insulin

clearance[2] or as a hyperbolic surface of insulin sensitivity, insulin secretion, and 1/insulin

clearance[2]. The deterioration of glucose tolerance is sometimes explained by the failure of

insulin secretion to increase sufficiently in compensation for a decline in insulin sensitivity,

with this failure being referred to as falling off the hyperbolic curve of DI (1,2). Our data now

suggest that, even when DI decreases, insulin effectiveness is preserved if insulin clearance is

also reduced. In general, moving up the hyperbolic curve of DI eventually positions

individuals at unstable extreme points of the curve where the risk of disease onset increases

(1), a phenomenon that has been referred to as canalization and decanalization (1,2) or as

glucose allostasis (32). Given that individuals with IGT are at high risk of developing T2DM,

the progression from NGT to IGT and T2DM can be referred to as canalization and

decanalization of the hyperbolic surface of insulin effectiveness.

Our data also indicate that IGT encompasses the condition of decreased insulin

clearance with a relatively well preserved insulin effectiveness and glucose set point

(Appendix 4), and that T2DM encompasses that of decreased insulin effectiveness with a

disrupted glucose set point. A modeling study with simulated data indicated that changes in

insulin clearance may affect glucose dynamics even when the normal glucose set point is

maintained, and can constitute a pathway for progression of glucose intolerance (19). Our

study based on mathematical models and observational data now indicates that the

phenomenon predicted by this previous model can occur in IGT. Our findings are also

consistent with those of a previous study indicating that an increase in insulin secretion is

followed by a decrease in first-pass hepatic insulin clearance when insulin resistance is induced (15).

Despite the apparent analogy of the moving up–falling off phenomenon, whether the body possesses the ability to reduce insulin clearance in response to a decline in DI and the potential physiological mechanism underlying such a response remain unknown. However, evidence suggests that changes in insulin clearance can be a determinant of glucose intolerance (21,33). Whereas the direction of causation of decreased insulin clearance and glucose intolerance remains unclear, a classification based on insulin effectiveness may provide new insight into the trajectories of glucose intolerance and T2DM.

Whereas we showed that insulin effectiveness can be estimated from a hyperinsulinemic-euglycemic clamp test alone, performance of even a single clamp is labor-intensive. Moreover, the correlation between insulin effectiveness estimated from the two clamps and that estimated from the single clamp was only 0.52, 0.67 or 0.71. It will therefore be important to develop more accurate and convenient methods for estimation of insulin effectiveness in order to validate its physiological relevance and clinical usefulness.

The current study has several limitations. For simplicity, we considered only insulin sensitivity, insulin secretion, and insulin clearance. However, insulin secretion has basal, first-phase, and second-phase components (34), and both insulin sensitivity and insulin clearance have hepatic and peripheral components (3,22). Saturation of or time-dependent changes in insulin clearance (6,22) were not considered. Whereas it is necessary to correct DI by insulin clearance for accurate assessment of the effect of insulin on blood glucose levels, individuals with the same insulin effectiveness can have different blood glucose levels depending on the input (Appendix 4). Several better approaches to evaluation of the hyperbolic law have also been presented, and a power function describing the relation between insulin sensitivity and insulin secretion gives a better representation (35–37). Given

that insulin sensitivity and insulin clearance are correlated (38), the plane representing

1/insulin clearance^2 is not necessarily horizontal (Fig. 2E), and the curve corresponding to DI

also departs from a hyperbola when the plane departs from the horizontal. Future research on

the coefficients of insulin sensitivity, secretion, and clearance in the formula defining the

effect of insulin on blood glucose levels is therefore warranted. In addition, the capacity for

glucose disposal is affected not only by insulin, but also by other hormones including

glucagon as well as by glucose (39,40). It is possible that differences in such factors may

explain the glucose dynamics of glucose tolerance. Further studies are also warranted to

develop extended equations that include glucose and glucagon to represent mechanisms of

blood glucose regulation.

In conclusion, we have here shown that insulin effectiveness, a new index that

corrects DI by insulin clearance, better reflects changes in glycemia compared with DI. By

analogy with falling off the hyperbolic curve of DI, the progression of glucose intolerance

may be considered as falling off the hyperbolic surface of insulin effectiveness. In theory,

insulin effectiveness represents the effect of insulin on blood glucose levels, and it may serve

as an indicator for the trajectory of changes in glucose tolerance. Further study is warranted

to understand the physiological and pathological relevance of this new index.

Acknowledgments

Personal Thanks. We thank our laboratory members for critically reading this manuscript

Author Contributions. H.S. and K.H. analyzed the data. H.S., K.H., T.Y., K.S., W.O., and

S.K. wrote the manuscript. W.O. and S.K. supervised the study. S.K. is the guarantor of this

work and, as such, had full access to all the data in the study and takes responsibility for the

integrity of the data and the accuracy of the data analysis.

Conflict of Interest. The authors declare no competing interests.

Funding and Assistance. This study was supported by the Japan Society for the Promotion

of Science (JSPS) KAKENHI (JP21H04759), CREST, the Japan Science and Technology

Agency (JST) (JPMJCR2123), and The Uehara Memorial Foundation.

References

1. Gibson G. Decanalization and the origin of complex disease. Nat Rev Genet 2009;10:134–140.
2. Kodama K, Tojjar D, Yamada S, Toda K, Patel CJ, Butte AJ. Ethnic differences in the relationship between insulin sensitivity and insulin response: a systematic review and meta-analysis. Diabetes Care 2013;36:1789–1796.
3. Ha J, Sherman A. Type 2 diabetes: one disease, many pathways. Am J Physiol Endocrinol Metab 2020;319:E410–E426.
4. De Gaetano A, Hardy TA. A novel fast-slow model of diabetes progression: Insights into mechanisms of response to the interventions in the Diabetes Prevention Program. PLoS One 2019;14:e0222833.
5. Topp B, Promislow K, Devries G, Miura RM, Finegood DT. A Model of β -Cell Mass, Insulin, and Glucose Kinetics: Pathways to Diabetes. J Theor Biol 2000;206:605–619.
6. Mari A, Tura A, Grespan E, Bizzotto R. Mathematical Modeling for the Physiological and Clinical Investigation of Glucose Homeostasis and Diabetes. Front Physiol 2020;11:575789.
7. Jones CN, Pei D, Staris P, Polonsky KS, Chen YD, Reaven GM. Alterations in the glucose-stimulated insulin secretory dose-response curve and in insulin clearance in nondiabetic insulin-resistant individuals. J Clin Endocrinol Metab 1997;82:1834–1838.
8. Kim SH, Reaven GM. Insulin clearance: an underappreciated modulator of plasma insulin concentration. J Investig Med 2016;64:1162–1165.
9. Goodarzi MO, Cui J, Chen YDI, Hsueh WA, Guo X, Rotter JI. Fasting insulin reflects heterogeneous physiological processes: role of insulin clearance. Am J Physiol Endocrinol Metab 2011;301:E402–E408.
10. de Oliveira DM, Tura A, Vasques ACJ, Camilo DF, Lima MM, de Lemos-Marini SHV, et al. Insulin Resistance in Congenital Adrenal Hyperplasia is Compensated for by Reduced Insulin Clearance. J Clin Endocrinol Metab 2021;106:e1574–e1585.
11. Mittelman SD, Van Citters GW, Kim SP, Davis DA, Dea MK, Hamilton-Wessler M, et al. Longitudinal compensation for fat-induced insulin resistance includes reduced insulin clearance and enhanced beta-cell response. Diabetes 2000;49:2116–2125.
12. Gastaldelli A, Abdul Ghani M, DeFronzo RA. Adaptation of Insulin Clearance to Metabolic Demand Is a Key Determinant of Glucose Tolerance. Diabetes 2021;70:377–385.
13. Haffner SM, Stern MP, Watanabe RM, Bergman RN. Relationship of insulin clearance and secretion to insulin sensitivity in non-diabetic Mexican Americans. Eur J Clin Invest 1992;22:147–153.

14. Galderisi A, Polidori D, Weiss R, Giannini C, Pierpont B, Tricò D, et al. Lower Insulin
Clearance Parallels a Reduced Insulin Sensitivity in Obese Youths and Is Associated
With a Decline in β-Cell Function Over Time. Diabetes 2019;68:2074–2084.

15. Kim SP, Ellmerer M, Kirkman EL, Bergman RN. β-Cell "rest" accompanies reduced
first-pass hepatic insulin extraction in the insulin-resistant, fat-fed canine model. Am J
Physiol Endocrinol Metab 2007;292:E1581–E1589.

16. Bizzotto R, Tricò D, Natali A, Gastaldelli A, Muscelli E, De Fronzo RA, et al. New
Insights on the Interactions Between Insulin Clearance and the Main Glucose
Homeostasis Mechanisms. Diabetes Care 2021;44:2115–2123.

17. DeFronzo RA, Tripathy D, Abdul-Ghani M, Musi N, Gastaldelli A. The disposition
index does not reflect β-cell function in IGT subjects treated with pioglitazone. J Clin
Endocrinol Metab 2014;99:3774–3781.

18. Ladwa M, Bello O, Hakim O, Shojaee-Moradie F, Boselli L, Charles-Edwards G, et al.
Insulin clearance as the major player in the hyperinsulinaemia of black African men
without diabetes. Diabetes Obes Metab 2020;22:1808–1817.

19. Karin O, Swisa A, Glaser B, Dor Y, Alon U. Dynamical compensation in physiological
circuits. Mol Syst Biol 2016;12:886.

20. Galderisi A, Giannini C, Weiss R, Kim G, Shabanova V, Santoro N, et al. Trajectories
of changes in glucose tolerance in a multiethnic cohort of obese youths: an observational
prospective analysis. Lancet Child Adolesc Health 2018;2:726–735.

21. Bergman RN, Piccinini F, Kabir M, Kolka CM, Ader M. Hypothesis: Role of Reduced
Hepatic Insulin Clearance in the Pathogenesis of Type 2 Diabetes. Diabetes
2019;68:1709–1716.

22. Piccinini F, Bergman RN. The Measurement of Insulin Clearance. Diabetes Care
2020;43:2296–2302.

23. Ohashi K, Komada H, Uda S, Kubota H, Iwaki T, Fukuzawa H, et al. Glucose
Homeostatic Law: Insulin Clearance Predicts the Progression of Glucose Intolerance in
Humans. PLoS One 2015;10:e0143880.

24. Ohashi K, Fujii M, Uda S, Kubota H, Komada H, Sakaguchi K, et al. Increase in hepatic
and decrease in peripheral insulin clearance characterize abnormal temporal patterns of
serum insulin in diabetic subjects. npj Systems Biology and Applications 2018;4:1–12.

25. Cobelli C, Dalla Man C. Minimal and Maximal Models to Quantitate Glucose
Metabolism: Tools to Measure, to Simulate and to Run in Silico Clinical Trials. J
Diabetes Sci Technol 2021;19322968211015268.

26. DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying
insulin secretion and resistance. Am J Physiol 1979;237:E214-E223.

27. Okuno Y, Komada H, Sakaguchi K, Nakamura T, Hashimoto N, Hirota Y, et al.
Postprandial serum C-peptide to plasma glucose concentration ratio correlates with oral

glucose tolerance test- and glucose clamp-based disposition indexes. Metabolism
2013;62:1470–1476.

28. Warton DI, Duursma RA, Falster DS, Taskinen S. smatr 3 - an R package for estimation
and inference about allometric lines. Methods Ecol Evol 2012;3:257–259.

29. Gastaldelli A, Ferrannini E, Miyazaki Y, Matsuda M, DeFronzo RA, San Antonio
metabolism study. Beta-cell dysfunction and glucose intolerance: results from the San
Antonio metabolism (SAM) study. Diabetologia 2004;47:31–39.

30. Kahn SE, Bergman RN, Schwartz MW, Taborsky GJ Jr, Porte D Jr. Short-term
hyperglycemia and hyperinsulinemia improve insulin action but do not alter glucose
action in normal humans. Am J Physiol 1992;262:E518-E523.

31. Sjaarda L, Lee S, Tfayli H, Bacha F, Bertolet M, Arslanian S. Measuring β-cell function
relative to insulin sensitivity in youth: does the hyperglycemic clamp suffice? Diabetes
Care 2013;36:1607–1612.

32. Stumvoll M, Tataranni PA, Stefan N, Vozarova B, Bogardus C. Glucose allostasis.
Diabetes 2003;52:903–909.

33. Sugiyama S, Jinnouchi H, Hieshima K, Kurinami N, Jinnouchi K, Yoshida A, et al.
Potential identification of type 2 diabetes with elevated insulin clearance. NEJM
Evidence 2022;1:EVIDoa2100052.

34. Cobelli C, Toffolo GM, Man CD, Campioni M, Denti P, Caumo A, et al. Assessment of
β-cell function in humans, simultaneously with insulin sensitivity and hepatic extraction,
from intravenous and oral glucose tests. Am J Physiol Endocrinol Metab 2007;293:E1–
E15.

35. Denti P, Toffolo GM, Cobelli C. The disposition index: from individual to population
approach. Am J Physiol Endocrinol Metab 2012;303:E576-E586.

36. Ferrannini E, Mari A. Beta cell function and its relation to insulin action in humans: a
critical appraisal. Diabetologia 2004;47:943–956.

37. Cobelli C, Dalla Man C, Toffolo G, Basu R, Vella A, Rizza R. The oral minimal model
method. Diabetes 2014;63:1203–1213.

38. Lorenzo C, Hanley AJG, Wagenknecht LE, Rewers MJ, Stefanovski D, Goodarzi MO,
et al. Relationship of insulin sensitivity, insulin secretion, and adiposity with insulin
clearance in a multiethnic population: the insulin Resistance Atherosclerosis study.
Diabetes Care 2013;36:101–103.

39. Dube S, Errazuriz-Cruzat I, Basu A, Basu R. The forgotten role of glucose effectiveness
in the regulation of glucose tolerance. Curr Diab Rep 2015;15:605.

40. Lee YH, Wang MY, Yu XX, Unger RH. Glucagon is the key factor in the development
of diabetes. Diabetologia 2016;59:1372–1375.

Figures

Figure 1. Relation between insulin-related parameters and blood glucose levels. (A) Scatter

plot for $a_a \log \text{PG120} + b_a$ versus $\log k_4 k_5$ (Eq (2)). Each point corresponds to the values for

a single subject. Green circles, red crosses, and blue squares indicate NGT, IGT, and T2DM

subjects, respectively. R^2 is the coefficient of determination, and the P value is for testing the

hypothesis of no correlation. The parameters a_a and b_a were estimated to minimize RSS

between $a_a \log \mathrm{PG}120 + b_a$ and $\log k_4 k_5$ in all subjects. (B) Scatter plots for $a \log \mathrm{PG}120 +$

b versus $\log k_4 k_5$ in NGT, IGT, and T2DM subjects (Eq (2)). The parameters a_n and b_n, a_i

and b_i, and a_t and b_t were estimated to minimize RSS between $a \log \mathrm{PG}120 + b$ and

$\log k_4 k_5$ in NGT, IGT, and T2DM subjects, respectively. (C) Scatter plot for $2 \log k_7 +$

$a_a \log \mathrm{PG}120 + b_a$ versus $\log k_4 k_5$ (Eq (3)). The parameters a_a and b_a were estimated to

minimize RSS between $2 \log k_7 + a_a \log \mathrm{PG}120 + b_a$ and $\log k_4 k_5$ in all subjects. (D)

Scatter plots for $2 \log k_7 + a \log \mathrm{PG}120 + b$ versus $\log k_4 k_5$ in NGT, IGT, and T2DM

subjects (Eq (3)). The parameters a_n and b_n, a_i and b_i, and a_t and b_t were estimated to

minimize RSS between $2 \log k_7 + a \log \mathrm{PG}120 + b$ and $\log k_4 k_5$ in NGT, IGT, and T2DM

subjects, respectively. The indices k_4, k_5, and k_7, which correspond to insulin sensitivity,

insulin secretion, and insulin clearance, respectively, were calculated by a previously

described method (23), and PG120 is the plasma glucose concentration at 120 min during an

OGTT.

Figure 2. Relation between DI and insulin effectiveness. (A) Box plots of

log (insulin effectiveness), corresponding to the logarithm of $k_4 k_5 / k_7^2$, in NGT, IGT, and

T2DM subjects. Each point corresponds to the value for a single subject, with green circles,

red crosses, and blue squares indicating NGT, IGT, and T2DM subjects, respectively. The

boxes denote the median and upper and lower quartiles. P values were determined by one-

way ANOVA and Tukey's post hoc test. (B) Scatter plots for k_4 and k_5 fitted with hyperbolic

curves (based on the function $xy = $ constant), and kernel density plots. The hyperbolic

curves correspond to the mean value of DI for NGT, IGT, or T2DM subjects. Each point

corresponds to the values for a single subject, with green, red, and blue indicating NGT, IGT,

and T2DM, respectively. The bandwidth of each kernel is based on the rule-of-thumb

bandwidth. (C) Scatter plots for $k_4 k_5$ and $1/k_7{}^2$ fitted with hyperbolic curves (based on the

function $xy = $ constant), and kernel density plots. The hyperbolic curves correspond to the

mean value of insulin effectiveness in NGT, IGT, or T2DM subjects. Each point corresponds

to the values for a single subject, with green, red, and blue indicating NGT, IGT, and T2DM,

respectively. The bandwidth of each kernel is based on the rule-of-thumb bandwidth. (D)

Hyperbolic surfaces (based on the function $xyz = $ constant) corresponding to the mean

value of insulin effectiveness in NGT, IGT, or T2DM subjects. (E) Hyperbolic surfaces

corresponding to the mean value of insulin effectiveness as well as planes corresponding to

the mean value of 1/insulin clearance[2] in NGT, IGT, or T2DM subjects. The axes correspond

to k_4, k_5, and $1/k_7{}^2$ as in (D), but the label for k_5 was omitted for clarity. The intersections

(solid lines) of the surfaces representing insulin effectiveness and the planes representing the

mean value of 1/insulin clearance[2] correspond to the hyperbolic curves in (B).

Figure 3. Modified mathematical model of the feedback loop that links glucose and insulin.

(A) Model diagram. The letters within circles indicate the variables of the model, the arrows

denote the fluxes, and the lines ending in circles or bars represent activation or inactivation,

respectively. ϕ indicates a fixed value. The variables G and I represent blood glucose and

insulin concentrations, respectively. The variable Y corresponds to the effective glucose

concentration for induction of variable X, which represents insulin secreted from pancreatic

β-cells. (B) Time courses of blood glucose, blood insulin, infused glucose, and infused

insulin during hyperinsulinemic-euglycemic clamp analysis for an NGT subject. The blue

lines indicate the simulated patterns, and the red circles indicate the measured time course

data. The simulated values of each variable were rescaled to absolute concentration and

plotted. (C) Scatter plot for simulated ISI versus measured ISI. Each point corresponds to the

values for a single subject, with green circles, red crosses, and blue squares indicating NGT,

IGT, and T2DM subjects, respectively. The measured and simulated ISI values were

calculated from the normalized time course and plotted. R^2 is the coefficient of

determination, and the P value is for testing the hypothesis of no correlation.

Figure 4. Relation between parameters estimated from a hyperinsulinemic-euglycemic clamp

alone and those estimated from both hyperglycemic and hyperinsulinemic-euglycemic

clamps. (A) Estimated parameters of the modified model based on a hyperinsulinemic-

euglycemic clamp alone. (B) Heatmap of Spearman's correlation coefficient, with the P

values corresponding to testing the hypothesis of no correlation. The parameters k_4_full,

k_5_full, k_7_full, $k_5/k_7{}^2$_full, and $k_4 k_5/k_7{}^2$_full correspond to insulin sensitivity, insulin

secretion, insulin clearance, the ratio of insulin secretion to the square of insulin clearance,

and insulin effectiveness estimated from both hyperglycemic and hyperinsulinemic-

euglycemic clamps by the previously described mathematical model (23) (Fig. S6A),

respectively. The parameters k_{m4}_ins, k_{m5}_ins, k_{m7}_ins, $k_{m5}/k_{m7}{}^2$_ins, and

$k_{m4} k_{m5}/k_{m7}{}^2$_ins correspond to insulin sensitivity, insulin secretion, insulin clearance, the

ratio of insulin secretion to the square of insulin clearance, and insulin effectiveness

estimated from the hyperinsulinemic-euglycemic clamp alone by the modified mathematical

model (Fig. 3A), respectively. The insulin sensitivity index (ISI) and area under the insulin

curve during the first 10 min of the hyperglycemic clamp were also calculated. (C) Simulated

time courses of blood glucose, blood insulin, infused glucose, and infused insulin during a

hyperinsulinemic-euglycemic clamp. G_{SS}, blood glucose level at the steady state during

insulin administration; G_0, blood glucose level in the fasting state; I_{SS}, blood insulin level at

the steady state during insulin administration; I_0, blood insulin level in the fasting state; f_g,

glucose infusion rate at the steady state; and f_i, insulin infusion rate at the steady state. (D)

Scatter plot for $\log k_{sen} k_{sec}/k_{cle}{}^2$ versus $\log k_4 k_5/k_7{}^2$_full. The parameters $k_{sen} k_{sec}/k_{cle}{}^2$

and $k_4 k_5/k_7{}^2$_full correspond to insulin effectiveness analytically estimated from the

hyperinsulinemic-euglycemic clamp alone (Appendix 7) and that estimated from the

hyperglycemic and hyperinsulinemic-euglycemic clamps by the previously described

mathematical model (23) (Fig. S6A), respectively. Each point corresponds to the values for a

single subject, with green circles, red crosses, and blue squares indicating NGT, IGT, and

T2DM subjects, respectively. r is Spearman's correlation coefficient, and the P value is for

testing the hypothesis of no correlation.