nucMACC: An optimized MNase-seq pipeline measures genome-wide nucleosome accessibility and stability

AUTHORS:

Wernig-Zorc Sara[1], Kugler Fabian[1,2], Schmutterer Leo[2], Räß Patrick[2], Hausmann Clemens[2],

Holzinger Simon[1], Gernot Längst[1,*], Uwe Schwartz[2]*

[1] Biochemistry III, University of Regensburg, Regensburg, Germany
[2] NGS Analysis Center Biology and Pre-clinical Medicine, University of Regensburg, Regensburg, Germany
*Co-corresponding authors

ABSTRACT:

Micrococcal nuclease sequencing (MNase-seq) is the state-of-the-art method for determining the chromatin structure and nucleosome positioning. Analysis of MNase-seq data is complex due to variable degrees of sample quality, sequence bias of the endonuclease, and high sequencing depth requirement. Therefore, we developed an optimized analysis pipeline written in nextflow and publicly available on GitHub. The nucMACC pipeline provides high-resolution nucleosome positions and associated properties such as nucleosome accessibility and stability. Hyper-/hypo-accessible nucleosomes or non-canonical and unstable nucleosome positions in the genome are called automatically. We provide recommendations for the minimum sequencing depth and show robust results using two differential MNase conditions.

Using the nucMACC pipeline, we characterized the nucleosomal properties in D. melanogaster and explored the correlation between these properties and gene expression. Intriguingly, we found a distinct group of unstable nucleosomes enriched in promoters marked with the M1BP motif. The expression rate of the genes harboring an unstable nucleosome in the promoter was not altered compared to genes with a nucleosome-depleted promoter region, whereas the RNA polymerase pausing rate was affected.

INTRODUCTION:

Nucleosomes are the basic unit of DNA packaging in eukaryotic cells. Electron microscopy of chromatin spreads showed regularly spaced nucleosome core particles separated by linker DNA, giving the appearance of a "beads-on-string" structure (Olins, Douglas Carlson and Olins, 1975). The nucleosome core particle consists of 147 bp of DNA wrapped around a histone octamer, composed of histone proteins H2A, H2B, H3, and H4 (Kornberg, 1974; Luger *et al.*, 1997). The stability of the nucleosome core is maintained by over 360 hydrogen bonds between the DNA and histone proteins (Davey *et al.*, 2002). Linker DNA varies in length from 15 to 95 bp depending on the species, cell type, and even the chromatin state within the same nucleus (Valouev *et al.*, 2011; Silberhorn *et al.*, 2016; Baldi *et al.*, 2018). DNA in the linker is generally accessible to DNA-binding factors, whereas access to DNA in the nucleosome core is hindered. Nucleosome positioning, structure, and modification play crucial roles in regulating gene expression and determining the accessibility of DNA for DNA-templated processes, thereby defining cell type-specific gene activity and repression.

In vitro experiments have revealed that nucleosome positioning depends on the bendability and structure of the underlying DNA sequence and the effect of steric hindrance between neighboring nucleosomes (Kaplan *et al.*, 2009; Struhl and Segal, 2013). On the other hand, *in vivo* nucleosome positioning is more complex and is affected by additional factors, such as chromatin remodelers, histone variants, transcription factors (TF), histone posttranslational modifications, or transcription machinery (Chereji and Clark, 2018). Regulatory regions, such as promoters, enhancers, and insulators, are marked by well-positioned nucleosomes, which is a common feature of eukaryotic chromatin (Schones *et al.*, 2008; Hu *et al.*, 2011).

Besides the positioning of nucleosomes, studying their structural characteristics is central to revealing the functions of chromatin. Nucleosomes vary in their histone composition and number. Histone variants have distinct structural and functional characteristics, serve specific roles in chromatin regulation, and are produced independently of the cell cycle. The histone H3 variant CENP-A is incorporated into nucleosomes at the centromeres. DNA histone interactions are weakened in CENP-A-containing nucleosomes, partially unwrapping DNA from the histone surface (Hasson *et al.*, 2013). Hexasomes, which miss one H2A-H2B dimer, can form through various mechanisms such as transcription, histone chaperones, and chromatin remodeling complexes (Kireeva *et al.*, 2002; Prasad *et al.*, 2016; Ramachandran, Ahmad and Henikoff, 2017; Liu *et al.*, 2019). The presence of hexasomes near transcription start sites was suggested in a recent study (Rhee *et al.*, 2014), highlighting their potential role

in gene regulation. However, it is still unclear whether these sub-nucleosomal structures are

stable or short-lived nucleosome intermediates *in vivo*.

MNase-seq has been widely used to study chromatin structure and nucleosome positioning.

It involves the use of micrococcal nuclease (MNase), an enzyme that selectively cleaves linker

DNA and leaves the nucleosome core particles intact. The resulting DNA fragments were

sequenced and analyzed to determine the location and distribution of nucleosomes along the

genome (Baldi, Korber and Becker, 2020). MNase-seq provides a high-resolution map of the

nucleosome landscape and has contributed to our understanding of the dynamic and

regulated nature of chromatin structure (Valouev *et al.*, 2011; Teif *et al.*, 2012; Diermeier *et*

al., 2014). Initially, high MNase digestions were applied, hydrolyzing chromatin completely to

the mono-nucleosomal level. Later, we and others combined different degrees of MNase

digestion to study chromatin accessibility and nucleosome stability (Henikoff *et al.*, 2011;

Chereji *et al.*, 2015; Mieczkowski *et al.*, 2016; Schwartz *et al.*, 2018; Chereji, Bryson and

Henikoff, 2019). As the data analysis of MNase titration experiments is complex, new metrics

have been introduced to provide a stable framework for accurate quantification of chromatin

accessibility and nucleosome occupancy (Mieczkowski *et al.*, 2016; Chereji, Bryson and

Henikoff, 2019). However, at least six titrations of MNase per sample and additional spike-ins

were required, resulting in a high workload and the required amount of material.

Furthermore, application to new experiments is challenging because complete analysis

pipelines are missing.

Here, we introduce a comprehensive and user-friendly nucleosome-based MNase

accessibility (nucMACC) analysis pipeline. The nucMACC workflow starts directly with raw

sequencing data, assesses the critical quality metrics relevant to MNase-seq experiments, and

summarizes the results into user-friendly reports. The workflow reveals the genome-wide

nucleosome positions and their corresponding nucleosome accessibility (nucMACC) scores.

Hyper- and hypo-accessible nucleosome positions with exceptionally high or low nucMACC

scores are determined. Additionally, as a unique feature of the nucMACC pipeline,

nucleosome stability scores are inferred from sub-nucleosomal fragments uncovering un-

stable or non-canonical nucleosomes (Figure S1). We show that the pipeline obtains robust

results using only two MNase conditions per sample. The nucMACC pipeline is written in

nextflow, available at GitHub (https://github.com/uschwartz/nucMACC), and integrates the

Docker software container, making it easy to use, portable, reproducible, and scalable.

RESULTS:

nucMACC pipeline overview

The nucMACC pipeline requires at least two MNase-seq titrations as input, preferentially in combination with a histone immunoprecipitation step to exclude non-histone protein-mediated DNA-binding events. MNase, which preferentially hydrolyzes accessible linker DNA between nucleosomes, releases nucleosome core particles from the chromatin. Depending on the MNase digestion conditions, nucleosomes are released at different rates (Figure 1A). Under mild MNase conditions, nucleosomes from accessible and active genomic sites are preferentially released, referred to as hyper-accessible nucleosomes (Figure 1A, green nucleosome) (Mieczkowski *et al.*, 2016; Schwartz *et al.*, 2018). Increasing the MNase concentration or digestion duration continuously releases more nucleosome core particles from the chromatin until the chromatin is fully hydrolyzed. Additionally, MNase is a processive enzyme. Once the nucleosome is released from chromatin, MNase progressively trims the linker DNA and hydrolyzes it within the realm of the nucleosomal DNA, resulting in sub-nucleosomal fragments. Canonical nucleosome core particles are resistant to internal MNase cleavage and remain mainly complete even under extensive chromatin digestion conditions. However, it has been suggested that a special group of nucleosomes, often referred to as un-stable or fragile nucleosomes, exists, which are MNase-sensitive and start to be degraded already under mild MNase conditions to sub-nucleosome-sized fragments (Figure 1A, blue nucleosome) (Weiner *et al.*, 2010; Kubik *et al.*, 2015; Voong *et al.*, 2016). We refer to such nucleosomes as un-stable nucleosomes in the remainder of this manuscript. Under extensive MNase conditions, un-stable nucleosomes are completely degraded and not any more detectable. Therefore, sub-nucleosomal fragments in MNase titration experiments can be used to measure nucleosome stability, as reflected by resistance against internal MNase cleavage.

Starting from raw sequencing files in the FASTQ format, the nucMACC pipeline first assesses the sequencing quality. Next, the reads are aligned to the provided reference genome and subsequently mapped reads are filtered based on mapping quality (Figure S2). Optionally

ambiguous genomic elements, such as blacklisted regions or mitochondrial chromosomes,

can be removed from the analysis. To obtain high-resolution nucleosome positions, fragments

between 140 and 200 bp in length, the typical size of mono-nucleosomal DNA, were selected

(Figure 1B). Size-selected fragments of all MNase titrations are pooled and used to derive a

comprehensive map of the genome-wide nucleosome positions.

Recently, a metric for measuring chromatin accessibility, termed MACC (MNase accessibility),

was introduced (Mieczkowski et al., 2016). As the MACC scores provide an elegant way to

measure chromatin accessibility based on MNase-seq data, we adapted the MACC scoring

system to increase resolution and accuracy. Instead of counting the MNase reads in arbitrary

genomic bins as in the original MACC version, we quantify the MNase accessibility directly at

the defined nucleosome positions, which results in a specific accessibility score for each

nucleosome (nucMACC). Next, linear regression is conducted on the normalized fragment

frequencies for each nucleosome (Figure S2, see Methods section for details). The slope of

the regression line multiplied by minus one is used as the nucMACC score. Since MNase

exhibits an increasing AT preference with higher concentrations, raw nucMACC scores are

normalized for the underlying GC content using a LOESS regression.

As an additional feature of the nucMACC pipeline fragments shorter than 140 bp, which

represent sub-nucleosomal DNA fragments, are used to measure nucleosome stability.

Here, we use only the lowest MNase concentration to call nucleosome positions in the

following called sub-nucleosomes, as in the higher MNase titrations un-stable nucleosomes

are completely hydrolyzed. Analogous to the nucMACC scores, sub-nucleosomal fragment

frequencies for each MNase titration are quantified at the sub-nucleosome positions. The

slope of the regression is used to measure nucleosome stability and normalized to the

underlying GC content. The normalized score is referred to as the sub-nucleosome MNase

accessibility score (sub-nucMACC).

The nucMACC pipeline scores nucleosome accessibility and stability

To establish the nucMACC pipeline, we used mainly the MNase chromatin digestions coupled

with H4 immunoprecipitation from the original MACC publication (Mieczkowski *et al.*, 2016).

The dataset collection contains four different MNase concentrations (1.5U, 6.25U, 25U, and

100U) done in two replicates each. The fragment size distribution is an important quality

control step as it clearly shows the degree of chromatin digestion in each sample (Figure 1B).

With increasing MNase concentrations the di-nucleosome fraction is disappearing, and the

levels of mono-nucleosomes are increasing. The main mono-nucleosomal peak is in the range

of 140–200 bp, while the peak summit is shifted with higher MNase concentrations closer to

147 bp, the length of DNA in a nucleosome core particle. Fragment sizes shorter than 140 bp

indicate sub-nucleosomal DNA fragments, which mainly arise from DNA cleavage within the

core nucleosome. Sub-nucleosomal DNA fragments are barely present in mild MNase

conditions but increase in extensive MNase samples. Based on the curve shape of the

fragment size distribution, we selected the fragments in the range of 140 – 200 bp in length

for mono-nucleosome analysis and shorter than 140 bp for sub-nucleosome analysis,

respectively (Figure 1B).

Plotting the fragment frequencies of mono- and sub-nucleosomes at the TSS shows that the

selected categories contain different information (Figure 1C and 1E). Looking at the mono-

nucleosomes of the two extensive MNase conditions (25U and 100U) reveals a well-known

nucleosome positioning pattern at the TSS: a nucleosome depleted region (NDR) followed

downstream by a phased nucleosome array, which starts with a well-positioned +1-

nucleosome and propagates into the gene. The mild MNase samples (1.5U and 6.25U) reveal

the same mono-nucleosome positions, however, distinct positions, such as the +1-

nucleosome, are markedly enriched compared to the extensive MNase samples indicating

that nucleosomes exhibit different release rates from chromatin. The nucMACC score

quantifies chromatin accessibility as mono-nucleosome fragment frequencies change with

MNase titrations (Figure 1D and S2).

Comparing the mono-nucleosome and the sub-nucleosome TSS profile the extensive MNase

samples are very similar showing that once the nucleosomes are released from chromatin,

MNase trims the DNA within the nucleosome core. Nevertheless, an additional dominant

peak is appearing at the denominated (NDR) in the sub-nucleosomal fraction of the mild

MNase condition, indicating that this position is not completely free of histones, but occupied

by MNase sensitive nucleosomes, which are completely degraded in extensive MNase

digestions (Figure 1E). The sub-nucMACC score quantifies nucleosome stability as changes in

sub-nucleosome fragment frequencies over MNase titrations (Figure 1F and S2).

Even though the way how nucMACC and sub-nucMACC scores are calculated is very similar,

their scores are not correlated at overlapping positions, confirming that both metrics show

distinct features of nucleosomes (Figure S3A). The correlation of the nucMACC scores with

the original MACC scores (Mieczkowski *et al.*, 2016) was modest (R=0.48), whereas the sub-

nucMACC scores showed no correlation (R=-0.24) indicating that our pipeline uncovers new

features in MNase titration experiments (Figure S3B-C).

Centering the nucMACC scores on the TSS shows that the +1 nucleosomes of actively

transcribed genes are in general hyper-accessible, whereas nucleosomes further downstream

in the gene body did not significantly change nucleosome accessibility (Figure 1G).

Remarkably, in genic regions, we observed no difference in accessibility between silent and

highly expressed genes, except at the +1 nucleosome (Figure S3D), confirming that chromatin

accessibility is not changed at a global scale but modulated at defined local sites, such as the

promoter, to regulate gene transcription (Schwartz *et al.*, 2018).

The nucleosome stability profile at the TSS of active genes was different compared to the

nucleosome accessibility score. The lowest nucleosome stability scores were obtained

upstream of the +1 nucleosome directly at the TSS of active genes. The sub-nucleosome

analysis uncovered un-stable nucleosomes directly at the denominated NDRs showing overall

low nucleosome occupancy in mono-nucleosome analysis (Figure 1G). In contrast, the

transcription end sites (TES) are nucleosome depleted in Drosophila, since we do not observe

un-stable nucleosomes there as recently proposed in yeast and mouse (Figure S3E) (Chereji,

Ocampo and Clark, 2016; Voong *et al.*, 2016).

Overall, the nucleosome features provided by the nucMACC pipeline are robust and

reproducible, as shown by a comparison with MNase titrations combined with H3-ChIP data

(Figure S4A-G). We observed the highest variation between H3 and H4 experiments in the

number and location of the sub-nucleosome positions (Figure S4C). This can be explained, as

the number of sub-nucleosomal fragments in the H3 mild MNase samples was significantly

lower compared to the H4 dataset revealing fewer sub-nucMACC positions (addressed in

more detail later). Nevertheless, the sub-nucMACC scores at overlapping positions showed a

high correlation (Figure S4C, R=0.8).

In summary, splitting up MNase titration data by fragment sizes into mono-nucleosomes and

sub-nucleosomes enables the nucMACC pipeline to reveal nucleosome positions, nucleosome

accessibility, and nucleosome stability scores within the same experiment.

Furthermore, we tested whether the additional histone immunoprecipitation step after

chromatin digestion is required to obtain reliable nucleosome features. Mono- and sub-

nucleosome positioning changed and overlapping positions showed only a modest correlation

with whole chromatin extractions (Fig S4H-K). Changes in the fragment profile were clearly

visible upstream of the TSS and most pronounced at the -1 nucleosome position affecting

nucMACC and sub-nucMACC scores (Figure S4L-N). The -1 nucleosome position exhibited the

highest accessibility in whole chromatin extractions and the sub-nucleosomal fraction in the

mild MNase sample showed a broad and pronounced enrichment at the promoter of

expressed genes reminiscent of other methods to measure chromatin accessibility such as

DNase- or ATAC-seq (Buenrostro *et al.*, 2013; Vierstra *et al.*, 2014). The differences between

whole chromatin and histone immunoprecipitated MNase datasets can be addressed by non-

histone proteins bound to DNA (Mieczkowski *et al.*, 2016). To obtain accessibility and stability

measurements specific to nucleosomes, an additional histone immunoprecipitation step is

necessary. Nevertheless, the nucMACC pipeline can be run at whole chromatin extractions

and measures overall chromatin accessibility and stability over all molecules bound to DNA.

Characterization of nucleosome accessibility scores

To identify functionally relevant nucleosomes, the nucMACC pipeline extracts nucleosomes

with exceptionally high or low accessibility, termed hyper- or hypo-accessible nucleosomes.

Therefore, the nucMACC scores of each nucleosome are plotted against their ranks. Ranks are

normalized by the total number of nucleosomes and the MACC score by the total range of

scores. In that way, both the x and y axes have a value range of 1 (Figure 2A). To geometrically

identify the positions, where the signal decreases/increases rapidly, we determined the

positions at the curve where the slope is the first/last time below one. Hypo-accessible

nucleosomes are defined as the nucleosome positions before the slope of the curve decreases

below 1 (Figure 2A, purple dots), and hyper-accessible nucleosomes after the slope increases

again above 1, respectively (Figure 2A, green dots). A similar strategy is applied to define

super-enhancer (Whyte *et al.*, 2013).

In total, we identified 6,763 (2 %) hypo-accessible and 16,342 (4,9 %) hyper-accessible

nucleosomes in the Drosophila S2 H4-ChIP data set. Hyper-accessible nucleosomes are

strongly enriched at DNase hypersensitive sites, whereas hypo-accessible nucleosomes are

depleted, confirming the nucMACC score as a measure for accessibility (Figure 2B). Genomic

feature characterization of hyper- and hypo-accessible nucleosomes revealed an enrichment

at the promoter and intergenic regions, respectively (Figure 2C). Hyper-accessible

nucleosomes are preferentially associated with H3K4me1, H3K4me3, and H3K27ac histone

modifications marking active TSS and enhancer, whereas hypo-accessible nucleosomes are

associated with H3K9me3 and H3K27me3 modifications indicative for heterochromatic

regions (Figure 2D). In agreement with the histone modification analysis, hyper-accessible

nucleosomes occupy mainly expressed genes, while hypo-accessible nucleosomes are present

on silent genes (Figure 2E). Nevertheless, we found a substantial fraction of hypo-accessible

nucleosomes at the exons of expressed genes potentially regulating differential exon usage.

The nucMACC pipeline identifies functional nucleosome positions in the genome regulating

DNA accessibility and therefore genome activity.

Characterization of nucleosome stability scores

To measure nucleosome stability, sub-nucleosome positions were derived from the sub-

nucleosome fragments in the lowest MNase condition. Hyper-accessible nucleosomes are

expected to have a higher MNase degradation rate as they are less protected once they are

released from chromatin. Therefore, sub-nucleosome positions are selected, which either

contained at least four-fold more normalized sub-nucleosome fragment counts than the

mono-nucleosome fraction or did not overlap with any mono-nucleosome position.

Similar to the selection of hyper- and hypo-accessible nucleosomes, nucleosomes exhibiting

extreme stability scores (sub-nucMACC) were determined (Figure 3A). This selection strategy

resulted in 4,233 (6.7 %) nucleosomes showing very low stability scores and therefore termed

un-stable nucleosomes. Additionally, a rather small subset of nucleosomes (n=355, 0.6 %)

were enriched in the sub-nucleosome fraction of the lowest MNase condition and remained

stable at higher MNase concentrations. This group represents most likely structurally altered

nucleosomes with additional MNase cleavage sites in the realm of the nucleosome core or

non-octamer histone compositions, such as hexa- or hemisomes (Dalal *et al.*, 2007; Hasson *et*

al., 2013; Rhee *et al.*, 2014). Nucleosomes enriched in the low sub-nucleosomal fractions and

showing extraordinarily high stability scores are referred here to as non-canonical

nucleosomes.

In contrast to non-canonical nucleosomes, un-stable nucleosomes positions are characterized

by high chromatin accessibility (Figure 3B) and are mainly located at gene promoter (53,8 %),

whereas non-canonical nucleosomes are enriched in intergenic and exonic regions (Figure

3C). Preferentially, active histone marks are associated with un-stable nucleosomes (Figure

3D) and accordingly, un-stable nucleosomes are present at expressed genes (Figure 3E).

Interestingly, we identify a subgroup of un-stable nucleosomes associated with active

H3K4me1 and repressive H3K27me3 marking bivalent enhancers. Conversely, non-canonical

nucleosomes showed enrichment in repressive H3K27me3 and H3K4me1 (Figure 3D).

Nevertheless, 30% of the non-canonical nucleosomes are associated with active histone

modifications, and therefore non-canonical nucleosomes are located at both expressed and

silent genes (Figure 3E).

Un-stable nucleosomes at the promoter and their function in the regulation of gene

expression

As we observed that un-stable nucleosomes are often occurring directly upstream of the TSS

(Figure 1G and Figure 3C) of expressed genes, we wanted to investigate their role in gene

regulation. Thus, we divided all expressed genes into two groups: 1) those which contained

an un-stable nucleosome 200 bp upstream of the TSS (n=1,588) and 2) those which had

instead a nucleosome depleted region (NDR) in the promoter (n=5,363) (Figure 4A). To define

NDR or un-stable nucleosome promoter it is absolutely required to use MNase chromatin

digestions coupled with histone immunoprecipitations, as non-histone proteins will dominate

the signal at the promoter in whole chromatin extractions (Figure S5). Overall, the expression

rate of the corresponding genes was similar in both groups and only slightly elevated for genes

carrying an un-stable nucleosome (Figure 4B, p-value = 2×10^{-15}). GO enrichment analysis

revealed that genes containing un-stable nucleosome in their promoter are stimulus or

developmentally regulated (Figure 4C). Furthermore, unstable nucleosome promoter were

enriched for Drosophila promoter motifs Unknown1-3 and M1BP motif, but depleted in

Initiator, TATA-box, and Unkown4 motif (Figure 4D). As M1BP is a known factor to regulate

RNA Pol II pausing depending on nucleosome positioning (Li and Gilmour, 2013), which

provides an efficient way to induce gene transcription rapidly or synchronously in response

to developmental and environmental signals, we tested whether Pol II pausing differed

between NDR or unstable nucleosome promoter using nascent RNA sequencing data (Fuda *et*

al., 2015). In agreement with M1BP motif enrichment, unstable nucleosome promoter show

a pronounced association with paused Pol II directly downstream of the TSS and higher

pausing index rates compared to NDR promoter enabling a fast transition into productive

transcription at responsive and developmental genes. (Figure 4E-D).

M1BP binds to un-stable nucleosomes

We next assessed the relationship between the binding of the transcription pausing factor

M1BP and nucleosome properties. Here, we investigated nucleosome positioning,

accessibility, and stability at annotated M1BP binding sites (Zouaz *et al.*, 2017). We found that

M1BP binding sites are surrounded by hyper-accessible nucleosomes (Figure 5A-C). Directly

in the binding center of M1BP, we observed a clear enrichment of un-stable nucleosomes

(Figure 5D-F) suggesting that M1BP does not bind to nucleosome-free DNA but rather to DNA

still co-occupied with histones. Nevertheless, histone DNA interactions seemed to be

weakened upon M1BP binding making them sensitive to internal MNase digestions.

Evaluating the robustness of the nucMACC pipeline

Previously four or more MNase digestion conditions have been used to determine chromatin

accessibility scores (Mieczkowski *et al.*, 2016; Chereji, Bryson and Henikoff, 2019). We sought

to determine the minimum number of titration points per sample to obtain reliable and

robust results with our pipeline. We compared the nucMACC and sub-nucMACC scores of two

or four titration points of the same experimental dataset. We observed that the results of

only two different MNase conditions are highly correlated with 4 titration points, as long as

the difference in the MNase concentration pairs is at least 10-fold (Figure 6A and B; Figure

S6A an B). In addition, we observed only consistent results using the lowest available MNase

concentration to detect un-stable nucleosomes using the sub-nucleosomal fragments. At

higher concentrations, fragments of unstable nucleosomes are often lost due to their fragile

nature.

Next, we addressed the importance of sequencing depth on the results of the nucMACC

pipeline, as MNase-seq analysis requires a high coverage of the genome. To obtain a genome

independent quantification we normalized the sequencing depth to the number of fragments

per kilobase of the mappable genome (FPKG):

$$Fragments\ per\ kilobase\ of\ genome\ (FPKG) = \frac{number\ of\ sequenced\ fragments}{\frac{effective\ genome\ size}{1,000}}$$

We found that using half of the fragments, with coverage of 122 FPKG preserves 73% of the

mono-nucleosome positions, and the nucMACC score correlation is reasonably high (R=0.88)

(Figure 6C and Figure S6C). Whereas further lowering the sequencing depth to 61 FPKG,

results in a considerable loss of called mono-nucleosome positions (25%), albeit the nucMACC

score correlation remains stable at overlapping nucleosome positions, with a correlation of

0.80. For sub-nucleosomes, starting with 244 FPKG results already at low coverage, since the

sub-nucleosomal fragments are underrepresented in mild MNase digestions. Reducing the

coverage to 183 FPKG (75% of sequencing fragments) amounts to a decrease in called sub-

nucleosomes to 49% (Figure 6D and Figure S6D). Nonetheless, the correlation between sub-

nucMACC scores remains very high at overlapping positions. Below 183 FPKG coverage, the

sub-nucleosome analysis produces an insufficient amount of called sub-nucleosome positions

(Figure 6D).

In summary, we recommend a minimum sequencing depth of 180 FPKG or higher to be used

when analyzing nucleosome stability.

Ultimately, we tested the reliability and robustness of nucMACC and sub-nucMACC scores

and tested if the differential amount of nucleosomes released at various MNase titrations

would bias the scores. Therefore, we ran our pipeline on a published dataset with spiked-in

nucleosomes and compared the results with or without spike-in information. Remarkably, we

find almost no difference in nucMACC scores when comparing the pipeline with or without

spike-in information for mono- (Figure 6E) and sub-nucleosomes (Figure 6F).

In conclusion, the nucMACC pipeline doesn't require spike-ins and is robust and reproducible

with only two MNase titrations.

Discussion:

In the current study, we set out to create an optimized MNase-seq pipeline and characterize

nucleosomes based on their accessibility and stability. We provide a unique way to select

nucleosome positions showing extraordinary accessibility and stability properties. Hyper-

accessible nucleosomes were enriched in promoter regions, whereas hypo-accessible

nucleosomes showed enrichment in intergenic regions. We conclude most nucleosomes have

similar levels of accessibility, with local effects of hyper- and hypo- accessibility, which has

been also previously observed in human cell lines (Schwartz *et al.*, 2018).

The nucMACC pipeline provides the first systematic and genome wide approach to detect un-
stable nucleosomes. We find un-stable nucleosomes enriched on active promoter of
expressed genes, while non-canonical nucleosomes are enriched at intergenic and exonic
regions of both silent and expressed genes. Furthermore, we show that some non-canonical
nucleosomes exhibit shorter DNA length and are resistant to high MNase digestions.
However, we cannot discriminate between nucleosome structures, such as hemisomes,
hexasomes or lexasomes. Intriguingly we find a special group of un-stable nucleosomes
enriched at TSS of promoters marked with M1BP TF-specific motif (Li and Gilmour, 2013).
Undeniably, we see a high enrichment of un-stable nucleosomes at M1BP binding sites. This
made us speculate that un-stable nucleosomes are involved in RNA pol-II pausing, which we
confirmed with GRO-seq data and we detected a clear difference in RNA pol-II pausing rate
compared to nucleosome depleted promoter. We hypothesize that un-stable nucleosomes
may stimulate binding properties of specific TF, despite nucleosomes normally occluding
binding. This may enable a rapid response to stimuli and developmentally regulated gene
expression. The un-stable nucleosome represents a gene regulation facet, adding to the
complexity of eukaryotic gene regulation.

Moreover, we provide a robust pipeline and guidelines for the experimental setup of the
MNase-seq protocol by analyzing different datasets with two or four MNase titrations per
sample, with or without spike-in information, different sequencing depths, and the
correlation between replicates, and correlation between different anti-histone antibody
pulldowns. In short, we show that only two MNase titrations are required, albeit at a high
sequencing depth. A minimum sequencing depth of 122 FPKG per sample for mono- and 244
FPKG for sub-nucleosomes is recommended to attain high-quality results. Of course, this
depends on the experimental design. For example, if mono- and sub-nucleosome fractions
are specifically enriched, such as through gel excision, the sequencing depth required per
sample will be lower.

For nucleosome analysis, applying an anti-histone ChIP after MNase treatment is crucial to
differentiate between histone and non-histone proteins bound to DNA, which cannot be
guaranteed just by fragment size selection.

We also analyzed the nucMACC scores with or without spike-in information. Spike-ins are
typically used to compare antibody pulldowns with different efficiencies or to make an
absolute quantification of RNA expression or, in our case, nucleosome occupancy. We see no

improvement or difference in nucMACC scores when using spike-in information. In
conclusion, for determining absolute nucleosome occupancy, spike-ins are still compulsory
and should be included in the experimental design, whereas using relative differences in the
MNase fragment frequencies is sufficient to calculate the nucleosome accessibility and
stability scores. Sequencing depth seems to have a more detrimental effect on the accuracy
of nucMACC scores than the absence of spike-in information. In a recent publication, authors
suggest spike-ins are necessary to accurately estimate the nucleosome digestion rate and
subsequently accessibility scores (Chereji, Bryson and Henikoff, 2019). Unfortunately, we
could not directly compare their scores and ours in the same dataset since they did not
provide analyzed data tracks. Furthermore, we could not reproduce their results with the
available information about the workflow they used, and an automated pipeline was not
provided.

In summary, we provide a publicly available Nextflow pipeline for analyzing nucleosome
positions and nucleosomal properties using MNase titrations. We provide guidelines for the
experimental design of MNase-seq experiments. Lastly, using our pipeline on public MNase-
seq datasets in D. melanogaster we find a novel mechanism of gene regulation in eukaryotes,
via nucleosome stability modulation and RNA pol-II pausing rate.

METHODS:

Public datasets used in this study

Whole chromatin MNase-seq, MNase-ChIP-H3-seq, MNase-ChIP-H4-seq and RNA-seq
(GSE78984), MNase-ChIP-H2B-seq, MNase-ChIP-H3-seq (GSE69336), M1BP-ChIP-seq
(SRX3011239), GRO-seq (GSM3274626), and MNase-seq with spike-ins in S2 and Kc167 cell
lines (PRJNA528497) were downloaded from the European nucleotide archive (ENA)
database. Histone modification and TF ChIP-seq or ChIP-seq data were retrieved through
modENCODE, with the following accession numbers: H3K27ac (296), H3K27me3 (298),
H3K4me1 (304), H3K9me3 (313), RNA pol-II ChIP-seq (329), H3K36me3 (3189), H3K4me3
(3761), CBP TF ChIP-seq (858), GAF TF ChIP-seq (858). DNAse-seq (GSM5258764) data were
retrieved from the GEO database.

Processing of MNase-seq data

Raw sequencing reads were mapped to the UCSC Drosophila melanogaster dm3 genome,

using Bowtie2 with the following parameters --very-sensitive-local --no-discordant. Aligned

reads were filtered for their mapping quality (MAPQ > 30), fragment size (length 140 – 200

bp for mono-nucleosomes and length < 140 bp for sub-nucleosomes), and ambiguous

chromosomes and blacklisted regions were removed using DeepTools2 alignmentSieve. Often

there is a shift in the fragment sizes due to variations in MNase enzyme activity; thus, each

dataset should be assessed individually. To achieve higher sequencing depth, mono-

nucleosome positions were called on pooled MNase titrations, using DANPOS, with dpos -m

1 --extend 70 -c 162367812 -u 0 -z 1 -a 1 -e 1, parameters. Sub-nucleosome positions were

called using only the lowest digestion samples, with the following parameters: dpos -m 1 - -

extend 70 -z 70 -c 162367812 -u 0 -e 1. Here library size was normalized to effective dm3

genome size (162367812), paired-end reads centered to their midpoint and extended to 70

bp to obtain nucleosome-sized positions. Sub-nucleosome positions were further filtered by

two parameters: a non-overlapping position with a mono-nucleosome or an overlapping

position with an enriched signal over a mono-nucleosome signal. Here a cut-off of > 4x higher

signal in sub-nucleosomal data as opposed to mono-nucleosomal data was chosen.

NucMACC score calculation

Reads per sample per nucleosome position were counted using featureCounts, using a

custom SAF file with nucleosome positions and the following parameters -SAF -p -B -C -

largestOverlap. Information on the GC content of each nucleosome was added to the read

count file using BEDtools nuc. Nucleosome positions were obtained by converting the

DANPOS annotation file to SAF format using AWK. NucMACC scores were calculated using R

and DESeq2 R package. Firstly, raw fragments were filtered for sequencing coverage, with cut-

off > 30 fragments per mono-nucleosome and > 5 fragments for sub-nucleosomes. Then,

fragments were normalized based on the library size (counts per million, CPM) or to the

calculated library size based on normalization factors for spike-in analysis. Linear regression

was calculated using normalized fragment counts and log2 MNase concentrations. Finally,

nucMACC scores were normalized for the inherent GC bias using the locally weighted

scatterplot smoothing (LOESS) algorithm. Henceforth, the loess function implemented in R

was applied to estimate the influence of GC content on MNase concentration titrations, using

the slope of the linear regression line. The function predicts a local score based on the GC

content and trained values. The predicted scores were used as a normalization factor for the
final nucMACC scores. The nucMACC score can be positive or negative. For mono-
nucleosomes, the nucMACC score was multiplied by -1 for a more intuitive interpretation.
Here, a negative score represents hypo-accessible DNA, whereas a positive score represents
hyper-accessible DNA. For sub-nucleosomes, a negative score represents unstable
nucleosomes, whereas a positive score characterizes nucleosomes with non-canonical
structures.

NucMACC score cut-off
A universal cut-off of the nucMACC score was obtained by ranking nucleosomes based on the
score and the first derivative and followed by a local regression, which predicts the slope
based on the scores. The cut-off for differentially accessible nucleosomes is when the slope =
1; this happens twice for positive and negative scores.

Spike-in analysis
Yeast reads were aligned to the sacCer3 Saccharomyces cerevisiae genome and treated
analogously to the nucMACC pipeline. The number of reads per mono- and sub-nucleosome
fraction was counted, and normalization factors were calculated by dividing the number of
reads by 10e6.

Nucleosome characterization
Characterization of nucleosomes and figures used in the manuscript were done in R.
Nucleosomes were assigned to genes using bitr R package and
TxDb.Dmelanogaster.UCSC.dm3.ensGene annotation R package. Pathway enrichment of
genes was performed with clusterProfiler R package. For expression correlation with
nucleosome positioning, nucleosomes on promoter and gene body (exon and intron) regions
were used. For heatmaps, LSD R package was used. For figures ChIPpeakAnno, ChIPseeker
and ggplot2 R packages were used. For TF enrichment, HOMER findMotifsGenome was used,
with a custom BED file containing the positions of TSS-un-stable nucleosomes and the *D.*
melanogaster TF database.

Pausing index

Pausing index is defined as the ratio of the GRO-seq read density at TSS to gene body density.

It was calculated as read density +/-100 nt from TSS divided by read density +150 nt from TSS

to the TES.

Minimum sequencing coverage analysis

Samples were sub-samples using SAMtools view -bs 42.X, where X represents the percentage

to which the data was reduced to, and 42 as the seed number to obtain reproducible results.

Processing of RNA-seq data

Sequencing reads were aligned to the UCSC Drosophila melanogaster dm3 genome using

STAR   v2.7.2a   with   the   following   parameters   --outFilterType   BySJout   --

outFilterMultimapNmax 20 --alignSJoverhangMin 8 --alignSJDBoverhangMin 1 --

outFilterMismatchNmax 999 --outFilterMismatchNoverReadLmax 0.04 --alignIntronMin 20 --

alignIntronMax 1000000 – alignMatesGapMax. Aligned reads were filtered for their mapping

quality (MAPQ > 30), using SAMtools v1.9. Reads per exon were counted using featureCounts

v1.6.2 and parameters -C -p -B -t exon -g gene_id.

{Bibliography}
Baldi, S. *et al.* (2018) 'array regularity and spacing by nanopore sequencing', *Nature*
Structural & Molecular Biology, 25(September). Available at:
https://doi.org/10.1038/s41594-018-0110-0.
Baldi, S., Korber, P. and Becker, P.B. (2020) 'Beads on a string—nucleosome array
arrangements and folding of the chromatin fiber', *Nature Structural and Molecular Biology*.
Nature Research, pp. 109–118. Available at: https://doi.org/10.1038/s41594-019-0368-x.
Buenrostro, J.D. *et al.* (2013) 'Transposition of native chromatin for fast and sensitive
epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position',
Nature Methods, 10(12), pp. 1213–1218. Available at: https://doi.org/10.1038/nmeth.2688.
Chereji, R. V., Bryson, T.D. and Henikoff, S. (2019) 'Quantitative MNase-seq accurately maps
nucleosome occupancy levels', *Genome biology*, 20(1), p. 198. Available at:
https://doi.org/10.1186/s13059-019-1815-z.
Chereji, R. V. and Clark, D.J. (2018) 'Major Determinants of Nucleosome Positioning',
Biophysical Journal, 114(10), pp. 2279–2289. Available at:
https://doi.org/10.1016/J.BPJ.2018.03.015.
Chereji, R. V, Ocampo, J. and Clark, D.J. (2016) 'MNase-sensitive Complexes in Yeast :
Nucleosomes and Non-histone Barriers', (December), pp. 565–577. Available at:
https://doi.org/10.1016/j.molcel.2016.12.009.
Chereji, V. *et al.* (2015) 'Genome-wide profiling of nucleosome sensitivity and chromatin
accessibility in Drosophila melanogaster', pp. 1–16. Available at:
https://doi.org/10.1093/nar/gkv978.

Dalal, Y. *et al.* (2007) 'Structure, dynamics, and evolution of centromeric nucleosomes',
Proceedings of the National Academy of Sciences of the United States of America, 104(41),
pp. 15974–15981. Available at: https://doi.org/10.1073/PNAS.0707648104.

Davey, C.A. *et al.* (2002) 'Solvent mediated interactions in the structure of the nucleosome
core particle at 1.9 a resolution', *Journal of molecular biology*, 319(5), pp. 1097–1113.
Available at: https://doi.org/10.1016/S0022-2836(02)00386-8.

Diermeier, S. *et al.* (2014) 'TNFα signalling primes chromatin for NF-κB binding and induces
rapid and widespread nucleosome repositioning.', *Genome biology*, 15(12), p. 536. Available
at: https://doi.org/10.1186/s13059-014-0536-6.

Fuda, N.J. *et al.* (2015) 'GAGA Factor Maintains Nucleosome-Free Regions and Has a Role in
RNA Polymerase II Recruitment to Promoters', *PLOS Genetics*, 11(3), p. e1005108. Available
at: https://doi.org/10.1371/JOURNAL.PGEN.1005108.

Hasson, D. *et al.* (2013) 'The octamer is the major form of CENP-A nucleosomes at human
centromeres', *Nature Structural & Molecular Biology 2013 20:6*, 20(6), pp. 687–695.
Available at: https://doi.org/10.1038/nsmb.2562.

Henikoff, J.G. *et al.* (2011) 'Epigenome characterization at single base-pair resolution'.
Available at: https://doi.org/10.1073/pnas.1110731108/-
/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1110731108.

Hu, G. *et al.* (2011) 'Regulation of nucleosome landscape and transcription factor targeting
at tissue-specific enhancers by BRG1', *Genome research*, 21(10), pp. 1650–1658. Available
at: https://doi.org/10.1101/GR.121145.111.

Kaplan, N. *et al.* (2009) 'The DNA-encoded nucleosome organization of a eukaryotic
genome', *Nature*, 458(7236), pp. 362–366. Available at:
https://doi.org/10.1038/nature07667.

Kireeva, M.L. *et al.* (2002) 'Nucleosome remodeling induced by RNA polymerase II: loss of
the H2A/H2B dimer during transcription', *Molecular cell*, 9(3), pp. 541–552. Available at:
https://doi.org/10.1016/S1097-2765(02)00472-0.

Kornberg, R.D. (1974) 'Chromatin structure: A repeating unit of histones and DNA', *Science*,
184(4139), pp. 868–871. Available at: https://doi.org/10.1126/science.184.4139.868.

Kubik, S. *et al.* (2015) 'Nucleosome Stability Distinguishes Two Different Promoter Types at
All Protein-Coding Genes in Yeast', *Molecular Cell*, 60(3), pp. 422–434. Available at:
https://doi.org/10.1016/j.molcel.2015.10.002.

Li, J. and Gilmour, D.S. (2013) 'Distinct mechanisms of transcriptional pausing orchestrated
by GAGA factor and M1BP, a novel transcription factor', *EMBO Journal*, 32(13), pp. 1829–
1841. Available at: https://doi.org/10.1038/EMBOJ.2013.111.

Liu, Y. *et al.* (2019) 'FACT caught in the act of manipulating the nucleosome', *Nature 2019
577:7790*, 577(7790), pp. 426–431. Available at: https://doi.org/10.1038/s41586-019-1820-
0.

Luger, K. *et al.* (1997) 'Crystal structure of the nucleosome core particle at 2.8 Å resolution',
Nature, 389(6648), pp. 251–260. Available at: https://doi.org/10.1038/38444.

Mieczkowski, J. *et al.* (2016) 'MNase titration reveals differences between nucleosome
occupancy and chromatin accessibility.', *Nature communications*, 7(May), p. 11485.
Available at: https://doi.org/10.1038/ncomms11485.

Olins, A.L., Douglas Carlson, R. and Olins, D.E. (1975) 'Visualization of chromatin
substructure: ν bodies', *Journal of Cell Biology*, 64(3), pp. 528–537. Available at:
https://doi.org/10.1083/jcb.64.3.528.

Prasad, R. *et al.* (2016) 'Coordinated Action of Nap1 and RSC in Disassembly of Tandem
Nucleosomes', *Molecular and cellular biology*, 36(17), pp. 2262–2271. Available at:
https://doi.org/10.1128/MCB.00195-16.
Ramachandran, S., Ahmad, K. and Henikoff, S. (2017) 'Transcription and Remodeling
Produce Asymmetrically Unwrapped Nucleosomal Intermediates', *Molecular Cell*, 68(6), pp.
1038-1053.e4. Available at: https://doi.org/10.1016/j.molcel.2017.11.015.
Rhee, H.S. *et al.* (2014) 'Subnucleosomal structures and nucleosome asymmetry across a
genome', *Cell*, 159(6), pp. 1377–1388. Available at:
https://doi.org/10.1016/j.cell.2014.10.054.
Schones, D.E. *et al.* (2008) 'Dynamic Regulation of Nucleosome Positioning in the Human
Genome', *Cell*, 132(5), pp. 887–898. Available at: https://doi.org/10.1016/j.cell.2008.02.022.
Schwartz, U. *et al.* (2018) 'Characterizing the nuclease accessibility of DNA in human cells to
map higher order structures of chromatin', *Nucleic Acids Research*, 47(3), pp. 1239–1254.
Available at: https://doi.org/10.1093/nar/gky1203.
Silberhorn, E. *et al.* (2016) 'Plasmodium falciparum Nucleosomes Exhibit Reduced Stability
and Lost Sequence Dependent Nucleosome Positioning', *PLOS Pathogens*. Edited by K.W.
Deitsch, 12(12), p. e1006080. Available at: https://doi.org/10.1371/journal.ppat.1006080.
Struhl, K. and Segal, E. (2013) 'Determinants of nucleosome positioning', *Nature Publishing
Group*, 20(3), pp. 267–273. Available at: https://doi.org/10.1038/nsmb.2506.
Teif, V.B. *et al.* (2012) 'Genome-wide nucleosome positioning during embryonic stem cell
development', *Nature Structural & Molecular Biology*, 19(11), pp. 1185–1192. Available at:
https://doi.org/10.1038/nsmb.2419.
Valouev, A. *et al.* (2011) 'Determinants of nucleosome organization in primary human cells',
Nature, 474(7352), pp. 516–520. Available at: https://doi.org/10.1038/nature10002.
Vierstra, J. *et al.* (2014) 'Coupling transcription factor occupancy to nucleosome architecture
with DNase-FLASH', 11(1). Available at: https://doi.org/10.1038/NMETH.2713.
Voong, L.N. *et al.* (2016) 'Insights into Nucleosome Organization in Mouse Embryonic Stem
Cells through Chemical Mapping', *Cell*, 167(6), pp. 1555-1570.e15. Available at:
https://doi.org/10.1016/j.cell.2016.10.049.
Weiner, A. *et al.* (2010) 'High-resolution nucleosome mapping reveals transcription-
dependent promoter packaging', *Genome research*, 20(1), pp. 90–100. Available at:
https://doi.org/10.1101/GR.098509.109.
Whyte, W.A. *et al.* (2013) 'Master transcription factors and mediator establish super-
enhancers at key cell identity genes', *Cell*, 153(2), pp. 307–319. Available at:
https://doi.org/10.1016/J.CELL.2013.03.035/ATTACHMENT/7BB94B8B-BF47-473D-9788-
A9CE552FB4BF/MMC9.XLSX.
Zouaz, A. *et al.* (2017) 'The Hox proteins Ubx and AbdA collaborate with the transcription
pausing factor M1BP to regulate gene transcription', *The EMBO Journal*, 36(19), pp. 2887–
2906. Available at: https://doi.org/10.15252/EMBJ.201695751.
635

Figures and Legends:

Figure 1

(A) Schematic representation of differential MNase chromatin digestion. Under mild MNase conditions, chromatin is partially digested and nucleosomes (green, hyper-accessible) from accessible regions are preferentially released (upper panel). At extensive MNase conditions, chromatin is completely digested to nucleosome core particles and MNase-sensitive nucleosomes (blue, un-stable) are lost. (B) Fragment size distribution of MNase titrations. Fragments are selected based on the fragment size and grouped into sub- (< 140 bp, light red section) or mono-nucleosomes (140-200 bp, purple section). (C) and (E) average fragment frequencies at the TSS. (D) and (J) Genome browser snapshot showing the output of the nucMACC pipeline. nucMACC scores are deduced from mono-nucleosomal fragments (D) and sub-nucMACC scores from sub-nucleosomal fragments, respectively (J). (E) Heatmaps sorted by gene expression showing nucleosome positioning (left), accessibility (middle, nucMACC scores), and stability (right, sub-nucMACC scores) at TSS.

Figure 2

(A) Selection of hyper- (green) and hypo-accessible (purple) nucleosomes based on nucMACC

scores. Nucleosomes were ranked by nucMACC score. X- and y-axis were normalized to a

range of one. Cut-offs were deduced from the slope of the LOESS curve fit. Nucleosomes

before the point, where the slope goes the first time below 1 (purple dashed line) were

classified as hypo-accessible. Nucleosomes after the point where the slope of the curve

exceeds one again (green dashed line), were classified as hyper-accessible nucleosomes. (B)

Correlation of DNase-seq sites with hyper-/hypo-accessible nucleosomes. (C) Enrichment of

hyper-/hypo-accessible nucleosomes at genomic features. (D) K-means clustering (n=4) of

histone modifications at hyper-/hypo-accessible sites. Centroid values of each cluster are

indicated by the color code. The relative number of nucleosomes represented in each cluster

is indicated on the right side. Repressive histone modifications are highlighted in orange and

active in green, respectively. The pie chart depicts the fraction of nucleosomes assigned to

either active or repressive histone clusters. (E) Gene expression levels of genes exhibiting

hyper-/hypo-accessible nucleosomes at distinct features. Promoter was defined as 500 bp

upstream of the TSS to 300 bp downstream.

Figure 3

(A) Selection of un-stable (blue) and non-canonical (red) nucleosomes based on nucMACC

scores. Nucleosomes were ranked by sub-nucMACC score. X- and y-axis were normalized to a

range of one. Cut-offs were deduced from the slope of the LOESS curve fit. Nucleosomes

before the point, where the slope goes the first time below 1 (blue dashed line) were classified

as un-stable. Nucleosomes after the point where the slope of the curve exceeds one again

(red dashed line), were classified as non-canonical nucleosomes. (B) Correlation of DNase-seq

sites with un-stable and non-canonical nucleosomes. (C) Enrichment of un-stable or non-

canonical nucleosomes at genomic features. (D) K-means clustering (n=5) of histone

modifications at un-stable or non-canonical sites. Centroid values of each cluster are indicated

by the color code. The relative number of nucleosomes represented in each cluster is

indicated on the right side. Repressive histone modifications are highlighted in orange, active

in green, respectively. The bivalent cluster enriched for both repressive and active histone

modifications is indicated in grey. The pie chart depicts the fraction of nucleosomes assigned

to either active, repressive, or bivalent histone clusters. (E) Gene expression levels of genes

exhibiting un-stable or non-canonical nucleosomes at distinct features. Promoter was defined

as 500 bp upstream of the TSS to 300 bp downstream.

Figure 4

(A) Average fragment frequencies at the TSS of expressed genes either containing an un-

stable nucleosome (n=1,588, right) or exhibiting a nucleosome depleted region (NDR)

(n=5,368, left) directly upstream of the TSS. (B) Violin plot showing expression levels of genes

either containing an un-stable nucleosome (n=1,588, right) or exhibiting a nucleosome

depleted region (NDR) (n=5,368, left) directly upstream of the TSS. (C) Enrichment of

biological processes associated with genes characterized by an un-stable nucleosome

promoter. (D) Motif enrichment analysis in the un-stable nucleosome (left) or NDR (right)

promoter. (E) Median GRO-seq profile showing nascent RNA abundance. Regions upstream

of TSS, downstream of TTS and the first 200 bp downstream of TSS are unscaled. The

remaining gene body is scaled to have the same length for each gene. (F) Boxplots showing

the pausing index based on the GRO-seq signal. Pausing index was calculated as the fold

change of the GRO-seq signal in the first 200 bp downstream of the TSS versus the remaining

gene body.

Figure 5

(A) and (D) average fragment frequencies of mono- (A) and sub-nucleosomal (D) fragments at

M1BP bound sites. (B) and (E) Heatmap showing nucMACC (B) or sub-nucMACC (E) scores at

M1BP bound sites. (C) and (F) Distribution of hyper-/hypo-accessible (C) or un-stable and non-

canonical (F) nucleosomes at M1BP bound sites.

Figure 6

(A-B) Correlation of nucMACC (A) or sub-nucMACC (B) scores between samples with two or

four MNase titrations. The pearson correlation coefficient is indicated (C-D) The number of

called nucleosomes based on sequencing depth (FPKG – fragments per kilobase of genome).

Correlation of nucMACC (C) or sub-nucMACC (D) scores between samples with different

sequencing depths is illustrated at the right. (E-F) Scatterdensityplot showing correlation of

nucMACC (E) or sub-nucMACC (F) scores with or without spike-in information. The Pearson

correlation coefficient is indicated.