Starvation sensing by mycobacterial RelA/SpoT homologue through constitutive surveillance of translation

Yunlong Li[1#], Soneya Majumdar[2#], Ryan Treen[1,3#], Manjuli R. Sharma[2], Jamie Corro[1,3], Howard B. Gamper[4], Swati R. Manjari[2], Jerome Prusa[5], Nilesh K. Banavali[2], Christina L. Stallings[5], Ya-Ming Hou[4], Rajendra K. Agrawal[2,3*], Anil K. Ojha[1,3*]

1. Division of Genetics, Wadsworth Center, New York State Department of Health, Albany, NY 12208, USA
2. Division of Translational Medicine, Wadsworth Center, New York State Department of Health, Albany, NY 12237, USA
3. Department of Biomedical Sciences, School of Public Health, University at Albany, Albany, NY 12208, USA
4. Department of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia, PA 19107, USA
5. Department of Molecular Microbiology, Washington University in St. Louis School of Medicine, St. Louis, MO 63110, USA

[#]Equal contribution

*Correspondence: anil.ojha@health.ny.gov;
                          rajendra.agrawal@health.ny.gov

Abstract:

The stringent response, which leads to persistence of nutrient-starved mycobacteria, is induced

by activation of the RelA/SpoT homologue (Rsh) upon entry of a deacylated-tRNA in a

translating ribosome. However, the mechanism by which Rsh identifies such ribosomes *in vivo*

remains unclear. Here, we show that conditions inducing ribosome hibernation result in loss of

intracellular Rsh in a Clp protease-dependent manner. This loss is also observed in non-starved

cells using mutations in Rsh that block its interaction with the ribosome, indicating that Rsh

association with the ribosome is important for Rsh stability. The cryo-EM structure of the Rsh-

bound 70S ribosome in a translation initiation complex reveals unknown interactions between

the ACT domain of Rsh and components of the ribosomal L7/L12-stalk base, suggesting that

the aminoacylation status of A-site tRNA is surveyed during the first cycle of elongation.

Altogether, we propose a surveillance model of Rsh activation that originates from its

constitutive interaction with the ribosomes entering the translation cycle.

Significance:

Bacteria persist under nutrient starvation by activating RelA/SpoT homologue (Rsh), which synthesizes a growth regulating alarmone, ppGpp. Rsh is activated specifically upon recognizing a translation elongation complex with deacylated tRNA at the A-site. It is however unclear how Rsh identifies such a complex in vivo. We show here that conditions inducing ribosome hibernation in mycobacteria cause loss of intracellular Rsh, implying that association with translating ribosomes is necessary for intracellular stability of Rsh. Using structural analysis of Rsh-bound 70S translation initiation complex, we propose here that mycobacterial Rsh identifies a Rsh-activating ribosomal complex by constitutively surveying the ribosome entering the translation cycle at the early elongation stage.

Introduction:

The stringent response in bacteria is a programmed adaptation to nutrient starvation facilitated by the secondary messengers, guanosine tetra- or penta-phosphates [collectively called (p)ppGpp] (1-4). This response promotes long-term survival and persistence of mycobacteria under various stresses including antibiotics (5-10). The primary function of (p)ppGpp is to reprogram the global gene expression in cells either directly by binding to RNA polymerase (3, 11) or by binding to other cellular targets (12-14).

The level of (p)ppGpp is primarily regulated by a GDP/GTP pyrophosphokinase and a GDP/GTP pyrophosphohydrolase, encoded either as two enzymes RelA and SpoT in β- and γ-proteobacteria (3), or as a single protein called Rel in other species (15). Further insights into the domain architectures of RelA/Rel and the mechanism of its activation have been obtained from the structures of either the free enzyme or in complex with the ribosome, mRNA and A- and P-site tRNAs (16-20). RelA/Rel has five distinct domains (Fig. 1A): i) N-terminal domain possessing the (p)ppGpp synthase and hydrolase activities, ii) TGS (ThrRS, GTPase, SpoT) domain that interacts with 3' CCA tail of deacylated-tRNA, iii) alpha helical domain (AHD), iv) zinc-binding domain (ZBD) that contacts the A-site finger (ASF) on the 23S rRNA and S19 at the inter-subunit interface, and v) the ACT domain (Aspartate Kinase, Chorismate and TyrA) occupying a cavity in the large 50S ribosomal subunit (or LSU) (17-19). The interaction between the TGS domain and 3' CCA tail of the deacylated-tRNA stabilizes the tRNA in a distorted state at the A-site, called the A/R state, while pinning the domain against the small 30S ribosomal subunit (or SSU) and extending the flexible catalytic N-terminal domain of RelA in an open and active conformation (17-19).

The mechanism by which RelA finds a stalled ribosome complex in a nutrient starved cell is unclear. In the currently adopted 'hopping model', diffusible RelA samples translation elongation complexes and binds to a stalled ribosome with deacylated A-site tRNA in the A/R state, and upon catalyzing (p)ppGpp synthesis the enzyme 'hops' to another stalled complex

(21-23). The hopping model further hypothesizes that an inactive RelA in non-starved cells

acquires a ribosome-free, closed conformation, in which the catalytic activity is autoinhibited by

the CTD, either through intermolecular (24-26) or intramolecular interactions (27-32). Winther et

al. further proposed that free cytosolic RelA in *E. coli* binds to deacylated-tRNA before binding

to the translating ribosome (22), although this model is challenged by the evidence of RelA

bound to the ribosome without any tRNA at the A-site (18). Moreover, single-molecule tracking

of RelA-fluorescent protein suggests constitutive association between RelA and ribosome in

non-starved *E. coli* cells (33). Recognizing these discrepancies, Takada et al. offered a refined

hopping model in *B. subtilis* in which activation of the stringent response appears to be initiated

by a weak interaction between Rel and the ribosome before the complex is stabilized by the

entry of deacylated-tRNA at the A-site, although they found RelA-ribosome interaction in *E. coli*

to be too weak to universally validate their model (23). These studies while revealing subtle

differences in the properties of RelA homologues across bacterial species, also underscore the

limitations of biochemical assays in modeling a highly dynamic *in vivo* interaction between the

stringent factor and the translation machinery.

Mycobacteria encode a single dual-function enzyme, RelA/SpoT homologue (henceforth

called Rsh), with both kinase and hydrolase activities located in distinct domains of the enzyme

(25, 26, 34-36). The optimal kinase activity of mycobacterial Rsh is observed in a Rsh Activating

Complex (RAC) comprising mRNA, deacylated-tRNA and the ribosome (37). The ribosome in

Mycobacterium smegmatis and *Mycobacterium tuberculosis* (Mtb) is remodeled under relatively

moderate zinc starvation, and subsequently targeted for hibernation under severe zinc

starvation (38, 39). Ribosome remodeling involves replacement of ribosomal proteins containing

the CXXC motif (called C+ r- proteins) with their paralogs without the motif (called C- r-proteins)

(38, 40). The C- ribosome is subsequently targeted for hibernation by mycobacterial protein Y

(Mpy), with assistance from a zinc-sensing protein called Mpy recruitment factor (Mrf) (38, 39).

A zinc-bound form of Mrf is sensitive to degradation by Clp protease, such that Mrf

accumulation increases with decreasing concentration of zinc (39).

Investigating the interplay between the stringent response and ribosome hibernation in

mycobacteria, we observed that interaction with the ribosome is necessary for the stability of

intracellular Rsh in non-starved cells. The structure of Rsh-bound 70S ribosome in a translation

initiation complex allows us to propose a new mechanism of starvation sensing by

mycobacterial Rsh, in which its activation arises from its unique interaction with the ribosome

entering the translation cycle.

Results:

Mrf attenuates the stringent response in zinc-starved mycobacteria

We tested how a zinc-limiting growth condition that induces ribosome hibernation in *M.*

smegmatis would impact ppGpp synthesis. After 96-hours of growth in high- or low-zinc

Sauton's medium, wild-type *M. smegmatis* cells were exposed for 3 hours to macronutrient

starvation in tris-buffered saline with Tween-80 (TBST) to induce the stringent response to a

measurable level (6). Rsh-dependent accumulation of ppGpp upon TBST starvation of the high-

zinc culture was significantly greater than the low-zinc culture (Fig. 1B). To distinguish the

contribution of macronutrient depletion from zinc limitation in TBST, we examined the stringent

response in cells from high-zinc cultures after exposure to metal-chelated or zinc-supplemented

TBST. Metal chelation did not impact the ppGpp synthesis, indicating that macronutrient

starvation in TBST was the primary inducer of the stringent response (Fig. 1C).

Interaction with ribosomes is critical for the stringent response (21), we therefore

hypothesized that the attenuated stringent response in low-zinc cultures was likely due to zinc-

responsive changes in the ribosome structure and/or function. We therefore analyzed the Rsh

occupancy on the ribosome in high- and low-zinc cultures. Rsh occupancy on the ribosomes

purified from low-zinc cells was significantly lower than that from high-zinc cells (Fig. 1D).

Unexpectedly, the difference in Rsh abundance on the ribosome correlated tightly with the

levels in the total cellular lysate (Fig. 1D). Moreover, lack of any other detectable smaller Rsh

isoforms (Fig S1) suggests total loss of intracellular Rsh. Together, figures 1B-D show that

physiological conditions developed during zinc starvation decreases the abundance of Rsh and

thus its interaction with the ribosome, thereby diminishing the stringent response.

We next investigated the effect of two zinc-responsive physiological changes in the

ribosome, remodeling and hibernation, in altered Rsh stability and Rsh-ribosome interaction.

The Zur-mediated C+ to C- remodeling of the ribosome induced by zinc starvation did not

impact the Rsh-ribosome interaction: Rsh occupancy on the ribosome from a high-zinc culture

of wildtype was comparable to Δzur and a strain constitutively synthesizing C- ribosomes (Fig.

S2). To determine the effect of zinc-responsive ribosome hibernation on Rsh-ribosome

interaction, we analyzed the impact of Mpy recruitment factor (Mrf) on Rsh stability, activity, and

its occupancy on the ribosome. Mrf binds to SSU and facilitates inhibition of translation and

growth retardation upstream to Mpy recruitment to the ribosome (39). In zinc-starved *M.*

smegmatis, deletion of *mrf* restored the levels of intracellular and ribosome-bound Rsh similar to

those in high-zinc wildtype cells*,* and the mutant phenotype was complemented by plasmid-

borne *mrf* (Fig. 2A-B). The increase in Rsh abundance in zinc-starved Δmrf also correlated with

the increase in ppGpp levels upon exposure to TBST (Fig. 2C). Together, these data suggest

that: i) the limiting intracellular free zinc concentration has little direct impact on the stringent

response, and ii) accumulation of Mrf induced by zinc depletion and likely downregulation of

translational activity decrease total Rsh abundance in the cell. Deletion of *rsh* did not impact

Mpy-ribosome interaction (Fig. S3), distinguishing mycobacteria from other species in which

ppGpp transcriptionally induces factors necessary for ribosome hibernation (41-43).

Ectopic expression of Mrf can reduce intracellular abundance of Rsh.

To further establish the relationship between Rsh abundance and Mrf accumulation, we

examined the impact of an ectopic expression of a Mrf variant (Mrf*) on Rsh levels in zinc-rich

cells. Mrf* has reduced zinc affinity and is recalcitrant to regulation by the Clp protease system

(39). Consequently, constitutive expression of Mrf* leads to its substantial accumulation in high-

zinc mycobacteria (39), presumably in a conformation that is capable of binding to the ribosome

and inhibiting both translation and cellular growth. The abundance of Rsh and its association

with the ribosome was determined upon acetamide-inducible expression of Mrf* in high-zinc

cultures of *M. smegmatis*. Figure 2D shows that the expression of Mrf* in high-zinc cells led to

concomitant loss of Rsh, both in the lysates (Fig. 2D) and on the ribosomes (Fig. 2E).

Because SpoT in *Salmonella enterica* was identified as a potential substrate of the Clp

protease system (44), we tested the possibility whether mycobacterial ClpP1P2 protease could

be involved in regulating the abundance of free Rsh in cells. We used a *M. smegmatis* strain,

which constitutively transcribed FLAG-tagged Mrf and harbored an anhydrotetracycline-

inducible CRISPR-Ca9 system for targeting ClpP1 (39), to test whether Rsh loss during Mrf

stabilization is Clp-dependent. Depletion of ClpP1 in high-zinc cells resulted in concomitant

accumulation of both Mrf and Rsh (Fig. 2F). Thus, loss of Rsh in cells during Mrf stabilization is

likely achieved through a Clp-dependent mechanism, although whether Rsh is a direct substrate

of Clp remains unknown.

Interaction with the ribosome is necessary for intracellular stability of Rsh

Above, we have established loss of Rsh in cells with progressively increasing amounts of Mrf

prior to ribosome hibernation, suggesting that availability of translating ribosome might be

important for Rsh stability. Moreover, similar levels of Rsh-bound ribosomes in high-zinc

cultures before and after nutrient starvation (Fig. S4), suggesting that there is a constant level of

Rsh-ribosome interaction regardless of the changes in the relative levels of (p)ppGpp. We

therefore tested whether interaction with the ribosome impacts the intracellular stability of Rsh.

E. coli RelA interacts with the ribosome primarily through its CTD containing a conserved CCHC

type zinc-finger in the ZBD, which makes extensive contacts with the inter-subunit region of the

70S ribosome, and ASF of the LSU (17, 45). Although the requirement of zinc in RelA-ribosome

interaction remains unclear, the zinc coordinating residues C612, C613, H634 and C638 (C666,

C667, H688 and C692 in *M. smegmatis Rsh*) of ZBD are conserved in mycobacteria and likely

acquire the same three-dimensional conformation as in *E. coli* RelA (Fig. 3A). Furthermore,

mutation of C633 in *M. tuberculosis* Rsh (C692 in *M. smegmatis*) impairs the enzyme *in vitro*

(25), supporting an important role of this residue in the Rsh activity. We therefore mutated the

key ZBD residues and studied the impact on the intracellular abundance of Rsh. The mutation

scheme closely mirrored the previous study (24), which also showed that mutation in the

neighboring D637 (D691 in *M. smegmatis*) had substantially less impact on protein function

compared to C638 (C692 in *M. smegmatis*). Substitution of C666 and C692 residues resulted in

substantial loss of Rsh in both the cell lysate and on the ribosome in actively growing *M.*

smegmatis cells in zinc-rich conditions (Fig. 3B). The presence of wildtype Rsh on the ribosome

was unambiguous (Fig. 3B), supporting the notion that Rsh-ribosome interaction in non-starved

cells is constitutive. Substitution in the adjacent D691 neither impacted the Rsh-ribosome

interaction nor the intracellular abundance of the protein (Fig. 3B). Recombinant expression of

these Rsh variants in *E. coli* did not significantly impact the protein yield (Fig. S5), indicating that

these mutations do not affect the intrinsic stability of the protein *in vivo*, and their properties are

likely determined by the intracellular environment of mycobacteria. To further test if the

mutations specifically disrupted Rsh-ribosome interaction in mycobacteria, we mixed

recombinant Rsh_{C666G} and Rsh_{C692F} variants with 70S ribosomes purified from Δ*rsh* mutant of *M.*

smegmatis and analyzed their occupancy on the ribosome. Ultracentrifugation on a 32%

sucrose cushion was used to separate ribosome-bound from unbound Rsh. Compared to the

wildtype Rsh, association with the ribosome was significantly compromised for both

recombinant proteins, Rsh_{C666G} and Rsh_{C692F} (Fig. 3C). We thus conclude that Rsh constitutively

interacts with the ribosome using its ZBD in non-starved cells, and this interaction is necessary

for the intracellular stability of the enzyme.

Rsh interacts with the IF2 bound 70S initiation complex

We next focused on understanding the stage of translation when Rsh would most likely interact

with the ribosome in a non-starved cell. We streamlined our approach based on the published

structures (17-19), which delineate permissive and non-permissive translation complexes for

RelA binding. While RelA can bind to a late translation-initiation intermediate (70S- fMet-

tRNA$_i^{Met}$ -mRNA with vacant A-site) (18), its binding to the ribosome during translation

elongation is expected to be sterically hindered by inter-subunit rotation (17-19). We therefore

hypothesized that mycobacterial Rsh would enter the translation initiation complex through one

of the free ribosomal subunits and likely remain on the complex until the start of the first cycle of

elongation. This hypothesis also implies that Rsh must remain associated with the ribosome

during its transition from initiation to pre-elongation stages, without affecting the overall

canonical conformation of the translation complex.

To investigate this possibility, we first determined the propensity of Rsh to interact with

free ribosomal subunits. Purified recombinant Rsh was mixed individually with each of the two

purified ribosomal subunits free of any canonical protein synthesis ligands and the

macromolecules were separated on a 32% sucrose cushion. Rsh appeared to preferentially co-

sediment with the LSU (Fig. 4A). We next asked if Rsh would remain in the translation complex

during its transitions through the various intermediate stages of initiation, especially given the

potential of a steric clash between Rsh and translation initiation factor 2 (IF2). The contact

points of IF2 on the 70S initiation complex significantly overlaps with those of Rsh (17-19, 46).

We therefore tested the ability of Rsh to interact with the ribosome in the presence of IF2, fMet-

tRNA$_i^{Met}$ and mRNA *in vitro*. Dissociated ribosomes (Fig. S6) were mixed with His$_6$-IF2, His$_6$-

Rsh, fMet-tRNA$_i^{Met}$, leadered mRNA, and either GTP or a non-hydrolysable analogue (GMP-

PNP). The assembled 70S complex purified on 10-40% sucrose density gradient (SDG) was

analyzed for the presence of IF2 and Rsh. The 70S ribosomes assembled in the presence of

GTP contained Rsh but had undetectable levels of IF2 (Fig. 4B), demonstrating that Rsh can

associate with the ribosome in the presence of the canonical ligands without interfering with

GTP-dependent release of IF2 (47). This property of mycobacterial Rsh is in sharp contrast to

B. subtilis Rel, which cannot form a stable complex with the ribosome without an A-site

deacylated-tRNA (23). Unexpectedly, both Rsh and IF2 were detected in 70S particles

assembled in the presence of GMP-PNP (Fig. 4C), suggesting that one or both proteins could

be occupying the ribosome in their non-canonical states.

Cryo-EM structure reveals unknown ribosome-Rsh interactions

To gain further insight into the Rsh-bound *M. smegmatis* 70S ribosome in a translation initiation

complex, we performed single-particle cryo-EM to determine the structure of the 70S ribosome

assembled in the presence of IF2, Rsh, fMet-tRNA$_i^{Met}$, leadered mRNA, and GMP-PNP. 3D

classification of the cryo-EM dataset (Figs. S7, S8, Table S3) revealed a unique class of Rsh-

bound ribosome in a classical non-rotated state, but none of the classes showed any density for

IF2. The Rsh-bound 3D class refined to 2.7 Å resolution showed densities only for two C-

terminal domains of Rsh, ZBD and ACT, along with a fMet-tRNA$_i^{fMet}$ bound at the ribosomal P

site, while all other domains were found to be disordered (Figs. 4D, 4E, S7 and S9). The ZBD

interacts with both the SSU and LSU: specifically, protein uS19 of the SSU and 23S rRNA helix

H38 of the LSU (also referred to as ASF) (Fig. 4E). These interactions resemble those reported

in the structures of *E. coli* RelA and *B. subtilis* Rsh on the ribosome (18, 20). The Rsh-ACT

domain occupies the inter-subunit space and interacts with LSU proteins uL16, uL11, and the

23S rRNA helices H43-44 at the base of the protein L7/L12 stalk (Fig. 4E).

Previous structural studies suggest IF2 and RelA binding to the ribosome would be

mutually exclusive (18, 46). However, our data shows co-existence of the two proteins in a

reconstituted 70S translation initiation complex (Fig. 4C). To begin to resolve this issue, we

superimposed either our mycobacterial structure or the *E. coli* 70S ribosome-RelA structure (18)

with 70S-IF2-GMP-PNP initiation complexes (46). Indeed, RelA's alpha-helical domain (AHD)

would directly clash with domain VI-C1 of IF2 (Fig. 4F). In tertiary structure, the AHD of *E. coli*

RelA lies adjacent to the ACT domain while the ZBD is anchored to the ribosome via its

interactions with uS19 and ASF (18). In our structure the rotational angle between the ACT and

ZBD domains was ~14 degrees greater than that in *E. coli* RelA (Fig. 4F) and AHD was

disordered, presumably displaced from its canonical position, thus avoiding a clash with domain

VI-C1 of IF2 (Fig. 4F). The ACT domain is known to exhibit some conformational variations with

respect to the ZBD domain in the previously published structures (18, 20), but an additional

rotation by 14 degrees is unprecedented. It is possible that the adopted conformation of the ACT

domain in our structure is to accommodate the exiting IF2 molecule upon completion of the

translation initiation phase.

Compared to the *E. coli* RelA and *B. subtilis* Rel, mycobacterial Rsh shows an overall

similarity in tertiary structures of ZBD and ACT domains, although the rotation of the ACT

domain translates into its shift by ~6 Å towards the LSU's L7/L12 stalk base (Fig. 5A, B). This

altered conformation of ACT in Rsh is stabilized through extensive interactions with the uL11

and the 23S rRNA helices H43/44 (Fig. 5C). Specifically, uL11 lysines, 89 and 96 interact with

Y791, D789 and H776 of Rsh-ACT. Further, Rsh-ACT S779, N783, D789 and Y788 interact with

the 23S rRNA helices H43-44 phosphate backbones. These interactions are in sharp contrast to

the *E. coli* ribosome-RelA structures, where uL11 interacts with the A/R tRNA but not directly

with RelA (18). Thus, our structure reveals a unique conformational state of the ribosome-bound

Rsh. Multiple sequence alignment of the amino acid sequences does not reveal any apparent

substitutions, insertions, or deletions that can explain the altered orientation of the ACT domain

in *M. smegmatis* Rsh (Fig. S10). Given that our structure was obtained in the presence of IF2-

GMP-PNP, a key difference from the *E. coli* ribosome-RelA complex, a possible role of IF2 in

inducing the rotation of ACT cannot be ruled out.

Interactions stabilizing Rsh in mycobacterial structure are primarily through LSU's ASF,

uL16, uL11, and rRNA helices H43/44, suggesting that Rsh associates primarily with the LSU,

corroborating the results of our binding experiments that show association of Rsh with free LSU.

The rRNA helices H43/44 and uL11 together constitute the L7/L12-stalk base of LSU and are

known to be mobile during elongation (48) and recycling (49) phases of translation. We thus

propose that intensive contacts between the L7/L12-stalk base and the Rsh-ACT domain would

transiently lock the ribosome in a stalled state before the start of the translation elongation

cycle, as a mechanism to survey the aminoacylation state of the incoming tRNA to the A site. A

direct contact of Rsh with uL11 is distinct from *E. coli* RelA, which binds to the ribosome in

uL11-independent manner, even though uL11 and a deacylated-tRNA at the A-site are required

for (p)ppGpp synthesis (21, 50-52). While the structural changes in *M. smegmatis* complex here

could be induced by the presence of IF2, we cannot exclude the possibility that these are

intrinsic *to M. smegmatis* Rsh.

The superimposed structures also reveal some species-specific differences in

Rsh/RelA/Rel, which could probably explain differences in their ribosome binding affinities (Fig.

S10). The ZBD domains interact with uS19 and ASF in all three cases, however, compared

to *M. smegmatis*, *B. subtilis* Rel has additional interactions — two with uS19, one with ASF, and

one with uS13 (Fig. S10). The loss of the additional interactions with uS19 and ASF in *M.*

smegmatis is likely due to substitution at corresponding positions with non-polar amino acids

(Fig. S10). However, *M. smegmatis* Rsh-ZBD could potentially interact with uS13 since it has an

Arg (R681) corresponding to K617 in *B. subtilis*. The *E. coli* RelA ZBD also has two additional

interactions compared to *M. smegmatis*, one with uS19 and the other with ASF (Fig. S10).

However, *E. coli* RelA ZBD has a Gln (Q627) corresponding to K617 and R681 in *B.*

subtilis and *M. smegmatis*, respectively, which might not interact with uS13 due to its shorter

side chain. Since the density corresponding to ZBD is relatively strong in our structure, we

designate it as the primary anchoring domain of mycobacterial Rsh.

Exclusion of Rsh from Mpy-bound ribosomes

The requirement of the ribosomes for the intracellular stability of Rsh, combined with its loss in

cells harboring Mrf and Mpy-bound hibernating ribosomes, suggests that Rsh is not able to bind

to hibernating ribosomes. Mpy and Rsh bind at two physically distinct sites on the unrotated

SSU of the ribosome (Fig. 6A). The Mpy N-terminal domain binds to the SSU spanning the

decoding center by interacting extensively with the 16S rRNA components that converge from

platform, body, and head regions at the SSU's neck (38). By contrast, the Rsh ZBD interacts

near the inter-subunit bridge B1a (53) forming components, including uS19 of the SSU and the

LSU's ASF (54) (Figs. 4E and 6A). Since the binding sites of the two factors do not overlap, we

superimposed the structures of Rsh- and Mpy-bound *M. smegmatis* 70S ribosomes to

understand the accessibility of Mpy-bound hibernating ribosome to Rsh binding. Both the

ribosomes are in an unrotated state, but the SSU head domain in the Rsh-bound structure is

tilted more towards the LSU in comparison to the Mpy-bound ribosome structure (Fig. 6B). We

predict that the untilted SSU head in the MPY-bound state will prevent stable interactions

between Rsh-ZBD and uS19 (Fig. 6C). ZBD is the primary anchoring domain of Rsh (18, 20),

thus, any alteration in the bridge B1a configuration in the Mpy-bound state is likely to directly

inhibit Rsh binding, thereby rendering ribosome-free labile Rsh. Thus, reduction in the

intracellular Rsh during ribosome hibernation could be a combined outcome of both a decrease

in the synthesis of new Rsh and the occlusion of pre-existing Rsh from the hibernating

ribosome.

Discussion:

The stringent response, characterized by Rsh-dependent synthesis of (p)ppGpp and

subsequent transcriptional reprogramming by the alarmone, is among the central adaptation

mechanisms in nutrient-starved mycobacteria. Yet, a significant knowledge gap exists in the

mechanism underlying starvation sensing by Rsh. Structural studies with *E. coli* RelA provide

mechanistic details on how Rsh is activated by the presence of deacylated-tRNA at the A- site

in a stalled translation-elongation complex. However, it remains unclear how Rsh would

encounter such a complex *in vivo*. Two models have emerged from recent studies: a) the

hopping model in which a Rel-deacylated-tRNA complex is formed prior to its entry at the A- site

of the elongation complex (22), and b) starvation-induced formation of a weak RelA-ribosome

complex prior to its stabilization by deacylated-tRNA (23). Both models however preclude RelA-

ribosome interaction in non-starved cells, which is inconsistent with the evidence of constitutive

RelA-ribosome interaction throughout the growth of *E. coli* and *P. aeruginosa* cells (33, 55).

Here, we propose a new model of Rsh recognition of a stalled translation complex in

mycobacteria. In this model, which we call the surveillance model, Rsh constitutively interacts

with ribosomes during the formation of the initiation complex, and its transition to the pre-

elongation stage (Fig. 7A). Consequently, entry of an amino acylated-tRNA would allow

elongation, accompanied by inter-subunit rotation that would displace Rsh from the complex

(Fig. 7B) (17-19). By contrast, entry of a deacylated tRNA into the A-site would trigger Rsh

activation and (p)ppGpp synthesis (Fig. 7D). Free Rsh appears to be labile (Fig. 7G). To

maintain steady state levels Rsh must therefore associate with free LSU entering the translation

cycle (Fig. 7C), or another stalled elongation complex with deacylated-tRNA at the A- site (Fig.

7E). The latter interaction is expected to be uncertain and less frequent given the low

abundance of stalled elongation complexes relative to large abundance of free LSU. Given that

Rsh abundance in an actively growing cell is ~200-fold less compared to the ribosome (21), a

rapid rate of translation would likely maintain a sufficiently large pool of free LSU transitioning

between termination and initiation cycles, thereby ensuring a stable population of Rsh-ribosome

complex. The predictable fates of Rsh in the surveillance model renders the model more

stringent than either the original or the refined hopping model, both of which rely on random

encounter of Rsh with a deacylated-tRNA and/or a stalled ribosome.

The lack of Rsh under conditions such as those inducing ribosome hibernation further

supports the model that optimum protein synthesis and accessibility to LSU entering the

translation cycle is critical for the intracellular stability of Rsh (Fig. 7F). This loss of Rsh further

differentiates the physiological conditions for ribosome hibernation and the stringent response.

The stringent response is an upstream response before ribosome hibernation; a logical order

given that the stringent response reprograms gene expression to fine-tune metabolism, while

ribosome hibernation directly shuts down global protein synthesis. Moreover, restoration of

(p)ppGpp synthesis in a zinc-depleted Δmrf strain implies that Rsh-ribosome interaction and

Rsh activation are relatively less sensitive to intracellular depletion of free zinc. We hypothesize

that the coordinating zinc residues of Rsh have high affinity for zinc, supporting continued

interaction with the ribosome even during the declining intracellular level of free zinc. This would

allow activation of Rsh under low-zinc conditions before the induction of Mpy-dependent

ribosome hibernation.

Zinc is an essential nutrient that binds to 6-10% of all proteins in bacteria (56). Notably,

several amino acyl tRNA synthetases are zinc binding proteins (56). A lack of zinc in the active

site of threonyl-tRNA synthetase inactivates the aminoacylation activity of the enzyme (57).

Moreover, after alanine (26%) threonine (17%) is the second-most frequent amino acid to follow

the initiator Met in the top 183 most translated genes in *M. smegmatis*. Limitation in zinc

therefore will directly increase the level of deacylated-Thr-tRNA, activating the stringent

response through the surveillance model. Thus, the intracellular availability of zinc in

mycobacteria could potentially be a shared signal for the sequential onset of the stringent

response and ribosome hibernation, mediated directly through sensing the translation activity of

the ribosome. Beyond zinc starvation as the documented condition leading to mycobacterial

ribosome hibernation, it is likely that other conditions inducing ribosome hibernation would also

attenuate the stringent response in a similar manner.

The Clp-dependent loss of ribosome-free intracellular Rsh, while unveiling a novel

mechanism of regulation of Rsh activity *in vivo*, does not entirely rule out the CTD-dependent

autoinhibition of the synthase activity through oligomerization: a model primarily proposed from

in vitro characterization of the enzyme (24-26). However, the evidence of ribosome-bound Rsh

as the dominant species *in vivo* suggests that Rsh oligomers, if any, would be a relatively minor

molecular species in the cell. The predominance of ribosome-bound Rsh *in vivo* further

suggests a reversible transition between the synthase and the hydrolase activities may occur

through differential allosteric effects of ppGpp and GDP/ATP binding to the hydrolase and the

synthase domain, respectively, as shown by Tamman et al. (28).

In summary, we propose a unique structure-based mechanism of mycobacterial Rsh

recognition of translating ribosomes stalled with deacylated A-site tRNA, in which the enzyme

predominantly surveys the amino acylation status of the tRNA during the first cycle of elongation

to activate the stringent response. This mechanism of Rsh activation is programmed to slow

down upon induction of ribosome hibernation. These findings will serve as a new molecular

frame of reference for further investigation of the relationship between the stringent response

and ribosome hibernation in mycobacteria.

Materials and Methods:

Bacterial growth medium

Mycobacterium smegmatis (mc^{2}155) and its variant strains were grown shaking at 200 rpm at 37 °C in Middlebrook 7H9 with ADC enrichment (5% albumin, 2% dextrose, 0.85% sodium chloride and 0.003% catalase) and 0.05% (v/v) Tween80. For ribosome purification, cells from saturated cultures were washed with PBS + 0.05% (v/v) Tween80 for three times and then inoculated at 1:100 dilution in Sauton's medium with either 1 mM ZnSO$_4$ for high-zinc condition or 1 µM zinc-chelator (TPEN, N,N,N',N'-Tetrakis(2-pyridylmethyl)ethylenediamine) for low-zinc condition. 7H10ADC agar plate was used for selection of recombinants and colony growth (*M. smegmatis*). *Escherichia coli* (GC5 or BL21) was grown in LB broth or LB agar at 37 °C. Zeocin (25 µg/mL), kanamycin (20 µg/mL), hygromycin (150 µg/mL for *M. smegmatis* and *E. coli*, carbenicillin (50 µg/mL), and apramycin (5 µg/mL) were used for selection as necessary. For recombinant strain harboring acetamide-inducible promoter, cells were cultured in high- or low -zinc Sauton's medium with 0.2% sodium succinate and induced with 0.2% acetamide.

Construction of recombinant plasmids and strains

A list of plasmids, bacterial strains used in this study is provided in Table S1. A list of oligonucleotides used in this study is provided in Table S2. For construction of Δ*rsh* in *M. smegmatis*, the gene was replaced with a zeocin-resistance marker using a PCR-based modified version of the recombineering strategy as previously described (38). For creating C$_{666}$G, D$_{691}$R and C$_{692}$F variants of Rsh, the Rsh ORF with 500 bp of upstream sequence was PCR-amplified and cloned between KpnI and XbaI sites of the integrative vector pMH94 (39), resulting in the recombinant plasmid pYL238. Mutations at the respective sites were introduced using pYL238 as a template by PCR-based mutagenesis using the primer pairs listed in Table S2, resulting in pYL240 (C$_{666}$G), pYL241 (D$_{691}$R) and pYL242 (C$_{692}$F). Mutations were sequence-confirmed and

the plasmids pYL238, pYL240, pYL241 and pYL242 were electroporated into the mc^{2}155:Δrsh

strain of *M. smegmatis*. Cells expressing the mutant variants were cultured in nutrient-rich

medium and analyzed for Rsh levels by immunoblotting. The integrative plasmid with

anhydrotetracycline(ATc)-inducible CRISPRi-dCas9 system targeting the *clpP1* expression was

constructed and donated by Dr. Keith Derbyshire as a part of the Mycobacterium Systems

Resources (MSR)(58). The *M. smegmatis* strain (YL9) harboring the plasmid was cultured in high-

and low-zinc and ClpP1 depletion was induced by ATc as described previously (39).

Recombinant protein expression and purification

Rsh from *M. smegmatis* (Msmeg_2965) and its variants (Rsh$_{C666G}$, Rsh$_{D691R}$ and Rsh$_{C692F}$) with an

N-terminal His tag fusion were subcloned in pET21b vector using NdeI and HindIII. The plasmids

were introduced into BL21 (DE3) pLysS cells for Rsh expression and purification. Briefly, the cells

were grown in LB medium to log phase (OD$_{600}$ of 0.7) at 37 °C, when RSH expression was induced

with 0.5 mM IPTG for 4 hours at 37 °C. Induced cells were harvested (8000 rpm, 20 minutes at 4

°C in Thermo Scientific™ Fiberlite™ F12-6 × 500 Fixed-Angle Rotor) and then resuspended in N-

I buffer (50 mM Tris (pH 8.0), 300mM NaCl, 5% glycerol, 10 mM imidazole, 1mM PMSF). The

cells were further sonicated (Amplitude 30%, 6 cycles of 10 sec on and 60 sec off for) at 4 °C and

the soluble protein in the supernatant was separated from the debris by centrifugation (13000

rpm, 20 minutes, 4 °C in Thermo Scientific™ Fiberlite™ F12-6 × 500 Fixed-Angle Rotor) and

incubated with pre-equilibrated Ni-NTA resin for 1 hour at 4 °C. After incubation, Ni-NTA resin

was washed three times with N-II buffer [50 mM Tris (pH 8.0), 300mM NaCl, 5% (v/v) glycerol, 30

mM imidazole]. Rsh was then eluted with stepwise increments of imidazole (from 60 mM, to

100mM and to 250 mM) in a buffer containing 50 mM Tris (pH 8.0), 300mM NaCl and 5% (v/v)

glycerol. Peak fractions with were pooled and imidazole was removed by dialysis in the buffer

containing 50 mM Tris (pH 8.0), 300mM NaCl and 30% (v/v) glycerol. For recombinant IF2

(MSMEG_2628) with N-terminal His_{6x} tag, the full-length codon optimized gene with N-terminal

6x-His-sequnce and NdeI-XhoI sites was synthesized (IDT DNA Inc.) (Table S2). The synthetic

fragment was cloned into a pET21a vector. Recombinant protein was expressed in BL21-DE3

(pLysS) and purified on Ni-NTA matrix as detailed for His_6-Rsh above.

Ribosome purification

Rsh bound ribosomes were purified with a few modifications from the previously described

protocol (38). *M. smegmatis* cells were grown for the specified time in 500 mL of Sauton's media

containing 0.05% Tween 80 and either 1 mM $ZnSO_4$ or 1 µM TPEN. Cells were harvested (8000

rpm for 20 minutes in a Thermo Scientific™ Fiberlite™ F12-6 × 500 Fixed-Angle Rotor) and

pulverized 6 times at 15 Hz for 3 minutes in a mixer mill (Retsch MM400). The lysates were mixed

with 20 mL of HMA-10 buffer (20 mM HEPES-K pH7.5, 30 mM NH_4Cl, 10 mM $MgCl_2$, 5 mM β-

mercaptoethanol) and centrifuged for 30 minutes at 30,000g in Thermo Scientific™ Fiberlite™

F12-8 × 50 Fixed-Angle Rotor. The supernatants were collected in the Beckman PC

ultracentrifuge tubes and further centrifuged for 2 hours and 15 minutes at 42800 rpm in a

Beckman rotor Type 70Ti. Pellets were soaked in 4 mL of HMA-10 buffer in an ice water bath

overnight, and then homogenized for 30 minutes. The homogenized pellets were treated with 3

units/mL Rnase-free Dnase (Ambion) for two hours at 4 °C. The contents were transferred to

Beckman PA tubes and centrifuged at 22,000 g for 15 minutes in Thermo Scientific™ Fiberlite™

F12-8 × 50 Fixed-Angle Rotor. The supernatants were collected into Beckman ultracentrifuge PC

tubes and centrifuged for 2 hours and 15 minutes at 42,800 rpm in a Beckman rotor Type 70Ti.

The pellets were dissolved in 1 mL HMA-10 buffer and centrifuged for 10 minutes at 10,000 g.

The supernatants containing the crude ribosomes were collected and quantified by measuring

their optical density at 260 nm. The crude ribosome preparations were then layered on top of 37

mL sucrose density gradients (10%-40%) and centrifuged for 16 hours at 24,000 rpm in a

Beckman rotor SW 28. The 70S ribosome fractions were collected after fractionating the sucrose

gradient in a 260 nm Brandel gradient analyzer as previously described (59). The pooled 70S

ribosome fractions pelleted by ultracentrifugation at 42,800 rpm for 3 hours in a Beckman rotor

Type 70Ti were suspended in HMA-10 buffer (20 mM HEPES-K pH7.5, 30 mM NH$_4$Cl, 10 mM

MgCl$_2$, 5 mM β-mercaptoethanol) and quantified by measuring absorbance at 260 nm.

MPY-bound ribosomes and other high-salt washed ribosomes were purified exactly as

described previously (38, 39). The steps up to homogenization were same as described above.

After homogenization, 4 mL of the homogenized suspension was treated with 3 units/mL Rnase-

free Dnase (Ambion) for one hour at 4 °C, then 4 mL of HMA-0.06 buffer (20 mM HEPES-K pH

7.5, 600 mM NH$_4$Cl, 10 mM MgCl$_2$, 5 mM β-mercaptoethanol) was added to the suspension, which

was further incubated at 4 °C for 2 hours. The content was transferred to Beckman PA tubes and

centrifuged at 22,000 g for 15 minutes in Thermo Scientific™ Fiberlite™ F12-8 × 50 Fixed-Angle

Rotor. Supernatants were collected and layered on top of 8 mL 32% sucrose solution in HMA-10

buffer and centrifuged for 16 hours at 37000 rpm in a Beckman rotor Type 70Ti. The supernatant

was discarded, and the pellet was rinsed with HMA-10 buffer to remove the brownish material.

The crude ribosome was resuspended in 1 mL HMA-10 buffer and centrifuged for 10 minutes at

10,000 g. Supernatant containing the crude ribosome was collected and quantified by measuring

optical density at 260nm. 70S ribosome from the crude ribosome was further separated on a 10-

40% sucrose density gradient and fractionated as indicated above.

In vitro reconstitution of Rsh with 70S initiation complex

Co-sedimentation of Rsh with the ribosomal subunits: 70S ribosomes were purified from low-zinc

culture of isogenic Δ*rsh*/Δ*mpy*/Δ*mrf* triple mutant of *M. smegmatis*. Subunits were dissociated from

the purified 70S ribosomes by incubation at 37°C for thirty minutes in HMA-1 buffer (20 mM

HEPES-K pH7.5, 30 mM NH$_4$Cl, 1 mM MgCl$_2$, 5 mM β-mercaptoethanol. The resulting subunits

were then purified from a 10-40% sucrose density gradient. Individual subunits equivalent to A_{260}

of 0.3 (7.2 pmol) were mixed with 1.8 pmol of purified recombinant Rsh in 100 μL binding buffer

(20 mM HEPES-K pH7.5, 100 mM NH4Cl, 10 mM $MgCl_2$, 5 mM β-mercaptoethanol, 50mM KCl),

and the mixture was incubated for thirty minutes at 37°C. The reactions were then layered on 100

μL of 32% sucrose cushion and centrifuged at 42,800 rpm for 3 hours in a Beckman TLA 100

rotor using the Optima-Max TL ultracentrifuge (Beckman Coulter). After centrifugation, the

supernatant was discarded, and the ribosome pellet was resuspended in 20 μL HMA-10 buffer.

The abundance of Rsh was further determined by immunoblotting using the protocol described

below Rsh, and S13, and L33.

Preparation of fMet-tRNA$_i^{Met}$: An *E. coli* culture containing a plasmid borne expression vector for

tRNA$_i^{Met}$ under control of the LacI repressor was grown to an optical density at 600 nm of 0.4-

0.6 at which time IPTG was added to induce expression of the tRNA. After 8 h the cells were

harvested, lysed by phenol, and total tRNA isolated using a series of differential precipitation

steps (60). Approximately half of the recovered tRNA was a substrate for charging by MetRS.

The charging reaction (2 mL) contained approximately 200 nmoles of total tRNA and 10 nmoles

of MetRS in the presence of 1 mM methionine, 2 mM ATP, 10 mM $MgCl_2$, 20 mM KCl, 4 mM

DTT, 50 μg/mL BSA, and 50 mM Tris-HCl pH 7.5. After 10 min at 37 °C a 4.0 μL aliquot of the

reaction was removed for determination of the charging efficiency as described by Gamper et *al*

(61). The remaining reaction was incubated for 10 min at 37 °C with 1.7 μmoles of 10-

formyltetrahydrofolate and 20 nmoles of methionyl formyl transferase to convert Met-tRNA$_i^{Met}$ to

fMet-tRNA$_i^{Met}$. The reaction was quenched by adding a 0.1 volume of 2.5 M NaOAc pH 5.0.

Following an equal volume pH 5 phenol-chloroform-isoamyl alcohol extraction (80:17:3), the

tRNA was ethanol precipitated, dissolved in 300 μL 25 mM NaOAc pH 5.0, and stored at -70 °C.

Approximately 200 nmoles of tRNA was recovered, of which 45% was fMet-tRNA$_i^{Met}$.

Reconstitution of Rsh-bound 70S initiation complex: 70S ribosomes prepared from a low-zinc

culture of isogenic Δ*rsh*/Δ*mpy*/Δ*mrf* triple mutant of *M. smegmatis* were dialyzed at 4°C with low-

magnesium buffer (20 mM HEPES-K pH7.5, 30 mM NH_4Cl, 0.5 mM $MgCl_2$, 5 mM β-

mercaptoethanol) to dissociate the 30S and 50S subunits. Two rounds of dialysis were

performed, with fresh buffer change every four hours. Subunit dissociation was confirmed by

loading 2.4 picomoles of the dialyzed ribosomes onto a 10-40% sucrose density gradient.

Another 2.4 pmoles of dissociated ribosomes were mixed with five-fold molar excess of fMet-

$tRNA_i^{fMet}$, a synthetic mRNA template (GGCAAGGAGGUAAAAAUGUUCAAAAAA-Flour) (IDT,

USA), recombinant His_6-IF2, recombinant His_6-Rsh, and either GTP or GMP-PNP (Sigma-

Aldrich) in a 100 μL reaction in buffer I (20 mM HEPES-K pH 7.5, 30 mM NH_4Cl, 1 mM $MgCl_2$, 5

mM β-mercaptoethanol), and the mixture was incubated at 37 °C for 15 minutes to allow

formation of the initiation complex on the 30S subunit. Then 25 μL of buffer II (20 mM HEPES-K

pH7.5, 30 mM NH4Cl, 50 mM MgCl2, 5 mM β-mercaptoethanol, 600mM KCl) was added to the

mixture, thereby bringing the final concentration of $MgCl_2$ and KCl to 10mM and 120 mM,

respectively. The reaction was then incubated for an additional 30 minutes at 37°C to allow

joining of the 50S subunit. Resulting complexes were either used for structural analysis or

resolved on a 15 mL 10-40% sucrose density gradient centrifuged at 35000 rpm for 135 minutes

using SW41 rotor on an Optima L-90K ultracentrifuge (Beckman Coulter), and the resolved

particles were fractionated as previously described (59). For immunoblotting analysis, content of

the indicated fractions were precipitated using Methanol-chloroform extraction as previously

described (39). Briefly, each fraction was vortexed with 4X volume of methanol for 5-10 seconds

and centrifuged for 10 seconds at 13,200 rpm on a benchtop centrifuge before adding one

volume of chloroform. The mixture was further centrifuged at 13,200 rpm for 60 seconds, after

which 3X volume of water was added and the mixture was vortexed before further centrifugation

at 13,200 rpm for 60 seconds. The upper aqueous layer was discarded while retaining the

interface and the organic layer, which was mixed with 3X volume of methanol, vortexed and

centrifuged at 13,200 rpm for 2 minutes. The supernatant was discarded, and the resulting

pellet was dried and resuspended in 0.1 M Tris containing 8M urea for further analysis of

proteins by immunoblotting.

Immunoblotting

Total cell lysates (10 µg) or purified 70S ribosomes (2.4 pmoles) were used for detecting S13,

MPY, Mrf-FLAG, and S14$_{C-}$, while 24 pmoles ribosome were used for detecting Rsh. Samples

were resolved on 8% SDS-PAGE (for FLAG, Rsh and GroEL analysis) or 12% SDS-PAGE (for

MPY, S14$_{C-}$ and S13), and proteins were probed with anti-FLAG (1:5000, Genescript), anti-GroEL

(1:5000, Enzo), anti-RelA (1:5000), anti-S13 of *E. coli* (1:100, The Developmental Studies

Hybridoma Bank), endogenously raised anti-MPY (1:5000) and anti-S14$_{C-}$ (1:2000) antibodies.

ppGpp extraction and analysis

Cells of specified *M. smegmatis* strains were grown in Sauton's medium with either 1 mM Zn or

1 µM TPEN for 96 hours, after which 1 mL of the culture was labelled with 100   Ci/mL of ^{32}P

KH$_2$PO$_4$ (Perkin Elmer, 900-1100mCi/mmol) for 6 hours on a shaker at 37 $^{\circ}$C. Labeled cells were

washed once with TBST (20mM Tris pH7.6 + 150 mM NaCl + 0.05% Tween 80). Half of the cells

(0.5 mL) were pelleted and resuspended in 25 µL of 4M formic acid and kept frozen at -20 $^{\circ}$C until

ready for extraction. Remaining 0.5 mL cells were starved for 3 hours in TBST, or metal-chelated

TBST, or zinc-supplemented TBST as specified. Following treatment, both pre- and post-starved

cells were pelleted and resuspended in 25   L of 4M formic acid and incubated on ice for 10

minutes. Cell suspensions were frozen on dry ice for 10 minutes and thawed at 37 $^{\circ}$C for 10

minutes. The freeze-thaw cycle was repeated 5 times, after which the cell lysates were

centrifuged at 13,500 rpm for 5 minutes at 4 $^{\circ}$C, and 5   L of the supernatants containing ^{32}P

labeled intracellular nucleotides were resolved on a 20 x 20 cm plastic PEI cellulose F TLC plate

(Millipore) using 1.5 M KH_2PO_4 pH 3.4 as a solvent. When the solvent front reached the top of the

plate, the plate was dried and exposed to X-ray film. 5 L of 100 mM nonradioactive nucleotides

[ppGpp (Trilink), ATP, GDP and GTP (Thermo Fisher)] were also resolved as positional markers,

which were visualized by handheld UV lamp (Stuart, Cole Palmer). For quantitative analysis, pixel

density of ppGpp spot were determined from densitometric scan of radiographs using Fiji

(ImageJ). Using the counts from low-zinc culture of wild-type strain in a radiograph as the common

denominator, relative counts in other samples from the same radiograph were calculated.

Cryo-electron microscopy and image processing   Quantifoil

holey carbon copper grids R 1.2/1.3 were coated with a continuous layer of carbon (~50 Å

thick). After glow discharge for 30 s on a plasma sterilizer, 4 µl of the 200 nM Rsh-bound pre-

elongation complex sample was placed on the grids. The sample was incubated on the grids for

15 s at 4 °C and 100% humidity, followed by blotting for 4 s before flash-freezing into the liquid

ethane using a Vitrobot IV (FEI). Data were collected on a Titan Krios electron microscope at

300 keV using a K3 direct electron-detecting camera (Gatan). −1.50 to −2.50 µm defocus range

was used at a magnification of 81,000×, yielding a pixel size of 0.846 Å. The dose rate of 23.3

electrons/pixel/second with a 2.05 second exposure time, with 41 frames of 0.05 second

duration each, resulted in a total dose of 66.59 e/Å^{-2}. RELION 4.0 was used for data

processing. Image stacks were gain-corrected, dose-weighted, and aligned using MotionCor2

(62) for 9,103 micrographs. The contrast transfer function of each aligned micrograph was

estimated using CTFFIND-4.1 (63). A subset of micrographs was used for particle picking

followed by 2D classification. Relevant 2D classes were used for reference-based particle

picking on all micrographs yielding 1,928,609 particles. These particles were 2D classified and

suitable class averages selected (389,456 particles) (Fig. S7). The selected particles were 3D

refined and followed by 3D classification. Classes corresponding to monosomes with 349,437

particles were selected for further processing. In order to resolve structural heterogeneity in the

monosome particles, we performed 3D refinement using a large subunit mask followed by three

rounds of fixed orientation 3D classification. In the first round of fixed orientation 3D

classification, we used an SSU mask to resolve the SSU conformations into six classes. Class I

with 158,489 particles showed density corresponding to P-site tRNA and Rsh, representing the

Rsh-bound pre-elongation complex. We then performed the second round of fixed orientation

3D classification on this class using a mask encompassing the entire inter-ribosomal subunit

space such as the L7/L12-stalk, A-, P-, and E-site tRNA binding sites, and the L1-stalk regions.

The second round of fixed orientation 3D classification resolved the conformational and

particularly compositional heterogeneity in the masked region. We derived four classes with P-

site tRNA density and two classes with P-site tRNA and Rsh densities. We performed the third

round of fixed orientation 3D classification combining the two classes with P-site tRNA and Rsh

densities, (49,058 particles) using a mask encompassing the P-site tRNA and Rsh. Two of the

classes, one with 16,013 particle and another with 20,308 particles, showed densities

corresponding to P-site tRNA and ZBD and ACT domains of Rsh. However, density

corresponding to ACT domain was slightly better in the class with 16,013 particles and was

refined to a global resolution of 2.98 Å and was used to model the Rsh-ACT domain. Since the

densities corresponding to ZBD were very similar in both classes, they were merged. The

combined class with a total of 36,321 particles showed significant density corresponding to P-

site tRNA and Rsh-ZBD. CTF refinement and Bayesian polishing of the combined class with a

total of 36,321 particles yielded a map with a global resolution of 2.7 Å and was used to model

the Rsh-ZBD.

Model building   Coordinates

of the large and small subunits from our published C- *M. smegmatis* ribosome structure

(PDB:6DZI) were docked as rigid bodies into the cryo-EM map using Chimera 1.14 (64). To

achieve optimal fitting, we adjusted the model based on the densities in Coot (65). The model

was subsequently refined using PHENIX 1.14 (66). A predicted structure of *M. smegmatis* Rsh

is available in the Alphafold Protein Structure database (67). This structure was docked into the

corresponding density in the cryo-EM map using Chimera 1.14 (64) and subsequently refined

with Coot (65) and PHENIX 1.14 (66). A validation report for the model was obtained from

PHENIX 1.14 (66). The overall statistics of EM reconstruction and molecular modeling are listed

in Table S3. ChimeraX-1.0 (68) and Chimera 1.14 (64) were used to generate the structural

figures in the manuscript.

Acknowledgments:

This work was supported by grants to AKO (NIH: AI132422, NIH: AI163599), CS (AI111696),

RKA (NIH:GM61576), and YMH (NIH: GM134931). R.K.A. also acknowledges support to his lab

through NIH R01 grants AI132422, GM139277 and AI155473. Support from the Wadsworth

Center core facilities–Applied Genomic Technologies and the media core facility– is

acknowledged. We also acknowledge Wadsworth Center's and New York Structural Biology

Center's (NYSBC's) 3D-EM facilities. NYSBC EM facilities are supported by grants from the

Simons Foundation (349247), NYSTAR, the NIH (GM103310) and the Agouron Institute

(F00316). Authors acknowledge technical support from Kelley-Hurst, and discussions with Drs.

Pallavi Ghosh, Keith Derbyshire, Todd Gray and Erica Lasek-Nesselquist.

Authors' Contribution:

YL, RT, JC and AKO designed and performed genetics and biochemical experiments; SM,

MRS, SRM, NKB and RKA designed and performed structural analysis and model construction;

JP, CS, HG and YMH provided reagents. All authors wrote the manuscript.

Financial Conflict of Interest:

Authors declare no financial conflict of interest.

Data Deposition:

The cryo-EM map and atomic coordinates of the *M. smegmatis* 70S-Rsh-fMet-tRNAiMet complex

are deposited in the Electron Microscopy and PDB Data Bank (wwPDB.org). Accession codes

are EMD-29397 and 8FR8 respectively.

References

1.   M. Cashel, J. Gallant, Two compounds implicated in the function of the RC gene of Escherichia coli. *Nature* **221**, 838-841 (1969).

2.   D. Chatterji, A. K. Ojha, Revisiting the stringent response, ppGpp and starvation signaling. *Curr Opin Microbiol* **4**, 160-165 (2001).

3.   K. Potrykus, M. Cashel, (p)ppGpp: still magical? *Annu Rev Microbiol* **62**, 35-51 (2008).

4.   J. Prusa, D. X. Zhu, C. L. Stallings, The stringent response and Mycobacterium tuberculosis pathogenesis. *Pathog Dis* **76** (2018).

5.   J. L. Dahl *et al.*, The role of RelMtb-mediated adaptation to stationary phase in long-term persistence of Mycobacterium tuberculosis in mice. *Proc Natl Acad Sci U S A* **100**, 10026-10031 (2003).

6.   C. L. Stallings *et al.*, CarD is an essential regulator of rRNA transcription required for Mycobacterium tuberculosis persistence. *Cell* **138**, 146-159 (2009).

7.   L. A. Weiss, C. L. Stallings, Essential roles for Mycobacterium tuberculosis Rel beyond the production of (p)ppGpp. *J Bacteriol* **195**, 5629-5638 (2013).

8.   J. P. Richards, W. Cai, N. A. Zill, W. Zhang, A. K. Ojha, Adaptation of Mycobacterium tuberculosis to Biofilm Growth Is Genetically Linked to Drug Tolerance. *Antimicrob Agents Chemother* **63** (2019).

9.   C. Vilcheze, W. R. Jacobs, Jr., The Isoniazid Paradigm of Killing, Resistance, and Persistence in Mycobacterium tuberculosis. *J Mol Biol* **431**, 3450-3461 (2019).

10.   N. K. Dutta *et al.*, Inhibiting the stringent response blocks Mycobacterium tuberculosis entry into quiescence and reduces persistence. *Sci Adv* **5**, eaav2104 (2019).

11.   W. Ross, C. E. Vrentas, P. Sanchez-Vazquez, T. Gaal, R. L. Gourse, The magic spot: a ppGpp binding site on E. coli RNA polymerase responsible for regulation of transcription initiation. *Mol Cell* **50**, 420-429 (2013).

12.   Z. D. Dalebroux, M. S. Swanson, ppGpp: magic beyond RNA polymerase. *Nat Rev Microbiol* **10**, 203-212 (2012).

13.   S. Diez, J. Ryu, K. Caban, R. L. Gonzalez, Jr., J. Dworkin, The alarmones (p)ppGpp directly regulate translation initiation during entry into quiescence. *Proc Natl Acad Sci U S A* **117**, 15565-15572 (2020).

14.   S. E. Irving, N. R. Choudhury, R. M. Corrigan, The stringent response and physiological roles of (pp)pGpp in bacteria. *Nat Rev Microbiol* **19**, 256-271 (2021).

15.   G. C. Atkinson, T. Tenson, V. Hauryliuk, The RelA/SpoT homolog (RSH) superfamily: distribution and functional evolution of ppGpp synthetases and hydrolases across the tree of life. *PLoS One* **6**, e23479 (2011).

16.   T. Hogg, U. Mechold, H. Malke, M. Cashel, R. Hilgenfeld, Conformational antagonism between opposing active sites in a bifunctional RelA/SpoT homolog modulates (p)ppGpp metabolism during the stringent response [corrected]. *Cell* **117**, 57-68 (2004).

17.   A. Brown, I. S. Fernandez, Y. Gordiyenko, V. Ramakrishnan, Ribosome-dependent activation of stringent control. *Nature* **534**, 277-280 (2016).

18.   A. B. Loveland *et al.*, Ribosome*RelA structures reveal the mechanism of stringent response activation. *Elife* **5** (2016).

19.   S. Arenz *et al.*, The stringent factor RelA adopts an open conformation on the ribosome
to stimulate ppGpp synthesis. *Nucleic Acids Res* **44**, 6471-6481 (2016).

20.   P. Pausch *et al.*, Structural Basis for Regulation of the Opposing (p)ppGpp Synthetase and
Hydrolase within the Stringent Response Orchestrator Rel. *Cell Rep* **32**, 108157 (2020).

21.   T. M. Wendrich, G. Blaha, D. N. Wilson, M. A. Marahiel, K. H. Nierhaus, Dissection of the
mechanism for the stringent factor RelA. *Mol Cell* **10**, 779-788 (2002).

22.   K. S. Winther, M. Roghanian, K. Gerdes, Activation of the Stringent Response by Loading
of RelA-tRNA Complexes at the Ribosomal A-Site. *Mol Cell* **70**, 95-105 e104 (2018).

23.   H. Takada *et al.*, Ribosome association primes the stringent factor Rel for tRNA-dependent
locking in the A-site and activation of (p)ppGpp synthesis. *Nucleic Acids Res* **49**, 444-457
(2021).

24.   M. Gropp, Y. Strausz, M. Gross, G. Glaser, Regulation of Escherichia coli RelA requires
oligomerization of the C-terminal domain. *J Bacteriol* **183**, 570-579 (2001).

25.   A. Avarbock *et al.*, Functional regulation of the opposing (p)ppGpp synthetase/hydrolase
activities of RelMtb from Mycobacterium tuberculosis. *Biochemistry* **44**, 9913-9923
(2005).

26.   V. Jain, R. Saleem-Batcha, A. China, D. Chatterji, Molecular dissection of the mycobacterial
stringent response protein Rel. *Protein Sci* **15**, 1449-1464 (2006).

27.   H. Takada *et al.*, The C-Terminal RRM/ACT Domain Is Crucial for Fine-Tuning the Activation
of 'Long' RelA-SpoT Homolog Enzymes by Ribosomal Complexes. *Front Microbiol* **11**, 277
(2020).

28.   H. Tamman *et al.*, A nucleotide-switch mechanism mediates opposing catalytic activities
of Rel enzymes. *Nat Chem Biol* **16**, 834-840 (2020).

29.   K. J. Turnbull, I. Dzhygyr, S. Lindemose, V. Hauryliuk, M. Roghanian, Intramolecular
Interactions Dominate the Autoregulation of Escherichia coli Stringent Factor RelA. *Front*
Microbiol **10**, 1966 (2019).

30.   U. Mechold, H. Murphy, L. Brown, M. Cashel, Intramolecular regulation of the opposing
(p)ppGpp catalytic activities of Rel(Seq), the Rel/Spo enzyme from Streptococcus
equisimilis. *J Bacteriol* **184**, 2878-2888 (2002).

31.   S. Ronneau *et al.*, Regulation of (p)ppGpp hydrolysis by a conserved archetypal regulatory
domain. *Nucleic Acids Res* **47**, 843-854 (2019).

32.   F. L. Gratani *et al.*, Regulation of the opposing (p)ppGpp synthetase and hydrolase
activities in a bifunctional RelA/SpoT homologue from Staphylococcus aureus. *PLoS Genet*
14, e1007514 (2018).

33.   B. P. English *et al.*, Single-molecule investigations of the stringent response machinery in
living bacterial cells. *Proc Natl Acad Sci U S A* **108**, E365-373 (2011).

34.   D. Avarbock, J. Salem, L. S. Li, Z. M. Wang, H. Rubin, Cloning and characterization of a
bifunctional RelA/SpoT homologue from Mycobacterium tuberculosis. *Gene* **233**, 261-269
(1999).

35.   T. P. Primm *et al.*, The stringent response of Mycobacterium tuberculosis is required for
long-term survival. *J Bacteriol* **182**, 4889-4898 (2000).

36.   A. K. Ojha, T. K. Mukherjee, D. Chatterji, High intracellular level of guanosine
tetraphosphate in Mycobacterium smegmatis changes the morphology of the bacterium.
Infect Immun **68**, 4084-4091 (2000).

37.   D. Avarbock, A. Avarbock, H. Rubin, Differential regulation of opposing RelMtb activities
by the aminoacylation state of a tRNA.ribosome.mRNA.RelMtb complex. *Biochemistry* **39**,
11640-11648 (2000).

38.   Y. Li *et al.*, Zinc depletion induces ribosome hibernation in mycobacteria. *Proc Natl Acad*
Sci U S A **115**, 8191-8196 (2018).

39.   Y. Li, J. H. Corro, C. D. Palmer, A. K. Ojha, Progression from remodeling to hibernation of
ribosomes in zinc-starved mycobacteria. *Proc Natl Acad Sci U S A* **117**, 19528-19537
(2020).

40.   S. Prisic *et al.*, Zinc regulates a switch between primary and alternative S18 ribosomal
proteins in Mycobacterium tuberculosis. *Mol Microbiol* **97**, 263-280 (2015).

41.   K. Izutsu, A. Wada, C. Wada, Expression of ribosome modulation factor (RMF) in
Escherichia coli requires ppGpp. *Genes Cells* **6**, 665-676 (2001).

42.   R. D. Hood, S. A. Higgins, A. Flamholz, R. J. Nichols, D. F. Savage, The stringent response
regulates adaptation to darkness in the cyanobacterium Synechococcus elongatus. *Proc*
Natl Acad Sci U S A **113**, E4867-4876 (2016).

43.   H. Schafer *et al.*, The alarmones (p)ppGpp are part of the heat shock response of Bacillus
subtilis. *PLoS Genet* **16**, e1008275 (2020).

44.   X. Gao, J. Yeom, E. A. Groisman, The expanded specificity and physiological role of a
widespread N-degron recognin. *Proc Natl Acad Sci U S A* **116**, 18629-18637 (2019).

45.   P. Kudrin *et al.*, The ribosomal A-site finger is crucial for binding and activation of the
stringent factor RelA. *Nucleic Acids Res* **46**, 1973-1983 (2018).

46.   T. Sprink *et al.*, Structures of ribosome-bound initiation factor 2 reveal the mechanism of
subunit association. *Sci Adv* **2**, e1501502 (2016).

47.   R. A. Marshall, C. E. Aitken, J. D. Puglisi, GTP hydrolysis by IF2 guides progression of the
ribosome into elongation. *Mol Cell* **35**, 37-47 (2009).

48.   P. P. Datta, M. R. Sharma, L. Qi, J. Frank, R. K. Agrawal, Interaction of the G ' domain of
elongation factor G and the C-terminal domain of ribosomal protein L7/L12 during
translocation as revealed by cryo-EM. *Molecular Cell* **20**, 723-731 (2005).

49.   T. Yokoyama *et al.*, Structural insights into initial and intermediate steps of the ribosome-
recycling process. *Embo J* **31**, 1836-1846 (2012).

50.   J. D. Friesen, N. P. Fiil, J. M. Parker, W. A. Haseltine, A new relaxed mutant of Escherichia
coli with an altered 50S ribosomal subunit. *Proc Natl Acad Sci U S A* **71**, 3465-3469 (1974).

51.   J. Parker, R. J. Watson, J. D. Friesen, A relaxed mutant with an altered ribosomal protein
L11. *Mol Gen Genet* **144**, 111-114 (1976).

52.   X. Yang, E. E. Ishiguro, Involvement of the N terminus of ribosomal protein L11 in
regulation of the RelA protein of Escherichia coli. *J Bacteriol* **183**, 6532-6537 (2001).

53.   I. S. Gabashvili *et al.*, Solution structure of the E. coli 70S ribosome at 11.5 A resolution.
Cell **100**, 537-549 (2000).

54.   H. Gao *et al.*, Study of the structural dynamics of the E coli 70S ribosome using real-space
refinement. *Cell* **113**, 789-801 (2003).

55.   D. Pletzer *et al.*, The Stringent Stress Response Controls Proteases and Global Regulators
under Optimal Growth Conditions in Pseudomonas aeruginosa. *mSystems* **5** (2020).

56.   C. Andreini, L. Banci, I. Bertini, A. Rosato, Zinc through the three domains of life. *J*
Proteome Res **5**, 3173-3178 (2006).

57.   R. Sankaranarayanan *et al.*, The structure of threonyl-tRNA synthetase-tRNA(Thr)
complex enlightens its repressor activity and reveals an essential zinc ion in the active
site. *Cell* **97**, 371-381 (1999).

58.   J. A. Judd *et al.*, A Mycobacterial Systems Resource for the Research Community. *mBio* **12**
(2021).

59.   Y. Li *et al.*, Purification of Hibernating and Active C- Ribosomes from Zinc-Starved
Mycobacteria. *Methods Mol Biol* **2314**, 151-166 (2021).

60.   v. Ehrenstein, Isolation of sRNA from intact Escherichia coli cells. *Methods in Enzymology*
12A, 588-596 (1967).

61.   H. Gamper, Y. M. Hou, A Label-Free Assay for Aminoacylation of tRNA. *Genes (Basel)* **11**
(2020).

62.   S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

63.   A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216-221 (2015).

64.   E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612 (2004).

65.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
Crystallogr D Biol Crystallogr **66**, 486-501 (2010).

66.   P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).

67.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021).

68.   E. F. Pettersen *et al.*, UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci* **30**, 70-82 (2021).

827
828

829   **Figure legends:**
830
Figure 1: Attenuated stringent response and loss of Rsh in zinc-starved mycobacteria. A.

A conserved domain architecture in mycobacterial RelA/SpoT homologue (Rsh) compared to

RelA nd Rel in other species. The N-terminal domain (NTD) containing the hydrolase (HYD) and

synthase (SYN) is distinguished from the C-terminal domain (CTD) by TGS (Thr-RS, GTPase and

SpoT) and alpha-helix domain (AHD). The CTD is comprised of ZBD (zinc binding domain) and

ACT (aspartokinase, chorismate mutase and TyrA) subdomains. **B.** Radio-TLC showing ppGpp

levels in cultures of wild-type (WT) *M. smegmatis* isolated from high-zinc (1mM $ZnSO_4$) and low-

zinc (1 µM TPEN) Sauton's medium after 96 hours of growth. ^{32}P-labeled sodium phosphate (100

µCi/mL) was added to the cultures six hours prior to starvation in TBST for three hours. After

starvation nucleotides were extracted in formic acid and amounts equivalent to 50000 cpm were

analyzed by TLC. Purified cold ppGpp was loaded as positional marker. The plot shows ppGpp

levels in high-zinc WT samples relative to their low-zinc counterparts from four biologically

independent experiments. **C.** Radio-TLC showing ppGpp levels in WT cells cultured for 96-hours

in high-zinc Sauton's medium, then labelled for six hours, and resuspended in either TBST, or

Chelex 100-treated TBST ($TBST_{ch}$), or zinc-supplemented $TBST_{ch}$ ($TBST_{Zn}$). A Δ*rsh* mutant was

used as control in panels B and C. **D**. Immunoblot analysis of Rsh levels in 70S ribosomes purified

from 96-hour-old high-and low-zinc cultures of *M. smegmatis* strains described in panel B. S14c-

was used as a marker for the remodeled C- ribosome, while S13 was used as the loading control

for ribosomes. In the set of immunoblots on the right, cell lysates from the samples used for

ribosome analysis were probed. The asterisk indicates a non-specific unknown protein cross-

reactive to Rsh antibody. The plots to the extreme right show the normalized average difference

in Rsh between high- and low-zinc cultures from three independent experiments.

Figure 2: Mrf accumulation causes loss of Rsh. A-B. Analysis of Rsh in 70S ribosomes (A)

and cell lysates (B) from high- or low-zinc cultures of Δ*mrf* and complemented (Δ*mrf*comp) strains

of *M. smegmatis.* High and low-zinc cultures of WT were controls. S14$_{c-}$, S13 and GroEL were

probed as loading controls for ribosomes or cell lysates as indicated. The asterisk in panel B

indicates a non-specific protein cross-reactive to Rsh antibody. **C.** Radio-TLC showing ppGpp

levels in 96-hour old high- or low-zinc cultures of WT and low-zinc cultures of Δ*mrf and* Δ*mrfcomp*

of *M. smegmatis* strains after 3-hours of starvation in TBST. Purified cold ppGpp was loaded as

positional marker. The plot shows relative ppGpp levels in each sample compared to the level in

low-zinc culture of WT from two biologically independent experiments. **D.** Ectopic Mrf expression

causes loss of Rsh. Levels of Rsh in the cell lysates of a *M. smegmatis* strain (pYL222) expressing

FLAG-tagged Mrf* under the acetamide-inducible promoter. Cells were cultured in high-zinc

Sauton's medium (1mM ZnSO4) containing 0.2% succinate as the carbon source until OD 0.6,

after which Mrf* expression was induced with 0.2% acetamide, and cells were grown for 96 hours

from inoculation. Uninduced cells were used as the reference. As controls, 96-hour old cells from

the wildtype (WT) and Δ*rsh* strains cultured in either high- or low-zinc (1µM TPEN) Sauton's

medium with succinate as the carbon source. The plot shows average normalized density of Rsh

from two independent experiments. **E.** Analysis of Rsh in ribosomes purified from 96-hour-old

induced or uninduced high-zinc cultures of pYL222 strain. **F.** Simultaneous accumulation of Rsh

and Mrf in YL9 cells (Li et al. 2020) upon conditional depletion of ClpP1 protease. Cells were

grown in high-zinc Sauton's medium until OD of 0.6, after which anhydrotetracycline (ATc) was

added to induce CRISPRi system to deplete ClpP1. Cell lysates were analyzed after 16 hours of

ClpP1 depletion. A parallel uninduced cells were used as control. YL9 cells grown for 96 hours in

high- or low-zinc Sauton's medium without ATc were used as controll. The plot shows average

normalized density of Rsh from two independent experiments.

Figure 3: **Ribosome binding residues in Rsh are critical for its intracellular stability**. **A.** (Top)

Sequence alignment of the C- terminal region of Rsh from *E. coli*, *M. smegmatis* and *M.*

tuberculosis depicting the CCHC like zinc-finger (highlighted yellow) in the ZBD. (Bottom)

Superimposition of the structure (PDBID: 5IQR) of the ZBD region from *E. coli* (yellow) and *M.*

smegmatis (red). Zinc atom coordinated (shown with dashed lines) by the CCHC motif (shown as

ball-and-stick model and numbered based on *E. coli* sequence) has been zoomed in the image

to the right. The corresponding residues from *M. smegmatis* (shown in red ball-and-stick model)

predict similarity in spatial orientation with respect to their *E. coli* counterparts. **B.** Immunoblot

analysis of Rsh variants ($C_{666}G$, $D_{691}R$ and $C_{692}F$) in cell lysates and 70S ribosomes purified from

96-hour-old high-zinc cultures of a Δ*rsh* strain *of M. smegmatis* expressing each Rsh variant

individually from an integrative plasmid (pYL238, pYL240, pYL241, pYL242 for Rsh, Rsh_{C666G},

Rsh_{D691R} and Rsh_{C692F}, respectively). Cultures of Δ*rsh* strain expressing WT Rsh were controls.

Normalized average levels of Rsh and its variants in cell lysates from three independent

experiments are shown in the plot. **C.** *In vitro* ribosome binding of recombinant Rsh and its

variants. 2.4 picomoles of 70S ribosomes purified from a 96-hour old high-zinc culture of Δ*rsh*

strain of *M. smegmatis* were mixed with 1.2 picomoles of Rsh or variant proteins (purified from

recombinant *E. coli* harboring pYL243, pYL246, pYL247, pY248 for Rsh, Rsh_{C666F}, Rsh_{D691R} and

Rsh_{C692F}, respectively) and the mixture was incubated for 30 minutes at 37 ∘C. The complex was

separated by ultracentrifugation through a 32% sucrose cushion. S13 was probed as the loading

control.   The plot shows quantitative analysis of Rsh from two biologically independent

experiments.

Figure 4: **Rsh interaction with a 70S translation initiation complex. A.** Co-sedimentation of

Rsh with the 50S ribosomal subunit. Pellets resulting from ultracentrifugation on a 32% sucrose

cushion were analyzed for S13 to confirm SSU, $L33_C$ to confirm LSU, and Rsh to test preferential

subunit binding. Rsh in the absence of ribosomes was included as control to rule out

sedimentation of free protein. Purified Rsh was loaded as marker. **B-C.** Reconstitution of Rsh-

bound pre-elongation complex in the presence of GTP (B) or GMP-PNP (C). Dissociated

ribosomes (Figure S6) were mixed with indicated ligands in the presence of GTP or GMP-PNP.

After incubation for 15 minutes in low-magnesium (1mM MgCl$_2$) buffer, recombinant Rsh was

added, and magnesium was adjusted to 10 mM. The resulting mixture was incubated for

additional 30 minutes. The complex was resolved in 10-40% SDG, and fractions corresponding

to the 70S-peak and its margin were analyzed for Rsh, IF2 and indicated ribosomal proteins by

immunoblotting using anti-Rsh, anti-His$_6$ (for IF2), anti-S13 and anti-L31 antibodies. All reagents

in the reactions in panels B and C were added in 5-fold molar excess relative to the ribosomal

subunits. **D.** 2.7 Å resolution cryo-EM structure of the 70S-Rsh complex in late initiation state of

translation, with a fMet-tRNA$_i^{Met}$ present in the P-site (green), mRNA (brown), and two C-terminal

domains of Rsh, ZBD (magenta) and ACT (blue). The 30S and 50S subunits are shown in yellow

and blue, respectively. **E**. Ribosomal components interacting with Rsh ZBD and ACT domain,

along with the fMet-tRNA$_f^{Met}$ in the P-site (green), are shown. ZBD interacts with components of

inter-subunit bridge B1a, uS19 (light green), and the A-site finger (ASF, sky blue). ACT interacts

with the 50S components uL16 (sky blue), L11, and 23S rRNA helices H43/44 (sky blue). F.

Superimposition of *M. smegmatis* 70S-Rsh, *E. coli* 70S-RelA (PDB ID: 5KPV) and *E. coli* 70S-IF2

(PDB ID: 6HTQ) structures, comparing their binding sites and conformations on the ribosome.

Position of IF2 (VI-C1, cyan) suggests a direct steric clash (highlighted by red dotted circle) with

the *E. coli* RelA's α-helical domain (RelA-AHD, light coral). *M. smegmatis* Rsh-ACT (Ms- ACT,

blue) is rotated by ~14° (indicated by a red arrow) relative to the *E. coli* RelA-ACT (Ec-ACT, light

blue), while the ZBD domains (Ms-ZBD, magenta and Ec-ZBD in light purple) have the same

position.

Figure 5: The Rsh-ACT domain interacts with the L7/L12 stalk base. A. Superimposition of

the Rsh/Rel A interacting components from the 70S-Rsh/RelA complex structures from *M.*

smegmatis (Ms) 70S-Rsh (ZBD, magenta; ACT, blue), *B. subtilis* (Bs) 70S-Rsh (ZBD, light pink;

ACT, gray), and *E. coli* (Ec) 70S-RelA (ZBD, light purple; ACT, steel blue). Only components

interacting with RelA/Rsh, L7/L12 stalk base (H43/H44 in blue), A-site finger (ASF in blue) uL16

and uL11 (light pink), the A/R tRNA (light coral) and uS19 (sea green), along with the P-site tRNA

(light gray), are displayed. **B.** Superimposition of the C-terminal ZBD and ACT domain from Ms,

Bs, and Ec (color-coded as in panel A) highlighting the rotation in Ms-ACT domain with respect

to Ms-ZBD. **C.** Interactions of Ms Rsh-ACT (blue) with L11 (light pink) and the 23S rRNA helices

H43/44 (light blue) are shown from two viewing directions.

Figure 6: Structural basis of Rsh exclusion from a Mpy-bound 70S ribosome. A.

Superimposition of Ms 70S-Mpy (gray, PDB ID:6DZI) and Ms 70S-Rsh (gold) structures. Rsh

(ZBD, magenta; ACT, blue) and Mpy (green) are shown in the space-filled rendering. **B**. The 16S

rRNAs corresponding to head domain of the 30S subunit in panel A has been enlarged to highlight

their relative position in the two functional states. **C**. The position of uS19 (gray) in the Mpy-bound

state would disrupt the observed interactions between Rsh-ZBD (magenta) and uS19 (orange).

Figure 7: **Model of a dynamic interaction of Rsh with the ribosome in mycobacteria,**

influenced by three distinct conditions. In a nutrient-rich condition supporting high translation

activity, labile Rsh predominantly remains associated with the free 50S subunit, which confers

stability to the protein. Rsh-50S complex joins with a 30S pre-initiation complex (pre-IC). Following

the release of initiation factors, Rsh likely remains in the pre-elongation intermediate (PEI), which

in an unrotated conformation is ready to elongate upon entry of the first A- site tRNA (step A).

Rsh is likely sterically displaced during inter-subunit rotation associated with the first cycle of

elongation (step B). Free Rsh associates with 50S subunit (step C) and the cycle continues until

a deacylated tRNA enters the A-site at the second codon (type I pause), which leads to activation

of Rsh (step D). Rsh could also potentially be activated if free Rsh interacts with a stalled ribosome

downstream of the second codon (Type II pause) (step E), although its frequency could be limited

by relatively low abundance of such stalled complexes compared to the free 50S subunit.

Conditions inducing hibernating ribosomes, which are likely less accessible to Rsh, would

sequester ribosomes from the translation cycle and displace Rsh (step F), thereby rendering Rsh

labile in a Clp protease dependent manner (step G).

Figure 1: Attenuated stringent response and loss of Rsh in zinc-starved mycobacteria. A. A conserved domain architecture in mycobacterial RelA/SpoT homologue (Rsh) compared to RelA nd Rel in other species. The N-terminal domain (NTD) containing the hydrolase (HYD) and synthase (SYN) is distinguished from the C-terminal domain (CTD) by TGS (Thr-RS, GTPase and SpoT) and alpha-helix domain (AHD). The CTD is comprised of ZBD (zinc binding domain) and ACT (aspartokinase, chorismate mutase and TyrA) subdomains. **B.** Radio-TLC showing ppGpp levels in cultures of wild-type (WT) *M. smegmatis* isolated from high-zinc (1mM $ZnSO_4$) and low-zinc (1 μM TPEN) Sauton's medium after 96 hours of growth. ^{32}P-labeled sodium phosphate (100 μCi/mL) was added to the cultures six hours prior to starvation in TBST for three hours. After starvation nucleotides were extracted in formic acid and amounts equivalent to 50000 cpm were analyzed by TLC. Purified cold ppGpp was loaded as positional marker. The plot shows ppGpp levels in high-zinc WT samples relative to their low-zinc counterparts from four biologically independent experiments. **C.** Radio-TLC showing ppGpp levels in WT cells cultured for 96-hours in high-zinc Sauton's medium, then labelled for six hours, and resuspended in either TBST, or Chelex 100-treated TBST (TBST$_{ch}$), or zinc-supplemented TBST$_{ch}$ (TBST$_{Zn}$). A Δ*rsh* mutant was used as control in panels B and C. **D**. Immunoblot analysis of Rsh levels in 70S ribosomes purified from 96-hour-old high-and low-zinc cultures of *M. smegmatis* strains described in panel B. S14c- was used as a marker for the remodeled C- ribosome, while S13 was used as the loading control for ribosomes. In the set of immunoblots on the right, cell lysates from the samples used for ribosome analysis were probed. The asterisk indicates a non-specific unknown protein cross-reactive to Rsh antibody. The plots to the extreme right show the normalized average difference in Rsh between high- and low-zinc cultures from three independent experiments.

Figure 2: Mrf accumulation causes loss of Rsh. A-B. Analysis of Rsh in 70S ribosomes (A) and cell lysates (B) from high- or low-zinc cultures of Δ*mrf* and complemented (Δ*mrf*comp) strains of *M. smegmatis.* High and low-zinc cultures of WT were controls. S14$_{c-}$, S13 and GroEL were probed as loading controls for ribosomes or cell lysates as indicated. The asterisk in panel B indicates a non-specific protein cross-reactive to Rsh antibody. **C.** Radio-TLC showing ppGpp levels in 96-hour old high- or low-zinc cultures of WT and low-zinc cultures of Δ*mrf and* Δ*mrfcomp* of *M. smegmatis* strains after 3-hours of starvation in TBST. Purified cold ppGpp was loaded as positional marker. The plot shows relative ppGpp levels in each sample compared to the level in low-zinc culture of WT from two biologically independent experiments. **D.** Ectopic Mrf expression causes loss of Rsh. Levels of Rsh in the cell lysates of a *M. smegmatis* strain (pYL222) expressing FLAG-tagged Mrf* under the acetamide-inducible promoter. Cells were cultured in high-zinc Sauton's medium (1mM ZnSO4) containing 0.2% succinate as the carbon source until OD 0.6, after which Mrf* expression was induced with 0.2% acetamide, and cells were grown for 96 hours from inoculation. Uninduced cells were used as the reference. As controls, 96-hour old cells from the wildtype (WT) and Δ*rsh* strains cultured in either high- or low-zinc (1μM TPEN) Sauton's medium with succinate as the carbon source. The plot shows average normalized density of Rsh from two independent experiments. **E.** Analysis of Rsh in ribosomes purified from 96-hour-old induced or uninduced high-zinc cultures of pYL222 strain. **F.** Simultaneous accumulation of Rsh and Mrf in YL9 cells (Li et al. 2020) upon conditional depletion of ClpP1 protease. Cells were grown in high-zinc Sauton's medium until OD of 0.6, after which anhydrotetracycline (ATc) was added to induce CRISPRi system to deplete ClpP1. Cell lysates were analyzed after 16 hours of ClpP1 depletion. A parallel uninduced cells were used as control. YL9 cells grown for 96 hours in high- or low-zinc Sauton's medium without ATc were used as controll. The plot shows average normalized density of Rsh from two independent experiments.

Figure 3: **Ribosome binding residues in Rsh are critical for its intracellular stability**. **A.** (Top) Sequence alignment of the C- terminal region of Rsh from *E. coli*, *M. smegmatis* and *M. tuberculosis* depicting the CCHC like zinc-finger (highlighted yellow) in the ZBD. (Bottom) Superimposition of the structure (PDBID: 5IQR) of the ZBD region from *E. coli* (yellow) and *M. smegmatis* (red). Zinc atom coordinated (shown with dashed lines) by the CCHC motif (shown as ball-and-stick model and numbered based on *E. coli* sequence) has been zoomed in the image to the right. The corresponding residues from *M. smegmatis* (shown in red ball-and-stick model) predict similarity in spatial orientation with respect to their *E. coli* counterparts. **B.** Immunoblot analysis of Rsh variants ($C_{666}G$, $D_{691}R$ and $C_{692}F$) in cell lysates and 70S ribosomes purified from 96-hour-old high-zinc cultures of a Δrsh strain *of M. smegmatis* expressing each Rsh variant individually from an integrative plasmid (pYL238, pYL240, pYL241, pYL242 for Rsh, Rsh_{C666G}, Rsh_{D691R} and Rsh_{C692F}, respectively). Cultures of Δrsh strain expressing WT Rsh were controls. Normalized average levels of Rsh and its variants in cell lysates from three independent experiments are shown in the plot. **C.** *In vitro* ribosome binding of recombinant Rsh and its variants. 2.4 picomoles of 70S ribosomes purified from a 96-hour old high-zinc culture of Δrsh strain of *M. smegmatis* were mixed with 1.2 picomoles of Rsh or variant proteins (purified from recombinant *E. coli* harboring pYL243, pYL246, pYL247, pY248 for Rsh, Rsh_{C666F}, Rsh_{D691R} and Rsh_{C692F}, respectively) and the mixture was incubated for 30 minutes at 37 °C. The complex was separated by ultracentrifugation through a 32% sucrose cushion. S13 was probed as the loading control. The plot shows quantitative analysis of Rsh from two biologically independent experiments.

Figure 4

Figure 4: **Rsh interaction with a 70S translation initiation complex. A.** Co-sedimentation of Rsh with the 50S ribosomal subunit. Pellets resulting from ultracentrifugation on a 32% sucrose cushion were analyzed for S13 to confirm SSU, L33$_{C-}$ to confirm LSU, and Rsh to test preferential subunit binding. Rsh in the absence of ribosomes was included as control to rule out sedimentation of free protein. Purified Rsh was loaded as marker. **B-C.** Reconstitution of Rsh-bound pre-elongation complex in the presence of GTP (B) or GMP-PNP (C). Dissociated ribosomes (Figure S6) were mixed with indicated ligands in the presence of GTP or GMP-PNP. After incubation for 15 minutes in low-magnesium (1mM MgCl$_2$) buffer, recombinant Rsh was added, and magnesium was adjusted to 10 mM. The resulting mixture was incubated for additional 30 minutes. The complex was resolved in 10-40% SDG, and fractions corresponding to 70S-peak and its margin were analyzed for Rsh, IF2 and indicated ribosomal proteins by immunoblotting using anti-Rsh, anti-His$_6$ (for IF2), anti-S13 and anti-L31 antibodies. All reagents in the reactions in panels B and C were added in 5-fold molar excess relative to the ribosomal subunits. **D.** 2.7 Å resolution cryo-EM structure of the 70S-Rsh complex in late initiation state of translation, with a fMet-tRNA$_i^{Met}$ present in the P-site (green), mRNA (brown), and two C-terminal domains of Rsh, ZBD (magenta) and ACT (blue). The 30S and 50S subunits are shown in yellow and blue, respectively. **E.** Ribosomal components interacting with Rsh ZBD and ACT domain, along with the fMet-tRNA$_f^{Met}$ in the P-site (green), are shown. ZBD interacts with components of inter-subunit bridge B1a, uS19 (light green), and the A-site finger (ASF, sky blue). ACT interacts with the 50S components uL16 (sky blue), L11, and 23S rRNA helices H43/44 (sky blue). F. Superimposition of *M. smegmatis* 70S-Rsh, *E. coli* 70S-RelA (PDB ID: 5KPV) and *E. coli* 70S-IF2 (PDB ID: 6HTQ) structures, comparing their binding sites and conformations on the ribosome. Position of IF2 (VI-C1, cyan) suggests a direct steric clash (highlighted by red dotted circle) with the *E. coli* RelA's α-helical domain (RelA-AHD, light coral). *M. smegmatis* Rsh-ACT (Ms- ACT, blue) is rotated by ~14° (indicated by a red arrow) relative to the *E. coli* RelA-ACT (Ec-ACT, light blue), while the ZBD domains (Ms-ZBD, magenta and Ec-ZBD in light purple) have the same position.

Figure 5: The Rsh-ACT domain interacts with the L7/L12 stalk base. A. Superimposition of the Rsh/Rel A interacting components from the 70S-Rsh/RelA complex structures from *M. smegmatis* (Ms) 70S-Rsh (ZBD, magenta; ACT, blue), *B. subtilis* (Bs) 70S-Rsh (ZBD, light pink; ACT, gray), and *E. coli* (Ec) 70S-RelA (ZBD, light purple; ACT, steel blue). Only components interacting with RelA/Rsh, L7/L12 stalk base (H43/H44 in blue), A-site finger (ASF in blue) uL16 and uL11 (light pink), the A/R tRNA (light coral) and uS19 (sea green), along with the P-site tRNA (light gray), are displayed. **B.** Superimposition of the C-terminal ZBD and ACT domain from Ms, Bs, and Ec (color-coded as in panel A) highlighting the rotation in Ms-ACT domain with respect to Ms-ZBD. **C.** Interactions of Ms Rsh-ACT (blue) with L11 (light pink) and the 23S rRNA helices H43/44 (light blue) are shown from two viewing directions.

Figure 6: Structural basis of Rsh exclusion from a Mpy-bound 70S ribosome. A. Superimposition of Ms 70S-Mpy (gray, PDB ID:6DZI) and Ms 70S-Rsh (gold) structures. Rsh (ZBD, magenta; ACT, blue) and Mpy (green) are shown in the space-filled rendering. **B**. The 16S rRNAs corresponding to head domain of the 30S subunit in panel A has been enlarged to highlight their relative position in the two functional states. **C**. The position of uS19 (gray) in the Mpy-bound state would disrupt the observed interactions between Rsh-ZBD (magenta) and uS19 (orange).

Figure 7: **Model of a dynamic interaction of Rsh with the ribosome in mycobacteria, influenced by three distinct conditions**. In a nutrient-rich condition supporting high translation activity, labile Rsh predominantly remains associated with the free 50S subunit, which confers stability to the protein. Rsh-50S complex joins with a 30S pre-initiation complex (pre-IC). Following the release of initiation factors, Rsh likely remains in the pre-elongation intermediate (PEI), which in an unrotated conformation is ready to elongate upon entry of the first A- site tRNA (step A). Rsh is likely sterically displaced during inter-subunit rotation associated with the first cycle of elongation (step B). Free Rsh associates with 50S subunit (step C) and the cycle continues until a deacylated tRNA enters the A-site at the second codon (type I pause), which leads to activation of Rsh (step D). Rsh could also potentially be activated if free Rsh interacts with a stalled ribosome downstream of the second codon (Type II pause) (step E), although its frequency could be limited by relatively low abundance of such stalled complexes compared to the free 50S subunit. Conditions inducing hibernating ribosomes, which are likely less accessible to Rsh, would sequester ribosomes from the translation cycle and displace Rsh (step F), thereby rendering Rsh labile in a Clp protease dependent manner (step G).

Supplementary materials

Figures S1-S10
Tables S1-S3

Figure S1: A full-gel immunoblot image of cells lysates from WT and Δ*rsh* strains cultured in high- (1mM ZnSO$_4$; abbreviated as Zn) and low-zinc Sauton's medium (1μM TPEN; abbreviated as T) and probed with anti-Rsh antibody. Data shows lack of any Rsh-derived smaller product in the low-zinc culture. GroEL from the lysate was probed as a loading control.

Figure S2: Ribosome remodeling has no effect on Rsh-ribosome interaction.
Immunoblot analysis of Rsh bound to 70S ribosomes from 96-hour old high-zinc cultures of WT *M. smegmatis* and its two recombinant strains, Δ*zur* and Δ*c*-:pYL53, which constitutively expressed the remodeled (C-) ribosomes. Low-zinc culture of WT was used as control.

Figure S3: Levels of Mpy in 70S ribosomes purified from high- and low-zinc cultures of WT and Δ*rsh*. S14$_{c-}$ and S13 were probed as controls as indicated.

Figure S4: Level of Rsh-bound ribosomes (A) and ppGpp (B) in *M. smegmatis* before and after exposure to starvation (TBST) from high- or low-zinc culture conditions after 28 hours of growth.

Figure S5: Purified recombinant 6xHis-tagged Rsh proteins expressed from an IPTG-inducible T7 promoter in the *E. coli* BL21 strain and visualized by SDS-PAGE and the Coomassie Blue stain. Each recombinant protein was purified from the same volume of starting cells using the same method, and equal volume of each sample was loaded on the gel.

Figure S6: Sucrose density gradient profile of dissociated ribosomes used in the reconstitution experiments described in figures 4B and C. The reconstituted ribosome complex in figure 4C was used for structural studies.

Figure S7: Image processing of M. smegmatis 70S-Rsh-fMet-tRNAiMet complex. A. A representative micrograph from the M. smegmatis 70S-Rsh-fMet-tRNAiMet complex dataset. **B.** Representative 2D class averages used for the 3D reconstructions. **C.** Flowchart showing the details of 3D classifications and refinements. **D.** Fourier-shell correlation (FSC) plot for the final map (36,321 particles) used in this study. Local resolution for this map is shown in figure S8. **E.** FSC plot for the map (16,013 particles) used in this study to model the Rsh-ACT domain.

Figure S8: Local resolution of the cryo-EM map of the *M. smegmatis* 70S-Rsh-fMet-tRNAiMet complex. Refined by using (A) 70S monosome, (B) 50S LSU, and (C) 30S SSU masks. In panels B and C, SSU and LSU appear to be partially disordered, as expected. A cutting plane has been applied to show the core of the ribosome in each case.

Rsh-ZBD

Rsh-ACT

Figure S9: Modeling of the ZB- and ACT domains into corresponding cryo-EM densities. Two different density threshold values were used for modeling the two domains, as density for the main anchoring ZBD (pink) was stronger than that for the ACT domain.

Figure S10: Species-specific differences in Rsh/RelA-ZBD interaction with uS19, uS13, and ASF. Top- Multiple sequence alignment of a short stretch in the ZBD domain of *M. smegmatis* (Ms) Rsh, *B. subtilis* (Bs) Rsh, and *E. coli* (Ec) RelA. Amino acid substitutions at four regions (indicated by red arrows) that result in species-specific differences in interactions (i1-i4) of the ZBD domain with ribosomal components. Bottom left- Superimposition of Ms 70S-Rsh (ZBD, magenta) and Bs 70S-Rsh (ZBD, light pink). Interactions of Bs-Rsh-ZBD with the ribosome that are absent in the case of Ms-Rsh-ZBD are displayed: two with uS19 (i1 and i2), one with uS13 (i3), and one with ASF (i4). Bottom right- Superimposition of Ms 70S-Rsh (ZBD, magenta) and Ec 70S-RelA (ZBD, light purple). Ec RelA-ZBD has two extra interactions with the ribosome, as compared to Ms Rsh-ZBD: One with uS19 (i2) and another with ASF (i4).

Table S1: List of plasmids and strains

Plasmids	Remarks	Reference
pJI37	P_{hsp60}-based expression vector for mycobacteria; *kan*^{r}	[1]
pMH94	L5-attp-based integrative vector for mycobacteria; *kan*^{r}	[1]
pYUB854	Cosmid vector, hygr	[1]
pTTPla	Tweety-attp-based integrative vector for mycobacteria; *kan*^{r}	[1]
pLam12	$P_{acatamidase}$-based expression vector for mycobacteria; *kan*^{r}	[1]
pJV53-SacB	Sucrose-sensitive marker SacB cloned in pJV53 @ SpeI site; *kan*^{r}	[1]
pET21b	Vectors carry an N-terminal T7-Tag sequence plus an optional C-terminal His Tag sequence, *amp*^{r}	[1]
pYL3	*kan*^{r} cassette in pMH94@EcoRV & NotI sites is replaced by *hyg*^{r} cassette from pYUB854; *hyg*^{r}	[2]
pYL40	*Msmeg_6069* internal deletion in pYL42 backbone @ SacI & XbaI, *hyg*^{r}	[1]
pYL53	MSMEG_6065-6070 cloned in pYL3 @ SacI+XbaI under a constitutive promoter (P^{const}): mutant variant of $P_{zur-box}$, *hyg*^{r}	[1]
pYL155	C-terminally FLAG-tagged *mrf* with P_{zurbox} cloned in pTTPla @ SacI & XbaI sites; *kan*^{r}	[1]
pYL181	C-terminally FLAG-tagged *mrf* with P_{zurbox} cloned in pYL180 @ SacI & XbaI sites; *apr*^{r}	[1]
pYL206	L5-attp-based integrative vector carrying tet-inducible dCas9 and gRNA complementary to Clpp1; *kan*^{r}	[1]
pYL222	*Msmeg_6069* with all six his/cys mutations (H to A, C to A) and Flag tag at C-terminal and $p^{zur-box}$ cloned in plam12 @ XbaI & HindIII sites; *kan*^{r}	This Study
pYL238	*MSMEG_2965* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan*^{r}	This Study
pYL240	*MSMEG_2965* with C666G mutation + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan*^{r}	This Study
pYL241	*MSMEG_2965 with D691R mutation* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan*^{r}	This Study
pYL242	*MSMEG_2965 with C692F mutation* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan*^{r}	This study
pYL243	*MSMEG_2965* with N-terminal His-tag cloned in pET21b @ NdeI & HindIII; *amp*^{r}	This study
pYL246	*MSMEG_2965* (C666G) with N-terminal His-tag cloned in pET21b @ NdeI & HindIII; *amp*^{r}	This study
pYL247	*MSMEG_2965* (D691R) with N-terminal His-tag cloned in pET21b @ NdeI & HindIII; *amp*^{r}	This study
pYL248	*MSMEG_2965* (C692F) with N-terminal His-tag cloned in pET21b @ NdeI & HindIII; *amp*^{r}	This study
pYL259	*MSMEG_2965* + 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg*^{r}	This study
pYL260	*MSMEG_2965* (C666G)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg*^{r}	This study
pYL261	*MSMEG_2965* (D691R)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg*^{r}	This study
pYL262	*MSMEG_2965* (C692F)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg*^{r}	This study
Strains	**Remarks**	**Reference**
mc²155 (WT)	High-Frequency Transformation strain of *M. smegmatis* as parent wild-type	[2]
Δ*msm*$_{c-}$	Unmarked Δ*Msmeg_6065-6070* in mc²155	[2]
Δ*mrf (YL1)*	Unmarked Δ*c-* operon harboring pYL40; *hyg*^{r}	[1]
Δ*mrf*comp (YL2)	Unmarked Δ*c-* operon harboring pYL40 and pYL155; *hyg*^{r}, *kan*^{r}	[1]
Δ*mpy* (YL3)	Unmarked Δ*Msmeg_1878* in mc²155	[2]
Δ*mpy*/Δ*mrf* (YL4)	Unmarked Δ*Msmeg_6065-6070*/Δ*Msmeg_1878* harboring pYL40; *zeo*^{r}, *hyg*^{r}	[1]
Δ*zur* (YL5)	Unmarked Δ*Msmeg_4487* in mc²155	[2]
Δ*zur*/Δmrf (YL6)	Unmarked Δ*Msmeg_6065-6070* /Δ*zur* harboring pYL40; *hyg*^{r}	[1]

| YL9 | Δ*zur* harboring pYL181 and pYL206; *apr*[r], *kan*[r] | [1] |
| Δ*rsh* | Δ*Msmeg_2965* in mc²155, *zeo*[r] | This Study |

1.   Li Y, Corro JH, Palmer CD, Ojha AK: **Progression from remodeling to hibernation of ribosomes in zinc-starved mycobacteria**. *Proc Natl Acad Sci U S A* 2020, **117**(32):19528-19537.
2.   Li Y, Sharma MR, Koripella RK, Yang Y, Kaushal PS, Lin Q, Wade JT, Gray TA, Derbyshire KM, Agrawal RK *et al*: **Zinc depletion induces ribosome hibernation in mycobacteria**. *Proc Natl Acad Sci U S A* 2018, **115**(32):8191-8196.

Table S2: List of oligonucleotides

Name	Sequence	Used in:
His1-R	GGTGCGGACGACGACAGCGCCGTCGAAGGTGGCGCTGACGAGCAGGGT	pYL222
His1-F	ACCCTGCTCGTCAGCGCCACCTTCGACGGCGCTGTCGTCGTCCGCACC	
His2-R	GGCCGAGACGGCTCCGGCCACCAGCTCAAGTAC	
His2-F	GTACTTGAGCTGGTGGCCGGAGCCGTCTCGGCC	
His3-R	CACGTCGGCGCGCCGGGCCAGACGGCGAAGCAGGAT	
His3-F	ATCCTGCTTCGCCGTCTGGCCCGGCGCGCCGACGTG	
p6069-CXXCR	ATCGTCACGCACGGTGGCCGAGACGGCTCCGTGCACCAGCTC	
p6069-CXXCF	GAGCTGGTGCACGGAGCCGTCTCGGCCACCGTGCGTGACGAT	
Pzurbox 6069-XF	CGCGCCGTTCGTCTAGACGCTGCACCAGTTCTCGCC	
p6069-FLAGHR	GGCCGGAAGCTTTCACTTATCGTCGTCATCCTTGTAATCCGATTGCTCTCCTGT	
pRelAMS-up500KpnIF	CTTCCCCGAACCGGTACCCCAGTTCAAGGACCTCAC	pYL236,240, 241,242,256, 259,260,261, 262
pRelAMS-XbaIFLAGR	CGAACCTCTAGATTACTTATCGTCGTCATCCTTGTAATCGGCCGCGCTGGTGACGCG	
RelAC666G-F	ACCAAGCTGGCCAAGGGCTGCACCCCGGTGCCG	pYL240,246, 260
RelAC666G-R	CGGCACCGGGGTGCAGCCCTTGGCCAGCTTGGT	
RelAD691R-F	AGCGTGCACCGCACCCGTTGCACCAACGCCGAG	pYL241,247, 261
RelAD691R-R	CTCGGCGTTGGTGCAACGGGTGCGGTGCACGCT	
RelAC692F-F	GTGCACCGCACCGACTTCACCAACGCCGAGTCG	pYL242,248, 262
RelAC692F-R	CGACTCGGCGTTGGTGAAGTCGGTGCGGTGCAC	
prelA-NdeIHISF	TGACACATATGCACCACCACCACCACCACGTCGACGAGCCAGGCAAG	pYL243,246, 247,248
prelA-HindIIItgaR	CGCCGAACCGCTAAGCTTTCAGGCCGCGCTGGTGACGCG	

> Sequence of codon optimized synthetic IF2:
ATACGAGAACCATatgCACCACCACCACCACCACCAGGCACCGGTCGTGGGAGGGGTAAGACTTCCTCA
CGGTAACGGCGAGACGATCCGTCTGGCACGCGGCGCTTCATTATCAGATTTTGCTGAAAAGATAAATG
CGAATCCGGCTTCACTTGTGCAGGCATTGTTCAATCTTGGAGAAATGGTTACTGCCACGCAAAGTGTCG
GTGATGAAACACTTGAGCTGTTAGGGTCGGAGATGAATTACAATGTACAGGTCGTATCACCAGAAGACG
AGGACCGTGAATTGCTTGAAAGCTTTGATTTAACCTACGGCGAAGACGCGGGTGACGAGGAGGACTTA
GAGGTGCGTCCACCCGTTGTAACCGTTATGGGCCATGTCGATCATGGTAAGACTAGACTTCTGGACAC
AATCCGGAAGGCAAATGTGAGAGAGGGCGAGGCCGGCGGTATAACGCAGCACATCGGGGCCTATCAG
GTTGAGGTTGATTTGGACGGGACGGTGCGTCCCATAACTTTTATTGATACTCCCGGTCACGAGGCCTTT
ACCGCAATGCGTGCCAGAGGTGCCAAGGCTACAGATATTGCTATTTTAGTGGTAGCCGCGGATGATGG
CGTGATGCCTCAAACGGTTGAAGCGATTAACCACGCACAGGCCGCGGACGTTCCGATAGTGGTAGCA
GTCAACAAAATTGATAAGGAAGGAGCGGACCCCGCTAAAATTCGTGGGCAGTTAACAGAATATGGATTA
ATACCGGAAGAGTATGGCGGAGACACCATGTTTGTAGATATCAGTGCGAAACAGGGAACAAATATTGAA
GCATTATTAGAAGCAGTGATTCTTACCGCAGACGCCTCTTTAGATTTGCGTGCAAACCCCGATATGGAA
GCTCAGGGAGTCGCTATAGAAGCCCATCTTGACAGAGGTCGGGGTCCTGTTGCTACTGTATTGATACA
GCGCGGCACCCTTCGGGTTGGTGATTCCGTTGTTGCGGGTGACGCTTATGGGAGAGTCCGGAGAATG
ATCGATGAACATGGAGAGGACGTCGAAGAAGCATTGCCGTCTCGGCCGGTCCAGGTAGTTGGTTTCAC
CTCGGTACCCGGCGCTGGGGACAATTTTTTAGTTGTAGATGAAGATCGGATCGCGCGTCAAATCGCGG
ATCGCCGGTCAGCGCGTAAACGGAACGCCCTTGCTGCACGTAGTCGCAAACGGATCAGCTTGGAGGA
CTTAGACTCCGCCCTTAAAGAGACGTCGCAATTAAACCTTATATTAAAAGGAGACAACGCGGGGACAGT
TGAAGCACTGGAAGAGGCCTTAATGGGAATCCAGGTGGACGACGAGGTAGAACTGCGCGTCATCGAC
CGCGGTGTCGGAGGCGTCACAGAAACGAATGTAAACTTAGCAAGTGCCTCGGACGCGATTATCATTGG
TTTTAACGTCCGGGCTGAAGGAAAGGCGACTGAATTGGCTAATAGAGAAGGAGTTGAAATACGCTATTA
TTCTGTAATCTACCAGGCGATTGATGAAATTGAAGCAGCACTTAAAGGAATGCTGAAGCCTGTATACGA
AGAGAAGGAGCTTGGGCGTGCAGAAATACGTGCCATCTTCCGCAGTTCTAAGGTGGGCAATATAGCTG
GCTGCCTTGTCACATCAGGAATAATGCGCCGCAACGCTAAGGCCCGTTTATTAAGAGACAATGTAGTAG
TTGCACAGAATTTGACGGTAAGTAGTTTAAGACGCGAAAAGGATGACGTAACGGAAGTTCGTGATGGTT
ACGAGTGCGGACTGACCTTAACCTATAATGACATCAAAGAAGGTGACGTAATTGAGACTTATGAATTAG
TTGAAAAAGCGCGTACTtgaCTCGAGTGGCCGATCC

Table. S3 Data collection, Refinement and Model Validation parameters.

Description	70S-Rsh-fMet-tRNA$_i$^{Met} complex
Data collection	
Microscope	FEI Titan Krios
Voltage (kV)	300
Pixel size (Å)	0.846
Defocus range (μm)	-1.50 to -2.50
Average e$^-$ dose per image (e$^-$/ Å^2)	66.59
Particles (initial)	1,928,609
Particles (final)	36,321
FSC-threshold	0.143
Resolution (Å)	2.7
Map-sharpening B factor (Å^2) overall	-44.3
Refinement	
RMS deviations	
Bonds (Å)	0.005
Angles (^{o})	0.616
MolProbity score	2.09
Clash score	13.51
Rotamer outliers (%)	0.41
Ramachandran plot	
Outliers (%)	0.13
Allowed (%)	6.82
Favored (%)	93.05
RNA	
Correct sugar puckers (%)	99.24
Angle outliers (%)	0.00
Bond outliers (%)	0.00
Good backbone conformations (%)	78.98
Model composition	
RNA bases	4843
Protein residues	6299