Starvation sensing by mycobacterial RelA/SpoT homologue through constitutive surveillance of translation

Yunlong Li[1#], Soneya Majumdar[2#], Ryan Treen[1,3#], Manjuli R. Sharma[2], Jamie Corro[1,3], Howard B. Gamper[4], Swati R. Manjari[2], Jerome Prusa[5], Nilesh K. Banavali[2], Christina L. Stallings[5], Ya-Ming Hou[4], Rajendra K. Agrawal[2,3*], Anil K. Ojha[1,3*]

1. Division of Genetics, Wadsworth Center, New York State Department of Health, Albany, NY 12208, USA

2. Division of Translational Medicine, Wadsworth Center, New York State Department of Health, Albany, NY 12237, USA

3. Department of Biomedical Sciences, School of Public Health, University at Albany, Albany, NY 12208, USA

4. Department of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia, PA 19107, USA

5. Department of Molecular Microbiology, Washington University in St. Louis School of Medicine, St. Louis, MO 63110, USA

[#]Equal contribution

*Correspondence:   anil.ojha@health.ny.gov;
       rajendra.agrawal@health.ny.gov

Summary:

The stringent response, which leads to persistence of nutrient-starved mycobacteria, is

induced by activation of RelA/SpoT homologue (Rsh) upon entry of a deacylated-tRNA

in a translating ribosome. However, the mechanism by which Rsh finds such ribosomes

in vivo remains unclear. Here we show that conditions inducing ribosome hibernation

cause loss of intracellular Rsh in a Clp protease-dependent manner. The loss is also

observed in non-starved cells when Rsh-ribosome interaction is abolished by mutation

in Rsh, indicating that interaction with translating ribosomes is necessary for

physiological stability of Rsh. The cryo-EM structure of the Rsh-bound 70S ribosome in

a translation initiation complex reveals hitherto unknown interactions between the ACT

domain of Rsh and components of the ribosomal L7/L12-stalk base, suggesting that the

aminoacylation status of A-site tRNA is surveyed during the first cycle of elongation.

Altogether, we propose a *surveillance model* of Rsh activation that originates from its

constitutive interaction with translating ribosomes.

Introduction:

The stringent response in bacteria is a programmed adaption to nutrient starvation facilitated by the secondary messengers, guanosine tetra- or penta-phosphates [collectively called (p)ppGpp] (Cashel and Gallant, 1969; Chatterji and Ojha, 2001; Potrykus and Cashel, 2008; Prusa et al., 2018). This adaptation mechanism promotes long-term survival and persistence of mycobacteria under various stresses including antibiotics (Dahl et al., 2003; Dutta et al., 2019; Richards et al., 2019; Stallings et al., 2009; Vilcheze and Jacobs, 2019; Weiss and Stallings, 2013). The primary function of (p)ppGpp is to reprogram the global gene expression in cells either directly by binding to RNA polymerase (Potrykus and Cashel, 2008; Ross et al., 2013) or by binding to other cellular targets (Dalebroux and Swanson, 2012; Diez et al., 2020; Irving et al., 2021).

The level of (p)ppGpp in *E. coli* is primarily regulated by a GDP/GTP pyrophosphokinase encoded by RelA, and GDP/GTP pyrophosphohydrolase encoded by SpoT (Potrykus and Cashel, 2008). Mycobacteria encode a single dual-function enzyme, RelA/SpoT homologue (henceforth called Rsh), with both kinase and hydrolase activities possessed by distinct domains of the enzyme (Avarbock et al., 2005; Avarbock et al., 1999; Jain et al., 2006; Ojha et al., 2000; Primm et al., 2000). In addition, mycobacterial Rsh also has a distinct C-terminal region with possible autoinhibitory function, inferred from *in vitro* biochemical studies (Avarbock *et al.,* 2005; Avarbock et al., 2000; Jain *et al.,* 2006), although occurrence of autoinhibition mechanism *in vivo* is unknown. The optimum kinase activity of mycobacterial Rsh is observed in a Rsh Activating Complex (RAC) comprising mRNA, deacylated-tRNA and

the ribosome (Avarbock *et al.*, 2000). Further insights into the domain architectures of

the enzyme and the mechanism of its activation can be derived from the structures of

free RelA from *Streptococcus dysgalactiae* (Hogg et al., 2004), or its *E. coli* and *B.*

subtilis homologues in complex with the ribosome, mRNA and A- and P-site tRNAs

(Arenz et al., 2016; Brown et al., 2016; Loveland et al., 2016). RelA has five distinct

domains (Fig. 1A): i) N-terminal domain possessing the (p)ppGpp synthase and

hydrolase activities, ii) TGS (ThrRS, GTPase, SpoT) domain that interacts with 3' CCA

tail of deacylated-tRNA, iii) alpha helical domain (AHD), iv) zinc-binding domain (ZBD)

that contacts the A-site finger (ASF) on the 23S rRNA and S19 at the inter-subunit

interface, and v) the ACT domain (Aspartate Kinase, Chorismate and TyrA) occupying a

cavity in the large 50S ribosomal subunit (or LSU) (Arenz *et al.*, 2016; Brown *et al.*,

2016; Loveland *et al.*, 2016). The interaction between the TGS domain and 3' CCA tail

of the deacylated-tRNA stabilizes the tRNA in the A/R state while pinning the domain

against the small 30S ribosomal subunit (or SSU) and extending the flexible catalytic N-

terminal of RelA in an open and active conformation (Arenz *et al.*, 2016; Brown *et al.*,

2016; Loveland *et al.*, 2016).

The mechanism by which RelA finds a stalled ribosome complex in a nutrient

starved cell remains unclear. According to a currently adopted 'hopping model',

diffusible RelA samples translation elongation complexes and binds to a stalled

ribosome with deacylated A-site tRNA in the A/R state, and upon catalyzing (p)ppGpp

synthesis the enzyme 'hops' to another stalled complex (Takada et al., 2021; Wendrich

et al., 2002; Winther et al., 2018). The hopping model further hypothesizes that an

inactive RelA in non-starved cells acquires a ribosome-free, closed conformation, in

which the catalytic activity is autoinhibited by the CTD, either through intermolecular

(Avarbock *et al.*, 2005; Gropp et al., 2001; Jain *et al.*, 2006) or intramolecular

interactions (Gratani et al., 2018; Mechold et al., 2002; Ronneau et al., 2019; Takada et

al., 2020; Tamman et al., 2020; Turnbull et al., 2019). Winther et al. later proposed that

free cytosolic RelA in *E. coli* binds to deacylated-tRNA before binding to the translating

ribosome (Winther *et al.*, 2018), although this model is challenged by: a) the evidence of

RelA bound to the ribosome without any A/R state tRNA (Loveland *et al.*, 2016), and b)

single-molecule tracking of RelA-fluorescent protein suggesting constitutive association

between RelA and ribosome in non-starved *E. coli* cells (English et al., 2011).

Recognizing these discrepancies, Takada et al. offered a refined hopping model in *B.*

subtilis in which activation of the stringent response appears to be initiated by a weak

interaction between Rsh and ribosome before the complex is stabilized by the entry of

deacylated-tRNA at the A-site, although they found RelA-ribosome interaction in *E. coli*

to be too weak to universally validate their model (Takada *et al.*, 2021). These studies

while revealing subtle differences in the properties of RelA/Rsh across bacterial

species, also underscore the limitations of biochemical assays in modeling a highly

dynamic *in vivo* interaction between the stringent factor and the translation machinery.

Besides the stringent response, nutrient starvation in bacteria also induces ribosome

hibernation, which is characterized by binding of a specialized protein near the decoding

center of the 70S ribosome to preserve the ribosome in an inactive and antibiotic

resistant state [reviewed in (Gohara and Yap, 2018; Li et al., 2021b; Prossliner et al.,

2018)]. *Mycobacterium smegmatis* and *Mycobacterium tuberculosis* (Mtb) induce

ribosome hibernation as a specific response to severe zinc starvation, a likely condition

in hosts during chronic TB infection (Lai et al., 2021; Li et al., 2020; Li et al., 2018).

Moderate zinc limitation also induces remodeling of ribosomes, in which ribosomal (r-)

proteins containing the CXXC motif (called C+ r- proteins) are replaced by their paralogs

without the motif (called C- r-proteins) (Li *et al.,* 2018; Prisic et al., 2015). The C-

ribosome is subsequently targeted for hibernation by mycobacterial protein Y (Mpy),

with assistance from a zinc-sensing protein called Mpy recruitment factor (Mrf) (Li *et al.,*

2020; Li *et al.,* 2018). A zinc-bound form of Mrf is sensitive to degradation by Clp

protease, such that Mrf accumulation increases with decreasing concentration of zinc

(Li *et al.,* 2020).

Investigating the interplay between the stringent response and ribosome hibernation

in mycobacteria, we show here that interaction with the ribosome is necessary for the

stability of intracellular Rsh in non-starved cells. The structure of Rsh-bound 70S

ribosome in a translation initiation complex allows us to propose a new mechanism of

starvation sensing by mycobacterial Rsh, in which its activation arises from its

engagement with translating ribosomes.

Results:

Mrf attenuates the stringent response in zinc-starved mycobacteria

We tested how a zinc limiting growth condition that induces ribosome hibernation in *M.*

smegmatis would impact ppGpp synthesis. After 96-hours of growth in high- or low-zinc

Sauton's medium, wildtype *M. smegmatis* cells were exposed for 3 hours to

macronutrient starvation in tris-buffered saline with Tween-80 (TBST) to induce the

stringent response to a measurable level (Stallings *et al.*, 2009). Rsh-dependent

accumulation of ppGpp upon TBST starvation of the high-zinc culture was significantly

greater than the low-zinc culture (Fig. 1B). To dissociate the contribution of

macronutrient depletion from zinc limitation in TBST, we examined the stringent

response in cells from high-zinc cultures after exposure to metal-chelated or zinc

supplemented TBST. Metal chelation did not impact the ppGpp synthesis, indicating

that macronutrient starvation in TBST was the primary inducer of the stringent response

(Fig. 1C).

Because interaction with ribosomes is critical for the stringent response, we

hypothesized that the attenuated stringent response in low-zinc cultures was likely due

to zinc-responsive changes in the ribosome structure and/or function. We therefore

analyzed the Rsh occupancy on the ribosome in high- and low-zinc cultures. Rsh

occupancy on the ribosomes purified from low-zinc cells was significantly lower than on

the ribosomes from high-zinc cells (Fig. 1D). Unexpectedly, the difference in Rsh

abundance on the ribosome tightly correlated with the levels in the total cellular lysate

(Fig. 1D). Moreover, lack of any other detectable antibody-recognizing Rsh isoforms

(Fig S1) suggests overall loss of intracellular Rsh. Together, figures 1B-D show that

physiological conditions developed during zinc starvation decreases the stability of Rsh

and its interaction with the ribosome, thereby diminishing the cellular potential to induce

the stringent response. Seeking further explanation, we investigated the effect of two

zinc-responsive physiological changes in the ribosome, remodeling and hibernation, in

altered Rsh stability and Rsh-ribosome interaction. The Zur-mediated C+ to C-

remodeling of the ribosome induced by zinc starvation did not impact the Rsh-ribosome

interaction: Rsh occupancy on the ribosome from a high-zinc culture of wildtype was

comparable to Δ*zur* and pYL53 strains, which constitutively synthesize C- ribosomes

(Fig. S2). To determine the effect of zinc-responsive ribosome hibernation on Rsh-

ribosome interaction, we analyzed the impact of Mpy recruitment factor (Mrf) on Rsh

stability, activity, and its occupancy on the ribosome. Mrf inhibits translation and growth

of cells, likely by sequestering SSU to promote Mpy recruitment (Li *et al.*, 2020).

Deletion of *mrf* in zinc-starved *M. smegmatis* restored the levels of intracellular and

ribosome bound Rsh to those in high-zinc wildtype cells, and the mutant phenotype was

complemented by plasmid-borne *mrf* (Fig. 2A-B). Increase in Rsh abundance in zinc-

starved Δ*mrf* also corresponded to an increase in ppGpp levels upon exposure to TBST

(Fig. 2C). Together, these data are consistent with figure 1C towards the conclusion that

the limiting intracellular free zinc concentration has little direct impact on the stringent

response, while accumulation of Mrf induced by zinc depletion and likely sequestration

of ribosomes for hibernation inhibits Rsh-ribosome interaction and decreases total Rsh

abundance in the cell. Interestingly, deletion of Rsh did not impact Mpy-ribosome

interaction (Fig. S3), distinguishing mycobacteria from other species in which ppGpp

transcriptionally induces factors necessary for ribosome hibernation (Hood et al., 2016;

Izutsu et al., 2001; Schafer et al., 2020).

Ectopic expression of Mrf can reduce intracellular abundance of Rsh.

To further validate Rsh instability, and thus its activity, by Mrf in *M. smegmatis*, we

asked how an ectopic expression of Mrf would impact Rsh levels in the cell. We

exploited a stable Mrf variant (Mrf*), which has reduced zinc affinity, and is recalcitrant

to regulation by the Clp protease system (Li *et al.*, 2020). Consequently, constitutive

transcription of Mrf* leads to its substantial accumulation in high-zinc mycobacteria,

presumably in a conformation that is capable of binding to the ribosome and inhibiting

growth (Li *et al.*, 2020). The effect of ectopic accumulation of Mrf* on Rsh association

with the ribosome was determined using an acetamide-based heterologous expression

system in high-zinc cultures of *M. smegmatis*. Figure 2D shows that expression of Mrf*

in high-zinc cells led to concomitant loss of Rsh, both in the lysates (Fig. 2D) and on the

ribosome (Fig. 2E).

Because SpoT in *Salmonella enterica* was identified as a potential substrate of

the Clp protease system (Gao et al., 2019), we tested the possibility whether

mycobacterial ClpP1P2 protease could be involved in regulating the abundance of free

Rsh in cells. The *M. smegmatis* strain YL9 (Li *et al.*, 2020) harboring CRISPRi-based

system for conditional depletion of ClpP1, while constitutively transcribing FLAG-tagged

Mrf, was used to test whether Rsh loss during Mrf stabilization is Clp-dependent.

Depletion of ClpP1 in high-zinc cells resulted in concomitant accumulation of both Mrf

and Rsh (Fig. 2F). Thus, loss of Rsh in cells during Mrf stabilization is likely achieved

through a Clp-dependent mechanism, although whether Rsh is a direct substrate of Clp

remains to be determined.

Interaction with the ribosome is necessary for intracellular stability of Rsh

The loss of intracellular Rsh abundance in cells with hibernating ribosomes reveals its

instability, while raising the question as to how it is stabilized . Interestingly, we

observed similar levels of Rsh-bound ribosomes in high-zinc cultures before and after

the TBST exposure (Fig. S4), which suggest an interaction between Rsh and ribosome

even during a basal level of ppGpp synthesis. We therefore tested whether interaction

with the ribosome impacts the intracellular stability of Rsh. *E. coli* RelA interacts with the

ribosome primarily through its CTD containing a conserved CCHC type zinc-finger in the

ZBD, which makes extensive contacts with the inter-subunit region of the 70S ribosome,

and ASF of LSU (Brown *et al.*, 2016; Kudrin et al., 2018). Although the requirement of

zinc in RelA-ribosome interaction remains unclear, the zinc coordinating residues

$C_{612}C_{613}H_{634}C_{638}$ ($C_{666}C_{667}H_{688}C_{692}$ in *M. smegmatis Rsh*) of ZBD are conserved in

mycobacteria and likely acquire the same three-dimensional conformation as in *E. coli*

RelA (Fig. 3A). Furthermore, Rsh_{C633A} variant in *M. tuberculosis* (C692 in *M. smegmatis*)

is catalytically impaired *in vitro* (Avarbock *et al.*, 2005), supporting its important role in

Rsh activity. We therefore mutated the key ZBD residues and studied the impact on the

intracellular abundance of Rsh. Substitution of C_{666} and C_{692} residues resulted in

substantial loss of Rsh in both the cell lysate and on the ribosome in actively growing *M.*

smegmatis cells in zinc-rich condition (Fig. 3B). Presence of wildtype Rsh on the

ribosome was unambiguous (Fig. 3B), supporting an argument that Rsh-ribosome

interaction in non-starved cells is constitutive. Substitution in the adjacent D_{691} neither

impacted the Rsh-ribosome interaction nor the intracellular abundance of the protein

(Fig. 3B). Recombinant expression of these Rsh variants in *E. coli* did not significantly

impact the protein yield (Fig. S5), implying that these mutations do not affect the

inherent stability of the protein *in vivo*, and their properties are likely determined by the

intracellular environment of mycobacteria. To further test if the mutations specifically

disrupted Rsh-ribosome interaction in mycobacteria, we mixed recombinant Rsh_{C666G}

and Rsh_{C692F} variants with 70S ribosomes purified from Δrsh mutant of *M. smegmatis*

and analyzed their occupancy on the ribosome separated from the free protein by

ultracentrifugation on a 32% sucrose cushion. Compared to the wild-type Rsh,

association with the ribosome was significantly compromised for both recombinant

proteins, $rRsh_{C666G}$ and $rRsh_{C692F}$ (Fig. 3C). We thus conclude that Rsh constitutively

interacts with the ribosome using its ZBD in non-starved cells, and this interaction is

necessary for the intracellular stability of the enzyme.

Rsh interacts with the IF2 bound 70S initiation complex

We next focused on understanding the stage of translation when Rsh would most likely

interact with the ribosome in a non-starved cell. We streamlined our approach based on

the published structures (Arenz *et al.*, 2016; Brown *et al.*, 2016; Loveland *et al.*, 2016),

which delineate permissive and non-permissive translation complexes for RelA binding.

While RelA can bind to a late translation initiation intermediate (70S- fMet-tRNA$_i^{Met}$ -

mRNA with vacant A-site) (Loveland *et al.*, 2016), its binding to the ribosome during

translation elongation is expected to be sterically hindered by inter-subunit rotation

(Arenz *et al.*, 2016; Brown *et al.*, 2016; Loveland *et al.*, 2016). We therefore

hypothesized that mycobacterial Rsh would enter the translation initiation complex

through one of the free ribosomal subunits and likely remain on the complex until the

start of the first cycle of elongation. This hypothesis also implies that Rsh must remain

associated with the ribosome during its transition from initiation to pre-elongation

stages, without affecting the canonical conformation of the translation complex.

To investigate this possibility, we first determined the propensity of Rsh to

interact with free ribosomal subunits. When purified recombinant Rsh was mixed

individually with each of the two purified factor-free ribosomal subunits and the

macromolecules were separated on 32% sucrose cushion, Rsh appeared to

preferentially co-sediment with LSU (Fig. 4A). We next asked if Rsh would remain in the

translation complex during its transitions through various intermediate stages of

initiation, especially given the potential of a steric clash between Rsh and translation

initiation factor 2 (IF2). The contact points of IF2 on the 70S initiation complex

significantly overlaps with those of Rsh (Arenz *et al.*, 2016; Brown *et al.*, 2016; Loveland

et al., 2016; Sprink et al., 2016). We therefore tested the ability of Rsh to interact with

the ribosome in the presence of IF2, fMet-tRNA$_i^{Met}$ and mRNA *in vitro.* Dissociated

ribosomes (Fig. S6) were mixed with His$_6$-IF2, His$_6$-Rsh, fMet-tRNA$_i^{Met}$, leadered mRNA,

and either GTP or a non-hydrolysable analogue (GMP-PNP). The assembled 70S

complex purified on 10-40% sucrose density gradient (SDG) was analyzed for the

presence of IF2 and Rsh. The 70S ribosomes assembled in the presence of GTP

contained Rsh, but undetectable levels of IF2 (Fig. 4B), demonstrating that Rsh can

associate with the ribosome in the presence of the canonical ligands without interfering

with GTP-dependent release of IF2 (Marshall et al., 2009). This property of

mycobacterial Rsh was in sharp contrast to *B. subtilis* Rsh, which cannot form a stable

complex with the ribosome without an A-site deacyl-tRNA (Takada *et al.*, 2021).

Unexpectedly, both Rsh and IF2 were detected in 70S particles assembled in the

presence of GMP-PNP (Fig. 4C), suggesting that perhaps one or both the proteins

could be occupying the ribosome in their non-canonical conformations.

Cryo-EM structure reveals a hitherto unknown ribosome-Rsh interactions

To gain further insight into the Rsh-bound *M. smegmatis* 70S ribosome in a translation

initiation complex, we performed single-particle cryo-EM to determine the structure of

the 70S ribosome assembled in the presence of IF2, Rsh, fMet-tRNA$_i^{Met}$, leadered

mRNA, and GMP-PNP. 3D classification of the cryo-EM dataset (Figs. S7, S8, Table

S3) revealed a unique Rsh-bound state in which the ribosome is in a classical non-

rotated state, but none of the classes showed any density for IF2. The Rsh-bound 3D

class refined to 2.7 Å resolution showed densities only for two C-terminal domains of

Rsh, ZBD and ACT, along with a fMet-tRNA$_i^{fMet}$ bound at the ribosomal P site, while

remainder of the Rsh domains were found to be disordered (Figs. 4D, 4E, S7 and S9).

The ZBD interacts with the components of both SSU and LSU: SSU protein uS19 and

the LSU 23S rRNA helix H38 (also referred to as ASF) (Fig. 4E). These interactions

resemble those reported in the structures of *E. coli* RelA and *B. subtilis* Rsh on the

ribosome (Loveland *et al.,* 2016; Pausch et al., 2020). The Rsh-ACT domain occupies

the inter-subunit space and interacts with LSU proteins uL16, uL11, and the 23S rRNA

helices H43-44 at the base of the protein L7/L12 stalk (Fig. 4E).

To understand a possible spatial interaction between IF2 and Rsh in *M.*

smegmatis 70S ribosome, we superimposed either our structure or *E. coli* 70S

ribosome-RelA (Loveland *et al.*, 2016) with 70S-IF2-GMP-PNP initiation complexes

(Sprink *et al.*, 2016). Indeed, RelA's alpha-helical domain (AHD) which is followed by

the ZBD, and the ACT domain in its canonical conformation bound to A/R tRNA

(Loveland *et al.*, 2016) would directly clash with domain VI-C1 of IF2 (Fig. 4F). In tertiary

structure, the AHD lies adjacent to the ACT domain while the ZBD is anchored to the

ribosome via its interactions with uS19 and ASF. Moreover, in our structure the ACT

domain was rotated with respect to ZBD by ~14 degrees, presumably to prevent the

clash between AHD and domain VI-C1 of IF2 (Fig. 4F). Across the reported structures

(Loveland *et al.*, 2016; Pausch *et al.*, 2020) the ACT domain is known to exhibit some

conformational variations as compared to the ZBD domain, although a rotation by 14

degrees is unprecedented. It is possible that the adopted conformation of the ACT

domain in our structure is to accommodate the exiting IF2 molecule immediately upon

completion of the translation initiation phase.

Compared to the *E. coli* RelA and *B. subtilis* Rsh, mycobacterial Rsh shows an

overall similarity in tertiary structures of ZBD and ACT domains, although the rotation of

the ACT domain translates into its shift by ~6 Å towards the LSU's L7/L12 stalk base

(Fig. 5A, B). This altered conformation of ACT is stabilized through extensive

interactions with the uL11 and the 23S rRNA helices H43/44 (Fig. 5C). Specifically,

uL11 lysines, 89 and 96 interact with Y791, D789 and H776 of Rsh-ACT. Further, Rsh-

ACT S779, N783, D789 and Y788 interact with the 23S rRNA helices H43-44

phosphate backbones. These interactions are in sharp contrast to the *E. coli* ribosome-

RelA structures, where uL11 interacts with the A/R tRNA but not directly with RelA

(Loveland *et al.*, 2016). Thus, our structure reveals a unique conformational state of the

ribosome-bound Rsh. Multiple sequence alignment of the amino acid sequences does

not reveal any apparent substitutions, insertions, or deletions that can explain the

altered orientation of the ACT domain in *M. smegmatis* Rsh (Fig. S10). Given that our

structure was obtained in the presence of IF2-GMP-PNP, a key difference from the *E.*

coli ribosome-RelA complex, a possible role of IF2 in inducing the rotation of ACT

cannot be ruled out.

The superimposition also reveals some species-specific differences, which could

manifest as differences in Rsh/RelA-ribosome binding affinities (Fig. S10). The ZBD

domains interact with uS19 and ASF in all three cases, however, compared to *M.*

smegmatis, *B. subtilis* Rsh has additional interactions — two with uS19, one with ASF,

and one with uS13 (Fig. S10). Loss of the additional interactions with uS19 and ASF in

M. smegmatis is likely due to substitution at corresponding positions with non-polar

amino acids (Fig. S10). However, *M. smegmatis* Rsh-ZBD could potentially interact with

uS13 since it has an Arg (R681) corresponding to K617 in *B. subtilis*. The *E. coli* RelA

ZBD also has two additional interactions compared to *M. smegmatis*, one with uS19 and

the other with ASF (Fig. S10). However, *E. coli* RelA ZBD has a Gln (Q627)

corresponding to R681 and K617 in *B. subtilis* and *M. smegmatis*, respectively, which

might not interact with uS13 due to its shorter side chain. Since the density

corresponding to ZBD is relatively strong in our structure, we designate it as the primary

anchoring domain of mycobacterial Rsh.

Interactions stabilizing Rsh in our structure are primarily through LSU's ASF,

uL16, uL11, and rRNA helices H43/44, suggesting that Rsh associates primarily with

the LSU, corroborating the results of our binding experiments that shows association of

Rsh with free LSU. Moreover, uL11 and rRNA helices H43/44 together constitute the

L7/L12-stalk base of LSU and are known to be mobile during elongation (Datta et al.,

2005) and recycling (Yokoyama et al., 2012) phases of translation. We thus surmise

that intensive contacts between L7/L12-stalk base and the Rsh-ACT domain would

transiently lock the ribosome in a stalled state before the start of translation elongation

cycle, as a mechanism to survey the aminoacylation state of the incoming tRNA to the A

site. A direct contact of Rsh with uL11 is distinct from *E. coli* RelA, binding of which is

independent of uL11 while (p)ppGpp synthesis requires uL11 and a deacylated-tRNA at

the A-site (Friesen et al., 1974; Parker et al., 1976; Wendrich *et al.*, 2002; Yang and

Ishiguro, 2001). While the changes in *M. smegmatis* complex here could be induced by

the presence of IF2, we cannot exclude the possibility that these interactions or the ACT

conformation are specific *to M. smegmatis* Rsh.

Can Rsh bind to a hibernating ribosome?

Given that interaction with the ribosome is necessary for the intracellular stability of Rsh,

its loss in cells harboring Mrf, and therefore Mpy-bound hibernating ribosomes, suggest

that these ribosomes are perhaps inaccessible to free Rsh. To further probe this

possibility, superimposed the structures of Mpy-bound and Rsh-bound 70S ribosomes.

Mpy and Rsh bind at two physically distinct sites on the unrotated SSU of the ribosome

(Fig. 6A). Mpy N-terminal domain binds to SSU spanning the decoding center by

interacting extensively with the 16S rRNA components that converge from platform,

body, and head regions at the SSU's neck (Li *et al.*, 2018). Rsh ZBD on the other hand

interacts near the inter-subunit bridge B1a (Gabashvili et al., 2000) forming

components, including uS19 of SSU and LSU's ASF (Gao *et al.*, 2019) (Figs. 4E and

6A). Since the binding sites of the two factors do not overlap, we superimposed the

structures of Rsh- and Mpy-bound *M. smegmatis* 70S ribosomes to understand the

accessibility of Mpy-bound hibernating ribosome to Rsh binding. Despite both the

ribosomes being in their unrotated state, the SSU head domain in the Rsh-bound

structure is tilted more towards the LSU in comparison to the Mpy-bound ribosome

structure (Fig. 6B). The untilted SSU head in the MPY-bound state will prevent

interactions of Rsh-ZBD with uS19 (Fig. 6C). Since ZBD is the primary anchoring

domain of Rsh (Loveland *et al.*, 2016; Pausch *et al.*, 2020), any alteration in bridge B1a

configuration in the Mpy-bound state is likely to directly inhibit Rsh binding, thereby

exposing Rsh to proteolytic degradation. Thus, loss in the overall pool of intracellular

Rsh during ribosome hibernation could be a combined outcome of both a decrease in

the synthesis of new Rsh and the occlusion of pre-existing Rsh from the hibernating

ribosome.

Discussion:

The stringent response, characterized by Rsh-dependent synthesis of (p)ppGpp and

subsequent transcriptional reprogramming by the alarmone, is among the central

adaptation mechanisms in nutrient-starved mycobacteria. Yet, a significant knowledge

gap exists in the mechanism underlying starvation sensing by Rsh. Structural studies

with *E. coli* RelA offer mechanistic details about Rsh activation by the presence of

deacylated-tRNA at the A- site in a stalled translation elongation complex, although it

remains unclear how Rsh would encounter such a complex *in vivo*. Two models have

emerged from recent studies: a) the hopping model in which Rel-deacylated-tRNA

complex is formed prior to its entry at the A- site of the elongation complex (Winther *et*

al., 2018), and b) starvation-induced formation of a weak RelA-ribosome complex prior

to its stabilization by deacylated-RNA (Takada *et al.*, 2021). Both models however

preclude RelA-ribosome interaction in non-starved cells, which is inconsistent with the

evidence of constitutive RelA-ribosome interaction throughout the growth of *E. coli* and

P. aeruginosa cells (English *et al.*, 2011; Pletzer et al., 2020). These discrepancies

suggest more complex scenarios that need further clarification. Here, we propose a new

model of Rsh recognition of a paused translation elongation complex in mycobacteria.

In this model, which we call the surveillance model, Rsh constitutively interacts with

translating ribosomes during the formation of the initiation complex, and its transition to

the pre-elongation stage (Fig. 7). Presence of Rsh in a pre-elongation translation

intermediate would arguably trigger (p)ppGpp synthesis upon entry of a deacylated-

tRNA at the A- site during the first cycle of elongation, while entry of a cognate amino

acylated-tRNA and normal cycle of elongation accompanied by inter-subunit rotation

would likely displace Rsh from the complex (Arenz *et al.*, 2016; Brown *et al.*, 2016;

Loveland *et al.*, 2016). Free Rsh appears to be labile and must associate with

ribosomes entering the translation cycle, or a paused elongation complex with

deacylated-tRNA at the A- site (Fig. 7), although the latter interaction would arguably be

less frequent. A rapid rate of translation in an actively growing cell would likely maintain

a sufficiently large pool of free ribosomes, transitioning between termination and

initiation cycles, to ensure a stable population of Rsh-ribosome complex. The

surveillance model of the stringent response activation is arguably less stochastic than

the hopping model.

Lack of Rsh under conditions such as those inducing ribosome hibernation

further supports the model that optimum protein synthesis and accessibility to

translating ribosomes is critical for the intracellular stability of Rsh. This finding further

differentiates the physiological conditions for ribosome hibernation and the stringent

response. The negative impact of Mpy-dependent ribosome hibernation on the stringent

response also highlights a unique relationship between the two central adaptation

mechanisms in mycobacteria in a zinc-responsive manner. This relationship further

implies that the stringent response is an upstream adaptation response relative to

ribosome hibernation; arguably a logical order given that the stringent response

reprograms the gene expression to fine-tune the metabolism, while ribosome

hibernation directly shuts down global protein synthesis altogether. Moreover,

restoration of (p)ppGpp synthesis in zinc-depleted Δ*mrf* or Δ*mpy* implies that Rsh-

ribosome interaction and Rsh activation are relatively more resilient to intracellular

depletion of free zinc, despite that the zinc coordinating residues are crucial for Rsh

function. Perhaps the affinity for zinc in Rsh is too strong to be affected by global

depletion of free zinc and may even support activation of the stringent response under

moderately low-zinc condition that is not severe enough to induce Mpy-dependent

ribosome hibernation. Zinc is an essential nutrient that binds to 6-10% of all proteomes

in bacteria (Andreini et al., 2006), likely impacting the function of multiple proteins

including RNA polymerase β' subunit (Hu and Liu, 2022), and several amino acyl tRNA

synthetases (aaRS) (Andreini *et al.*, 2006). Lack of zinc in the active site of ThrRS

inactivates the aminoacylation activity of the enzyme (Sankaranarayanan et al., 1999).

Thus, intracellular availability of zinc in mycobacteria could be a shared signal for

sequential onset of the two adaptive responses mediated directly through sensing the

translation activity of the ribosome. The sequential relationship between the stringent

response and ribosome hibernation could possibly occur even in other nutrient-starved

conditions that induce both the adaptation pathways.

The Clp-dependent loss of ribosome-free intracellular Rsh, while reaffirming the

tight regulation of ppGpp synthesis in cells, does not entirely rule out the CTD-

dependent autoinhibition of the synthase activity through oligomerization: a model

primarily proposed from *in vitro* characterization of the enzyme (Avarbock *et al.*, 2005;

Gropp *et al.*, 2001; Jain *et al.*, 2006). However, the evidence of ribosome-bound Rsh as

the dominant species suggests that Rsh oligomers, if any, would be a minor molecular

species *in vivo*. Moreover, the requirement of CTD in oligomerization *in vitro* (Avarbock

et al., 2005) suggest that formation of the oligomeric form is perhaps outcompeted *in*

vivo by a stronger affinity of the CTD for the ribosome than for another Rsh CTD. The

likely predominance of ribosome-bound Rsh *in vivo* further raises the possibility that a

reversible transition between synthase and hydrolase activities may occur without the

CTD-dependent autoinhibition mechanism, perhaps through differential allosteric effects

of ppGpp and GDP/ATP binding to the hydrolase and the synthase domain,

respectively, as shown by Tamman et al. (Tamman *et al.*, 2020). The CTD however

could modulate the enzymatic activity of Rsh through its interaction with the ribosome,

perhaps like the *B. subtilis* Rsh (Takada *et al.*, 2020). In summary, this study highlights

that the *in vitro* biochemical studies may have limitations in understanding the function

and regulation of mycobacterial Rsh, while offering new framework for further

investigation of the interrelationship between the stringent response and ribosome

hibernation in mycobacteria.

Experimental Procedures:

Bacterial growth medium

Mycobacterium smegmatis (mc^{2}155) and its variant strains were grown on a shaker at 200 rpm at 37 °C in Middlebrook 7H9 + 10%ADC + 0.05% Tween80 until required. For experiments related to ribosome purification, cells from saturated culture were washed with PBS + 0.05% Tween80 for three times and then inoculated at 1:100 dilution in Sauton's medium with either 1 mM ZnSO$_4$ or 1 µM zinc-chelator (TPEN, N,N,N',N'-Tetrakis(2-pyridylmethyl)ethylenediamine) as specified. 7H10ADC agar plate was used for selection of recombinants and colony growth (*M. smegmatis*). Pellicle biofilms of *Mycobacterium tuberculosis* (mc^{2}7000) were cultured in detergent-free Sauton's medium either with 1mM ZnSO$_4$ or without supplemental zinc. Cells from a 40 µL aliquot of a saturated planktonic culture in 7H9 medium with 10% OADC and 0.05% Tween-80 were inoculated into 4 mL of detergent-free Sauton's medium in a 12-well polystyrene plate and incubated without shaking for 7-weeks at 37 °C. 100 µg/mL pantothenate was added for cultures of *Mycobacterium tuberculosis* (mc^{2}7000). *Escherichia coli* (GC5 or BL21) was grown in LB broth or LB agar at 37 °C. Zeocin (25 µg/mL), kanamycin (20 µg/mL), hygromycin (150 µg/mL for *M. smegmatis* and *E. coli*, or 50 µg/mL for *M. tuberculosis*), carbenicillin (50 µg/mL), and apramycin (5 µg/mL) were used for selection as necessary.

Construction of recombinant plasmids and strains

List of plasmids, bacterial strains in this study are provided in Table S1. List of oligonucleotides are provided in Table S2. For construction of Δ*rsh* in *M. smegmatis*, the

gene was replaced with a zeocin-resistance marker using a PCR-based modified version

of the recombineering strategy as previously described (Li *et al.,* 2018). For analyzing

the stability of wildtype or $C_{666}G$, $D_{691}R$ and $C_{692}F$ variants of Rsh, wildtype Rsh ORF with

500 bp ustream sequence was PCR-amplified and cloned into KpnI-XbaI sites of an

integrative vector pMH94, resulting in a recombinant plasmids pYL238. Mutations at

respective sites were introduced using pYL238 as template by PCR-based mutagenesis

using primers listed in Table S2, resulting in pYL240 ($C_{666}G$), pYL241 ($D_{691}R$) and pYL242

($C_{692}F$). The plasmids pYL238, pYL240, pYL241 and pYL242 were electroporated into

Δ*rsh* strain of *M. smegmatis*. The recombinant strains were cultured in nutrient-rich

medium and analyzed for Rsh levels by immunoblotting as described.

Recombinant protein expression and purification

Rsh from *M. smegmatis* (Msmeg_2965) and its variants (Rsh_{C666G}, Rsh_{D691R} and

Rsh_{C692F}) with N-terminal His tag fusion were cloned in pET21b vector using NdeI and

HindIII. The recombinant plasmids were introduced into BL21 (DE3) pLysS cells for

recombinant Rsh expression and purification. Briefly, the cells were grown in LB medium

to log phase (OD_{600} around 0.7) at 37 °C when RSH expression was induced with 0.5 mM

IPTG for 4 hours at 37 °C. Induced cells were harvested (8000 rpm, 20 minutes, RT in

Thermo Scientific™ Fiberlite™ F12-6 × 500 Fixed-Angle Rotor) and then resuspended in

N-I buffer (50 mM Tris (pH 8.0), 300mM NaCl, 5% glycerol, 10 mM imidazole, 1mM

PMSF). The cells were further sonicated (Amplitude 30%, 10 sec on/off for 60 minutes)

at 4 °C and the soluble protein in the supernatant was separated from the debris by

centrifugation (13000 rpm 20 minutes, 4 °C in Thermo Scientific™ Fiberlite™ F12-6 × 500

Fixed-Angle Rotor) and incubated with pre-equilibrated Ni-NTA resin for 1 hour at 4 °C.

After incubation, Ni-NTA resin was washed three times with N-II buffer [50 mM Tris (pH

8.0), 300mM NaCl, 5% (v/v) glycerol, 30 mM imidazole]. Rsh was then eluted with

stepwise increments of imidazole (from 60 mM to 100mM and to 250 mM) in a buffer

containing 50 mM Tris (pH 8.0), 300mM NaCl and 5% (v/v) glycerol. Fractions with the

purest proteins were pooled and imidazole was removed by dialysis in the buffer

containing 50 mM Tris (pH 8.0), 300mM NaCl and 30% (v/v) glycerol. For recombinant

IF2 with N-terminal His_{6x} tag, the full-length codon optimized gene with N-terminal 6X-

His-sequnce and NdeI-XhoI sites was synthesized (IDTDNA Inc.) (see the sequence after

Table S2), and clone into a pET21a vector. Recombinant protein was expressed in BL21-

DE3 pLysS) and purified on Ni-NTA matrix as detailed for His_6-Rsh above.

Ribosome purification

Rsh bound ribosomes were purified with few modifications from the previously described

protocol (Li *et al.*, 2018). *M. smegmatis* cells were grown for specified time in 500 mL of

Sauton's media containing 0.05% Tween 80 and either 1 mM $ZnSO_4$ or 1 μM TPEN. Cells

were harvested (8000 rpm for 20 minutes in Thermo Scientific™ Fiberlite™ F12-6 × 500

Fixed-Angle Rotor) and pulverized 6 times at 15 Hz for 3 minutes in a mixer mill (Retsch

MM400). The lysates were mixed with 20 mL of HMA-10 buffer (20 mM HEPES-K pH7.5,

30 mM NH_4Cl, 10 mM $MgCl_2$, 5 mM β-mercaptoethanol) and centrifuged for 30 minutes

at 30,000 g in Thermo Scientific™ Fiberlite™ F12-8 × 50 Fixed-Angle Rotor. The

supernatants were collected in the Beckman PC ultracentrifuge tubes and further

centrifuged for 2 hours and 15 minutes at 42800 rpm in a Beckman rotor Type 70Ti.

Pellets were soaked in 4 mL of HMA-10 buffer in an ice water bath overnight, and then

homogenized for 30 minutes. Then the homogenized pellets were treated with 3 units/mL

Rnase-free Dnase (Ambion) for two hours at 4 °C. The contents were transferred to

Beckman PA tubes centrifuged at 22,000 g for 15 minutes in Thermo Scientific™

Fiberlite™ F12-8 × 50 Fixed-Angle Rotor. Supernatants were collected into Beckman

ultracentrifuge PC tubes and centrifuged for 2 hours and 15 minutes at 42,800 rpm in a

Beckman rotor Type 70Ti. Pellets were dissolved in 1 mL HMA-10 buffer and centrifuged

for 10 minutes at 10,000 g. Supernatants containing the crude ribosomes were collected

and quantified by measuring their optical density at 260 nm. The crude ribosome

preparations were then layered on top of 37 mL sucrose density gradients (10%-40%)

and centrifuged for 16 hours at 24,000 rpm in a Beckman rotor SW 28. The 70S ribosome

fractions were collected after fractionating the sucrose gradient in a 260 nm Brandel

gradient analyzer as previously described (Li et al., 2021a). The pooled 70S ribosome

fractions pelleted by ultracentrifugation at 42,800 rpm for 3 hours in a Beckman rotor Type

70Ti were suspended in HMA-10 buffer (20 mM HEPES-K pH7.5, 30 mM NH_4Cl, 10 mM

$MgCl_2$, 5 mM β-mercaptoethanol) and quantified by measuring absorbance at 260 nm.

MPY-bound ribosomes and other high-salt washed ribosomes were purified

exactly as described previously (Li *et al.*, 2020; Li *et al.*, 2018). The steps up to

homogenization were same as described above. After homogenization, 4 mL of the

homogenized suspension was treated with 3 units/mL Rnase-free Dnase (Ambion) for

one hour at 4 °C, then 4 mL of HMA-0.06 buffer (20 mM HEPES-K pH 7.5, 600 mM NH$_4$Cl,

10 mM MgCl$_2$, 5 mM β-mercaptoethanol) was added to the suspension, which was further

incubated at 4 °C for 2 hours. The content was transferred to Beckman PA tubes and

centrifuged at 22,000 g for 15 minutes in Thermo Scientific™ Fiberlite™ F12-8 × 50 Fixed-

Angle Rotor. Supernatants were collected and layered on top of 8 mL 32% sucrose

solution in HMA-10 buffer and centrifuged for 16 hours at 37000 rpm in a Beckman rotor

Type 70Ti. The supernatant was discarded, and the pellet was rinsed with HMA-10 buffer

to remove the brownish material. The crude ribosome was resuspended in 1 mL HMA-10

buffer and centrifuged for 10 minutes at 10,000 g. Supernatant containing the crude

ribosome was collected and quantified by measuring optical density at 260nm. 70S

ribosome from crude ribosome was further separated on a 10-40% sucrose density

gradient and fractionated as indicated above.

In vitro reconstitution of Rsh with 70S initiation complex

Co-sedimentation of Rsh with the ribosomal subunits: 70S ribosomes were purified from

low-zinc culture of isogenic Δ*rsh*/Δ*mpy*/Δ*mrf* triple mutant of *M. smegmatis*. Subunits were

dissociated from the purified 70S ribosomes by incubation at 37°C for thirty minutes in

HMA-1 buffer (20 mM HEPES-K pH7.5, 30 mM NH$_4$Cl, 1 mM MgCl$_2$, 5 mM β-

mercaptoethanol. The resulting subunits were then purified from a 10-40% sucrose

density gradient. Individual subunits equivalent to A$_{260}$ of 0.3 (7.2 pmol) were mixed with

1.8 pmol of purified recombinant Rsh in 100 μL binding buffer (20 mM HEPES-K pH7.5,

100 mM NH4Cl, 10 mM MgCl$_2$, 5 mM β-mercaptoethanol, 50mM KCl), and the mixture

was incubated for thirty minutes at 37°C. The reactions were then layered on 100 μL of

32% sucrose cushion and centrifuged at 42,800 rpm for 3 hours in a Beckman TLA 100

rotor using the Optima-Max TL ultracentrifuge (Beckman Coulter). After centrifugation,

the supernatant was discarded, and the ribosome pellet was resuspended in 20 μL HMA-

10 buffer. The abundance of Rsh was further determined by immunoblotting using the

protocol described below Rsh, and S13, and L33.

Preparation of fMet-tRNA$_i^{Met}$: An *E. coli* culture containing a plasmid borne expression

vector for tRNA$_i^{Met}$ under control of the LacI repressor was grown to an optical density at

600 nm of 0.4-0.6 at which time IPTG was added to induce expression of the tRNA.

After 8 h the cells were harvested, lysed by phenol, and total tRNA isolated using a

series of differential precipitation steps (Ehrenstein, 1967). Approximately half of the

recovered tRNA was a substrate for charging by MetRS. The charging reaction (2 mL)

contained approximately 200 nmoles of total tRNA and 10 nmoles of MetRS in the

presence of 1 mM methionine, 2 mM ATP, 10 mM MgCl$_2$, 20 mM KCl, 4 mM DTT, 50

μg/mL BSA, and 50 mM Tris-HCl pH 7.5. After 10 min at 37 °C a 4.0 μL aliquot of the

reaction was removed for determination of the charging efficiency as described by

Gamper et *al* (Gamper and Hou, 2020)*.* The remaining reaction was incubated for 10

min at 37 °C with 1.7 *μ*moles of 10-formyltetrahydrofolate and 20 nmoles of methionyl

formyl transferase to convert Met-tRNA$_i^{Met}$ to fMet-tRNA$_i^{Met}$. The reaction was quenched

by adding a 0.1 volume of 2.5 M NaOAc pH 5.0. Following an equal volume pH 5

phenol-chloroform-isoamyl alcohol extraction (80:17:3), the tRNA was ethanol

precipitated, dissolved in 300 μL 25 mM NaOAc pH 5.0, and stored at -70 $^{\circ}$C.

Approximately 200 nmoles of tRNA was recovered, of which 45% was fMet-tRNA$_i$^{Met}.

Reconstitution of Rsh-bound 70S initiation complex: 70S ribosomes prepared from a

low-zinc culture of isogenic Δ*rsh*/Δ*mpy*/Δ*mrf* triple mutant of *M. smegmatis* were

dialyzed at 4°C with low-magnesium buffer (20 mM HEPES-K pH7.5, 30 mM NH$_4$Cl, 0.5

mM MgCl$_2$, 5 mM β-mercaptoethanol) to dissociate the 30S and 50S subunits. Two

rounds of dialysis were performed, with fresh buffer change every four hours. Subunit

dissociation was confirmed by loading 2.4 picomoles of the dialyzed ribosomes onto a

10-40% sucrose density gradient. Another 2.4 pmoles of dissociated ribosomes were

mixed with five-fold molar excess of fMet-tRNA$_i$^{fMet}, a synthetic mRNA template

(GGCAAGGAGGUAAAAAUGUUCAAAAAA-Flour) (IDT, USA), recombinant His$_6$-IF2,

recombinant His$_6$-Rsh, and either GTP or GMP-PNP (Sigma-Aldrich) in a 100 μL

reaction in buffer I (20 mM HEPES-K pH 7.5, 30 mM NH$_4$Cl, 1 mM MgCl$_2$, 5 mM β-

mercaptoethanol), and the mixture was incubated at 37 °C for 15 minutes to allow

formation of the initiation complex on the 30S subunit. Then 25 μL of buffer II (20 mM

HEPES-K pH7.5, 30 mM NH4Cl, 50 mM MgCl2, 5 mM β-mercaptoethanol, 600mM KCl)

was added to the mixture, thereby bringing the final concentration of MgCl$_2$ and KCl to

10mM and 120 mM, respectively. The reaction was then incubated for an additional 30

minutes at 37°C to allow joining of the 50S subunit. Resulting complexes were either

used for structural analysis or resolved on a 15 mL 10-40% sucrose density gradient

centrifuged at 35000 rpm for 135 minutes using SW41 rotor on an Optima L-90K

ultracentrifuge (Beckman Coulter), and the resolved particles were fractionated as

previously described (Li *et al.*, 2021a). For immunoblotting analysis, content of the

indicated fractions were precipitated using Methanol-chloroform extraction as previously

described (Li *et al.*, 2020). Briefly, each fraction was vortexed with 4X volume of

methanol for 5-10 seconds and centrifuged for 10 seconds at 13,200 rpm on a benchtop

centrifuge before adding one volume of chloroform. The mixture was further centrifuged

at 13,200 rpm for 60 seconds, after which 3X volume of water was added and the

mixture was vortexed before further centrifugation at 13,200 rpm for 60 seconds. The

upper aqueous layer was discarded while retaining the interface and the organic layer,

which was mixed with 3X volume of methanol, vortexed and centrifuged at 13,200 rpm

for 2 minutes. The supernatant was discarded, and the resulting pellet was dried and

resuspended in 0.1 M Tris containing 8M urea for further analysis of proteins by

immunoblotting.

Immunoblotting

Total cell lysates (10 μg) or purified 70S ribosomes (2.4 pmoles) were used for detecting

S13, MPY, Mrf-FLAG, and $S14_{C-}$, while 24 pmoles ribosome were used for detecting Rsh.

Samples were resolved on 8% SDS-PAGE (for FLAG, Rsh and GroEL analysis) or 12%

SDS-PAGE (for MPY, $S14_{C-}$ and S13), and proteins were probed with anti-FLAG (1:5000,

Genescript), anti-GroEL (1:5000, Enzo), anti-RelA (1:5000), anti-S13 of *E. coli* (1:100,

The Developmental Studies Hybridoma Bank), endogenously raised anti-MPY (1:5000)

and anti-$S14_{C-}$ (1:2000) antibodies.

ppGpp extraction and analysis

Cells of specified *M. smegmatis* strains were grown in Sauton's medium with either 1 mM

Zn or 1 µM TPEN for 96 hours, after which 1 mL of the culture was labelled with 100

µCi/mL of ^{32}P KH$_2$PO$_4$ (Perkin Elmer, 900-1100mCi/mmol) for 6 hours on a shaker at 37

^{o}C. Labeled cells were washed once with TBST (20mM Tris pH7.6 + 150 mM NaCl +

0.05% Tween 80). Half of the cells (0.5 mL) were pelleted and resuspended in 25 µL of 4

M Formic acid (and kept frozen at -20 ^{o}C until ready for extraction). Remaining 0.5 mL

cells were starved for 3 hours in TBST, or metal-chelated TBST, or zinc-supplemented

TBST as specified. Following treatment, both pre- and post-starved cells were pelleted

and resuspended in 25 µL of 4 M formic acid and incubated on ice for 10 minutes. Cell

suspensions were frozen on dry ice for 10 minutes and thawed at 37 ^{o}C for 10 minutes.

The freeze-thaw cycle was repeated 5 times, after which the cell lysates were centrifuged

at 13,500 rpm for 5 minutes at 4 ^{o}C, and 5 µL of supernatants containing ^{32}P labeled

intracellular nucleotides were resolved on a 20 x 20 cm plastic PEI cellulose F TLC plate

(Millipore) using 1.5 M KH$_2$PO$_4$ pH 3.4 as a solvent. When the solvent front reached the

top of the plate, the plate was dried and exposed to X-ray film. 5 µL of 100 mM

nonradioactive nucleotides [ppGpp (Trilink), ATP, GDP and GTP (Thermo Fisher)] were

also resolved as positional markers, which were visualized by handheld UV lamp (Stuart,

Cole Palmer).

Cryo-electron microscopy and image processing

Quantifoil holey carbon copper grids R 1.2/1.3 were coated with a continuous layer of

carbon (~50 Å thick). After glow discharge for 30 s on a plasma sterilizer, 4 µl of the 200

nM Rsh-bound pre-elongation complex sample was placed on the grids. The sample

was incubated on the grids for 15 s at 4 °C and 100% humidity, followed by blotting for

4 s before flash-freezing into the liquid ethane using a Vitrobot IV (FEI). Data were

collected on a Titan Krios electron microscope at 300 keV using a K3 direct electron-

detecting camera (Gatan). −1.50 to −2.50 μm defocus range was used at a

magnification of 81,000×, yielding a pixel size of 0.846 Å. The dose rate of 23.3

electrons/pixel/second with a 2.05 second exposure time, with 41 frames of 0.05 second

duration each, resulted in a total dose of 66.59 e/Å−2. RELION 4.0 was used for data

processing. Image stacks were gain-corrected, dose-weighted, and aligned using

MotionCor2 (Zheng et al., 2017) for 9,103 micrographs. The contrast transfer function of

each aligned micrograph was estimated using CTFFIND-4.1 (Rohou and Grigorieff,

2015). A subset of micrographs was used for particle picking followed by 2D

classification. Relevant 2D classes were used for reference-based particle picking on all

micrographs yielding 1,928,609 particles. These particles were 2D classified and

suitable class averages selected (389,456 particles) (Fig. S7). The selected particles

were 3D refined and followed by 3D classification. Classes corresponding to

monosomes with 349,437 particles were selected for further processing. In order to

resolve structural heterogeneity in the monosome particles, we performed 3D

refinement using a large subunit mask followed by three rounds of fixed orientation 3D

classification. In the first round of fixed orientation 3D classification, we used an SSU

mask to resolve the SSU conformations into six classes. Class I with 158,489 particles

showed density corresponding to P-site tRNA and Rsh, representing the Rsh-bound

pre-elongation complex. We then performed the second round of fixed orientation 3D

classification on this class using a mask encompassing the entire inter-ribosomal

subunit space such as the L7/L12-stalk, A-, P-, and E-site tRNA binding sites, and the

L1-stalk regions. The second round of fixed orientation 3D classification resolved the

conformational and particularly compositional heterogeneity in the masked region. We

derived four classes with P-site tRNA density and two classes with P-site tRNA and Rsh

densities. We performed the third round of fixed orientation 3D classification combining

the two classes with P-site tRNA and Rsh densities, (49,058 particles) using a mask

encompassing the P-site tRNA and Rsh. Two of the classes with a total of 36,321

particles showed significant density corresponding to P-site tRNA and Rsh. CTF

refinement and Bayesian polishing of these particles yielded a map with a resolution of

2.7 Å.

Model building

Coordinates of the large and small subunits from our published C- *M. smegmatis*

ribosome structure (PDB:6DZI) were docked as rigid bodies into the cryo-EM map using

Chimera 1.14 (Pettersen et al., 2004). To achieve optimal fitting, we adjusted the model

based on the densities in Coot (Emsley et al., 2010). The model was subsequently

refined using PHENIX 1.14 (Adams et al., 2010). A predicted structure of *M. smegmatis*

Rsh is available in the Alphafold Protein Structure database (Jumper et al., 2021). This

structure was docked into the corresponding density in the cryo-EM map using Chimera

1.14 (Pettersen et al., 2004) and subsequently refined with Coot (Emsley *et al.*, 2010)

and PHENIX 1.14 (Adams *et al.*, 2010). A validation report for the model was obtained

from PHENIX 1.14 (Adams et al., 2010). The overall statistics of EM reconstruction and

molecular modeling are listed in Table S3. ChimeraX-1.0 (Pettersen et al., 2021) and

Chimera 1.14 (Pettersen et al., 2004) were used to generate the structural figures in the

manuscript.

Acknowledgments:

This work was supported by grants to AKO (NIH: AI132422, NIH: AI163599), CS
(AI111696), RKA (NIH:GM61576), and YMH (NIH: GM134931). R.K.A. also
acknowledges support to his lab through NIH R01 grants AI132422, GM139277 and
AI155473. Support from the Wadsworth Center core facilities–Applied Genomic
Technologies and the media core facility– is acknowledged. We also acknowledge
Wadsworth Center's and New York Structural Biology Center's (NYSBC's) 3D-EM
facilities. NYSBC EM facilities are supported by grants from the Simons Foundation
(349247), NYSTAR, the NIH (GM103310) and the Agouron Institute (F00316). Authors
acknowledge technical support from Kelley-Hurst, and discussions with Drs. Pallavi
Ghosh, Keith Derbyshire and Todd Gray.

Authors' Contribution:

YL, RT, JC and AKO designed and performed genetics and biochemical experiments;
SM, MRS, SRM, NKB and RKA designed and performed structural analysis and model
construction; JP, CS, HG and YMH provided reagents. All authors wrote the manuscript.

Financial Conflict of Interest:

Authors declare no financial conflict of interest.

Data Deposition:

The cryo-EM map and atomic coordinates are being deposited in the Electron Microscopy
and PDB Data Bank (wwPDB.org). Accession codes for these will be updated.

References

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr *66*, 213-221. 10.1107/S0907444909052925.

Andreini, C., Banci, L., Bertini, I., and Rosato, A. (2006). Zinc through the three domains of life. J Proteome Res *5*, 3173-3178. 10.1021/pr0603699.

Arenz, S., Abdelshahid, M., Sohmen, D., Payoe, R., Starosta, A.L., Berninghausen, O., Hauryliuk, V., Beckmann, R., and Wilson, D.N. (2016). The stringent factor RelA adopts an open conformation on the ribosome to stimulate ppGpp synthesis. Nucleic Acids Res *44*, 6471-6481. 10.1093/nar/gkw470.

Avarbock, A., Avarbock, D., Teh, J.S., Buckstein, M., Wang, Z.M., and Rubin, H. (2005). Functional regulation of the opposing (p)ppGpp synthetase/hydrolase activities of RelMtb from Mycobacterium tuberculosis. Biochemistry *44*, 9913-9923. 10.1021/bi0505316.

Avarbock, D., Avarbock, A., and Rubin, H. (2000). Differential regulation of opposing RelMtb activities by the aminoacylation state of a tRNA.ribosome.mRNA.RelMtb complex. Biochemistry *39*, 11640-11648. 10.1021/bi001256k.

Avarbock, D., Salem, J., Li, L.S., Wang, Z.M., and Rubin, H. (1999). Cloning and characterization of a bifunctional RelA/SpoT homologue from Mycobacterium tuberculosis. Gene *233*, 261-269. 10.1016/s0378-1119(99)00114-6.

Brown, A., Fernandez, I.S., Gordiyenko, Y., and Ramakrishnan, V. (2016). Ribosome-dependent activation of stringent control. Nature *534*, 277-280. 10.1038/nature17675.

Cashel, M., and Gallant, J. (1969). Two compounds implicated in the function of the RC gene of Escherichia coli. Nature *221*, 838-841. 10.1038/221838a0.

Chatterji, D., and Ojha, A.K. (2001). Revisiting the stringent response, ppGpp and starvation signaling. Curr Opin Microbiol *4*, 160-165. 10.1016/s1369-5274(00)00182-x.

Dahl, J.L., Kraus, C.N., Boshoff, H.I., Doan, B., Foley, K., Avarbock, D., Kaplan, G., Mizrahi, V., Rubin, H., and Barry, C.E., 3rd (2003). The role of RelMtb-mediated adaptation to stationary phase in long-term persistence of Mycobacterium tuberculosis in mice. Proc Natl Acad Sci U S A *100*, 10026-10031. 10.1073/pnas.1631248100.

Dalebroux, Z.D., and Swanson, M.S. (2012). ppGpp: magic beyond RNA polymerase. Nat Rev Microbiol *10*, 203-212. 10.1038/nrmicro2720.

Datta, P.P., Sharma, M.R., Qi, L., Frank, J., and Agrawal, R.K. (2005). Interaction of the G ' domain of elongation factor G and the C-terminal domain of ribosomal protein L7/L12 during translocation as revealed by cryo-EM. Molecular Cell *20*, 723-731. 10.1016/j.molcel.2005.10.028.

Diez, S., Ryu, J., Caban, K., Gonzalez, R.L., Jr., and Dworkin, J. (2020). The alarmones (p)ppGpp directly regulate translation initiation during entry into quiescence. Proc Natl Acad Sci U S A *117*, 15565-15572. 10.1073/pnas.1920013117.

Dutta, N.K., Klinkenberg, L.G., Vazquez, M.J., Segura-Carro, D., Colmenarejo, G., Ramon, F., Rodriguez-Miquel, B., Mata-Cantero, L., Porras-De Francisco, E., Chuang, Y.M., et al. (2019). Inhibiting the stringent response blocks Mycobacterium tuberculosis entry into quiescence and reduces persistence. Sci Adv *5*, eaav2104. 10.1126/sciadv.aav2104.

Ehrenstein, v. (1967). Isolation of sRNA from intact Escherichia coli cells. Methods in
Enzymology *12A*, 588-596.
Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of Coot.
Acta Crystallogr D Biol Crystallogr *66*, 486-501. 10.1107/S0907444910007493.
English, B.P., Hauryliuk, V., Sanamrad, A., Tankov, S., Dekker, N.H., and Elf, J. (2011). Single-
molecule investigations of the stringent response machinery in living bacterial cells. Proc
Natl Acad Sci U S A *108*, E365-373. 10.1073/pnas.1102255108.
Friesen, J.D., Fiil, N.P., Parker, J.M., and Haseltine, W.A. (1974). A new relaxed mutant of
Escherichia coli with an altered 50S ribosomal subunit. Proc Natl Acad Sci U S A *71*, 3465-
3469. 10.1073/pnas.71.9.3465.
Gabashvili, I.S., Agrawal, R.K., Spahn, C.M., Grassucci, R.A., Svergun, D.I., Frank, J., and Penczek,
P. (2000). Solution structure of the E. coli 70S ribosome at 11.5 A resolution. Cell *100*,
537-549. 10.1016/s0092-8674(00)80690-x.
Gamper, H., and Hou, Y.M. (2020). A Label-Free Assay for Aminoacylation of tRNA. Genes (Basel)
11. 10.3390/genes11101173.
Gao, X., Yeom, J., and Groisman, E.A. (2019). The expanded specificity and physiological role of
a widespread N-degron recognin. Proc Natl Acad Sci U S A *116*, 18629-18637.
10.1073/pnas.1821060116.
Gohara, D.W., and Yap, M.F. (2018). Survival of the drowsiest: the hibernating 100S ribosome
in bacterial stress management. Curr Genet *64*, 753-760. 10.1007/s00294-017-0796-2.
Gratani, F.L., Horvatek, P., Geiger, T., Borisova, M., Mayer, C., Grin, I., Wagner, S., Steinchen,
W., Bange, G., Velic, A., et al. (2018). Regulation of the opposing (p)ppGpp synthetase
and hydrolase activities in a bifunctional RelA/SpoT homologue from Staphylococcus
aureus. PLoS Genet *14*, e1007514. 10.1371/journal.pgen.1007514.
Gropp, M., Strausz, Y., Gross, M., and Glaser, G. (2001). Regulation of Escherichia coli RelA
requires oligomerization of the C-terminal domain. J Bacteriol *183*, 570-579.
10.1128/JB.183.2.570-579.2001.
Hogg, T., Mechold, U., Malke, H., Cashel, M., and Hilgenfeld, R. (2004). Conformational
antagonism between opposing active sites in a bifunctional RelA/SpoT homolog
modulates (p)ppGpp metabolism during the stringent response [corrected]. Cell *117*,
57-68. 10.1016/s0092-8674(04)00260-0.
Hood, R.D., Higgins, S.A., Flamholz, A., Nichols, R.J., and Savage, D.F. (2016). The stringent
response regulates adaptation to darkness in the cyanobacterium Synechococcus
elongatus. Proc Natl Acad Sci U S A *113*, E4867-4876. 10.1073/pnas.1524915113.
Hu, Y., and Liu, B. (2022). Roles of zinc-binding domain of bacterial RNA polymerase in
transcription. Trends Biochem Sci *47*, 710-724. 10.1016/j.tibs.2022.03.007.
Irving, S.E., Choudhury, N.R., and Corrigan, R.M. (2021). The stringent response and
physiological roles of (pp)pGpp in bacteria. Nat Rev Microbiol *19*, 256-271.
10.1038/s41579-020-00470-y.
Izutsu, K., Wada, A., and Wada, C. (2001). Expression of ribosome modulation factor (RMF) in
Escherichia coli requires ppGpp. Genes Cells *6*, 665-676. 10.1046/j.1365-
2443.2001.00457.x.

Jain, V., Saleem-Batcha, R., China, A., and Chatterji, D. (2006). Molecular dissection of the mycobacterial stringent response protein Rel. Protein Sci *15*, 1449-1464. 10.1110/ps.062117006.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583-589. 10.1038/s41586-021-03819-2.

Kudrin, P., Dzhygyr, I., Ishiguro, K., Beljantseva, J., Maksimova, E., Oliveira, S.R.A., Varik, V., Payoe, R., Konevega, A.L., Tenson, T., et al. (2018). The ribosomal A-site finger is crucial for binding and activation of the stringent factor RelA. Nucleic Acids Res *46*, 1973-1983. 10.1093/nar/gky023.

Lai, R.P.J., Cortes, T., Marais, S., Rockwood, N., Burke, M.L., Garza-Garcia, A., Horswell, S., Sesay, A.K., O'Garra, A., Young, D.B., and Wilkinson, R.J. (2021). Transcriptomic Characterization of Tuberculous Sputum Reveals a Host Warburg Effect and Microbial Cholesterol Catabolism. mBio *12*, e0176621. 10.1128/mBio.01766-21.

Li, Y., Corro, J.H., Palmer, C.D., and Ojha, A.K. (2020). Progression from remodeling to hibernation of ribosomes in zinc-starved mycobacteria. Proc Natl Acad Sci U S A *117*, 19528-19537. 10.1073/pnas.2013409117.

Li, Y., Keshavan, P., Corro, J.H., Koripella, R.K., Agrawal, R.K., and Ojha, A.K. (2021a). Purification of Hibernating and Active C- Ribosomes from Zinc-Starved Mycobacteria. Methods Mol Biol *2314*, 151-166. 10.1007/978-1-0716-1460-0_5.

Li, Y., Sharma, M.R., Koripella, R.K., Banavali, N.K., Agrawal, R.K., and Ojha, A.K. (2021b). Ribosome hibernation: a new molecular framework for targeting nonreplicating persisters of mycobacteria. Microbiology (Reading) *167*. 10.1099/mic.0.001035.

Li, Y., Sharma, M.R., Koripella, R.K., Yang, Y., Kaushal, P.S., Lin, Q., Wade, J.T., Gray, T.A., Derbyshire, K.M., Agrawal, R.K., and Ojha, A.K. (2018). Zinc depletion induces ribosome hibernation in mycobacteria. Proc Natl Acad Sci U S A *115*, 8191-8196. 10.1073/pnas.1804555115.

Loveland, A.B., Bah, E., Madireddy, R., Zhang, Y., Brilot, A.F., Grigorieff, N., and Korostelev, A.A. (2016). Ribosome*RelA structures reveal the mechanism of stringent response activation. Elife *5*. 10.7554/eLife.17029.

Marshall, R.A., Aitken, C.E., and Puglisi, J.D. (2009). GTP hydrolysis by IF2 guides progression of the ribosome into elongation. Mol Cell *35*, 37-47. 10.1016/j.molcel.2009.06.008.

Mechold, U., Murphy, H., Brown, L., and Cashel, M. (2002). Intramolecular regulation of the opposing (p)ppGpp catalytic activities of Rel(Seq), the Rel/Spo enzyme from Streptococcus equisimilis. J Bacteriol *184*, 2878-2888. 10.1128/JB.184.11.2878-2888.2002.

Ojha, A.K., Mukherjee, T.K., and Chatterji, D. (2000). High intracellular level of guanosine tetraphosphate in Mycobacterium smegmatis changes the morphology of the bacterium. Infect Immun *68*, 4084-4091. 10.1128/IAI.68.7.4084-4091.2000.

Parker, J., Watson, R.J., and Friesen, J.D. (1976). A relaxed mutant with an altered ribosomal protein L11. Mol Gen Genet *144*, 111-114. 10.1007/BF00277313.

Pausch, P., Abdelshahid, M., Steinchen, W., Schafer, H., Gratani, F.L., Freibert, S.A., Wolz, C., Turgay, K., Wilson, D.N., and Bange, G. (2020). Structural Basis for Regulation of the

855   Opposing (p)ppGpp Synthetase and Hydrolase within the Stringent Response
856   Orchestrator Rel. Cell Rep *32*, 108157. 10.1016/j.celrep.2020.108157.

857 Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
858   Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and
859   analysis. J Comput Chem *25*, 1605-1612. 10.1002/jcc.20084.

860 Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., Morris, J.H., and
861   Ferrin, T.E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators,
862   and developers. Protein Sci *30*, 70-82. 10.1002/pro.3943.

863 Pletzer, D., Blimkie, T.M., Wolfmeier, H., Li, Y., Baghela, A., Lee, A.H.Y., Falsafi, R., and Hancock,
864   R.E.W. (2020). The Stringent Stress Response Controls Proteases and Global Regulators
865   under Optimal Growth Conditions in Pseudomonas aeruginosa. mSystems *5*.
866   10.1128/mSystems.00495-20.

867 Potrykus, K., and Cashel, M. (2008). (p)ppGpp: still magical? Annu Rev Microbiol *62*, 35-51.
868   10.1146/annurev.micro.62.081307.162903.

869 Primm, T.P., Andersen, S.J., Mizrahi, V., Avarbock, D., Rubin, H., and Barry, C.E., 3rd (2000). The
870   stringent response of Mycobacterium tuberculosis is required for long-term survival. J
871   Bacteriol *182*, 4889-4898. 10.1128/JB.182.17.4889-4898.2000.

872 Prisic, S., Hwang, H., Dow, A., Barnaby, O., Pan, T.S., Lonzanida, J.A., Chazin, W.J., Steen, H., and
873   Husson, R.N. (2015). Zinc regulates a switch between primary and alternative S18
874   ribosomal proteins in Mycobacterium tuberculosis. Mol Microbiol *97*, 263-280.
875   10.1111/mmi.13022.

876 Prossliner, T., Skovbo Winther, K., Sorensen, M.A., and Gerdes, K. (2018). Ribosome
877   Hibernation. Annu Rev Genet *52*, 321-348. 10.1146/annurev-genet-120215-035130.

878 Prusa, J., Zhu, D.X., and Stallings, C.L. (2018). The stringent response and Mycobacterium
879   tuberculosis pathogenesis. Pathog Dis *76*. 10.1093/femspd/fty054.

880 Richards, J.P., Cai, W., Zill, N.A., Zhang, W., and Ojha, A.K. (2019). Adaptation of Mycobacterium
881   tuberculosis to Biofilm Growth Is Genetically Linked to Drug Tolerance. Antimicrob
882   Agents Chemother *63*. 10.1128/AAC.01213-19.

883 Rohou, A., and Grigorieff, N. (2015). CTFFIND4: Fast and accurate defocus estimation from
884   electron micrographs. J Struct Biol *192*, 216-221. 10.1016/j.jsb.2015.08.008.

885 Ronneau, S., Caballero-Montes, J., Coppine, J., Mayard, A., Garcia-Pino, A., and Hallez, R. (2019).
886   Regulation of (p)ppGpp hydrolysis by a conserved archetypal regulatory domain. Nucleic
887   Acids Res *47*, 843-854. 10.1093/nar/gky1201.

888 Ross, W., Vrentas, C.E., Sanchez-Vazquez, P., Gaal, T., and Gourse, R.L. (2013). The magic spot:
889   a ppGpp binding site on E. coli RNA polymerase responsible for regulation of
890   transcription initiation. Mol Cell *50*, 420-429. 10.1016/j.molcel.2013.03.021.

891 Sankaranarayanan, R., Dock-Bregeon, A.C., Romby, P., Caillet, J., Springer, M., Rees, B.,
892   Ehresmann, C., Ehresmann, B., and Moras, D. (1999). The structure of threonyl-tRNA
893   synthetase-tRNA(Thr) complex enlightens its repressor activity and reveals an essential
894   zinc ion in the active site. Cell *97*, 371-381. 10.1016/s0092-8674(00)80746-1.

895 Schafer, H., Beckert, B., Frese, C.K., Steinchen, W., Nuss, A.M., Beckstette, M., Hantke, I., Driller,
896   K., Sudzinova, P., Krasny, L., et al. (2020). The alarmones (p)ppGpp are part of the heat
897   shock response of Bacillus subtilis. PLoS Genet *16*, e1008275.
898   10.1371/journal.pgen.1008275.

Sprink, T., Ramrath, D.J., Yamamoto, H., Yamamoto, K., Loerke, J., Ismer, J., Hildebrand, P.W.,
Scheerer, P., Burger, J., Mielke, T., and Spahn, C.M. (2016). Structures of ribosome-
bound initiation factor 2 reveal the mechanism of subunit association. Sci Adv *2*,
e1501502. 10.1126/sciadv.1501502.
Stallings, C.L., Stephanou, N.C., Chu, L., Hochschild, A., Nickels, B.E., and Glickman, M.S. (2009).
CarD is an essential regulator of rRNA transcription required for Mycobacterium
tuberculosis persistence. Cell *138*, 146-159. 10.1016/j.cell.2009.04.041.
Takada, H., Roghanian, M., Caballero-Montes, J., Van Nerom, K., Jimmy, S., Kudrin, P., Trebini,
F., Murayama, R., Akanuma, G., Garcia-Pino, A., and Hauryliuk, V. (2021). Ribosome
association primes the stringent factor Rel for tRNA-dependent locking in the A-site and
activation of (p)ppGpp synthesis. Nucleic Acids Res *49*, 444-457. 10.1093/nar/gkaa1187.
Takada, H., Roghanian, M., Murina, V., Dzhygyr, I., Murayama, R., Akanuma, G., Atkinson, G.C.,
Garcia-Pino, A., and Hauryliuk, V. (2020). The C-Terminal RRM/ACT Domain Is Crucial for
Fine-Tuning the Activation of 'Long' RelA-SpoT Homolog Enzymes by Ribosomal
Complexes. Front Microbiol *11*, 277. 10.3389/fmicb.2020.00277.
Tamman, H., Van Nerom, K., Takada, H., Vandenberk, N., Scholl, D., Polikanov, Y., Hofkens, J.,
Talavera, A., Hauryliuk, V., Hendrix, J., and Garcia-Pino, A. (2020). A nucleotide-switch
mechanism mediates opposing catalytic activities of Rel enzymes. Nat Chem Biol *16*,
834-840. 10.1038/s41589-020-0520-2.
Turnbull, K.J., Dzhygyr, I., Lindemose, S., Hauryliuk, V., and Roghanian, M. (2019).
Intramolecular Interactions Dominate the Autoregulation of Escherichia coli Stringent
Factor RelA. Front Microbiol *10*, 1966. 10.3389/fmicb.2019.01966.
Vilcheze, C., and Jacobs, W.R., Jr. (2019). The Isoniazid Paradigm of Killing, Resistance, and
Persistence in Mycobacterium tuberculosis. J Mol Biol *431*, 3450-3461.
10.1016/j.jmb.2019.02.016.
Weiss, L.A., and Stallings, C.L. (2013). Essential roles for Mycobacterium tuberculosis Rel beyond
the production of (p)ppGpp. J Bacteriol *195*, 5629-5638. 10.1128/JB.00759-13.
Wendrich, T.M., Blaha, G., Wilson, D.N., Marahiel, M.A., and Nierhaus, K.H. (2002). Dissection
of the mechanism for the stringent factor RelA. Mol Cell *10*, 779-788. 10.1016/s1097-
2765(02)00656-1.
Winther, K.S., Roghanian, M., and Gerdes, K. (2018). Activation of the Stringent Response by
Loading of RelA-tRNA Complexes at the Ribosomal A-Site. Mol Cell *70*, 95-105 e104.
10.1016/j.molcel.2018.02.033.
Yang, X., and Ishiguro, E.E. (2001). Involvement of the N terminus of ribosomal protein L11 in
regulation of the RelA protein of Escherichia coli. J Bacteriol *183*, 6532-6537.
10.1128/JB.183.22.6532-6537.2001.
Yokoyama, T., Shaikh, T.R., Iwakura, N., Kaji, H., Kaji, A., and Agrawal, R.K. (2012). Structural
insights into initial and intermediate steps of the ribosome-recycling process. Embo J *31*,
1836-1846. 10.1038/emboj.2012.22.
Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. Nat Methods *14*, 331-332. 10.1038/nmeth.4193.
941
942

Starvation sensing by mycobacterial RelA/SpoT homologue

A

NTD   CTD

| HYD | SYN | TGS | AHD | ZBD | ACT | RelA/Rsh

B

C

D

Figure 1: Attenuated stringent response and loss of Rsh in zinc-starved mycobacteria. A. Domain architecture of *E. coli* RelA or mycobacterial RelA/SpoT homologue (Rsh). The N-terminal domain (NTD) containing the hydrolase (HYD) and synthase (SYN) is separated from the C-terminal domain (CTD) by TGS (Thr-RS, GTPase and SpoT) and alpha-helix domain (AHD). The CTD is comprised of ZBD (zinc binding domain) and ACT (aspartokinase, chorismate mutase and TyrA) subdomains. **B.** Radio-TLC showing ppGpp levels in cultures of *M. smegmatis* wildtype (WT) from high-zinc (1mM $ZnSO_4$) and low-zinc (1 μM TPEN) Sauton's medium after 96 hours of growth. ^{32}P-labeled sodium phosphate (10 μCi/mL) was added to the cultures six hours prior to starvation, after which nucleotides were extracted in formic acid and amounts equivalent to 50000 cpm were analyzed by TLC. Purified cold ppGpp was loaded as positional marker. The plot shows quantitative analysis of the ppGpp levels in high-zinc WT samples relative to their low-zinc counterparts from four biologically independent experiments. **C.** Radio-TLC showing ppGpp levels in WT cells cultured for 96-hours in high-zinc Sauton's medium, then labelled for six hours, and resuspended in either TBST, or Chelex-100 treated TBST (TBST$_{ch}$), or zinc-supplemented TBST$_{ch}$ (TBST$_{Zn}$). Δrsh mutant was analyzed as control in panels B and C. **D**. Immunoblot analysis of Rsh levels in 70S ribosomes purified from 96-hour-old high-and low-zinc cultures of *M. smegmatis* strains described in panel B. S14c- was probed for the remodeled C- ribosome, while S13 was probed as the loading control. In the set of immunoblots on the right, cell lysates from the samples used for ribosome analysis were probed. The asterisk indicates a non-specific unknown protein cross-reactive to Rsh antibody. The plots to the extreme right show normalized average difference in Rsh between high- and low-zinc cultures from three independent experiments.

Figure 2: Mrf accumulation causes loss of Rsh. A-B. Analysis of Rsh in 70S ribosomes (B) and cell lysates (C) from high- or low-zinc cultures of Δ*mrf* and complemented (Δ*mrf*comp) strains of *M. smegmatis.* High and low-zinc cultures of WT were controls. $S14_{c-}$, S13 and GroEL were probed as loading controls for ribosomes or cell lysates as indicated. The asterisk in panel B indicates a non-specific protein cross-reactive to Rsh antibody. **C.** Radio-TLC showing ppGpp levels in 96-hour old high- or low-zinc cultures of WT and low-zinc cultures of Δ*mrf and* Δ*mrfcomp* of *M. smegmatis* strains after 3-hours of starvation in TBST medium. Purified cold ppGpp was loaded as positional marker. The plot shows quantitative analysis of the relative ppGpp levels in each sample compared to the level in WT_TPEN culture from two biologically independent experiments. **D.** Ectopic Mrf accumulation causes loss of Rsh. Levels of Rsh in the cell lysates of a *M. smegmatis* strain (pYL222) expressing FLAG-Tagged Mrf* under the acetamide-inducible promoter. Cells were cultured in high-zinc Sauton's medium (1mM ZnSO4; denoted Zn) containing 0.2% succinate (denoted S) as the carbon source until OD 0.6, after which Mrf* expression was induced with 0.2% acetamide (denoted SA), and cells were grown for 96 hours from inoculation. Uninduced cells (S) were used as the reference. As controls, 96-hour old cells from the wildtype (WT) and Δ*rsh* strains cultured in either high- or low-zinc (1μM TPEN) Sauton's medium with succinate (S) or succinate + acetamide-induced (SA) conditions, as indicated. The plot shows average normalized density of Rsh from two independent experiments. **E.** Analysis of Rsh in ribosomes purified from 96-hour-old induced (SA) or uninduced (S) high-zinc cultures of pYL222 strain. **F.** Simultaneous accumulation of Rsh and Mrf in YL9 cells (Li et al. 2020) upon conditional depletion of ClpP1 protease (lane 3). Cells were grown in high-zinc Sauton's medium until OD of 0.6, after which anhydrotetracycline (Atc) was added to induce CRISPRi system to deplete ClpP1. Cell lysates were analyzed after 16 hours of ClpP1 depletion. A parallel uninduced cells were used as control (lane 4). YL9 cells grown for 96 hours in high- or low-zinc Sauton's medium without ATc (lane 1 and 2, respectively) were used as reference. The plot shows average normalized density of Rsh from two independent experiments.

A

```
Ec_RelA   QNRSKDNGRVVVEGVGNLMHHIAR CC QPIPGDEIVGFITQGRGISV H RAD C EQLAELRSH   647
Mtb_Rsh   RPRSTDDVGVSVPGAPGVLTKLAK CC TPVPGDVIMGFVTRGGGVSV H RTD C TNAASLQQ-   693
Msm_Rsh   RQRSNDDTGVAVPGAPGVLTKLAK CC TPVPGDTIMGFVTRGGGVSV H RTD C TNAESLQQ-   700
```

B

C

Figure 3: **Ribosome binding residues in Rsh are critical for its intracellular stability**. **A.** (Top) Sequence alignment of the C- terminal region of Rsh from *E. coli*, *M. smegmatis* and *M. tuberculosis* depicting the CCHC like zinc-finger (highlighted yellow) in the ZBD of Rsh. (Bottom) Superimposition of the structure (PDBID: 5IQR) of the ZBD region from *E. coli* and *M. smegmatis* (red). Zinc atom coordinated (shown with dashed lines) by the CCHC motif (shown as ball-and-stick model and numbered based on *E. coli* sequence) has been zoomed in the image to the right. The corresponding residues from *M. smegmatis* (shown in red ball-and-stick model) predict similarity in spatial orientation with respect to their *E. coli* counterparts. **B.** Immunoblot analysis of Rsh variants ($C_{666}G$, $D_{691}R$ and $C_{692}F$) in cell lysates and 70S ribosomes purified from 96-hour-old high-zinc cultures of a Δ*rsh* strain *of M. smegmatis* expressing each Rsh variant individually from an integrative plasmid (pYL238, pYL240, pYL241, pYL242 for Rsh, Rsh_{C666G}, Rsh_{D691R} and Rsh_{C692F}, respectively). Cultures of Δ*rsh* strain expressing WT RSH were controls. Normalized average levels of Rsh and its variants in cell lysates from three independent experiments are shown in the plot. **C.** *In vitro* ribosome binding of recombinant Rsh and its variants. 2.4 picomoles of 70S ribosomes purified from a 96-hour old high-zinc culture of Δ*rsh* strain of *M. smegmatis* were mixed with 1.2 picomoles of Rsh or variant proteins (purified from recombinant *E. coli* harboring pYL243, pYL246, pYL247, pY248 for Rsh, Rsh_{C666F}, Rsh_{D691R} and Rsh_{C692F}, respectively) and the mixture was incubated for 30 minutes at 37 °C. The complex was separated by ultracentrifugation through a 32% sucrose cushion. S13 was probed as the loading control. The plot shows quantitative analysis of Rsh from two biologically independent experiments.

Figure 4: **Rsh interaction with a 70S translation initiation complex. A.** Co-sedimentation of Rsh with the 50S ribosomal subunit. Pellets resulting from centrifugation on a 32% sucrose cushion were analyzed for S13 to confirm small subunit, $L33_C$ to confirm large subunit, and Rsh to test preferential subunit binding. Rsh in the absence of ribosomes was included as control to rule out sedimentation of free protein. Purified Rsh was loaded as marker. **B-C.** Reconstitution of Rsh-bound pre-elongation complex in the presence of GTP (B) or GMP-PNP (C). Dissociated ribosomes (Figure S6) were mixed with indicated ligands in the presence of GTP or GMPPNP. After incubation for 15 minutes in low-magnesium (1mM $MgCl_2$) buffer, recombinant Rsh was added, and magnesium was adjusted to 10 mM. The resulting mixture was incubated for additional 30 minutes. The complex was resolved in 10-40% SDG, and fractions corresponding to 70S-peak and its margin were analyzed for Rsh, IF2 and indicated ribosomal proteins by immunoblotting using anti-Rsh, anti-His$_6$ (for IF2), anti-S13 and anti-L31 antibodies. All reagents in the reactions in panels C and D were added in 5-fold molar excess relative to the ribosomal subunits. **D.** 2.7 Å resolution cryo-EM structure of the 70S-Rsh complex in pre-elongation state of translation, with a fMet-tRNA$_i^{Met}$ present in the P-site (green), mRNA (brown), and Rsh' two C-terminal domains, ZBD (magenta) and ACT (blue). The 30S and 50S subunits are shown in yellow and blue, respectively. **E.** Ribosomal components interacting with Rsh ZBD and ACT domain, along with the fMet-tRNA$_f^{Met}$ in the P-site (green), are shown. ZBD interacts with components of inter-subunit bridge B1a, uS19 (light green), and the A-site finger (ASF, sky blue). ACT interacts with the 50S components uL16 (sky blue), L11, and 23S rRNA helices H43/44 (sky blue). **F.** Superimposition of *M. smegmatis* 70S-Rsh, *E. coli* 70S-RelA (PDB ID: 5KPV) and *E. coli* 70S-IF2 (PDB ID: 6HTQ) structures, comparing their binding sites and conformations on the ribosome. Position of IF2 (VI-C1, cyan) suggests a direct steric clash (highlighted by red dotted circle) with the *E. coli* RelA's α-helical domain (RelA-AHD, purple). *M. smegmatis* Rsh-ACT (Ms-ACT, blue) is rotated by ~14° (indicated by a red arrow) relative to the *E. coli* RelA-ACT (Ec-ACT, light blue), while the ZBD domains (Ms-ZBD, magenta and Ec-ZBD in light purple) have the same position.

Figure 5: Conformation and interactions of the Rsh-ACT domain in a pre-elongation ribosome complex. A. Superimposition of the Rsh/Rel A interacting components from the 70S-Rsh/RelA complex structures from *M. smegmatis* (Ms) 70S-Rsh (ZBD, magenta; ACT, blue), **B.** *subtilis* (Bs) 70S-Rsh (ZBD, light pink; ACT, gray), and *E. coli* (Ec) 70S-RelA (ZBD, light purple; ACT, steel blue). Only components interacting with RelA/Rsh, L7/L12 stalk base (H43/H44 in blue), A-site finger (ASF in blue) uL16 and uL11 (light pink), the A/R tRNA (light coral) and uS19 (sea green), along with the P-site tRNA (light gray), are displayed. B. Superimposition of the C-terminal ZBD and ACT domain from Ms, Bs, and Ec (color-coded as in panel A) highlighting the rotation in Ms-ACT domain with respect to Ms-ZBD. **C.** Interactions of Ms Rsh-ACT (blue) with L11 (light pink) and the 23S rRNA helices H43/44 (light blue) are shown from two viewing directions.

Figure 6: A structural explanation of inhibition of Rsh binding to a Mpy-bound 70S ribosome. A. Superimposition of Ms 70S-Mpy (gray, PDB ID:6DZI) and Ms 70S-Rsh (gold) structures. Rsh (ZBD, magenta; ACT, blue) and Mpy (green) are shown in the space-filled rendering. **B**. The 16S rRNAs corresponding to head domain of the 30S subunit in panel A has been enlarged to highlight the their relative position in the two functional states. **C**. The position of uS19 (gray) in the Mpy-bound state would disrupt the observed interactions between Rsh-ZBD (magenta) and uS19 (orange).

Figure 7: **Model of a dynamic interaction of Rsh with the ribosome in mycobacteria, influenced by three distinct conditions**. In a nutrient-rich condition supporting normal translation, labile Rsh predominantly remains associated with the free 50S subunit, which confers stability to the protein. Rsh-50S complex joins with a 30S pre-initiation complex (pre-IC). Following the release of initiation factors, Rsh likely remains in the pre-elongation intermediate (PEI), which in an unrotated conformation is ready to elongate upon entry of the first A- site tRNA. Rsh is likely sterically displaced during intersubunit rotation associated with the first cycle of elongation. Free Rsh associates with 50S subunit and the cycle continues until a deacylated tRNA enters the A-site at the second codon (type I pause), which will lead to activation of Rsh. Rsh could also potentially be activated if free Rsh interacts with a stalled ribosome downstream of the second codon (Type II pause), although its frequency could be limited by the efficiency of free Rsh in finding such complexes. Conditions inducing hibernating ribosomes, which appear to be less permissive to Rsh *in vivo*, would sequester ribosomes from translation cycle, thereby exposing Rsh to degradation in a Clp protease dependent manner.

Figure S1: A full-gel immunoblot of cells lysates from WT and Δ*rsh* strains cultured in high-(1mM ZnSO$_4$; abbreviated as Zn) and low-zinc Sauton's medium (1μM TPEN; abbreviated as T) and probed with anti-Rsh antibody. Data shows lack of any Rsh-derived smaller product in the low-zinc culture. GroEL from the lysate was probed as a loading control.

Figure S2: Effect of ribosome remodeling on Rsh-ribosome interaction.
Immunoblot analysis of Rsh bound to 70S ribosomes from 96-hour old high-zinc cultures of WT *M. smegmatis* and its two recombinant strains, Δ*zur* and *pYL53,* which constitutively expressed the remodeled (C-) ribosomes. Low-zinc culture of WT was used as control.

Figure S3: Levels of Mpy in the purified 70S ribosomes from high- and low-zinc cultures of WT and Δ*rsh*. S14$_{c-}$ and S13 were probed as controls as indicated.

A

B

Figure S4: Level of Rsh-bound ribosomes (A) and ppGpp (B) in *M. smegmatis* before and after exposure to starvation (TBST) from high- or low-zinc culture conditions after 28 hours of growth.

Figure S5: Purified recombinant 6XHis-tagged Rsh proteins expressed from an IPTG-inducible T7 promoter in the *E. coli* BL21 strain and visualized by SDS-PAGE and the Coomassie Blue stain. Each recombinant protein was purified from the same volume of starting cells using the same method, and equal volume of each sample was loaded on the gel.

Figure S6: Sucrose density gradient profile of dissociated ribosomes used in the reconstitution experiments described in figure 4. The reconstituted ribosomal were also used for structural analysis.

A

B

C

Selected 2D classes
(389,456 particles)
↓
Consensus refinement
↓
3D classification
↓
Selected monosome classes
(349,437 particles)
↓

Large subunit
refinement

Small subunit
fixed orientation
3D classification

| Class I | Class II | Class III | Class IV | Class V | Class VI |
| (158,489 particles) | (71,984 particles) | (96,654 particles) | | | |

Empty
22,308 particles

P-site tRNA and Rsh
densities

Mask:L7/L12-stalk+APE-site+L1-stalk
Fixed orientation
3D classification

| Class I | Class II | Class III | Class IV | Class V | Class VI |
| (25,230 particles) | (23,828 particles) | | | | |

P-site tRNA density
109,854 particles

Mask:Rsh+P-tRNA
Fixed orientation
3D classification

Class I	Class II	Class III
		weak Rsh densities
		12,739 particles

Refined map
(2.7 Å)
36,321 particles

D

FSC

2.7 Å

Resolution 1/Å

Figure S7: Image processing of *M. smegmatis* 70S-Rsh-fMet-tRNA$_i$^Met complex. Image processing of *M. smegmatis* 70S-Rsh-fMet-tRNAi^Met complex. A. a representative micrograph from the *M. smegmatis* 70S-Rsh-fMet-tRNAi^Met complex dataset. **B.** Representative 2D class averages used for the 3D reconstructions. **C.** Flowchart showing the details of 3D classifications and refinements. **D.** Fourier-shell correlation (FSC) plot for the final map (36,321 particles) used in this study. Local resolution for this map is shown in figure S8.

Figure S8: Local resolution of the cryo-EM map of the *M. smegmatis* 70S-Rsh-fMet-tRNAi^Met complex. Refined by using (A) 70S monosome, (B) 50S LSU, and (C) 30S SSU masks. In panels B and C, SSU and LSU appear to be partially disordered, as expected. A cutting plane has been applied to show the core of the ribosome in each case.

Rsh-ZBD

Rsh-ACT

Figure S9: Modeling of the ZB- and ACT domains into corresponding cryo-EM densities. Two different density threshold values were used for modeling the two domains, as density for the main anchoring ZBD (pink) was stronger than that for the ACT domain.

Figure S10: Species-specific differences in Rsh/RelA-ZBD interaction with uS19, uS13, and ASF. Top- Multiple sequence alignment of a short stretch in the ZBD domain of *M. smegmatis* (Ms) Rsh, *B. subtilis* (Bs) Rsh, and *E. coli* (Ec) RelA. Amino acid substitutions at four regions (indicated by red arrows) that result in species-specific differences in interactions (i1-i4) of the ZBD domain with ribosomal components. Bottom left- Superimposition of Ms 70S-Rsh (ZBD, magenta) and Bs 70S-Rsh (ZBD, light pink). Interactions of Bs-Rsh-ZBD with the ribosome that are absent in the case of Ms-Rsh-ZBD are displayed: two with uS19 (i1 and i2), one with uS13 (i3), and one with ASF (i4). Bottom right- Superimposition of Ms 70S-Rsh (ZBD, magenta) and Ec 70S-RelA (ZBD, light purple). Ec RelA-ZBD has two extra interactions with the ribosome, as compared to Ms Rsh-ZBD: One with uS19 (i2) and another with ASF (i4).

Table S1: List of plasmids and strains

Plasmids	Remarks	Reference
pJl37	P_{hsp60}-based expression vector for mycobacteria; *kan^r*	[1]
pMH94	L5-attp-based integrative vector for mycobacteria; *kan^r*	[1]
pYUB854	Cosmid vector, hygr	[1]
pTTPla	Tweety-attp-based integrative vector for mycobacteria; *kan^r*	[1]
pLam12	$P_{acatamidase}$-based expression vector for mycobacteria; *kan^r*	[1]
pJV53-SacB	Sucrose-sensitive marker SacB cloned in pJV53 @ SpeI site; *kan^r*	[1]
pET21b	Vectors carry an N-terminal T7-Tag sequence plus an optional C-terminal His Tag sequence, *amp^r*	[1]
pYL3	*kan^r* cassette in pMH94@EcoRV & NotI sites is replaced by *hyg^r* cassette from pYUB854; *hyg^r*	[2]
pYL40	*Msmeg_6069* internal deletion in pYL42 backbone @ SacI & XbaI, *hyg^r*	[1]
pYL53	MSMEG_6065-6070 cloned in pYL3 @ SacI+XbaI under a constitutive promoter (P^{const}): mutant variant of $P^{zur-box}$, *hyg^r*	[1]
pYL155	C-terminally FLAG-tagged *mrf* with P^{zurbox} cloned in pTTPla @ SacI & XbaI sites; *kan^r*	[1]
pYL181	C-terminally FLAG-tagged *mrf* with P^{zurbox} cloned in pYL180 @ SacI & XbaI sites; *apr^r*	[1]
pYL206	L5-attp-based integrative vector carrying tet-inducible dCas9 and gRNA complementary to Clpp1; *kan^r*	[1]
pYL222	*Msmeg_6069* with all six his/cys mutations (H to A, C to A) and Flag tag at C-terminal and p^{zur-box} cloned in plam12 @ XbaI & HindIII sites; *kan^r*	This Study
pYL236	*MSMEG_2965* (T518A) + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This Study
pYL238	*MSMEG_2965* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This Study
pYL239	*MSMEG_2965* (S519A) + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This Study
pYL240	*MSMEG_2965* with C666G mutation + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This Study
pYL241	*MSMEG_2965 with D691R mutation* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This Study
pYL242	*MSMEG_2965 with C692F mutation* + 500 bp UPS cloned in pMH94 @ KpnI & XbaI; *kan^r*	This study
pYL243	*MSMEG_2965* with N-terminal His-tag cloned in pET21b @ NdeI & HindIII; *amp^r*	This study
pYL246	*MSMEG_2965* (C666G) with N-terminal His-tag cloned in pET21a @ NdeI & HindIII; *amp^r*	This study
pYL247	*MSMEG_2965* (D691R) with N-terminal His-tag cloned in pET21a @ NdeI & HindIII; *amp^r*	This study
pYL248	*MSMEG_2965* (C692F) with N-terminal His-tag cloned in pET21a @ NdeI & HindIII; *amp^r*	This study
pYL259	*MSMEG_2965* + 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg^r*	This study
pYL260	*MSMEG_2965* (C666G)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg^r*	This study
pYL261	*MSMEG_2965* (D691R)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg^r*	This study
pYL262	*MSMEG_2965* (C692F)+ 500 bp UPS cloned in pYL3 @ KpnI & XbaI; *hyg^r*	This study
Strains	**Remarks**	**Reference**
mc²155 (WT)	High-Frequency Transformation strain of *M. smegmatis* as parent wild-type	[2]
Δmsm_c-	Unmarked Δ*Msmeg_6065-6070* in mc²155	[2]
Δmrf (YL1)	Unmarked Δ*c-* operon harboring pYL40; *hyg^r*	[1]
*Δmrf*comp (YL2)	Unmarked Δ*c-* operon harboring pYL40 and pYL155; *hyg^r, kan^r*	[1]
Δmpy (YL3)	Unmarked Δ*Msmeg_1878* in mc²155	[2]
Δmpy/*Δmrf* (YL4)	Unmarked Δ*Msmeg_6065-6070*/Δ*Msmeg_1878* harboring pYL40; *zeo^r, hyg^r*	[1]

Δ*zur* (YL5)	Unmarked Δ*Msmeg_4487* in mc^{2}155	[2]
Δ*zur*/Δmrf (YL6)	Unmarked Δ*Msmeg_6065-6070* /Δ*zur* harboring pYL40; *hyg^r*	[1]
YL9	Δ*zur* harboring pYL181 and pYL206; *apr^r, kan^r*	[1]
Δ*rsh*	Δ*Msmeg_2965* in mc^{2}155, *zeo^r*	This Study
mc^{2}7000	*M. tuberculosis* H37Rv:*ΔRD1:ΔpanCD* as parent wild-type	[2]
Δ*mpy$_{mtb}$*	*ΔRv_3241c* in mc^{2}7000 or Mtb(Erd)	[1]
Δ*mrf$_{mtb}$*	*ΔRv_0106* in mc^{2}7000	[1]
Δ*mrf$_{mtb}$*comp	*ΔRv_0106* harboring pYL149 in mc^{2}7000	[1]

1.   Li Y, Corro JH, Palmer CD, Ojha AK: **Progression from remodeling to hibernation of ribosomes in zinc-starved mycobacteria**. *Proceedings of the National Academy of Sciences of the United States of America* 2020, **117**(32):19528-19537.
2.   Li Y, Sharma MR, Koripella RK, Yang Y, Kaushal PS, Lin Q, Wade JT, Gray TA, Derbyshire KM, Agrawal RK *et al*: **Zinc depletion induces ribosome hibernation in mycobacteria**. *Proc Natl Acad Sci U S A* 2018, **115**(32):8191-8196.

Table S2: Synthetic oligonucleotides and IF2 sequences

Name	Sequence	Used in:
His1-R	GGTGCGGACGACGACAGCGCCGTCGAAGGTGGCGCTGACGAGCAGGGT	pYL222
His1-F	ACCCTGCTCGTCAGCGCCACCTTCGACGGCGCTGTCGTCGTCCGCACC	
His2-R	GGCCGAGACGGCTCCGGCCACCAGCTCAAGTAC	
His2-F	GTACTTGAGCTGGTGGCCGGAGCCGTCTCGGCC	
His3-R	CACGTCGGCGCGCCGGGCCAGACGGCGAAGCAGGAT	
His3-F	ATCCTGCTTCGCCGTCTGGCCCGGCGCGCCGACGTG	
p6069-CXXCR	ATCGTCACGCACGGTGGCCGAGACGGCTCCGTGCACCAGCTC	
p6069-CXXCF	GAGCTGGTGCACGGAGCCGTCTCGGCCACCGTGCGTGACGAT	
Pzurbox 6069-XF	CGCGCCGTTCGTCTAGACGCTGCACCAGTTCTCGCC	
p6069-FLAGHR	GGCCGGAAGCTTTCACTTATCGTCGTCATCCTTGTAATCCGATTGCTCTCCTGT	
pRelAMS-up500KpnIF	CTTCCCCGAACCGGTACCCCAGTTCAAGGACCTCAC	pYL236,238, 239,240,241, 242,256,259, 260,261,262
pRelAMS-XbaIR	CGCCGAACCTCTAGATTAGGCCGCGCTGGTGACGCG	
RelAC666G-F	ACCAAGCTGGCCAAGGGCTGCACCCCGGTGCCG	pYL240,246, 260
RelAC666G-R	CGGCACCGGGGTGCAGCCCTTGGCCAGCTTGGT	
RelAD691R-F	AGCGTGCACCGCACCCGTTGCACCAACGCCGAG	pYL241,247, 261
RelAD691R-R	CTCGGCGTTGGTGCAACGGGTGCGGTGCACGCT	
RelAC692F-F	GTGCACCGCACCGACTTCACCAACGCCGAGTCG	pYL242,248, 262
RelAC692F-R	CGACTCGGCGTTGGTGAAGTCGGTGCGGTGCAC	
preIA-NdeIHISF	TGACACATATGCACCACCACCACCACCACGTCGACGAGCCAGGCAAG	pYL243,246, 247,248
preI-HindIIItgaR	CGCCGAACCGCTAAGCTTTCAGGCCGCGCTGGTGACGCG	

> Sequence of codon optimized synthetic IF2:
ATACGAGAACCATatgCACCACCACCACCACCACCAGGCACCGGTCGTGGGAGGGGTAAGACTTCCTCA
CGGTAACGGCGAGACGATCCGTCTGGCACGCGGCGCTTCATTATCAGATTTTGCTGAAAAGATAAATG
CGAATCCGGCTTCACTTGTGCAGGCATTGTTCAATCTTGGAGAAATGGTTACTGCCACGCAAAGTGTCG
GTGATGAAACACTTGAGCTGTTAGGGTCGGAGATGAATTACAATGTACAGGTCGTATCACCAGAAGACG
AGGACCGTGAATTGCTTGAAAGCTTTGATTTAACCTACGGCGAAGACGCGGGTGACGAGGAGGACTTA
GAGGTGCGTCCACCCGTTGTAACCGTTATGGGCCATGTCGATCATGGTAAGACTAGACTTCTGGACAC
AATCCGGAAGGCAAATGTGAGAGAGGGCGAGGCCGGCGGTATAACGCAGCACATCGGGGCCTATCAG
GTTGAGGTTGATTTGGACGGGACGGTGCGTCCCATAACTTTTATTGATACTCCCGGTCACGAGGCCTTT
ACCGCAATGCGTGCCAGAGGTGCCAAGGCTACAGATATTGCTATTTTAGTGGTAGCCGCGGATGATGG
CGTGATGCCTCAAACGGTTGAAGCGATTAACCACGCACAGGCCGCGGACGTTCCGATAGTGGTAGCA
GTCAACAAAATTGATAAGGAAGGAGCGGACCCCGCTAAAATTCGTGGGCAGTTAACAGAATATGGATTA
ATACCGGAAGAGTATGGCGGAGACACCATGTTTGTAGATATCAGTGCGAAACAGGGAACAAATATTGAA
GCATTATTAGAAGCAGTGATTCTTACCGCAGACGCCTCTTTAGATTTGCGTGCAAACCCCGATATGGAA
GCTCAGGGAGTCGCTATAGAAGCCCATCTTGACAGAGGTCGGGGTCCTGTTGCTACTGTATTGATACA

GCGCGGCACCCTTCGGGTTGGTGATTCCGTTGTTGCGGGTGACGCTTATGGGAGAGTCCGGAGAATG
ATCGATGAACATGGAGAGGACGTCGAAGAAGCATTGCCGTCTCGGCCGGTCCAGGTAGTTGGTTTCAC
CTCGGTACCCGGCGCTGGGGACAATTTTTTAGTTGTAGATGAAGATCGGATCGCGCGTCAAATCGCGG
ATCGCCGGTCAGCGCGTAAACGGAACGCCCTTGCTGCACGTAGTCGCAAACGGATCAGCTTGGAGGA
CTTAGACTCCGCCCTTAAAGAGACGTCGCAATTAAACCTTATATTAAAAGGAGACAACGCGGGGACAGT
TGAAGCACTGGAAGAGGCCTTAATGGGAATCCAGGTGGACGACGAGGTAGAACTGCGCGTCATCGAC
CGCGGTGTCGGAGGCGTCACAGAAACGAATGTAAACTTAGCAAGTGCCTCGGACGCGATTATCATTGG
TTTTAACGTCCGGGCTGAAGGAAAGGCGACTGAATTGGCTAATAGAGAAGGAGTTGAAATACGCTATTA
TTCTGTAATCTACCAGGCGATTGATGAAATTGAAGCAGCACTTAAAGGAATGCTGAAGCCTGTATACGA
AGAGAAGGAGCTTGGGCGTGCAGAAATACGTGCCATCTTCCGCAGTTCTAAGGTGGGCAATATAGCTG
GCTGCCTTGTCACATCAGGAATAATGCGCCGCAACGCTAAGGCCCGTTTATTAAGAGACAATGTAGTAG
TTGCACAGAATTTGACGGTAAGTAGTTTAAGACGCGAAAAGGATGACGTAACGGAAGTTCGTGATGGTT
ACGAGTGCGGACTGACCTTAACCTATAATGACATCAAAGAAGGTGACGTAATTGAGACTTATGAATTAG
TTGAAAAAGCGCGTACTtgaCTCGAGTGGCCGATCC

Table. S3 Data collection, Refinement and Model Validation parameters.

Description	70S-Rsh-fMet-tRNA$_i^{Met}$ complex
Data collection	
Microscope	FEI Titan Krios
Voltage (kV)	300
Pixel size (Å)	0.846
Defocus range (μm)	-1.50 to -2.50
Average e$^-$ dose per image (e$^-$/ Å^2)	66.59
Particles (initial)	1,928,609
Particles (final)	36,321
FSC-threshold	0.143
Resolution (Å)	2.7
Map-sharpening B factor (Å^2) overall	-44.3
Refinement	
RMS deviations	
Bonds (Å)	0.007
Angles (°)	0.923
MolProbity score	1.80
Clash score	6.43
Rotamer outliers (%)	0.61
Ramachandran plot	
Outliers (%)	0.16
Allowed (%)	6.72
Favored (%)	93.12
RNA	
Correct sugar puckers (%)	99.4
Angle outliers (%)	0.02
Bond outliers (%)	0.00
Good backbone conformations (%)	75.9
Model composition	
RNA bases	4843
Protein residues	11,068