LHX2 in germ cells control tubular organization in the developing mouse testis

Neha Singh[1], Domdatt Singh[1], Anshul Bhide[1], Richa Sharma[1], Shilpa Bhowmick[2], Vainav
Patel[2] and Deepak Modi[1*]

[1]Molecular and Cellular Biology Laboratory, ICMR-National Institute for Research in
Reproductive and Child Health, Indian Council of Medical Research (ICMR), JM Street,
Parel, Mumbai 400012, India

[2]Viral Immunopathogenesis Laboratory, ICMR-National Institute for Research in
Reproductive and Child Health, Indian Council of Medical Research (ICMR), JM Street,
Parel, Mumbai 400012, India

ORCiD ID:
Neha Singh (ORCiD ID: 0000-0001-7088-839X)
Domdatt Singh (ORCiD ID: 0000-0002-3361-0388)
Anshul Bhide (ORCiD ID: 0000-0003-4592-7451)
Richa Sharma (ORCiD ID: 0000-0001-8490-9761)
Shilpa Bhowmick (ORCiD ID: 0000-0002-9586-3989)
Vainav Patel (ORCiD ID: 0000-0001-6661-3238)
Deepak Modi (ORCiD ID: 0000-0002-4230-4219)

Running Title: LHX2 in germ cell controls testicular organization

* Corresponding author
Dr. Deepak Modi
Scientist F and Head
Molecular and Cellular Biology Laboratory
ICMR-National Institute for Research in Reproductive and Child Health
Indian Council of Medical Research (ICMR)
J.M. Street, Parel, Mumbai 400 012
India
Tel no: 91-22-24192034
Fax no: 91-22-24139412
Email: deepaknmodi@yahoo.com, modid@nirrch.res.in

Keywords: Homeobox, LIM-HD gene, Knockout, Sex determination, Gonad, development,
Interstitial cell, Endothelial cell, Infertility, Male, Seminiferous tubules

Interstitial cells
Sertoli Cells
Germ Cells
Endothelial Cells

- Disrupted tubular organization
- Altered angiogenesis
- Interstitial cell expansion
- Altered Sertoli cell functions

Wild type testis

***Lhx2* knockout testis**

Highlight

- *Lhx2* is expressed in germ cells of developing mouse testis but is dispensable for germ cell and Sertoli cell specification
- Loss of *Lhx2* disrupts testicular vascularization, leads to the expansion of interstitial cells, and alters the tubular organization
- Germ cells govern the tubular organization in the developing testis

Abstract

In the gonads of mammalian XY embryos, the organization of cords is the hallmark of testis

development. This organization is thought to be controlled by interactions of the Sertoli cells,

endothelial and interstitial cells with little or no role of germ cells. Challenging this notion,

herein we show that the germ cells play an active role in the organization of the testicular

tubules. We observed that the LIM-homeobox gene, *Lhx2* is expressed in the germ cells of

the developing testis between E12.5-E15.5. In *Lhx2* knockout- fetal testis there was altered

expression of several genes not just in germ cells but also in the supporting (Sertoli) cells,

endothelial cells, and interstitial cells. Further, loss of *Lhx2* led to disrupted endothelial cell

migration and expansion of interstitial cells in the XY gonads. The cords in the developing

testis of *Lhx2* knockout embryos are disorganized with a disrupted basement membrane.

Together, our results show an important role of *Lhx2* in testicular development and imply the

involvement of germ cells in the tubular organization of the differentiating testis.

Introduction

Mammalian sex determination is characterized by the differentiation of the bipotential gonad into the testis or an ovary [1]. In the XY embryos, testicular development requires Sertoli cell specification and tubular organization, vascularization, and differentiation of the interstitial cells to Leydig cells, peritubular myoid cells, and other cell types. In the XY embryos, SRY-positive precursors differentiate into Sertoli cells that act as a scaffold for the germ cells to aggregate, organize and start forming the testicular cords [2,3]. The number of Sertoli cells is the key determinant of tubulogenesis and reduced numbers of the Sertoli cells compromise tubular organization [4].

The tubular organization is further supported by the migrating vascular endothelial cells from the adjacent mesonephros [5]. Blocking mesonephric cell migration in developing testis either directly by a barrier [6,7] or by inhibiting VE-Cadherin [8,9] leads to defective cord formation. Sertoli cells secrete a plethora of factors that activate multiple signal transduction events in the endothelial cells to aid in their migration from the mesonephros [10–12]. Inhibition of the PI3K signaling pathway [13], PDGF signaling [14,15], or VEGF signaling [8,16–18] disrupts the organization of tubules in the developing testis. These observations imply the involvement of Sertoli and endothelial cell communication for the tubular organization.

In addition to endothelial cells, interstitial progenitors, fetal Leydig cells, peritubular myoid cells, and macrophages also aid in tubule organization. Blocking the macrophage migration into the testis [9], genetic disruption of gene encoding activin A from fetal Leydig cells [19], and defective or reduced numbers of peritubular myoid cells [20–22] result in altered tubular morphology. These results imply the involvement of interstitial cells in the structural organization of tubules in testis development.

Somatic cells have an unequivocal role, while the role of germ cells (if any) in the tubulogenesis organization is underexplored. A single study has demonstrated that fetal mice lacking germ cells (due to a point mutation in the cKIT gene) have defective tubule organization due to compromised Sertoli cell functions [23]. However, it is unclear if the tubules were hypoplastic due to the physical absence of the germ cell core or due to the absence of some factors secreted by the germ cells. Thus, the involvement of germ cells in the tubular organization is unclear.

Lim homeobox gene 2 (*Lhx2*), is a transcription factor of the LIM-HD family, having roles in

brain patterning, eye development, olfaction path, hair folliculogenesis, limb bud outgrowth,

and several others [24,25]. Though the functions of *Lhx2* are well-reported in the developing

nervous system, its expression and function in the developing gonads is not explored.

Members of the LIM-HD family are expressed in the developing gonads of multiple species

including humans, of them the roles of *Lhx1*, *Lhx8,* and *Lhx9* are well reported [26,27].

Recently we reported that the Lim-HD gene *Lhx2* is expressed in the germ cells of the

developing XX gonads, and loss of *Lhx2* promotes endothelial cell migration in the

developing ovary [26]. To our knowledge, this is the first study demonstrating the

involvement of germ cells in controlling endothelial cell migration and vascularization.

Whether *Lhx2* is also expressed in the germ cells of XY gonads and if it governs

vascularization is unknown. Further, the functions of *Lhx2* in the developing gonads beyond

the suppression of vascularization are not known.

In the present study, we aimed to investigate the expression of *Lhx2* and its function in

developing testis. The results revealed that *Lhx2* is expressed by germ cells of the fetal testis

and loss of *Lhx2* leads to disrupted tubular organization.

Materials and Methods

Animal Ethics, Mice Strain and nomenclature

The study was approved by the institutional animal ethics committee (IAEC Project No.,

27/14, 2/16, 16/16). *Oct4*-GFP transgenic mice [28] and cKit$^{W/+}$ mice [29] were kindly

provided by Dr. K.V.R. Reddy, NIRRCH. FVB mice constitutively expressing Green

fluorescent protein (GFP) were provided by the National Institute of Immunology (NII), New

Delhi, India. Heterozygous mice with a targeted disruption of *Lhx2* [30] and those with

conditional *Lhx2* allele [31] were kindly provided by Prof. Shubha Tole, TIFR, Mumbai,

India. All the animals were bred and maintained at the Experimental Animal Facility of the

Institute.

In the present study, pups of pregnant dams with *Lhx2* flox alleles along with tamoxifen-

inducible CreER and treated with tamoxifen are referred to as *Lhx2$^{flox/flox}$*. Pregnant dams with

Lhx2 flox alleles without the tamoxifen-inducible CreER and treated with tamoxifen are

labeled as wild-type (WT) controls. In the case of standard *Lhx2* knockouts, the homozygous

knockout pups are referred to as *Lhx2*-/- (or mutants) and the homozygous wild-type pups as

WT controls. For the germ cell-depleted mice, the cKit$^{+/+}$ are WT and cKit$^{W/W}$ are
homozygous mutants.

Animals were mated and the day of vaginal plug was considered as 0.5; the pups were
collected from E11.5 to E15.5 as per experimental requirement. The gonads were dissected
under a stereomicroscope (Olympus; Tokyo, Japan); the mesonephros was removed and the
gonad was stored in RNALater for RNA extraction. For histology and immunolabeling
experiments, the whole pups or isolated gonads were fixed overnight in 4%
paraformaldehyde and processed for paraffin embedding and sectioning.

Genotyping

Primer sequences for all the genotyping are mentioned in Supplementary Table. 1.
Genotyping was done using the Terra™ PCR Direct kit (Takara Bio, Kusatsu, Shiga Japan).
Embryonic sex and *Lhx2* allele genotyping were performed as mentioned [26]. For germ cell-
depleted gonads, the embryos from cKit$^{W/+}$ crosses were collected and genotyping was done
by PCR [29]. In brief, the PCR product of 231 bp was digested with Hph1 restriction enzyme
and resolved on 15% polyacrylamide gel. cKIT$^{W/W}$ mutant pups lacked fragments of 40 bp
and 70 bp bands while homozygous WT lack a fragment of 111 bp.

RNA extraction, cDNA synthesis, and real-time PCR

RNA extraction was done using RNeasy mini kit (Qiagen, Germany). cDNA was synthesized
using High Capacity cDNA synthesis kit (Applied Biosystems, ThermoFisher Scientific, MA,
USA) as mentioned earlier [26]. Real-time PCR was performed using Eva Green chemistry
(BioRad Laboratories Inc., CA, USA) in the CFX96 Real-Time PCR System (BioRad) as
detailed previously [26]. PCR was carried out in duplicates, for all biological replicates. Gene
expression was normalized to *Gapdh* and data were analyzed using the Pfaffl method [32].
The primer sequences, their annealing temperatures, and expected product sizes are given in
Supplementary Table 1.

Section *In Situ* hybridization

Section *in situ* hybridization was performed on 7 μm thick paraffin sections of wild-type fetal
testes at E13.5 as detailed previously [33,34]. In brief, deparaffinized sections were
acetylated, treated in proteinase-K, and pre-hybridized at 42^{0}C. The plasmids for *Lhx2* sense

and antisense probe preparation were a kind gift from Prof. Shubha Tole [35]. Digoxigenin

(DIG)-labeled sense and antisense riboprobes were prepared using the *in vitro* transcription

DIG labeling kit (Roche, IN, USA). Hybridization was carried out at 60^0C overnight and the

slides were washed and probed with an alkaline phosphatase-conjugated anti-DIG antibody

(Roche) at 1:2000 dilution. The slides were washed and detection was performed at pH 9.5

by incubating in nitro blue tetrazolium and 5-bromo-4-cholro-2-indoyl phosphate solution

(Roche) as per manufacturer instructions. The reaction was stopped, the slides were

dehydrated, air-dried, and mounted. The slides were observed under a brightfield microscope

(Olympus) and representative areas were photographed.

Fluorescence-Activated Cell Sorting (FACS) from *Oct4*-GFP testis

XY gonads from *Oct4*-GFP transgenic pups were collected on E13.5 and E15.5 and cell

sorting was done [36]. Briefly, the gonads were incubated in 0.25% trypsin-EDTA (Gibco,

Thermo Fisher Scientific, MA, USA) at 37^0C for 20 min and the reaction was stopped by

adding an equal volume of DMEM medium supplemented with 10% fetal bovine serum (both

from Gibco), followed by mechanical disruption. After gating for live cells cell sorting was

performed based on GFP on BD FACS Aria fusion Flow Cytometer (BD Biosciences). The

GFP positive (germ cells) and negative (somatic cells) live cells (propidium iodide negative)

were sorted (Supplementary Figure. 1A), and subjected to RNA extraction, cDNA synthesis,

and qPCR as described above.

Histology and Immunohistochemistry/Immunofluorescence

Hematoxylin-Eosin staining and immunohistochemistry were performed on 5 μm thick

paraffin sections as mentioned [37,38]. For immunohistochemistry, antigens were retrieved

by boiling the sections in Tris-EDTA Buffer (pH 9), followed by blocking in 5% bovine

serum albumin (Sigma-Aldrich). Sections were probed overnight with primary antibodies.

Negative controls were incubated either with isotype controls or Phosphate Buffered Saline

instead of the primary antibody. Detection was done using the Horse Radish Peroxidase

(HRP) streptavidin method using ABC staining system (Santa Cruz Biotechnology; Texas,

USA). All sections were briefly counterstained with Mayer's hematoxylin and mounted in

DPX liquid mountant (Himedia, Maharashtra, India). Slides were viewed under a brightfield

microscope (Olympus) and representative areas were photographed using a digital camera

(Olympus). For VE-cadherin, the signals were detected using a donkey anti-rabbit secondary

antibody conjugated with Alexa Fluor 568 dye (Invitrogen, A10042; 1:1000 dilution) as
described previously [39].

For multiplex immunofluorescence, we used the Tyramide signal amplification (TSA)
detection system (PerkinElmer, MA, USA, or Biotium, Inc. CA, USA) as per the
manufacturer's protocol. In brief, the sections were deparaffinized, rehydrated, and re-fixed
in 10% neutral buffered formalin. Antigen retrieval was done in Tris-EDTA buffer (pH 9).
Slides were washed in 1X tris-buffered saline with 0.1% Tween-20 followed by incubation in
blocking solution for 10 min at room temperature. Sections were probed with primary
antibody against (anti-DDX4 for germ cells), for 30 min at room temperature. Incubation
with anti-rabbit-secondary antibody tagged with HRP was done for 10 min and signal
amplification was performed using fluorophore 570. The antibody complex was stripped in
boiling in antigen retrieval treatment for 30 minutes, blocking was done and sections were
probed with anti-SOX9 antibody (for Sertoli cells) for 30 min at room temperature.
Downstream steps were repeated and fluorophore 690 was used to amplify the signals.
Eventually, sections were mounted with EverbriteTM hardest mounting medium (Biotium).
Laminin staining was done using the same procedure except that the detection was done
using the fluorophore 520. Details of all the antibodies and their optimized concentrations
used are given in Supplementary Table 2

Slides were viewed and images were captured using a fluorescence microscope equipped
with an sCMOS camera (Leica Microsystems DMi8, Mannheim, Germany). The contrast was
enhanced digitally but raw images were used for the quantification (Fiji version of ImageJ).

RNA Sequencing

The strategy of gonad collection from $Lhx2^{flox/flox}$ pups for RNAseq is detailed previously
[26]. A single dose of tamoxifen (75µg/g) was given orally to gravid females on E9.5 and
gonads were collected on E12.5 from the WT and $Lhx2^{flox/flox}$ embryos. This concentration
and time were chosen as they gave maximum efficiency of knockout [26,31]. Approximately,
30 pairs of gonads were pooled, RNA extracted, and outsourced for RNA sequencing on the
Illumina NextSeq 500 platform (Sandor Biosciences, Hyderabad, India). An average of, 45
million raw reads were generated per sample. Alignment was done using Tophat 2.0.13
software and mm84 sequence as reference (ftp://ftp.ensembl.org/pub/release-
84/fasta/mus_musculus/dna/).

Analysis of differentially expressed genes in cell-type-enriched transcriptome of the developing testis

Genes with fragments per kilobase per million (FPKM) values >0.5 were selected and the fold change between the *Lhx2^{flox/flox}* and WT was calculated. Genes with fold change ≤0.5 were taken as downregulated whereas those with fold change ≥1.5 were taken as upregulated. The list of genes enriched in the supporting cells, germ cells, endothelial cells, and interstitial cells were retrieved from the dataset GSE27715 [40] and the FPKM values from the WT and *Lhx2^{flox/flox}* testis were compared. The fold change values were calculated as above. The differentially expressed genes for the total and cell type-specific transcriptomes were subjected to gene ontology analysis using DAVID (https://david.ncifcrf.gov/tools.jsp). In addition, the expression profile of LIM-HD co-regulators in the different cell types of developing testis during the window of sex determination was determined from the dataset GSE27715 [40].

Gonad Recombination Assay and assessment of cell migration

The scheme of the gonad recombination assay is described in detail previously [26]. In brief, the gonads and mesonephros were carefully dissected from WT and *Lhx2-/-*. XY embryos at E12.5. The dissected gonad was carefully overlayed onto the donor mesonephros from GFP XY embryos of the same gestational age and the assembly was cultured at 37^0C. The. Recombinants were imaged for the migration of the GFP-positive cells at 24, 48, and 72 hours post-recombination.

Quantification of angiogenic patterns was done using the angiogenesis analyzer macro in ImageJ software as detailed previously [26]. In brief, the background of desired areas was corrected and the images were subjected to the angiogenesis analyzer (ImageJ) that quantified various branching parameters.

Imaging, Cell number counting, and Morphometric Analysis

Quantification of all signal intensities in immunofluorescence images was done using the LasX software (Leica Microsystems) or ImageJ (https://ij.imjoy.io/). For quantification, 5 random areas were selected per section and 3 non-serial sections per animal were analyzed. For quantifying tubule numbers and abnormal tubules, five hematoxylin and eosin-stained testis sections (each 20μm apart) were analyzed. The total number of tubules was counted in

each section and the number was divided by the area of the section. Tubules appearing close
to each other without a clear interstitium were defined as fused tubules. Incomplete tubules
were those which did not have well-defined boundaries and appeared interspersed with the
stroma. Histologically well-defined tubules without an outer layer of cells were defined as
tubules without peritubular myoid cells.

The numbers of DDX4-positive and SOX9-positive cells per testis cord were counted using
Image J software. Briefly, the putative tubule area was marked using the region of interest
(ROI) tool and the number of SOX9 or DDX4 positive cells in that area were quantified in
single-channel images. Three to five non-serial sections per animal were quantified.

ImageJ was used to measure the length (the longest diameter), width (the shortest diameter),
and rete testis length (testicular region attached to mesonephros) of WT and mutant testis.

Statistical analysis and data presentation

247 Data were analyzed statistically using an unpaired Student-t-test or multiple t-test with Holm-
248 Sidak corrections considering random distribution and unequal variance. All results were
249 presented as mean $\pm$ standard deviation (SD) and graphs were plotted on GraphPad Prism
250 (Version 8, CA) or Microsoft Excel.

Results:

Lhx2 **is expressed in germ cells of developing mouse testis**

Fig. 1A shows the quantification of *Lhx2* mRNA in the testis of wild-type (WT) embryos
from E11.5 to E15.5 (n=6 testis/time point). *Lhx2* transcripts were detected in all the samples
irrespective of the gestational age, their levels were highest at E11.5 which gradually and
significantly declined at all the time points tested. To determine the cellular localization of
Lhx2, RNA *in situ* hybridization (n=5) and immunohistochemistry (n=7) were performed on
fetal testis sections at E13.5. *Lhx2* transcripts (Fig. 1B) were localized in cells within the
cords of developing testis; the interstitium and the celomic epithelium were devoid of any
signals indicating that *Lhx2* is either in the developing Sertoli cells or in the germ cells. No
signals were detected in the sections incubated with the sense probe. Immunohistochemical
analysis (Fig. 1C) revealed that LHX2 was localized in large-size germ cells (open arrow)
enclosed within the testis cords. SOX9, a positive control, stained the Sertoli cells around the

germ cells (open arrow). No staining was detected in the negative control sections. These
results implied that *Lhx2*/LHX2 may be expressed in the germ cells.

To determine if *Lhx2* is enriched in the germ cells, we flow-sorted the somatic and germ cell
of the fetal testis (pools of 14-16 testes) at E13.5 (3 pools) and E15.5 (2 pools). These time
points were chosen because germ cells are committed to the gametogenic fate by E12.5 and
have exited mitosis and committed to the spermatogenic fate by E15.5. The purity of the
fractions was confirmed where the GFP+ fraction had a higher abundance of *Ddx4* (germ cell
marker) with negligible amounts in the GFP- somatic cell fraction. Conversely, the somatic
fraction had abundant *Amh* or *Sox9* mRNA (Sertoli cell markers) with negligible amounts in
the germ cell fraction. In these fractions at E13.5 (Fig 1D) and E15.5 (Fig1E), *Lhx2*
transcripts were detected in the GFP+ germ cell fraction with negligible amounts in the GFP-
somatic cell fraction.

We further validated these findings using a germ cell-depleted fetal mice testis ($cKIT^{W/W}$)
model. RNA from WT ($cKIT^{+/+}$) and germ cell-depleted testis ($cKIT^{W/W}$) at E12.5
(n=3/genotype) and E15.5 (n=3/genotype) were subjected to qPCR (Fig. 1F). Results
revealed that at both the time points, *Lhx2* and *Ddx4* were detected in the WT testis but
negligible signals were detected in the mutant testis. These results together confirm that *Lhx2*
is expressed in the germ cells of the developing testis.

Lhx2 requires a group of co-regulators for its appropriate functions and the relative
abundance of these factors in the cells is a determinant of the activity of LHX2 [27]. We
observed that of the *Lhx2* co-regulators, *Ldb1* was abundantly expressed in all the cell types
of the developing testis, while *Ldb2* was enriched in the endothelial and interstitial lineages
with lower expression in supporting cells and least in germ cells (Supplementary Figure. 1B).

Of the Lim-only factors (Supplementary Figure. 1B), *Lmo3* was not reported in the dataset.
Lmo1 had higher expression in the germ cells as compared to the other three cell types. *Lmo2*
was enriched in endothelial cells as compared to other cell types. *Lmo4* was expressed in
most somatic cells with the highest expression in supporting lineage and the lowest in the
germ cells.

Fig. 1: *Lhx2 is expressed in germ cells of developing mouse testis* (A) mRNA expression of *Lhx2* in XY gonads at E11.5-E15.5. Data is presented as mean ± SD (n=6 testis/timepoint). Y axis is fold change of normalized expression value where the mean of E11.5 was taken as 1. (B) Section *in situ* hybridization for *Lhx2* at E13.5 testis (n=5). Purple staining is a positive reaction. Inset is sense probe control. Bar is 50μm. (C) Immunostaining for LHX2 in XY gonads at E13.5 (n=7). SOX9 is a positive control. Arrowheads indicate germ cells. Inset is negative control without primary antibody. Bar is 50μm. Detection of *Lhx2* mRNA in flow-sorted germ cells and somatic cells of XY gonads at E13.5 (D) & E15.5 (E). Germ cells were flow-sorted from XY mice expressing GFP under the *Pou5f1* promoter. Germ cell and somatic cell fractions from several mice at E13.5 and E15.5 were pooled and qPCR for *Ddx4* (germ cell marker), *Amh/Sox9* (somatic cell markers), and *Lhx2* was carried out for three independent batches at E13.5 and two batches at E15.5. Y axis is fold change of normalized relative expression where the mean of the data for germ cell fraction was taken as 1 (for *Ddx4*

and *Lhx2*) and somatic cell fraction was taken as 1 for *Amh/Sox9*. (F) qPCR for *Ddx4* and *Lhx2* in XY gonads of cKit knockout (cKit$^{W/W}$) mice at E12.5 and E15.5 (n=3 /timepoint/genotype). Data is presented as mean $\pm$ SD. Y axis is the fold change of normalized expression values where the mean value of WT is taken as 1. In all qPCR experiments, *Gapdh* was used for normalization. *, **, **** are p<0.05, p< 0.01, and p<0.0001 respectively. In all graphs, each dot represents one biological replicate.

Loss of *Lhx2* alters the transcriptome of the developing testis

To reveal the role of *Lhx2* in the developing testis, RNA-seq of WT and *Lhx2$^{flox/flox}$* testis at E12.5 was performed (n=30 pairs each). This time point was chosen as most cell types are committed to the male lineage in the testis. The levels of *Lhx2* mRNA was negligible in *Lhx2$^{flox/flox}$* testis while it was detected in the WT testis (Supplementary Figure. 2A) indicating successful floxing of the gene. A total of 2993 genes (21% of the transcriptome) were differentially expressed (Supplementary Table. 3). Amongst these, 718 (5.10%) genes were downregulated while 2275 (16.16%) genes were upregulated in *Lhx2$^{flox/flox}$* testis as compared to WT (Fig. 2A). The list of all differentially up and downregulated genes are given in Supplementary Table. 4. Of the upregulated genes, 73 genes were exclusively expressed in *Lhx2$^{flox/flox}$* testis. Of the downregulated genes, 92 genes were exclusively expressed in WT testis and not detected in the *Lhx2$^{flox/flox}$* testis (Supplementary Table. 4). *Ifi27l2a* gene was the maximally upregulated while *Tmem100* was the most downregulated gene in absence of *Lhx2* (Supplementary Table. 4).

Gene Ontology analysis revealed that the differentially expressed genes were associated with biological processes like regulation of transcription, anterior/posterior patterning, cell proliferation, cell differentiation, cell migration, axon guidance, apoptosis, proximal/distal pattern formation, lipoprotein metabolic process, and angiogenesis (Fig. 2B). The associated molecular functions were protein/DNA binding, transcription factor activity, transcriptional repressor activity, chromatin binding, frizzled & β-catenin binding, ECM structural constituent, cadherin binding, GPCR binding, and actin-binding activity (Fig. 2C). In pathway analysis, there was enrichment of signaling pathways (MAPK, WNT, PI3K-Akt, Hippo, Ras, p53, Rap1, and NF-kappa B), pathways in cancer, complement and coagulation cascades, Focal/Cell adhesion, ECM-receptor interactions, and regulation of pluripotency (Fig. 2D).

Fig. 2: Loss of *Lhx2* alters the transcriptome of the developing testis RNAseq for wild type (WT) and *Lhx2^flox/flox^* XY embryos done at E12.5 (A) The percentage of upregulated (fold change ≥1.5) and downregulated genes (fold change ≤0.5) in the testis of *Lhx2^flox/flox^* embryos as compared to WT. The false discovery rate (FDR) of the enriched biological processes (B), molecular functions (C) and, pathways (D) for the differentially expressed genes are given.

Loss of *Lhx2* in the developing XY gonads alters the transcriptome in multiple cell types

The effects of loss of *Lhx2* in the germ cells on the expression of genes enriched in different testicular cell types were examined (Supplementary Table. 5). Information on 871/961 germ cell-enriched genes was available in our dataset. Nearly 13% of germ cell-enriched genes

14

were differentially expressed, of which 8% genes were upregulated while 5% genes were downregulated (Fig. 3A). GO analysis revealed that differentially regulated genes were associated with regulation of histone methylation, regulation of transcription, and cell differentiation.

Of the 540 supporting lineage-enriched genes, information on 505 genes was available in our dataset. Approximately 16% of genes were differentially expressed in absence of *Lhx2*. Almost 11% of genes were upregulated and 5% were downregulated (Fig. 3B). The differentially expressed genes in the supporting lineage were associated with biological processes such as cell differentiation and cell-cell adhesion.

Of the 435 endothelial cell-enriched genes, information for 392 was available in our dataset. Nearly, 23% were differentially expressed in the *Lhx2^{flox/flox}* testis. Of these genes, 16% were upregulated and 7% were downregulated. The differentially expressed endothelial cell-enriched genes were associated with biological processes like angiogenesis, integrin-mediated signaling, regulation of immune response, apoptosis, nervous system development, and osteoclast differentiation (Fig. 3C).

There are 164 genes highly enriched in the interstitial cells of which information for 141 genes was available. Approximately, 22% were differentially expressed, where 19% were upregulated and 3% of genes were downregulated in *Lhx2^{flox/flox}* testis (Fig. 3D). These differentially expressed genes were associated with organ morphogenesis, cell migration, axon guidance, and proteolysis.

15

Fig. 3: Loss of *Lhx2* in the developing XY gonads alters the transcriptome of multiple cell types The list of genes enriched in each of the four cell types in the developing testis was extracted from the dataset GSE27715. The pie chart represents the proportion of differentially expressed genes in the Germ cells (A), Supporting cells (B), Endothelial cells (C), and Interstitial cells (D) of the testis of wild-type and *Lhx2^flox/flox^* embryos at E12.5. The percentage of up and downregulated genes are given. The false discovery rate (FDR) of the

375 enriched biological processes for the differentially expressed genes in each of the cell-types

376 are shown as bar plots.

Germl cell numbers and gametogenic fate is not compromised in *Lhx2-/-* XY gonads

The mRNA levels of the germ cell marker *Ddx4* (Fig. 4A) was similar in WT and *Lhx2-/-*

testis at E13.5 (n=15 for both the genotypes), E14.5 (WT n=15, *Lhx2-/-* n=12) and E15.5

(WT n=8, *Lhx2-/-* n=6). The overall intensity of immunostained DDX4 protein was also

similar in WT (n=6) and *Lhx2-/-* (n=8) XY gonads at E14.5 (Fig. 4B & C). The numbers of

DDX4-positive cells and their intensity per cord were also similar in the WT and *Lhx2-/-*

testis (Supplementary Figure. 2B and C).

The DDX4-positive germ cells are pluripotent and acquire the gametogenic fate by

expressing *Dazl* and the spermatogenic fate by expressing *Nanos2* and this process is

completed by E15.5 [41]. We analyzed their mRNA levels at E15.5 (WT n=8, *Lhx2-/-* n=6)

and observed that *Dazl* and *Nanos2* mRNA were similar in the WT and *Lhx2-/-* testis.

However, the expression of pluripotency marker *Pou5f1* was reduced significantly (p<0.05)

in *Lhx2* knockout testis as compared to controls (Fig. 4D). The expression of germ cell-

specific genes (*Ddx4*, *Pou5f1*, *Dazl*, *Piwil2,* and *Dnmt3L*) were unchanged in the WT and

Lhx2$^{flox/flox}$ testis at E12.5 (Supplementary Figure. 2D). These observations imply that in

absence of *Lhx2*, the number of germ cells and their commitment to gametogenic fate is not

compromised though there are changes in the expression of a fraction of genes in germ cells

(Fig. 3A).

Fig. 4: Germ cell numbers and gametogenic fate is not compromised in the developing testis in absence of *Lhx2* (A) qPCR for *Ddx4* mRNA in wild type (WT) and *Lhx2-/-* testis at E13.5 (n=15/genotype), E14.5 (WT n=15, *Lhx2-/-* n=12) and E15.5 (WT n=8, *Lhx2-/-* n=6). (B) Immunostaining for DDX4 (germ cell marker) on sections of WT (n=5 pairs) and *Lhx2-/-* (n=7 pairs) testis at E14.5. The scale bar is 100μm. Negative control without primary antibody is shown in inset (C) Quantification of DDX4 protein intensity in WT (n=6, *Lhx2-/-* n=8) testis at E14.5. Each dot represent a single region of interest and data from different regions in multiple sections of WT (n=6) and *Lhx2-/-* (n=8) testis are plotted. (D) qPCR for *Pou5f1*, and *Nanos2* mRNA in WT (n=8) and *Lhx2-/-* (n=6) testis E15.5. Graphs are plotted as mean $\pm$ SD. Y axis is fold change where the mean value of WT was taken as 1. *Gapdh* is used for normalization in all qPCR experiments. In qPCR graphs, each dot represents one replicate. * indicates p<0.05.

Loss of *Lhx2* does not alter Sertoli cell specification in the developing XY gonads

Loss of germ cells from developing testis due to a mutation in the cKIT gene (cKIT$^{W/W}$) leads to a reduction in Sertoli cell number [23]. This prompted us to investigate the role of germ cell-specific *Lhx2* on the numbers and differentiation of Sertoli cells in the developing testis. Histologically, the overall testis appeared similar in *Lhx2-/-* (n=4) and WT (n=8) XY embryos (Fig. 5A). The mRNA levels of *Amh* (Sertoli cell marker) was not significantly

different in the *Lhx2-/-* (n=12) and WT (n=15) testis at E14.5 (Fig. 5B) indicating that the Sertoli cells differentiated appropriately in absence of *Lhx2*. The mRNA of the female supporting cell marker, *FoxL2* was not detected in both WT (n=6) and *Lhx2-/-* (n=6) testis at E14.5 (Fig. 5C) indicating that there was no somatic cell sex reversal. Similar results were also seen at E15.5 (not shown). Somatic sex reversal was not observed at E12.5 as the expression profiles of male and female sex-determining marker genes were similar in the WT and *Lhx2$^{flox/flox}$* XY gonads (Supplementary Figure .2E). To test if Sertoli cell numbers or their differentiation was altered in absence of *Lhx2*, we profiled the expression of *Sox9*/SOX9 between E13.5 to E15.5, during the entire course of the tubular organization (Fig. 5D-G). The results revealed that levels of *Sox9* mRNA at E13.5 (WT n=15, *Lhx2-/-* n=15), E14.5 (WT n=15, *Lhx2-/-* n=12) and E15.5 (WT n=8, *Lhx2-/-* n=6) and SOX9 protein (n=6/timepoint/genotype) did not alter between WT and *Lhx2-/-* testis. However, at E15.5 the SOX9 protein intensities were marginally but significantly lower in the *Lhx2-/-* testis as compared to controls (Fig. 5D-F). Also, the number of SOX9-positive Sertoli cells per tubule was significantly reduced at both E13.5 and E15.5 (Fig. 5G). However, the total numbers of SOX9+ cells per unit area were comparable in the WT and *Lhx2-/-* testis (Supplementary Figure. 2F). These results imply that *Lhx2* is dispensable for male sex determination and Sertoli cell differentiation although a proportion of the Sertoli cell transcriptome and numbers of Sertoli cells per tubule are altered (Fig. 3B).

Fig 5: Loss of *Lhx2* does not alter Sertoli cell specification in the developing XY gonads
(A) Histology of the wild type (n=8) and *Lhx2-/-* (n=4) XY gonads at E14.5. The scale bar is
50μm. (B & C) qPCR for *Amh* (Sertoli cell marker) and *FoxL2* (granulosa cell marker) in the
developing gonads at E14.5 in WT (n=15) and *Lhx2-/-* (n=12) XY gonads and WT ovaries
(n=6). *Gapdh* is used for normalization in all qPCR experiments. (D) Immunostaining for
SOX9 (Sertoli cell marker) in WT and *Lhx2-/-* XY at E13.5 and E15.5
(n=6/genotype/timepoint). Scale bar is 100μm. Inset is isotype control. (E) qPCR for *Sox9*
mRNA at E13.5 (n=15/genotype), E14.5 (WT n=15, *Lhx2-/-* n=12) and E15.5 (WT n=8,
Lhx2-/- n=8). Quantification of SOX9 protein (F) and SOX9 positive cell number (G) in WT
and *Lhx2-/-* testis at E13.5 & E15.5 as indicated. Each dot represent a single region of interest
and data from multiple independent regions in different sections of WT and *Lhx2-/-* testis are
plotted. All graphs are plotted as mean ± SD. Y axis is fold change where the mean value for

446 WT is taken as 1. In qPCR graphs, each dot represents one replicate. * and **** indicate

447 p<0.05 and p<0.0001 respectively.

Testicular vascularization is disrupted in absence of *Lhx2*

The observation that angiogenesis is one of the enriched biological processes in the

differential transcriptome of the *Lhx2$^{flox/flox}$* testis prompted us to investigate the phenomenon

further. We observed that of the 30 pro-angiogenic factors, 30% were upregulated while 10%

were downregulated in absence of *Lhx2* (Fig. 6A; Supplementary Table 7) suggesting that

vascularization may be altered in absence of *Lhx2*. We performed the immunoexpression of

endothelial cell marker VE-Cadherin in testis sections at E13.5 when the microvasculature is

organizing [42,43]. As compared to WT (n=3), there was a higher proportion of VE-Cadherin

positive cells in the gonadal stroma of *Lhx2-/-* (n=3) testis (Fig. 6B). In the knockouts, many

of these cells appeared in clusters in between the tubules indicating that endothelial cell

organization is disrupted in the *Lhx2-/-* testis. The same was also reflected in whole mounts

where we observed that in the wild-type testis, microvasculature was well organized which

culminated in the coelomic blood vessel; in the *Lhx2-/-* testis there were hemorrhagic areas

and the blood vessels were clumped in the intertubular space with no clear coelomic blood

vessel (Fig. 6C). These results together indicate that testis vascularization is disrupted in

absence of *Lhx2* in the germ cells.

Since the testicular transcriptome revealed enrichment of pro-angiogenic genes and there was

disruption of testicular vascularization, we quantified the mRNA expression of pro-

angiogenic factors *Vegfa*, *Vegfb*, and *Pdgfa* by qPCR at E15.5 when the process of testicular

vascularization is completed. The results (Fig. 6D) revealed that the levels of all these genes

were significantly upregulated in *Lhx2-/-* (n=6) as compared to the WT (n=8) testis. Previous

studies have reported that knocking out of *Wnt4*, *Rspo1*, or *Ctnnb1* causes ectopic

vascularization in XX gonads [44] and the overexpression of *Wnt4* disrupts vascularization in

the XY gonads [45]. As Wnt/ β-catenin signaling pathways were enriched in the

transcriptome *Lhx2* knockouts testis (Fig. 2), we hypothesized that altered expression of

Wnt4/Rspo1/β-catenin may have caused the disrupted vascularization in absence of *Lhx2*.

However, at E15.5, the mRNA expression of *Wnt4*, *Rspo1,* and *Ctnnb1* were identical in

Lhx2-/- (n=6) and WT (n=8) testis (Fig. 6E). These results imply that disrupted angiogenesis

in absence of *Lhx2* is not due to alterations in the *Wnt4*, *Rspo1,* and *Ctnnb1* genes but due to

elevated expression of pro-angiogenic factors like *Vegfa*, *Vegfb*, and *Pdgfa*.

Testicular vascularization involves the migration of endothelial cells from the mesonephric vascular plexus into the gonads [8]. To investigate if the loss of *Lhx2* in the germ cells affects endothelial cell migration, we carried out gonad recombination assays and quantified the migratory pattern of endothelial cells (Fig. 6F & G). The results revealed that in the WT XY gonads (n=3), GFP-positive cells had a linear path and the cells migrated to the anterior aspect of the gonads towards the coelomic epithelium. However, in the *Lhx2-/-* gonads (n=6), GFP-positive cells had a very distorted path that deviated from their course not reaching towards the coelomic epithelium (Fig. 6F). Quantitatively, as compared to WT gonads, the measure of branching such as extremes, nodes, and junctions twigs, segments, and branches were significantly increased in absence of *Lhx2* (Fig. 6G). These results imply that the absence of *Lhx2* in the germ cells alters the intra-testicular milieu to disrupt endothelial cell migration and vascularization.

Fig. 6: Vascularization is disrupted in the developing testis in absence of *Lhx2*. (A) The proportion of pro-angiogenic factors altered in *Lhx2^flox/flox^* testis at E12.5. (B) Immunostaining

of VE-cadherin (endothelial cell marker) in wild type (WT) and *Lhx2-/-* testis at E13.5 (n=3 /genotype). Scale bar is 50μm. Inset is negative control without primary antibody. (C) Brightfield image of WT (n=6) and *Lhx2-/-* (n=6) whole testis at E14.5. Scale bar is 200μm qPCR for selected pro-angiogenic factors (D) and for Wnt-βcatenin genes (E) in WT (n=8) and *Lhx2-/-* (n=6) testis at E15.5. *Gapdh* is used for normalization in all qPCR experiments. (F) Gonad recombination assay of WT and *Lhx2-/-* testis overlayed on GFP (green) mesonephros from WT embryos. The recombinants were imaged 72 hrs after co-culture. Scale bar is 400μm. (G) Images were analysed using ImageJ with angiogenesis analyser plugin and the branching parameters (X axis) of the GFP-positive cells were quantitatively assessed in WT (n=3 recombinants) and *Lhx2-/-* (n=6 recombinants). All graphs are plotted as mean $\pm$ SD. Y axis is fold change where the mean value for WT is taken as 1. In all graphs, each dot represents one replicate. * and ** indicate $p<0.05$ and $p<0.01$ respectively.

Expansion of interstitial cells in absence of *Lhx2* in developing XY gonads

In the interstitial compartment, 22% of the enriched genes were differentially expressed and these were associated with positive regulation of cell proliferation and cell migration (Fig. 3D). To test if the interstitial compartment was altered in absence of *Lhx2*, we immunostained for the interstitial cell marker PDGFRα [46] and proliferation marker MKI67 at E14.5 when the interstitial compartment is well demarcated from the tubules. The results revealed that the PDGFRα+ cells were localized through the interstitial compartment in the wild-type testis (n=8). In the *Lhx2-/-* (n=8), PDGFRα immunoreactivity was more intense specifically in the area below the celomic epithelium (Fig. 7A). MKI67 positive cells were localized in the interstitium, their numbers appeared to be higher in the *Lhx2-/-* testis as compared to wild type (n=8/genotype). The cells below the celomic epithelium were intensely labeled with MKI67 in the *Lhx2-/-* testis (Fig. 7B). At E15.5, the mRNA levels of interstitial progenitor cell markers [46] viz *Lhx9, Nr5a1*, and *Pdgfra* were increased in the *Lhx2-/-* testis (n=6) as compared to WT (n=8). The levels of the early progenitor marker *Gata4* was, however, unaltered (Fig. 7C). As compared to WT (n=8), the mRNA levels of fetal Leydig cell markers *Hoxb2*, and *Cyp17a1* were decreased in *Lhx2-/-* (n=6) testis at E15.5 (Fig. 7D). The reduction in levels of *Hoxb2* but not in *Cyp17a1* was statistically significant. These results imply that in the absence of *Lhx2* there is interstitial cell expansion and defective Leydig cell differentiation

524
Fig. 7: Expansion of interstitial cells in absence of *Lhx2* in the developing XY gonads.
Immunostaining for (A) PDGFRα (interstitial marker) and (B) MKI67 (proliferation marker)
in wild type (WT) and *Lhx2-/-* (n=8/genotype) testis at E14.5. Scale bar is 50μm. qPCR for
(C) interstitial progenitors *(Gata4, Lhx9, Nr5a1,* and *Pdgfra)* and (D) fetal Leydig cell
markers (*Hoxb2*, and *Cyp17a1*) in the WT (n=8) and *Lhx2-/-* (n=6) testis at E15.5. Graphs are
mean $\pm$ SD. Y axis is fold change where mean value of WT was taken as 1. *Gapdh* is used for
normalization in all qPCR experiments. In all graphs, each dot represents one replicate. * ,**
and *** are p value<0.05, <0.01, and <0.001 respectively.

The absence of *Lhx2* in germ cells leads to a defective tubular organization

Since there was a disruption in the supporting, endothelial and interstitial lineages (all of
which aid in the tubular organization) we asked if the absence of *Lhx2* in the germ cells
disrupts tubulogenesis. Towards this, co-staining of SOX9 and DDX4 was performed to
visualize the testicular cords organization (Fig. 8A). As evident, the pattern of tubule
organization was different in the WT (n=6) and mutant testis (n=8) at E14.5. In the WT, the
cords were uniformly sized and had similar shapes while in *Lhx2* mutants, many tubules were
abnormally shaped. Some tubules were fused, some were depleted of germ cells and others
had mis-spaced Sertoli cells. The number of Sertoli cells per tubule was also reduced (Fig.
5G). Laminin staining was done to demarcate the basement membrane (Fig. 8B), and we
observed that in the WT testis (n=5), laminin was uniformly detected at the base of the
tubules. In the *Lhx2-/-* testis (n=5) there was irregular laminin deposition and it did not
appear well demarcated as in controls.

We quantified the numbers of tubules per unit area in each of the histological sections of WT

and *Lhx2-/-* testis at E13.5 (WT n=14, *Lhx2-/-* n=13), E14.5 (WT= n=14, *Lhx2-/-* n=12) and

E15.5 (WT n=14, *Lhx2-/-* n=13) (Fig. 8C) and observed that overall there was a significant

increase in the number of tubules in the *Lhx2-/-* testis as compared to WT. However, most of

the tubules in the *Lhx2-/-* testis were not normal and histological analysis revealed that

(Supplementary Figure. 3) in some areas the testicular cords were not lined by peritubular

myoid cells, many appeared fused or incomplete, and some even lacked germ cells. These

phenotypes were quantified (Fig. 8D) and we observed an increase in the numbers of tubules

without peritubular myoid cells layer, fused appearing tubules, and incomplete appearing

tubules in the *Lhx2-/-* (n=4) testis as compared to the WT controls (n=8). The empty tubules

were rarely detected in the WT, but several tubules were devoid of germ cells in the *Lhx2*

knockouts. These results imply that the organization of tubules is altered in absence of *Lhx2*.

Analysis of the whole testis from WT (n=6) and *Lhx2-/-* (n=5) embryos at E15.5 (when the

testis cords are completely organized), revealed that overall cords arrangements and cords

length were significantly altered in *Lhx2-/-* testis (Fig. 8E). While the WT testis consistently

showed well-organized tubules reaching the coelomic epithelium with the longest tubules in

the center and the shortest at the poles, in the *Lhx2-/-* testis most tubules did not reach up to

the coelomic region, and they were of different heights with shorter tubules interspersed

between tall tubules resulting in irregular tubular arrangements. Quantification of the testis

dimension (Fig. 8F) showed that as compared to WT (n=6), the length (left to right distance

in the mid-region) and the rete length (left to right distance at the site of attachment to

mesonephros) were significantly increased in the *Lhx2-/-* (n=4), the width was significantly

reduced. These results imply that the organization of the tubule and testicular dimensions are

altered in absence of *Lhx2*.

Fig. 8: Absence of *Lhx2* leads to defect in the organization of testis cords (A) Co-immunostaining of SOX9 and DDX4 in wild type (WT n=6) and *Lhx2-/-* (n=8) testis at E14.5. Bar is 100μm (B) Immunofluorescent detection of Laminin in the WT and *Lhx2-/-* testis (n=5/genotype) at E14.5. Scale bar is 100μm. (C) Quantification of the number of testicular cords in WT and *Lhx2-/-* testis at E13.5 (WT n=14, *Lhx2-/-* n=13), E14.5 (WT n=14, *Lhx2-/-*n=12) and E15.5 (WT n=14, *Lhx2-/-* n=13). (D) Quantification of tubular phenotypes in testicular sections of the WT (n=8) and *Lhx2-/-* (n=4) embryos at E14.5. PMCs=peritubular myoid cells. (E) Brightfield image of WT and *Lhx2-/-* whole testis at E15.5 (WT n=6, *Lhx2-/-* n=5). White asterisks mark the distal tip of cords. Scale bar is

580   200µm in (D) (E) (F). Quantification of length, width and rete length of WT and *Lhx2-/-*

581   (n=4/genotype) testis at E15.5. In all graphs, data is mean $\pm$ SD of the fold change where the

582   mean value of WT was taken as 1. In all graphs, each dot represents one replicate and *, **,

583   *** and **** are p value <0.05, <0.01, <0.001 and <0.0001 respectively.

Discussion

The present study demonstrates that *Lhx2* is expressed in the germ cells of developing testis

and the absence of *Lhx2* alters the endothelial and interstitial cell compartments without

affecting germ cell and Sertoli cell specification resulting in the disrupted tubular

organization.

Lhx2 along with the other twelve members of the LIM-HD family and their co-factors are

expressed in the developing gonads of both sexes during the window of sex determination

[27]. Of these, only *Lhx1, Lhx8,* and *Lhx9* are known to have roles in the development of the

reproductive system, the functions of others are not explored [27]. Previously, we have

shown that *Lhx2* is expressed by the germ cells of the developing ovary [26]. Extending these

findings, herein we show that *Lhx2* mRNA and protein are also expressed in the germ cells of

the developing testis. Interestingly, *Ldb1* and *Lmo1* which are coregulators for *Lhx2* [47–49]

are also enriched in the germ cells of the developing testis implying that the *Lhx2* co-

regulator complex system may be functional in the developing germ cells.

The closest homolog of *Lhx2* is *Lhx9* which has overlapping expression patterns and

functions in the brain [50]. However, in the developing gonads, *Lhx9* is expressed in somatic

cells [51] while *Lhx2* is in the germ cells suggesting their divergent role in gonad

development. Indeed, in the *Lhx9* knockouts, the gonad development is arrested before the

onset of sex determination; the ovary [26] and the testis (present study) develop normally in

absence of *Lhx2*. Also, the specification of germ cells and Sertoli cells is unaltered in *Lhx2-/-*

testis, and neither the number of germ cells and Sertoli cells are compromised in *Lhx2-/-* XY

gonads. Together with our previous findings [26], our results imply that *Lhx2* is dispensable

for gonad development, germ cell specification, and sex determination of the supporting

lineage in both sexes.

Germ cell-specific expression of *Lhx2* in developing testis intrigued us to investigate its role

in testis development. RNAseq analysis revealed that the expression of almost 21% of genes

are altered in the developing testis in absence of *Lhx2*. The enriched biological processes

27

included anterior/posterior patterning, proximal/distal axis formation, limb bud
morphogenesis, cell proliferation, cell differentiation, cell migration, axon guidance, and
angiogenesis. The involvement of *Lhx2* in the proliferation and migration of neural stem and
progenitor cells and in various cancers is well established [52,53]. *Lhx2* is involved in
anterior/posterior and proximal/distal pattern formation and morphogenesis, neuron
migration, and limb bud morphogenesis in many species [25,54]. Since these processes are
also enriched in the testis (present study) and ovary [26] of *Lhx2* knockouts, it is tempting to
propose that there would be a core transcriptome regulated by *Lhx2* which has overlapping
functions irrespective of the tissue.

Since *Lhx2* is expressed by the germ cells, we first determined the changes in the expression
of germ cell-enriched genes and observed that nearly 13% of them were altered. These genes
were associated with germ cell development, chromatin remodeling, epithelial cell
proliferation, and cell differentiation. However, the number of germ cells and their
acquisition of gametogenic and spermatogenic fate is not altered in absence of *Lhx2*. These
results imply that *Lhx2* is dispensable for the development and differentiation of male germ
cells in the fetal testis although a proportion of its transcriptome involved in the chromatin
remodeling is compromised. How this will contribute to germ cell functions needs to be
investigated.

Sertoli cells are drivers of male sex determination and testicular morphogenesis [3,5]. A
previous study has shown that the lack of germ cells in the developing testis affects Sertoli
cell numbers and functions [23]. However, in the present study, we observed that loss of *Lhx2*
does not compromise the total Sertoli cell numbers nor alters the expression of key Sertoli
cell genes. To our surprise, we noticed that loss of *Lhx2* in the germ cells alters the
expression of 16% of Sertoli cell-enriched genes that had functions in cell-cell adhesion and
cell differentiation Indeed we did observe disruptions in the alignment of Sertoli cells of the
developing testicular tubules resulting indefective tubulogenesis. These observations imply
the involvement of intricate communication between the germ cells to the Sertoli cells in the
developing testis.

Angiogenesis is one of the biological processes enriched in the transcriptome of *Lhx2$^{flox/flox}$*
testis. These results prompted us to investigate if angiogenesis was also disrupted in the testis
of *Lhx2* knockout embryos. Indeed we observed increased numbers of VE-Cadherin positive
endothelial cells in the testis of *Lhx2-/-* embryos, these cells were often found in clusters and

stuck in the interstitial space suggestive of disrupted vasculature. Previous studies have implicated the involvement of WNT-βcatenin pathways for angiogenesis both in the testis and ovaries [44,45,55–57]. However, we did not observe altered expression of *Wnt4*, *Rspo1*, and *β-catenin* in the testis of the *Lhx2-/-* embryos implying that the disrupted vasculature in the *Lhx2-/-* testis is not due to the altered WNT/β-catenin signaling. Instead, we observed that the testicular transcriptome in absence of *Lhx2* had a heightened pro-angiogenic response as there was an elevated expression of multiple angiogenesis-promoting factors and reduced expression of anti-angiogenic factors. This altered ratio of the angiogenic group of molecules disrupted the path of the migrating endothelial cells in the developing testis as evident from the gonad recombination assays. Consequently, the coelomic blood vessel was not correctly developed in the mutant testis. A similar alteration in the ratio of pro and anti-angiogenic factors and excessive endothelial cell migration was observed in the developing ovaries of *Lhx2-/-* embryos [26]. These findings imply a role of germ cell-enriched *Lhx2* in the control of endothelial cell migration in the developing testis and the ovary. In a broader sense, the results from the present study and that published earlier in the ovary [26] prompt us to propose that germ cells control the endothelial cell migration and vascularization of the developing gonads of both the sexes.

The interstitial cells of the developing testis differentiate into peritubular cells, vascular smooth muscle cells, and steroidogenic Leydig cells [58]. Absence of *Lhx2* in the germ cells altered the expression of 22% of genes enriched in the interstitial cells which are involved in cell proliferation and migration. Indeed the proliferation of coelomic epithelium was significantly increased upon loss of *Lhx2* and this expansion seems to be of the undifferentiated precursors as there was increased expression of early interstitial markers *Lhx9* and *Nr5a1* in mutant testis and the expression of the Leydig cell-enriched genes was reduced. These results imply that the germ cells control the proliferation and differentiation of the interstitial precursors in the developing testis. Whether this is due to a direct effect of the germ cell secretions or indirectly via modulation of the secretome of other cell types in the testis needs to be investigated.

In the developing testis, the organization of the tubules is governed by Sertoli cells, the correct migratory path of the endothelial cells, and the proportion of the cells in the interstitium. Since all the major cell types were affected by the loss of *Lhx2*, we investigated the tubular organization in the developing testis. To our surprise, we found that the *Lhx2-/-* testis appeared more elongated and the tubules were not appropriately organized. Also, the

total numbers of tubules were higher in the *Lhx2* knockout testis; these tubules were abnormal as the numbers of SOX9 positive Sertoli cells were fewer and mis-aligned in most of the tubules. Consequently, some tubules were fused and a proportion of tubules were without the layer of peritubular myoid cells. Several tubules were also devoid of germ cells in the *Lhx2* knockout testis. In the developing testis, there is a male-specific pattern of deposition of the extracellular matrix, to ensure tubular organization. The deposition of this male-specific extracellular matrix deposition was thought to be governed by the coordinated function of Sertoli, endothelial, and interstitial cells [3,8,10,58–60]. Herein we observed that in the absence of *Lhx2* in germ cells, genes having functions in cell-cell adhesion, integrin-mediated signaling, and cell migration were differentially expressed in these three compartments and there was a disrupted organization of the basement membrane as judged by Laminin staining. These results for the first time provide conclusive evidence that germ cells govern the transcriptome of somatic cells for the organization of tubules in the developing testis. It will be of interest to study the *Lhx2*-regulated germ cell secretome to determine the mechanistic basis of this phenomenon.

To conclude, the present study has identified a crucial role of germ cells through a transcription factor, *Lhx2* in regulating the transcriptome of the supporting, vascular, and interstitial cells for governing the testicular tubule organization. The present study challenges the existing paradigm of the somatic cell-centric control of testis development and highlights an important role of germ cells in the process.

Funding and Acknowledgement

DM lab is funded by grants from the Indian Council of Medical Research (ICMR) and the study was funded by the Department of Biotechnology (DBT) Government of India under the Grant ID: BT/PR10368/MED/97/223/2014. NS is thankful to ICMR-Senior Research Fellowship. DD is grateful to UGC-Junior Research Fellowship and AB is thankful to DBT-project Junior Research Fellowship. RS is thankful to DST-INSPIRE for the Junior and Senior Research Fellowship. We thank Prof. Shubha Tole, TIFR for providing *Lhx2*-/- and *Lhx2* floxed animals and *Lhx2* riboprobe. We thank Dr. Reddy, ICMR-NIRRCH for providing *Oct4*-GFP & cKit mice and Dr. Swanand Koli for assisting with their genotyping. We also thank NII, New Delhi for providing FVB-GFP mice. The manuscript bears the NIRRCH ID: RA/1290/07-2022

Conflict of Interest/ Declaration of interest

The authors declare that there is no conflict of interest.

References

[1]   N. Singh, D. Modi, The Molecular Genetics of Testis Determination, in: Arafa M. et al. (Ed.), Genet. Male Infertil., Springer Cham, 2020: pp. 3–17. https://doi.org/doi.org/10.1007/978-3-030-37972-8_1.

[2]   J. Cool, T. Defalco, B. Capel, Testis formation in the fetal mouse: Dynamic and complex de novo tubulogenesis, Wiley Interdiscip. Rev. Dev. Biol. 1 (2012) 847–859. https://doi.org/10.1002/wdev.62.

[3]   T. Svingen, P. Koopman, Building the mammalian testis: Origins, differentiation, and assembly of the component cell populations, Genes Dev. 27 (2013) 2409–2426. https://doi.org/10.1101/gad.228080.113.

[4]   H.H.-C. Yao, E. Ungewitter, H. Franco, B. Capel, Establishment of fetal Sertoli cells and their role in testis morphogenesis, Elsevier Inc., 2015. https://doi.org/10.1016/b978-0-12-417047-6.00002-8.

[5]   S.P. Windley, D. Wilhelm, Signaling Pathways Involved in Mammalian Sex Determination and Gonad Development, Sex. Dev. 9 (2016) 297–315. https://doi.org/10.1159/000444065.

[6]   C. Tilmann, B. Capel, Mesonephric cell migration induces testis cord formation and Sertoli cell differentiation in the mammalian gonad, Development. 126 (1999) 2883–2890. https://doi.org/https://doi.org/10.1242/dev.126.13.2883.

[7]   M. Buehr, S. Gu, A. McLaren, Mesonephric contribution to testis differentiation in the fetal mouse, Development. 117 (1993) 273–281. https://doi.org/https://doi.org/10.1242/dev.117.1.273.

[8]   A.N. Combes, D. Wilhelm, T. Davidson, E. Dejana, V. Harley, A. Sinclair, P. Koopman, Endothelial cell migration directs testis cord formation, Dev. Biol. 326 (2009) 112–120. https://doi.org/10.1016/j.ydbio.2008.10.040.

[9] T. DeFalco, I. Bhattacharya, A. V. Williams, D.M. Sams, B. Capel, Yolk-sac-derived
macrophages regulate fetal testis vascularization and morphogenesis, Proc. Natl. Acad.
Sci. U. S. A. 111 (2014) E2384–E2393. https://doi.org/10.1073/pnas.1400057111.

[10] L. O'Donnell, L.B. Smith, D. Rebourcet, Sertoli cells as key drivers of testis function,
Semin. Cell Dev. Biol. 121 (2022) 2–9. https://doi.org/10.1016/j.semcdb.2021.06.016.

[11] P. Fan, L. He, D. Pu, X. Lv, W. Zhou, Y. Sun, N. Hu, Testicular Sertoli cells influence
the proliferation and immunogenicity of co-cultured endothelial cells, Biochem.
Biophys.   Res.   Commun.   404   (2011)   829–833.
https://doi.org/10.1016/j.bbrc.2010.12.068.

[12] D. Rebourcet, P.J. O'Shaughnessy, L.B. Smith, The expanded roles of Sertoli cells:
lessons from Sertoli cell ablation models, Curr. Opin. Endocr. Metab. Res. 6 (2019)
42–48. https://doi.org/10.1016/j.coemr.2019.04.003.

[13] M. Uzumcu, S.D. Westfall, K.A. Dirks, M.K. Skinner, Embryonic testis cord
formation and mesonephric cell migration requires the phosphotidylinositol 3-kinase
signaling   pathway,   Biol.   Reprod.   67   (2002)   1927–1935.
https://doi.org/10.1095/biolreprod.102.006254.

[14] C.A. Smith, P.J. McClive, Q. Hudson, A.H. Sinclair, Male-specific cell migration into
the developing gonad is a conserved process involving PDGF signalling, Dev. Biol.
284 (2005) 337–350. https://doi.org/10.1016/j.ydbio.2005.05.030.

[15] J. Brennan, C. Tilmann, B. Capel, Pdgfr-α mediates testis cord organization and fetal
Leydig cell development in the XY gonad, Genes Dev. 17 (2003) 800–810.
https://doi.org/10.1101/gad.1052503.

[16] J. Cool, T.J. DeFalco, B. Capel, Vascular-mesenchymal cross-talk through Vegf and
Pdgf drives organ patterning, Proc. Natl. Acad. Sci. U. S. A. 108 (2011) 167–172.
https://doi.org/10.1073/pnas.1010299108.

[17] K.M. Sargent, R.M. McFee, R.S. Gomes, A.S. Cupp, Vascular endothelial growth
factor A: Just one of multiple mechanisms for sex-specific vascular development
within the testis?, J. Endocrinol. 227 (2015) R31–R50. https://doi.org/10.1530/JOE-
15-0342.

[18] R.C. Bott, R.M. McFee, D.T. Clopton, C. Toombs, A.S. Cupp, Vascular endothelial
growth factor and kinase domain region receptor are involved in both seminiferous
cord formation and vascular development during testis morphogenesis in the rat, Biol.
Reprod. 75 (2006) 56–67. https://doi.org/10.1095/biolreprod.105.047225.

[19] D.R. Archambeault, H.H.C. Yao, Activin A, a product of fetal Leydig cells, is a unique
paracrine regulator of Sertoli cell proliferation and fetal testis cord expansion, Proc.
Natl. Acad. Sci. U. S. A. 107 (2010) 10526–10531.
https://doi.org/10.1073/pnas.1000318107.

[20] A.M. Clark, K.K. Garland, L.D. Russell, Desert hedgehog (Dhh) gene is required in
the mouse testis for formation of adult-type Leydig cells and normal development of
peritubular cells and seminiferous tubules, Biol. Reprod. 63 (2000) 1825–1838.
https://doi.org/10.1095/biolreprod63.6.1825.

[21] F. Pierucci-Alves, A.M. Clark, L.D. Russell, A developmental study of the desert
hedgehog-null mouse testis, Biol. Reprod. 65 (2001) 1392–1402.
https://doi.org/10.1095/biolreprod65.5.1392.

[22] J.J. Meeks, S.E. Crawford, T.A. Russell, K.I. Morohashi, J. Weiss, J.L. Jameson, Dax1
regulates testis cord organization during gonadal differentiation, Development. 130
(2003) 1029–1036. https://doi.org/10.1242/dev.00316.

[23] C. Rios-Rojas, C. Spiller, J. Bowles, P. Koopman, Germ cells influence cord formation
and leydig cell gene expression during mouse testis development, Dev. Dyn. 245
(2016) 433–444. https://doi.org/10.1002/dvdy.24371.

[24] S.J. Chou, S. Tole, Lhx2, an evolutionarily conserved, multifunctional regulator of
forebrain development, Brain Res. 1705 (2019) 1–14.
https://doi.org/10.1016/j.brainres.2018.02.046.

[25] I. Tzchori, T.F. Day, P.J. Carolan, Y. Zhao, C.A. Wassif, L.Q. Li, M. Lewandoski, M.
Gorivodsky, P.E. Love, F.D. Porter, H. Westphal, Y. Yang, LIM homeobox
transcription factors integrate signaling events that control three-dimensional limb
patterning and growth, Development. 136 (2009) 1375–1385.
https://doi.org/10.1242/dev.026476.

[26]   N. Singh, D. Singh, A. Bhide, R. Sharma, S. Sahoo, M.K. Jolly, D. Modi, Lhx2 in
793   germ cells suppresses endothelial cell migration in the developing ovary, Exp. Cell
794   Res. 415 (2022) 113108. https://doi.org/10.1016/J.YEXCR.2022.113108.

795   [27]   N. Singh, D. Singh, D. Modi, LIM Homeodomain (LIM-HD) Genes and Their Co-
796   Regulators in Developing Reproductive System and Disorders of Sex Development,
797   Sex. Dev. (2021) 1–15. https://doi.org/10.1159/000518323.

798   [28]   T. Yoshimizu, N. Sugiyama, M. De Felice, Y. Yeom, K. Ohbo, K. Masuko, M.
799   Obinata, A. Kuniya, H.R. Schöler, Y. Matsui, Germline-specific expression of the Oct-
800   4/green fluorescent protein (GFP) transgene in mice, Dev. Growth Differ. 41 (1999)
801   675–684. https://doi.org/10.1046/j.1440-169X.1999.00474.x.

802   [29]   C. Waskow, S. Paul, C. Haller, M. Gassmann, H.R. Rodewald, Viable c-KitW/W
803   mutants reveal pivotal role for c-Kit in the maintenance of lymphopoiesis, Immunity.
804   17 (2002) 277–288. https://doi.org/10.1016/S1074-7613(02)00386-2.

805   [30]   F.D. Porter, J. Drago, Y. Xu, S.S. Cheema, C. Wassif, S.P. Huang, E. Lee, A.
806   Grinberg, J.S. Massalas, D. Bodine, F. Alt, H. Westphal, Lhx2, a LIM homeobox gene,
807   is required for eye, forebrain, and definitive erythrocyte development, Development.
808   124 (1997) 2935–2944. https://doi.org/https://doi.org/10.1242/dev.124.15.2935.

809   [31]   V.S. Mangale, K.E. Hirokawa, P.R.V. Satyaki, N. Gokulchandran, S. Chikbire, L.
810   Subramanian, A.S. Shetty, B. Martynoga, J. Paul, M. V. Mai, Y. Li, L.A. Flanagan, S.
811   Tole, E.S. Monuki, Lhx2 selector activity specifies cortical identity and suppresses
812   hippocampal   organizer   fate,   Science   (80-.   ).   319   (2008)   304–309.
813   https://doi.org/10.1126/science.1151695.

814   [32]   G. Godbole, D. Modi, Regulation of decidualization, interleukin-11 and interleukin-15
815   by homeobox A 10 in endometrial stromal cells, J. Reprod. Immunol. 85 (2010) 130–
816   139. https://doi.org/10.1016/j.jri.2010.03.003.

817   [33]   D. Modi, S. Sane, D. Bhartiya, Over-expression of müllerian inhibiting substance
818   mRNA   in   the   turner   syndrome   ovary,   Sex.   Dev.   3   (2009)   245–252.
819   https://doi.org/10.1159/000261659.

820   [34]   G. Godbole, A. Roy, A.S. Shetty, S. Tole, Novel functions of LHX2 and PAX6 in the

developing telencephalon revealed upon combined loss of both genes, Neural Dev. 12 (2017) 1–8. https://doi.org/10.1186/s13064-017-0097-y.

[35] S. Bulchand, E.A. Grove, F.D. Porter, S. Tole, LIM-homeodomain gene Lhx2 regulates the formation of the cortical hem, Mech. Dev. 100 (2001) 165–175. https://doi.org/10.1016/S0925-4773(00)00515-3.

[36] L. Wojtasz, K. Daniel, A. Toth, Fluorescence activated cell sorting of live female germ cells and somatic cells of the mouse fetal gonad based on forward and side scattering, Cytom. Part A. 75 (2009) 547–553. https://doi.org/10.1002/cyto.a.20729.

[37] A. Mishra, M. Galvankar, N. Singh, D. Modi, Spatial and temporal changes in the expression of steroid hormone receptors in mouse model of endometriosis, J. Assist. Reprod. Genet. 37 (2020) 1069–1081. https://doi.org/10.1007/s10815-020-01725-6.

[38] A. Mishra, M. Galvankar, S. Vaidya, U. Chaudhari, D. Modi, Mouse model for endometriosis is characterized by proliferation and inflammation but not epithelial-to-mesenchymal transition and fibrosis, J. Biosci. 45 (2020) 105. https://doi.org/10.1007/s12038-020-00073-y.

[39] A. Tiwari, N. Ashary, N. Singh, S. Sharma, D. Modi, Modulation of E-Cadherin and N-Cadherin by ovarian steroids and embryonic stimuli, Tissue Cell. 73 (2021) 101670. https://doi.org/10.1016/j.tice.2021.101670.

[40] S.A. Jameson, A. Natarajan, J. Cool, T. DeFalco, D.M. Maatouk, L. Mork, S.C. Munger, B. Capel, Temporal transcriptional profiling of somatic and germ cells reveals biased lineage priming of sexual fate in the fetal mouse gonad, PLoS Genet. 8 (2012) e1002575. https://doi.org/10.1371/journal.pgen.1002575.

[41] R. Saba, Y. Kato, Y. Saga, NANOS2 promotes male germ cell development independent of meiosis suppression, Dev. Biol. 385 (2014) 32–40. https://doi.org/10.1016/j.ydbio.2013.10.018.

[42] D. Coveney, J. Cool, T. Oliver, B. Capel, Four-dimensional analysis of vascularization during primary development of an organ, the gonad, Proc. Natl. Acad. Sci. U. S. A. 105 (2008) 7212–7217. https://doi.org/10.1073/pnas.0707674105.

[43]   and P.K. N, Combes AAlexander N. Combes1, Emmanuelle Lesieur1, Vincent R.
Harley2, Andrew H. Sinclair3, Melissa H. Little1, Dagmar Wilhelm1, Three-
dimensional visualisation of testis cord morphogenesis, a novel tubulogenic
mechanism in development, Dev Dyn. 238 (2009) 1033–1041.
https://doi.org/10.1002/dvdy.21925.Three-dimensional.

[44] A.A. Chassot, I. Gillot, M.C. Chaboissier, R-spondin1, WNT4, and the ctnnb1
signaling pathway: Strict control over ovarian differentiation, Reproduction. 148
(2014) R97–R110. https://doi.org/10.1530/REP-14-0177.

[45] B.K. Jordan, J.H.C. Shen, R. Olaso, H.A. Ingraham, E. Vilain, Wnt4 overexpression
disrupts normal testicular vasculature and inhibits testosterone synthesis by repressing
steroidogenic factor 1/β-catenin synergy, Proc. Natl. Acad. Sci. U. S. A. 100 (2003)
10866–10871. https://doi.org/10.1073/pnas.1834480100.

[46] L. Garcia-Alonso, V. Lorenzi, C.I. Mazzeo, J.P. Alves-Lopes, K. Roberts, C. Sancho-
Serra, J. Engelbert, M. Marečková, W.H. Gruhn, Rachel A. Botting, T. Li, B. Crespo,
Stijn van Dongen, V.Y. Kiselev, E. Prigmore, M. Herbert, A. Moffett, A. Chédotal,
O.A. Bayraktar, A. Surani, M. Haniffa, R. Vento-Tormo, Gonadal, Single-cell
roadmap of human gonadal development, Nature. 607 (2022) 540–547.
https://doi.org/10.1038/s41586-022-04918-4.

[47] V. Kinare, A. Iyer, H. Padmanabhan, G. Godbole, T. Khan, Z. Khatri, U. Maheshwari,
B. Muralidharan, S. Tole, An evolutionarily conserved Lhx2-Ldb1 interaction
regulates the acquisition of hippocampal cell fate and regional identity, Development.
147 (2020) 1–6. https://doi.org/10.1242/dev.187856.

[48] V. Lakhina, L. Subramanian, D. Huilgol, A.S. Shetty, V.A. Vaidya, S. Tole, Seizure
evoked regulation of LIM-HD genes and co-factors in the postnatal and adult
hippocampus, F1000Research. 2 (2013) 1–12.
https://doi.org/10.12688/f1000research.2-205.v1.

[49] J. De Melo, B.S. Clark, A. Venkataraman, F. Shiau, C. Zibetti, S. Blackshaw, Ldb1-
and Rnf12-dependent regulation of Lhx2 controls the relative balance between
neurogenesis and gliogenesis in the retina, Development. 145 (2018) dev159970.
https://doi.org/10.1242/dev.159970.

[50]   D. Peukert, S. Weber, A. Lumsden, S. Scholpp, Lhx2 and Lhx9 determine neuronal
880   differentiation and compartition in the caudal forebrain by regulating Wnt signaling,
881   PLoS Biol. 9 (2011) e1001218. https://doi.org/10.1371/journal.pbio.1001218.

882   [51]   O.S. Birk, D.E. Caslano, C.A. Wassif, T. Cogilatl, L. Zhaos, Y. Zhao, A. Grinberg,
883   S.P. Huang, J.A. Kreidberg, K.L. Parker, F.D. Porter, H. Westphal, The LIM
884   homeobox gene Lhx9 is essential for mouse gonad formation, Nature. 403 (2000) 909–
885   913. https://doi.org/10.1038/35002622.

886   [52]   R.F. Hevner, Evolution of the mammalian dentate gyrus, J. Comp. Neurol. 524 (2016)
887   578–594. https://doi.org/10.1002/cne.23851.

888   [53]   X. Li, X. Wu, H. Chen, Z. Liu, H. He, L. Wang, LHX2 Enhances the Malignant
889   Phenotype of Esophageal Squamous Cell Carcinoma by Upregulating the Expression
890   of   SERPINE2,   Genes   (Basel).   13   (2022)   1457–1469.
891   https://doi.org/10.3390/genes13081457.

892   [54]   S.J. Chou, D.D.M. O'Leary, Role for Lhx2 in corticogenesis through regulation of
893   progenitor   differentiation,   Mol.   Cell.   Neurosci.   56   (2013)   1–9.
894   https://doi.org/10.1016/j.mcn.2013.02.006.

895   [55]   S.G. Tevosian, N.L. Manuylov, To β or not to β: Canonical β-catenin signaling
896   pathway   and   ovarian   development,   Dev.   Dyn.   237   (2008)   3672–3680.
897   https://doi.org/10.1002/DVDY.21784.

898   [56]   K. Jeays-Ward, M. Dandonneau, A. Swain, Wnt4 is required for proper male as well as
899   female   sexual   development,   Dev.   Biol.   276   (2004)   431–440.
900   https://doi.org/10.1016/j.ydbio.2004.08.049.

901   [57]   A. Heinrich, T. DeFalco, Essential roles of interstitial cells in testicular development
902   and function, Andrology. 8 (2020) 903–914. https://doi.org/10.1111/andr.12703.

903   [58]   A. Heinrich, T. Defalco, Essential roles of interstitial cells in testicular development
904   and function, Andrology. 8 (2020) 903–914. https://doi.org/10.1111/andr.12703.

905   [59]   P.S. Tung, M.K. Skinner, I.B. Fritz, Cooperativity between Sertoli Cells and
906   Peritubular Myoid Cells in the Formation of the Basal Lamina in the Seminiferous

907   Tubule, Ann. N. Y. Acad. Sci. 438 (1984) 435–446. https://doi.org/10.1111/j.1749-
908   6632.1984.tb38304.x.

909   [60] S.R. Chen, Y.X. Liu, Testis cord maintenance in mouse embryos: Genes and signaling,
910   Biol. Reprod. 94 (2016) 1–7. https://doi.org/10.1095/biolreprod.115.137117.

Author Statement

Neha Singh: Conceptualization, Investigation, Formal analysis, Data Curation, Writing original draft, Reviewing and Editing. **Domdatt Singh:** Data Curation, Resources, Reviewing and Editing. **Anshul Bhide:** Investigation, Reviewing and Editing. **Richa Sharma:** Investigations. **Shilpa Bhowmick:** Investigation, Reviewing and Editing. **Vainav Patel:** Resources, Reviewing and Editing. **Deepak Modi:** Conceptualization, Reviewing and Editing, Supervision and Funding acquisition.