Common core respiratory bacteriome of the blue whale *Balaenoptera musculus*, in the Gulf

of California

Carlos A. Domínguez-Sánchez[1], Roberto C. Álvarez-Martínez[1], Diane Gendron[3], Karina

Acevedo-Whitehouse[4,*]

[1] Unit for Basic and Applied Microbiology. School of Natural Sciences. Autonomous University of Queretaro,

Queretaro, Mexico.

[2] Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, La Paz, Baja California Sur, Mexico.

[*] Address correspondence to: Karina Acevedo-Whitehouse, karina.acevedo.whitehouse@uaq.mx

Abstract

The number of strandings and unusual mortality events that involve cetaceans may have

increased, and potential pathogens of the respiratory tract have been found during the

examination of individuals in many of these events. However, investigating the health of free-

ranging large whales is logistically complex. Given that the core microbiome is key to

understanding host-bacteria relationships and to identifying their relevance for individual and

population health, we characterized the core respiratory bacteriome of the Eastern North Pacific

blue whale, *Balaenoptera musculus*, using blow samples collected by a small quadracopter

drone. 16S rRNA gene high-throughput sequencing revealed 1,326 amplicon sequence variants

(ASVs), of which 11 were shared by more than 50% of all blue whales and had a relative

abundance higher than 0.02%. *Cutibacterium, Oceanivirga, Tenacibaculum*, and *Psychrobacter*

composed the common core respiratory bacteriome of the blue whale. Additionally,

compositional analysis identified 15 bacterial classes dominated by Gammaproteobacteria

(27.14%), Bacteroidea (19.83%), and Clostridia (12.89%) as the most representative classes in

the respiratory tract of blue whales. However, two whales had a high abundance of bacteria with

pathogenic potential, namely M*ycoplasma* spp. and *Streptococcus* spp. in their blow. Both

bacterial genera have been associated with pulmonary diseases in mammals. Ours is the first

study to characterize the respiratory bacteriome of apparently healthy blue whales and is a

baseline for future long-term studies on blue whale health, an endangered species of conservation

concern.

Keywords: microbiome, blue whale, *Balaenoptera musculus,* respiratory tract, bacteria, 16S rRNA gene

Introduction

The advent of modern technologies that allow the identification of all bacteria present in environmental or clinical samples (Haegeman et al., 2013; Salter et al., 2014; Rhodes et al., 2022) has led to a myriad of studies on the abundance, diversity, and structure of the microbiome of different species (Nelson et al., 2015; Watkins et al., 2022). One of the reasons why it is paramount to increase our knowledge about the microbiome of a given species is because the microbial communities associated with a particular epithelium can impact the host's physiology (Foster et al., 2017), and even influence its health status (Huang, Yang, & Li, 2016; Bierlich et al., 2018; Panthee et al., 2022; Watkins et al., 2022). For example, respiratory infections can occur when opportunistic microorganisms, which are normally part of the microbiome of a healthy respiratory mucosa, preferentially flourish under certain conditions (Dickson et al., 2016; Huffnagle, Dickson, & Lukacs, 2017; Rhodes et al., 2022), and *de novo* infections may occur if individuals are exposed to pathogens. In turn, infections can trigger changes in the diversity and composition of the original microbial communities, an event known as dysbiosis (Gagliardi et al., 2018; Robles-Malagamba et al., 2020; Sehnal et al., 2021). As a result, the composition of the microbiome may be a better predictor of disease progression than the presence of a specific pathogen commonly associated with the disease. Thus, understanding the composition of the microbiome and how it differs between healthy and sick animals could become an important tool for assessing an individual's health status (Shreiner, Kao, & Young, 2015). Such knowledge, in turn, is paramount for animal conservation efforts, as the impact of diseases on wild animal populations is increasingly becoming evident (Hu, Han, & He, 2023). This has led to a recent surge in studies that have begun to examine the microbiome of species of conservation concern (e.g. Dallas & Warne, 2023; Jiang et al., 2022; Vecchioni et al., 2022).

When attempting to use the microbiome to assess health status, commensal, opportunistic, and transient bacteria must be identified (Huang, Yang, & Li, 2016; Infante-Villamil, Huerlimann, & Jerry, 2021). To do this, it is necessary to identify the bacterial taxa that predominate in the community and are shared by healthy individuals (Huse et al., 2012; Willis, Bunge, & Whitman,

2017; Risely, 2020). This is known as the core microbiome and is well known to play an
important role in maintaining functional stability and homeostasis of a specific habitat within the
host (Hernandez-Agreda, Gates, & Ainsworth, 2017; Thomas et al., 2017; Björk et al., 2018;
Ross, Rodrigues-Hoffmann, & Neufeld, 2019). The definition of the core microbiome varies
across authors, although they tend to overlap in many of the components of the microbial
community (Risely, 2020). Different approaches to define the core microbiome have included
temporal stability (Ozkan et al., 2017), functional level (Dinsdale et al., 2008), ecological
influence (Coyte et al., 2019), host fitness (Shapira, 2016), and bacterial occupancy frequency,
which refers to the most common bacterial taxa that are shared by a considerable proportion of
hosts (Ingala et al., 2018; Nishida & Ochman, 2018; Risely, 2020; Toro et al., 2021).

Identifying the common core bacteriome (i.e. the bacterial community of the microbiome)
requires setting the detection threshold (relative abundance) and the percentage of occurrence
(prevalence) of bacterial taxa (Astudillo-García et al., 2017), criteria which have varied widely
among published studies (Risely, 2020), with common core annotations ranging from as low as
30% (e.g., Ainsworth et al., 2015) to 100% prevalence (e.g. (Huse et al., 2012; Apprill et al.,
2017; Hernandez-Agreda, Gates, & Ainsworth, 2017; Antwis et al., 2018), and detection
thresholds varying from 0.001% to 0.1% (Astudillo-García et al., 2017; Antwis et al., 2018).
Given that biological justifications for such prevalence and threshold values are rare (Risely,
2020), it is important to recognize the arbitrary aspect of the common core definition and to be
cautious when interpreting the results. However, the bacteriome common core tends to be robust
despite varying definitions, particularly when samples from closely related individuals are
analysed (Risely, 2020; Vendl et al., 2020).

Bacteria of the mammalian microbiome are found in composite communities (Lee &
Mazmanian, 2010; Rhodes et al., 2022), whose diversity and abundance are determined by
multiple interactions between species (Shade & Handelsman, 2012; Stubbendieck, Vargas-
Bautista, & Straight, 2016). The cetacean microbiome has recently begun to be studied and
important initial assessments of microbial diversity have been made for a few species (Johnson et
al., 2009a; Hooper et al., 2019; Apprill et al., 2020; Van Cise et al., 2020; Toro et al., 2021;
Rhodes et al., 2022). As long-lived cetaceans that contribute to the movement and storage of

carbon in the ocean, baleen whales play a vital role in the marine ecosystem (Pershing et al.,
2010), and are considered sentinels of ocean health (Sharp et al., 2019; Palmer et al., 2022).
However, to date, little is known about the respiratory microbiome of baleen whales, and while
some opportunistic pathogens of the respiratory tract have been described for live free-ranging
individuals of a few baleen whale species (Acevedo-Whitehouse et al., 2010), to the best of our
knowledge, there is only one published study on the core respiratory microbiome of a baleen
whale, the humpback whale *Megaptera novaeangliae* (Apprill et al., 2017).

Blue whales are among the world's largest and most iconic animals and play an essential role in
marine ecosystems (Attard, Beheregaray, & Möller, 2016). Owing to their long migration and
diverse habitats, they are also important indicators of the health of oceans worldwide (Rice et al.,
2022). Studies on the respiratory microbiome of blue whales are crucial for understanding their
health and well-being in their natural environment because the bacteria in their respiratory
system can provide valuable information about their exposure to pathogens (Raverty et al.,
2017). This can aid in identifying potential diseases and developing effective conservation
strategies for marine mammals (Raverty et al., 2020), especially considering global and local
environmental changes affecting oceans (Ferguson et al., 2021; Aguilar & Borrell, 2022; Barlow
et al., 2023). Therefore, studying the respiratory microbiome is key to conserving this iconic
species and protecting its long-term survival. In this study, we characterized the common core
respiratory bacteriome of the Eastern North Pacific blue whale using next-generation sequencing
on 17 blow samples collected from adult blue whales by a non-invasive drone-based technique
(Domínguez-Sánchez et al., 2018) during the boreal winter months in the Gulf of California.

Materials and Methods
Sample collection. Using a small Phantom 3 quadracopter drone (DJI Innovations, China) with
floaters and sterile Petri dishes, we collected 17 blow samples from 17 individual blue whales
sampled between February and March 2016 and 2017 in Loreto Bay National Park
(25°51'51''N, 111°07'18''O) within the Gulf of California, Mexico. The number of sampled
whales represents 17% of the estimated 100 blue whales that reside during winter/spring in the
southwestern Gulf of California (mark-recapture data from 1994-2006; Gendron & Ugalde de la
Cruz, 2012; SEMARNAT, 2018). Each whale was photo-identified prior to sample collection

(Gendron & Ugalde de la Cruz, 2012). The approach to the whale with the drone was made from

the caudal fin heading towards the head to minimize disturbance, and sampling was conducted at

a height between 3 and 4 m above the blowhole (Domínguez-Sánchez et al., 2018).

For each sample, the blow droplets were swabbed directly from the Petri dish using sterile

cotton-tipped swabs. These were then transferred to a sterile 1.5 mL cryogenic microtube

containing 500 µL of 96% molecular grade ethanol and kept frozen in a liquid nitrogen container

until processing. To address potential contamination, all necessary precautions were taken,

always including the use of sterile gloves and face masks during sample handling and

processing. In addition, we included three technical controls:1) LabControl (DNA extraction

from kit reagents), 2) HumanSneeze (human sneeze sampled from the person who collected and

processed the samples), and 3) SeaWater (seawater collected at a depth of 0.10 m in the exact

location where we sampled the whale blows).

DNA extraction, PCR amplification and sequencing. Total DNA was isolated from the whale

blow and technical control samples in one batch using a QIAamp DNA Mini Kit (QIAGEN,

Germany). The primers used for sequencing the 16SrRNA V3 and V4 regions were 341F (5'-

CCTACGGGNGGCWGCAG) and 785R (5' -GACTACHVGGGTATCTAATCC), which

amplified a single product of approximately 460 bp (Thijs et al., 2017). The PCR program used

an initial denaturation step at 95°C for 3 min; 25 cycles of 95°C for 30 s, 55°C for 30 s, and

72°C for 30s; and a final extension step at 72°C for 5 min. Each 25 µL-reaction contained 12.5

ng of extracted DNA, 5 µM of barcoded primers and 2x KAPA HiFi HotStart Ready Mix

(KAPABIOSYSTEM, Cape Town, South Africa). 1 µl of each sample was run on a 2100

Bioanalyzer (Agilent Technologies, CA, USA) with an Agilent DNA 1000 chip (Agilent

Technologies, CA, USA) to verify amplicon size. AMPure XP beads (New England BioLabs,

USA) were used to remove unused primers and primer dimers. Amplicons were sequenced over

2- by 250-bp MiSeq at the Unit of Sequencing and Identification of Polymorphisms of the

National Institute of Genomic Medicine (Instituto Nacional de Medicina Genómica, Unidad de

Secuenciación e Identificación de Polimorfismos) in Mexico.

16S rRNA sequence data processing. Quality control overview was performed using FASTQC (Andrews, 2010). This allowed us to obtain a quick impression of the data and avoid downstream problems. The raw sequences were then imported into R v.4.2.1 (R Core Team, 2022), where all subsequent analyses were carried out. We used the Divisive Amplicon Denoising Algorithm 2 (*dada2*) v.1.26.0 (Callahan et al., 2016) to infer exact amplicon sequence variants (ASVs) instead of the rough and less precise 16S rRNA OTU clustering approach (Callahan, McMurdie, & Holmes, 2017; Dahan et al., 2018) that groups the sequences with a 97% identity (Edgar, 2018). First, we filtered by quality (trunQ = 25) and discarded the sequences if they presented more than two Ns (maxN = 2) or more than two expected errors (maxEE = 2). Next, the forward and reverse reads for each sample were combined into a single merged contig sequence, and we grouped all identical reads into unique sequences to determine their abundance. After building the ASVs table and removing chimeras (detected using self-referencing), sequences were classified and identified with *Decipher* v.2.26.0 (Wright, 2016), using the SILVA rRNA sequence database v.138 as the taxa reference (Quast et al., 2013). We used *phyloseq* v.1.42.0 (McMurdie & Holmes, 2013) to classify and remove any sequence not classified at the kingdom level or belonging to Archaea, Eukaria, chloroplasts, or mitochondria. For the last steps of the sequence cleaning process, we used *metagMisc* v.0.5.0 (Mikryukov, 2023) to eliminate ASVs with less than ten reads (minabund = 10). Additionally, we used *SourceTracker* (Knights et al., 2013) to estimate the proportion of the bacterial community in the blue whales blows samples that comes from the set of technical controls. ASVs detected in the LabControl sample were considered potential contaminants if their mean abundance was equivalent to or greater than 25% of their mean abundance in the whale samples (Caruso et al., 2019; Minich et al., 2019). Finally, we plotted Venn diagrams to show the logical relation of ASVs between blow and technical control samples using *MicEco* version 0.9.19 (Russel, 2023).

Respiratory microbiome analysis. To get a sense of the community composition in the samples, we used *phyloseq* to identify the distribution of read counts from all the samples, and to plot rarefaction curves and the relative abundance stacked bar plot at the major level class and genus. Using *microbiome* v.1.20. (Lahti et al., 2017), we identified the common core bacteriome (threshold detection = 2/100, prevalence = 50/100). We selected these values because we wanted a more conservative approach and did not want to consider "rare bacteria" in the analysis. In

addition, we analysed how the pattern of the common central microbiome changed based on a
sliding prevalence range (10% - 100%) and a detection threshold of 0.02%. Using these criteria,
we ran a linear regression model to determine the number of ASVs detected, given a detection
threshold value for a specific prevalence. We calculated alpha diversity indices: richness (S) and
Simpson´s diversity index (D) using *vegan* v.2.6.4 (Oksanen et al., 2019). All graphs were
rendered using Tableau (Murray, 2013).

Results

A total of 20 samples were collected and analysed in this study. These samples included 17
photo-identified blue whales and three technical controls. Exhaled breath samples were collected
from animals in the Gulf of California using a small drone. No adverse behaviour was detected
before, during, or after sampling (see Domínguez-Sánchez et al., 2018). After filtering,
denoising, merging, and chimera elimination (2.33% of the reads), we recovered 72,254
sequences corresponding to 1,367 amplicon sequence variants (ASVs). Because of their low
relative abundance, 677 of these ASVs were categorized as "others". Additionally, we eliminated
41 ASVs classified as Archea (n = 2), chloroplasts (n = 31), or mitochondria (n = 8), and 9 ASVs
that were considered putative contaminants from LabControl. The sample coverage (i.e., the
proportion of the total number of individuals in a community that belong to the species
represented in the sample; (Chao & Chiu, 2016) exceeded 97.6% in all cases.

The logical relation analysis (Fig. 1) showed a total of 651 ASVs, of which 489 (76%) were
identified solely in blow samples, 37 (6%) in SeaWater, 28 (4%) in HumanSneeze, and 11 (2%)
in the LabControl sample. Alpha diversity measures revealed that species richness (S) in blow
samples ranged from 70 to 440 (mean = 253.06) and Simpson's index (D) ranged from 0.90 to
0.99 (mean = 0.97). The Compositional analysis identified 15 bacterial classes in the blow and
technical control samples (Fig. 2), dominated by Gammaproteobacteria (27.14%), Bacteroidea
(19.83%), and Clostridia (12.89%) as the most representative classes in the respiratory tract of
blue whales. We also calculated the relative abundances of the 15 most representative bacterial
genera in the blow samples (Fig. 3). *Psychrobacter* spp. (8.01%), *Oceanovirga* spp. (6.47%),
Tenacibaculum spp. (5.92%), *Streptococcus* spp. (5.03%), and *Cutibacterium* spp. (4.49%) were
the most abundant. Interestingly, two whale blows (Bm057 and Bm044) had a high relative

abundance of opportunistic pathogens, *Mycoplasma* spp. (26.8%), and *Streptococcus* spp.
(69.35%), respectively.

Eleven ASVs were considered the common core bacteriome of the respiratory tract of blue
whales in the Gulf of California (Fig. 4a). The common core spanned nine classified ASVs in
four bacterial genera (*Cutibacterium, Oceanivirga, Tenacibaculum*, and *Psychrobacter*) and two
ASVs that were not classified at the genus level but belonged to the families Lachnospiraceae
and Arcobacteraceae. Additionally, analysing changes in the pattern of the common central
bacteriome, based on a range of prevalence and detection threshold values (Fig. 4b), it was
possible to identify that the common central bacteriome of blue whale blow can vary from 232
ASVs (10% prevalence and 0.02 detection threshold) to 1 ASV (70% prevalence and 0.02
detection threshold), in all cases revealing *Cutibacterium* as the genus with the highest
prevalence (70%) in blue whale blow samples. This bacterium was also identified in the
technical controls, with a mean relative abundance of 4.73%. *Herbaspirillum* spp., the most
abundant genus found in the seawater sample with 19.39% abundance, was also found in the
blow samples with an average abundance of 2.38% and in three samples with higher abundances:
Bm023 (8.21%), Bm043 (14.03%), and Bm059 (13.60%).

Discussion
Animal conservation can benefit from knowledge on disease and health status of threatened and
endangered species (Heard et al., 2013; Pedersen et al., 2007). One of the factors that influence
both disease and health of animals is the microbiome (Matthewman et al., 2023), and
conservation science has recently begun to include studies of the microbiome to increase our
understanding of population health (see Comizzoli & Power, 2019). Here, we examined the
respiratory microbiome of the Eastern North Pacific blue whale, the planet's largest extant
animal and currently an endangered species for which there is extremely limited knowledge
about their health and disease.

Recent studies of the human respiratory microbiome have shown that bacterial communities of
the respiratory tract are key to maintaining respiratory health (Bond et al., 2017; Glendinning,
McLachlan, & Vervelde, 2017; Watson, De Koff, & Bogaert, 2019), not only in terms of their

metabolic contribution (Van Treuren & Dodd, 2020), but also because they prevent the
colonization of the epithelium by environmental pathogens (Man, De Steenhuijsen Piters, &
Bogaert, 2017; Yamamoto et al., 2021). Identifying the composition and abundance of the
bacterial communities that constitute the microbiome of healthy individuals is an important step
to establishing a baseline that will help identify bacteria associated with respiratory diseases
(Lemon et al., 2010; N. Lima et al., 2012), assess chronic states of suboptimal health (Mackenzie
et al., 2017) and predict community changes due to perturbation (Shade & Handelsman, 2012;
Yamamoto et al., 2021)

The results of our study demonstrate that the blow of the Eastern North Pacific blue whale
supports a diverse and rich bacterial community. We identified 1,367 ASVs with high sample
coverage and varying levels of richness and relative abundance among samples. This could be an
indicator of temporary fluctuations in the composition of the microbiome (Eloe-fadrosh &
Rasko, 2013). The richness and relative abundance of the microbiome vary among healthy
animals (Coyte et al., 2019; Toro et al., 2021), due mainly to bacterial immigration from the
environment to the lungs during inhalation, bacterial elimination via mucociliary clearance, and a
relatively small contribution of the growth rate of each bacterial community (Dickson, Erb-
Downward, & Huffnagle, 2014; Dickson et al., 2016; Huffnagle, Dickson, & Lukacs, 2017).
Evidently, we cannot rule out that the variation in richness and relative abundance that we
observed was also related to differences in the volume of blow collected, which due to the nature
of the samples collection technique used, could not be standardized. On one hand, whales are
likely to differ in the amount of blow exhaled, depending on the whale's size, and the depth and
duration of the dive. Furthermore, although drones are safe, minimally invasive, and do not seem
to affect the whales during blow collection (Domínguez-Sánchez et al., 2018), inherent
limitations such as flight height and different wind conditions, could potentially result in
different volumes of blow being collected (Apprill et al., 2017).

Various indices have been used to estimate the diversity of microbial communities. In our study,
we used Simpson's diversity index because it considers both richness and evenness (Johnson &
Burnet, 2016) and was previously identified as one of the most accurate estimators of diversity in
an unknown bacterial community (Haegeman et al., 2014). In our study, Simpson's diversity

index averaged 0.97 (minimum = 0.90, maximum = 0.99), demonstrating high bacterial diversity
in the blow of the whales that we sampled. Some studies have shown that the microbiome of a
healthy animal tends to have high diversity, which presumably allows it to tolerate or counteract
changes arising from extrinsic challenges (Chan, Estaki, & Gibson, 2013; Van Cise et al., 2020).
Bacterial diversity was high in nearly all of the blue whale blow samples, and the composition of
the microbiome was dominated mainly by members of the phyla Proteobacteria, Firmicutes,
Bacteroidota, Actinobacteria, and Fusobacteriota, which have been reported to be major
components of the healthy respiratory microbiome of other mammals (Chaban et al., 2013;
Rhodes et al., 2022). Simpson´s diversity index was similar to that reported for humpback whale
blows (Apprill et al., 2017) and bottlenose dolphin blowholes (Johnson et al., 2009b; Bik et al.,
2016). It is likely that these results are evidence that the respiratory microbiome diversity (based
on Simpson´s index) is conserved among cetaceans. However, the taxonomic richness of the blue
whale blow was nearly twice that reported for humpback whales. In our samples, the percentage
of "others" (sum of bacteria that did not reach the detection threshold of 0.02%) was higher than
that in humpback whales (Apprill et al., 2017). Based on this result, we suggest that the
microbiome of the Eastern North Pacific blue whale respiratory tract may be more complex than
that of humpback whales. At this stage we can only speculate about the reasons that could
explain such a difference. They may be due to the use of DADA2 for resolving ASVs rather than
minimum entropy decomposition (MEDs; Eren et al., 2015), which has been used previously,
given that although similar in what both approaches resolve, ASVs are better at removing
erroneous sequences (Callahan et al., 2016; Ahlgren et al., 2019). As the algorithm DADA2
allows for the independent analysis or grouping of samples, we conducted pooled analyses to
increase the sensitivity to detect ASVs that could be present at very low frequencies in multiple
samples (Callahan et al., 2016). This approach could explain why a higher percentage (mean =
30.91%) of "rare bacterial biosphere" (Pedrós-Alió, 2012) was identified in our study than in the
humpback whale study (see (Apprill et al., 2017). This "rare bacterial biosphere", formed by
bacteria that are present at low relative abundances, is particularly important for dealing with
dysbiosis, as they could be considered as a seed bank of genetic resources that can lead to the
restoration of the core microbiome (Pedrós-Alió, 2012; Skopina et al., 2016; Jousset et al., 2017).
A more complex respiratory microbial community is likely to be beneficial to whales, given that
microbiomes with a higher richness of species have more synergistic interactions between

bacterial taxa, which improve the functioning of the ecosystem (Bell et al., 2005). It is worth
mentioning that most studies published to date do not consider the bacterial taxa found in low
abundances to be relevant; however, these small populations are now thought to play an
important role for the functioning of the ecosystem (Willis, Bunge, & Whitman, 2017) and host
health (Jousset et al., 2017). It has been demonstrated that a high diversity of low-abundance
bacteria is correlated with less severe bacterial infections in human lungs (Van Der Gast et al.,
2010).

Sixteen bacterial classes were identified in the blow samples, some of which were also found in
the seawater samples. This is unsurprising, as there is likely to be some seawater carried over
when whales exhale. However, the blow samples harboured ASVs belonging to bacterial genera
that were not found in the seawater sample, such as *Psychrobacter, Oceanivirga,*
Tenacibaculum, Helcococcus, Porphyromonas, Mycoplasma, Dielma, Synechococcus, and
Suttonella. This shows that despite the potential carryover of seawater to the blow during
exhalation, the bacterial communities of the blue whale respiratory tract are different from those
of seawater.

Eleven   ASVs   belonging   to   the   families   Propionibacteriacea,   Flavobacteriaceae,
Lachnospiraceae, Leptotrichiaceae, Moraxellaceae, Porphyromonadaceae, and Arcobacteraceae;
were shared in more than half of the blue whale samples and were considered to be the common
core respiratory bacteriome of the Eastern North Pacific blue whale. The interindividual
variability of the respiratory microbiome appears to be higher than that of the humpback whales,
as 25 distinct bacteria were found to be shared among all the animals sampled (Bierlich et al.,
2018). *Psychrobacter* spp. and *Tenacibaculum* spp. have been reported to be key members of the
core microbiome in other baleen whales (Hooper, Littman, & Macpherson, 2012; Bierlich et al.,
2018; Van Cise et al., 2020; Toro et al., 2021). *Porphyromonas* spp. have previously been
isolated from humpback whale skin (Apprill et al., 2014) and humpback whale blows (Apprill et
al., 2017). Furthermore, *Porphyromonas* spp. and *Fusobacterium* spp. have been described as
bacteria of the core pulmonary microbiome in humans (Erb-Downward et al., 2011; Charlson et
al., 2012; Cui et al., 2014; Huang, Yang, & Li, 2016). These bacteria have also been reported
sporadically and in low abundance in the sheep respiratory tract (Glendinning et al., 2016). We

also identified *Moraxella* spp. in blue whale blows. This bacterial genus is present in humpback
whale blows (Apprill et al., 2017) and is commonly found in the lungs of healthy dogs (Tress et
al., 2017). However, it has also been reported in humans (Yi et al., 2014) and cattle with
respiratory diseases (Lima et al., 2016). The bacteria identified in the blue whale respiratory tract
are similar to those reported in other cetaceans and terrestrial mammals, and some are known to
cause diseases. At this stage, we were unable to unequivocally establish that the health of the
whales sampled was not compromised; however, given that they were present in most of the
whales, we can assume that they are part of their respiratory microbiome and that they are likely
to reflect a healthy respiratory tract.

Interestingly, five bacterial genera (*Psychrobacter, Tenacibaculum, Staphylococcus,*
Cutinebacterium, and *Corinebacterium*) identified in the blue whale blow are associated with the
skin of humans and other terrestrial mammals (Grice & Segre, 2011; Byrd, Belkaid, & Segre,
2018; Worthing et al., 2018) and were recently identified as part of the skin microbiota of captive
bottlenose dolphins *Tursiops truncatus*, killer whales *Orcinus orca*, and free-ranging humpback
whales (Hooper, Littman, & Macpherson, 2012; Apprill et al., 2014; Chiarello et al., 2017;
Rhodes et al., 2022). It is likely that their presence in the blow samples indicates that they
colonize the blowhole epithelial lining of blue whales and are forcefully expelled during
exhalation (Apprill et al., 2017), leading to their presence in the blow samples.

Mycoplasma spp. was detected in one sample (Bm057) with a 26.84% relative abundance. The
presence of these bacteria could be indicative of a transient bloom within the respiratory tract or
an active respiratory infection, considering that these bacteria are typically present in the
respiratory tract of mammals at a low abundance, but during active pathological processes, such
as pneumonia and other respiratory conditions, their relative abundance increases (Dai et al.,
2018). In marine mammals, *Mycoplasma* spp. has been associated with respiratory disease and
has been detected in the lungs of stranded harbour porpoises *Phocoena phocoena*, Sowerby's
beaked whales *Mesoplodon bidens* (Foster et al., 2011) and California sea lions *Zalophus*
californianus (Haulena et al., 2013) during unusual mortality events. Furthermore, although this
genus has been found in healthy killer whales (Raverty et al., 2017; Rhodes et al., 2022), the role
of *Mycoplasma* spp. during disease episodes in cetaceans, as well as their host specificity and

association with cetacean stranding events, remains poorly understood (Foster et al., 2011; Rhodes et al., 2022). It is certainly possible that the whale from which sample Bm057 was collected was experiencing a respiratory infection that involved *Mycoplasma* spp. Gaining clinical information that would allow us to establish this beyond any doubt is not feasible, but we propose that future studies of blue whale populations consider the presence of *Mycoplasma* spp. as an indicator of suboptimal respiratory health.

Infectious diseases are one of the leading causes of death in marine mammals (Numberger et al., 2021; Morick et al., 2022), and *Streptococcus* spp. is one of the most reported pathogens of pinnipeds and cetaceans (Bianchi et al., 2018). We identified one sample (Bm044) with a high relative abundance (69.35%) of *Streptococcus* spp. Overall, ten streptococcal species have been isolated and identified multiple times from twenty-three pinniped and cetacean species worldwide (Numberger et al., 2021). *Streptococcus agalactiae*, for example, was isolated from a wound and navel infection in grey seals *Halichoerus grapus* (Baker, 1988), as well as lung and spleen samples from Antarctic fur seals *Arctocephalus gazella* (Baker & McCann, 1989). *Streptococcus phocae* is frequently found in lung lesions, implying commensal colonization of the oropharynx and subsequent opportunistic respiratory tract infections (Taurisano et al., 2018), and *Streptococcus iniae* is a significant aquatic pathogen of farmed fish species, an important zoonotic pathogen, and an identified cause of disease in captive Amazon River dolphins *Inia geoffrensis* (Souter et al., 2021), bottlenose dolphins (Evans et al., 2006), and death in short-beaked common dolphins (Souter et al., 2021).

Having detected two potentially pathogenic bacterial genera in this study could mean that the Eastern North Pacific blue whale is not commonly in contact with coastal areas where spillover of pathogens from humans or domestic animals could occur. This would be in stark contrast to killer whales, which live in areas where there are a large number of environmental anthropic stressors (Raverty et al., 2017). However, a previous study reported *Entamoeba* spp., *Giardia* spp., and *Balantidium* spp., most likely from sewage discharge, in faeces of blue whales from the Gulf of California (Pacheco-Armenta, 2019). Therefore, it is plausible that rather than limited exposure, the presence of *Mycoplasma* spp. and *Streptococcus* spp. in two individuals reflect a

suboptimal immune status or an underlying upper or lower respiratory condition, which could

allow respiratory colonization of this pathogen.

It is important to note that the respiratory microbiome of the blue whales analysed in our study

harboured bacteria that are commonly found in the oropharynx, nasopharynx, and mouth of

different terrestrial mammals (German & Palmer, 2006; Guglielmetti et al., 2010), in which

those anatomical structures are interconnected in the upper respiratory tract. In contrast,

cetaceans have no anatomical connection between the nasopharynx and mouth (Apprill et al.,

2017; Smith, Tang, & Reidenberg, 2017). This finding provides strong evidence that the core

microbiome that we described belongs to the respiratory system of blue whales and does not

include their oral bacteria.

Given the current state of our oceans, which face habitat degradation, pollution, and other

anthropogenic stressors (Melcón et al., 2012; Mouton & Botha, 2012; Palmer et al., 2022);

suboptimal immune responses can occur in top-predator marine animals (Acevedo-Whitehouse

& Duffus, 2009; Van Bressem et al., 2014; Hall et al., 2018), in turn increasing the risk of

diseases in their populations (Sós et al., 2013; Van Bressem et al., 2015; Reisfeld et al., 2019)

The study of the respiratory bacteriome of blue whales provides an excellent opportunity to

develop conservation plans for this endangered species. Researchers can develop effective

strategies to protect blue whales from disease, pollution, and other threats by better

understanding the microbial communities inhabiting their respiratory tracts. To use the

respiratory bacteriome as a tool to help assess the health of large whales (Apprill et al., 2017;

Toro et al., 2021; Rhodes et al., 2022), it is imperative to first identify the composition of the

microbiome of the respiratory tract of different species, which is what we have done for the

Eastern North Pacific blue whales.

Conclusion

To the best of our knowledge, this is the first study to characterize the microbiome of the

respiratory tract of blue whales. We found that the Eastern North Pacific blue whales sampled in

the Gulf of California harboured a similar respiratory bacterial composition among individuals.

Additionally, richness and relative abundance results were comparable with those reported in the

microbiome of healthy animals and humans; therefore, we propose that the core respiratory bacteriome identified here could be used as a baseline for future long-term studies aimed at identifying shifts in the composition and co-occurrence patterns of the respiratory microbiome and identify ASVs related to changes in body condition, as a proxy for poor health condition. This knowledge can help conservation efforts to protect blue whales during their long migrations and ensure that future generations will continue to marvel these incredible creatures and appreciate their ecological role in maintaining the health of our oceans.

Ethical approval and consent to participate. This study complied with the recommendations and methods for approaching blue whales provided by Mexican legislation (NOM-059-SEMARNAT-2010). All procedures were approved by the Bioethics Committee of the Universidad Autónoma de Queretaro (Mexico), and sampling was conducted under permits SGPA/DGVS/00255/16 and SGPA/DGVS/01832/17 issued by the Dirección General de Vida Silvestre to D. Gendron.

Acknowledgements. We thank Manuel Antonio Zamarrón Nunez for his assistance during navigation, and Ana Sofia Merino, Aurora Paniagua, Madeleine Gauthier, Daniel Valdivia and Ricardo Mirsha Mata Cruz for their help during fieldwork. CADS was funded by a CONACYT PhD Studentship (558253). Fieldwork (sampling and navigation) was funded by The Instituto Politécnico Nacional (SIP20160496 and 2017014), Rufford Foundation 2nd small grant for Nature Conservation (2017), and the Program for the Conservation of Species at Risk (Programa de Conservación de Especies en Riesgo, Comisión Nacional de Áreas Naturales Protegidas). Molecular analysis was partly financed by a Small Grant in Aid of Research from the Society for Marine Mammalogy.

Competing interests. The authors declare that they do not have competing interests.

Availability of data and materials. The datasets used and analysed during the current study are available from the corresponding author upon reasonable request.

Authors' contributions. CAD collected the samples, performed molecular analyses, analysed
the data, and drafted the manuscript. RCA conducted statistical programming for microbiome
analysis and network construction and helped interpret the results. DG conducted fieldwork,
collected samples, and co-supervised the research. KAW conceived, designed, and supervised
the research. All authors read and commented the final draft of the manuscript and gave approval
for publication.
469
470 **References**
471

Acevedo-Whitehouse, K. A., & Duffus, A. L. J. (2009). Effects of environmental change on
wildlife health. *Philosophical Transactions of the Royal Society B: Biological Sciences* **364**,
3429–3438.

Aguilar, A., & Borrell, A. (2022). Unreported catches, impact of whaling and current status of
blue whales in the South European Atlantic Shelf. *Sci Rep* **12**.

Ahlgren, N. A., Perelman, J. N., Yeh, Y. C., & Fuhrman, J. A. (2019). Multi-year dynamics of
fine-scale marine cyanobacterial populations are more strongly explained by phage
interactions than abiotic, bottom-up factors. *Environ Microbiol* **21**, 2948–2963.

Ainsworth, T. D., Krause, L., Bridge, T., Torda, G., Raina, J. B., Zakrzewski, M., Gates, R. D.,
Padilla-Gamiño, J. L., Spalding, H. L., Smith, C., Woolsey, E. S., Bourne, D. G., Bongaerts,
P., Hoegh-Guldberg, O., & Leggat, W. (2015). The coral core microbiome identifies rare
bacterial taxa as ubiquitous endosymbionts. *ISME Journal* **9**, 2261–2274.

Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data
[Online].

Antwis, R. E., Lea, J. M. D., Unwin, B., & Shultz, S. (2018). Gut microbiome composition is
associated with spatial structuring and social interactions in semi-feral Welsh Mountain
ponies. *Microbiome* **6**, 1–11.

Apprill, A., Miller, C. A., van Cise, A. M., U'Ren, J. M., Leslie, M. S., Weber, L., Baird, R. W.,
Robbins, J., Landry, S., Bogomolni, A., & Waring, G. (2020). Marine mammal skin
microbiotas are influenced by host phylogeny. *R Soc Open Sci* **7**, 192046.

Apprill, A., Miller, C., Moore, M. J., Durban, J. W., Fearnbach, H., & Barrett-Lennard, L. G.
(2017). Extensive Core Microbiome in Drone-Captured Whale Blow Supports a Framework
for Health Monitoring. *mSystems* **2**, e00119-17.

Apprill, A., Robbins, J., Eren, A., Pack, A., Reveillaud, J., Mattila, D., Moore, M. J., Niemeyer,
M., Moore, K., & Mincer, T. (2014). Humpback whale populations share a core skin
bacterial community: Towards a health index for marine mammals? *PLoS One* **9**, e90785.

Astudillo-García, C., Bell, J. J., Webster, N. S., Glasl, B., Jompa, J., Montoya, J. M., & Taylor,
M. W. (2017). Evaluating the core microbiota in complex communities: A systematic
investigation. *Environ Microbiol* **19**, 1450–1462.

Attard, C., Beheregaray, L., & Möller, L. (2016). Towards population-level conservation in the
critically endangered Antarctic blue whale: The number and distribution of their
populations. *Sci Rep* **6**, 22291.

Baker, J. R. (1988). Further studies on grey seal (Halichoerus grypus) pup mortality on North
Rona. *Br Vet J* **144**, 497–506.

Baker, J. R., & McCann, T. S. (1989). Pathology and bacteriology of adult male Antarctic fur
seals, Arctocephalus gazella, dying at Bird Island, South Georgia. *Br Vet J* **145**, 263–275.

Barlow, D. R., Klinck, H., Ponirakis, D., Branch, T. A., & Torres, L. G. (2023). Environmental
conditions and marine heatwaves influence blue whale foraging and reproductive effort.
Ecol Evol **13**.

Bianchi, M. V., Ehlers, L. P., Vargas, T. P., Lopez, B. C., Taunde, P. A., De Cecco, B. S.,
Henker, L. C., Vielmo, A., Lorenzett, M. P., Riboldi, C. I., Lopes, C. E., De Amorin, D. B.,
Diaz-Delgado, J., Snel, G. G., Siqueira, F. M., & Sonne, L. (2018). Omphalitis,
urachocystitis and septicemia by Streptococcus dysgalactiae in a Southern right whale calf
(Eubalaena australis), Brazil. *Diseases of Aquatic Organisms* **131**, 227–232.

Bierlich, K. C., Miller, C., DeForce, E., Friedlaender, A. S., Johnston, D. W., & Apprill, A.
(2018). Temporal and regional variability in the skin microbiome of humpback whales
along the Western Antarctic Peninsula. *Appl Environ Microbiol* **84**, e02574-17.

Bik, E. M., Costello, E. K., Switzer, A. D., Callahan, B. J., Holmes, S. P., Wells, R. S., Carlin, K.
P., Jensen, E. D., Venn-watson, S., & Relman, D. A. (2016). Marine mammals harbor
unique microbiotas shaped by and yet distinct from the sea. *Nat Commun* **7**, 1–13.

Björk, J. R., O'Hara, R. B., Ribes, M., Coma, R., & Montoya, J. M. (2018). The dynamic core
microbiome: Structure, dynamics and stability. *bioRxiv* 137885.

Bond, S. L., Timsit, E., Workentine, M., Alexander, T., & Léguillette, R. (2017). Upper and
lower respiratory tract microbiota in horses: Bacterial communities associated with health
and mild asthma (inflammatory airway disease) and effects of dexamethasone. *BMC*
Microbiol **17**, 184.

Byrd, A. L., Belkaid, Y., & Segre, J. A. (2018). The human skin microbiome. *Nat Rev Microbiol*
16, 143–155.

Callahan, B. J., McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME Journal* **11**, 2639–2643.

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nat*
Methods **13**, 581–583.

Caruso, V., Song, X., Asquith, M., & Karstens, L. (2019). Performance of Microbiome Sequence
Inference Methods in Environments with Varying Biomass. *mSystems* **4**, e00163-18.

Chaban, B., Albert, A., Links, M. G., Gardy, J. L., Tang, P., & Hill, J. E. (2013).
Characterization of the Upper Respiratory Tract Microbiomes of Patients with Pandemic
H1N1 Influenza. *PLoS One* **8**, e69559.

Chan, Y. K., Estaki, M., & Gibson, D. L. (2013). Clinical consequences of diet-induced
dysbiosis. *Ann Nutr Metab* **63**, 28–40.

Chao, A., & Chiu, C.-H. (2016). Nonparametric Estimation and Comparison of Species
Richness. *eLS* 1–11.

Charlson, E. S., Bittinger, K., Chen, J., Diamond, J. M., Li, H., Collman, R. G., & Bushman, F.
D. (2012). Assessing Bacterial Populations in the Lung by Replicate Analysis of Samples
from the Upper and Lower Respiratory Tracts. (S. K. Highlander, Ed.)*PLoS One* **7**, e42786.

Chiarello, M., Villéger, S., Bouvier, C., Auguet, J. C., & Bouvier, T. (2017). Captive bottlenose
dolphins and killer whales harbor a species-specific skin microbiota that varies among
individuals. *Sci Rep* **7**, 15269.

Comizzoli, P., & Power, M. (2019). Reproductive Microbiomes in Wild Animal Species: A New
Dimension in Conservation Biology. *Adv Exp Med Bio.* **1200**, 225–240.

Coyte, K., Rao, C., Rakoff-Nahoum, S., & Foster, K. (2019). Community assembly in the
microbiome: ecological insights into infant microbiome development. *Access Microbiol* **1**,
241.

Cui, L., Morris, A., Huang, L., Beck, J. M., Twigg, H. L., von Mutius, E., & Ghedin, E. (2014).
The Microbiome and the Lung. *Ann Am Thorac Soc* **11**, S227–S232.

Dahan, D., Jude, B. A., Lamendella, R., Keesing, F., & Perron, G. G. (2018). Exposure to arsenic
alters the microbiome of larval zebrafish. *Front Microbiol* **9**, 1–12.

Dai, W., Wang, H., Zhou, Q., Feng, X., Lu, Z., Li, D., Yang, Z., Liu, Y., Li, Y., Xie, G., Shen,
K., Yang, Y., Zheng, Y., & Li, S. (2018). The concordance between upper and lower
respiratory microbiota in children with Mycoplasma pneumoniae pneumonia article. *Emerg*
Microbes Infect **7**, 92.

Dallas, J. W., & Warne, R. W. (2023). Captivity and Animal Microbiomes: Potential Roles of
Microbiota for Influencing Animal Conservation. *Microb Ecol* **85**, 820-838.

Dickson, R. P., Erb-Downward, J. R., & Huffnagle, G. B. (2014). Towards an Ecology of the
Lung: New Conceptual Models of Pulmonary Microbiology and Pneumonia Pathogenesis.
Lancet Respiratory Medicine **2**, 238–246.

Dickson, R. P., Erb-Downward, J. R., Martinez, F. J., & Huffnagle, G. B. (2016). The
Microbiome and the Respiratory Tract. *Annu Rev Physiol* **78**, 7.1-7.24.

Dinsdale, E. A., Edwards, R. A., Hall, D., Angly, F., Breitbart, M., Brulc, J. M., Furlan, M.,
Desnues, C., Haynes, M., Li, L., McDaniel, L., Moran, M. A., Nelson, K. E., Nilsson, C.,
Olson, R., Paul, J., Brito, B. R., Ruan, Y., Swan, B. K., Stevens, R., Valentine, D. L.,
Thurber, R. V., Wegley, L., White, B. A., & Rohwer, F. (2008). Functional metagenomic
profiling of nine biomes. *Nature* **452**, 629–632.

Domínguez-Sánchez, C. A., Acevedo-Whitehouse, K. A., Gendron, D., & Hoo. (2018). Effect of
drone-based blow sampling on blue whale (*Balaenoptera musculus*) behavior. *Mar Mamm*
Sci **34**, 841–850.

Edgar, R. C. (2018). Sequence analysis Updating the 97 % identity threshold for 16S ribosomal
RNA OTUs. *Bioinformatics* **34**, 2371–2375.

Eloe-fadrosh, E. A., & Rasko, D. A. (2013). The Human Microbiome□: From Symbiosis to
Pathogenesis. *Annu Rev Med* **64**, 145–163.

Erb-Downward, J. R., Thompson, D. L., Han, M. K., Freeman, C. M., McCloskey, L., Schmidt,
L. A., Young, V. B., Toews, G. B., Curtis, J. L., Sundaram, B., Martinez, F. J., &
Huffnagle, G. B. (2011). Analysis of the Lung Microbiome in the "Healthy" Smoker and in
COPD. *PLoS One* **6**, e16384.

Evans, J. J., Pasnik, D. J., Klesius, P. H., & Al-Ablani, S. (2006). First report of Streptococcus
agalactiae and Lactococcus garvieae from a wild bottlenose dolphin (Tursiops truncatus). *J*
Wildl Dis **42**, 561–569.

Ferguson, S. H., Higdon, J. W., Hall, P. A., Hansen, R. G., & Doniol-Valcroze, T. (2021).
Developing a Precautionary Management Approach for the Eastern Canada-West
Greenland Population of Bowhead Whales (Balaena mysticetus). *Front Mar Sci* **8**, 709989.

Foster, G., McAuliffe, L., Dagleish, M. P., Barley, J., Howie, F., Nicholas, R. A. J., & Ayling, R.
D. (2011). Mycoplasma Species Isolated from Harbor Porpoises (Phocoena phocoena) and a
Sowerby's Beaked Whale (Mesoplodon bidens) Stranded in Scottish Waters. *J Wildl Dis*
47, 206–211.

Foster, K. R., Schluter, J., Coyte, K. Z., & Rakoff-Nahoum, S. (2017). The evolution of the host
microbiome as an ecosystem on a leash. *Nature* **548**, 43–51.

Gagliardi, A., Totino, V., Cacciotti, F., Iebba, V., Neroni, B., Bonfiglio, G., Trancassini, M.,
Passariello, C., Pantanella, F., & Schippa, S. (2018). Rebuilding the Gut Microbiota
Ecosystem. *International Journal of Environmental Research and Publick Health* **15**,
e1679.

Gendron, D., & Ugalde de la Cruz, A. (2012). A new classification method to simplify blue
whale photo- identification technique. *Journal of Cetacean Resource Management* **12**, 79–
84.

German, R. Z., & Palmer, J. B. (2006). Anatomy and development of oral cavity and pharynx.
GI Motility online 1–8.

Glendinning, L., McLachlan, G., & Vervelde, L. (2017). Age-related differences in the
respiratory microbiota of chickens. *PLoS One* **12**, e0188455.

Glendinning, L., Wright, S., Pollock, J., Tennant, P., & Collie, D. (2016). Variability of the
Sheep Lung Microbiota. *Appl Environ Microbiol* **82**, 3225–3238.

Grice, E. A., & Segre, J. A. (2011). The skin microbiome. *Nat Rev Microbiol* **9**, 244–253.

Guglielmetti, S., Taverniti, V., Minuzzo, M., Arioli, S., Stuknyte, M., Karp, M., & Mora, D.
(2010). Oral bacteria as potential probiotics for the pharyngeal mucosa. *Appl Environ*
Microbiol **76**, 3948–3958.

Haegeman, B., Hamelin, J., Moriarty, J., Neal, P., Dushoff, J., & Weitz, J. S. (2013). Robust
estimation of microbial diversity in theory and in practice. *ISME J* **7**, 1092–1101.

Haegeman, B., Sen, B., Godon, J. J., & Hamelin, J. (2014). Only Simpson Diversity can be
Estimated Accurately from Microbial Community Fingerprints. *Microb Ecol* **68**, 169–172.

Hall, A. J., McConnell, B. J., Schwacke, L. H., Ylitalo, G. M., Williams, R., & Rowles, T. K.
(2018). Predicting the effects of polychlorinated biphenyls on cetacean populations through
impacts on immunity and calf survival. *Environmental Pollution* **233**, 407–418.

Haulena, M., Gulland, F. M. D., Lawrence, J. A., Fauquier, D. A., Jang, S., Aldridge, B.,
Spraker, T., Thomas, L. C., Brown, D. R., Wendland, L., & Davidson, M. K. (2013).
Lesions Associated With a Novel Mycoplasma Sp. in California Sea Lions (*Zalophus*
Californianus) Undergoing Rehabilitation. *J Wildl Dis* **42**, 40–45.

Heard, M. J., Smith, K. F., Ripp, K., Berger, M., Chen, J., Dittmeier, J., Goter, M., McGarvey, S.
T., & Ryan, E. (2013). The threat of disease increases as species move toward extinction.
Conserv Biol **27**, 1378–1388.

Hernandez-Agreda, A., Gates, R. D., & Ainsworth, T. D. (2017). Defining the Core Microbiome
in Corals' Microbial Soup. *Trends Microbiol* **25**, 125–140.

Hooper, R., Brealey, J. C., van der Valk, T., Alberdi, A., Durban, J. W., Fearnbach, H.,
Robertson, K. M., Baird, R. W., Bradley Hanson, M., Wade, P., Gilbert, M. T. P., Morin, P.
A., Wolf, J. B. W., Foote, A. D., & Guschanski, K. (2019). Host-derived population
genomics data provides insights into bacterial and diatom composition of the killer whale
skin. *Mol Ecol* **28**, 484–502.

Hooper, L. V., Littman, D., & Macpherson, A. (2012). Interactions Between the Microbiota and
the Immune System. *Science (1979)* **336**, 1223490.

Huang, Y., Yang, B., & Li, W. (2016). Defining the normal core microbiome of conjunctival
microbial communities. *Clinical Microbiology and Infection* **22**, 643.e7-643.e12.

Huffnagle, G. B., Dickson, R. P., & Lukacs, N. W. (2017). The respiratory tract microbiome and
lung inflammation: A two-way street. *Mucosal Immunol* **10**, 299–306.

Huse, S. M., Ye, Y., Zhou, Y., & Fodor, A. A. (2012). A core human microbiome as viewed through 16S rRNA sequence clusters. *PLoS One* **7**, e34242.

Infante-Villamil, S., Huerlimann, R., & Jerry, D. R. (2021). Microbiome diversity and dysbiosis in aquaculture. *Rev Aquac* **13**, 1077–1096.

Ingala, M. R., Simmons, N. B., Wultsch, C., Krampis, K., Speer, K. A., & Perkins, S. L. (2018). Comparing microbiome sampling methods in a wild mammal: Fecal and intestinal samples record different signals of host ecology, evolution. *Front Microbiol* **9**, 803.

Jiang, H., Luo, S., Zhou, J., Huang, W., Li, L., Zhang, X., He, J., & Chen, J. (2022) Skin Microbiota Was Altered in Crocodile Lizards (*Shinisaurus crocodilurus*) With Skin Ulcer. *Front Vet Sci* **9**, 817490.

Johnson, K. V. A., & Burnet, P. W. J. (2016). Microbiome: Should we diversify from diversity? *Gut Microbes* **7**, 455–458.

Johnson, W. R., Torralba, M., Fair, P. A., Bossart, G. D., Nelson, K. E., & Morris, P. J. (2009a). Novel diversity of bacterial communities associated with bottlenose dolphin upper respiratory tracts. *Environ Microbiol Rep* **1**, 555–562.

Johnson, W. R., Torralba, M., Fair, P. A., Bossart, G. D., Nelson, K. E., & Morris, P. J. (2009b). Novel diversity of bacterial communities associated with bottlenose dolphin upper respiratory tracts. *Environ Microbiol Rep* **1**, 555–562.

Jousset, A., Bienhold, C., Chatzinotas, A., Gallien, L., Gobet, A., Kurm, V., Küsel, K., Rillig, M. C., Rivett, D. W., Salles, J. F., Van Der Heijden, M. G. A., Youssef, N. H., Zhang, X., Wei, Z., & Hol, G. W. H. (2017). Where less may be more: how the rare biosphere pulls ecosystems strings. *The ISME Journal 2017 11:4* **11**, 853–862.

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G., Bushman, F. D., Knight, R., & Kelley, S. T. (2013). Bayesian community-wide culture-independent microbial source tracking. *Nat Methods* **8**, 761–763.

Lahti, L., Shetty, S., Blake, T., & Salajarvi, J. (2017). Tools for microbiome analysis in R.

Lee, Y. K., & Mazmanian, S. K. (2010). Has the microbiota played a critical role in the evolution of the adaptive immune system? *Science (1979)* **330**, 1768–1773.

Lemon, K. P., Klepac-Ceraj, V., Schiffer, H. K., Brodie, E. L., Lynch, S. V, & Kolter, R. (2010). Comparative Analyses of the Bacterial Microbiota of the Human Nostril and Oropharynx. *mBio* **1**, e00129-10.

Lima, N., Rogers, T., Acevedo-Whitehouse, K. A., & Brown, M. V. (2012). Temporal stability and species specificity in bacteria associated with the bottlenose dolphins respiratory system. *Environ Microbiol Rep* **4**, 89–96.

Lima, S. F., Teixeira, A. G. V, Higgins, C. H., Lima, F. S., & Bicalho, R. C. (2016). The upper
respiratory tract microbiome and its potential role in bovine respiratory disease and otitis
media. *Sci Rep* **6**, 29059.

Man, W. H., De Steenhuijsen Piters, W. A. A., & Bogaert, D. (2017). The microbiota of the
respiratory tract: gatekeeper to respiratory health. *Nature Reviews Microbiology 2017 15:5*
15, 259–270.

Matthewman, C., Narin, A., Huston, H., Hopkins, C. E. (2023) Systems to model the
personalized aspects of microbiome health and gut dysbiosis. *Mol Aspects Med* **91**, 101115.

McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217.

Melcón, M. L., Cummins, A. J., Kerosky, S. M., Roche, L. K., Wiggins, S. M., & Hildebrand, J.
A. (2012). Blue Whales Respond to Anthropogenic Noise. *PLoS One* **7**, 32681.

Mikryukov, V. (2023, February 13). vmikk/metagMisc: v.0.5.0.

Minich, J. J., Sanders, J. G., Amir, A., Humphrey, G., Gilbert, J. A., & Knight, R. (2019).
Quantifying and Understanding Well-to-Well Contamination in Microbiome Research.
mSystems **4**, e0018619.

Morick, D., Davidovich, N., Zemah-Shamir, Z., Bigal, E., Rokney, A., Ron, M., Blum, S. E.,
Fleker, M., Soto, E., Heckman, T. I., Lau, S. C. K., Wosnick, N., Tchernov, D., & Scheinin,
A. P. (2022). First Isolation and Characterization of Streptococcus agalactiae From a
Stranded Wild Common Dolphin (Delphinus delphis). *Front Mar Sci* **9**, 824071.

Mouton, M., & Botha, A. (2012). Cutaneous lesions in cetaceans: An indicator of ecosystem
status? In A. Romero & E. O. Keith (Eds.), *New Approaches to the Study of Marine*
Mammals. pp. 123–144. Croatia: InTech.

Murray, D. (2013). Tableau Your Data! Fast and Easy Visual Analysis with Tableau Software.
1st ed. Wiley Publishing.

Nelson, T., Apprill, A., Mann, J., Rogers, T. L., & Brown, M. V. (2015). The marine mammal
microbiome: current knowledge and future directions. *Microbiol Aust* **36**, 8.

Nishida, A. H., & Ochman, H. (2018). Rates of gut microbiome divergence in mammals. *Mol*
Ecol **27**, 1884–1897.

Numberger, D., Siebert, U., Fulde, M., & Valentin-Weigand, P. (2021). Streptococcal Infections
in Marine Mammals. *Microorganisms* **9**, 350.

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., Minchin, P. R.,
O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, H. H., & Szoecs, E. (2019). vegan:
Community Ecology Package.

Ozkan, J., Nielsen, S., Diez-Vives, C., Coroneo, M., Thomas, T., & Willcox, M. (2017).
Temporal Stability and Composition of the Ocular Surface Microbiome. *Sci Rep* **7**, 9880.

Pacheco-Armenta, M. J. (2019). Identificación morfológica y molecular de endoparásitos de la
ballena azul (Balaenoptera musculus) de vida libre en el suroeste del Golfo de California. .
Instituto Politécnico Nacional. Centro Interdisciplinario de Ciencias Marinas, La Paz.

Palmer, E., Alexander, A., Liggins, L., Guerra, M., Bury, S. J., Hendriks, H., Stockin, K. A., &
Peters, K. J. (2022). A piece of the puzzle: analyses of recent strandings and historical
records reveal new genetic and ecological insights on New Zealand sperm whales. *Mar Ecol*
Prog Ser **690**, 201–217.

Panthee, B., Gyawali, S., Panthee, P., & Techato, K. (2022). Environmental and Human
Microbiome for Health. *Life 2022, Vol. 12, Page 456* **12**, 456.

Pedersen, A. B., Jones, K. E., Nunn, C. L., & Altizer, S. (2007). Infectious diseases and
extinction risk in wild mammals. *Conserv Biol* **21**, 1269–1279.

Pedrós-Alió, C. (2012). The Rare Bacterial Biosphere. *Ann Rev Mar Sci* **4**, 449–466.

Pershing, A. J., Christensen, L. B., Record, N. R., Sherwood, G. D., & Stetson, P. B. (2010). The
impact of whaling on the ocean carbon cycle: Why bigger was better. *PLoS One* **5**, e12444.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Rg Peplies, J., & Glö
Ckner, F. O. (2013). The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590–D596.

R Core Team. (2022). R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing.

Raverty, S., Leger, J. S., Noren, D. P., Huntington, K. B., Rotstein, D. S., Gulland, F. M. D.,
Ford, J. K. B., Bradley Hanson, M., Lambourn, D. M., Huggins, J., Delaney, M. A., Spaven,
L., Rowles, T., Barre, L., Cottrell, P., Ellis, G., Goldstein, T., Terio, K., Duffield, D., Rice,
J., & Gaydos, J. K. (2020). Pathology findings and correlation with body condition index in
stranded killer whales (Orcinus orca) in the northeastern Pacific and Hawaii from 2004 to
2013. *PLoS One* **15**.

Raverty, S., Rhodes, L. D., Zabek, E., Eshghi, A., Cameron, C. E., Hanson, M. B., & Schroeder,
J. P. (2017). Respiratory Microbiome of Endangered Southern Resident Killer Whales and
Microbiota of Surrounding Sea Surface Microlayer in the Eastern North Pacific. *Sci Rep* **7**,
1–12.

Reisfeld, L., Sacristán, C., Canedo, P., Schwarz, B., Ewbank, A. C., Esperón, F., & Catão-Dias,
J. L. (2019). Fusariosis in a Captive South American Sea Lion (*Otaria flavescens*): A Case
Report. *Mycopathologia* **184**, 187–192.

Rhodes, L. D., Emmons, C. K., Wisswaesser, G. S., Wells, A. H., & Hanson, M. B. (2022).
Bacterial microbiomes from mucus and breath of southern resident killer whales (Orcinus
orca). *Conserv Physiol* **10**.

Rice, A., Širović, A., Hildebrand, J. A., Wood, M., Carbaugh-Rutland, A., & Baumann-Pickering, S. (2022). Update on frequency decline of Northeast Pacific blue whale (Balaenoptera musculus) calls. *PLoS One* **17**.

Risely, A. (2020). Applying the core microbiome to understand host–microbe systems. *Journal of Animal Ecology* 1–10.

Robles-Malagamba, M. J., Walsh, M. T., Ahasan, M. S., Thompson, P., Wells, R. S., Jobin, C., Fodor, A. A., Winglee, K., & Waltzek, T. B. (2020). Characterization of the bacterial microbiome among free-ranging bottlenose dolphins (Tursiops truncatus). *Heliyon* **6**, e03944.

Ross, A. A., Rodrigues-Hoffmann, A., & Neufeld, J. D. (2019). The skin microbiome of vertebrates. *Microbiome* **7**, 1–14.

Russel, J. (2023). MicEco: Various functions for microbial community data.

Salter, S. J., Cox, M. J., Turek, E. M., Calus, S. T., Cookson, W. O., Moffatt, M. F., Turner, P., Parkhill, J., Loman, N. J., & Walker, A. W. (2014). Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biol* **12**, 87.

Sehnal, L., Brammer-Robbins, E., Wormington, A. M., Blaha, L., Bisesi, J., Larkin, I., Martyniuk, C. J., Simonin, M., & Adamovsky, O. (2021). Microbiome Composition and Function in Aquatic Vertebrates: Small Organisms Making Big Impacts on Aquatic Animal Health. *Front Microbiol* **12**, 567408.

SEMARNAT. (2018). Programa de Acción para la Conservación de la Especie Ballena Azul (Balaenoptera musculus),. México, D.F.

Shade, A., & Handelsman, J. (2012). Beyond the Venn diagram: The hunt for a core microbiome. *Environ Microbiol* **14**, 4–12.

Shapira, M. (2016). Gut Microbiotas and Host Evolution: Scaling Up Symbiosis. *Trends Ecol Evol* **31**, 539–549.

Sharp, S. M., McLellan, W. A., Rotstein, D. S., Costidis, A. M., Barco, S. G., Durham, K., Pitchford, T. D., Jackson, K. A., Daoust, P. Y., Wimmer, T., Couture, E. L., Bourque, L., Frasier, T., Frasier, B., Fauquier, D., Rowles, T. K., Hamilton, P. K., Pettis, H., & Moore, M. J. (2019). Gross and histopathologic diagnoses from north atlantic right whale eubalaena glacialis mortalities between 2003 and 2018. *Dis Aquat Organ* **135**, 1–31.

Shreiner, A. B., Kao, J. Y., & Young, V. B. (2015). The gut microbiome in health and in disease. *Current Opinion Gastroenterology* **31**, 69–75.

Skopina, M. Y., Vasileva, A. A., Pershina, E. V, & Pinevich, A. V. (2016). Diversity at low abundance: The phenomenon of the rare bacterial biosphere. *Microbiology (N Y)* **85**, 272–282.

Smith, C. M., Tang, C. Y., & Reidenberg, J. S. (2017). Visualizing the Anatomy and Position of
the Larynx in Balaenopterid Whales. *The Faseb Journal* **31**.

Sós, E., Molnár, V., Lajos, Z., Koroknai, V., & Gál, J. (2013). Successfully Treated
Dermatomycosis in California Sea Lions (*Zalophus Californianus*). *Journal of Zoo and*
Wildlife Medicine **44**, 462–465.

Souter, R., Chaber, A. L., Lee, K., Machado, A., Lam, J., & Woolford, L. (2021). Fatal
Streptococcus iniae Infection in a Juvenile Free-Ranging Short-Beaked Common Dolphin
(Delphinus delphis). *Animals 2021, Vol. 11, Page 3123* **11**, 3123.

Stubbendieck, R. M., Vargas-Bautista, C., & Straight, P. D. (2016). Bacterial communities:
Interactions to scale. *Front Microbiol* **7**, 1–19.

Taurisano, N. D., Butler, B. P., Stone, D., Hariharan, H., Fields, P. J., Ferguson, H. W., Haulena,
M., Cotrell, P., Nielsen, O., & Raverty, S. (2018). Streptococcus phocae in marine
mammals of northeastern Pacific and Arctic Canada: A retrospective analysis of 85
postmortem investigations. *J Wildl Dis* **54**, 101–111.

Thomas, S., Izard, J., Walsh, E., Batich, K., Chongsathidkiet, P., Clarke, G., Sela, D. A., Muller,
A. J., Mullin, J. M., Albert, K., Gilligan, P., Diguilio, K., Dilbarova, R., Alexander, W., &
George, C. (2017). Primer and Perspective for Non-Microbiologists. *Cancer Res.* **77**, 1783–
1812.

Toro, F., Alarcón, J., Márquez, S., Capella, J., Bahamonde, P., Esperón, F., Moreno-Switt, A., &
Castro-Nallar, E. (2021). Composition and structure of the skin microbiota of rorquals off
the Eastern South Pacific. *FEMS Microbiol Ecol* **97**, 1–10.

Tress, B., Dorn, E. S., Suchodolski, J. S., Nisar, T., Ravindran, P., Weber, K., Hartmann, K., &
Schulz, B. S. (2017). Bacterial microbiome of the nose of healthy dogs and dogs with nasal
disease. *PLoS One* **12**, e0176736.

Van Bressem, M. F., Duignan, P. J., Banyard, A., Barbieri, M., Colegrove, K. M., de Guise, S.,
di Guardo, G., Dobson, A., Domingo, M., Fauquier, D., Fernandez, A., Goldstein, T.,
Grenfell, B., Groch, K. R., Gulland, F., Jensen, B. A., Jepson, P. D., Hall, A., Kuiken, T.,
Mazzariol, S., Morris, S. E., Nielsen, O., Raga, J. A., Rowles, T. K., Saliki, J., Sierra, E.,
Stephens, N., Stone, B., Tomo, I., Wang, J., Waltzek, T., & Wellehan, J. F. X. (2014).
Cetacean morbillivirus: Current knowledge and future directions. *Viruses* **6**, 5145–5181.

Van Bressem, M. F., Minton, G., Collins, T., Willson, A., Baldwin, R., & Van Waerebeek, K.
(2015). Tattoo-like skin disease in the endangered subpopulation of the Humpback Whale,
Megaptera novaeangliae, in Oman (Cetacea: Balaenopteridae). *Zool Middle East* **61**, 1–8.

Van Cise, A. M., Wade, P. R., Goertz, C. E. C., Burek-Huntington, K., Parsons, K. M., Clauss,
T., Hobbs, R. C., & Apprill, A. (2020). Skin microbiome of beluga whales: spatial,
temporal, and health-related dynamics. *Anim Microbiome* **2**, 39.

Van Der Gast, C. J., Walker, A. W., Stressmann, F. A., Rogers, G. B., Scott, P., Daniels, T. W.,
Carroll, M. P., Parkhill, J., & Bruce, K. D. (2010). Partitioning core and satellite taxa from
within cystic fibrosis lung bacterial communities. *The ISME Journal 2011 5:5* **5**, 780–791.

Van Treuren, W., & Dodd, D. (2020). Microbial Contribution to the Human Metabolome:
Implications for Health and Disease. *Annu Rev Pathol* **15**, 345.

Vecchioni, L., Pace, A., Sucato, A., Berlinghieri, F., Cambera, I., Visconti, G., Hochscheid, S.,
Arculeo, M., & Alduina, R. (2022) Unveiling the egg microbiota of the loggerhead sea
turtle Caretta caretta in nesting beaches of the Mediterranean Sea. *PLoS One* **17**, e0268345.

Vendl, C., Nelson, T., Ferrari, B., Thomas, T., & Rogers, T. (2020). Highly abundant core taxa
in the blow within and across captive bottlenose dolphins provide evidence for a temporally
stable airway microbiota. *BMC Microbiol* 1–29.

Watkins, C. A., Gaines, T., Strathdee, F., Baily, J. L., Watson, E., Hall, A. J., Free, A., &
Dagleish, M. P. (2022). A comparative study of the fecal microbiota of gray seal pups and
yearlings - a marine mammal sentinel species. *Microbiologyopen* **11**, e1281.

Watson, R. L., De Koff, E. M., & Bogaert, D. (2019). Characterising the respiratory microbiome.
European Respiratory Journal **53**, 1–11.

Willis, A., Bunge, J., & Whitman, T. (2017). Improved detection of changes in species richness
in high diversity microbial communities. *J R Stat Soc Ser C Appl Stat* **66**, 963–977.

Worthing, K. A., Abraham, S., Coombs, G. W., Pang, S., Saputra, S., Jordan, D., Trott, D. J., &
Norris, J. M. (2018). Clonal diversity and geographic distribution of methicillin-resistant
Staphylococcus pseudintermedius from Australian animals: Discovery of novel sequence
types. *Vet Microbiol* **213**, 58–65.

Wright, E. S. (2016). Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R. *R J*
8, 352–359.

Yamamoto, S., Saito, M., Tamura, A., Prawisuda, D., Mizutani, T., & Yotsuyanagi, H. (2021).
The human microbiome and COVID-19: A systematic review. *PLoS One* **16**.

Yi, H., Yong, D., Lee, K., Cho, Y., & Chun, J. (2014). Profiling bacterial community in upper
respiratory tracts. *BMC Infect Dis* **14**, 1–10.

Figure legends

Figure 1. Representation of the ASVs shared among the Eastern North Pacific blue whale blow

samples and technical control samples. The Venn diagram displays the ASV quantity shared

among blows (blue), human sneeze (yellow), seawater (green) and Laboratory control (red).

Figure 2. Total abundance (%) of the 15 bacterial classes identified in all samples. ASVs with

less than 0.02% of relative abundance are grouped as "others" along with the unclassified.

Figure 3. Stacked bar plot depicting relative abundance of the top 15 bacterial genera detected in

the Eastern North Pacific blue whale blow samples. Each vertical bar depicts the relative

abundance of adjusted sequence variants (ASVs) and associated taxa that were recovered per

sample. Plot shows the top fifteen identified bacterial genera, unclassified are grouped along

"others" (sum of bacteria that did not reach the detection threshold of 0.02%).

Figure 4. a) Regression model shows changes in the pattern of the common central respiratory

bacteriome of the Eastern North Pacific blue whale, based on a range of prevalence and detection

threshold values. **b)** Relative abundances of bacteria genera belonging to the core respiratory

bacteriome of the blue whale (nine classified and two unclassified ASVs present in more than

50% of the samples and more than 0.02% of relative abundance).