A ubiquitin-proteasome pathway degrades the inner nuclear membrane protein Bqt4 to

maintain nuclear membrane homeostasis

Toan Khanh Le[1], Yasuhiro Hirano[1], Haruhiko Asakawa[1], Koji Okamoto[2], Tokuko

Haraguchi[1] and Yasushi Hiraoka[1,*]

[1] Nuclear Dynamics Group, Graduate School of Frontier Biosciences, Osaka University, Suita

565-0871, Japan;

[2] Laboratory of Mitochondrial Dynamics, Graduate School of Frontier Biosciences, Osaka

University, Suita 565-0871, Japan

Running title: Bqt4 degradation pathway in NE

[*] **Corresponding to:**

Yasushi Hiraoka (hiraoka@fbs.osaka-u.ac.jp), Graduate School of Frontier Biosciences,

Osaka University, 1-3 Yamadaoka, Suita 565-0871, Japan. Tel.: +81-6-6879-4621, FAX:

+81-6-6789-4622

Keywords: Inner nuclear membrane, Protein degradation, Fission Yeast, Bqt3, Bqt4

Summary statement

The appropriate level of the inner nuclear membrane protein Bqt4 is maintained by ubiquitin-

proteasome-mediated degradation. Aberrant accumulation of Bqt4 disturbs the nuclear

membrane homeostasis.

Abstract

Aberrant accumulation of inner nuclear membrane (INM) proteins has been associated with

deformed nuclear morphology and certain mammalian diseases. However, the mechanisms

by which INM homeostasis is maintained remain poorly understood. In this study, we

explored the degradation mechanisms of the INM protein Bqt4 in the fission yeast

Schizosaccharomyces pombe. We have previously shown that Bqt4 interacts with the

transmembrane protein Bqt3 at the INM and is degraded in the absence of Bqt3. Here, we

revealed that excess Bqt4 unassociated with Bqt3 was targeted for degradation by the

ubiquitin-proteasome system localized in the nucleus and that Bqt3 antagonized this process.

The degradation process involves the Doa10 E3 ligase complex at the INM. Bqt4 is a tail-

anchored protein and extraction from the membrane by the Cdc48 complex is required for its

degradation. The C-terminal transmembrane domain of Bqt4 is necessary and sufficient for

proteasome-dependent protein degradation. Accumulation of Bqt4 at the INM impaired cell

viability with nuclear envelope deformation, suggesting that the quantity control of Bqt4

plays an important role in nuclear membrane homeostasis.

Introduction

The eukaryotic nucleus is enclosed by the nuclear envelope (NE), which separates the

nucleoplasm from the cytoplasm. The NE comprises the inner and outer nuclear membranes

(INM and ONM, respectively). The ONM is continuous with the endoplasmic reticulum

(ER). The INM is involved in the organization and regulation of chromosomes by modulating

the chromatin potential for gene expression, DNA replication, DNA damage repair, and

genome stability (Mekhail and Moazed, 2010; Hirano et al., 2020a; Pawar and Kutay, 2021).

Excess amounts of INM proteins cause aberrant accumulation at the NE, often causing

nuclear deformation and malfunctioning (Gonzalez et al., 2012). The aberrant accumulation

of INM proteins has been linked to certain mammalian diseases. For example, accumulation

of the INM protein SUN1 was found to be pathogenic in human laminopathies, such as

Emery-Dreifuss muscular dystrophy and Hutchinson-Gilford progeria syndrome (Chen et al.,

2012). Therefore, the amounts of INM protein must be regulated. However, the mechanisms

that regulate levels of resident INM proteins to maintain INM homeostasis remain poorly

understood.

Mechanisms for ER membrane homeostasis have been well studied: ER membrane

proteins are regulated through the ER-associated degradation (ERAD) system, which is

involved in degrading misfolded or mislocalized ER proteins as a means of protein quality

control (Vembar and Brodsky, 2008; Ruggiano et al., 2014; Christianson and Carvalho,

2022). ERAD involves the ubiquitin-proteasome system (UPS), in which polyubiquitin

chains are covalently attached to a substrate protein, leading to its subsequent recognition and

degradation by the 26S proteasome complex. Ubiquitination reactions involve three different

classes of enzymes: E1 ubiquitin-activating enzymes, E2 ubiquitin-conjugating enzymes, and

E3 ubiquitin ligases such as Hrd1 and Doa10 (human MARCHF6/TEB4). INM-associated

degradation (INMAD) has been proposed as an extension of ERAD to maintain INM

homeostasis, which also depends on the UPS (Foresti et al., 2014; Khmelinskii et al., 2014;

Koch and Yu, 2019; Smoyer and Jaspersen, 2019; Mannino and Lusk, 2022). It has been

shown that in the budding yeast *Saccharomyces cerevisiae*, ERAD and INMAD require

additional machinery for the degradation of integral membrane proteins, involving

Cdc48/p97, an ATPase associated with diverse cellular activities (AAA-ATPase), which

removes the ubiquitinated proteins from the membrane before transferring them to the

proteasome for degradation (Ruggiano et al., 2014; Christianson and Carvalho, 2022; Koch

and Yu, 2019; Smoyer and Jaspersen, 2019; Mannino and Lusk, 2022).

In the fission yeast *Schizosaccharomyces pombe*, several INM proteins have been

studied for their roles in the organization and regulation of chromosome stability, including

Ima1, Lem2, Man1, Bqt3, and Bqt4 (Chikashige et al., 2006; Chikashige et al., 2009; Hiraoka

et al., 2011; Steglich et al., 2012; Gonzalez et al., 2012; Tange et al., 2016; Hirano et al.,

2020b). In particular, we found that Bqt4 was degraded in the absence of Bqt3 (Chikashige et

al., 2009). Bqt3 is predicted to be a transmembrane INM protein composed of 7

transmembrane domains (TMDs) along its entire length; Bqt4, which interacts with Bqt3, is

predicted as a tail-anchored protein (Chikashige et al., 2009). Bqt3 and Bqt4 are required for

anchoring telomeres to the NE during vegetative growth and for clustering telomeres to the

spindle-pole body through the meiosis-specific telomere-binding proteins, Bqt1 and Bqt2

(Chikashige et al., 2006; Chikashige et al., 2009). Importantly, Bqt4 interacts with Lem2

(Hirano et al., 2018), and the depletion of Bqt4 is synthetic-lethal with the depletion of Lem2

(Tange et al., 2016; Kinuagsa et al., 2019). Therefore, it is expected that Bqt4 plays important

roles in cell viability. However, the non-telomeric vital roles of Bqt4 and the mechanisms

underlying its degradation remain unknown. In this study, we identified the factors required

for Bqt4 degradation via an INM-associated degradation mechanism. We also demonstrated

that excess amounts of Bqt4 result in impaired nuclear morphology and growth defects,

indicating the importance of regulated degradation of Bqt4 for membrane homeostasis.

Results

INM protein Bqt4 is degraded via the ubiquitin-proteasome system

Because the ubiquitin-proteasome system accounts for the selective degradation of many

proteins in the nucleus (Enam et al., 2018; Franić et al., 2021), we first tested the requirement

of the proteasome for Bqt4 degradation. We treated cells expressing N-terminally GFP-

tagged Bqt4 (GFP-Bqt4) under its own promoter with bortezomib (BZ), a chemical

proteasome inhibitor working in *S. pombe* (Takeda et al., 2011), and assessed the

fluorescence levels of GFP-Bqt4 at 4 h after drug addition in the presence or absence of Bqt3.

As a result of the treatment, GFP-Bqt4 fluorescence in the nucleus of both *bqt3+* and *bqt3*

cells was elevated (Figure. 1A). To confirm this result, we checked protein levels by Western

blotting, which consistently showed an increase in GFP-Bqt4 protein levels in these strains

upon proteasomal inhibition by BZ (Figure. 1B). Notably, the GFP-Bqt4 signal was restored

upon proteasome inhibition in the absence of Bqt3, and also elevated even in the presence of

Bqt3. This suggests that Bqt4 is constitutively turned over by the proteasome system and its

association with Bqt3 antagonizes this process. We sought to confirm this phenomenon by

introducing temperature-sensitive *mts2-1* and *mts3-1* proteasome mutations (Gordon and

McGurk, 1993; Gordon et al., 1996). Using both fluorescence microscopy and Western

blotting, we found that GFP-Bqt4 levels increased in *bqt3*[+] and *bqt3Δ* cells carrying these

proteasome mutations at the restrictive temperature of 36ºC, as compared to the

corresponding wild-type strains (Figure. 1C, D; Figure. S1). These results indicate that Bqt4

degradation proceeds via the proteasome system.

Since proteins are ubiquitinated as a requirement for recognition by the proteasome,

we examined whether Bqt4 is modified by polyubiquitination. We immunoprecipitated GFP-

Bqt4 under denaturing conditions and detected associated ubiquitin by immunoblotting.

Ubiquitinated forms of Bqt4 were detected in *bqt3Δ* cells and were enriched in both *bqt3⁺*

and *bqt3Δ* cells when proteasomal activity was inhibited (Figure. 1E), suggesting that Bqt4 is

targeted to the proteasome by polyubiquitin modification.

Vacuolar autophagy is another pathway through which cells degrade proteins. Thus,

we next examined whether Bqt4 degradation is regulated by the vacuolar autophagy pathway

by analyzing GFP-Bqt4 levels in *bqt3Δ* cells lacking *isp6* and *atg1* genes. Isp6 is a serine

protease necessary for vacuolar function, and Atg1 is a serine/threonine protein kinase that is

the upstream initiator of the autophagy machinery. Deletion of these two genes has been

shown to completely impair the vacuolar autophagy pathway (Kohda et al., 2007). We found

that the deletion of *isp6* and *atg1* had no detectable effect on the quantity of GFP-Bqt4 in the

absence of Bqt3, suggesting that Bqt4 degradation does not involve the vacuolar autophagy

system (Figure. 1F). Collectively, we concluded that the Bqt4 protein is degraded by the

ubiquitin-proteasome system, without involvement of the vacuolar autophagy pathway.

Bqt4 degradation occurs in the nucleus

Next, we attempted to determine whether Bqt4 was degraded in the nucleus by the nuclear

proteasome. In *S. pombe*, proper targeting of proteasomes to the nucleus requires the

tethering factor Cut8 (Tatebe and Yanagida, 2000). In *cut8* mutants, proteasomes fail to

localize in the nucleus and become mostly cytoplasmic (Tatebe and Yanagida, 2000).

Because nuclear protein degradation depends on the localization of proteasomes in the

nucleus (Takeda and Yanagida, 2005), we examined the level of GFP-Bqt4 in *bqt3⁺* and

bqt3Δ cells carrying the temperature-sensitive *cut8-563* mutation. We found that GFP-Bqt4

degradation was suppressed in *cut8-563* cells shifted to the nonpermissive temperature of

36°C (Figure. 2A, B). Taken together, our results strongly suggest that Bqt4 is degraded

mainly by the proteasome within the nucleus.

Bqt4 degradation involves the Doa10 E3 ligase complex

We next sought to identify the specific E3 ubiquitin ligase that directly recognizes Bqt4. As

Bqt4 is a protein associated with the INM, we investigated whether membrane-localized

Hrd1 and Doa10 E3 ligases participate in the Bqt4 degradation pathway. Deletion of *doa10⁺*

partially restored GFP-Bqt4 levels in the absence of Bqt3, whereas deletion of *hrd1⁺* did not

(Figure. 3A and B, comparing *hrd1Δ* to *doa10Δ*). These results are consistent with previous

studies that showed that Doa10 partially localizes to the INM and is involved in the

degradation of certain nuclear and INM substrates, whereas Hrd1 is found exclusively in the

ER (Deng and Hochstrasser, 2006; Boban et al., 2014). Additionally, the double deletion of

doa10⁺ and *hrd1⁺* showed no obvious synthetic effect on GFP-Bqt4 restoration compared to

the deletion of *doa10⁺* alone (Figure. 3A, B). We also checked the quantity of GFP-Bqt4 in

mutants lacking Ubc6 and Ubc7, which are E2 ubiquitin-conjugating enzymes associated

with Doa10 (Swanson et al., 2001). We found that GFP-Bqt4 levels increased to some extent

in *ubc6Δ*, *ubc7Δ* single, and *ubc6Δ ubc7Δ* double mutants (Figure. 3C, D). These findings

indicate that the Doa10 complex contributes to the degradation of Bqt4 in the INM.

Although degradation of Bqt4 depended on Doa10, the level of GFP-Bqt4 was only

partially restored in the *doa10Δ* mutant (Figure. 3A, B). Therefore, we attempted to search

for other redundant E3 ligases involved in Bqt4 degradation. We constructed mutants of E3

ligases and their components that are suggested to localize or function in the ER and nucleus,

namely *meu34Δ, dsc1Δ, ubr1Δ, ubr11Δ, san1Δ, hul5Δ, ptr1-1*, and *cut9-665* (PomBase,

Matsuyama et al., 2006, Nielsen et al., 2014; Enam et al., 2018; Franić et al., 2021), and

observed GFP-Bqt4 fluorescence intensity in these mutants. Except for the *hul5Δ* mutant,

which exhibited a slight increase in the GFP-Bqt4 level, the other mutants showed no

detectable changes by fluorescence microscopy (Figure. S2). Thus, unidentified E3 ligases

appear to function in the INM in cooperation with Doa10 to ubiquitinate Bqt4.

Bqt4 is an integral membrane protein

We have previously shown that the Bqt4 protein localizes exclusively to the nucleoplasmic

side of the INM, and this localization requires its C-terminal helix (Chikashige et al., 2009).

However, as this C-terminal helix is predicted to be a weak candidate for the TMD (Figure.

S3), we examined whether Bqt4 is an integral or a peripheral membrane protein to understand

how Bqt4 is removed from the INM before degradation. Therefore, we performed a

membrane protein extraction experiment. In brief, cells were lysed, treated under various

conditions, and then separated into pellet and supernatant fractions by ultracentrifugation,

followed by Western blotting; a protein was expected to enter the pellet fraction if it

remained in the lumen or associate with the membrane after the treatment. As a control for a

luminal and an integral membrane protein, we used GFP-ADEL (Rossanese et al., 2001) and

GFP-Bqt3 (Chikashige et al., 2009), respectively. As shown in Figure. 4, all these proteins

were extracted from the membrane by treatment with Triton X-100 and 0.5 M NaCl.

However, similar to Bqt3, Bqt4 was not dissociated from the membrane by sodium carbonate

(pH 11), urea, or NaCl treatments, which can strip peripheral membrane proteins. In contrast,

GFP-ADEL was completely released from the membrane by sodium carbonate treatment, as

expected for luminal proteins. Thus, Bqt4 is likely to behave as a tail-anchored protein and its

C-terminal sequence can act as a TMD that is integrally embedded in the INM.

Degradation of Bqt4 requires the Cdc48 retrotranslocation complex

The ATPase Cdc48 has been shown to facilitate the degradation of ubiquitinated ERAD

membrane substrates by extracting them from the ER membrane in *S. cerevisiae* (Ye et al.,

2001; Jarosch et al., 2002; Rabinovich et al., 2002; Wolf and Stolz, 2012). Therefore, we

investigated whether Cdc48 is necessary for Bqt4 degradation. To this end, we generated a

cdc48 mutant bearing mutations in both the D1 (P267L) and D2 (A550T) ATPase domains,

based on the temperature-sensitive *cdc48-6* allele in *S. cerevisiae* (Schuberth and Buchberger,

2005; Ruggiano et al., 2016) (Figure. 5A). The *cdc48* mutant (*cdc48-P267L/A550T*)

exhibited cold-sensitive growth defects (Figure. 5B), suggesting that these mutations

impaired Cdc48 activity in *S. pombe*. Inactivation of *cdc48-P267L/A550T* by shifting to the

nonpermissive temperature of 16°C elevated the level of GFP-Bqt4 in $bqt3^+$ and $bqt3\Delta$ cells

(Figure. 5C, D, Figure. S4). As the ATPase activity of Cdc48 is required for extracting

membrane proteins, our results suggest that Bqt4 is extracted from the INM by the Cdc48

complex prior to its degradation.

The C-terminal TMD of Bqt4 acts as a degradation signal

We previously showed that Bqt4 degradation depends on its C-terminal TMD (residues 414-

432) (Chikashige et al., 2009). To identify the minimal region of Bqt4 that is sufficient for its

degradation, we performed an analysis of truncated C-terminal fragments based on the

structure of Bqt4 predicted by AlphaFold2 (Figure. 6, Figure. S5). The Bqt4 fragment

(Bqt4C$^{369-432}$) containing the helix domain and the adjacent intrinsically disordered sequence

reproduced behaviors of full-length Bqt4, as shown in Figure. 1A, that is, localization and

responses to BZ, both in the presence and absence of Bqt3 (Figure. 6). The helix domain

alone (Bqt4C$^{394-432}$) showed weaker localization at the NE and somewhat diffused to the

membrane compartments in the cytoplasm; nevertheless, this fragment was degraded in the

absence of Bqt3, and its level increased with BZ treatment (Figure. 6). The TMD alone

(Bqt4C$^{414-432}$) showed even weaker localization at the NE with diffusion to the cytoplasm, but

showed the same responses, that is, degradation in the absence of Bqt3, and increased levels

upon BZ treatment (Figure. 6). These results indicate that the C-terminal fragment containing

residues 369–432 was necessary and sufficient for the NE localization of Bqt4, and its

truncation successively reduced the ability of NE localization. These results also indicated

that the C-terminal TMD of Bqt4 is sufficient for its degradation in the absence of Bqt3.

To further narrow the TMD of Bqt4 for proteasome-dependent degradation and Bqt3-

dependent protection against degradation, we generated constructs with a shorter TMD

truncated of the last seven residues (Bqt4C$^{369-425}$ and Bqt4C$^{414-425}$). The fragments Bqt4C^{369-}

^{425} and Bqt4C$^{414-425}$ lost their dependency on Bqt3 for degradation and showed similar

behaviors in the presence or absence of Bqt3. While Bqt4C$^{369-425}$ retained slight localization

at the NE and Bqt4C$^{414-425}$ showed diffuse localization in the cytoplasm, both fragments

showed increased signal levels in the presence of BZ. These fragments, lacking the last seven

residues, are also substrates of proteasomal degradation in a Bqt3-independent manner

(Figure. 6). These results suggest that the fragment of residues 414–425 is sufficient for

proteasomal degradation and that the fragment of residues 426–432 is required for Bqt3-

dependent protection against proteasomal degradation.

Excess Bqt4 causes deformation of the NE

We observed that inhibition of proteasomal degradation caused striking NE expansion and

deformation with irregular protrusions and invaginations, especially in *bqt3$^+$* cells, in which

Bqt4 levels were excessively high (Figure. 1A, C). We hypothesized that the high levels of

Bqt4 in these cells were responsible for this abnormal NE morphology. To test this

possibility, we inhibited proteasomal degradation in the presence or absence of endogenous

Bqt4 and visualized the NE using Ish1-GFP as an NE marker, and then nuclear deformation

was assessed by roundness of the nucleus. Only when Bqt4 was present was strong

deformation of NE observed upon proteasomal inhibition (see $bqt4^+$+BZ; Figure. 7A),

suggesting that Bqt4 was an essential factor causing this phenomenon. To ascertain whether

this abnormal phenotype was caused by an accumulation of Bqt4, we overexpressed GFP-

Bqt4 under the *nmt1* promoter using YAM2 to control the expression level. YAM2, a

chemical compound, suppresses *nmt1* promoter activity depending on its concentration

(Nakamura et al., 2011). Overexpression of GFP-Bqt4 at levels similar to proteasomal

inhibition reproduced the observed nuclear-deformed morphology, and maximal

overexpression of GFP-Bqt4 resulted in even more striking NE proliferation and deformation

in both $bqt3^+$ and $bqt3\Delta$ cells, indicating that abnormal NE morphology depended on the

amount of Bqt4 (Figure. 7B). Taken together, these results support our hypothesis that Bqt4

is the only protein whose accumulation upon degradation inhibition induces aberrant nuclear

morphology. Finally, we investigated whether excess GFP-Bqt4 was toxic. We found that

overproduction of GFP-Bqt4 caused a growth defect in both $bqt3^+$ and $bqt3\Delta$ strains

compared to cells expressing endogenous levels of Bqt4 (Figure. 7C). Altogether, these

observations suggest that Bqt4 plays an important role in regulating NE morphology and that

a nuclear proteasomal degradation pathway controlling the quantity of Bqt4 is crucial for

nuclear membrane homeostasis.

Discussion

In this study, we identified the molecular factors required for the degradation of the INM

integral protein Bqt4 in *S. pombe* (Figure. 8). Bqt4 undergoes degradation by an INMAD

mechanism involving the Doa10 E3 ligase, which possibly recognizes the C-terminal TMD

of Bqt4 as a degron. Subsequently, the AAA-ATPase Cdc48 removes Bqt4 from the INM,

and Bqt4 is finally degraded via the nuclear proteasome. We also demonstrated that

undegraded Bqt4 accumulating in the INM caused strong deformation of the NE and resulted

in growth defects. Bqt4 escapes degradation when interacting with Bqt3 through its C-

terminal TMD. Excess amounts of Bqt4 that are unbound to Bqt3 are subjected to INMAD-

mediated degradation to avoid the accumulation of Bqt4 at the NE. These findings contribute

to our understanding of the significance of INM protein degradation in INM homeostasis.

Bqt4 is a native INM degradation substrate containing an intramembrane degron

Studies in mammalian cells have suggested that integral INM proteins are largely stable

(Toyama et al., 2013; Buchwalter et al., 2019) and it has been difficult to study their turnover.

In the budding yeast *S. cerevisiae*, most of the INMAD substrates characterized thus far are

not native INM proteins, but rather nuclear soluble proteins (Swanson et al., 2001; Ravid et

al., 2006) and ER membrane proteins that are unassembled or mislocalized to the INM

(Foresti et al., 2014; Khmelinskii et al., 2014; Natarajan et al., 2020). Thus, our study of Bqt4

provides an opportunity for a native INM substrate to obtain further insight into the

mechanisms governing INM protein degradation. Along with several previous studies that

reported a handful of native INM substrates (Boban et al., 2014; Pantazopoulou et al., 2016;

Smoyer et al., 2019; Koch et al., 2019), our results on the degradation of the INM protein

Bqt4 demonstrate that INMAD is responsible for not only maintaining INM identity by

removing foreign proteins but also for maintaining the proteostasis of INM integral

components.

An important question is whether INMAD machinery specifically recognizes its

substrates. Although a degradation signal has yet to be identified for integral INM substrates

of INM-localized E3 ubiquitin ligases, such as Doa10 and the Asi complex, we identified that

Bqt4 has an intramembrane degron. This suggests that INMAD may not simply detect and

degrade misfolded or mislocalized proteins at the INM, but instead specifically recognizes a

substrate to regulate its protein levels.

An INM protein degradation pathway in *S. pombe*

In *S. cerevisiae*, the Doa10 E3 ubiquitin ligase cooperates with the Asi complex in INMAD

(Koch and Yu, 2019; Smoyer and Jaspersen, 2019; Mannino and Lusk, 2022). As homologs

of Asi proteins are absent in *S. pombe* as in other species, INMAD mechanisms may be

diverse among organisms. However, increasing evidence has shown that the degradation of

INM proteins in plants and mammals also employs the UPS and Cdc48/p97 (Tsai et al., 2016;

Huang et al., 2020; Krshnan et al., 2022), raising a general mechanism for INM protein

degradation.

In *S. cerevisiae*, Doa10 E3 ligase can passively diffuse from the ER to the INM and

functions in both ERAD and INMAD (Deng and Hochstrasser, 2006). In particular, Doa10

has been implicated in mediating the degradation of the INM protein Asi2, which itself is a

component of INMAD (Boban et al., 2014). In *S. pombe*, Bqt4 degradation also depends on

Doa10, indicating that Bqt4 is subject to regulation by INMAD. In addition, our data indicate

that *S. pombe* might possess additional unidentified redundant E3 ligases cooperating with

Doa10 at the INM. Because Bqt4 is supposed to be recognized by these E3 ligases at its

transmembrane degron, it is expected that the unidentified E3 ligases also localize to the INM

membranes ("X(E3)" in Figure 8). Similar situations, in which unidentified E3 ligases

degrade an INM substrate, have been reported in *S. cerevisiae* (Pantazopoulou et al. 2016).

Likewise, INM-localized E3 ligases have yet to be identified in higher eukaryotes. We

speculate that the uncharacterized E3 ligases functioning at the INM in *S. pombe* might

operate with E2 ubiquitin-conjugating enzymes other than Ubc6 and Ubc7 because

degradation of Bqt4 exhibits limited dependence on these enzymes. Thus, it would be of

great interest to identify and characterize other E3 ligases and E2 conjugating enzymes that

mediate protein degradation at the INM.

The Cdc48 complex plays a central role in the ERAD degradation of ER integral

membrane proteins by exerting a pulling force on the ubiquitinated membrane protein via its

ATPase activity (Braun, 2002; Rabinovich et al., 2002; Wolf and Stolz, 2012; Bodnar and

Rapoport, 2017). Cdc48 has also been implicated in the degradation of the INM proteins Asi1

and Msp3 in *S. cerevisiae* (Pantazopoulou et al. 2016; Koch et al., 2019). In *S. pombe*, Cdc48

mutations at its ATPase active center impair Bqt4 degradation, indicating that

ERAD/INMAD degradation also utilizes the Cdc48 complex to extract membrane-embedded

substrates and deliver them to the proteasome for proteolysis. Thus, our study demonstrated

the presence of a conserved INMAD system in *S. pombe*.

The biological significance of Bqt4 degradation

The INM protein Bqt4 has been suggested to have non-telomeric functions that are currently

unknown, yet vital for cell growth, as double deletion of Bqt4 and Lem2 results in synthetic

lethality (Tange et al., 2016; Hirano et al., 2018; Kinugasa et al., 2019). In cells lacking both

Bqt4 and Lem2, nuclear envelope rupture occurs in association with the leakage of nuclear

proteins into the cytoplasm, suggesting that Bqt4, in cooperation with Lem2, plays essential

roles in maintaining nuclear membrane integrity (Kinugasa et al., 2019). The nuclear

membrane rupture phenotype can be rescued by an increased expression of Elo2, a fatty acid

elongase that catalyzes the synthesis of very-long-chain fatty acids (Kinugasa et al., 2019). In

addition, depletion of Bqt4 and Lem2 also leads to a reduction in the quantity of t20:0/24:0

phytoceramide, the most abundant ceramide species in *S. pombe*, and this reduction can be

prevented by the overexpression of Elo2 (Kinugasa et al., 2019). These findings strongly

suggest that Bqt4 is involved in nuclear membrane lipid metabolism. Thus, we propose that

abnormal accumulation of Bqt4 at the INM may result in excessive lipid synthesis, ultimately

leading to nuclear membrane expansion and deformation. One possibility is that Bqt4

promotes the synthesis of phospholipids, a major component of the nuclear membrane.

Several lines of evidence suggest that increased phospholipid synthesis from phosphatidic

acid induces striking NE membrane deformation (Santos-Rosa et al., 2005; Barbosa et al.,

2015). The exact mechanisms leading to the deformed nuclear morphology upon the

accumulation of Bqt4 remain to be elucidated. Further investigations will advance our

understanding of the role of INM protein degradation in the maintenance of nuclear

membrane homeostasis.

Materials and methods

Yeast strains and culture media

All *S. pombe* strains used in this study are listed in Table S1. Unless otherwise specified, all *S. pombe* strains were cultured in yeast extract with supplements (YES) medium. Malt extract medium (ME) was used to induce sporulation. G418 disulfate (Nacalai Tesque, Japan), Hygromycin B (Wako, Japan), nourseothricin sulfate (Werner BioAgents, Germany), and Blasticidin S (Wako, Japan) were used at concentrations of 100, 200, 100, and 100 µg/ml, respectively, to select strains bearing the respective selection marker.

Construction of plasmids and strains

All plasmids were constructed using the NEBuilder system (New England BioLabs, MA, USA) and enzymatic digestion and ligation (TaKaRa, Japan) systems, according to the manufacturer's instructions. Gene disruption and GFP tagging were performed using the direct chromosomal integration method (Wach, 1996; Bähler et al., 1998). The integration cassettes were amplified using a two-step PCR. In the first PCR, fragments (~500 bp genomic regions upstream and downstream of the gene of interest) were amplified from the wild-type *S. pombe* genome using the KOD One DNA polymerase (TaKaRa Bio Inc., Japan). The first PCR products were used as primers in the second step of the PCR to amplify a cassette for integration, which contained a selection marker. For gene disruption, pFA6-kanMX6 (Bähler et al., 1998), pCR2.1-hph, pCR2.1-bsd, and pCR2.1-nat (Sato et al., 2005) were used to generate the integration cassettes. For GFP tagging, pFA6-GFP(S65T)-kanMX6 (Bähler et al., 1998) was used to generate the GFP-containing integration cassettes. The cassettes were then transformed into *S. pombe* cells and the desired integrant strains were selected on YES plates containing the appropriate selection agent. Correct disruptions and

integrations were confirmed by genomic PCR at the 5′ and 3′ ends using KOD FX Neo

(TaKaRa Bio Inc., Japan).

Fluorescence microscopy

Fluorescence microscopy data of intracellular GFP fusion proteins in living cells were

collected using a DeltaVision microscope system (GE Healthcare Inc., USA) equipped with a

pco.edge 4.2 sCMOS camera (PCO, Germany) and a ×60 PlanApo N OSC oil-immersion

objective lens (numerical aperture NA=1.4, Olympus, Japan). For live-cell imaging, cells

were precultured from a fresh colony, followed by culturing in YES medium at 30°C to reach

the logarithmic growth phase ($5\text{-}10\times10^6$ cells/ml). The cells were then attached to glass-

bottomed culture dishes coated with soybean lectin (Sigma-Aldrich, USA), covered with

Edinburgh minimal medium with glutamate (EMMG) medium, and observed at specified

temperatures.

Image quantification and processing

An identical fluorescence intensity range was applied to all images presented in each panel

using the Fiji software (Schindelin et al., 2012).

To quantify the fluorescence intensity in the nucleus, we used Fiji software. First, we

generated the sum projection image from the raw z-stack images and then subtracted the

background signal from the entire image by measuring the mean intensity of a region where

no cells exist. Second, the nuclear region was manually or semi-automatically selected. For

semi-automatic selection, individual nuclei were selected (original image), and the selected

image was blurred using Gaussian blur (blurred image). An appropriate sigma value was

applied depending on the image contrast. By subtracting the blurred image from the original

image, an edge-enhanced image was generated. The nuclear region was thresholded using the

appropriate minimum and maximum intensity ranges. Third, the threshold region was applied

to the original image and the total fluorescence intensity of the region was quantified. The

nuclear intensities obtained from *bqt3*[+] cells were averaged and the values were used to

calculate the relative fluorescence intensity. The relative fluorescence intensities were

plotted, and the results are presented as the mean ± standard deviation.

For the roundness index measurement, after selecting individual nuclei, the nuclear

membrane signal was thresholded using the appropriate fluorescence intensity ranges. The

nuclear region was filled in, and the roundness index was measured using the built-in

function in Fiji.

Western blotting

Cells were precultured in YES medium at 30°C overnight, inoculated in fresh YES, and then

cultured until the cell concentration reached the mid-log phase ($5\text{-}10\times10^6$ cells/ml).

Approximately 5×10^7 cells were harvested by centrifugation at 4°C, and the cell pellet was

washed with 1 ml ice-cold water containing 1 mM phenylmethylsulfonyl fluoride (PMSF).

Cells were lysed in 150 μl of ice-cold 1.85 M NaOH containing 7.5% β-mercaptoethanol (β-

ME) on ice for 15 min, followed by the addition of 150 μl of ice-cold 55% (v/v)

trichloroacetic acid (TCA) to precipitate the proteins, incubation on ice for 15 min, and

centrifugation (10 min, 14,000 ×g, 4°C). The protein pellet was then washed twice with ice-

cold acetone and dissolved in sample solubilization buffer [50 mM Tris-HCl pH 8.0, 1%

SDS, 1× Protease Inhibitor Cocktail (P8215, Sigma-Aldrich, USA), 1 mM PMSF, 1 mM

bortezomib]. Total protein concentrations were determined using the BCA assay (Thermo

Fisher Scientific, USA) according to the manufacturer's protocol. The samples were further

diluted in 2× Laemmli SDS sample buffer to a concentration of 1 μg/μl. After the addition of

dithiothreitol to a concentration of 50 mM, samples were heated at 95°C for 10 min before

immunoblotting. Identical amounts of protein (10 μg) were separated by SDS–PAGE and

transferred to a polyvinylidene difluoride membrane. The membranes were blocked for 1 h at

room temperature (approximately 26-28°C) in blocking solution PBST (10 mM Na_2HPO_4,

137 mM NaCl, 2.7 mM KCl, 1.76 mM KH_2PO_4, and 0.1% Tween 20) containing 5% skim

milk (Nacalai Tesque, Japan). To detect the GFP fusion proteins and β-actin as a loading

control, the membranes were probed with anti-GFP rabbit polyclonal antibody (Rockland,

USA; 1:2,000 dilution) or anti-β-actin mouse monoclonal antibody (Abcam, UK; 1:2,000

dilution) in blocking solution overnight at 4°C. After three rounds of washing with PBST, the

membranes were incubated with the appropriate horseradish peroxidase-conjugated

secondary antibodies (1:2,000 dilution, Cytiva, USA) in a blocking solution at room

temperature for 2 h. After three rounds of washing with PBST, immunoreactive bands were

detected using a chemiluminescence system (ImmunoStar LD, Wako, Japan) in a ChemiDoc

imaging system (Bio-Rad, USA).

Proteasome inhibition

Cells were grown in YES medium to the early mid-log phase ($2\text{-}5\times10^6$ cells/ml), as described

above. The culture was then divided into two, and the proteasome inhibitor bortezomib (LC

Laboratories, USA) dissolved in dimethyl sulfoxide (DMSO) was added to one culture to a

final concentration of 1 mM, and the same amount of DMSO was added to the other culture.

Cells were incubated at 26°C for 4 h before harvesting for fluorescence microscopy or

immunoblotting.

Temperature sensitive mutant experiments

For the temperature-sensitive mutant experiments, cells harboring *ts* or *cs* mutations were

grown in YES medium to the early mid-log phase ($2\text{-}5\times10^6$ cells/ml) at 26°C and 33°C,

respectively. The culture temperature was shifted up to 36°C or down to 16°C, and then the

cells were incubated for 4 h and 15 h before harvesting, respectively.

Ubiquitination detection assay

The ubiquitination detection assay was performed as described previously (Boban et al.,

2014) with modifications. Cells were grown in YES medium at 30°C until the mid-log phase

(5-10×10^6 cells/ml), and approximately 5×10^8 cells were harvested by centrifugation at 4°C.

Where indicated, the proteasome inhibitor bortezomib was added to a final concentration of 1

mM for 4 h before the cells were collected. The same volume of DMSO was added to the

control samples. The cells were washed with cold water containing 1 mM PMSF and lysed in

250 μl of 1.85 N NaOH solution containing 7.5% β-ME for 15 min on ice. Subsequently, the

proteins were precipitated by adding 250 μl of 55% (v/v) TCA, followed by incubation for 15

min on ice. The proteins were pelleted by centrifugation (10 min, 14,000 ×g, 4°C) and

washed with 500 μl of 50 mM Tris before being dissolved in 200 μl denaturation solution [25

mM Tris-HCl pH 7.4, 2% SDS, 2 mM ethylenediaminetetraacetic acid (EDTA), 1× Protease

Inhibitor Cocktail (P8215, Sigma-Aldrich, USA), 1 mM PMSF, 1 mM bortezomib, 10 μM

PR-619 (SML0430, Sigma-Aldrich, USA), and 50 mM 2-chloroacetamide (032-09762, Fuji

film Wako, Japan)]. Total protein concentrations were determined using the BCA assay

(Thermo Fisher Scientific, USA) according to the manufacturer's protocol. The samples were

incubated at 65°C for 10 min and clarified by centrifugation (5 min, 15,000 ×g, room

temperature). Equal amounts of protein lysates were diluted five-fold with cold IP dilution

buffer (50 mM Tris-HCl pH 7.5, 1.2% Triton X-100, 2 mM EDTA, 100 mM NaCl, 1 ×

Protease Inhibitor Cocktail (P8215, Sigma-Aldrich, USA), 1 mM PMSF, 1 mM bortezomib,

10 μM PR-619, and 50 mM 2-chloroacetamide), incubated with gentle rotation at 4°C for 1 h,

and cleared by centrifugation at 14,000 ×g for 10 min. A portion (20 μl) of the sample (2%

input) was mixed with 2× Laemmli SDS sample buffer (20 µl) and frozen until analysis. The

remaining sample was mixed with antibody-coupled beads prepared from 2 µg anti-GFP

rabbit polyclonal antibody (Rockland) and 50 µl Protein-A-coupled Dynabeads (Life

Technologies, USA) according to the manufacturer's instructions. The samples were

immunoprecipitated at 4°C for 3 h with gentle rotation and pelleted using a magnet. The

pellets were then washed as follows: twice with washing solution A (50 mM Tris-HCl pH

7.4, 2 mM EDTA, 100 mM NaCl, 1% Triton X-100, 0.4% SDS), once with washing solution

B (50 mM Tris-HCl pH 7.4, 2 mM EDTA, 350 mM NaCl, 1% Triton X-100) including a 10-

min incubation at 4°C with gentle rotation, and finally once with washing solution A. To

elute the immunoprecipitated protein, 50 µl of 1× Laemmli SDS sample buffer was added to

the pellet, and the mixture was incubated for 10 min at 65°C. Ten microliters of each sample

were separated on SDS-PAGE gels (8%). For the analysis of immunoprecipitated GFP-Bqt4,

the eluate was diluted 1:20 with 1× Laemmli SDS sample buffer. Immunoblotting analysis

was performed using an anti-GFP mouse monoclonal antibody (JL8; 1:1000; TaKaRa Bio

Inc., Japan) and an anti-ubiquitin mouse monoclonal antibody (P4D1, 1:1000; Santa Cruz

Biotechnology, USA). Protein levels in the input samples were assessed using an anti-β-actin

antibody (1:2000; Abcam).

Membrane association assay

The membrane association assay was performed according to a previously published protocol

(Habeck et al., 2015) with slight modifications. *S. pombe* cells expressing GFP-Bqt4 and

control proteins (GFP-Bqt3 and GFP-ADEL) were grown to the mid-log phase and mixed at

a 1:1:1 ratio before cell harvesting by centrifugation at 4°C. The harvested cells (~$5×10^8$

cells) were washed with cold water containing 1 mM PMSF and resuspended in 500 µl cold

fractionation buffer (FB; 0.7 M sorbitol, 50 mM Tris-HCl pH 7.4, 1× Protease Inhibitor

Cocktail, 1mM PMSF, 1 mM bortezomib). Resuspended cells were lysed using a bead

shocker (2,700 rpm for 10 cycles of 60 s on and 60 s off; Yasui Kikai, Co., Japan), followed

by washing the beads three times with 200 μl FB. The cell extracts were combined and debris

was removed by centrifugation (600 ×g for 5 min at 4°C). The lysate was divided into five

separate aliquots that were subjected to one of the following treatments: (1) FB alone; (2) 0.1

M Na_2CO_3, pH 11; (3) 2.5 M urea; (4) 0.5 M NaCl; or (5) 1% Triton X-100 and 0.5 M NaCl.

After incubation for 1 h on ice with occasional mixing, the samples were separated into pellet

(P) and supernatant (S) fractions by ultracentrifugation at 50,000 ×g for 45 min at 4°C. The

pellets were washed once with appropriate supplemented FB by centrifugation (50,000 ×g for

45 min at 4°C) before being resuspended in FB. Proteins were then precipitated from the

pellet and supernatant fractions by adding 150 μl of 55% (v/v) TCA. After 15 min of

incubation on ice, the proteins were pelleted by centrifugation (16,000 ×g for 10 min at 4°C),

washed twice with ice-cold acetone, and dissolved in dye-free 2× Laemmli SDS sample

buffer. Total protein concentrations were measured using the BCA assay, as described above.

The samples were incubated for 10 min at 95°C before immunoblotting.

Spot assay

Cells expressing GFP-Bqt4 under the control (*bqt4p*) or nmt1 promoter (*nmt1p*) were

precultured in YES medium overnight. Five-fold dilution series of the cells (starting

concentration was $8.0×10^5$ /mL) were spotted on the EMMG medium plates supplemented

with or without 10 μM thiamine, and then the plates were incubated at 30°C for 4 days.

For the temperature-sensitivity growth assay, wild-type and Cdc48 mutant cells were

spotted in a five-fold dilution series and incubated at 16, 20, 26, 30, 33 and 37°C for 3-6

days.

Acknowledgements

We thank Chizuru Ohtsuki and Yoshino Kubota for their technical assistance; We also thank

to Professor Mitsuhiro Yanagida (OIST) and the National BioResource Project Yeast in Japan

for providing the *S. pombe* strains.

Competing interests

The authors declare no competing financial or non-financial interests.

Funding

This study was supported by JSPS KAKENHI Grant Numbers JP20H05891 (to Y. Hirano),

JP19H03222 and JP20H05324 (to K.O.), JP18H05528 (to T.H.), and JP18H05533,

JP19K22389, and JP20H00454 (to Y. Hiraoka)).

Data availability

Further information and requests for resources and reagents should be directed to Yasushi

Hiraoka (hiraoka@fbs.osaka-u.ac.jp)

References

Asakawa, H., Hirano, Y., Shindo, T., Haraguchi, T., Hiraoka, Y. (2022). Fission yeast Ish1
and Les1 interact with each other in the lumen of the nuclear envelope. *Genes Cells* **27**,
643–656. doi: 10.1111/gtc.12981

Barbosa, A.D., Sembongi, H., Su, W.-M., Abreu, S., Reggiori, F., Carman, G.M.,
Siniossoglou, S. (2015). Lipid partitioning at the nuclear envelope controls membrane
biogenesis. *Mol. Biol. Cell* **26**, 3641–3657. doi: 10.1091/mbc.E15-03-0173

Bähler, J., J.Q. Wu, M.S. Longtine, N.G. Shah, A. McKenzie, A.B. Steever, A. Wach, P.
Philippsen, and J.R. Pringle. (1998). Heterologous modules for efficient and versatile
PCR-based gene targeting in *Schizosaccharomyces pombe*. *Yeast*. 14, 943-951. doi:
10.1002/(SICI)1097-0061(199807)14:10<943::AID-YEA292>3.0.CO;2-Y

Boban, M., Pantazopoulou, M., Schick, A., Ljungdahl, P.O., Foisner, R. (2014). A nuclear
ubiquitin-proteasomal pathway targets inner nuclear membrane protein Asi2 for
degradation. *J. Cell Sci.* **127**, 3603-3613. doi: 10.1242/jcs.153163

Bodnar N.O., Rapoport T.A. (2017). Molecular Mechanism of Substrate Processing by the
Cdc48 ATPase Complex. *Cell* **169**, 722-735. doi: 10.1016/j.cell.2017.04.020

Braun, S., Matuschewski, K., Rape, M., Thoms, S., Jentsch S. (2002). Role of the ubiquitin-
selective CDC48$^{UFD1/NPL4}$ chaperone (segregase) in ERAD of OLE1 and other substrates.
EMBO J. **15**, 615-621. doi: 10.1093/emboj/21.4.615

Buchwalter, A., Schulte, R., Tsai, H., Capitanio, J., Hetzer, M. (2019). Selective clearance of
the inner nuclear membrane protein emerin by vesicular transport during ER stress. *eLife*
8, e49796. doi: 10.7554/eLife.49796

Chen, C.-Y., Chi, Y.-H., Mutalif, R.A., Starost, M.F., Myers, T.G., Anderson, S.A., Stewart,
C.L., Jeang, K.-T. (2012). Accumulation of the Inner Nuclear Envelope Protein Sun1 Is
Pathogenic in Progeric and Dystrophic Laminopathies. *Cell* **149**, 565–577. doi:
10.1016/j.cell.2012.01.059

Chikashige, Y., Tsutsumi, C., Yamane, M., Okamasa, K., Haraguchi, T., Hiraoka, Y., 2006.
Meiotic Proteins Bqt1 and Bqt2 Tether Telomeres to Form the Bouquet Arrangement of
Chromosomes. Cell 125, 59–69. doi: /10.1016/j.cell.2006.01.048

Chikashige, Y., Yamane, M., Okamasa, K., Tsutsumi, C., Kojidani, T., Sato, M., Haraguchi,
T., Hiraoka, Y. (2009). Membrane proteins Bqt3 and -4 anchor telomeres to the nuclear
envelope to ensure chromosomal bouquet formation. *J. Cell Biol.* **187**, 413–427. doi:
10.1083/jcb.200902122

Christianson, J.C., Carvalho, P. (2022). Order through destruction: how ER-associated
protein degradation contributes to organelle homeostasis. *EMBO J.* **41**, e109845 doi:
10.15252/embj.2021109845

Deng, M., Hochstrasser, M. (2006). Spatially regulated ubiquitin ligation by an ER/nuclear
membrane ligase. *Nature* **443**, 827–831. doi: 10.1038/nature05170

Enam, C., Geffen, Y., Ravid, T., Gardner, R.G. (2018). Protein Quality Control Degradation
in the Nucleus. *Annu. Rev. Biochem.* **87**, 725–749. doi: 10.1146/annurev-biochem-
062917-012730

Foresti, O., Rodriguez-Vaello, V., Funaya, C., Carvalho, P. (2014). Quality control of inner
nuclear membrane proteins by the Asi complex. *Science* **346**, 751–755. doi:
10.1126/science.1255638

Franić, D., Zubčić, K., Boban, M. (2021). Nuclear Ubiquitin-Proteasome Pathways in
Proteostasis Maintenance. *Biomolecules* **11**, 54. doi: 10.3390/biom11010054

Gonzalez, Y., Saito, A., Sazer, S. (2012). Fission yeast Lem2 and Man1 perform fundamental
functions of the animal cell nuclear lamina. *Nucleus* **3**, 60–76. doi: 10.4161/nucl.18824

Gordon, C., McGurk, G., Dillon, P., Rosen, C., Hastie, N.D. (1993). Defective mitosis due to
a mutation in the gene for a fission yeast 265 protease subunit. *Nature* **366**, 355-357. doi:
10.1038/366355a0

Gordon, C., McGurk, G., Wallace, M., Hastie, N.D., 1996. A Conditional Lethal Mutant in
the Fission Yeast 26 S Protease Subunit *mts3*[+] Is Defective in Metaphase to Anaphase
Transition. *J. Biol. Chem.* **271**, 5704–5711. doi: 10.1074/jbc.271.10.5704

Habeck, G., Ebner, F.A., Shimada-Kreft, H., Kreft, S.G. (2015). The yeast ERAD-C
ubiquitin ligase Doa10 recognizes an intramembrane degron. *J. Cell Biol.* **209**, 261–273.
doi: 10.1083/jcb.201408088

Hirano, Y., Asakawa, H., Sakuno, T., Haraguchi, T., Hiraoka, Y. (2020a). Nuclear Envelope
Proteins Modulating the Heterochromatin Formation and Functions in Fission Yeast.
Cells **9**, 1908. doi: 10.3390/cells9081908

Hirano, Y., Y. Kinugasa, H. Osakada, T. Shindo, Y. Kubota, S. Shibata, T. Haraguchi, and Y.
Hiraoka. (2020b). Lem2 and Lnp1 maintain the membrane boundary between the nuclear
envelope and endoplasmic reticulum. *Commun. Biol.* **3**, 276. doi: 10.1038/s42003-020-
0999-9

Hirano, Y., Kinugasa, Y., Asakawa, H., Chikashige, Y., Obuse, C., Haraguchi, T., Hiraoka,
Y. (2018). Lem2 is retained at the nuclear envelope through its interaction with Bqt4 in
fission yeast. *Genes Cells* **23**, 122–135. doi: 10.1111/gtc.12557

Hiraoka, Y., Maekawa, H., Asakawa, H., Chikashige, Y., Kojidani, T., Osakada, H.,
Matsuda, A., Haraguchi, T. (2011). Inner nuclear membrane protein Ima1 is dispensable
for intranuclear positioning of centromeres: *S. pombe* inner nuclear membrane proteins.
Genes Cells **16**, 1000–1011. doi: 10.1111/j.1365-2443.2011.01544.x

Huang, A., Tang, Y., Shi, X., Jia, M., Zhu, J., Yan, X., Chen, H., Gu, Y. (2020). Proximity
labeling proteomics reveals critical regulators for inner nuclear membrane protein
degradation in plants. *Nat. Commun.* **11**, 3284. doi: 10.1038/s41467-020-16744-1

Jarosch, E., Taxis, C., Volkwein, C., Bordallo, J., Finley, D., Wolf, D.H., Sommer, T. (2002).
Protein dislocation from the ER requires polyubiquitination and the AAA-ATPase
Cdc48. *Nat. Cell Biol.* **4**, 134–139. doi: 10.1038/ncb746

Khmelinskii, A., Blaszczak, E., Pantazopoulou, M., Fischer, B., Omnus, D.J., Le Dez, G., Brossard, A., Gunnarsson, A., Barry, J.D., Meurer, M., Kirrmaier, D., Boone, C., Huber, W., Rabut, G., Ljungdahl, P.O., Knop, M. (2014). Protein quality control at the inner nuclear membrane. *Nature* **516**, 410–413. doi: 10.1038/nature14096

Kinuagsa, Y., Hirano, Y., Sawai, M., Ohno, Y., Shindo, T., Asakawa, H., Chikashige, Y., Shibata, S., Kihara, A., Haraguchi, T., Hiraoka, Y. (2019). The very-long-chain fatty acid elongase Elo2 rescues lethal defects associated with loss of the nuclear barrier function. *J. Cell Sci.* jcs.229021. doi: 10.1242/jcs.229021

Koch, B., Yu, H.-G. (2019). Regulation of inner nuclear membrane associated protein degradation. *Nucleus* 10, 169–180. doi: /10.1080/19491034.2019.1644593

Koch, B.A., Jin, H., Tomko, R.J., Yu, H.-G., 2019. The anaphase-promoting complex regulates the degradation of the inner nuclear membrane protein Mps3. *J. Cell Biol.* **218**, 839–854. doi: 10.1083/jcb.201808024

Kohda, T.A., Tanaka, K., Konomi, M., Sato, M., Osumi, M., Yamamoto, M. (2007). Fission yeast autophagy induced by nitrogen starvation generates a nitrogen source that drives adaptation processes. *Genes Cells* **12**, 155–170. doi: 10.1111/j.1365-2443.2007.01041.x

Krshnan, L., Siu, W.S., Van de Weijer, M., Hayward, D., Guerrero, E.N., Gruneberg, U., Carvalho, P. (2022). Regulated degradation of the inner nuclear membrane protein SUN2 maintains nuclear envelope architecture and function. *eLife* **11**, e81573. doi: 10.7554/eLife.81573.

Lord, S.J., K.B. Velle, R.D. Mullins, and L.K. Fritz-Laylin. (2020). SuperPlots: Communicating reproducibility and variability in cell biology. *J. Cell Biol.* **219**, e202001064. doi: 10.1083/jcb.202001064

Mannino, P.J., Lusk, C.P. (2022). Quality control mechanisms that protect nuclear envelope identity and function. *J. Cell Biol.* **221**, e202205123. doi: 10.1083/jcb.202205123

Matsuyama, A., Arai, R., Yashiroda, Y., Shirai, A., Kamata, A., Sekido, S., Kobayashi, Y., Hashimoto, A., Hamamoto, M., Hiraoka, Y., Horinouchi, S., Yoshida, M., 2006. ORFeome cloning and global analysis of protein localization in the fission yeast Schizosaccharomyces pombe. *Nat. Biotechnol.* **24**, 841–847. doi: 10.1038/nbt1222

Mekhail, K., Moazed, D. (2010). The nuclear envelope in genome organization, expression and stability. *Nat. Rev. Mol. Cell Biol.* **11**, 317–328. doi: 10.1038/nrm2894

Nakamura, Y., Arai, A., Takebe, Y., Masuda, M. (2011). A chemical compound for controlled expression of nmt1-driven gene in the fission yeast *Schizosaccharomyces pombe*. *Anal. Biochem.* **412**, 159–164. doi: 10.1016/j.ab.2011.01.039

Natarajan, N., Foresti, O., Wendrich, K., Stein, A., Carvalho, P. (2020). Quality Control of Protein Complex Assembly by a Transmembrane Recognition Factor. *Mol. Cell* **77**, 108-119. doi: 10.1016/j.molcel.2019.10.003

Nielsen, S., Poulsen, E., Rebula, C., Hartmann-Petersen, R., 2014. Protein Quality Control in the Nucleus. *Biomolecules* **4**, 646–661. doi: 10.3390/biom4030646

Pantazopoulou, M., Boban, M., Foisner, R., Ljungdahl, P.O. (2016). Cdc48 and Ubx1 participate in an inner nuclear membrane associated degradation pathway that governs the turnover of Asi1. *J. Cell Sci.* **129**, 3770-3780. doi: 10.1242/jcs.189332

Pawar, S., Kutay, U. (2021). The Diverse Cellular Functions of Inner Nuclear Membrane Proteins. *Cold Spring Harb. Perspect. Biol.* **13**, a040477. doi: 10.1101/cshperspect.a040477

Rabinovich, E., Kerem, A., Fröhlich, K.-U., Diamant, N., Bar-Nun, S. (2002). AAA-ATPase p97/Cdc48p, a Cytosolic Chaperone Required for Endoplasmic Reticulum-Associated Protein Degradation. *Mol. Cell. Biol.* **22**, 626–634. doi: 10.1128/MCB.22.2.626-634.2002

Ravid, T., Kreft, S.G., Hochstrasser, M. (2006). Membrane and soluble substrates of the Doa10 ubiquitin ligase are degraded by distinct pathways. *EMBO J.* **25**, 533–543. doi: 10.1038/sj.emboj.7600946

Rossanese, O.W., Reinke, C.A., Bevis, B.J., Hammond, A.T., Sears, I.B., O'Connor, J., Glick, B.S. (2001). A Role for Actin, Cdc1p, and Myo2p in the Inheritance of Late Golgi Elements in *Saccharomyces cerevisiae*. *J. Cell Biol.* **153**, 47–62. doi: 10.1083/jcb.153.1.47

Ruggiano, A., Foresti, O., Carvalho, P. (2014). ER-associated degradation: Protein quality control and beyond. *J. Cell Biol.* **204**, 869–879. doi: 10.1083/jcb.201312042

Ruggiano, A., Mora, G., Carvalho, P. (2016). Spatial control of lipid droplet proteins by the ERAD ubiquitin ligase Doa10. *EMBO J.* **35**, 1644-1655. doi: 10.15252/embj.201593106

Santos-Rosa, H., Leung, J., Grimsey, N., Peak-Chew, S., Siniossoglou, S. (2005). The yeast lipin Smp2 couples phospholipid biosynthesis to nuclear membrane growth. *EMBO J.* **24**, 1931–1941. doi: 10.1038/sj.emboj.7600672

Sato, M., Dhut, S., Toda, T. (2005). New drug-resistant cassettes for gene disruption and epitope tagging in *Schizosaccharomyces pombe*. *Yeast.* **22**, 583-591. doi: 10.1002/yea.1233

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods.* **9**, 676-682. doi: 10.1038/nmeth.2019

Schuberth, C., Buchberger, A. (2005). Membrane-bound Ubx2 recruits Cdc48 to ubiquitin ligases and their substrates to ensure efficient ER-associated protein degradation. *Nat. Cell Biol.* **7**, 999-1006. doi: 10.1038/ncb1299.

Smoyer, C.J., Jaspersen, S.L. (2019). Patrolling the nucleus: inner nuclear membrane-associated degradation. *Curr. Genet.* **65**, 1099–1106. doi: 10.1007/s00294-019-00971-1

Smoyer, C.J., Smith, S.E., Gardner, J.M., McCroskey, S., Unruh, J.R., Jaspersen, S.L. (2019). Distribution of Proteins at the Inner Nuclear Membrane Is Regulated by the Asi1 E3 Ligase in *Saccharomyces cerevisiae*. *Genetics* **211**, 1269–1282. doi: 10.1534/genetics.119.301911

Steglich, B., Filion, G., van Steensel, B., Ekwall, K. (2012). The inner nuclear membrane
proteins Man1 and Ima1 link to two different types of chromatin at the nuclear periphery
in *S. pombe*. *Nucleus* **3**, 77–87. doi: 10.4161/nucl.18825

Swanson, R., Locher, M., Hochstrasser, M. (2001). A conserved ubiquitin ligase of the
nuclear envelope/endoplasmic reticulum that functions in both ER-associated and Matα2
repressor degradation. *Genes Dev.* **15**, 2660-2674. doi: 10.1101/gad.933301.

Takeda, K., Mori, A., Yanagida, M. (2011). Identification of Genes Affecting the Toxicity of
Anti-Cancer Drug Bortezomib by Genome-Wide Screening in S. pombe. *PLoS ONE* **6**,
e22021. doi: 10.1371/journal.pone.0022021

Takeda, K., Yanagida, M. (2005). Regulation of Nuclear Proteasome by Rhp6/Ubc2 through
Ubiquitination and Destruction of the Sensor and Anchor Cut8. *Cell* **122**, 393–405. doi:
10.1016/j.cell.2005.05.023

Tange, Y., Chikashige, Y., Takahata, S., Kawakami, K., Higashi, M., Mori, C., Kojidani, T.,
Hirano, Y., Asakawa, H., Murakami, Y., Haraguchi, T., Hiraoka, Y. (2016). Inner
nuclear membrane protein Lem2 augments heterochromatin formation in response to
nutritional conditions. *Genes Cells* **21**, 812–832. doi: 10.1111/gtc.12385

Tatebe, H., Yanagida, M. (2000). Cut8, essential for anaphase, controls localization of 26S
proteasome, facilitating destruction of cyclin and Cut2. *Curr. Biol.* **10**, 1329–1338. doi:
10.1016/S0960-9822(00)00773-9

Toyama, B.H., Savas, J.N., Park, S.K., Harris, M.S., Ingolia, N.T., Yates, J.R., Hetzer, M.W.
(2013). Identification of Long-Lived Proteins Reveals Exceptional Stability of Essential
Cellular Structures. *Cell* **154**, 971–982. doi: 10.1016/j.cell.2013.07.037

Tsai, P.-L., Zhao, C., Turner, E., Schlieker, C. (2016). The Lamin B receptor is essential for
cholesterol synthesis and perturbed by disease-causing mutations. *eLife* **5**, e16011. doi:
10.7554/eLife.16011

Vembar, S.S., Brodsky, J.L. (2008). One step at a time: endoplasmic reticulum-associated
degradation. *Nat. Rev. Mol. Cell Biol.* **9**, 944–957. doi: 10.1038/nrm2546

Wach, A. (1996). PCR-synthesis of marker cassettes with long flanking homology regions for
gene disruptions in *S. cerevisiae*. *Yeast*. **12**, 259-265. doi: 10.1002/(SICI)1097-
0061(19960315)12:3%3C259::AID-YEA901%3E3.0.CO;2-C

Wolf, D.H., Stolz, A. (2012). The Cdc48 machine in endoplasmic reticulum associated
protein degradation. *Biochim. Biophys. Acta.* **1823**, 117–124. doi:
10.1016/j.bbamcr.2011.09.002

Ye, Y., Meyer, H.H., Rapoport, T.A. (2001). The AAA ATPase Cdc48/p97 and its partners
transport proteins from the ER into the cytosol. *Nature* **414**, 652–656. doi:
10.1038/414652a

Figure legends

Figure 1. Bqt4 protein is degraded by the ubiquitin-proteasome system

(A, B) Cells expressing GFP-Bqt4 in the *bqt3$^+$* or *bqt3Δ* background were treated with vehicle (+DMSO) or the proteasome inhibitor bortezomib (+BZ) for 4 h and subjected to microscopic observation (A) or Western blotting (B). (A) Fluorescence microscopic images of GFP-Bqt4 (upper panels); bright-field images (lower panels). Bar, 10 μm. (B) Protein amounts of GFP-Bqt4 and β-actin were detected by anti-GFP and anti-β-actin antibodies, respectively. Molecular weight markers are shown on the left.

(C, D) Cells harboring *mts3-1* in the *bqt3$^+$* or *bqt3Δ* background were cultured at a nonpermissive temperature (36°C) for 4 h and subjected to microscopic observation (C) or Western blotting (D). (C) Fluorescence microscopic images of GFP-Bqt4 (upper panels); bright-field images (lower panels). Bar, 10 μm. (D) Protein amounts of GFP-Bqt4 and β-actin were detected as in (B). The asterisk in (D) indicates non-specific bands.

(E) Polyubiqutination of Bqt4 protein. Cells expressing GFP or GFP-Bqt4 in the *bqt3$^+$* or *bqt3Δ* background were treated with vehicle (+DMSO) or the proteasome inhibitor bortezomib (+BZ) for 4 h. After denaturing the proteins, GFP and GFP-Bqt4 were immunoprecipitated with anti-GFP antibody (IP). The precipitated proteins were detected by anti-ubiquitin or anti-GFP antibodies, respectively (IB). β-actin was used as a loading control. Molecular weight markers are shown on the left.

(F) No Bqt4 degradation by the lysosome-autophagy system. Deletion of *atg1* (*atg1Δ*) or *isp6* (*isp6Δ*) was introduced into the *bqt3Δ* cells. Protein amounts of GFP-Bqt4 and β-actin in these cells were detected as in (B). The asterisk indicates non-specific bands. Molecular weight markers are shown on the left.

Figure 2. Bqt4 degradation occurs in the nucleus

Cells harboring *cut8-655 ts* mutation in the *bqt3⁺* or *bqt3Δ* background were cultured at a

nonpermissive temperature (36°C) for 4 h and subjected to microscopic observation (A) or

Western blotting (B). (A) Fluorescence images of GFP-Bqt4 (upper panels); bright-field

images (lower panels). Bar, 10 μm. (B) Protein levels of GFP-Bqt4 and β-actin were detected

using anti-GFP and anti-β-actin antibodies, respectively. The asterisk indicates non-specific

bands. Molecular weight markers are shown on the left.

Figure 3. Bqt4 is a substrate of the Doa10 complex

Bqt4 protein levels in *doa10, ubc6* and *ubc7* mutants. (A, B) Cells harboring *hrd1* (*hrd1Δ*),

doa10 (*doa10Δ*) single deletion, or *hrd1* and *doa10* double deletion (*hrd1Δ doa10Δ*) in *bqt3Δ*

background were cultured at 30°C for 16 h, and then subjected to microscopic observation

(A) or Western blotting (B). (A) Fluorescence microscopic images of GFP-Bqt4 (upper

panels); bright-field images (lower panels). Bar, 10 μm. (B) Protein levels of GFP-Bqt4 and

β-actin were detected using anti-GFP and anti-β-actin antibodies, respectively. (C, D) Cells

harboring *ubc6* (*ubc6Δ*), *ubc7* (*ubc7Δ*) single deletion, or *ubc6* and *ubc7* double deletion

(*ubc6Δ ubc7Δ*) in *bqt3Δ* background were cultured at 30°C for 16 h, and then subjected to

microscopic observation (C) or Western blotting (D). (C) Fluorescence microscopic images

of GFP-Bqt4 (upper panels); bright-field images (lower panels). Bar, 10 μm. (D) Protein

levels of GFP-Bqt4 and β-actin were detected using anti-GFP and anti-β-actin antibodies,

respectively. The asterisk indicates non-specific bands. Molecular weight markers are shown

on the left.

Figure 4. Bqt4 is an integral membrane protein

Cells expressing GFP-Bqt4, GFP-Bqt3, or GFP-ADEL were mixed at a 1:1:1 ratio and lysed

by bead beating. After removing cell debris, the extracts were treated under various

conditions, as shown on the top. The extracts were fractionated by ultracentrifugation into

supernatant (S) and pellet (P), and then GFP-tagged proteins in each fraction were detected

by Western blotting. Molecular weight markers are shown on the left.

Figure 5. Removal of Bqt4 requires the Cdc48 retrotranslocation complex

(A) The Cdc48 mutant used in this study. Left panel: the amino acid sequence alignment

around the D1 and D2 domains of *S. pombe* (*Sp*) and *S. cerevisiae* (*Sc*) Cdc48. Gray and light

gray shades denote identical and similar amino acids, respectively (top and bottom). The

amino acids indicated by red (P267 and A550; correspond to red stars in the schematic

diagram in the middle) are both mutated in this study (P267L/A550T). Right panel: the

mutation sites are shown in the predicted three-dimensional structure of Cdc48 by

AlphaFold2 (https://alphafold.ebi.ac.uk/entry/Q9P3A7).

(B) Temperature sensitivity of *cdc48-P267L/A550T* mutant. Five-fold serially diluted cells

harboring *cdc48-P267L/A550T* mutant were spotted on the YES plate, and cultured at

different temperatures as indicated at the bottom.

(C, D) Effect of hypomorphic mutation of *cdc48* on Bqt4 degradation. Cells harboring the

cdc48-P267L/A550T mutant in the *bqt3⁺* or *bqt3Δ* background were cultured at a

nonpermissive temperature (16°C) for 15 h and subjected to microscopic observation (C) or

Western blotting (D). (C) Left panels: Fluorescence microscopic images of GFP-Bqt4 (upper

panels); bright-field images (lower panels). Bar: 10 μm. Right panel: the fluorescence

intensities in the nuclei were quantified and the relative values were plotted. The mean values

are plotted with standard deviation on the right panel. The number of the cells analyzed is

shown at the bottom. ****$p < 0.0001$ via two-tailed unpaired student's *t*-test. (D) Protein

amounts of GFP-Bqt4 and β-actin were detected by anti-GFP and anti-β-actin antibodies,

respectively. Molecular weight markers are shown on the left.

Figure 6. The C-terminal sequence of Bqt4 is a degron

The C-terminal fragments of Bqt4 are shown as a cartoon on the left. These fragments were

expressed as GFP-fused proteins in the $bqt3^+$ or $bqt3\Delta$ cells. The cells were treated with

vehicle (+DMSO) or the proteasome inhibitor bortezomib (+BZ) for 4 h and then observed

by fluorescence microscopy. Bars: 10 μm.

Figure 7. Excess Bqt4 causes nuclear envelope deformation

(A) Left panels: Fluorescence microscopic images of Ish1-GFP in $bqt4^+$ or $bqt4\Delta$ cells. Ish1-

GFP was used as a nuclear membrane marker (Asakawa et al., 2022). The cells were treated

with vehicle (+DMSO) or the proteasome inhibitor bortezomib (+BZ) for 4 h. The

images were deconvoluted using the SoftWoRx software (v7.0.0). Bar: 10 μm. Right panel:

The roundness index of the nucleus in these cells was measured from three independent

experiments (shown in blue, green, and red, respectively) and plotted as a superplot (Lord et

al., 2020) (right panel). The roundness index one indicates a perfect circle. The solid blue

circle, green triangle, and red square represent the mean value of the measurements in each

experiment, respectively. Black horizontal lines represent the mean value for all data that is

shown with standard deviation. *n* at the bottom right represents the total number of cells

analyzed. ****$p < 0.0001$ via two-tailed unpaired student's *t*-test.

(B) Fluorescence microscopic images of GFP-Bqt4 in the $bqt3^+$ or $bqt3\Delta$ cells. GFP-Bqt4

was overproduced under the *nmt1* promoter in the presence or absence of 2.5 μM YAM2 in

EMMG for 20 and 18 h, respectively. For comparison, *bqt4p*-GFP-Bqt4 fluorescence images

of cells treated with BZ (*bqt4p*+BZ) are reproduced in Figure. 1A with different display

scales. Bar: 10 μm.

(C) Growth defects by Bqt4 overproduction. GFP-Bqt4 was overproduced under *bqt4* (*bqt4p*)

or *nmt1* promoter (*nmt1p*) in the *bqt3⁺* or *bqt3Δ* cells. Five-fold serially diluted cells were

spotted on the EMMG plate in the presence or absence of thiamine for 4 days. WT indicates

wildtype.

Figure 8. A model of INM protein degradation of Bqt4

Excess Bqt4, with exposed TMD as a degradation signal, is recognized and ubiquitinated by

INM-associated E3 ubiquitin ligase complexes, including Doa10 and additional unidentified

E3 ligases. Ubiquitinated Bqt4 is then actively dislocated from the INM by the AAA-ATPase

Cdc48 complex and transferred to the nuclear proteasome for proteolysis.

Figure 1

Figure 2

Figure 3

A

$bqt3\Delta$

$bqt3^+$ - $hrd1\Delta$ $doa10\Delta$ $doa10\Delta\ hrd1\Delta$

GFP-Bqt4

BF

B

$bqt3\Delta$

$bqt3^+$ - $hrd1\Delta$ $doa10\Delta$ $doa10\Delta\ hrd1\Delta$

(kDa)

＊

75 —   ◄ GFP-Bqt4

50 —   ◄ β-actin

C

$bqt3\Delta$

$bqt3^+$ - $ubc6\Delta$ $ubc7\Delta$ $ubc6\Delta\ ubc7\Delta$

GFP-Bqt4

BF

D

$bqt3\Delta$

$bqt3^+$ - $ubc6\Delta$ $ubc7\Delta$ $ubc6\Delta\ ubc7\Delta$

＊

75 —   ◄ GFP-Bqt4

50 —   ◄ β-actin

Figure 4

Figure 5

Figure 6

Figure 7

A Ish1-GFP

bqt4+   bqt4Δ

+ DMSO

+ BZ

Roundness index

bqt4	+	+	Δ	Δ
BZ	−	+	−	+
	150	125	113	70 (=n)

B GFP-Bqt4

bqt4p +BZ   nmt1p+YAM2

2.5 µM   0 µM

bqt3+

bqt3Δ

0-3000   0-3000   0-10000

Display scale

C

+ thiamine   −thiamine (OP)

WT

GFP-Bqt4

nmt1p bqt4p

bqt3+

bqt3Δ

bqt3+

bqt3Δ

Figure 8

Supplementary Table 1. *S. pombe* strains used in this study

Strain	Genotype	Source	Figure
TL204	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4*	This study	1A-F, 2A, B, 3A-D, 4, 5C, D 7B, C, S1, S4
TL205	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 bqt3Δ::NAT*	This study	1A-F, 2A, B, 3A-D, 4, 5C, D, 7B, C, S1, 2, 4
TL288	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 mts3-1*	This study	1C, D
TL289	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 mts3-1*	This study	1C, D
H1N95	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP*	This study	1E
TL233	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 isp6Δ::bsd*	This study	1F
TL234	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 atg1Δ:: kanʳ*	This study	1F
TL290	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 cut8-563*	This study	2A, B
TL291	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 cut8-563*	This study	2A, B
TL215	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 doa10Δ::bsd*	This study	3A, B, S2
TL219	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 hrd1Δ:: kanʳ*	This study	3A, B, S2
TL236	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 doa10Δ::bsd hrd1Δ:: kanʳ*	This study	3A, B
TL220	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ubc7Δ:: kanʳ*	This study	3C, D
TL221	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ubc6Δ:: kanʳ*	This study	3C, D
TL266	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ubc6Δ:: kanʳ ubc7Δ::bsd*	This study	3C, D
HA1382-3D	*h⁻ lys1-131 ura4-D18 leu1-32 bqt3Δ::LEU2 aurlʳ::bqt3p-GFP-bqt3*	This study	4
H1N789	*h⁻ lys1+::nmt41p-SS-GFP-ADEL*	This study	4
h-972	*h⁻*	Laboratory stock	5B, 7C
TL311	*h⁻ lys1-131 cdc48Δ:: kanʳ aurlʳ::cdc48p-cdc48-P267L/A550T*	This study	5B
TL306	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 cdc48Δ:: kanʳ aurlʳ::cdc48p-cdc48-P267L/A550T*	This study	5C, D, S4
TL307	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 bqt3Δ::NAT cdc48Δ::kanʳ aurlʳ::cdc48p-cdc48-P267L/A550T*	This study	5C, D, S4
TL242	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4C³⁶⁹⁻⁴³²*	This study	6
TL243	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4C³⁶⁹⁻⁴³²*	This study	6
TL246	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4C³⁹⁴⁻⁴³²*	This study	6
TL247	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4C³⁹⁴⁻⁴³²*	This study	6
TL248	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4C⁴¹⁴⁻⁴³²*	This study	6
TL249	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4C⁴¹⁴⁻⁴³²*	This study	6
TL250	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4C³⁶⁹⁻⁴²⁵*	This study	6
TL251	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4C³⁶⁹⁻⁴²⁵*	This study	6
TL252	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4C⁴¹⁴⁻⁴²⁵*	This study	6
TL253	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4C⁴¹⁴⁻⁴²⁵*	This study	6
H1N429	*h⁻ lys1⁺::ish1-GFP*	This study	7A
H1N546	*h⁺ bqt4Δ::hph lys1⁺::ish1-GFP*	This study	7A
TL206	*h⁻ bqt4Δ::hph lys1⁺::nmt1p-GFP-bqt4*	This study	7B, C
TL207	*h⁻ bqt4Δ::hph lys1⁺::nmt1p-GFP-bqt4 bqt3Δ::NAT*	This study	7B, C
TL286	*h⁻ bqt4Δ::hph lys1⁺::bqt4p-GFP-bqt4 mts2-1*	This study	S1
TL287	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 mts2-1*	This study	S1
TL222	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 meu34Δ:: kanʳ*	This study	S2
TL224	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 san1Δ:: kanʳ*	This study	S2
TL225	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ubr1Δ:: kanʳ*	This study	S2
TL226	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ubr11Δ:: kanʳ*	This study	S2
TL227	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 hul5Δ:: kanʳ*	This study	S2
TL267	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 dsc1Δ:: kanʳ*	This study	S2
TL292	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 cut9-665*	This study	S2
TL294	*h⁻ bqt4Δ::hph bqt3Δ::NAT lys1⁺::bqt4p-GFP-bqt4 ptr1-1*	This study	S2

Supplementary Figure S1. The *mts2-1* mutation prevents Bqt4 degradation

Cells harboring *mts2-1* in the *bqt3*⁺ or *bqt3Δ* background were cultured at a nonpermissive temperature (36°C) for 4 h and then observed by fluorescence microscopy. (Upper panels) Fluorescence images of GFP-Bqt4 (lower panels) bright-field images (BF). Bar: 10 μm.

Supplementary Figure S2. Most nuclear- or ER-associated E3 ligases have a minimal role in Bqt4 degradation

Fluorescence images of GFP-Bqt4 in mutants are shown. Deletion or mutation of E3 ligases localized in the ER (*meu34Δ*, *dsc1Δ*) or functioning in the nucleus (*san1Δ*, *ubr1Δ*, *ubr11Δ*, *hul5Δ*, *ptr1-1*, *cut9-665*) was introduced in *bqt3Δ* cells expressing GFP-Bqt4. The cells were cultured at 30°C for 16 h or shifted to 36°C for 4 h for *ts* mutants and then observed using fluorescence microscopy. Bars: 10 μm.

Supplementary Figure S3. Prediction of Bqt4 TMD properties

The transmembrane helices of Bqt4 were predicted by the TMHMM-2.0 prediction server (https://services.healthtech.dtu.dk/service.php?TMHMM-2.0). The horizontal and vertical axes indicate the prediction probability and amino acid residues of Bqt4, respectively.

Supplementary Figure S4. Degradation of Bqt4 is not significantly affected in Cdc48 mutant at the permissive temperature

Cells harboring the cdc48-P267L/A550T mutant in the *bqt3⁺* or *bqt3Δ* background were cultured at 33°C for 15 h and subjected to microscopic observation (A) or Western blotting (B). (A) Fluorescence microscopic images of GFP-Bqt4 (upper panels); bright-field images (lower panels). Bar: 10 μm. (B) Protein amounts of GFP-Bqt4 and β-actin were detected by anti-GFP and anti-β-actin antibodies, respectively. Molecular weight markers are shown on the left.

Supplementary Figure S5. Structure of Bqt4 as predicted by AlphaFold2

The three-dimensional structure of Bqt4 predicted by AlphaFold2 was obtained from the website (https://alphafold.ebi.ac.uk/entry/O60158).