The deubiquitinase Ubp3/Usp10 constrains glucose-mediated mitochondrial

repression via phosphate budgeting

Vineeth Vengayil[1,2], Shreyas Niphadkar[1,2], Swagata Adhikary[1,2], Sriram Varahan[1], Sunil

Laxman[1]*

1. Institute for Stem Cell Science and Regenerative Medicine (DBT-inStem)

GKVK Post Bellary Road

Bangalore 560065

2. Manipal Academy of Higher Education, Manipal, Karnataka 576104, India

*correspondence: sunil@instem.res.in

Keywords: mitochondria, glycolysis, phosphate, deubiquitinase, Crabtree

Abstract

Many cells in high glucose repress mitochondrial respiration, as observed in the Crabtree and

Warburg effects. Our understanding of biochemical constraints for mitochondrial activation

is limited. Using a *Saccharomyces cerevisiae* screen, we identified the conserved

deubiquitinase Ubp3 (Usp10), as necessary for mitochondrial repression. Ubp3 mutants have

increased mitochondrial activity despite abundant glucose, along with decreased glycolytic

enzymes, and a rewired glucose metabolic network with increased trehalose production.

Utilizing *Δubp3* cells, along with orthogonal approaches, we establish that the high glycolytic

flux in glucose continuously consumes free Pi. This restricts mitochondrial access to inorganic

phosphate (Pi), and prevents mitochondrial activation. Contrastingly, rewired glucose

metabolism with enhanced trehalose production and reduced GAPDH (as in *Δubp3* cells)

restores Pi. This collectively results in increased mitochondrial Pi and derepression, while

restricting mitochondrial Pi transport prevents activation. We therefore suggest that

glycolytic-flux dependent intracellular Pi budgeting is a key constraint for mitochondrial

repression.

Introduction

Rapidly proliferating cells have substantial metabolic and energy demands in order to increase biomass [1, 2]. This includes a high ATP demand, obtained from cytosolic glycolysis or mitochondrial oxidative phosphorylation (OXPHOS), to fuel multiple reactions [3]. Interestingly, many rapidly proliferating cells preferentially rely on ATP from glycolysis/fermentation over mitochondrial respiration even in oxygen-replete conditions, and is the well-known Warburg effect [4, 5]. Many such cells repress mitochondrial processes in high glucose, termed glucose-mediated mitochondrial repression or the Crabtree effect [6, 7]. This is observed in some tumors [5], neutrophils [8], activated macrophages [9], stem cells [10–12], and famously *Saccharomyces cerevisiae* [7]. Numerous studies have identified signaling programs or regulators of glucose-mediated mitochondrial repression. However, biochemical programs and regulatory processes in biology evolve around key biochemical constraints [13]. The biochemical constraints for mitochondrial repression remain unresolved [14, 15].

There are two hypotheses on the biochemical principles driving mitochondrial repression. The first proposes direct roles for glycolytic intermediates in driving mitochondrial respiration, by inhibiting specific mitochondrial outputs [16, 17]. The second hypothesizes that a competition between glycolytic and mitochondrial processes for mutually required metabolites/co-factors such as pyruvate, ADP or inorganic phosphate (Pi) could determine the extent of mitochondrial repression [14, 15, 18]. These are not all mutually exclusive, and a combination of these factors might dictate mitochondrial repression in high glucose. However, any hierarchies of importance are unclear [19], and experimental data for the necessary constraints for mitochondrial repression remains incomplete.

One approach to resolve this question has been to identify regulators of metabolic state under high glucose. Post-translational modifications (PTMs) regulated by signaling systems can regulate mitochondrial repression [20–22]. Ubiquitination is a PTM that regulates global proteostasis [23, 24], but the roles of ubiquitination-dependent processes in regulating mitochondrial repression are poorly explored. Ubiquitination itself is determined by the balance between ubiquitination, and deubiquitinase (DUB) dependent deubiquitination [25]. Little is known about the roles of DUBs in regulating metabolic states, making the DUBs interesting candidate regulators of mitochondrial repression.

In this study, using an *S. cerevisiae* DUB knock-out library-based screen, we identified the evolutionarily conserved deubiquitinase Ubp3 (mammalian Usp10) as required for mitochondrial repression in high glucose. Loss of Ubp3 resulted in mitochondrial activation, along with a reduction in the glycolytic enzymes-phosphofructokinase 1 (Pfk1) and GAPDH (Tdh2 and Tdh3). This consequently reroutes glucose flux and increases trehalose biosynthesis. This metabolic rewiring increases Pi release from trehalose synthesis, and decreases Pi consumption in glycolysis, to cumulatively increase Pi pools. Using *ubp3Δ* cells along with independent analysis of wild-type cells, and isolated mitochondrial fractions, we establish that glycolytic-flux dependent Pi allocations to mitochondria determines mitochondrial activity. Through these data, we propose how intracellular Pi balance as controlled by glycolytic flux, is a biochemical constraint for mitochondrial repression.

Results

A deubiquitinase deletion screen identifies Ubp3 as a regulator of glucose-mediated mitochondrial repression

In cells such as *S. cerevisiae*, high glucose represses mitochondrial activity as well as OXPHOS-dependent ATP synthesis (7, 26) (illustrated in Figure 1A). Our initial objective was to identify proteostasis regulators of glucose-mediated mitochondrial repression. We generated and used a deubiquitinase (DUB) deletion strain library of *S. cerevisiae* (Figure S1A), to unbiasedly identify regulators of mitochondrial repression by measuring the fluorescence intensity of a potentiometric dye Mitotracker CMXRos (illustrated in Figure 1B). Using this screen, we identified DUB mutants with altered mitochondrial membrane potential (Figure 1C, Figure S1A). Note: WT cells in a respiratory medium (2% ethanol) were used as a control to estimate maximum mitotracker fluorescence intensity (Figure S1B).

A prominent 'hit' was the evolutionarily conserved deubiquitinase Ubp3 (Figure 1C), (homologous to mammalian Usp10) (Figure S1C). Due to its high degree of conservation across eukaryotes (Figure S1C) as well as putative roles in metabolism or mitochondrial function (27–29), we focused our further attention on this DUB. Cells lacking Ubp3 showed an ~1.5-fold increase in mitotracker fluorescence (Figure 1C, Figure S1A). Cells with catalyticaly inactive Ubp3 (Ubp3^{C469A}), showed increased mitochondrial potential comparable

to *ubp3Δ* (Figure 1D). This data confirmed that Ubp3 catalytic activity of is required to fully

repress mitochondrial activity under high glucose. The catalytic site mutation did not affect

steady-state Ubp3 levels (Figure S1D).

Next, to assess the requirement of Ubp3 for mitochondrial function, we quantified the

electron transport chain (ETC) complex IV subunit Cox2 (30)(Figure 1E). *ubp3Δ* had higher

Cox2 than WT (Figure 1E). There was no increase in mitochondrial outer membrane protein

Tom70 (Figure S1E), suggesting that the increased Cox2 is not merely because of higher total

mitochondrial content. We next measured the basal oxygen consumption rate (OCR) of

ubp3Δ, and basal OCR was higher in *ubp3Δ*, indicating higher respiration (Figure 1F).

Next, we asked if mitochondrial ATP synthesis was higher in *ubp3Δ*. The total ATP

levels in WT and *ubp3Δ* were comparable (Figure 1G). However, upon treatment with the ETC

complex IV inhibitor sodium azide, ATP levels in WT were higher than *ubp3Δ*, contributing to

~60% of the total ATP (Figure 1H). In contrast, the ATP levels in *ubp3Δ* after sodium azide

treatment were ~ 40% of the total ATP (Figure 1H). These data suggest a higher contribution

of mitochondrial ATP synthesis towards the total ATP pool in *ubp3Δ*.

We next asked if *ubp3Δ* required higher mitochondrial activity for growth. In high

glucose, WT show minimal growth inhibition in the presence of sodium azide, indicating lower

reliance on mitochondrial function (Figure 1I). Contrastingly, *ubp3Δ* or Ubp3^{C469A} show a

severe growth defect in the presence of sodium azide (Figure 1I). Deletion of ETC components

Cox2, and the ATP synthase subunits Atp1 and Atp10 results in a severe growth defect in

ubp3Δ compared to WT (Figure S1F). Together, these results indicate that the loss of Ubp3

makes cells dependent on mitochondrial ATP synthesis in high glucose.

Collectively these data show that in 2% glucose, *ubp3Δ* have high mitochondrial

activity, respiration, and rely on this mitochondrial function for ATP production and growth.

We therefore decided to use *ubp3Δ* cells to start delineating requirements for glucose-

mediated mitochondrial repression.

Key glycolytic enzymes decrease and glucose flux is rerouted in *ubp3Δ* cells

Glucose-6 phosphate (G6P) is the central node in glucose metabolism where carbon-
allocations are made towards distinct metabolic arms, primarily: glycolysis, the pentose
phosphate pathway (PPP), and trehalose biosynthesis (Figure 2A). We first compared
amounts of two key ('rate-controling') glycolytic enzymes - Phosphofructokinase 1 (Pfk1),
GAPDH isozymes (Tdh2, Tdh3) (3), along with the enolase isozymes (Eno1, Eno2) in WT and
ubp3Δ cells. Pfk1, Tdh2, and Tdh3 substantially decreased in *ubp3Δ* (but not Eno1 and Eno2)
(Figure 2B, S2A). Since deubiquitinases can control protein amounts by regulating
proteasomal degradation, we asked if the decrease in Pfk1, Tdh2 and Tdh3 in *ubp3Δ* is due to
proteasomal degradation. To test this, we measured the levels of these enzymes in *ubp3Δ*
after treatement with proteasomal inhibitor MG132. We did not observe any rescue in the
levels of these enzymes in MG132 treated samples, suggesting that the decreased levels of
these enzymes were not due to increased proteasomal degradation (Figure S2B). To further
understand if these enzyme transcripts are altered in *ubp3Δ*, we measured the mRNA levels
of PFK1, TDH2 and TDH3 in WT, *ubp3Δ* and Ubp3[C469A] cells. The transcripts of all the three
genes in *ubp3Δ* and Ubp3[C469A] cells decreased (Figure S2C), suggesting that Ubp3 regulates
the transcripts of these glycolytic enzyme genes. The reduction in the Pfk and GAPDH enzyme
amounts was intriguing, because the Pfk and GAPDH steps are critical in determining
glycolytic flux (3, 31). A reduction in these enzymes could therefore decrease glycolytic flux,
and would reroute glucose (G6P) allocations via mass action towards other branches of
glucose metabolism, primarily the pentose phosphate pathway as well as trehalose
biosynthesis (Figure 2A). To assess this, we first measured the steady-state levels of key
glycolytic and pentose phosphate pathway (PPP) intermediates, and trehalose in WT or *ubp3Δ*
using targeted LC-MS/MS. Glucose-6/fructose-6 phosphate (G6P/F6P) increased in *ubp3Δ*
(Figure 2C). Concurrently, trehalose, and the PPP intermediates ribose 5-phosphate (R5P) and
sedoheptulose 7-phosphate (S7P), increased in *ubp3Δ* (Figure 2C).

Since steady-state metabolite amounts cannot separate production from utilization,
in order to unambiguously assess if glycolytic flux is reduced in *ubp3Δ* cells, we utilized a pulse
labeling of $^{13}C_6$ glucose, following which the label incorporation into glycolytic and other
intermediates was measured. Note that because glycolytic flux is very high in yeast, this
experiment would require rapid pulsing and extraction of metabolites in order to stay in a
linear range and avoid label saturation. We therefore established a very short time point of

label-addition, quenching and metabolite extraction post ^{13}C glucose pulse. Since flux

saturates/reaches steady-state in seconds, we first ensured that the label incorporation into

individual metabolites after the ^{13}C glucose pulse was in the linear range, and for early

glycolytic intermediates this was seconds after glucose addition. This new methodology is

extensively described in materials and methods, with required controls shown in Figure S2E.

WT and *ubp3Δ* cells were grown in high glucose, pulsed with $^{13}C_6$ glucose, and the relative ^{13}C

label incorporation into glycolytic intermediates and trehalose were measured, as shown in

the schematic (Figure S2D). In *ubp3Δ,* ^{13}C label incorporation into G6P/F6P as well as trehalose

substantially increased (Figure 2D). Contrastingly, ^{13}C label incorporation into glycolytic

intermediates F1,6BP, G3P, 3PG and PEP decreased, indicating decreased glycolytic flux

(Figure 2D). We next measured ethanol concentrations and production rates, as an additional

output of relative glycolytic rates. We observed decreased steady-state ethanol levels, as well

as ethanol production rates in *ubp3Δ* (Figure 2F).

Collectively, these results reveal that that reduced Pfk1 and GAPDH (in *ubp3Δ*)

decrease glucose flux via glycolysis, which results in rewired glucose flux towards trehalose

biosynthesis and the PPP.

Rerouted glucose flux results in phosphate (Pi) accumulation

We therefore asked if the proteomic state, as observed in *ubp3Δ* cells, could provide

clues to explain the coupling between mitochondrial derepression and rerouted glucose flux.

A recent study by Isasa et al., had systematically quantified the changes in protein levels in

ubp3Δ cells (28). We therefore reanalyzed this extensive dataset, looking for changes in

proteins that would correlate with these metabolic processes. Notably, we observed

increased levels of proteins of the mitochondrial ETC and respiration, and decreased amounts

of glucose metabolizing enzymes in *ubp3Δ* (Figure 3A). Additionally, multiple proteins

involved in regulating phosphate (Pi) homeostasis were decreased in *ubp3Δ* (Figure 3A) (28).

We had recently uncovered a reciprocal coupling of Pi homeostasis with the different arms of

glucose metabolism, particularly trehalose biosynthesis (32, 33), and therefore wondered if a

glycolytic-flux dependent change in Pi homeostatsis had any role in mitochondrial respiration.

Our hypothesis was refined based on the reasoning given below.

One explanation for mitochondrial repression can be an internal competition for

shared metabolites/co-factors between (cytosolic) glycolytic and mitochondrial processes,

that mitochondria might not be sufficiently able to access (14, 18, 19). In this context, a

plausible role for inorganic phosphate (Pi) in regulating mitochondrial repression can be

hypothesized. Cytosolic glycolysis requires rapid, high consumption of net Pi (19, 34, 35), and

this could possibly limit the Pi that is continuously available for mitochondrial use (18, 36),

thereby repressing mitochondria. Contextually, the balance between reactions releasing vs

consuming Pi could explain changes in global Pi levels (33). A continuous hub of Pi

consumption is glycolysis. Here, GAPDH catalyzes G3P to 1,3BPG, converting ADP to ATP,

while concurrently consuming a molecule of Pi (35, 37). This Pi that goes into ATP will

subsequently be used for nucleotide biosynthesis, polyphosphate biosynthesis and protein

phosphorylation (33, 38). Therefore, we can surmise that in high glycolytic flux, the

production of ATP, nucleotides and polyphosphates is concurrent with Pi consumption (37,

39, 40). Could this reaction therefore limit cytosolic Pi for the mitochondria (as illustrated in

Figure 3B)? Notably, *ubp3Δ* have reduced GAPDH levels and decreased glycolytic flux. Second,

trehalose synthesis is a Pi releasing reaction, and a major source of free Pi that is critical for

Pi homeostasis (32, 35). Flux through this reaction is also substantially higher in *ubp3Δ*.

Therefore, these cells might have increased Pi release (via trehalose), coupled with

decreased Pi consumption (via decreased GAPDH). We therefore asked if total Pi increases in

ubp3Δ (Figure 3B)?

To test this, we directly assessed total Pi levels in *ubp3Δ* and WT. *ubp3Δ* cells had

higher Pi than WT, and this was comparable to Pi in WT grown in excess Pi (Figure 3C).

Similarly, Pi amounts also increased in Ubp3[C469A] (Figure S3A). Therefore, the loss of Ubp3

increases intracellular Pi levels. Next, we asked if *ubp3Δ* cells exhibit signatures of a 'high Pi'

state. *S. cerevisiae* maintains internal Pi balance by controlling the expression of multiple

genes collectively known as the Pho regulon (41). The Pho regulon is induced under Pi

limitation, and repressed during Pi sufficiency (32, 41). We assessed two major Pho proteins

(Pho84: a high-affinity membrane Pi transporter, and Pho12: an acid phosphatase) in WT and

ubp3Δ in high glucose. *ubp3Δ* cells have lower amounts of Pho84 and Pho12 (Figure 3D).

Further, upon shifting to low Pi for one hour, Pho84 and Pho12 increased in *ubp3Δ*. These

data suggest that reduced Pho84 and Pho12 amounts in *ubp3Δ* are because of increased Pi,

and not due to altered Pho regulon function itself (Figure 3D). These data clarify earlier

observations from *ubp3Δ* which noted reduced Pho proteins (28). Therefore, *ubp3Δ*

constitutively have higher Pi, and likely a consequent decrease in Pho proteins.

We next asked if the increased Pi in *ubp3Δ* is because of altered G6P allocations

towards different end-points, particularly trehalose synthesis, which can be a major node of

Pi restoration (32, 35). We assessed the contribution of increased trehalose synthesis towards

the high Pi in *ubp3Δ*, by estimating Pi levels in the absence of trehalose 6-phosphate

phosphatase (Tps2), which catalyzes the Pi-releasing step in trehalose synthesis. Notably, loss

of Tps2 in *ubp3Δ* decreased Pi (Figure 3E). There was no additive difference in Pi between

tps2Δ and *ubp3Δtps2Δ* (Figure 3E). As an added control, we assessed trehalose in WT and

ubp3Δ in the absence of Tps2, and found no difference (Figure S3B). Therefore, increasing

G6P flux towards trehalose biosynthesis is a major source of the increased Pi in *ubp3Δ*.

Since the major GAPDH isozymes, Tdh2 and Tdh3, are reduced in *ubp3Δ*, we directly

asked if reducing GAPDH can decrease Pi consumption and increase Pi. To assess this, we

generated *tdh2Δtdh3Δ* cells, which exhibit a growth defect, but are viable, permitting further

analysis. *tdh2Δtdh3Δ* had higher Pi in high glucose (Figure 3F). Expectedly, we observed a

decrease in ethanol in *tdh2Δtdh3Δ* (Figure S3C), along with an accumulation of F1,6 BP and

G3P, and decreased 3PG and PEP (Figure S3D). However, G6P and trehalose levels between

WT and *tdh2Δtdh3Δ* were comparable (Figure S3D, S3E). These data sugest that unlike in

ubp3Δ, the increased Pi in *tdh2Δtdh3Δ* comes mainly from decreased Pi consumption (GAPDH

step). To further assess the role of reduced glycolytic flux in increasing Pi (*ubp3Δ* and

tdh2Δtdh3Δ), we measured the Pi in these cells growing in a gluconeogenic medium – 2%,

ethanol. In this scenario, the GAPDH catalyzed reaction will be reversed, converting 1,3BPG

to G3P, which should release and not consume Pi. Compared to WT, the Pi levels decreased

in *ubp3Δ* and *tdh2Δtdh3Δ* (Figure 3G), suggesting that the changes in Pi in these mutants is

driven by the relative change in Pi release vs consumption.

These results collectively indicate that the combined effect of increased Pi release

coming from trehalose synthesis, and decreased Pi consumption from reduced GAPDH

increase Pi levels in *ubp3Δ*.

Mitochondrial Pi availability correlates with mitochondrial activity in *ubp3Δ*

We therefore wondered if this observed Pi increase from the combined rewiring of

glucose metabolism resulted in more Pi becoming accessible to the mitochondria. This would

effectively result in Pi budgeting between cytosolic glycolysis and mitochondria based on the

relative flux in different arms of glucose metabolism, by increasing Pi availability for the

mitochondria (Figure 4A). To test if this were so, we first asked if the mitochondria in *ubp3Δ*

have increased Pi. Mitochondria were isolated by immunoprecipitation from WT and *ubp3Δ*,

the isolation efficiency was analyzed (Figure S4A), and relative Pi levels compared. Pi levels

were normalised to Idh1 (isocitrate dehydrogenase) in isolated mitochondria, since Idh1

protein levels did not decrease in *ubp3Δ* (Figure S4B), and consistent with (28). Mitochondrial

Pi was higher in *ubp3Δ* (Figure 4B). We next asked if the increased mitochondrial activity in

ubp3Δ is a consequence of high Pi. If this were so, *tdh2Δtdh3Δ* should partially phenocopy

ubp3Δ with respect to mitochondrial activity. Consistently, *tdh2Δtdh3Δ* have higher

mitotracker intensity as well as Cox2 levels compared to WT (Figure S4C, Figure 4C). Also

consistent with this, a high basal OCR in *tdh2Δtdh3Δ* was observed (Figure 4D), together

indicating high mitochondrial activity in *tdh2Δtdh3Δ*, which is comparable to *ubp3Δ*.

We next asked if higher Pi is necessary to increase mitochondrial activity in *ubp3Δ* and

tdh2Δtdh3Δ. Since glycolysis is defective in *ubp3Δ* and *tdh2Δtdh3Δ*, it is necessary to

distinguish the effect of high Pi vs. only the effect of low glycolysis in activating mitochondria.

Logically, if decreased glycolysis (independent of Pi) is sufficient to activate mitochondria,

bringing down the Pi levels in *ubp3Δ* to that of WT should not affect mitochondrial activity.

Notably, *ubp3Δ* grown in low (1 mM) Pi have Pi levels similar to WT in standard (normal Pi)

medium (Figure S4D). *ubp3Δ* grown in low Pi also had decreased ethanol, suggesting reduced

glycolysis (Figure S4E). Therefore, we used this condition to further understand the role of Pi

in inducing mitochondrial activity. Mitotracker fluorescence decreased in both *ubp3Δ* and

tdh2Δtdh3Δ in low Pi (Figure S4F). Note: basal mitotracker fluorescence in WT also decreases

in low Pi, which is consistent with a required role of Pi for mitochondrial activity (Figure S4F).

Similarly, Cox2 levels were reduced in both *ubp3Δ* and *tdh2Δtdh3Δ* (Figure 4E). Consistent

with both reduced mitotracker intensity and Cox2 levels, basal OCR also decreased in both

ubp3Δ and *tdh2Δtdh3Δ* in low Pi (Figure 4F). These data collectively suggest that high

intracellular Pi is necessary to increase mitochondrial activity in *ubp3Δ* and *tdh2Δtdh3Δ*.

Next, we asked how mitochondrial Pi transport regulates mitochondrial activity. Mir1

and Pic2 are mitochondrial Pi transporters, with Mir1 being the major Pi transporter (42, 43).

We first limited mitochondrial Pi availability in *ubp3Δ* by knocking-out *MIR1*. In *mir1Δ*, the

increased Cox2 observed in *ubp3Δ* was no longer observed (Figure 4G). Consistent with this,

we observed no further increase in the basal OCR in *mir1Δubp3Δ* compared to *mir1Δ* (Figure

4H). Furthermore, *mir1Δ* showed decreased Cox2 as well as basal OCR even in WT cells (Figure

4G, Figure 4H). These data together suggest that mitochondrial Pi transport is critical for

increasing mitochondrial activity in *ubp3Δ*, and in maintaining basal mitochondrial activity

even in high glucose.

As a control, no significant increase in Mir1 and Pic2 was observed in *ubp3Δ* (Figure

S4G), suggesting that *ubp3Δ* do not increase mitochondrial Pi by merely increasing the Pi

transporters, but rather by increasing available Pi pools.

Taken together, these data suggest the possibility that increased cytosolic Pi increases

the mitochondrial Pi pool, which increases mitochondrial activity. Decreasing mitochondrial

Pi by either reducing total Pi, or by reducing mitochondrial Pi transport decreases

mitochondrial activity.

Mitochondrial Pi availability constrains mitochondrial activity under high glucose

So far, these data suggest that the cytosolic Pi available for the mitochondria can

dermine the extent of mitochondrial activity. Therefore, we further investigated if

mitochondrial Pi allocation was a necessary constraint for glucose-mediated mitochondrial

repression.

To test this, we first asked how important mitochondrial Pi transport was to switch to increased respiration. In WT yeast, low glucose or glycolytic inhibition will result in increased respiration (22). What happens therefore if we restrict mitochondrial Pi in this context? For this, we measured the basal OCR in WT and *mir1Δ* after switching from high (2%) to low (0.1%) glucose. We observed a significant increase in the basal OCR in WT but not in *mir1Δ* (Figure 5A). The alternate scenario is after glycolytic inhibition. We assessed the role of mitochondrial Pi in this context, by inhibiting glycolysis using 2-deoxyglucose (2DG). WT, but not *mir1Δ*, increased their OCR (respiration) upon a 1-hour treatment with 2DG (Figure 5B). Consistent with this, mitotracker fluorescence increased with an increase in 2DG in WT, but not in *mir1Δ* (Figure S5A). We further asked if the mitochondrial Pi transporter itself glucose repressed, and therefore assessed Mir1 amounts in high and low glucose. Mir1 levels are higher upon a shift to low glucose, and in cells grown in 2% ethanol, suggesting that Mir1 is glucose repressed (Figure 5C, S5B). These data suggest that mitochondrial Pi transport is necessary for increasing mitochondrial activity after glucose derepression.

We next asked whether just adding external Pi was sufficient to increase mitochondrial activity, when cells are in high glucose. In medium supplemented with excess Pi, the internal Pi increases as seen earlier (Figure 3C). Therefore, a simplistic assumption would be that the addition of external Pi to cells in high glucose would also increase mitochondrial Pi. However, an alternate possibility presents itself whererin since glycolytic flux is already high in glucose, supplementing Pi will continue to fuel glycolysis. Indeed, this was originally obseved by Harden and Young in 1908, where adding Pi increased fermentation (44). In such a scenario, mitochondrial Pi will not increase. We estimated mitochondrial Pi in this condition where excess Pi was externally supplemented. Notably, cells grown in high Pi had decreased mitochondrial Pi (Figure S5C). Furthermore, directly adding Pi to cells growing in high glucose decreased basal OCR (Figure S5D), consistent with decreased mitochondrial Pi. These data indicate that in high glucose, simply supplementing Pi will not increase Pi access to the mitochondria. We therefore now asked, if we inhibit glycolytic flux and then supplement Pi, what would happen to mitochondrial activity. For this, we treated cells with 2DG, and subsequently added Pi and measured the OCR (Figure S5E). In this case, supplementing Pi increased the basal OCR (Figure S5E). Collectively, these data suggest that a combination of decreasing glycolysis and increasing Pi can together increase respiration.

Next, in order to directly test mitochondrial activation based on external Pi availability, we

isolated mitochondria, and estimated activity *in vitro* upon adding increasing Pi.

Mitochondrial activity increased with increased Pi, with maximum activity observed with 25

mM Pi supplemented (Figure 5D). In a complementary experiment, we overexpressed the

Mir1 transporter in wild-type cells, to increase Pi within mitochondria (Figure S5F). Mir1-OE

cells have higher Cox2 levels and basal OCR (Figure 5E, 5F). Therefore, increasing Pi transport

to mitochondria is sufficient to increase mitochondrial activity in high glucose.

Mitochondrial pyruvate transport is also required for mitochondrial respiration (45).

We asked where Pi availability stands in a hierarchy of constraints for mitochondrial

derepression, as compared to mitochondrial pyruvate transport. We measured the amounts

of the Mpc3 subunit of the mitochondrial pyruvate carrier (MPC) complex (45, 46). Mpc3

protein increases in *ubp3Δ* (Figure S5G), correlating with the increased mitochondrial activity.

Interestingly, in *ubp3Δ* grown in low Pi, Mpc3 further increased (Figure S5G), but as shown

earlier this condition cannot increase OCR or mitochondrial activity (Figure 4F, S4F). Basal

Mpc3 levels decrease in *mir1Δ*, but upon shifting to 0.1% glucose, Mpc3 increases in both WT

and *mir1Δ*, with higher levels in *mir1Δ* (Figure S5H). Therefore, even where Mpc3 is high

(*ubp3Δ* in low Pi, and *mir1Δ* in low glucose), mitochondrial activity remains low if Pi is

restricted (Figure 4F, 4H). There was also no decrease in basal OCR in *mpc3Δ* in high glucose,

and the basal OCR increased to the same level as of WT after shifting to 0.1% glucose (Figure

S5I). Since Mpc3 changes with mitochondrial Pi availability (Figure S5G, S5H), we also

measured Mpc3 in the Mir1OE. No further changes in Mpc3 were observed in Mir1OE (Figure

S5J), indicating that increasing mitochondrial Pi alone need not increase Mpc3. Overall,

although Mpc3 levels corelate with decreased glycolysis (*ubp3Δ* -Figure S5G, *ubp3Δ* in low Pi-

Figure S5G, low glucose -Figure S5H), increased Mpc3 alone cannot increase mitochondrial

activity and respiration in the absence of adequate mitochondrial Pi.

Finally, we asked how important mitochondrial Pi transport was for growth, under

2DG treatment. In the presence of 2DG, WT cells have only slightly decreased growth. In

contrast, *mir1Δ* show a severe growth defect upon 2DG treatment (Figure 5G), revealing a

synergetic effect of combining 2DG with inhibiting mitochondrial Pi transport. Therefore, the

combined inhibition of glycolysis and mitochondrial Pi transport restricts the growth of

glycolytic cells.

Collectively, mitochondrial Pi availability constrains glucose-mediated mitochondrial

repression. Increasing available pools of Pi to enter the mitochondria is sufficient to induce

mitochondrial activity.

Discussion

In this study, we highlight a role for Pi budgeting between cytosolic glycolysis and

mitochondrial processes (which compete for Pi) in constraining mitochondrial repression

(Figure 5H). Ubp3 controls this process (Figure 1), by maintaining the amounts of the glycolytic

enzymes Pfk1 and GAPDH (Tdh2 and Tdh3) and thereby allows high glycolytic flux. At high

fermentation rates, glycolytic enzymes levels at maximal activity maintain high glycolytic rates

(effectively following zero-order kinetics), and therefore, changes in the enzyme levels will

have a direct, proportionate effect on flux (47). The loss of Ubp3 decreases glycolytic flux,

resulting in a systems-level, mass-action based rewiring of glucose metabolism where more

G6P is routed towards trehalose synthesis and PPP (Figure 2, Figure 5H). Indeed, inhibiting

just phosphofructokinase can reroute glucose flux from glycolysis to PPP (48–50), and our

study now permits contextualized interpretations of these results. Such reallocations of

glucose flux will collectively increase overall Pi, coming from the combined effect of increased

Pi release from trehalose synthesis, and decreased Pi consumption via reduced GAPDH. This

altered intracellular Pi economy increases Pi pools available to mitochondria, and increasing

mitochondrial Pi is necessary and sufficient to increase respiration in high glucose (Figure 4,

Figure 5). Mitochondrial Pi transport maintains mitochondrial activity in high glucose, and

increases mitochondrial activity in low glucose (Figure 4, Figure 5). Finally, the mitochondrial

Pi transporter Mir1 itself decreases in high glucose (Figure 5). Therefore, this glucose-

dependent repression of Mir1 also restricts mitochondrial Pi availability (and thereby activity)

in high glucose.

Traditionally, loss-of-function mutants of metabolic enzymes are used to understand

metabolic state regulation. This approach negates nuanced investigations, since metabolic

enzymes are often essential for viability. Furthermore, metabolic pathways have multiple,

contextually regulated nodes, through which cells maintain their metabolic state. Therefore,

alternate approaches to identify global regulators of metabolic states (as opposed to single enzymes), might uncover ways via which multiple nodes are simultaneously tuned, and can reveal unanticipated systems-level principles of metabolic state rewiring. In this study of glucose-mediated mitochondrial repression, the loss of the DUB Ubp3 decreases glycolytic flux by reducing the enzymes at two critical nodes in the pathway - Pfk1 and GAPDH. Unlike loss of function mutants, a reduction in amounts will only rewire metabolic flux. By 'hitting' multiple steps in glycolysis simultaneously, *ubp3Δ* have decreased Pi consumption, as well as increased Pi release. Such a cumulative phenomenon reveals more than inhibiting only GAPDH, where increased Pi comes only from reduced Pi consumption, and not from increased trehalose biosynthesis. Our serendipitous identification of a regulator which regulates multiple steps in glucose metabolism to change the metabolic environment, now suggests a general basis of mitochondrial regulation that would have otherwise remained hidden. Separately, finding the substrates of Ubp3 and whether Ubp3 directly regulates glycolytic enzymes are exciting future research questions requiring concurrent innovations in accessible chemical-biological approaches to study DUBs.

Because phosphates are ubiquitous, it is challenging to identify hierarchies of Pi-dependent processes in metabolic state regulation (33). Phosphate transfer reactions are the foundation of metabolism, driving multiple, thermodynamically unfavorable reactions (51, 52). Contextually, the laws of mass action predict that the relative rates of these reactions will regulate overall Pi balance, and contrarily the Pi allocation to Pi-dependent reactions will determine reaction rates (33, 35). Additionally, cells might control Pi allocations for different reactions via compartmentalizing Pi in organelles, to spatially restrict Pi availability (53–55). Our data collectively suggest a paradigm where the combination of factors regulates mitochondrial Pi and thereby activity. In glycolytic yeast cells growing in high glucose, mitochondrial Pi availability becomes restricted due to higher utilization of Pi in glycolysis compared to mitochondria. Consistent with this, a rapid decrease in Pi upon glucose addition has been observed (18, 19, 37). Interesting, *in vitro* studies with isolated mitochondria from tumor cells also find that decreasing Pi levels decreases respiration (19), which would be consistent with this scenario. Further, supplementing Pi correlates with decreased mitochondrial repression in tumors (18, 56). By increasing Pi through a systems-level rewiring of glucose metabolism (such as in *ubp3Δ* cells), cells can collectively increase mitochondrial

access to Pi. This Pi budgeting determines mitochondrial activity. Supplementing Pi under conditions of low glycolysis (where mitochondrial Pi transport is enhanced), as well as directly supplementing Pi to isolated mitochondria, increases respiration (Figure 5, S5). Therefore, in order to deprepress mitochondria, a combination of increased Pi along with decreased glycolysis is required. An additional systems-level phenomenon that might regulate Pi transport to the mitochondria is the decrease in cytosolic pH upon decreased glycolysis (57, 58). Since mitochondrial Pi transport itself dependeds on the proton gradient, a low cytosolic pH would favour mitochondrial Pi transport (59). Therefore, decreasing glycolysis (2DG treatment, deletion of Ubp3, and decreased GAPDH activity), and increasing cytosolic Pi might synergestically increase mitochondria Pi transport, and increase respiration. Alternately, increasing mitochondrial Pi transporter amounts can achieve the same result, as seen by overexpressing Mir1 (Figure 5). A similar observation has been reported in *Arabidopsis*, reiterating an evolutionarily conserved role for mitochondrial Pi in controlling respiration (60). Relatedly, glycolytic inhibition can suppress cell proliferation in Warburg-positive tumors (61, 62), or inflammatory responses (63). However, these cells survive by switching to mitochondrial respiration (64, 65), requiring alternate approaches to prevent their proliferation (66). Inhibiting mitochondrial Pi transport in combination with glycolytic inhibition could restrict the proliferation of Warburg/Crabtree positive cells.

Since its discovery in the 1920s, the phenomenon of accelerated glycolysis with concurrent mitochondrial repression has been intensely researched. Yet, the biochemical constraints for glucose-mediated mitochondrial repression remains unresolved. One hypothesis suggests that the availability of glycolytic intermediates might determine the extent of mitochondrial repression. F1,6BP inhibits complex III and IV of the electron transport chain (ETC) in Crabtree positive yeast (14–17). Similarly, the ratio between G6P and F1,6BP regulates the extent of mitochondrial repression (16, 17). Although G6P/F6P accumulates in *ubp3Δ* (Figure 2C), this is not the case in *tdh2Δtdh3Δ* (GAPDH mutant) (Figure S3D), suggesting that G6P/F6P accumulation in itself is not the criterion to increase mitochondrial activity. Separately, the competition for common metabolites/co-factors between glycolysis and respiration (such as ADP, Pi or pyruvate) could drive this phenomenon (14, 18). Here we observe that Mpc3 mediated mitochondrial pyruvate transport alone cannot increase respiration. An additional consideration is the possible contribution of changes in ADP in

regulating mitochondrial activity, where the use of ADP in glycolysis might limit mitochondrial

ADP. Therefore, when Pi changes as a consequence of glycolysis, it coud be imagined that a

change in ADP balance can coincidentally occur. However, prior studies show that even

though cytosolic ADP decreases in the presence of glucose, this does not limit mitochondrial

ADP uptake, or decrease respiration, due to the very high affinity of the mitochondrial ADP

transporter (14, 19). These collectively reiterate the importance of Pi access and transport to

mitochondria in constraining mitochondrial respiration. Indeed, this interpretation can also

contextually explain observations from other model systems where mitochondrial Pi

transport seems to regulate respiration (67, 68).

We parsimoniously suggest that Pi access to the mitochondria as a key constraint for

mitochondrial repression under high glucose. In a hypothetical scenario, a single-step event

in evolution, reducing mitochondrial Pi transporter amounts, and/or increasing glycolytic flux

(to deplete cytosolic Pi), will result in whole-scale metabolic rewiring to repress mitochondria.

More elaborate regulatory events can easily be imagined as subsequent adaptations to

enforce mitochondrial repression. Given the central role played by Pi, something as

fundamental as access to Pi will constrain mitochondrial repression. Concurrently, the rapid

incorporation of Pi into faster glycolysis can give cells a competitive advantage, while also

sequestering Pi in the form of usable ATP. Over the course of evolution, this could conceivably

drive other regulatory mechanisms to enforce mitochondrial repression, leading to the

currently observed complex regulatory networks and signaling programs observed in the

Crabtree effect, and other examples of glucose-dependent mitochondrial repression.

Materials and methods

Statistics and graphing

Unless otherwise indicated, statistical significance for all indicated experiments were

calculated using unpaired Student's t-tests (GraphPad prism 9.0.1). Graphs were plotted using

GraphPad prism 9.0.1

Yeast strains, media, and growth conditions

A prototrophic CEN.PK strain of *Saccharomyces cerevisiae* (WT) (69) was used. Strains are listed in appendix table S1. Gene deletions, chromosomal C-terminal tagged strains were generated by PCR mediated gene deletion/tagging (70). Media compositions, growth conditions and CRISPR-Cas9 based mutagenesis are described in extended methods (SI appendix).

Mitotracker fluorescence

Mitotracker fluorescence was measured using Thermo Varioscan™ LUX multimode plate reader (579/599 excitation/emission). Detailed protocol is described in extended methods. Mitotracker fluorescence were normalized using OD_{600} of each sample and relative fluorescence intensity calculated.

Protein extraction and Western blotting

Total protein was precipitated, extracted using trichloroacetic acid (TCA) as described earlier (71). Blots were quantified using ImageJ software. Detailed protocol is described in extended methods.

Basal OCR measurement

The basal OCR was measured using Agilent Seahorse XFe24 analyzer. Basal OCR readings were normalized for cell number (using OD_{600} of samples) in each well. The detailed methods are described in extended methods.

RNA extraction and RT-qPCR

The RNA extraction was done using the hot phenol extraction method as described in (71). The isolated RNA was DNase treated, and used for cDNA synthesis. Superscript III reverse transcriptase enzyme (Invitrogen) was used for cDNA synthesis and RT-qPCR was performed using KAPA SYBR FAST qRT PCR kit (KK4602, KAPA Biosystems). Taf10 was used as a control for normalisation and the fold change in mRNA levels were calculated by 2^-ddct method.

ATP, ethanol and Pi measurements

501 ATP levels were measured by ATP estimation kit (Thermo Fisher A22066). Ethanol
502 concentration in the medium was estimated using potassium dichromate based assay
503 described in (72) with modifications. Pi was estimated using a malachite green phosphate
504 assay kit (Cayman chemicals, 10009325). Detailed sample collection and assay protocols are
505 described in extended methods.

Metabolite extraction and analysis by LC-MS/MS

507 The steady state levels and relative ^{13}C label incorporation into metabolites were estimated
508 by quantitative LC-MS/MS methods as described in (73). Detailed methodology is extensively
509 described in extended methods. Peak area measurements are listed in supplement file 1.

Mitochondrial isolation

511 Mitochondria was isolated by immunoprecipitation as described in (74, 75) with
512 modifications. The detailed protocol is described in extended methods (SI appendix). The
513 eluted mitochondria were used in malachite green assay for Pi estimation, boiled with SDS-
514 glycerol buffer for western blots or incubated with mitotracker CMXROS with mitochondria-
515 activation buffer for mitotracker assays.

Data availability

518 The LC-MS/MS peak area for all the metabolites can be found in supplement file 1. This study
519 includes no data deposited in external repositories.

Acknowledgements

522 We thank all SL lab members, Benjamin Tu, Anand Bachhawat and Vijay Jayaraman for
523 comments and suggestions. We acknowledge the NCBS/inStem/CCAMP mass spectrometry
524 facility for LC-MS/MS support. We thank Aayushee Khanna for help with experiments. V.V
525 acknowledges funding support from DST-INSPIRE fellowship (IF170236) from the Department
526 of Science and Technology (DST), Govt. of India, S.N and SA acknowledges intramural funding

from the DBT-inStem PhD program. S.L acknowledges support from the DST SERB CRG grant CRG/2019/004772, a DBT-Wellcome India Alliance Senior Fellowship (IA/S/21/2/505922), and institutional support from inStem.

References

1. J. Zhu, C. B. Thompson, Metabolic regulation of cell growth and proliferation. *Nat. Rev. Mol. Cell Biol.* **20**, 436–450 (2019).

2. L. Cai, B. P. Tu, Driving the Cell Cycle Through Metabolism. *Annu. Rev. Cell Dev. Biol.* **28**, 59–87 (2011).

3. D. L. (David L. Nelson, D. L. (David L. Nelson, A. L. Lehninger, M. M. Cox, Lehninger principles of biochemistry (2008) (March 10, 2022).

4. O. Warburg, The metabolism of carcinoma cells 1. *J. Cancer Res.* **9**, 148–163 (1925).

5. M. G. Vander Heiden, L. C. Cantley, C. B. Thompson, Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. *Science (80-.).* **324**, 1029–1033 (2009).

6. H. G. Crabtree, Observations on the carbohydrate metabolism of tumours. *Biochem. J.* **23**, 536–545 (1929).

7. R. H. De Deken, The Crabtree effect: a regulatory system in yeast. *J. Gen. Microbiol.* **44**, 149–156 (1966).

8. L. Xia, *et al.*, The cancer metabolic reprogramming and immune response. *Mol. Cancer* **20**, 1–21 (2021).

9. M. D. Kornberg, The immunologic Warburg effect: Evidence and therapeutic opportunities in autoimmunity. *Wiley Interdiscip. Rev. Syst. Biol. Med.* **12**, 1–17 (2020).

10. N. Pacini, F. Borziani, Cancer stem cell theory and the warburg effect, two sides of the same Coin? *Int. J. Mol. Sci.* **15**, 8893–8930 (2014).

11. A. M. Abdel-Haleem, *et al.*, The emerging facets of non-cancerous Warburg effect. *Front. Endocrinol. (Lausanne).* **8**, 1–7 (2017).

12. E. Tsogtbaatar, C. Landin, K. Minter-Dykhouse, C. D. L. Folmes, Energy Metabolism Regulates Stem Cell Pluripotency. *Front. Cell Dev. Biol.* **8**, 1–16 (2020).

13. A. Cornish-Bowden, Biochemical evolution : the pursuit of perfection. *Garl. Sci.* (2016).

14.   R. Diaz-Ruiz, M. Rigoulet, A. Devin, The Warburg and Crabtree effects: On the origin

of cancer cell energy metabolism and of yeast glucose repression. *Biochim. Biophys.*

Acta - Bioenerg. **1807**, 568–576 (2011).

15.   N. Hammad, M. Rosas-Lemus, S. Uribe-Carvajal, M. Rigoulet, A. Devin, The Crabtree

and Warburg effects: Do metabolite-induced regulations participate in their

induction? *Biochim. Biophys. Acta - Bioenerg.* **1857**, 1139–1146 (2016).

16.   R. Díaz-Ruiz, *et al.*, Mitochondrial Oxidative Phosphorylation Is Regulated by Fructose

1,6-Bisphosphate: A POSSIBLE ROLE IN CRABTREE EFFECT INDUCTION? *. *J. Biol.*

Chem. **283**, 26948–26955 (2008).

17.   M. R. Lemus, *et al.*, The role of glycolysis-derived hexose phosphates in the induction

of the Crabtree effect. *J. Biol. Chem.* **293**, 12843–12854 (2018).

18.   D. H. Koobs, Phosphate mediation of the Crabtree and Pasteur effects. *Science (80-.).*

178, 127–133 (1972).

19.   S. Rodríguez-Enríquez, O. Juárez, J. S. Rodríguez-Zavala, R. Moreno-Sánchez, Multisite

control of the Crabtree effect in ascites hepatoma cells. *Eur. J. Biochem.* **268**, 2512–

2519 (2001).

20.   F. Tripodi, R. Nicastro, V. Reghellin, P. Coccetti, Post-translational modifications on

yeast carbon metabolism: Regulatory mechanisms beyond transcriptional control.

Biochim. Biophys. Acta - Gen. Subj. **1850**, 620–627 (2015).

21.   T. Hitosugi, J. Chen, Post-translational modifications and the Warburg effect.

Oncogene **33**, 4279–4285 (2014).

22.   J. R. Broach, Nutritional control of growth and development in yeast. *Genetics* **192**,

73–105 (2012).

23.   A. Hershko, A. Ciechanover, THE UBIQUITIN SYSTEM. *Annu. Rev. Biochem.* **67**, 425–79

(1998).

24.   D. Komander, M. Rape, The Ubiquitin Code. *Annu. Rev. Biochem.* **81**, 203–229 (2012).

25.   C. M. Pickart, M. J. Eddins, Ubiquitin: Structures, functions, mechanisms. *Biochim.*

Biophys. Acta - Mol. Cell Res. **1695**, 55–72 (2004).

26.   E. Postma, C. Verduyn, W. A. Scheffers, J. P. Van Dijken, Enzymic analysis of the

crabtree effect in glucose-limited chemostat cultures of Saccharomyces cerevisiae.

Appl. Environ. Microbiol. **55**, 468–77 (1989).

27.   B. Ossareh-Nazari, *et al.*, Cdc48 and Ufd3, new partners of the ubiquitin protease

Ubp3, are required for ribophagy. *EMBO Rep.* **11**, 548–554 (2010).

28. M. Isasa, *et al.*, Multiplexed, Proteome-Wide Protein Expression Profiling: Yeast
Deubiquitylating Enzyme Knockout Strains. *J. Proteome Res.* **14**, 5306–5317 (2015).

29. R. Nostramo, S. N. Varia, B. Zhang, M. M. Emerson, P. K. Herman, The Catalytic
Activity of the Ubp3 Deubiquitinating Protease Is Required for Efficient Stress Granule
Assembly in Saccharomyces cerevisiae. *Mol. Cell. Biol.* **36**, 173 (2016).

30. F. Fontanesi, I. C. Soto, D. Horn, A. Barrientos, Assembly of mitochondrial cytochrome
c-oxidase, a complicated and highly regulated cellular process. *Am. J. Physiol. - Cell*
Physiol. **291**, 1129–1147 (2006).

31. A. A. Shestov, *et al.*, Quantitative determinants of aerobic glycolysis identify flux
through the enzyme GAPDH as a limiting step. *Elife* **3**, 1–18 (2014).

32. R. Gupta, *et al.*, A tRNA modification balances carbon and nitrogen metabolism by
regulating phosphate homeostasis. *Elife* **8**, 1–33 (2019).

33. R. Gupta, S. Laxman, Cycles, sources, and sinks: Conceptualizing how phosphate
balance modulates carbon flux using yeast metabolic networks. *Elife* **10**, 1–14 (2021).

34. P. W. Mason, D. P. Carbone, R. A. Cushman, A. S. Waggoner, The importance of
inorganic phosphate in regulation of energy metabolism of Streptococcus lactis. *J.*
Biol. Chem. **256**, 1861–1866 (1981).

35. J. H. Van Heerden, *et al.*, Lost in transition: Start-up of glycolysis yields subpopulations
of nongrowing cells. *Science (80-.).* **343** (2014).

36. P. C. Brazy, L. J. Mandel, V. W. Dennis, S. R. Gullans, Metabolic Requirement for
Inorganic Phosphate by the Rabbit Proximal Tubule: EVIDENCE FOR A CRABTREE
EFFECT Metabolic Requirement for Inorganic Phosphate by the Rabbit Proximal
Tubule EVIDENCE FOR A CRABTREE EFFECT. *J Clin Invest* **70**, 53 (1982).

37. S. Hohmann, W. Bell, M. J. Neves, D. Valckx, J. M. Thevelein, Evidence for trehalose-6-
phosphate-dependent and-independent mechanisms in the control of sugar influx
into yeast glycolysis. *Mol. Microbiol.* **20**, 981–991 (1996).

38. T. Hunter, Why nature chose phosphate to modify proteins. *Philos. Trans. R. Soc. B*
Biol. Sci. **367**, 2513–2516 (2012).

39. P. O. Ljungdahl, B. Daignan-Fornier, Regulation of amino acid, nucleotide, and
phosphate metabolism in Saccharomyces cerevisiae. *Genetics* **190**, 885–929 (2012).

40. S. Austin, A. Mayer, Phosphate Homeostasis – A Vital Metabolic Equilibrium

Maintained Through the INPHORS Signaling Pathway. *Front. Microbiol.* **11**, 1367
(2020).

41.   J. M. Mouillon, B. L. Persson, New aspects on phosphate sensing and signalling in
Saccharomyces cerevisiae. *FEMS Yeast Res.* **6**, 171–176 (2006).

42.   H. Murakami, G. Blobel, D. Pain, Isolation and characterization of the gene for a yeast
mitochondrial import receptor. *Nat. 1990 3476292* **347**, 488–491 (1990).

43.   V. Zara, *et al.*, Yeast mitochondria lacking the phosphate carrier/p32 are blocked in
phosphate transport but can import preproteins after regeneration of a membrane
potential. *Mol. Cell. Biol.* **16**, 6524–6531 (1996).

44.   A. Harden, W. J. Young, The alcoholic Ferment of Yeast-juice.Part III.-The Function of
Phosphates in the Fermentation o f Glucose hy Yeast-juice. *Proc. R. Soc. London. Ser.*
B, **80**, 299–311 (1908).

45.   A. Timoń-Gómez, M. Proft, A. Pascual-Ahuir, Differential Regulation of Mitochondrial
Pyruvate Carrier Genes Modulates Respiratory Capacity and Stress Tolerance in Yeast.
PLoS One **8**, e79405 (2013).

46.   T. Bender, G. Pena, J.-C. Martinou, Regulation of mitochondrial pyruvate uptake by
alternative pyruvate carrier complexes. *EMBO J.* **34**, 911–924 (2015).

47.   P. Grigaitis, B. Teusink, An excess of glycolytic enzymes under glucose-limited
conditions may enable Saccharomyces cerevisiae to adapt to nutrient availability.
FEBS Lett. (2022) https:/doi.org/10.1002/1873-3468.14484 (November 28, 2022).

48.   W. Yi, *et al.*, PFK1 Glycosylation Is a Key Regulator of Cancer Cell Growth and Central
Metabolic Pathways. *Science* **337**, 975 (2012).

49.   H. Miyazawa, *et al.*, Rewiring of embryonic glucose metabolism via suppression of
PFK-1 and aldolase during mouse chorioallantoic branching. *Dev.* **144**, 63–73 (2017).

50.   W. D. Hollinshead, *et al.*, Examining Escherichia coli glycolytic pathways, catabolite
repression, and metabolite channeling using Δpfk mutants. *Biotechnol. Biofuels* **9**, 1–
13 (2016).

51.   F. H. Westheimer, Why Nature Chose Phosphates. *Science (80-.).* **235**, 1173–1178
(1987).

52.   S. C. L. Kamerlin, P. K. Sharma, R. B. Prasad, A. Warshel, Why nature really chose
phosphate. *Q. Rev. Biophys.* **46**, 1–132 (2013).

53.   J. W. Booth, G. Guidotti, Phosphate Transport in Yeast Vacuoles *. *J. Biol. Chem.* **272**,

20408–20413 (1997).

54.   T. Vila, S. Frases, F. M. Gomesid, Lessons from protozoans: Phosphate sensing and

polyphosphate storage in fungi. *PLOS Pathog.* **18**, e1010298 (2022).

55.   M. E. Solesio, *et al.*, Depletion of mitochondrial inorganic polyphosphate (polyP) in

mammalian cells causes metabolic shift from oxidative phosphorylation to glycolysis.

Biochem. J. **478**, 1631–1646 (2021).

56.   M. Brin, R. W. McKee, The Inhibition of Respiration by Glucose, Fructose, and

Mannose in the Ehrlich Mouse Ascites Tumor. *Cancer Res.* **16**, 364–368 (1956).

57.   R. Orij, S. Brul, G. J. Smits, Intracellular pH is a tightly controlled signal in yeast.

Biochim. Biophys. Acta - Gen. Subj. **1810**, 933–944 (2011).

58.   R. Dechant, *et al.*, Cytosolic pH is a second messenger for glucose and regulates the

PKA pathway through V-ATPase. *EMBO J.* **29**, 2515–2526 (2010).

59.   P. Hamel, *et al.*, Redundancy in the function of mitochondrial phosphate transport in

Saccharomyces cerevisiae and Arabidopsis thaliana. *Mol. Microbiol.* **51**, 307–317

(2004).

60.   F. Jia, *et al.*, Overexpression of Mitochondrial Phosphate Transporter 3 Severely

Hampers Plant Development through Regulating Mitochondrial Function in

Arabidopsis. *PLoS One* **10**, e0129717 (2015).

61.   H. Pelicano, D. S. Martin, R. Xu, P. Huang, Glycolysis inhibition for anticancer

treatment. 4633–4646 (2006).

62.   S. O'Neill, R. K. Porter, N. McNamee, V. G. Martinez, L. O'Driscoll, 2-Deoxy-D-Glucose

inhibits aggressive triple-negative breast cancer cells by targeting glycolysis and the

cancer stem cell phenotype. *Sci. Rep.* **9**, 1–11 (2019).

63.   G. Soto-Heredero, M. M. Gómez de las Heras, E. Gabandé-Rodríguez, J. Oller, M.

Mittelbrunn, Glycolysis – a key player in the inflammatory response. *FEBS J.* **287**,

3350–3369 (2020).

64.   R. Shiratori, *et al.*, Glycolytic suppression dramatically changes the intracellular

metabolic profile of multiple cancer cell lines in a mitochondrial metabolism-

dependent manner. *Sci. Rep.* **9**, 1–15 (2019).

65.   C. L. Lu, *et al.*, Tumor Cells Switch to Mitochondrial Oxidative Phosphorylation under

Radiation via mTOR-Mediated Hexokinase II Inhibition - A Warburg-Reversing Effect.

PLoS One **10**, e0121046 (2015).

66.   G. Cheng, *et al.*, Mitochondria-targeted drugs synergize with 2-deoxyglucose to
trigger breast cancer cell death. *Cancer Res.* **72**, 2634–2644 (2012).

67.   M. Scheibye-Knudsen, B. Quistorff, Regulation of mitochondrial respiration by
inorganic phosphate; comparing permeabilized muscle fibers and isolated
mitochondria prepared from type-1 and type-2 rat skeletal muscle. *Eur. J. Appl.*
Physiol. **105**, 279–287 (2009).

68.   E. L. Seifert, E. Ligeti, J. A. Mayr, N. Sondheimer, G. Hajnóczky, The mitochondrial
phosphate carrier: Role in oxidative metabolism, calcium handling and mitochondrial
disease. *Biochem. Biophys. Res. Commun.* **464**, 369–375 (2015).

69.   J. B. J.P. van Dijken, *et al.*, An interlaboratory comparison of physiological and genetic
properties of four Saccharomyces cerevisiae strains. *Enzyme Microb. Technol.* **26**,
706–714 (2000).

70.   M. S. Longtine, *et al.*, Additional modules for versatile and economical PCR-based
gene deletion and modification in Saccharomyces cerevisiae. *Yeast* **14**, 953–961
(1998).

71.   V. Vengayil, Z. Rashida, S. Laxman, The E3 ubiquitin ligase Pib1 regulates effective
gluconeogenic shutdown upon glucose availability. *J. Biol. Chem.* **294**, 17209–17223
(2019).

72.   M. Sriariyanun, P. Mutrakulcharoen, S. Tepaamorndech, K. Cheenkachorn, K.
Rattanaporn, A Rapid Spectrophotometric Method for Quantitative Determination of
Ethanol in Fermentation Products. *Orient. J. Chem.* **35**, 744–750 (2019).

73.   A. Walvekar, Z. Rashida, H. Maddali, S. Laxman, A versatile LC-MS / MS approach for
comprehensive , quantitative analysis of central metabolic pathways [version 1 ; peer
review : 2 approved]. 1–15 (2018).

74.   P. C. Liao, I. R. Boldogh, S. E. Siegmund, Z. Freyberg, L. A. Pon, Isolation of
mitochondria from Saccharomyces cerevisiae using magnetic bead affinity
purification. *PLoS One* **13**, 1–15 (2018).

75.   W. W. Chen, E. Freinkman, D. M. Sabatini, Rapid immunopurification of mitochondria
for metabolite profiling and absolute quantification of matrix metabolites. *Nat.*
Protoc. **12**, 2215 (2017).

Figure 1: A deubiquitinase deletion screen identifies Ubp3 as a regulator of glucose-mediated mitochondrial repression

Figure 1: A deubiquitinase deletion screen identifies Ubp3 as a regulator of glucose-mediated mitochondrial repression

A) Schematic depicting glucose-mediated mitochondrial repression (Crabtree effect).

B) Schematic describing the screen with a yeast DUB KO library to identify regulators of Crabtree effect.

C) Identifying DUB knockouts with altered mitochondrial potential. Heat map shows relative mitochondrial membrane potential of 19 deubiquitinase deletions in high glucose, from 2 biological replicates. Also see figures S1A, S1B.

D) The deubiquitinase activity of Ubp3 and repression of mitochondrial membrane potential. WT, *ubp3Δ,* and Ubp3^{C469A} were grown in high glucose and relative mitochondrial membrane potential was measured. Data represent mean ± SD from three biological replicates (n=3). Also see figure S1D.

E) Effect of loss of Ubp3 on ETC complex IV subunit Cox2. WT and *ubp3Δ* were grown in high glucose, and Cox2 was measured (western blot using an anti-Cox2 antibody). A representative blot (out of 3 biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD.

F) Basal oxygen consumption rate (OCR) in high glucose in *ubp3Δ*. WT and *ubp3Δ* were grown in high glucose, and OCR was measured. Basal OCR corresponding to ~3 x 10^5 cells, from two independent experiments (n=2), normalized to the OD$_{600}$ is shown. Bar graph representations are shown in the inset. Data represent mean ± SD.

G) Total ATP levels in *ubp3Δ* and WT. WT and *ubp3Δ* were grown in high glucose, and total ATP were measured. Data represent mean ± SD from three biological replicates (n=3).

H) Dependence of *ubp3Δ* on mitochondrial ATP. WT and *ubp3Δ* cells were grown in high glucose, and treated with 1 mM sodium azide for 45 minutes. Total ATP levels in sodium azide treated and untreated cells were measured. Data represent mean ± SD (n=3).

I) Requirement for mitochondrial respiration in high glucose in *ubp3Δ*. Serial dilution growth assay in high glucose in the presence/absence of sodium azide (1 mM) are shown. Also see figure S1F.

Data information: **p<0.01, ***p<0.001.

Figure 2: Key glycolytic enzymes decrease and glucose flux is rerouted in *ubp3Δ* cells

Figure 2: Key glycolytic enzymes decrease and glucose flux is rerouted in *ubp3Δ* cells

A) A schematic illustrating directions of glucose-6-phosphate (G6) flux in cells. Glucose is converted to G6P, a precursor for trehalose, the pentose phosphate pathway (PPP), and glycolysis.

B) Effect of loss of Ubp3 on key glycolytic enzymes. WT and *ubp3Δ* were grown in high glucose and the Pfk1, Tdh2, and Tdh3 levels were measured by western blot using an anti-FLAG antibody. A representative blot (out of three biological replicates, n=3) and their quantification are shown. Data represent mean ± SD. Also see figure S2A.

C) Steady-state metabolite amounts in WT and *ubp3Δ* in high glucose. Relative steady-state levels of trehalose, major glycolytic, and PPP intermediates were estimated in WT and *ubp3Δ*. Data represent mean ± SD from three biological replicates (n=3). Also see Appendix Table S3.

D) Relative glycolytic and trehalose synthesis flux in WT and *ubp3Δ*. Relative ^{13}C-label incorporation into trehalose and glycolytic intermediates, after a pulse of 1% $^{13}C_6$ glucose is shown. Data represent mean ± SD from three biological replicates (n=3). Also see Appendix Table S3, figures S2B, S2C.

E) Ethanol production in *ubp3Δ*. WT and *ubp3Δ* were grown in high glucose and ethanol in the media was measured. Data represent mean ± SD from three biological replicates (n=3).

F) Relative rate of ethanol production in WT vs *ubp3Δ*. WT and *ubp3Δ* were grown in high glucose (to OD_{600} ~ 0.6), equal numbers of cells were shifted to fresh medium (high glucose) and ethanol concentration in the medium was measured temporally. Data represent mean ± SD from three biological replicates (n=3)

Data information: *p<0.05, **p<0.01, ***p<0.001.

Figure 3: Rerouted glucose flux results in phosphate (Pi) accumulation

Figure 3: Rerouted glucose flux results in phosphate (Pi) accumulation

A) Changes in protein levels in *ubp3Δ* (dataset from Isasa et al., 2015)*. ubp3Δ* cells have an increase in proteins involved in mitochondrial respiration and decrease in proteins involved in glucose and phosphate metabolism.

B) Schematic showing maintenance of Pi balance during glycolysis. Trehalose synthesis from G6P releases Pi, and the conversion of G3P to 1,3BPG by GAPDH consumes Pi. In *ubp3Δ*, trehalose biosynthesis (which releases Pi) increases. *ubp3Δ* have decreased GAPDH, which will decrease Pi consumption. This increase in Pi release along with decreased Pi consumption could increase cytosolic Pi.

C) Intracellular Pi levels in WT and *ubp3Δ*. WT and *ubp3Δ* were grown in high glucose and the total free phosphate (Pi) levels were estimated. WT in high Pi (2% glucose, 10mM Pi) was a positive control. Data represent mean ± SD from three biological replicates (n=3). Also see figure S3A.

D) Pho regulon responses in WT and *ubp3Δ*. Protein levels of Pho84-FLAG and Pho12-FLAG were compared between WT grown in high glucose and in high Pi, *ubp3Δ* in high glucose with or without a shift to a no-Pi medium for one hour, by western blot. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD.

E) Contribution of trehalose synthesis as a Pi source. WT, *tps2Δ, ubp3Δ,* and *ubp3Δtps2Δ* were grown in high glucose and the total Pi levels were estimated. Data represent mean ± SD from three biological replicates (n=3). Also see figure S3B.

F) Loss of GAPDH isozymes Tdh2 and Tdh3 and effect on Pi. WT, *ubp3Δ,* and *tdh2Δtdh3Δ* were grown in high glucose and total Pi was estimated. Data represent mean ± SD from three biological replicates (n=3).

G) Pi levels in *ubp3Δ* and *tdh2Δtdh3Δ* cells in ethanol medium. WT, *ubp3Δ,* and *tdh2Δtdh3Δ* cells were grown in ethanol medium and the total Pi levels were estimated. Data represent mean ± SD from three biological replicates (n=3).

Data information: *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 4: Mitochondrial Pi availability correlates with mitochondrial activity in *ubp3Δ*

Figure 4: Mitochondrial Pi availability correlates with mitochondrial activity in *ubp3Δ*

A) A hypothetical mechanism of cytosolic free Pi controling mitochondrial activity by regulating mitochondrial Pi availability.

B) Mitochondrial Pi amounts in WT vs *ubp3Δ*. Mitochondria were isolated by immunoprecipitation from WT and *ubp3Δ* and mitochondrial Pi estimated. Mitochondrial Pi levels (normalised to Idh1) are shown. Data represent mean ± SD from three biological replicates (n=3). Also see figure S4A, B.

C) Cox2 protein in *tdh2Δtdh3Δ*. WT, *ubp3Δ,* and *tdh2Δtdh3Δ* were grown in high glucose and Cox2 protein was estimated. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD.

D) Basal OCR levels in *tdh2Δtdh3Δ*. WT, *ubp3Δ,* and *tdh2Δtdh3Δ* were grown in high glucose and basal OCR was measured from two independent experiments (n=2). Data represent mean ± SD. Also see figure S4C.

E) Comparative Pi amounts and Cox2 levels in *ubp3Δ*, *tdh2Δtdh3Δ,* WT cells. WT cells were grown in high glucose, *ubp3Δ* and *tdh2Δtdh3Δ* were grown in high glucose and low Pi, and Cox2 protein was estimated. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD. Also see figure S4D, S4F.

F) Pi amounts and basal OCR in *ubp3Δ* and *tdh2Δtdh3Δ* vs WT cells. WT cells were grown in high glucose, *ubp3Δ* and *tdh2Δtdh3Δ* were grown in high glucose and low Pi, and basal OCR was measured from two independent experiments (n=2). Data represent mean ± SD.

G) Effect of loss of mitochondrial Pi transporter Mir1 on Cox2protein. WT, *ubp3Δ*, *mir1Δ,* and *mir1Δubp3Δ* were grown in high glucose and Cox2 amounts compared. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD.

H) Relationship of mitochondrial Pi transport and basal OCR in WT vs *ubp3Δ*. WT, *ubp3Δ*, *mir1Δ,* and *mir1Δubp3Δ* cells were grown in high glucose and basal OCR was measured from two independent experiments (n=2). Data represent mean ± SD.

Data information: *p<0.05, **p<0.01, ****p<0.0001.

Figure 5: Mitochondrial Pi availability constrains mitochondrial activity under high glucose

Figure 5: Mitochondrial Pi availability constrains mitochondrial activity under high glucose

A) Relationship of mitochondrial Pi transport and respiration after glucose removal. WT and *mir1Δ* cells were cultured in high (2%) glucose and shifted to low (0.1%) glucose for 1 hour. The normalized basal OCR, from two independent experiments (n=2) are shown. Data represent mean ± SD.

B) Requirement of mitochondrial Pi transport for switch to respiration upon glycolytic inhibition by 2DG. WT and *mir1Δ* cells were cultured in high glucose and treated with or without 0.25% 2DG for 1 hour. Basal OCR was measured from two independent experiments (n=2). Data represent mean ± SD. Also see figure S5A.

C) Glucose-dependent regulation of Mir1. Cells (with Mir1-HA) were grown in high glucose and shifted to low glucose (0.1% glucose) for 1 hour, and Mir1 levels compared. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD. Also see figure S5B.

D) Increasing Pi concentrations and mitochondrial activity in isolated mitochondria. Mitochondria were isolated from WT cells grown in high glucose, incubated with 1 mM pyruvate, 1 mM malate, 0.5 mM ADP and 0-50 mM KH_2PO_4. The mitochondrial activity was estimated by mitotracker fluorescence intensity, and intensities relative to the sample with 0 mM KH_2PO_4 is shown. Data represent mean ± SD from three biological replicates (n=3).

E) Effect of overexpressing Mir1 on Cox2 protein. WT (containing empty vector) and Mir1 overexpressing (Mir1OE) cells were grown in high glucose and Cox2 levels were estimated. A representative blot (out of three biological replicates, n=3) and their quantifications are shown. Data represent mean ± SD. Also see figure S5F.

F) Effect of overexpressing Mir1 on basal OCR. The basal OCR in WT (containing empty vector) and Mir1OE in high glucose was measured from three independent experiments (n=3). Data represent mean ± SD.

G) Mitochondrial Pi transport requirement for growth after 2DG treatment. Shown are serial dilution growth assays in high glucose in the presence and absence of 0.1% 2DG, using WT and *mir1Δ* cells. The results after 40hrs incubation/30°C are shown.

H) A model illustrating how mitochondrial Pi availability controls mitochondrial activity. In high glucose, the decreased Pi due to high Pi consumption in glycolysis, along with the glucose-mediated repression of mitochondrial Pi transporters, decreases mitochondrial Pi availability. This reduces mitochondrial activity. In low glucose, increased mitochondrial Pi transporters and lower glycolytic flux increases mitochondrial Pi, leading to enhanced mitochondrial activity. In *ubp3Δ* cells in high glucose, high trehalose synthesis and lower glycolytic flux results in an increase in Pi. This increases mitochondrial Pi availability and thereby the mitochondrial activity.

Data information: *p<0.05, **p<0.01, ***p<0.001.