How benthic sediment microbial communities respond to glyphosate and its metabolite: A microcosm experiment

Christine M Cornish[1] (0000-0002-4600-0669), Peter Bergholz[2,3] (0000-0002-3420-5261), Kaycie Schmidt[2], and Jon Sweetman[1,4] (0000-0002-9849-7355)

[1]Department of Biological Sciences and [2]Department of Microbiological Sciences, North Dakota State University, Fargo, ND 58105, United States

[3]Institute for Quantitative Health Sciences & Engineering, and Department of Microbiology & Molecular Genetics, Michigan State University, East Lansing, MI 48823, United States

[4]Department of Ecosystem Science and Management, The Pennsylvania State University, University Park, PA 16802, United States

Corresponding author: Christine Cornish, christine.cornish@ndsu.edu

Acknowledgments

Thank you to the North Dakota Established Program to Stimulate Competitive Research for funding this research. Thank you to Dr. López-Martínez for providing support and usage of equipment. Additionally, thank you to the anonymous reviewers for their time and feedback of our manuscript.

Abstract

Glyphosate is the most commonly used agricultural herbicide in the world. In aquatic

ecosystems, glyphosate often adsorbs to benthic substrates or is metabolized and degraded by

microorganisms. The effects of glyphosate on microbial communities varies widely as

microorganisms respond differently to exposure. To help understand the impacts of glyphosate

on the sediment microbiome we conducted a microcosm experiment examining the responses of

benthic sediment microbial communities to herbicide treatments. Sediments from a prairie

pothole wetland were collected and 16S rRNA gene sequencing was used to analyze community

composition 2-hours and 14-days after a single treatment of low (0.07 ppm), medium (0.7 ppm),

or high (7 ppm) glyphosate, aminomethylphosphonic acid (glyphosate metabolite), or a

glyphosate-based commercial formula. We found no significant differences in microbial

community composition between treatments, concentration levels, or time. These findings

suggest that microbial species in the Prairie Pothole Region of North America may be tolerant to

glyphosate exposure.

Keywords: glyphosate, AMPA, sediments, microorganisms, wetlands, microcosms

Introduction

Agrochemical contamination of aquatic ecosystems is an ongoing concern due to the direct and indirect risks to environmental health. Glyphosate (i.e., Roundup®) is a non-selective, systemic herbicide that has become the most commonly used herbicide in the world since the 1990s [1]. The substantial use of glyphosate has resulted in its widespread and frequent detection in surface waters and groundwater [2, 3], where benthic sediments often become sinks [4]. Microbial metabolism is the primary degradation mechanism of glyphosate [5] resulting in its metabolites, aminomethylphosphonic acid (AMPA) or phosphate and sarcosine [6].

Glyphosate targets higher plants through the shikimate pathway [7], but this pathway is also present in some microorganisms [8]. Therefore, it was initially presumed that glyphosate may have inadvertent effects on microbial communities. Studies have reported that glyphosate can have a wide range of negative, positive, or neutral impacts on microbial community composition and function in terrestrial and aquatic ecosystems. Aquatic microbial communities have been shown to utilize glyphosate as a nutrient source resulting in increased activity, whereas in others, microbes are inhibited by toxicological effects. Pérez et al. [9] found 6 mg L^{-1} and 12 mg L^{-1} Roundup® caused a significant decrease in average microphytoplankton and nanophytoplankton abundance, whereas Lu et al. [10] found significantly enhanced gene expression in mechanisms potentially related to glyphosate tolerance, but did not find overall community structure shifts from 2.5 mg L^{-1} glyphosate. Several species are known to exhibit glyphosate tolerance [6], in addition to species capable of using glyphosate directly as a nutrient source [11, 12]. These examples demonstrate the variety of impacts glyphosate can have on aquatic microbial communities.

In addition to the impacts of glyphosate, AMPA, glyphosate's main metabolite, can also have direct and indirect effects on aquatic ecosystems [13]. Similar to glyphosate AMPA can be degraded by microorganisms, but it is more mobile and persistent [14–16]. It is also a known phytotoxin [17] with additional concerns regarding its potential to bioaccumulate [18]. AMPA is highly dependent on the presence and concentration of glyphosate [19], where both compounds frequently co-occur in areas of high agricultural intensity [5, 20]. Glyphosate and AMPA are most often detected in surface soil [21], and frequently transported to surface waters, including wetlands [2, 5] where they can impact benthic microorganisms.

The Prairie Pothole Region (PPR) is a large complex landscape covered with shallow wetlands and prairies [22]. This region is surrounded by agriculture, primarily corn and soybean croplands, which are the predominant crops glyphosate is used on [1, 2]. Consequently, wetlands in the PPR are subject to prolonged glyphosate contamination, where glyphosate has been reported as the most frequently detected and the highest detected herbicide in this region [23]. These wetlands are key ecosystems providing many economical and ecological services. For example, biogeochemical processes including C turnover and sequestration, N and P capture, and remediation of agrochemicals are essential ecosystem functions [24, 25], where benthic sediment microbial communities play a fundamental role in these ecosystem processes.

Understanding the impacts of agrochemical contamination on microbial community structure is vital because shifts in microbial communities impact the whole ecosystem. This is especially important in a region with high agricultural intensity, like the PPR, because microorganisms are chronically exposed to chemical stressors like glyphosate. The objective of our research was to assess the responses of benthic sediment microbial communities from the PPR to glyphosate-based herbicide treatments. We hypothesized that 1) observed Operational

Taxonomic Units (OTUs) and microbial diversity would decrease at high herbicide concentration

due to toxicological effects, and 2) microbial community composition would shift at low

herbicide concentration in favor of species capable of metabolizing glyphosate.

Materials and Methods

Microcosm preparation

Surface sediment (~ 10 cm) was collected from a wetland (P1) within the Cottonwood

Lake Study Area (U.S. Fish and Wildlife service managed Waterfowl Production Area) of the

PPR located in Stutsman County, North Dakota. This area has minimal agricultural influence,

where native prairie grasslands and wetlands cover over 80% [26]. After collection, sediment

was stored refrigerated (~ 4 °C) until initiation of the experiment. Thirty-two 5.7 L microcosms

were prepared with the following contents: 1 cm layer of homogenized sediment, 2.5 L of

dechlorinated tap water, and covered with "no-see-um" mesh (Duluth Sport Nets, Duluth, MN).

Microcosms were then stored in incubators at 20 °C with a 16:8 (light:dark) hour photoperiod

and left to acclimate for one month prior to treatment. Over the entirety of the 6-week

experiment, microcosms were monitored for water evaporation and were filled back to the 2.5 L

volume when necessary.

Herbicide treatment and sediment sampling

We conducted a 3 x 4 factorial experiment with three replicates per treatment and five

controls ($n = 32$). Herbicide treatments consisted of analytical grade glyphosate (98.1% purity)

or AMPA (99% purity), or 41% glyphosate concentrate (commercial formula) at concentrations

of 0.07 parts per million (ppm), 0.7 ppm, or 7 ppm. Our concentrations were chosen based on the

U.S. maximum contaminant level (MCL) of glyphosate in drinking water (U.S. EPA 2015),

which is equivalent to our medium concentration of 0.7 ppm (or 7 mg L^{-1}). Treatments were

added to the water, lightly stirred to evenly distribute, and then allowed to settle for two hours. A

50 mL polypropylene corer (needle-less syringe) was used to collect sediment samples pre-

treatment, two hours post-treatment, and two weeks post-treatment. All samples were stored in

50 mL polypropylene tubes at -80 °C immediately after sampling until analyses were performed.

Analyses: Herbicide residues and microbial 16S sequencing

Sediment subsamples were shipped frozen on dry ice to the Agriculture and Food

Laboratory at the University of Guelph (Ontario, Canada) for glyphosate and AMPA residue

analysis. Liquid chromatography tandem mass spectrometry (LC-MS/MS) was used to quantify

glyphosate and AMPA residues with a limit of detection of 0.005 ppm and a limit of

quantification of 0.02 ppm for both compounds. Matrix blanks and standards were analyzed

alongside sediment samples for quality control (QC).

Sediment microbial communities were analyzed after extraction of environmental DNA

with the Qiagen DNeasy® PowerSoil® kit. Briefly, 0.25 g sediment (wet weight) was lysed using

PowerBead Tubes in a bead-beater (Biospec Mini-beadbeater-24). Kit solutions (C1 – C6) were

added stepwise to purify, wash, and elute DNA into 100 µL final volume. Microbial 16S rRNA

gene was PCR amplified using universal primers, 27F and U1492R, for QC to verify suitability

for sequencing. Sequencing library preparation and sequencing were performed according to

Oxford Nanopore Technologies 16S Barcoding Kit (SQK-RAB204) protocol and reagents.

Briefly, amplicons were cleaned using AMPure XP bead cleanup, quantified via PicoGreen™

analysis (Quant-iT PicoGreen™ Kit, modified from Invitrogen's Quant-iT PicoGreen™ dsDNA

Reagent and Kits protocol), and pooled to obtain a final DNA concentration of 50 – 100 ng.

Pooled samples contained up to twelve separately barcoded samples for sequencing using an

Oxford Nanopore minION™. Each library was run for approximately four hours. Raw fast5

reads were basecalled and demultiplexed using Guppy v3.4. EPI2ME 16S workflow

(https://epi2me.nanoporetech.com, rev 2.1.0) was used for QC and initial characterization. For

QIIME2 analysis, sequences were pre-processed using MetONTIIME [27] and QIIME2 was

used to filter and analyze the resulting OTU table abundances. The SILVA v138 database was

downloaded and utilized as a reference database for taxonomic identification [28, 29]. Taxonomy

barplots, describing the composition of each sample at the desired taxonomic level were

visualized at QIIME2 view (https://view.qiime2.org/). Sequence tables were filtered to remove

taxonomically unassigned sequences. A table of absolute OTU abundances was exported in

BIOM format for further analysis in R.

Statistical analysis

A feature-table containing absolute abundances of Family level OTUs from QIIME2 was

imported into R (version 4.0.2) using the R/biomformat package (1.24.0) [30]. Unassigned taxa

and taxa that were present in $\leq$ 10% of samples were removed to minimize the risk of sequencing

error carryover and due to low statistical power to analyze underrepresented OTUs, respectively.

All subsequent analyses were conducted using the R/stats (3.6.2) and R/vegan (2.6-2) packages

[31, 32]. Shannon's H. diversity index was calculated using the "diversity" function and

observed OTUs were calculated using the "specnumber" function. Differences in diversity and

observed OTUs between herbicide treatments, concentration levels, and time of sampling were

analyzed with a Kruskal-Wallis Rank Sum test using the "kruskal.test" function. Lastly, Bray-

Curtis dissimilarity distance measures were calculated and square root transformed for non-

metric multidimensional scaling (NMDS) using the "metaMDS" function to display community

structure.

Results

After unassigned and rare taxa were removed, a total of 430 OTUs were detected across

all sediment samples ($n = 96$). Relative abundances across treatments were not significantly

different (Fig. 1). The three most abundant families in all treatments were *Hydrogenophilaceae*

in the *Pseudomonadota* (*Gammaproteobacteria*) Phylum, and *Sulfurovaceae* and

Desulfobacteraceae, which are assigned to the *Proteobacteria* (*Deltaproteobacteria*) Phylum.

Prolixibacteraceae, a family within the *Bacteroidota* Phylum thought to be associated with

sediment, was only detected in the control and glyphosate treatments.

Fig. 1 Relative abundance of microbial Families among all samples ($n = 96$) in each herbicide
treatment; legend only shows the top three most abundant Families across all herbicide
treatments

Shannon H index and observed OTUs were similar among all treatments and controls

over the entirety of the experiment (Fig. 2 and Fig. 3). Kruskal-Wallis tests showed no

significant differences in alpha diversity metrics between herbicide treatments, concentration

levels, or time of sampling (Table 1). A two-dimensional ordination solution was reached (stress

= 0.1793708) using NMDS. Microbial communities were similar in multivariate space, which

showed there were no significant compositional differences between treatments (Fig. 4).

Fig. 2 Shannon H diversity index quantified from microcosm sediments of herbicide treatments

and treatment levels at each sampling timepoint

Fig. 3 Observed OTUs quantified from microcosm sediments of herbicide treatments and

treatment levels at each sampling timepoint

Table 1 Summary of Kruskal-Wallis Rank Sum test on the effects of treatments on Shannon

diversity and observed OTU. χ^2 = Chi-square value, df = degrees of freedom, and P = the p-

value where the significance level is 0.05

Quantified Metric	Treatment	χ^2	df	P
	Time of Sampling	1.81	2	0.40
Shannon Diversity	Herbicide Treatment	4.67	3	0.20
	Concentration	3.48	3	0.32
	Time of Sampling	2.25	3	0.52
Observed OTU	Herbicide Treatment	0.22	2	0.90
	Concentration	2.45	3	0.48

Fig. 4 NMDS plot of sediment microbial community structure across all samples ($n = 96$) where

shapes represent day of sampling and colors represent herbicide treatment

Both glyphosate and AMPA concentrations in sediments increased over time within each

herbicide treatment, but not in control treatments. Additionally, AMPA was detected in

microcosms where glyphosate or commercial formula only was added, confirming that

glyphosate degradation did occur (Online Resource 1).

Discussion

The current study used microcosms to examine the impact of glyphosate-based herbicide

treatments on benthic sediment microbial communities from a prairie pothole wetland. Both

glyphosate and AMPA were detected in microcosm sediments shortly following treatment

indicating that residues rapidly dissipate out of the water column [5, 15]. AMPA was also

detected in sediments of microcosms that only received glyphosate (i.e., glyphosate or

commercial formula treatment), but no direct AMPA treatment, indicating that glyphosate

degradation occurred [33]. In our experimental microcosms, we found that microbial community

diversity and composition were not significantly affected by glyphosate, AMPA, or commercial

formula at any concentration, which was contrary to our original hypotheses. Our highest

treatment level (7 ppm) was a magnitude higher than the U.S. EPA's MCL for drinking water,

and approximately 41 times higher than the average concentration that have been reported in

wetland sediments in the PPR [23]. Some research suggests that microbial responses are more

dependent on previous glyphosate exposure history and application rates [18, 34]. The present

experiment reflects conditions of acute glyphosate and AMPA exposure, rather than long-term

exposure on the sediment community. Sediments used in our study were collected from a

permanent wetland in North Dakota with no routine pesticide usage in the immediate catchment.

Thus, agricultural inputs within this area are minimal to none compared to other areas of the

PPR. However, we found no differences between treatments and controls suggesting that

glyphosate-based herbicides may not have adverse effects on sediment microbial communities

from wetlands in this area of the PPR. This observed lacked effect may be the result of

glyphosate- and AMPA-based selection pressure.

A lack of a response from microbial communities following glyphosate exposure has also

been observed previously in the literature. For example, Pesce et al. [35] exposed natural spring-

and summer-collected riverine microbial communities to 10 μg L^{-1} glyphosate and found no

effect on bacterial activity or diversity. Lane et al. [36] conducted a six-month soil incubation

experiment with monthly glyphosate treatments at 59 µg g^{-1} and 118 µg g^{-1}, and found no

significant glyphosate effect on community structure represented by the relative abundances of

functional microbial groups. Muturi et al. [37] used 20 mg L^{-1} glyphosate treatment in

microcosms and found no differences in water microbial diversity or richness after 3- and 7-

days. Dennis et al. [38] found no significant effects of a single glyphosate treatment at the

recommended field application rate (33.03 mg L^{-1}) on bacterial or archaeal evenness, richness,

and composition after 60-days incubation. These studies in addition to the current study

represented a wide range of glyphosate concentrations, which all showed that glyphosate does

not always have direct toxic effects on microbial communities. Presumably, this may be due to

the presence of glyphosate-tolerant microbes.

Several studies have, however, shown shifts in microbial communities after either short-

or long-term glyphosate exposure. Lu et al. [10], for example, found increases in Shannon and

Simpson diversity of lacustrine microbial communities, in addition to differences in community

structure 10- and 15-days post-treatment of 2.5 mg L^{-1} glyphosate. Widenfalk et al. [39] also

found that an environmentally relevant glyphosate concentration (150 µg kg^{-1} dry weight) caused

significant shifts in lake sediment bacterial community composition in treated microcosms

relative to controls after 31-days. Sura et al. [40] found pelagic and biofilm bacterial production

in outdoor mesocosms was significantly inhibited by 225 µg L^{-1} glyphosate compared to

controls. Microbial communities in our study and Sura et al. [40] were both collected from

wetlands within the PPR, however our results were not consistent potentially due to differences

in sediment- versus water-associated communities, land use within our collection site's

watersheds, or incubation versus outdoor experimental design. These studies all showed that

glyphosate can have various negative and positive effects on microbial communities, whereas

our results showed glyphosate can also have no effect. This discrepancy indicates that there are

many underlying complexities on the effects of glyphosate at the microbial level, which may be

more related to molecular mechanisms and/or environmental variables.

Some microorganisms naturally express a glyphosate-resistant form of the EPSP enzyme

and there are many glyphosate-tolerant microbial species listed in the literature [41, 42].

Specifically, *Agrobacterium* sp. strain CP4 was the bacteria used for the original glyphosate-

resistant EPSP gene in glyphosate-resistant crops [43]. Whereas, other species have evolved

tolerance through mutations of the EPSP synthase, or metabolic or detoxifying processes [44],

which can be a result of prolonged or repeated glyphosate exposure. For example, Tang et al.

[34] added 100 mg L^{-1} glyphosate to sediments with high, low, and no previous glyphosate

exposure, and found that microbes degraded glyphosate quicker in sediments with high exposure

history compared to low exposure history. Additionally, Lane et al. [36] found significantly

higher microbial respiration in soils after glyphosate treatment, specifically in soils with previous

glyphosate exposure history. The sediments used in the present study were collected from an

agriculturally-undisturbed wetland located in an area with no known extensive glyphosate use

(David Mushet, Research Wildlife Biologist, USGS Northern Prairie Wildlife Research Center,

email communication June 21, 2022), thus microbial communities should not have any

prolonged exposure history. We did find multiple species across all treatments that are capable of

glyphosate degradation and mineralization including *Cyanobiaceae*, *Enterobacteriaceae*,

Pseudomonadaceae, and *Rhizobiaceae* (e.g., genus *Agrobacterium*). While our sediment

collection site has no known history of glyphosate use within the immediate watershed, the

majority of the PPR is in an agriculturally intensive area where glyphosate use is common.

Therefore, microbial species from the regional species pool may have evolved tolerance despite

our study wetland and its catchment having no existing agricultural pesticide use.

Glyphosate is highly water soluble, which also allows it to be easily transported into

wetlands, where environmental variables can play a role in its bioavailability and toxicity to

microorganisms. Glyphosate has an affinity for oxides and organic matter content resulting in its

rapid adsorption to sediments [45]. However, glyphosate and phosphate can compete for

sediment binding sites due to their chemical structural similarities [46]. Thus, lower phosphate

can increase glyphosate's sediment adsorption capacity [47], subsequently decreasing its

bioavailability. Temperature can also affect glyphosate's environmental persistence, where

higher temperatures facilitate degradation due to increased microbial activity [48, 49]. While the

present study did not measure sediment physicochemical characteristics, temperature and light

conditions were controlled for, and homogenized sediments were used for all microcosms to help

minimize variability.

Conclusion

We evaluated the responses of benthic sediment microbial communities to a single

addition of low, medium, or high glyphosate-based herbicide treatment after 14-days incubation.

We expected to see toxicological-induced shifts potentially in favor of tolerant species, however

we found no differences in sediment microbial communities among treatments or concentration

after 2 weeks. Our results may be explained by the lower concentration of bioavailable

glyphosate in the sediments compared to the actual concentrations added. Additionally, our

results may suggest glyphosate-tolerance in the benthic sediment microbial communities, but that

is inconclusive without further investigation of the presence of the glyphosate-resistant EPSP

gene within the microbiome. Our research suggests that in the PPR the direct effects of

glyphosate on sediment microorganisms may not be as severe as initially presumed. However,

the literature continues to reveal new implications of the extensive use of glyphosate on aquatic

ecosystems. Therefore, further research is still necessary to determine the full range of potential

effects of glyphosate on sediment microbial communities.

References

1. Benbrook CM (2016) Trends in glyphosate herbicide use in the United States and globally. Environ Sci Eur 28:3. https://doi.org/10.1186/s12302-016-0070-0

2. Coupe RH, Kalkhoff SJ, Capel PD, Gregoire C (2012) Fate and transport of glyphosate and aminomethylphosphonic acid in surface waters of agricultural basins. Pest Manag Sci 68:16–30. https://doi.org/10.1002/ps.2212

3. Belden JB, Hanson BR, McMurry ST, et al (2012) Assessment of the Effects of Farming and Conservation Programs on Pesticide Deposition in High Plains Wetlands. Environ Sci Technol 46:3424–3432. https://doi.org/10.1021/es300316q

4. Warren N, Allan IJ, Carter JE, et al (2003) Pesticides and other micro-organic contaminants in freshwater sedimentary environments—a review. Applied Geochemistry 18:159–194. https://doi.org/10.1016/S0883-2927(02)00159-2

5. Battaglin WA, Meyer MT, Kuivila KM, Dietze JE (2014) Glyphosate and Its Degradation Product AMPA Occur Frequently and Widely in U.S. Soils, Surface Water, Groundwater, and Precipitation. J Am Water Resour Assoc 50:275–290. https://doi.org/10.1111/jawr.12159

6. Zhan H, Feng Y, Fan X, Chen S (2018) Recent advances in glyphosate biodegradation. Appl Microbiol Biotechnol 102:5033–5043. https://doi.org/10.1007/s00253-018-9035-0

7. Cobb AH, Reade JPH (2010) Herbicides and Plant Physiology: Cobb/Herbicides and Plant Physiology. Wiley-Blackwell, Oxford, UK

8. Herrmann KM, Weaver LM (1999) THE SHIKIMATE PATHWAY. 32

9. Pérez GL, Torremorell A, Mugni H, et al (2007) Effects of the Herbicide Roundup on Freshwater Microbial Communities: A Mesocosm Study. Ecological Applications 17:2310–2322. https://doi.org/10.1890/07-0499.1

10. Lu T, Xu N, Zhang Q, et al (2020) Understanding the influence of glyphosate on the structure and function of freshwater microbial community in a microcosm. Environmental Pollution 260:114012. https://doi.org/10.1016/j.envpol.2020.114012

11. Forlani G, Pavan M, Gramek M, et al (2008) Biochemical Bases for a Widespread
Tolerance of Cyanobacteria to the Phosphonate Herbicide Glyphosate. Plant and Cell
Physiology 49:443–456. https://doi.org/10.1093/pcp/pcn021

12. Shushkova TV, Vasilieva GK, Ermakova IT, Leontievsky AA (2009) Sorption and
microbial degradation of glyphosate in soil suspensions. Appl Biochem Microbiol 45:599–
603. https://doi.org/10.1134/S0003683809060040

13. Grandcoin A, Piel S, Baurès E (2017) AminoMethylPhosphonic acid (AMPA) in natural
waters: Its sources, behavior and environmental fate. Water Research 117:187–197.
https://doi.org/10.1016/j.watres.2017.03.055

14. Blake R, Pallett K (2018) The Environmental Fate and Ecotoxicity of Glyphosate. outlook
pest man 29:266–269. https://doi.org/10.1564/v29_dec_08

15. Degenhardt D, Humphries D, Cessna AJ, et al (2012) Dissipation of glyphosate and
aminomethylphosphonic acid in water and sediment of two Canadian prairie wetlands.
Journal of Environmental Science and Health, Part B 47:631–639.
https://doi.org/10.1080/03601234.2012.668459

16. Kjaer J, Olsen P, Ullum M, Grant R (2005) Leaching of Glyphosate and Amino-
Methylphosphonic Acid from Danish Agricultural Field Sites. J Environ Qual 34:608–620.
https://doi.org/10.2134/jeq2005.0608

17. Reddy KN, Rimando AM, Duke SO, Nandula VK (2008) Aminomethylphosphonic Acid
Accumulation in Plant Species Treated with Glyphosate. J Agric Food Chem 56:2125–
2130. https://doi.org/10.1021/jf072954f

18. Bai SH, Ogbourne SM (2016) Glyphosate: environmental contamination, toxicity and
potential risks to human health via food contamination. Environ Sci Pollut Res 23:18988–
19001. https://doi.org/10.1007/s11356-016-7425-3

19. Beecraft L, Rooney R (2021) Bioconcentration of glyphosate in wetland biofilms. Science
of The Total Environment 756:143993. https://doi.org/10.1016/j.scitotenv.2020.143993

20. Malaj E, Liber K, Morrissey CA (2020) Spatial distribution of agricultural pesticide use and
predicted wetland exposure in the Canadian Prairie Pothole Region. Science of The Total
Environment 718:134765. https://doi.org/10.1016/j.scitotenv.2019.134765

21. Okada E, Costa JL, Bedmar F (2016) Adsorption and mobility of glyphosate in different
soils under no-till and conventional tillage. Geoderma 263:78–85.
https://doi.org/10.1016/j.geoderma.2015.09.009

22. Dahl TE (2014) Status and trends of prairie wetlands in the United States 1997 to 2009.
U.S. Department of the Interior, Fish and Wildlife Service, Ecological Services,
Washington, D.C.

23. McMurry ST, Belden JB, Smith LM, et al (2016) Land use effects on pesticides in
347   sediments of prairie pothole wetlands in North and South Dakota. Science of The Total
348   Environment 565:682–689. https://doi.org/10.1016/j.scitotenv.2016.04.209

349   24. Holloway JM, Goldhaber MB, Mills CT (2011) Carbon and nitrogen biogeochemistry of a
350   Prairie Pothole wetland, Stutsman County, North Dakota, USA. Applied Geochemistry
351   26:S44–S47. https://doi.org/10.1016/j.apgeochem.2011.03.025

352   25. Dalcin Martins P, Hoyt DW, Bansal S, et al (2017) Abundant carbon substrates drive
353   extremely high sulfate reduction rates and methane fluxes in Prairie Pothole Wetlands.
354   Glob Change Biol 23:3107–3120. https://doi.org/10.1111/gcb.13633

355   26. Winter TC (2003) Hydrological, chemical, and biological characteristics of a prairie pothole
356   wetland complex under highly variable climate conditions: the Cottonwood Lake area, east-
357   central North Dakota. U.S. Dept. of the Interior, U.S. Geological Survey : U.S. Geological
358   Survey, Information Services [distributor], Denver, CO

359   27. Maestri, Cosentino, Paterno, et al (2019) A Rapid and Accurate MinION-Based Workflow
360   for Tracking Species Biodiversity in the Field. Genes 10:468.
361   https://doi.org/10.3390/genes10060468

362   28. Pruesse E, Quast C, Knittel K, et al (2007) SILVA: a comprehensive online resource for
363   quality checked and aligned ribosomal RNA sequence data compatible with ARB. Nucleic
364   Acids Research 35:7188–7196. https://doi.org/10.1093/nar/gkm864

365   29. Quast C, Pruesse E, Yilmaz P, et al (2012) The SILVA ribosomal RNA gene database
366   project: improved data processing and web-based tools. Nucleic Acids Research 41:D590–
367   D596. https://doi.org/10.1093/nar/gks1219

368   30. McMurdie PJ, Paulson JN (2015) biomformat: An interface package for the BIOM file
369   format

370   31. Oksanen J, Simpson G, Blanchet F, et al (2017) Vegan: Community Ecology Package

371   32. R Core Team (2022) R: A language and environment for statistical computing

372   33. Gill JPK, Sethi N, Mohan A (2017) Analysis of the glyphosate herbicide in water, soil and
373   food using derivatising agents. Environ Chem Lett 15:85–100.
374   https://doi.org/10.1007/s10311-016-0585-z

375   34. Tang FHM, Jeffries TC, Vervoort RW, et al (2019) Microcosm experiments and kinetic
376   modeling of glyphosate biodegradation in soils and sediments. Science of The Total
377   Environment 658:105–115. https://doi.org/10.1016/j.scitotenv.2018.12.179

378   35. Pesce S, Batisson I, Bardot C, et al (2009) Response of spring and summer riverine
379   microbial communities following glyphosate exposure. Ecotoxicology and Environmental
380   Safety 72:1905–1912. https://doi.org/10.1016/j.ecoenv.2009.07.004

36. Lane M, Lorenz N, Saxena J, et al (2012) The effect of glyphosate on soil microbial activity, microbial community structure, and soil potassium. Pedobiologia 55:335–342. https://doi.org/10.1016/j.pedobi.2012.08.001

37. Muturi EJ, Donthu RK, Fields CJ, et al (2017) Effect of pesticides on microbial communities in container aquatic habitats. Sci Rep 7:44565. https://doi.org/10.1038/srep44565

38. Dennis PG, Kukulies T, Forstner C, et al (2018) The effects of glyphosate, glufosinate, paraquat and paraquat-diquat on soil microbial activity and bacterial, archaeal and nematode diversity. Sci Rep 8:2119. https://doi.org/10.1038/s41598-018-20589-6

39. Widenfalk A, Bertilsson S, Sundh I, Goedkoop W (2008) Effects of pesticides on community composition and activity of sediment microbes – responses at various levels of microbial community organization. Environmental Pollution 152:576–584. https://doi.org/10.1016/j.envpol.2007.07.003

40. Sura S, Waiser M, Tumber V, et al (2012) Effects of Glyphosate and Two Herbicide Mixtures on Microbial Communities in Prairie Wetland Ecosystems: A Mesocosm Approach. J Environ Qual 41:732–743. https://doi.org/10.2134/jeq2011.0376

41. Priestman MA, Funke T, Singh IM, et al (2005) 5-Enolpyruvylshikimate-3-phosphate synthase from Staphylococcus aureus is insensitive to glyphosate. FEBS Letters 579:728–732. https://doi.org/10.1016/j.febslet.2004.12.057

42. Firdous S, Iqbal S, Anwar S, Jabeen H (2018) Identification and analysis of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) gene from glyphosate-resistant Ochrobactrum intermedium Sq20: Identification of EPSPS from Ochrobactrum intermedium Sq20. Pest Manag Sci 74:1184–1196. https://doi.org/10.1002/ps.4624

43. Padgette SR, Kolacz KH, Delannay X, et al (1995) Development, Identification, and Characterization of a Glyphosate-Tolerant Soybean Line. Crop Sci 35:1451–1461. https://doi.org/10.2135/cropsci1995.0011183X003500050032x

44. Hertel R, Gibhardt J, Martienssen M, et al (2021) Molecular mechanisms underlying glyphosate resistance in bacteria. Environmental Microbiology 23:2891–2905. https://doi.org/10.1111/1462-2920.15534

45. Dion HM, Harsh JB, Hill Jr. HH (2001) Competitive sorption between glyphosate and inorganic phosphate on clay minerals and low organic matter soils. Journal of Radioanalytical and Nuclear Chemistry 249:385–390. https://doi.org/10.1023/A:1013222704311

46. de Jonge H, de Jonge LW, Jacobsen OH, et al (2001) Glyphosate sorption in soils of different pH and phosphorus content. Soil Science 166:230–238. https://doi.org/10.1097/00010694-200104000-00002

47. Guijarro KH, Aparicio V, De Gerónimo E, et al (2018) Soil microbial communities and glyphosate decay in soils with different herbicide application history. Science of The Total Environment 634:974–982. https://doi.org/10.1016/j.scitotenv.2018.03.393

48. Helander M, Saloniemi I, Saikkonen K (2012) Glyphosate in northern ecosystems. Trends in Plant Science 17:569–574. https://doi.org/10.1016/j.tplants.2012.05.008

49. Muskus AM, Miltner A, Hamer U, Nowak KM (2022) Microbial community composition and glyphosate degraders of two soils under the influence of temperature, total organic carbon and pH. Environmental Pollution 297:118790. https://doi.org/10.1016/j.envpol.2022.118790

Statements and Declarations

Funding

This work was supported by a STEM Research and Education Seed Grant from North Dakota Established Program to Stimulate Competitive Research (FAR0032100).

Competing interests

The authors have no competing interests to declare that are relevant to the content of this study.

Author contributions

All authors contributed to the study conception and design. Material preparation, data collection, and microbial analysis were performed by Christine Cornish and Kaycie Schmidt. The first draft of the manuscript was written by Christine Cornish with additions from Jon Sweetman and Peter Bergholz. All authors commented on previous versions of the manuscript, and read and approved the final manuscript.

Availability of data and materials

The data used in this study are available from the corresponding author upon reasonable request.

Ethics approval

443   Not applicable.

444   *Consent to participate*

445   All authors have given consent to participate.

446   *Consent for publication*

447   All authors have given consent to publish.