MANUSCRIPT TITLE

Differential roles of positive and negative supercoiling in organizing the *E. coli* genome

AUTHORS

Ziqi Fu[1], Monica S. Guo[2,*], Weiqiang Zhou[1,*] and Jie Xiao[3,*]

[1] Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205, U.S.A.
[2] Department of Microbiology, University of Washington, Seattle, WA 98198, U.S.A.
[3] Department of Biophysics and Biophysical Chemistry, Johns Hopkins School of Medicine, Baltimore, MD 21205, U.S.A.

* To whom correspondence should be addressed. Email: msguo@uw.edu, wzhou14@jhu.edu, and xiao@jhmi.edu

ABSTRACT

This study aims to explore whether and how positive and negative supercoiling contribute to the three-dimensional (3D) organization of the bacterial genome. We used recently published *Escherichia coli* GapR ChIP-seq and TopoI ChIP-seq (also called EcTopoI-seq) data, which marks positive and negative supercoiling sites, respectively, to study how positive and negative supercoiling correlates with the corresponding contact frequencies obtained from chromosome conformation capture sequencing (Hi-C and 5C). We found that supercoiled chromosomal loci have overall higher Hi-C contact frequencies than sites that are not supercoiled, with positive supercoiling surprisingly corresponding to higher spatial contacts than negative supercoiling. Additionally, Hi-C contact frequencies alone could identify positive, but not negative, supercoiling with high accuracy. The majority of positive and negative supercoils coincide with highly active transcription units, with a minor group likely associated with replication and other genomic processes. Our results suggest that both positive and negative supercoiling enhance chromosome interactions, but positive supercoils contribute more than negative supercoils to bring distant chromosomal loci closer in space. Based on these results, we propose new physical models of how the *E. coli* chromosome is organized differentially by positive and negative supercoils.

INTRODUCTION

  All living cells contain DNA that is compacted 1000-10,000-fold in a nuclear space. In bacterial cells, DNA is packaged mainly through supercoiling, the twisting and crossing of the double-stranded

DNA about itself, and the binding of nucleoid-associated proteins (NAPs), while eukaryotic cells use

histones to organize their DNA within a nuclear envelope (1, 2). Supercoiling can be categorized as

negative or positive, depending on the direction of the twists with respect to that of the DNA double

helix, with both types of supercoiling forming coils (twists) or plectonemes (writhes) (3). Recently,

chromosome conformation capture technologies such as Hi-C allow high-resolution interrogation of

genome organization by measuring the physical interaction frequency of any two DNA fragments (4, 5, 6,

7). Despite revolutionizing our understanding of the overall architecture of chromosomes and revealing

chromosome sub-structures such as DNA loops and topologically associated domains (TADs), Hi-C, like

most other genomic technologies, captures DNA sequences but not DNA shapes. Consequently, how

supercoils underlie the three-dimensional organization of chromosomes remains poorly understood.

In *E. coli*, previous work has revealed that the chromosome is organized into six ~800 kilobase-

long macrodomains, analogous to eukaryotic TADs, and that NAPs play differential roles in making short

and long-range interactions (4, 5, 8). Our understanding of supercoiling in chromosome organization,

however, is more limited. Supercoiling occurs during transcription as unwinding of the DNA leads to

formation of positive supercoiling ahead of RNA polymerase and negative supercoiling behind it, referred

to as the "twin-domain model" (9, 10). Replication also generates significant positive supercoiling ahead

of the replisome (22). Recent experiments using psoralen, a DNA intercalator that specifically binds to

undertwisted, negatively supercoiled DNA, indicates that negative supercoiling is pervasive in *E. coli*

chromosomes (11) and organizes the local DNA (± 25 kb) around transcription units (12). These psoralen

studies further suggest that the *E. coli* macrodomains may be organized by supercoiling. However,

because psoralen only interrogates negatively supercoiled DNA, the contribution of positive supercoiling

to these organizational effects can only be indirectly inferred.

Two recent supercoiling-mapping technologies, GapR-seq (13) and EcTopoI-seq (14), have

largely overcome the technical limits of the previous psoralen approach, generating high-resolution (<1

kb) snapshots of the location and magnitude of positive and negative supercoiling, respectively. Both

technologies rely on chromatin immunoprecipitation sequencing (ChIP-seq) to probe the genomic

locations at which two shape-recognizing proteins, *Caulobacter crescentus* GapR and *E. coli*

Topoisomerase I (EcTopoI) bind, respectively. GapR was shown to preferentially bind to overtwisted,

positively supercoiled DNA and serve as a sensor for such DNA in *Caulobacter*, *E. coli* and yeast (13,

15). Similarly, EcTopoI, an enzyme that binds and removes excessive negative supercoils (16), was

shown to localize to regions of negative supercoiling (14). GapR-seq and EcTopo-seq discriminate

between these two DNA shapes and have an inverted enrichment pattern around transcription units, where

EcTopo-seq labels negative supercoiling upstream of transcription while GapR-seq identifies positive

supercoiling downstream of transcription, validating the twin-domain model and providing a whole-

genome portrait of supercoiling (14).

In this study we computationally integrate GapR-seq, EcTopoI-seq, RNA-seq and Hi-C/5C

datasets to describe how positive and negative supercoiling contributes to the three-dimensional

organization of the *E. coli* genome (**Fig. 1A**). Our statistical analyses reveal that genomic regions

containing supercoiling are associated with higher-frequency spatial interactions than those not containing

supercoiling. These supercoiling-associated contacts can span entire macrodomains and have not been

previously reported. Strikingly, we discovered that positive supercoiling is correlated with higher spatial

contact than negative supercoiling. Using a machine-learning model, we found that positively, but not

negatively, supercoiled loci can be accurately predicted from Hi-C data alone, suggesting that positive

supercoiling is a major driver of chromosome architecture. Lastly, we observed that while most

supercoiled loci colocalize with active transcription units as expected, a minor set does not, suggesting

additionally genomic processes in generating supercoils and contributing to chromosome packaging. Our

results suggest that positive supercoiling is pervasive and plays a dominant role in global genome

organization.

MATERIAL AND METHODS

Hi-C/5C data processing

We align the Hi-C/5C sequence reads to the *E. coli* reference genome NC_000913.3 and follow

the Hi-C-Pro 3.1.0 pipeline (17). The resolution is 5 Kb for the data set generated in this study and the

one by Lioy et. al., and it is 0.5 Kb for the data set generated by Cockram et al. The contact count

matrices are normalized using the iterative correction and eigenvector decomposition (ICED) method

(18). The ICED contact counts are log2 transformed after adding a pseudo count. Then, we use the R

package ComplexHeatmap (19) to produce all heatmaps.

ChIP-seq data processing and analysis

For the two ChIP-seq data sets, we first perform a quality check to the sequence reads using

FastQC (20) and eliminate adapters using Trimmomatics (21). Bowtie2 (22) is used to align all sequence

reads to the *E. coli* reference genome NC_000913.3, follow by conversion of SAM to BAM files using

Samtools (23). ChIP-seq peaks are detected using MACS2's callpeak function (24). To obtain a set of

reproducible ChIP-seq peaks, a customized R script (see **Data Availability**) is used to obtain peak signals

across replicates.

Hi-C/5C bin classification based on ChIP-seq signals

A 5 Kb genomic bin in the Hi-C/5C data is classified as PS (positive supercoiling site) or NS (negative supercoiling site) if it contains the center of at least one GapR or TopoI peak, respectively. The complementary set of genomic bins to the set of PS and NS bins are labeled as aPS (absent of a PS-site) and aNS (absent of a NS-site), respectively. Bins that contain only GapR peak centers but no Topo peak center are defined as PS-only (positive supercoiling only) bins, while NS-only (negative supercoiling only) bins contain only Topo peak centers but no GapR peak center. Bins that lack any peak center are labeled as aa (absent of either). Similarly, a genomic bin of size 0.5 Kb is classified as PS or NS if it has a non-empty overlap with at least one GapR or TopoI peak, respectively. Conversely, aPS and aNS are assigned to the set of bins that do not overlap with any GapR or TopoI peaks, respectively. This assignment strategy, which is more relaxed compared with the one used for the 5 Kb bins, is implemented to resolve severe class imbalance issues caused by false negative identification of PS and NS. **Supplementary Data** Table S1 and Figure S1D display the summary statistics for the number and the width of the peaks.

Training the random forest models

We align Hi-C/5C sequence reads to the E. coli reference genome NC_000913.3, following the Hi-C-Pro 3.1.0 pipeline (27) with iterative correction and eigenvector decomposition (ICED) normalization. The resulting contact counts are log2-transformed with a pseudo count to form a contact frequency matrix. To prepare the feature matrix, we transform the contact frequency matrix such that its diagonal becomes the middle column, representing the self-contacts. Each column represents a genomic separation distance from a given locus (row), which serves as a feature for our model. The dimension of the feature matrix is 928 by 928, determined by the maximum Hi-C resolution. The response columns, or target vectors, are labeled according to each locus' classification as PS/aPS and NS/aNS, as described in the bin classification step. Each row represents a genomic locus with a binning size of 5 Kb, and the model predicts whether this locus contains any GapR or TopoI peaks. It is worth noting that the genomic location of each bin (or row) remains inaccessible to the models, as we generated the testing sets and their corresponding training sets randomly. In other words, each row is considered as an independent data point by the models.

We then build separate models for predicting positive and negative supercoiling. To train these models, we partition the feature matrix into four balanced and non-overlapping sets and build four series of models using each of the quarters as testing data. For each testing set, we build random forest models by using 25%, 50%, or 75% data from its complement in the feature matrix as the training data. The

process was repeated 10 times for each training data set. We use the R package caret (25) for random

forest training, and each random forest is trained using 5-fold cross-validation with 10 repeats. We

optimize the mTry parameter (from 2 to 42 by a step of 5) and nTrees parameter (from 200 to 1200 by a

step of 100) using the receiver operating characteristic (ROC) metric.

Evaluating random forest model prediction performance

After training each model, we evaluated its performance by making predictions on an

independent testing set. That is, we ask the model to predict the PS/aPS and NS/aNS labels of an unseen

set of loci using only their Hi-C frequencies. For each testing dataset and prediction percentage (25%,

50%, and 75%), we made predictions using the 10 trained models and computed the average AUROC and

AUPRC. The AUPRC metric was used in addition to AUROC to account for any potential class

imbalance issues (26).

AUROC (Area Under the Receiver Operating Characteristic Curve) is a created by plotting the

true positive rate against the false positive rate for different threshold values. While AUROC may take

any values between 0 to 1, AUROC = 1 indicates a perfect model and AUROC = 0.5 indicates a random

classifier, which typically serves as a theoretical baseline performance. AUPRC (Area Under the

Precision-Recall Curve) is created by plotting the ratio of true positive predictions to the total number of

positive predictions (precision) against the ratio of true positive predictions to the total number of actual

positive cases (recall). The AUPRC ranges between 0 and 1. AUPRC = 1 indicates a perfect classifier,

while its baseline score is the ratio of positive cases. AUPRC = 0 indicates that the model's precision is

always zero, regardless of the recall. Both metrics measure the classification performance of the random

forest model, and AUPRC is a more reliable metric in class imbalance cases.

To provide a baseline for comparison, we generated a set of models using the same feature

matrices as the actual models, but with randomly permuted response/target vectors. These models were

then tested on unseen testing sets that were correctly labeled (the same as the actual models). The training

process for these baseline models was the same as that used for the actual models. We calculated the

AUROC and AUPRC for these baseline models and use them as references.

Growth conditions and chemical treatment for *E. coli* used in Hi-C

E. coli was grown in LB (10 g/L NaCl, 10 g/L tryptone, 5 g/L yeast extract) at 37 °C with shaking

at 200 rpm. Optical density was measured at 600 nm using a Genesys 10 Bio Spectrophotometer.

Chromosome conformation capture deep-sequencing (Hi-C)

Hi-C experiments were performed as described previously with the following modifications (7).

Wild-type *E. coli* K-12 MG1655 was grown to OD600 ~0.2 and then fixed with 3% formaldehyde for 10

min at room temperature before quenching with 0.375 M final concentration glycine for 15 min. Fixed

cells were pelleted, washed twice with 1X M2 salts (6.1 mM Na_2HPO_4, 3.9 mM KH_2PO_4, 9.3 mM

NH_4Cl), then washed 2 more times with 1X TE buffer (10 mM Tris-HCl [pH 8.0] and 1 mM EDTA),

before resuspension in 1X TE at ~10^7 cells / µL. Resuspended cells were divided into 25 µL aliquots

before storage at -80 °C. Each Hi-C experiment was performed with 2 x 25 µL aliquots. To each 25 µL

aliquot, 20 000 U of Ready-Lyse Lysozyme (Lucigen) was added, and the mixture was incubated for 30

min before addition of SDS to a final concentration of 0.25% for 15 additional minutes to completely

dissolve cell membranes and release chromosomal DNA.

Each aliquot of DNA was then digested with HpaII (NEB) as follows. A 50 uL reaction was

assembled containing lysed cells, rCutSmart buffer, and 5 µL of 10% Triton-X 100, which was allowed to

sit for 15 min at RT to inactivate SDS. Subsequently, 100 U of HpaII was added and the DNA was

digested at 37 °C for 2 hr. The reaction was cooled on ice before labelling cleavage overhangs with

Biotin-14-dCTP (ThermoFisher) in a 60 uL reaction (digestion reaction, 0.9 µL dGTP, dTTP and dATP at

2 mM each, 4.5 µL 0.4 mM biotin-14-dCTP, 1.2 µL of Klenow Large fragment [5000 units/ml]) for 45

min at room temperature. Reactions were quenched by adding 0.125% final concentration of SDS and

incubated for 5 min at room temperature. Blunted DNAs were ligated in dilute conditions as follows: A

master mix containing 75 µL 10% Trition-X 100, 100 µL 10X T4 Ligase Buffer (NEB), 2.5 µL of 20 mg

/ mL BSA (NEB), and 800 µL water per blunting reaction was assembled and chilled on ice. This chilled

ligase mix was added to the blunted DNAs and incubated for 15 min at room temperature to inactivate

SDS. Ligation reactions were then chilled on ice for 15 min before addition of 3 µL T4 DNA ligase

(NEB, 2,000,000 U / mL). Reactions were mixed well and incubated at 10-16°C for 4 hr.

After ligation, 10 mM final concentration EDTA and 2.5 µL of 20 mg / mL Proteinase K (NEB)

was added, and reactions incubated at 65 °C overnight to digest proteins and reverse crosslinks. DNA was

then extracted twice with phenol:chloroform:isoamyl alcohol, isopropanol precipitated, and then the

entirely of the DNAs from the two original aliquots was resuspended in 60 µL of water. Next, non-ligated

biotinylated DNAs were eliminated in a 100 µL reaction (60 µL purified DNA, 10 µL Buffer #2 (NEB),

0.5 µL 20 mg / mL BSA (NEB), 5 µL each of 2 mM dATP and dTTP) with 0.5 µL of T4 DNA

polymerase for 2 hr at 12 °C. DNA was sheared using a Bioruptor (Diagenode) and size-selected by gel

extraction to be between 200 and 500 bp.

Paired-end Illumina libraries were generated as in (Guo et al, 2021) as follows using DNA

LoBind tubes (Eppendorf) (13). First, DNA was end-repaired with 4 µL T4 DNA polymerase (NEB), 4

µL T4 PNK (NEB), and 0.75 µL Klenow large fragment (NEB) in 100 µL T4 DNA ligase buffer with

0.25 mM final concentration dNTPs for 30 min at room temperature. Repaired DNA was recovered by

Ampure XP (Beckman Coulter) bead purification, using 100 µL beads in 300 µL 20% PEG/NaCl

solution. Beads were washed twice with 80% ethanol, dried, and resuspended in 32 µL EB (10 mM Tris-

HCl [pH 8.0]). The bead slurry was directly treated with 3 µL Klenow (3'→5' exo-) (NEB) in 40 µL

reaction with 0.2 mM ATP at 37 °C for 30 min to add 3' overhangs to DNA. Repaired DNA was

recovered by Ampure XP capture and resuspended in 15 µL EB without beads. Y-shaped adaptors were

prepared by annealing Illumina PE adapter 1 and Illumina PE adapter 2. Y-shaped adapters were added to

bead slurry, and the mix was ligated in 25 µL total volume with 5 µL 15 µM ligated adapters and 1.5 µL

T4 DNA Ligase (NEB, 400,000 U / mL) for 30 hr at room temperature, with occasional mixing. Biotin-

labeled junctions were purified from non-labelled junctions with Dynabeads MyOne Streptavidin C1

(ThermoFisher). Dynabeads were first washed twice with 1X NTB (5 mM Tris-HCl [pH 8.0], 0.5 mM

EDTA and 1 M NaCl) then added to ligation and mixed well. Biotin-labeled DNAs were allowed bind for

30 min at room temperature with occasional mixing before pulldown with a magnet. The beads were

washed three times with 100 µL 1X NTB, then washed with 100 µL water, resuspended in 10 µL water,

transferred a new tube, and stored at -20 °C. DNA libraries were amplified in 50 µL final volume with 2X

KAPA HiFi Master Mix (Roche) supplemented with 5 % final concentration DMSO (Sigma) and

appropriate barcoded primers. The total number of cycles was optimized for each sample to minimize the

number of cycles required for library generation. Libraries were purified by two-step Ampure XP capture

to recover 200-500 bp amplified libraries. DNA was recovered from Ampure beads by resuspending in 20

µL 10 mM Tris-HCl (pH 8.0). Paired-end sequencing was performed with a 91nt kit on a NextSeq500 at

the MIT Bio Micro Center.

RESULTS

Positive supercoiling sites correlate with higher-than-average Hi-C contact frequencies in *E. coli*

Although supercoiling is known to alter DNA shape, how supercoiling underlies the three-

dimensional organization of bacterial chromosomes is unknown. To understand how supercoiling

contributes to chromosomal organization, we first sought to compare *E. coli* chromosome architecture

deduced from Hi-C mapping with a recently published map of *E. coli* positive supercoiling using GapR-

seq (LB, 37°C) (13). From the GapR-seq data, we identified 609 reproducible GapR-binding peaks (see

Material and Methods, ChIP-seq data processing) along the *E. coli* genome. Because GapR

preferentially binds to overtwisted double-stranded DNA (dsDNA), these peak positions correspond to

positively supercoiled sites (13, 15) and hereafter we term them as PS (positive supercoiling) sites. Next,

we integrated these PS-sites with three chromosome architecture datasets: two publicly available 5C and

Hi-C datasets (referred to as Hi-C data below) (4, 5), in addition to one independent Hi-C experiment we

performed in this study. We divided the *E. coli* genome of 4.6 million base (Mb) pairs into either 928 (5

Kb per bin) or 9284 (0.5 Kb per bin) bins based on the maximum resolution of the respective Hi-C

datasets (**Fig. 1B**). Each 5 Kb bin and 0.5 Kb bin was then assigned a tag, PS/aPS (presence/absence of a

PS-site, see **Material and Methods**, HiC/5C bin classification based on ChIP-seq signals and

Supplement Fig. S1D, Table 1 for summary statistics of ChIP-seq peaks).

When we compared the chromosome contacts of bins containing positive supercoiling (PS, **Fig.**

1C, D, dense black bars) to those lacking positive supercoiling (aPS, **Fig. 1C, D**, sparse black bars), we

found that PS bins appeared to have high contact frequencies with genomic loci hundreds of kilobases

away than aPS bins. To qualitatively illustrate this observation, we calculated the sum of the contacts for

each locus within a 100 Kb to 1 Mb window (purple lines in **Fig. 1C, 1D**) and observed that the total

contacts are higher in genomic regions with dense PS-sites (at ~3 Mb in the genome) and lower in the

region with sparse PS-sites (at ~1 Mb in the genome) (**Fig. 1E, 1F**). We validated our observation in three

other 5C/Hi-C experimental replicates at both 5 Kb and 0.5 Kb resolution (**Supplement Fig. S1A**). The

length scale (~ Mb) of the separation between PS and aPS sites matches the scale of the *E. coli*

macrodomains (8).

To quantify? the differences in DNA association between positively supercoiled and non-

positively supercoiled loci, we computed the inverse of the normalized Hi-C contact frequency between

any two bins and plotted the averages against the linear genomic distance between the loci pairs (**Fig. 1G,**

H). This inversed Hi-C contact count serves as a proxy for the spatial separation because it is proportional

to the spatial distance between two chromosomal sites (27). Our results show that this apparent spatial

separation between two PS bins (**Fig. 1G, 1H,** PS-PS, blue) was smaller than that of all bins at all

separation ranges (all-all, red). Conversely, the spatial separation between two aPS bins (**Fig. 1G, 1H**,

aPS-aPS, green) was larger than the spatial separation of all bins at all distances. The relative difference

in the spatial separation between the PS-PS and aPS-aPS curves (([aPS-aPS] - [PS-PS])/max([aPS-aPS] -

[PS-PS])) peaks near the 500 Kb loci separation distance, remaining high until 1 Mb, and then gradually

decreases (**Fig. 1I, 1J**). We observed a similar trend and pattern on other 5C/Hi-C replicates conducted in

different growth conditions and with different resolutions (**Supplement Fig. S1A-B**). Taken together,

these data support our observation that positively supercoiled loci interact more frequently than sites that

are not positively supercoiled and that this effect is most significant at a macrodomain scale (500 Kb – 1

Mb).

Next, to verify that the observed pattern is indeed dependent on positive supercoiling, we re-

analyzed the Hi-C data using a mutant GapR (GapR[1-76]) dataset. GapR[1-76] lacks the GapR dimerization

domain required for tetramerizing on positively supercoiled DNA, and hence binds to DNA

nonspecifically (13, 15). GapR[1-76] binding sites are distributed uniformly along the *E. coli* genome (**Fig.**

1C, 1D, green bars) unlike GapR, which preferentially binds surrounding the *oriC* half of the

chromosome (see **Supplemental Table 1A** for definitions of *ori*C half and *ter* half). The difference in

spatial separation between GapR[1-76]-bound bins (PS[1-76]-PS[1-76]) and GapR[1-76] unbound bins (aPS[1-76]-aPS[1-

76]) is substantially diminished, validating that PS sites are specific to GapR binding and that they are

associated with long-distance contact (**Supplement Fig. S1C**). In summary, our analyses reveal that

positively supercoiled loci make more frequent spatial contacts than non-positively supercoiled loci, with

the largest effect on spatial association at a macrodomain scale.

Negative supercoiling sites correlate with higher-than-average Hi-C contact frequencies in *E. coli*

Bacterial genomes are typically net-negatively supercoiled (27), and previous research suggests

that this type of supercoiling is a main contributor for genome compaction (28). To explore how negative

supercoiling affects chromosome organization, we used an *Ec*TopoI-seq data set in which all *E. coli*

TopoI binding sites were identified using immunoprecipitation under the same growth condition as the 5

Kb Hi-C datasets (LB, 37°C) (14). *E. coli* TopoI removes excessive negative supercoiling and was

proposed to mark sites of negative supercoiling in *E. coli* (14). Because TopoI can also bind to RNA

polymerase during transcription (16), we excluded TopoI binding sites within a transcription unit (TU), in

order to focus exclusively on TopoI binding to negative supercoiling. Using the EcTopoI-seq dataset, we

identified 446 reproducible TopoI-binding sites and assigned accordingly the 5K *E. coli* genome bins as

either NS (negative supercoiling, 38% of total bins) or aNS (absent of a NS-site, 64% of total bins)

similarly as that for PS sites (**Fig. 1A, B**, see **Material and Methods**, ChIP-seq data processing &

HiC/5C bin classification based on ChIP-seq). NS bins along the chromosome have a largely similar

distribution as PS bins in that they are densely distributed surrounding the *ori*C half of the chromosome

(**Supplemental Table 1A**, *ori*C half and *ter* half), which contains multiple *rrn* operons, and sparsely

surrounding the *ter* half of the chromosome. Similar to what we observed for PS bins, genomic regions

with more frequent NS bins (**Fig. 1C, 1D,** dense blue bars) also showed a higher Hi-C contact frequencies

with other genomic loci a few hundred Kb away than the ones with sparse NS bins (sparse blue bars).

To quantify differences in DNA association between negatively supercoiled loci (NS) and those

absent of negatively supercoiled sites (aNS), we calculated the apparent spatial separation between two

negatively supercoiled bins along the chromosome (**Fig. 2A-2B**). Akin to our previous observations with

positive supercoiling, we found that the spatial separation between two bins containing negatively

supercoiled DNA (NS-NS, blue) is the lowest. This is followed by the spatial separation between all

genomic locations (all-all, red), with the largest spatial separation occurring between two non-negatively

supercoiled bins (aNS-aNS, green). Furthermore, the relative difference in spatial separation between NS-

NS and aNS -aNS sites follows the same trend as observed in the GapR analysis (**Fig 2C, D**). These

observations are also reproducible across all Hi-C experimental replicates, despite differences in the data

resolution, experimental conditions, and technologies. (**Supplement Fig. S2A-B**). Our analyses here

suggest that, as with positive supercoiling, two negatively supercoiled loci make more frequent spatial

contacts than non-negatively supercoiled loci, again with the largest effects on spatial association at

macrodomain scales.

Positive supercoiling has a larger effect on chromosomal DNA interaction than negative

supercoiling

Our analyses so far suggest that both positive and negative supercoiling play significant roles in

E. coli chromosome compaction (**Fig. 1G, H, 2C, D**). Interestingly, when we compared the apparent

spatial separations (for example, compare the y axes of **Fig. 1G** and **Fig. 2A**), positive supercoiling

appeared to have a greater effect than negative supercoiling, i.e. PS-PS spatial separations appear to be

closer than those of NS-NS sites. To quantify this difference, we assigned each 5 Kb or 0.5 Kb genomic

bin a label depending on what type of supercoiling, if any, dominants within that bin (PS-only, NS-only),

or is absent of either (aa) (see **Material and Methods**, HiC/5C bin classification based on ChIP-seq

signals). Bins containing both GapR and TopoI peaks were not used in the analysis (a total of 162 out of

928 5 Kb bins and 539 out of 9284 0.5 Kb bins). This mutually exclusive bin assignment allows us to

isolate the effects of each supercoiling type in our analysis. Plotting the apparent spatial separation within

these three groups confirmed that positively supercoiled bins (**Fig. 2E, F,** PSonly-PSonly, blue) associate

more frequently with each other than negatively supercoiled bins (NSonly-NSonly, green), with non-

supercoiled loci displaying the lowest frequency of association (aa-aa, red). Interestingly, at very long

distances (>1.25 Mb), the spatial separation of non-supercoiled bins (aa-aa) was slightly smaller than

negatively supercoiled bins and approached or became even smaller than positively supercoiled bins at

the largest linear distances (> 2 Mb). The significance of these differences is unclear and may require

further investigation. Importantly, the spatial separation between two positively supercoiled loci is always

lower than that of two negatively supercoiled loci, as demonstrated by the positive difference between

PSonly and NSonly's apparent spatial separation (**Fig. 2G, 2H**). Similar patterns were found in all other

independent Hi-C replicates (**Supplement Fig. S2B**). Based on these analyses, we conclude that while

positive and negative supercoils both package the chromosome, positive supercoiling leads to higher

spatial contacts than negative supercoiling, especially for contacts that occur on the scale of

macrodomains (~500 Kb to ~1.5 Mb, **Fig. 2G, 2H**).

Positive, but not negative, supercoiling sites can be inferred from Hi-C contact map

Our results suggest that positive and negative supercoiling generate distinct chromosome

organizational patterns that are captured in the Hi-C map. We wondered if supercoiling is a strong enough

underlying feature of chromosome organization that the locations of supercoils can be inferred from Hi-C

data. To investigate whether the Hi-C contact matrix encodes such supercoiling information, we utilized a

random forest model to predict positively and negative supercoiling loci based on the corresponding

contact frequencies. We chose random forest model because its prediction performance is robust under

limited sample points and a large number of covariates.

Our random forest models were first trained using the transformed Hi-C contact matrices and the

correct supercoiling label of each genomic loci (bin size = 5 Kb) obtained from the ChIP-seq data sets

(see **Material and Methods**, Training the random forest models and **Fig. 3A**, left). In the training, each

row of the input feature is the contact frequencies of a genomic locus with itself (brown square) and all

other flanking loci on both sides (blue squares); the output, or response, is whether this locus contains a

PS peak, NS peak, both, or none (green and purple boxes on the right). After the models were trained with

varying amounts of data, we ask the models to predict PS/aPS and NS/aNS bins for a set of unseen,

randomly distributed bins, based solely on their Hi-C contact frequencies with all other genomic loci (**Fig.**

3A, right). To evaluate the model prediction performance, we compared the average AUROC and

AUPRC computed on the testing set with those of the baseline models, which were trained on randomly

assigned supercoiling labels. The ROC (Receiver Operating Characteristic) curve plots the true positive

rates against the false positive rates at different decision thresholds for making a prediction, and AUROC

is the area under the ROC curve, which is commonly used to evaluate the performance of a prediction

model. An AUROC value of 1 means a perfect prediction model, and values between 0.5 and 1 indicates

the prediction is better than a random prediction model. The precision-recall curve (PRC) plots the

positive predictive values (precision) against the true positive rates (recall) for different threshold values,

and AUPRC summarizes the area under the precision-recall curve, which is an alternate metric for

evaluating the prediction performance. The AUPRC ranges between 0 and 1, where higher values indicate

better model performance. A value of 0 indicates that the model's precision is always zero, regardless of

the recall, and a value of 1 indicates perfect precision-recall trade-off. (see **Material and Methods**,

Evaluating random forest model prediction performance). The combination of the two parameters allows

the model classification to be robust against potential class imbalance issues, for example, the numbers of

NS and aNS bins are not equal.

We observed that the random forest model performed well for predicting positive supercoiling

using 75% of data for training, achieving an AUROC value of ~0.74. The model's prediction performance

remains consistently robust (AUROC = ~0.74) even when less data was used for training (50% and 25%),

outperforming the baseline models (where the target labels were permuted) where a low AUROC value of

0.43 was observed (**Fig. 3B,** and see baseline models definition **in Material and Methods,** Evaluating

random forest model prediction performance, Training the random forest models). Using AUPRC as the

evaluation metric yielded similar results, where our models trained with different data amounts achieved

an AUPRC of ~0.68, as compared to AUPRC of 0.44 for the baseline models (**Fig. 3C**). These

consistently robust performances across models with different training-to-testing data ratios suggest that

positively supercoiled loci have highly uniform, strong interaction patterns with other genomic loci, and

that such patterns distinguish them from loci without positive supercoiling.

To gain a deeper understanding of how the models can distinguish between PS (containing

positive supercoiling sites) and aPS (absent of positive supercoiling sites) bins, we computed the feature

importance using the Mean Decrease in Accuracy (MDA), a metric that measures how much accuracy the

model losses by excluding each feature. Features with high MDA values (in this case, Hi-C contact

frequencies at a certain linear distance along the chromosome) significantly contributes to the

classification of the associated genomic locus to PS or aPS. The MDA plot peaked at ~500 Kb, indicating

that spatial contact between loci separated by this distance is the greatest contributor to the correct

categorization of PS and aPS bins (**Fig. 3D**). This result confirms our finding that positively supercoiled

and non-positively supercoiled bins differ the most in their interactions with loci ~500 Kb away, and the

spatial interaction patterns at these separation ranges can robustly predict the existence of positive

supercoiling at any genomic loci.

Next, to determine whether Hi-C data can be used to predict negative supercoiling in a similar

fashion, we trained the random forest models using TopoI-seq data, following the same procedure as

described above. Surprisingly, we found that the model's performance was not as good as predicting

positive supercoiling, with AUROC ranging from 0.54 to 0.58 at different training-to-testing data ratios.

AUPRC scores from these models ranged from 0.41 to 0.44, which are indistinguishable from or slightly

worse than the baseline model with AURPC at 0.44 (**Fig. 3E, 3F**). Furthermore, the MDA plot did not

show a clear pattern as the positive supercoiling prediction model (**Fig. 3G**). These results indicate that

Hi-C contact frequencies contain some information but cannot effectively determine whether a locus is

negatively supercoiled, in contrast to what we observed with positive supercoiling (**Fig. 3B-3D**).

Together, these observations indicate that positive supercoiling enhances chromosomal loci spatial

interactions more than negative supercoiling.

We conclude this section by examining the reasons that explain the different prediction

performance of PS and NS peaks. The presence of significant difference between the interactions of PS

and aPS loci with all genomic loci (**Fig. 3H**, green and blue lines, **Supplement Fig. S3A**) explained the

robust prediction performance of PS sites. The poor performance of TopoI peak prediction could be

attributed to the lack of significant difference between the interaction pattern of NS and aNS with all

genomic loci (**Fig. 3I**, overlapping green and blue lines, **Supplement Fig. S3B**). Although there is a

significant difference between NS-NS and aNS-aNS interactions (**Supplement Fig. S3B**, green and

purple lines), this NS-all versus aNS-all interaction difference was not different from each other

accessible to the models as the features were not labeled as peak or non-peak. Instead, the models rely on

the difference between peak and non-peak with all genomic loci to make a classification (**Supplement**

Fig. S3B, yellow and blue lines). Since such a difference (peak-to-all versus non-peak-to-all) is

significant in PS sites but not NS sites, we conclude that recovering positive supercoiling signals, but not

negative supercoiling, is possible from using Hi-C datasets alone. Our findings suggest that positive

supercoiling has a characteristic signature that is reflected in the Hi-C contact map.

Transcription-independent and -dependent supercoiling

Transcription is a major supercoil-generator in bacteria (29, 30). During transcription, an

elongating RNA polymerase (RNAP) under-twists upstream DNA and over-twists downstream DNA,

generating negative and positive supercoiling, respectively (10). By examining the positions of genomic

supercoiling relative to the transcriptional activity of nearby transcription units (TUs), we can access the

capacity of transcription to generate persistent supercoiling and consequently its contribution to genome

packaging. We used a set of RNA-seq data (LB, 37°C) with 2640 TUs annotated on the *E.*

coli chromosome. We calculated the distance between a transcription end site (TES) and the nearest PS

site indicated by a GapR peak downstream, and the distance between a transcription start site (TSS) and

its nearest NS site (TopoI peak) upstream. The distributions of the distance between NS/PS site to its

nearest TSS/TES do not differ from each other significantly (p=0.13, Wilcoxon, **Fig. 4A, B**). The

majority (94.7% and 94.8%) of PS and NS sites are within 1.0 Kb of its nearest TES and TSS,

respectively. For comparison, on average 90.0% and 77.0% of random chromosomal sites are within the

same 1.0 Kb distance from the nearest TES or TSS, respectively, due to the close spacing between TUs

along the *E. coli* chromosome (**Fig. 4C, D** & **Supplement Fig. S4B, S4C**). This comparison demonstrates that the majority of supercoils are associated with transcription activities, and that it makes up to ~95% of total sustained genomic supercoils, hence transcription contributes significantly to genome organization.

Because all TUs make up to 92% of the *E. coli* genome in length (31), ~ 45% of the TU coverage overlap with either PS or NS sites (identified by GapR and TopoI ChIP-seq binding sites, respectively), but 65% of these overlapping regions are associated only with PS sites. Previous studies have shown that DNA supercoiling generated by active transcription may modulate the activity of RNA polymerase molecules on the same gene as well as impact the transcription of adjacent genes (31, 32, 33). Therefore, the observation that positive supercoils extend into adjacent TUs more frequently than negative supercoils suggests that positive supercoiling may play a larger role than negative supercoiling in regulating *E. coli* transcription. As transcription is largely suppressed by positive supercoiling but promoted by negative supercoiling, this differential regulation could maintain most TUs in a repressed state, which only turns on when needed to minimizing potential transcription cost.

We also observed that a minor subset of GapR and TopoI peaks (3.8% and 4.1% of all peaks, respectively) are located more than 1 Kb from any TUs (indicated by hex bins above the orange line, **Fig. 4A, 4B**). Approximately 72% and 79% of this subset associate with TUs with low transcriptional activities, respectively (expression level below the 75[th] percentile of all TUs, purple-highlighted quadrant, **Fig. 4A, 4B**). Furthermore, we observed that this minor group tend to be located near the *Ori* and *Ter* regions but are sparse in between (**Fig. 4E, 4F**). These patterns were not observed in the control analysis of randomly selected chromosomal sites, which were uniformly distributed across the genome as expected (**Fig. 4G, 4H**). These observations suggest that these supercoils may be introduced by genomic processes other than transcription since they are far from highly active TUs (expression above the 25[th] percentile of all TUs), but likely by replication and/or the actions of NAPs and topoisomerases (34).

Taken together, our analyses show that supercoils are mostly associated with transcription activities, in line with previous findings and suggest that transcription is a main driving force for genome structuring.

DISCUSSION

In this work, we integrated two recently published maps of *E. coli* positive and negative

supercoiling (GapR-seq and TopoI-seq, respectively) with three independent Hi-C data sets to explore

computationally the roles of positive and negative supercoiling in *E. coli* genome organization (**Fig. 1C-**

J). We observed a consistent, robust pattern across various Hi-C data sets that supercoiled genomic loci

have shorter spatial separation than non-supercoiled sites, with two positively supercoiled loci having

shorter spatial separations than two negatively supercoiled loci (**Fig. 2A-D, 2E-H**). Both positive and

negative supercoiling have the largest effects on spatial association of loci are separated by 0.5-1.5 Mb in

the linear chromosome, on the order of the *E. coli* macrodomains. Building upon these observations, we

developed a supervised machine learning model and found that Hi-C data can accurately predict

positively supercoiled sites but not negatively supercoiled sites, suggesting that the spatial interactions

mediated by positive supercoiling dominate Hi-C measurement (**Fig. 3A-I**). Finally, we found ~ 95% of

GapR and TopoI sites reside next to a TU, and that more positive supercoiling sites than negative

supercoiling sites are within the body of an adjacent TU. These observations suggest that transcription is

the major driver of supercoiling and genome organization. Taken together, our results support the

longstanding idea that supercoiling leads to chromosome packaging and suggest that, although positive

and negative supercoiling can be reductively considered to structure DNA similarly, their effects on

chromosome organization are not equivalent. Our work suggests that positive supercoiling enhances

spatial contact more than negative supercoiling, which has not been proposed (**Fig. 4A-F**).

The differential roles of positive and negative supercoiling in chromosome organization

The *E. coli* chromosome is naturally negatively supercoiled (27), which is necessary to allow

genomic processes such as replication or transcription to proceed without being energetically costly to

open the double-stranded chromosomal DNA. As such, it is reasonable to expect that positive

supercoiling is necessary to package the genome and compensate naturally existing negative supercoiling,

which may destabilize the chromosome by randomly introducing ssDNA regions due to stochastic

dsDNA melting.

Why would positive and negative supercoiling have differential effects on chromosome

organization? The mechanisms that regulate positive and negative supercoiling in *E. coli* are complex.

However, we note that there are many more abundant DNA structuring proteins in *E. coli* (e.g., HU, H-

NS, Fis) that are known to constrain negative supercoiling than those constrain positive supercoiling,

suggesting that the effects of negative supercoiling may be buffered by these DNA binding proteins (35,

36, 37). Hence, the role of negative supercoiling may be more pertinent to genomic processes while that

of positive supercoiling may be more pertinent to genome organization. Further investigation will be

needed to address this question.

Genomic locations of positive and negative supercoiling

476   When we arranged PS and NS sites along the *E. coli* chromosome, we found that they share

similar patterns. Both PS and NS sites are enriched in *Ori* and *ter* regions, but less in between (**Fig. 1C,**

1D). As the *Ori* macrodomain contains most of the active genes, and both *Ori* and *Teri* are involved in

DNA replication (8, 28), the dense location of PS and NS sites in these two regions is consistent with the

expectation that transcription and replication are major supercoil generators. Between the two,

transcription appears to have a more significant role in generating supercoils, as over 95% of GapR and

TopoI peaks are associated with TUs. Of note, there are more GapR peaks (609) compared to TopoI

peaks (446) (see **Supplementary Table 1** for statistics), which could be due to different capturing

efficiency of the corresponding antibody in the ChIP-Seq experiments.

Short-distance and long-distance compaction

486   We show that the spatial distance separation between two bins containing PS or NS sites is

smaller than those without these sites, with the largest effects on compaction occurring for loci 0.5-1 Mb

apart in a linear chromosome (**Fig. 1I-J, 2C-D**). We hypothesize that the fact that supercoiling-dependent

compaction peaks at 0.5-1 Mb is a result of the chromosome conformation and is dictated by the

macrodomain organization of the *E. coli* genome. Short-distance interactions such as TADs occur on only

~100 Kb scales (8), which are too short to explain our observations. In contrast, the observed

supercoiling-dependent compaction matches the size of macrodomains, which are estimated to be ~1 Mb

(8). While the actual peak is slightly different for each experiment, they all fall in the 0.5-1 Mb genomic

distance range. Such a difference may be due to experimental variations since it has an association with

the number of valid paired reads from the Hi-C/5C experiment. As such, we hypothesize that positive and

negative supercoiling contributes to organizing the chromosome by enhancing the interaction within a

macrodomain.

Physical models of the supercoiling-organized *E. coli* chromosome

499   What would be the physical model of the *E. coli* chromosome architecture that explains our

observations? There are two challenges to forming a unified model. First, most models of chromosome

architecture are based on the underlying assumption that physical distance and contact frequency are

inversely correlated (7, 38, 39). However, this relationship is not necessarily valid for all loci and, in

experiments that test this relationship, the variation between contact frequency and physical distance can

be large (7, 39, 40, 41). Second, Hi-C captures population-averaged contact, meaning that contact could

result from distant loci being physically closer but static, or from dynamic (and potentially rare), frequent

collisions between loci. Therefore, we propose two models, one static and one dynamic. In the static

model (**Fig. 5A**), both positive and negative supercoiling lead to the compaction of the chromosome but

positive supercoiling has a stronger effect than negative supercoiling. As such, chromosomal sites

containing positive supercoils may form the core of the condensed chromosome, while sites containing

negative supercoils are on the periphery of the core, allowing genomic process such as replication and

transcription to occur. In this sense, positively supercoiled and negatively supercoiled chromosomal sites

may be analogous to heterochromatin and euchromatin in eukaryotes. In the dynamic model (**Fig. 5B**), the

chromosome may be an amorphous network of DNA that is being dynamically reshaped due to the

transitory, nonspecific interactions with nucleoid-associated proteins such as HU (42). Sites containing

positive supercoils may be more mobile due to the lack of NAP binding and hence have a higher chance

to interact with each other over long linear distances in the nucleoid (**Fig. 5B**, blurry lines). Sites

containing negative supercoils and no supercoils, however, are relatively immobile due to the stable

association with NAPS, some of them having a higher binding tendency to negatively supercoiled DNA

(14), and the transcription/translation machineries, and hence have less chances to interact with each other

(**Fig. 5B**, sharp lines and small blobs). Both models should give rise to similar Hi-C contact maps. Further

investigation to visualize the spatial localization and dynamics of positively and negatively supercoiled

sites in live *E. coli* cells is required to differentiate between these models.

DATA AVAILABILITY

The sequencing data from this study have been deposited at the Gene Expression Omnibus (GEO):

GSE214856. Scripts used for data processing, analysis, trained machine learning models are available

from the Xiao Laboratory GitHub repository (XiaoLab JHU: https://github.com/XiaoLabJHU).

SUPPLEMENTARY DATA

Supplementary Data are available at NAR online.

ACKNOWLEDGEMENTS

The authors thank all members of the Xiao, Ji laboratories for their valuable discussions and advice on

this project. All computations were performed on the Joint High Performance Computing Exchange

(JHPCE), organized by the Department of Biostatistics at the Johns Hopkins Bloomberg School of Public

Health, and the Maryland Advanced Research Computing Center (MARCC) Rockfish cluster. Z.F. wants

to thank Jiafang Song for creating the illustration cartoons and Dr. Hongkai Ji for offering statistical

insights.

FUNDING

This work was supported by the National Institutes of General Medical Science [1R35GM136436 to J.X.

and R00GM134153 to M.S.G.]; and the National Science Foundation [MCB1817551 to J.X.]. Funding

for open access charge: National Institutes of Health.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

548   References

1. Cavalli,G. and Misteli,T. (2013) Functional implications of genome topology. *Nat Struct Mol Biol,* **20,**
290-299.

2. Ali Azam,T., Iwata,A., Nishimura,A., Ueda,S. and Ishihama,A. (1999) Growth phase-dependent
variation in protein composition of the escherichia coli nucleoid. *J. Bacteriol.,* **181,** 6361-6370.

3. White,J.H. and Bauer,W.R. (1986) Calculation of the twist and the writhe for representative models of
DNA. *J. Mol. Biol.,* **189,** 329-341.

4. Lioy,V.S., Cournac,A., Marbouty,M., Duigou,S., Mozziconacci,J., Espéli,O., Boccard,F. and Koszul,R.
(2018) Multiscale structuring of the E. coli chromosome by nucleoid-associated and condensin proteins.
Cell, **172,** 771-783.e18.

5. Cockram,C., Thierry,A. and Koszul,R. (2021) Generation of gene-level resolution chromosome contact
maps in bacteria and archaea. *STAR Protoc,* **2,** 100512.

6. van Berkum,N.L., Lieberman-Aiden,E., Williams,L., Imakaev,M., Gnirke,A., Mirny,L.A., Dekker,J.
and Lander,E.S. (2010) Hi-C: A method to study the three-dimensional architecture of genomes. *J. Vis.*
Exp., **(39):1869. doi,** 10.3791/1869.

7. Le,T.B.K., Imakaev,M.V., Mirny,L.A. and Laub,M.T. (2013) High-resolution mapping of the spatial
organization of a bacterial chromosome. *Science,* **342,** 731-734.

8. Valens,M., Penaud,S., Rossignol,M., Cornet,F. and Boccard,F. (2004) Macrodomain organization of
the escherichia coli chromosome. *Embo J.,* **23,** 4330-4341.

9. Wu,H.Y., Shyy,S.H., Wang,J.C. and Liu,L.F. (1988) Transcription generates positively and negatively
supercoiled domains in the template. *Cell,* **53,** 433-440.

10. Liu,L.F. and Wang,J.C. (1987) Supercoiling of the DNA template during transcription. *Proc. Natl.*
Acad. Sci. U. S. A., **84,** 7024-7027.

11. Lal,A., Dhar,A., Trostel,A., Kouzine,F., Seshasayee,A.S.N. and Adhya,S. (2016) Genome scale
patterns of supercoiling in a bacterial chromosome. *Nat Commun,* **7,** 11055.

12. Bates,A.D. and Maxwell,A. (2005) DNA Topology. OUP Oxford, .

13. Guo,M.S., Kawamura,R., Littlehale,M.L., Marko,J.F. and Laub,M.T. (2021) High-resolution,
genome-wide mapping of positive supercoiling in chromosomes. *Elife,* **10,** 10.7554/eLife.67236.

14. Sutormin,D., Galivondzhyan,A., Musharova,O., Travin,D., Rusanova,A., Obraztsova,K., Borukhov,S.
and Severinov,K. (2022) Interaction between transcribing RNA polymerase and topoisomerase I prevents
R-loop formation in E. coli. *Nat. Commun.,* **13,** 4524-5.

15. Guo,M.S., Haakonsen,D.L., Zeng,W., Schumacher,M.A. and Laub,M.T. (2018) A bacterial
chromosome structuring protein binds overtwisted DNA to stimulate type II topoisomerases and enable
DNA replication. *Cell,* **175,** 583-597.e23.

16. Tan,K., Zhou,Q., Cheng,B., Zhang,Z., Joachimiak,A. and Tse-Dinh,Y. (2015) Structural basis for
suppression of hypernegative DNA supercoiling by E. coli topoisomerase I. *Nucleic Acids Res.,* **43,**
11031-11046.

17. Servant,N., Varoquaux,N., Lajoie,B.R., Viara,E., Chen,C., Vert,J., Heard,E., Dekker,J. and Barillot,E.
(2015) HiC-pro: An optimized and flexible pipeline for hi-C data processing. *Genome Biol.,* **16,** 259.

18. Imakaev,M., Fudenberg,G., McCord,R.P., Naumova,N., Goloborodko,A., Lajoie,B.R., Dekker,J. and
Mirny,L.A. (2012) Iterative correction of hi-C data reveals hallmarks of chromosome organization. *Nat
Methods,* **9,** 999-1003.

19. Gu,Z., Eils,R. and Schlesner,M. (2016) Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics,* **32,** 2847-2849.

20. Andrews,S. (2010) FastQC: A quality control tool for high throughput sequence data.

21. Bolger,A.M., Lohse,M. and Usadel,B. (2014) Trimmomatic: A flexible trimmer for illumina sequence
data. *Bioinformatics,* **30,** 2114-2120.

22. Langmead,B. and Salzberg,S.L. (2012) Fast gapped-read alignment with bowtie 2. *Nat Methods,* **9,**
357-359.

23. Li,H., Handsaker,B., Wysoker,A., Fennell,T., Ruan,J., Homer,N., Marth,G., Abecasis,G., Durbin,R.
and 1000 Genome Project Data Processing Subgroup. (2009) The sequence alignment/map format and
SAMtools. *Bioinformatics,* **25,** 2078-2079.

24. Zhang,Y., Liu,T., Meyer,C.A., Eeckhoute,J., Johnson,D.S., Bernstein,B.E., Nusbaum,C., Myers,R.M.,
Brown,M., Li,W., et al. (2008) Model-based analysis of ChIP-seq (MACS). *Genome Biol.,* **9,** R137.

25. Kuhn,M. (2008) Building predictive models in R using the caret package. *J. Stat. Soft.,* **28,** 1.

26. Davis,J. and Goadrich,M. (2006) The relationship between precision-recall and ROC curves. 233–
240.

27. Wang,S., Su,J., Beliveau,B.J., Bintu,B., Moffitt,J.R., Wu,C. and Zhuang,X. (2016) Spatial
organization of chromatin domains and compartments in single chromosomes. *Science,* **353,** 598-602.

28. Verma,S.C., Qian,Z. and Adhya,S.L. (2019) Architecture of the escherichia coli nucleoid. *PLoS
Genet.,* **15,** e1008456.

29. Racko,D., Benedetti,F., Dorier,J. and Stasiak,A. (2018) Are TADs supercoiled? *Nucleic Acids Res.,*
47, 521-532.

30. Ma,J. and Wang,M.D. (2016) DNA supercoiling during transcription. *Biophys. Rev.,* **8,** 75-87.

31. Herzel,L., Stanley,J.A., Yao,C. and Li,G. (2022) Ubiquitous mRNA decay fragments in E. coli
redefine the functional transcriptome. *Nucleic Acids Res,* **50,** 5029-5046.

32. Johnstone,C.P. and Galloway,K.E. (2022) Supercoiling-mediated feedback rapidly couples and tunes
transcription. *Cell Reports,* **41,** .

33. Geng,Y., Bohrer,C.H., Yehya,N., Hendrix,H., Shachaf,L., Liu,J., Xiao,J. and Roberts,E. (2022) A
spatially resolved stochastic model reveals the role of supercoiling in transcription regulation. *PLoS
Comput. Biol.,* **18,** e1009788.

34. Alberts,B. (2003) DNA replication and recombination. *Nature,* **421,** 431-435.

35. Rouvière-Yaniv,J., Yaniv,M. and Germond,J.E. (1979) E. coli DNA binding protein HU forms
nucleosomelike structure with circular double-stranded DNA. *Cell,* **17,** 265-274.

36. Broyles,S.S. and Pettijohn,D.E. (1986) Interaction of the escherichia coli HU protein with DNA.
evidence for formation of nucleosome-like structures with altered DNA helical pitch. *J Mol Biol,* **187,** 47-
60.

37. Schneider,R., Travers,A. and Muskhelishvili,G. (1997) FIS modulates growth phase-dependent
topological transitions of DNA in escherichia coli. *Mol Microbiol,* **26,** 519-530.

38. Conin,B., Billault-Chaumartin,I., El Sayyed,H., Quenech'Du,N., Cockram,C., Koszul,R. and
Espéli,O. (2022) Extended sister-chromosome catenation leads to massive reorganization of the E. coli
genome. *Nucleic Acids Res.,* **50,** 2635-2650.

39. Imakaev,M.V., Fudenberg,G. and Mirny,L.A. (2015) Modeling chromosomes: Beyond pretty
pictures. *FEBS Lett.,* **589,** 3031-3036.

40. Fudenberg,G. and Imakaev,M. (2017) FISH-ing for captured contacts: Towards reconciling FISH and
3C. *Nat. Methods,* **14,** 673-678.

41. Williamson,I., Berlivet,S., Eskeland,R., Boyle,S., Illingworth,R.S., Paquette,D., Dostie,J. and
Bickmore,W.A. (2014) Spatial genome organization: Contrasting views from chromosome conformation
capture and fluorescence in situ hybridization. *Genes Dev.,* **28,** 2778-2791.

42. Bettridge,K., Verma,S., Weng,X., Adhya,S. and Xiao,J. (2021) Single-molecule tracking reveals that
the nucleoid-associated protein HU plays a dual role in maintaining proper nucleoid volume through
differential interactions with chromosomal DNA. *Mol. Microbiol.,* **115,** 12-27.

FIGURES AND TABLES

Figure 1.

Figure 1. *E. coli* **chromosomal loci containing positive supercoiling sites exhibit higher-than-average spatial contacts in Hi-C maps.**

(**A**) A cartoon illustration of the study design. We correlate the spatial contact frequency of any two *E. coli* chromosomal loci (green and cyan dots on the wiggle lines) as depicted in the Hi-C map (red square, bottom) with the supercoiling states of the two loci as identified by GapR (indicating positively supercoiled DNA) or Topo I (indicating negatively supercoiled DNA) chromatin-immunoprecipitation sequencing data (ChIP-seq, right).

(**B**) A cartoon illustration of GapR/Topo I ChIP-seq data bin assignment. The 4.6 Mb *E. coli* genome is divided into 5 Kbp or 0.5 Kbp bins, with each bin assigned as PS (positive supercoiling site marked by GapR-peak), aPS (absent of a positive supercoiling site, no GapR peak), NS (negative supercoiling site marked by Topo I-peak), or aNS (absent of a negative supercoiling site, no TopoI peak), based on GapR-Seq and TopoI-seq data. A bin can contain either one, both, or none.

(**C-D**) Comparison of genomic locations of GapR and TopoI peaks with respect to their spatial contacts with other genomic locations in the Hi-C contact heatmaps of *E. coli* MG 1655. Normalized (log2) contact map are plotted as symmetric halves from the 5C data set (LB, 37 ℃, 5 Kbp resolution) generated by Lioy et. al (left) and Hi-C data set (LB, 37℃, 5 Kb resolution) generated in this study (right), with a 45% counterclockwise rotation. Vertical bars at the bottom of the contact maps mark the ChIP-seq peaks for GapR (black), GapR[1-76] (green), and Topo I (blue). The two horizontal lines in purple (10Kb, 1Mb) represents the window used to compute the contact sum (in **E-F**) at each genomic locus in the contat map (5 Kb bin). The two reference rules provide the *E. coli* genomic coordinates (bottom, increment = 500 Kb) and locations of *Ori*C and *ter*C.

(**E-F**) Normalized (log2) sum of the total contact counts in the 100 Kb and 1 Mb window of the contact map (C or D) at each locus (black dots) is plotted along the *E. coli* genomic coordinates. Blue curve shows the moving average (window size = 15) of the normalized contact sum and peaks near ~3-4 Mb where *Ori*C resides.

(**G-H**) Inversed average contact frequencies of two genomic loci (*y*-axis) at each linear genomic separation distance (*x*-axis) with simple moving averages (colored curves, window size = 15) for loci pairs both containing positive supercoiling sites (PS-PS, blue), not containing positive supercoiling sites (aPS-aPS, green), and all loci pairs (all-all, red) show that PS-PS loci pairs have on average lowest inversed average contact frequencies (highest spatial contact) at all linear genomic distances than aPS-aPS or all-all loci pairs.

(**I-J**) Numeric difference in the inversed average contact frequencies (y-axis) between PS-PS loci and aPS-aPS loci at each genomic separation distance (x-axis) reaches maximum in the range of 500- 1000 Kbp.

Figure 2.

Figure 2. *E. coli* **chromosomal loci containing negative supercoiling sites exhibit higher-than-average but lower spatial contacts in Hi-C maps than loci containing positive supercoiling.**

(A-B) Inversed average contact frequencies of two genomic loci (*y*-axis) at each linear genomic separation distance (*x*-axis) with simple moving averages (colored curves, window size = 15) for loci pairs both containing negative supercoiling sites (NS-NS, blue), not containing positive supercoiling sites (aNS-aNS, green), and all loci pairs (all-all, red) show that NS-NS loci pairs have on average lowest inversed average contact frequencies (highest spatial contact) at all linear genomic distances than aNS-aNS or all-all loci pairs.

(C-D) Numeric difference in the inversed average contact frequencies (y-axis) between NS-NS loci and aNS-aNS loci at each genomic separation distance (x-axis) reaches maximum in the range of 500- 1000 Kbp Kbp.

(E-F) Inversed average contact frequencies of two genomic loci (*y*-axis) at each linear genomic separation distance (*x*-axis) with simple moving averages (colored curves, window size = 15) for loci pairs containing positive supercoiling site only (PS only -PS only, blue), negative supercoiling site only (NS only – NS only, green), and none (aa – aa, red) show that PS only-PS only loci pairs have on average lowest inversed average contact frequencies

(G-H) Numeric difference in the inversed average contact frequencies (y-axis) between PS only-PS only loci and aa-aa loci at each genomic separation distance (x-axis) reaches maximum in the range of 500- 1000 Kbp.

Figure 3.

Figure 3. Machine learning models can predicate **positive but not negative supercoiling sites accurately from Hi-C contact map.**

(**A**) A cartoon illustration of the machine learning model design. The input matrix is a rearrangement of the conventional contact matrix[7] (left), in which each input feature (row) is the contact frequency of a genomic locus (one 5 Kbp bin, brown) with flanking chromosome on both sides (-464 to + 463 bins, blue). The responses, or whether this 5 Kb bin contains any Gap (green) or TopoI (purple) peaks, are indicated by the check and error marks

(middle). A set of randomly chosen loci (25% out of 928 bins) is preserved as a fixed testing set for all random forests models (right).

(**B-C**) ROC and PR curves of 10 independent experiments (light-shaded curves) with 25% (red), 50% (green), and 75% (blue) training data predicted positive supercoiling (marked by GapR peaks) locations robustly with similar AUROC (0.74) and AUPRC (0.68) values, while the baseline models (black), trained with randomized peak labels, are at ~ 0.43 and 0.44 respectively.

(**D**) Average feature importance (over 10 independent experiments) measured in mean decrease accuracy (colored dots, with simple moving averages of window size = 15 as solid curves) from experiments with 25% (red), 50% (green), and 75% (blue) training data peaked at ~ 500 Kb genomic separation distance, indicating that these features contribute the most to the effective classification of GapR peak-containing bins.

(**E-F**) ROC and PR curves of 10 independent experiments (light-shaded curves) with 25% (red), 50% (green), and 75% (blue) training data did not predicate negative supercoiling (marked by Topo I peaks) locations robustly. The AUROC (0.54-0.58) and AUPRC (0.41-0.44) values are not significantly different from those of the baseline models (black curves), indicating that the models cannot effectively recover negative supercoiling spatial patterns from Hi-C contact maps.

(**G**) Average feature importance (over 10 independent experiments) measured in mean decrease accuracy (colored dots, with simple moving averages of window size = 15 as solid curves) from experiments with 25% (red), 50% (green), and 75% (blue) training data showed uniform patterns across different genomic separation distances

(**H**) The average contact frequencies (y-axis, inversed contact counts representing the apparent spatial separation between two loci) at each signed genomic separation distance (x-axis) are plotted (colored dots) with simple moving averages (colored curves, window size = 15). Different colors indicate the interaction between loci pairs with at least one containing positive supercoiling site (PS-all, blue) and that with at least one containing no positive supercoiling site (aPS-all, green). The average contact frequencies at each genomic separation are plotted in red as a reference.

(**H**) The inverse average contact frequencies among PS-all (blue), aPS-all (green), and all-all (red) loci pairs in the range of $\pm$ 500 – 1000 Kbp are significantly different from each other, explaining the effective prediction of positive supercoiling sites on *E. coli* genome using Hi-C contacts as the input.

(**I**) The inverse average contact frequencies among NS-all (blue), aNS-all (green), and all-all (red) loci pairs in all ranges are not significantly different from each other, explaining the ineffective prediction of negative supercooling sites on *E. coli* genome using Hi-C contacts as the input.

Figure 4.

Figure 4. Transcription is a strong driving force for supercoiling generation.
(**A**) 2D-histrogram showing the distribution of supercoiling sites (hexagon bins colored by density) with respect to the distance (bp, y-axis) to their nearest upstream transcription end site (TES) and the normalized (log2) transcription activity (x-axis, RPK) of that transcription unit (TU). Orange vertical line indicates the 75th percentile of all TU activities (x-intercept = 29.34). Orange horizontal line indicates 1 Kb away from the nearest TES (y-intercept = 1000 bp). Positive supercoiling sites located in the purple block are 1 Kb away from the nearest TES and

the associated TUs have transcription activities below the 75th percentile, suggesting that those sites are not associated with transcription.

(**B**) The negative supercoiling counterpart of (**A**).

(**C-D**) Control experiments repeating the analysis in (**A and B**) by randomly generating a set of genomic locations matching the number of PS and NS sites. More randomized sites reside within the purple blocks (> 1 Kb away from TUs of low transcription activities) than those in (**A-B**).

(**E**) PS sites (colored dots) greater than 1 Kb away from their nearest TES (x-axis) are plotted to visualize their location along the *E. coli* chromosome (y-axis). Position of *Ori*C and *Ter*C are indicated by two horizontal lines. Positive supercoiling sites are colored in red if they associate with a highly transcribed TU (top 25% percentile) and in green otherwise.

(**F**) The NS counterpart of (**E**).

(**G-H**) Control experiments repeating the analysis in (**E**) using the randomly generated sets of genomic sites.

SUPPLEMENTAL FIGURES AND TABLES

Figure S1.

Figure S1.

(A) Replicates of the corresponding Hi-C and 5C experiments illustrated in (Fig. 1C-D, E-F) verified the reproducibility of the patterns across different experimental conditions and resolutions as described in the article.

(B) Replicates of the corresponding Hi-C and 5C experiments illustrated in (Fig. 1G-H, I-J) verified the reproducibility of the patterns described in the article.

(C) GapR[1-76] counterpart of the analysis illustrated in (Fig. 1G-H, I-J).

(D) Density plots for lengths (log2 normalized, x-axis) of identified reproducible GapR peaks (salmon, N=609), GapR[1-76] peaks (green, N=482), and TopoI peaks (blue, N=446).

Table S1.

1A. The *E. coli* genome is divided into 928 5Kb bins and 4642 0.5 Kb bins. We identified 609 positive supercoiling sites (marked by reproducible GapR peaks) and 446 negative supercoiling sites (marked by reproducible TopoI peaks) (see methods, ChIP-seq analysis). The *OriC* half is defined as the half of the *E. coli* genome centering at the 5 Kb or 0.5 Kb bin which contains *OriC* (coordinate = 3,925,744, EcoCyc). The *ter* half is defined by the set of bins complement to the *OriC* half.

	GapR	Topo
Total peak number	609	446
5Kb bins (928)	454	352
0.5Kb bins (9284)	3672	1890
5Kb bins in *ter* half (464)	147	150
5Kb bins in *OriC* half (464)	307	201
0.5Kb bins in *ter* half (4642)	714	716
0.5Kb bins in *OriC* half (4642)	2958	1174

1B: Statistics of three sets of identified, reproducible peak widths

	Min	**1st Qu.**	**Median**	**Mean**	**3rd Qu.**	**Max.**
GapR	278	654	1193	2532	3112	35457
TopoI	330	558.2	946.5	1647.2	1942	15616
GapR[1-76]	257	583.8	820	1201.8	1302.5	14331

Figure S2.

Fig S2.
(A) Replicates of the corresponding Hi-C and 5C experiments illustrated in (**Fig. 2A-B, C-D**) verified the reproducibility of the patterns described in the article.
(B) Replicates of the corresponding Hi-C and 5C experiments illustrated in (**Fig. 2F-G, H-I**) verified the reproducibility of the patterns described in the article

Figure S3.

Fig S3.
(A) Average contact frequencies (y-axis, inversed contact counts representing the apparent spatial separation between two loci) at each signed genomic separation distances (x-axis) are plotted (colored dots) with moving averages (colored curves, window size = 15). Different colors indicate the interaction between all loci pairs (all-all, red), when at least one locus does not contain positive supercoiling site (aPS-all, yellow), when both loci do not contain positive supercoiling site (aPS-aPS, green), when at least one locus contains positive supercoiling site (PS-all, blue), or when both loci contain positive supercoiling site (PS-PS, pink).
(B) The negative supercoiling counterpart of (A).

Figure S4.

Fig S4.
(A) Density plots of the log2 normalized transcription unit length (left) and activity (right 2, measured in Reads Per Kilobase).

(B) Density plots of the distance from PS v.s. random-PS (609 sites) to their nearest TES (top left) and from NS v.s. random-NS (364 sites) to their nearest TSS (top right). The cumulative distribution functions (bottom left) of the four distributions in the above two figures and with log2 normalization (bottom right).

(C) Four sets of peaks are plotted to visualize their location along the *E. coli* genome (y-axis) and distance to their nearest TU (x-axis). Two horizontal lines in blue indicate the position of *ori*C and *ter*. Positive supercoiling sites are colored in red if they associate with a highly transcribed TU and in green otherwise.