Novel *emm*4 lineage associated with an upsurge in invasive group A streptococcal disease in the Netherlands, 2022

Boas C.L. van der Putten[1, 5]*, Wendy C. Bril-Keijzers[1,5], Lidewij W. Rümke[2], Stefan M.T. Vestjens[2], Linda A.M. Koster[1], Marloes Willemsen[1#], Bart J.M. Vlaminckx[3], Brechje de Gier[4], Nina M. van Sorge[1,5&]

[1]Department of Medical Microbiology and Infection Prevention, Amsterdam UMCs location University of Amsterdam, Amsterdam, the Netherlands

[2]Medical Microbiology, UMC Utrecht, Utrecht, the Netherlands

[3]Department of Medical Microbiology and Immunology, St Antonius Hospital, Nieuwegein, the Netherlands

[4]National Institute for Public Health and the Environment, Bilthoven, the Netherlands

[5]Netherlands Reference Laboratory for Bacterial Meningitis, Amsterdam UMCs location AMC, Amsterdam, the Netherlands

*Current affiliation: National Institute for Public Health and the Environment, Bilthoven, the Netherlands.

#Current affiliation: Hartwig Medical Foundation, Amsterdam, the Netherlands.

&Corresponding author full contact information: Department of Medical Microbiology and Infection Prevention, Amsterdam UMCs location University of Amsterdam, Meibergdreef 9, IWO building IA3.159, 1105 AZ Amsterdam, the Netherlands; email address: n.m.vansorge@amsterdamumc.nl; phone number +31205664862

Running title: Emerging *S. pyogenes emm*4 lineage in the Netherlands

Keywords: *Streptococcus pyogenes*, public health surveillance, outbreak, *emm* typing, whole-genome sequencing

Abstract

Invasive group A streptococcal (iGAS) disease cases increased in the first half year of 2022 in the Netherlands with a remarkably high proportion of *emm4* isolates. Whole-genome sequence analysis of 66 *emm*4 isolates from 2009-May 2022 identified a novel GAS lineage (M4$_{NL22}$), which accounted for 85% of *emm*4 iGAS cases in 2022. Surprisingly, we detected few isolates of the *emm*4 hypervirulent clone, which has replaced nearly all other *emm*4 lineages in the USA and the UK. M4$_{NL22}$ displayed genetic differences compared to other *emm*4 although these are currently of unclear biological significance. In publicly-available data, we identified a single Norwegian isolate belonging to M4$_{NL22}$ that was sampled after the isolates from this study, possibly suggesting export of M4$_{NL22}$ to Norway. In conclusion, our study identified a novel lineage associated with an increase of iGAS disease in early 2022 in the Netherlands

Impact statement

Group A *Streptococcus* (mostly comprised by *Streptococcus pyogenes*) has a long history of causing disease, including epidemic outbreaks, in humans. Over the past decades, researchers have described various changes in the *S. pyogenes* population that changed its virulence potential. In this study, we describe a shift in the *S. pyogenes* population within type *emm*4 in the Netherlands in early 2022. A novel lineage, named M4$_{NL22}$, became the predominant *emm*4 lineage during this period. This was surprising, as in other countries such as the USA and the UK, another recently emerged hypervirulent *emm*4 lineage became predominant. Our work informed public health officials on the possible causes of increased

S. pyogenes cases in the Netherlands and underlines the added value of continued

bacteriological surveillance.

Data summary

Code is freely available from https://github.com/boasvdp/GAS_M4. All sequencing data is

freely available from BioProject PRJEB58654.

Introduction

Streptococcus pyogenes can cause a wide variety of infections in humans, including non-

invasive manifestations such as scarlet fever, pharyngitis and impetigo but also invasive

manifestations such as fasciitis necroticans, streptococcal toxic shock syndrome, and

puerperal fever. These three invasive group A streptococcal (iGAS) infections are notifiable

by law in the Netherlands. An upsurge in both notifiable and non-notifiable iGAS infections

was observed in the Netherlands compared to pre-COVID-19 levels in spring 2022, shortly

after restrictions were lifted. Alarmingly, this increase disproportionally affected children

under 5 years of age[1,2]. Changes or shifts in dominance of *emm*-types, which can be assessed

through bacteriological surveillance, may indicate changes in virulence or transmission

within the *S. pyogenes* population.

   Bacteriological surveillance for *S. pyogenes* was initiated as part of a research project

in 2019. *S. pyogenes* isolates cultured from normally-sterile compartments have been

voluntarily submitted by medical microbiology laboratories to the Netherlands Reference

Laboratory for Bacterial Meningitis (NRLBM) for *emm*-typing (see appendix for details).

Between 1 January to 13 May 2022, 25 isolates were received from children under the age of

5 years of which *emm*4 and *emm*12 were the most dominant *emm* types (28% each),

followed by *emm*1.0 (16%). Among isolates from all ages (n=134), *emm*4 ranked third (13%). The dominance of *emm*4 among iGAS cases was unexpected, since data from a retrospective study (2009-2019) indicated that this *emm*-type was associated with carriage rather than invasive disease in the Netherlands (L.W. Rümke, unpublished data).

Recently, a hypervirulent *emm*4 clone (henceforth referred to as the "hypervirulent clone") was described, which has replaced nearly all other *emm*4 *S. pyogenes* strains in the USA and UK[3]. We hypothesized that this hypervirulent *emm*4 clone was responsible for the increase of *emm*4 iGAS cases in the Netherlands.

Methods

Sample selection

We included all 26 *emm*4 *S. pyogenes* isolates from iGAS patients that were received by the NRLBM between January 1st and May 14th, 2022 (see Table 1 and Appendix for details). Additionally, we included 40 *emm*4 *S. pyogenes* isolates from the 2009-2019 period (pre-COVID samples), which were from invasive disease or from asymptomatic carriage (Table 1). All 66 isolates were subjected to whole-genome sequencing (WGS).

Whole-genome sequencing

Short read sequencing

DNA extraction was performed using a magnetic bead-based protocol on an automated Maxwell machine (Promega). Whole-genome sequencing of 62 (of total 66) isolates was conducted by the Core Facility Genomics of the Amsterdam UMC. Library preparation was performed using the Kapa HTP Library Preparation kit and libraries were sequenced on an Illumina Hiseq4000 using a PE150 kit, all according manufacturer's instructions. The

remaining four isolates were sequenced by MicrobesNG, United Kingdom

(https://microbesng.com/). Isolates were suspended in DNA Shield buffer (Zymo Research,

USA) and shipped to MicrobesNG. The DNA extraction, library preparation, sequencing and

bioinformatics QC protocols of MicrobesNG are available online (v20210419,

https://microbesng.com/documents/24/MicrobesNG_Sequencing_Service_Methods_v2021

0419.pdf).

Long read sequencing

Isolate 2220758 was selected for long read sequencing using Oxford Nanopore Technologies

(ONT). Reads from three sequencing experiments were combined due to low yields of

individual experiments. DNA extractions for the first two sequencing experiments were

performed according to a method described previously[4]. DNA extraction for the third

sequencing experiment was performed using the Maxwell RSC Cultured Cells DNA Kit (Cat.#

AS1620) according to manufacturer's instructions. Library preparation and MinION

sequencing methodologies for all three experiments were described before[4].

WGS analysis

Short read analysis

Illumina reads were processed with an adapted version of the NRLBM genomic pipeline

(v0.3, https://github.com/NRLBM/assembly/releases/tag/v0.3). All analyses were run with

default settings unless otherwise noted. Sequence reads were trimmed and filtered using

Trimmomatic v0.36[5]. Sequence read qualities were assessed before and after Trimmomatic

using FastQC v0.11.9[6]. Contamination was assessed using Kraken2 v2.1.2[7] and the

MiniKraken v1 database. Filtered sequence reads were assembled using the Shovill v1.1.0

wrapper (https://github.com/tseemann/shovill) for SPAdes v3.15.4[8]. Depth of coverage was

assessed by mapping filtered reads on de novo assemblies using minimap2 v2.17-r941[9], samtools v1.9[10] and bedtools v2.29.0[11]. Assembly quality was assessed using Quast v4.6.3[12]. Multi-locus sequence types were assigned using the PubMLST API[13]. Virulence genes were detected using ABRicate v1.0.1 (https://github.com/tseemann/abricate) and VFDB[14]. *emm*-types were confirmed using emmtyper v0.2.0 (https://github.com/MDU-PHL/emmtyper).

A maximum-likelihood (ML) phylogeny was constructed by mapping filtered sequence reads on the Duke strain reference genome (GenBank accession number CP031770) using Snippy v4.6.0 (https://github.com/tseemann/snippy). The previously described hypervirulent clone was identified according to 36 SNPs typical for this lineage[3]. A genome alignment was created from Snippy mappings and recombination was identified using ClonalFrameML v1.12[15]. Recombinatory regions were masked using maskrc-svg v0.5 (https://github.com/kwongj/maskrc-svg) and a ML phylogeny was inferred from the recombination-free alignment using IQ-TREE v2.0.3[16], with the substitution model HKY+F.

Draft assemblies were annotated using Bakta v1.4.1[17] with database v3.1. The pan-genome of this dataset was defined using Panaroo v1.2.10[18] in strict mode and lineage-associated genetic elements were identified using PySEER v1.3.9[19]. A Bonferroni correction was applied to define statistically significant associations.

Long read analysis

Leading and trailing 80 bp of all ONT reads were trimmed using NanoFilt v2.8.0[20]. Trimmed reads were subsequently filtered using filtlong v0.2.1 (https://github.com/rrwick/Filtlong), retaining reads of at least 5,000 bp long and the best 90% of reads matched by nucleotide identity based on Illumina reads. Illumina and ONT reads were assembled using Unicycler v0.5.0[21] resulting in a completely resolved genome. The complete genome of 2220758 was

annotated as described above. Additionally, the PHASTER webserver[22] was queried to annotate prophages and the Artemis Comparison Tool v18.2.0[23] was used to identify large-scale genomic rearrangements.

Public data

To relate the Dutch data to other recent surveillance data of *S. pyogenes*, we downloaded reads from BioProjects PRJNA395240 and PRJEB42599 comprising *S. pyogenes* data from USA and Norwegian surveillance programmes, respectively. Downloaded reads were processed together with Dutch data using the standard pipeline.

Results

Patient characteristics

For 25 of the 26 iGAS cases from 2022, the median patient age was 34 years (IQR: 5-43 years; Table 1) and 15 (60%) were male. For one patient no information was available on age or sex. Site of isolation is summarized in Table 1. Of the pre-COVID isolates, 26 out of 40 (65%) were from 0-5-year olds, while the remaining 14 isolates were from adults.

Table 1. *S. pyogenes* isolate characteristics from 2009-2019 (pre-COVID) and 2022 periods.

	2009-2019 (pre-COVID)		2022
	Carriage	Invasive	Invasive
Number of isolates	29	11	26
Median year of isolation (IQR)	2012 (2009-2016)	2013 (2013-2015)	2022
Cultured from adult*, no. (%)	6 (21%)	8 (72%)	17 (65%)

Site of isolation			
Blood, no. (%)	0 (0%)	11 (100%)	15 (58%)
Nose, no. (%)	23 (79%)	0 (0%)	0 (0%)
Throat, no. (%)	6 (21%)	0 (0%)	0 (0%)
Female genital tract, no. (%)	0 (0%)	0 (0%)	5 (19%)
Other#, no. (%)	0 (0%)	0 (0%)	6 (23%)
M4$_{NL22}$ lineage, no. (%)	1 (3%)	1 (3%)	22 (85%)
Hypervirulent clone, no. (%)	3 (10%)	1 (9%)	2 (8%)

*For one group only age categories are available instead of exact age
#Other sources include wound, abscess, general tissue and a dialysis machine

Bacterial characteristics: new *emm*4 clone

The hypervirulent clone represented less than 10% among recent and older *S. pyogenes* isolates (Table 1), in contrast to the strong emergence of this clone in the USA and the UK. However, 22 out-of-26 (84.6%) of 2022 iGAS isolates clustered together, representing a new *emm*4 lineage M4$_{NL22}$ (Figure). Fifteen out-of-22 (68%) M4$_{NL22}$ isolates showed very limited genetic variation with a median of 6 SNPs between them (range 0-16). This suggests a very recent emergence and potentially a high transmission rate. In contrast, only two out-of-40 (5%) isolates from 2009-2019 were identified as M4$_{NL22}$ strains (isolated in 2017 and 2018). Analysis of public surveillance databases, identified one isolate (NCBI BioSample SAMEA111521236) from 2022 from Norway (Norwegian Public Institute of Health surveillance, NCBI BioProject PRJEB42599) that clustered in the clonal part of the M4$_{NL22}$ lineage and differed 9 SNPs from the closest Dutch isolate. No M4$_{NL22}$ isolates were identified from USA surveillance data from 2021-2022 (CDC Active Bacterial Core surveillance, NCBI BioProject PRJNA395240).

Figure. Phylogeny of 66 Dutch *emm*4 isolates and 14 reference genomes. Reference

genomes originate from Debroy *et al*. (2021) (*2*) and are shown without metadata. The

hypervirulent clone lineage is colored blue and the M4$_{NL22}$ lineage is colored red.

Characteristics of M4$_{NL22}$ lineage strains

The M4$_{NL22}$ lineage displayed several genetic variations compared to other *S. pyogenes*

*emm*4 strains, including six unique SNPs and a 306 bp deletion in the *proA* gene, which is

involved in amino acid biosynthesis (see Table S2 in the Appendix for details).

Characteristically, M4$_{NL22}$ lacked the *hasABC* genes for synthesis of a hyaluronic acid

capsule[24]. Further characterisation through *in vitro* and *in vivo* experiments should reveal whether M4$_{NL22}$ is more transmissible than other *emm*4 *S. pyogenes*. No changes in known virulence determinants were observed (*nga-ifs-slo* region, superantigens, known virulence genes) in isolates from this M4$_{NL22}$ lineage.

The complete genome of isolate 2220758 harboured two intact prophages (closest PHASTER hits: Streptococcus phage 315.3, NCBI Nucleotide accession NC_004586, and Streptococcus phage 315.5, NC_004588), one possibly intact prophage (closest hit: Streptococcus phage 315.2, NC_004585, marked "questionable" by PHASTER) and two incomplete prophages. Streptococcus phage 315.3 harbours virulence-associated genes *hylP* and *ssa*, while Streptococcus phage 315.5 harbours *fbp*54 and *mf*3. In comparison to isolate ABC221, representing the genome most closely related to M4$_{NL22}$, isolate 2220758 did not harbour Streptococcus phage 315.4, which typically encodes a streptococcal DNase *sdn*. However, 14 of the M4$_{NL22}$ isolates did harbour the *sdn* gene, and possibly also the Streptococcus phage 315.4.

Discussion

Several novel *S. pyogenes* clones have emerged over the past years, including *emm*4[3], *emm*89[25], and *emm*1 lineages[26]. Although most emerged lineages seem endowed with increased capacity for transmission or virulence, there may be additional explanations for the emergence of M4$_{NL22}$ reported in the current study. As with many other infectious diseases[27], we have observed a marked shift in the epidemiology of *S. pyogenes* infections after COVID-19 pandemic restrictions were lifted, with a surprising absolute and proportional increase of *emm*4 isolates among iGAS cases.

There were only a few genetic differences between M4$_{NL22}$ and other *emm*4 *S. pyogenes*, including six SNPs and a unique in-frame deletion of *proA*, a gene involved in proline biosynthesis. The biological implications of these six SNPs and the *proA* deletion are unclear and should be investigated in more detail. No large-scale genomic rearrangements were observed for M4$_{NL22}$. However, 54% of M4$_{NL22}$ isolates harboured the streptococcal DNase *sdn*, encoded by Streptococcus phage 315.4. Since this prophage is lacking in the complete genome of 2220758, this may suggest the phage is able to move through the *S. pyogenes emm*4 population within a short time span. Remmington *et al*. previously showed that the hypervirulent clone of *emm*4 *S. pyogenes* contained many degraded and probably immobile prophages[28]. In contrast, prophages in M4$_{NL22}$ did not seem to be degraded or immobile based on our sequence analysis, which is expected as M4$_{NL22}$ has not descended from the hypervirulent clone.

In conclusion, we identified an emerging *emm*4 lineage in the Netherlands, which descended from a predominantly carriage-associated type. Although iGAS cases in late 2022 and early 2023 in the Netherlands are dominated by *emm*1 (more specifically M1$_{UK}$)[2], *emm*4 isolates are continuously detected in invasive and carriage sampling studies in the Netherlands (unpublished data). Continued bacteriological surveillance of isolates from both invasive disease and asymptomatic carriers remains critical to timely inform public health officials of circulating clones with possibly altered transmission or virulence capacities. Genomic analysis is required to do this effectively, as classical typing techniques (e.g. *emm*-typing), while valuable, have insufficient resolution to provide valuable information for strain comparison.

Acknowledgments

The authors would like to thank Agaath Arends-van 't Klooster, Ilse de Beer-Schuurman and Claudia Burger for their continued contributions as part of the NRLBM and the medical microbiology laboratories for continues submission of *S. pyogenes* isolates.

We thank SURF (www.surf.nl) for the support in using the Lisa Compute Cluster.

We thank Jeroen Bos, Sietze Brandes, Fabian Landman, Sjoerd Kuiling and Rob Mariman of the National Institute for Public Health and the Environment (RIVM) for the ONT sequencing of strain 2220758.

Appendix

-   Supplementary information: description of underlying datasets.

-   Table S1: Isolates included in this study.

-   Table S2: Annotations of SNPs associated with M4$_{NL22}$

References

1. Kempen EB van, Bruijning-Verhagen PCJ, Borensztajn D, et al. Increase in invasive group A streptococcal infections in children in the Netherlands, a survey among 7 hospitals in 2022. *medRxiv*. Published online December 13, 2022:2022.12.13.22283399. doi:10.1101/2022.12.13.22283399

2. Gier B de, Marchal N, Beer-Schuurman I de, et al. Increase in invasive group A streptococcal (*Streptococcus pyogenes*) infections (iGAS) in young children in the Netherlands, 2022. *Eurosurveillance*. 2023;28(1):2200941. doi:10.2807/1560-7917.ES.2023.28.1.2200941

3. DebRoy S, Sanson M, Shah B, et al. Population Genomics of *emm*4 Group A *Streptococcus* reveals progressive replacement with a hypervirulent clone in North America. *mSystems*. 2021;6(4):e00495-21. doi:10.1128/mSystems.00495-21

4. Hendrickx APA, Landman F, de Haan A, et al. Plasmid diversity among genetically related *Klebsiella pneumoniae* blaKPC-2 and blaKPC-3 isolates collected in the Dutch national surveillance. *Sci Rep*. 2020;10:16778. doi:10.1038/s41598-020-73440-2

5. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30(15):2114-2120. doi:10.1093/bioinformatics/btu170

6. Andrews S. FastQC a quality control tool for high throughput sequence data. *URL: https://www bioinformatics babraham ac uk/projects/fastqc*. Published online 2010.

7. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. *Genome Biol*. 2019;20(1):257. doi:10.1186/s13059-019-1891-0

8. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo Assembler. *Current Protocols in Bioinformatics*. 2020;70(1):e102. doi:10.1002/cpbi.102

9. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. 2018;34(18):3094-3100. doi:10.1093/bioinformatics/bty191

10. Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*. 2009;25(16):2078-2079. doi:10.1093/bioinformatics/btp352

11. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*. 2010;26(6):841-842. doi:10.1093/bioinformatics/btq033

12. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies. *Bioinformatics*. 2013;29(8):1072-1075. doi:10.1093/bioinformatics/btt086

13. Jolley KA, Bray JE, Maiden MCJ. A RESTful application programming interface for the PubMLST molecular typing and genome databases. *Database*. 2017;2017:bax060. doi:10.1093/database/bax060

14. Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic platform with an interactive web interface. *Nucleic Acids Research*. 2019;47(D1):D687-D692. doi:10.1093/nar/gky1080

15. Didelot X, Wilson DJ. ClonalFrameML: Efficient inference of recombination in whole bacterial genomes. *PLOS Computational Biology*. 2015;11(2):e1004041. doi:10.1371/journal.pcbi.1004041

16. Minh BQ, Schmidt HA, Chernomor O, et al. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Molecular Biology and Evolution*. 2020;37(5):1530-1534. doi:10.1093/molbev/msaa015

17. Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. Bakta: rapid and standardized annotation of bacterial genomes via alignment-free sequence identification. *Microb Genom*. 2021;7(11):000685. doi:10.1099/mgen.0.000685

18. Tonkin-Hill G, MacAlasdair N, Ruis C, et al. Producing polished prokaryotic pangenomes with the Panaroo pipeline. *Genome Biology*. 2020;21(1):180. doi:10.1186/s13059-020-02090-4

19. Lees JA, Croucher NJ, Goldblatt D, et al. Genome-wide identification of lineage and locus specific variation associated with pneumococcal carriage duration. Cobey S, ed. *eLife*. 2017;6:e26255. doi:10.7554/eLife.26255

20. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics*. 2018;34(15):2666-2669. doi:10.1093/bioinformatics/bty149

21. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology*. 2017;13(6):e1005595. doi:10.1371/journal.pcbi.1005595

22. Arndt D, Grant JR, Marcu A, et al. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res*. 2016;44(W1):W16-21. doi:10.1093/nar/gkw387

23. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. ACT: the Artemis comparison tool. *Bioinformatics*. 2005;21(16):3422-3423. doi:10.1093/bioinformatics/bti553

24. Flores AR, Chase McNeil J, Shah B, Van Beneden C, Shelburne SA III. Capsule-negative *emm* types are an increasing cause of pediatric Group A streptococcal infections at a large pediatric hospital in Texas. *Journal of the Pediatric Infectious Diseases Society*. 2019;8(3):244-250. doi:10.1093/jpids/piy053

25. Turner CE, Abbott J, Lamagni T, et al. Emergence of a new highly successful acapsular Group A *Streptococcus* clade of genotype *emm*89 in the United Kingdom. *mBio*. 2015;6(4):e00622-15. doi:10.1128/mBio.00622-15

26. Lynskey NN, Jauneikaite E, Li HK, et al. Emergence of dominant toxigenic M1T1 *Streptococcus pyogenes* clone during increased scarlet fever activity in England: a population-based molecular epidemiological study. *The Lancet Infectious Diseases*. 2019;19(11):1209-1218. doi:10.1016/S1473-3099(19)30446-3

27. Chow EJ, Uyeki TM, Chu HY. The effects of the COVID-19 pandemic on community respiratory virus activity. *Nat Rev Microbiol*. Published online October 17, 2022:1-16. doi:10.1038/s41579-022-00807-9

28. Remmington A, Haywood S, Edgar J, Green LR, de Silva T, Turner CE. Cryptic prophages within a *Streptococcus pyogenes* genotype emm4 lineage. *Microb Genom*. 2020;7(1):mgen000482. doi:10.1099/mgen.0.000482