F-actin and Myosin F control apicoplast elongation dynamics which drive
apicoplast-centrosome association in *Toxoplasma gondii*

Parvathi Madhavi Devarakonda[a], Valeria Sarmiento[a], Aoife T. Heaslip [a, #]

[a] Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT,
06269

Running Title: Mechanisms of apicoplast division

[#]Address correspondence to Aoife Heaslip, aoife.heaslip@uconn.edu

12
13

ABSTRACT

Toxoplasma gondii contains an essential plastid organelle called the apicoplast that is necessary for fatty acid, isoprenoid, and heme synthesis. Perturbations affecting apicoplast function or inheritance lead to parasite death. The apicoplast is a single copy organelle and therefore must be divided so that each daughter parasite inherits an apicoplast during cell division. In this study we identify new roles for F-actin and an unconventional myosin motor, TgMyoF, in this process. First, loss of TgMyoF and actin lead to an accumulation of apicoplast vesicles in the cytosol indicating a role for this actomyosin system in apicoplast protein trafficking or morphological integrity of the organelle. Second, live cell imaging reveals that during division the apicoplast is highly dynamic, exhibiting branched, U-shaped and linear morphologies that are dependent on TgMyoF and actin. In parasites where movement was inhibited by the depletion of TgMyoF, the apicoplast fails to associate with the parasite centrosomes. Thus, this study provides crucial new insight into mechanisms controlling apicoplast-centrosome association, a vital step in the apicoplast division cycle, which ensures that each daughter inherits a single apicoplast.

INTRODUCTION

The phylum Apicomplexa contains over 6,000 parasites, many of which are of significant medical and veterinary importance, including *Toxoplasma gondii* the causative agent of Toxoplasmosis and *Plasmodium spp.* which causes malaria [1]. These parasites are distinguished by the presence of vestigial non-photosynthetic plastid organelle named the apicoplast (<u>Apic</u>omplexan <u>plast</u>id), that is found in most Apicomplexa.

The apicoplast is an essential organelle required for fatty acid, isoprenoid, iron-sulfur cluster and heme synthesis [2–8]. Genetic manipulations or perturbation of the apicoplast with pharmacological compounds results in apicoplast inheritance defects and parasite death [9–11]. Despite containing a small 35kb genome, the majority of apicoplast localized proteins are encoded in the nucleus. Therefore, trafficking of these proteins to the apicoplast is a critical cellular process and our understanding of the mechanisms of apicoplast protein trafficking is incomplete. Many apicoplast proteins contain a bipartite N-terminal signal sequence that includes a signal peptide (SP), which directs protein translation to the endoplasmic reticulum (ER), and an apicoplast transit peptide which is revealed after removal of the SP [12,13]. It is unclear how these proteins translocate from the ER to the apicoplast. A subset of apicoplast proteins that localize within the apicoplast membranes, including APT1, Atrx1 and FtsH1, do not contain a signal peptide or apicoplast targeting sequence and have been observed in vesicles that are thought to directly traffic these proteins from the ER to the apicoplast in

a cell cycle dependent manner [14–19]. Proteins that localize to both the apicoplast and the mitochondria, are translocated via the Golgi and trafficking of these proteins to the apicoplast is perturbed by the addition of the ER retention motif HDEL [20]. Overall, there are numerous mechanisms controlling the trafficking of proteins to the apicoplast and the molecular details of each system are poorly understood.

Evolutionarily, the apicoplast was acquired by the parasite via a secondary endosymbiosis of red algae and contains four membranes [21]. The inner two membranes are derived from the progenitor chloroplast (inner most), the third from the algae plasma membrane (periplastid) and outermost membrane from plasma membrane of the Apicomplexan ancestor which first acquired this organelle (Fig. 1A). Protein translocons Tic and Toc in the inner two apicoplast membranes facilitate protein transport from the periplastid to the apicoplast lumen [22,23]. Repurposed ER-associated degradation (ERAD) proteins translocate proteins across the periplastid membrane [24,25]. How proteins traverse the outer membrane is not understood but, in some instances, may involve fusion of the outer membrane with ER derived vesicles [26].

The apicoplast is a single-copy organelle, thus apicoplast division and inheritance must occur with high accuracy to ensure parasite survival. Apicoplast division is a multistep process. During interphase the apicoplast has a circular morphology and is located at the parasite's apical end (Fig. 1A). At the beginning of cell division, the apicoplast interacts with the duplicated centrosomes and elongates [27].

This interaction between centrosomes and the apicoplast is dependent on the autophagy related protein Atg8 which localizes to the outer membrane of the apicoplast and disruption of Atg8 in both *Toxoplasma* and *Plasmodium* results in apicoplast inheritance defects [28–31]. While the apicoplast is elongated, daughter cell construction within the parasite cytosol is initiated [32]. The apical tip of the growing daughters is connected to the centrosomes via a fiber containing striated fiber assemblins (SFAs) [33], ensuring that each daughter cell inherits a single centrosome. As the centrosome migrates into the growing daughter scaffolds, the elongated apicoplast then extends to a U-shaped structure, whereupon apicoplast fission is mediated by dynamin-related protein A (DrpA) [34].

Three cytoskeletal proteins, actin (TgAct1), Myosin F (TgMyoF: an unconventional apicomplexan specific myosin motor), and the actin nucleator formin-2 also play major roles in apicoplast division [35–38]. Since TgMyoF has also been implicated in centrosome positioning, it was previously proposed that TgMyoF maintains the duplicated centrosomes in close proximity, thus allowing the apicoplast to associate with both structures [37]. However, it has also been shown that TgMyoF and actin control the transport of a variety of vesicles, including dense granules and Rab6 vesicles [38,39], thus an alternative hypothesis is that TgMyoF and actin are needed for the trafficking of vesicles to the apicoplast, which could provide new material (proteins and lipids) to the apicoplast prior to division.

In this study, we investigate the roles of TgMyoF and actin in apicoplast division.

We demonstrate that vesicles containing the apicoplast protein APT1 are formed in a

cell cycle dependent manner consistent with previous work [14]. Loss of MyoF and actin

lead to an increased number of vesicles in the cytosol but this cytoskeletal system does

not play a major role in vesicle movement. While APT1 protein translation at the ER

coincides with vesicle formation, we could not definitely determine that these vesicles

were ER-derived vesicles trafficking to the apicoplast.

Live cell imaging revealed that elongated apicoplasts are highly dynamic,

exhibiting linear, branched and U-shaped morphologies which are driven by both

TgMyoF and actin. Loss of this actomyosin system results in reduced apicoplast length

and defects in apicoplast-centrosome association. Thus, we conclude that MyoF and

actin play a vital role in recruiting the apicoplast to the duplicated centrosomes during

apicoplast division, a vital step that ensures each daughter inherits a single apicoplast.

MATERIAL AND METHODS

Cell culture

RH parasites were maintained by passage in human foreskin fibroblast (HFF) cells

obtained from ATCC (Catalog # CRL-2429). HFFs were grown to confluence in

Dulbecco's modified Eagle's medium containing 10% (v/v) heat inactivated fetal bovine

serum (FBS) and 1% (v/v) antibiotic/antimycotic at 37°C, 5% CO_2. Before parasite

addition, confluent HFF monolayers were maintained in Dulbecco's modified Eagle's

medium containing 1% (v/v) heat inactivated fetal bovine serum (FBS) and 1% (v/v)

antibiotic/antimycotic 37°C, 5% CO_2.

Drug treatment

Cytochalasin D treatment. For fixed cell imaging, parasites invaded confluent HFF

monolayer for 4 hours and then treated with 0.2 μM cytochalasin D (CD) or equivalent

volume of DMSO for 12 hours before fixation with 4% paraformaldehyde. For live cell

imaging, parasites were grown for 12 hours before treatment with 2 μM CD or

equivalent volume of DMSO for 30 minutes. Imaging was carried out in the presence of

DMSO or CD.

IAA Treatment: To deplete TgMyoF protein levels, *TgMyoF-mAID:TgAPT1-EmGFP*

parasites were grown in confluent HFF monolayer for 5 or 12 hours and then treated

with a final concentration of 500 μM indole acetic acid (IAA) or equivalent volume of

100% ethanol for 6 or 15 hours before fixation for immunohistochemistry or live cell

imaging [40,41].

Creation of transgenic parasite lines

A list of parasite lines, primers and gene accession numbers used in this study can be

found in Tables 1, 2 and 3 respectively.

Parasite transfection

Parasite transfection was conducted as previously described [42]. Briefly, linearized or

circular plasmid DNA or CRISPR HR oligo (produced by PCR) was ethanol precipitated,

washed twice 70% ethanol pre-chilled to -20°C and resuspended in cytomix (120mM

KCl, 10mM potassium phosphate (pH 7.6), 25mM HEPES (pH 7.6), 2mM EGTA (pH

7.6), 5mM MgCl$_2$, 2mM potassium ATP, 150uM CaCl$_2$, 5mM reduced glutathione) and

electroporated using a BTX electroporator set as follows: voltage 1500 V, capacitance

25 µF and resistance 25 Ω.

Creation of TgAPT1-EmGFP parasite line

In the *TgAPT1-EmGFP* parasite line the endogenous TgAPT1 gene is tagged with

EmeraldGFP (EmGFP). The pTKOII_APT1EmGFP plasmid was created from

pTKOII_MyoF-EmGFP plasmid [39] by digesting with AvrII and BglII restriction enzymes

to remove the TgMyoF genomic sequence. 2 kb of genomic DNA upstream of the

TgAPT1 stop site was amplified by PCR using Q5 high fidelity polymerase (New

England BioLabs) using the Ku80 genomic DNA as a template and the primers TgAPT1

F1 and TgAPT1 R1 (Table 2). The plasmid backbone and the PCR product were gel

purified. Ligation of the plasmid was carried out via Gibson assembly using NEB builder

HiFi DNA assembly master mix as per manufacturer's instructions (New England

Biolabs; catalog # M5520AA2) and transfected into NEB5α bacteria. Positive clones

were screened by colony PCR and verified by sanger sequencing. pTKOII-TgAPT1-

EmGFP plasmid was linearized for homologous recombination using the PstI restriction

enzyme and 25µg was transfected into 1x10^7 *ΔKu80: ΔHXGPRT* parental parasites

[43,44]. 24 hours after transfection mycophenolic acid in ethanol and Xanthine in 0.3M

NaOH were added to growth media to a final concentration of 25µg/ml and 50µg/ml

respectively. Drug treatment continued until at least 75% of the parasites were EmGFP

positive. Clonal parasites lines were FACS sorted into a single 96-well plate (2 per well)

and grown for 7 days to isolate clonal parasite lines resulting in the creation of TgAPT1-

EmGFP parasite line. To verify integration of this construct into the APT1 genomic

locus, genomic DNA was isolated using Qiagen DNeasy blood and tissue kit as per

manufacturer's instructions (Qiagen, Germantown, MD) and confirmed via PCR using

the primers TgAPT1-Promoter F1, APT1 CDS R, EmGFP R (Table 2) (Fig. S1).

Creation of TgMyoF-mAID::TgAPT1-EmGFP parasite line

In the *TgMyoF-mAID::TgAPT1-EmGFP* parasite line, the endogenous TgMyoF gene is

tagged with HA-mAID [39] and TgAPT1-EmGFP is expressed under the control of the

APT1 promoter from the dispensable UPRT locus. For the creation of 5'UPRT-pAPT1-

APT1-EmGFP-3'UPRT plasmid (which will be used as a template for created of the HR

oligo via PCR), 5'UPRT-pMyoF-mcherry-TgMyoF-3'UPRT (unpublished plasmid) was

digested with AvrII and AflII restriction enzyme to remove the pMyoF-mCherry-TgMyoF

sequence and replace with pAPT1-TgAPT1-EmGFP fragment, containing 1Kb of

upstream of the APT1 start codon and the APT1 coding sequence. TgAPT1-EmGFP

sequence was amplified from the pTKOII_APT1-EmGFP plasmid and primers 5'-APT1-

UPRT-F and 3'-APT1-UPRT-R (Table 2). The plasmid backbone and the PCR product

were gel purified. Ligation of the plasmid was carried out via Gibson assembly as per

manufacturer's instructions (New England Biolabs; catalog # M5520AA2) and

transfected into NEB5alpha bacteria. Positive clones were screened by colony PCR and

verified by sanger sequencing.

The CRISPR HR oligo was generated by PCR using UPRT F and UPRT R

primers and the 5'UPRT-pAPT1-APT1-EmGFP-3'UPRT plasmid as a template with Q5

high fidelity polymerase and 50ul of PCR reaction was cotransfected with 25 µg of

pSag1::Cas9::U6::sgUPRT plasmid [45] (AddGene plasmid # 54467) into 1x10^7

TgMyoF-mAID parasites [39]. 24 hours post-transfection parasites were treated with 10

µM fluorodeoxyribose (FUDR) (which selects for loss of the UPRT gene [46]) until at

least 75% of the parasites were EmGFP positive. Parasites were FACS sorted for

EmGFP positivity into a single 96-well plate (2 per well) and grown for 7 days. Single

clones were selected for further analysis. Genomic DNA was isolated using Qiagen

DNeasy blood and tissue kit as per manufacturer's instructions (Qiagen, Germantown,

MD). Genomic DNA was analyzed for correct insertion into the UPRT locus via PCR

using primers 5'-UPRT F, APT1 CDS R and UPRT-Intron R (Table 2) (Fig. S2).

Creation of TgMyoF-mAID; TgAPT1-Halo parasite line

In the *TgMyoF-mAID:TgAPT1-Halo* parasite line, the endogenous TgMyoF gene

is tagged with HA-mAID [39] and TgAPT1-Halo is expressed under the control of the

APT1 promoter from the dispensable UPRT locus. For the creation of 5'UPRT-pAPT1-

APT1-Halo-3'UPRT plasmid (which will be used as a template for created of the HR

oligo via PCR), the 5'UPRT-pAPT1-APT1-EmGFP-3'UPRT plasmid was digested with

AflII and BglII to remove the EmGFP coding sequence. The Halo coding sequence was

amplified using Halo-F1 and Halo-R2 primers using pmin-ActinCB-Halo as a template

(Table 2). The plasmid backbone and the PCR product were gel purified. Ligation of the

plasmid was carried out via Gibson assembly as per manufacturer's instructions (New

England Biolabs; catalog # M5520AA2) and transfected into NEB5alpha bacteria.

Positive clones were screened by colony PCR and verified by sanger sequencing.

The CRISPR HR oligo was generated by PCR using UPRT F and UPRT R

primers and the 5'UPRT-pAPT1-APT1-Halo-3'UPRT plasmid as a template with Q5

high fidelity polymerase and 50ul of PCR reaction was cotransfected with 25 µg of

pSag1::Cas9::U6::sgUPRT plasmid [45] (AddGene plasmid # 54467) into $1x10^7$

TgMyoF-mAID parasites [39]. 24 hours post-transfection parasites were treated with 10

µM fluorodeoxyribose (FUDR) (which selects for loss of the UPRT gene [46]) until at

least 75% of the parasites were EmGFP positive. Parasites were cloned by limited

dilution into a single 96-well plate (2 per well) and grown for 7 days. Single clones were

screen for positives by IFA.

Creation of ectopic ATG8 expression plasmid

Generation of pAtg8-3xTy-Atg8 plasmid for ectopic expression of TgAtg8 under the

endogenous promoter was a 2-step process. First, we created the pAtg8-tdTomato-Atg8

plasmid by digesting the pTub_tdTomato-Atg8 [28] with SfoI and BglII restriction

enzymes to remove the tubulin promoter sequence. TgAtg8 promoter was amplified

from the Ku80 parental genomic DNA using the primers Atg8 promoter F and Atg8

promoter R. Plasmid backbone and the PCR product was gel purified. Ligation of the

plasmid was carried out via Gibson assembly as per manufacturer's instructions (New

England Biolabs; catalog # M5520AA2) and transfected into NEB5alpha bacteria.

Positive clones were screened by colony PCR using Atg8 promoter F and Atg8

promoter R (Table 2) and verified by sanger sequencing. In the second step, pAtg8-

tdTomato-Atg8 was digested with AvrII and BglII restriction enzymes to remove

tdTomato coding sequence. The3xTy tag was created by PCR using primers 3xTy F

and 3xTy R (Table 2) and the pTKOII_MyoL-3xTy-mAID plasmid as a template

(unpublished plasmid; Table 1). Ligation of the plasmid with 3Xty insert was carried out

via Gibson assembly as per manufacturer's instructions (New England Biolabs; catalog

M5520AA2) and transfected into NEB5alpha bacteria. Positive clones were screened

by colony PCR using primers Atg8 promoter F and 3xTy R and verified by sanger

sequencing.25 μg of the pAtg8-3xTy-Atg8 plasmid was transiently transfected into the

TgMyoF-mAID parasite line and were grown for 12h before treating the parasites with

either ethanol or IAA for 6h.

Fluorescence microscopy

All live and fixed images were acquired on a DeltaVision Elite microscope system built

on an Olympus base with a 100x 1.39 NA objective in an environmental chamber

heated to 37 ˚C. This system is equipped with a 15-bit scientific CMOS camera and DV

Insight solid state illumination module with excitation wavelengths DAPI = 390/18nm,

FITC = 475/28nm, TRITC = 475/28nm and Alexa 647 = 632/22nm. Single band pass

emission filters had the following wavelengths DAPI 435/48nm, FITC = 525/48nm,

TRITC = 597/45nm and Alexa 647 = 679/34nm. Image acquisition speeds for the live

cell imaging were determined on a case-by-case basis and are indicated in the figure

legends. All cells were imaged in multiple focal planes and Z-series were captured at

0.2 µm (fixed) or 0.4 µm (live) steps. Images presented in the figures represent

maximum intensity Z-slice projections. Brightness and contrast of images were

optimized for print.

Immunocytochemistry and live cell imaging

Parasites were fixed with freshly made 4% paraformaldehyde (Electron

microscopy sciences, Catalog # 15714) in 1x PBS for 15 minutes at room temperature.

Cells were washed three times in 1x PBS and permeabilized in 0.25% Triton X-100

(ThermoFisher) in 1x PBS for 15 minutes at room temperature. Cells were washed

three times in 1x PBS and were blocked in 2% BSA in 1x PBS and incubated for 15

minutes. All antibodies were diluted in 2% BSA/1x PBS at the concentrations listed in

the Table 4. Cells were incubated sequentially with primary and secondary antibody at

room temperature for 30 minutes each, with 3 washes with 1x PBS between

incubations. To stain DNA parasites were incubated in 10 µM DAPI diluted in 1x PBS

for 10 minutes and then washed three times in 1x PBS. Coverslips were mounted onto

slides using either Prolong Gold (ThermoFisher Catalog # P36930) or Prolong diamond

(ThermoFisher Catalog # P36970) anti-fade reagent and allowed to dry overnight before

imaging.

For live cell imaging, growth media was replaced with Fluorobrite DMEM

(ThermoFisher) containing 1% FBS and 1x antimycotic/antibiotic pre-warmed to 37 ˚C.

The halo pulse chase assay was carried out as per manufacturer's instructions

(Promega, Inc). Parasites were grown overnight in confluent HFF monolayers in MetTak

dishes. Parasites were incubated for 15 minutes with 200nM Halo-ligand JF646 diluted

in growth media. Cells were wash three times in growth media and incubated for 30

minutes. Cells were washed again two times in growth media and incubated for 5.5

hours. Cells were than labeled in 5µM Halo-ligand TMR for 15 minutes. Cells were wash

three times in growth media and incubated for a further 30 minutes in growth media.

Cells were washed twice in Fluorobrite DMEM with 1% FBS before imaging cells live.

Image analysis

Quantification of TgAPT1 Vesicle number

To determine the number of APT1 vesicles during parasite division. The apicoplasts

were categorized according to their morphology to distinct stages of apicoplast division

stage (interphase, elongation, and post fission) by visualizing apicoplast morphology

with anti-Cpn60 IFA and expression of TgAPT1-EmGFP. Cell cycle stage by determined

by expressing ptub-mCherry-TubulinA1 a marker for the pellicle of daughter and mother

parasites. Number of vesicles were counted manually on average intensity projection

images using cell counter plugin in Fiji (Fiji is Just ImageJ; http://imageJ.net). Three

independent experiments were performed with 25 parasites per experiment counted for

each stage of division.

Characterization of TgAPT1 vesicle motion

To characterize TgAPT1 vesicle motion, vesicle tracking analysis was performed using

Imaris v9.9.0 microscopy image analysis software (Bitplane AG, Oxford Instruments). Z-

stack movies that were bleach corrected in ImageJ using the histogram matching

algorithm were uploaded to Imaris and brightness and contrast was auto-adjusted using

the "normalize time point" extension. Movies were cropped to remove areas that did not

contain apicoplast and vesicles. To prevent aberrant tracking inside the apicoplast itself,

the apicoplasts were demarcated from vesicles using the surface tool with a lower

surface threshold of 1500 arbitrary units (a.u.). Vesicles were identified using the spots

creation wizard. Size parameter (estimated XY diameter) was 0.3 µm. To distinguish

between vesicles and background fluorescence, only objects with a fluorescence

intensity above 150 a.u. were tracked. Tracks were created using autoregressive motion

algorithm. To be considered true vesicles, vesicles must be located within 5 µm from the

apicoplast surface, to prevent the aberrant detection of fluorescent puncta outside the

parasite, especially at later time points when photobleaching reduced the fluorescence

intensity of the vesicles. A maximum gap of 2 frames was permitted, (i.e., maximum

number of the consecutive time points that a track segment can be missed to be

considered continuous). Tracks with less than 5 data points were excluded from the

analysis as it was not possible to accurately assign a straightness value. Two

independent experiments with a total of 14-17 videos were used for this analysis.

Straightness values were calculated by determining the ratio of track displacement (the

distance between first and last positions of the vesicle) to track length (total distance

traveled).

Measuring apicoplast to centrosome distance

Distance between the elongated apicoplast and centrosomes was determined using the

straight-line tool in ImageJ by measuring the closest distance from each centrosome to

the apicoplast. The distance was considered as 0 when there was an overlap between

fluorescence pixels in each channel. Three independent experiments were performed

(N=108 and 93 from control and knockdown respectively).

Measuring apicoplast length

Elongated apicoplast length was measured using segmented/straight line tool in Fiji.

Three independent experiments were performed (N=60 parasites total).

Daughter cell-apicoplast proximity categories

For determining the apicoplast proximity to growing daughter cells, we visualized

apicoplast localization (Cpn60 staining) in growing daughter parasites (based on eGFP-

tubulin). Daughter proximity to the apicoplast was determined visually and categorized

as being near to both the daughters (2D), near to one of the daughters (1D) or not close

to either daughter (none). Three independent experiments were performed (N=60 total).

TgAtg8 localization

TgAtg8 localization during different stages of apicoplast division was determined using

anti-Ty antibody staining (kind donation of Dr. Chris de Graffenried: Brown University).

Two independent experiments were performed (25 parasites per experiment).

Statistical analysis

Statistical analysis was performed using GraphPad Prism software. Statistics on data sets were performed using unpaired t-tests. P-values <0.05 were considered statistically significant. Superplots were made as described in Lord et al., [47].

RESULTS

The number of TgAPT1-EmGFP vesicles in the cytosol increases during apicoplast elongation

To investigate the mechanisms of apicoplast division and apicoplast protein trafficking from the ER, we generated a parasite line where the endogenous TgAPT1 gene was tagged with EmGFP (Emerald green fluorescent protein) (Fig. 1A and 1B; Fig S1) as TgAPT1 localizes to vesicles in the parasite cytosol and the apicoplast [14,16,48]. Previously published data indicated that APT1 vesicles are formed in a cell cycle dependent manner so we transiently expressed mCherry-TubulinA1 which demarcates the parasite periphery of both the mother and growing daughter parasites to enable quantification of vesicle number at each stage of the parasite cell division cycle. Parasites were also stained with an anti-Cpn60 antibody, a marker of the apicoplast lumen. Based on the apicoplast shape and the presence and size of daughters, parasites were categorized as interphase (no daughter parasites, circular apicoplast), elongating (small daughters, elongated apicoplast) or post-fission (large daughters, circular apicoplast in each daughter) and the number of vesicles per parasite was determined at each cell cycle stage. In interphase parasites, TgAPT1 forms a ring that surrounds the TgCpn60 signal consistent with its localization at the apicoplast

membrane (Fig. 1B) and the number of TgAPT1 positive (TgAPT1+) vesicles per parasite is low (1.15±0.07, mean±SEM) (Fig. 1B and 1C). During elongation, the number of TgAPT1 vesicles in the cytosol increased 7-fold to 8.08±0.5 vesicles per parasite (Fig. 1B and 1C). After apicoplast fission and segregation into the daughter cells, the number of TgAPT1 vesicles per parasite decreased to 3.68±0.2 (Fig. 1B and 1C). Notably, Cpn60, a nuclear encoded protein that localizes to the apicoplast lumen, was not found in TgAPT1+ vesicles as previously reported [48] (Fig. 1B, inset). Consistent with the observations and interpretations made by [14], we hypothesize that these vesicles are ER-derived and transport newly synthesized proteins to the apicoplast in a cell cycle dependent manner.

To further investigate the origin of the APT1 vesicles, and determine if these vesicles are derived from or trafficking to the apicoplast, we designed a fluorescent pulse chase experiment using a parasite line expressing APT1-Halo under the endogenous APT1 promoter. Parasites were grown overnight and then pulse labeled with Halo-Ligand JF646 for 15 minutes (Fig. 1D). After this "pulse" labeling parasites were grown without ligand for 6 hours, labeled with Halo-Ligand TMR for 15 minutes (chase) and imaged immediately. Interphase parasites containing a circular apicoplast were labeled with JF646 only indicating that proteins synthesis did not occur after the first pulse labeling. In contrast, during apicoplast division, indicated by elongated apicoplasts, the apicoplast was labeled with both JF646 and TMR. In addition, Halo-Ligand TMR labeled newly synthesized protein that localized to the ER/nuclear

envelope and accumulated at the apical and basal ends of the nucleus (Fig. 1D; green

arrowheads). We did not observe Halo-JF646 labeled vesicles in the cytosol during

apicoplast elongation, as would be expected if the APT1 vesicles originated at the

apicoplast. Unfortunately, due to the prevalent ER localization of Halo-TMR it was not

possible to discern if TMR labeled vesicles were also present in the cell as would be

expected if the APT1 vesicles were derived from the ER. Notably, the ER localization of

APT1 was never observed in the APT1-EmGFP construct, likely due to inefficient

fluorescent protein maturation in the ER. Collectively, these results show that APT1

protein synthesis and the appearance of the APT1 vesicles occur in a cell cycle

dependent manner.

Actin depolymerization and TgMyoF knockdown leads to an increased number of
TgAPT1-EmGFP vesicles in the cytoplasm

TgAct1 and TgMyoF are cytoskeletal proteins with previously established roles in

vesicle transport [38,39]. Therefore, we wanted to determine if these proteins play a role

in vesicle trafficking to the apicoplast. To do this a previously generated auxin inducible

TgMyoF knockdown parasite line TgMyoF-mAID [39] was modified to express TgAPT1-

EmGFP under the endogenous TgAPT1 promoter from the dispensable UPRT locus to

create TgMyoF-mAID::TgAPT1-GFP parasite line (Fig. S2A).

To knockdown TgMyoF, TgMyoF-mAID::TgAPT1-EmGFP parasites were treated

for 15 hours with ethanol (control) or 500 μM IAA, which results in depletion of TgMyoF

to undetectable levels after 4 hours [39]. To depolymerize actin, TgAPT1-EmGFP

parasites were treated with DMSO (control) or 0.2 μM cytochalasin D (CD) for 12h prior

to fixation. Parasites transiently expressing mCherry-Tubulin were categorized as

interphase, elongation or post-fission as described for figure 1. Both actin

depolymerization and TgMyoF depletion resulted in an approximately 2-fold increase in

the number of TgAPT1-EmGFP positive vesicles at all stages of the apicoplast division

cycle (Fig. 2A-F; Fig. S3 & S4). In both treatment conditions, an apicoplast inheritance

defect was observed as expected [37] (Fig. 2A&2D; magenta arrow). "Apicoplast-less"

parasites contained the highest accumulation of vesicles in comparison to all other

stages of apicoplast division (Fig. 2C and F).

TgAPT1 vesicles exhibit predominately diffusive-like motion

Next, we sought to describe the motion characteristics of TgAPT1+ vesicles, as

previous work demonstrated that other vesicle types are transported throughout the

parasite cytosol in an actin and TgMyoF dependent manner [38,39]. TgAPT1-EmGFP

parasites in the elongation phase of apicoplast division were imaged by live cell

microscopy. Visual inspection of these recordings revealed two striking observations.

First, TgAPT1-EmGFP vesicles were found in the parasite cytosol, confirming

observations from the fixed cell experiments (Fig. 1 and 2). The motion of these vesicles

was predominately diffusive (Fig 3A; inset 2), although a small number of vesicles

exhibited directed motion, i.e., – sustained movement in a single direction for

continuous frames (Fig. 3A; inset 1) (Video S1). Secondly, elongated apicoplasts were

not static, linear structures, but exhibited changes in morphology throughout the 2-5min

imaging period (Fig. 5; Video S2).

To quantify vesicle movement in an automated and unbiased manner, vesicle

motion analysis was performed using the spot creation wizard in Imaris image analysis

software (https://imaris.oxinst.com/) (Fig. 3B). To prevent aberrant tracking of

fluorescent signal inside the apicoplast itself, a surface mask was created over the

apicoplast (Fig 3B; raw vs. surface). Vesicles were automatically detected based on

size and fluorescent intensity (Fig. 3B; surface and vesicles) (see materials and

methods for details). Vesicles that are transported in an active, motor driven manner

typically exhibit "directed trajectories", i.e.-vesicles move in approximately a straight

trajectory [38], while diffusive vesicles exhibit random movements. Thus, to determine

the number of vesicles exhibiting diffusive or directed movement, track straightness was

calculated by determining the ratio of track displacement (the distance between first and

last positions of the vesicle) to track length (total distance traveled). A straightness

value close to zero indicates random motion, i.e. – track length is greater than the

distance between the first and last positions indicating random motion. A straightness

value close to 1 indicates that the total distance travelled and the distance between the

first and last positions were similar indicating that the vesicle moved in approximately a

straight line. Vesicle tracks observed in the first 20 seconds of imaging are color-coded

according to straightness (Fig. 3B). A frequency distribution of the straightness values

indicates that over 80% of vesicles have a straightness value less than 0.5 and the

mean straightness value was 0.17±0.15 confirming that apicoplast vesicle motion is

predominantly diffusive in nature (Fig. 3C). We then compared the track displacement

length for trajectories that had a straightness value greater or less than 0.5. Randomly

moving vesicles (with a straightness value <0.5) had a mean track displacement value

of 0.3μm. While vesicles with straighter trajectories moved twice as far with a mean

displacement of 0.59μm (Fig. 3D).

Loss of TgMyoF and F-actin has only a minor impact TgAPT1-EmGFP vesicle
motion

Next, we sought to determine if actin depolymerization or TgMyoF knockdown

altered vesicle motion. TgAPT1-EmGFP parasites were grown for 18h prior to treatment

with 2 μM CD or equivalent volume of DMSO for 30 min before live cell imaging (Fig.

3E) (Video S3), while to deplete TgMyoF, TgMyoF-mAID::TgAPT1-EmGFP parasites

were grown for 15 hours with ethanol or 500μm IAA before live cell imaging (Fig. 3F)

(Video S4).

Vesicle movements were again tracked with Imaris as described above. Actin

depolymerization or TgMyoF knockdown had no effect on the mean straightness values

for DMSO, CD, Ethanol and IAA treated parasites were 0.18±0.18, 0.14±0.13, 0.2±0.17

and 0.17±0.17 respectively (Fig. 3G and 3H). Although, upon loss of F-actin and

TgMyoF, the number of vesicles in the cytosol was increased compared to controls. In

all cases (treatment and control) the percentage of vesicles with a straightness value

<0.5 was ~80%. To determine if the movement of the remaining ~20% of vesicles was

TgMyoF and actin driven, vesicles with a straightness value greater than 0.5 were

analyzed further. Since vesicles with a straightness value with greater than 0.5 exhibited

longer track displacement (Mean length was 0.59μm (Fig. 3D)) we defined "directed

trajectories" as those with a straightness value >0.5 and a track displacement length of

at least 0.5µm. In no treatment, DMSO- and ethanol-treated parasites, the percentage

of vesicles which met these criteria were 9.5%, 10.3% and 11% respectively. Upon actin

depolymerization with CD or TgMyoF knockdown this percentage was reduced to 4.7%

and 6.8% respectively (Fig. 3I). Although these changes are statistically significant,

overall our data indicates that only a small portion of APT1 vesicles are transported in

active, actomyosin dependent manner. There are two possible origins for these

vesicles: those which have bud from the ER and are destined for the apicoplast or

vesicles that have originated from the apicoplast.

TgMyoF knockdown results in reduced apicoplast length

If vesicle accumulation was due to either a defect in protein trafficking to the

apicoplast or fragmentation of the apicoplast, we predict that this would alter apicoplast

morphology. Therefore, we assessed apicoplast length in the elongation phase of

apicoplast division in the absence of TgMyoF. In order to compare parasites at the

same stage of cell division, parasites were transiently transfected with a GFP-tubulin

expression cassette in order to fluorescently label the pellicle of mother and daughter

parasites. Parasites were grown for 12 hours prior to initiating a 6-hour treatment with

ethanol or 500 µM IAA. This shortened IAA treatment time compared with previous

experiments allowed us to visualize the apicoplast in the first cell division cycle after

depletion of TgMyoF. Parasites were then stained with an anti-Cpn60 antibody and

DAPI before image acquisition (Fig. 4A). During elongation, TgMyoF knockdown

resulted in a ~25% reduction in apicoplast length (1.54$\pm$0.05µm, mean$\pm$SEM) when

compared to the control parasites (2.11±0.05 µm, mean±SEM) (Fig. 4B). Apicoplast

morphology was not disrupted in TgMyoF knockdown parasites in interphase and post-

fission stages of division (Fig. S5).

The apicoplast undergoes morphological changes during elongation

Live cell imaging of TgAPT1-EmGFP parasites demonstrated that the apicoplast

morphology in the elongation phase of apicoplast division was highly variable between

individual vacuoles. In some vacuoles, the apicoplast appeared to be highly dynamic

exhibiting alternating linear, U-shaped and branched morphologies throughout the ~2-

minute imaging period (Fig. 4A; top panel) (Video S2), while others were static in the

elongation phase of apicoplast division (Figure 5A; bottom panel) (Video S2). Given this

variability, we quantified the percentage of apicoplasts that were highly dynamic and

exhibited rapid changes in morphology during the course of imaging (102 seconds) (Fig.

5A and B). ""Dynamic" apicoplasts were defined as those that exhibited one or more of

the following morphological changes: extending and/or retracting, bending, or branch

formation, or branch movement (Fig. 5A; top panel). 63±6.1% (Mean±SEM) of the

apicoplasts were highly dynamic (Fig. 5B). Whereas 37±6.1% (Mean±SEM) of parasites

contained either less dynamic apicoplasts, exhibiting subtle changes in morphology, or

were static with no observable changes in the morphology throughout the ~2-minute

imaging time (Fig 5A and B).

Apicoplast dynamics requires actin and TgMyoF proteins

Considering the dynamic nature of the apicoplast, we wanted to determine if these dynamics are dependent on actin and TgMyoF. To depolymerize actin, TgAPT1-EmGFP parasites were again treated with 2 µM cytochalasin D or equivalent volume of DMSO for 30 min before live cell imaging (Fig. 5C). 82±2.9% (Mean±SEM) of the apicoplasts in control DMSO-treated parasites were highly dynamic, while only 9±6.6% (Mean±SEM) of the apicoplasts in the cytochalasin D treated parasites were highly dynamic (Fig. 5D). To test the role of TgMyoF in apicoplast dynamics, TgMyoF-mAID; TgAPT1-EmGFP parasites treated with 500 µM indole acetic acid (IAA) to induce TgMyoF knockdown for 15h prior to live cell imaging (Fig. 5E). Similar to the cytochalasin D treatment, TgMyoF knockdown resulted in reduced apicoplast dynamics. The percentage of parasites with a highly dynamic apicoplast was significantly reduced from 86±4% (Mean±SEM) in control parasites to 22±13% in the TgMyoF KD parasites (Fig. 5F).

The apicoplast dynamically associates with static centrosomes

Apicoplast association with the centrosomes is a crucial step in the apicoplast division cycle. In the first step in the cell division cycle, the centrosome migrates to the basal end of the nucleus and duplicates, then both centrosomes return to the apical side of the nucleus where they associate with the apicoplast [27,49]. The dynamics of the elongated apicoplast (Fig. 5) have not previously been described and it is unclear if or how the apicoplast interaction with the centrosomes, which are embedded in the nuclear envelope [50], is maintained during apicoplast movement.

To further explore this phenomenon, wild type parasites were transiently

transfected with plasmids containing FNR-RFP (a marker for the apicoplast; [27]) and

Centrin 1-GFP (a marker for the centrosome [51]) expression cassettes and visualized

using live-cell microscopy. In parasites with highly dynamic, elongated apicoplasts the

centrosomes were not always associated with the tips of the apicoplast but were

frequently associated with the sides of the elongated apicoplast (Fig. 6A; inset 1 and 3)

(Video S5). The apicoplast appears to slide in relation to the static centrosome (Fig. 6A,

Inset 1, Video S6). The apicoplast was observed to undergo branching when the

centrosome was associated with the side rather than the tip of the apicoplast (Fig. 6A;

Inset 3, magenta arrow) (Video S7). The most dramatic morphological changes

occurred when the apicoplast was associated with only a single centrosome (Fig. 6A,

Inset 2). The apicoplast was observed extending towards the second centrosome

before retracting backwards. In contrast to the dynamics observed in the elongation

phase of apicoplast division, interphase and post-fission apicoplasts were static. In

interphase parasites the apicoplast was circular and did not change shape over the

course of imaging (Fig. 6B; upper panel) (Video S8). After apicoplast fission, the

apicoplast remains elongated for a time before retraction into a circular morphology.

Post-fission elongated apicoplasts are static over the course of the imaging period. The

apicoplast movement and branching that were observed prior to fission are no longer

observed and centrosomes are statically associated with the apicoplast tips (Fig. 6B;

lower panel) (Videos S9).

TgMyoF knockdown disrupts apicoplast and centrosome interaction

The live cell imaging of apicoplast and centrosome interactions, suggest that the actomyosin-driven apicoplast dynamics are important for initiating the interaction between these two structures and positioning of the centrosomes at the apicoplast tips (Fig. 6). Therefore, we directly investigated if the loss of TgMyoF, which disrupted apicoplast dynamics (Fig. 5), also disrupted apicoplast-centrosome interactions. TgMyoF-mAID parasites transiently expressing the centrosome marker Centrin1-GFP were grown for 12 hours post invasion and then treated with ethanol or 500 μM IAA for 6 hours before fixation and staining with an anti-Cpn60 antibody and DAPI. In control parasites, at the early stages of apicoplast division when the length of the elongated apicoplast is short (less than 1.5μm), the centrosomes and the apicoplast are frequently not associated with one another (Fig. 7A) and the mean apicoplast to centrosome distance is 0.99±0.18μm (Mean±SEM) (Fig. 7B&C). Later in the apicoplast division cycle, when the length of the apicoplast is greater than 1.5μm, the mean centrosome-apicoplast distance is significantly shorter at only 0.36±0.05μm (Fig. 7A-C) and the centrosomes were close with either the sides or the tips of the apicoplast (Fig. 7A). However, in TgMyoF knockdown parasites the apicoplast to centrosome distances were 1.2±0.12 and 1.0±0.09 μm when the apicoplast length was greater than or less than 1.5μm respectively. (Fig. 7A-C) indicating that TgMyoF is needed for recruiting the apicoplast to the centrosomes.

We next assessed how the loss of centrosome interaction affected apicoplast

inheritance. In 100% of control parasites, elongated apicoplasts were found to be in

close proximity to both of the growing daughter parasites. In early stages of division, the

apicoplast was adjacent to the daughter parasites (Fig. 8A; white arrow head). Later in

division, both daughters contain one end of the elongated apicoplast, even when one

the daughters was in the "upside down" orientation, (i.e. the apical ends of the daughter

parasites have opposite orientations within the mother) (Fig 7C; magenta arrowhead).

Upon depletion of TgMyoF only 13.3% of elongated apicoplasts were in close proximity

to both the growing daughter parasites (2D) while 70% of the elongated apicoplasts

were in close proximity to only one of the daughter parasites (1D) (Fig. 8A; white arrow),

and 16.6% of the elongated apicoplasts were not found to be in close proximity to either

of the growing daughter parasites (none) (Fig. 8B and Fig. S6). Thus, the disruption of

the centrosome-apicoplast interaction leads to asymmetric segregation of the apicoplast

into a single daughter cell or to the residual body.

TgMyoF knockdown did not alter TgAtg8 localization with the apicoplast

TgAtg8 is a protein that localizes to the apicoplast outer membrane. Loss of

TgAtg8 disrupts the interaction between the centrosomes and apicoplast resulting in

asymmetric distribution of the apicoplast to the growing daughter parasites, reminiscent

of the phenotype observed after TgMyoF depletion (Fig. 8A-B) [28]. To determine if loss

of TgMyoF affected the localization of TgAtg8 at the apicoplast, TgMyoF-mAID

parasites transiently expressing 3xTy-TgAtg8 under the endogenous TgAtg8 promoter

were treated with ethanol or 500 µM IAA for 6h before fixation and staining with anti-

Cpn60 antibody. TgAtg8 localized to the apicoplast at all stages of the apicoplast

division cycle and knockdown of TgMyoF did not disrupt TgAtg8 localization (Fig. 8C

and Fig. S7) indicating that the loss of interaction between centrosomes and the

elongated apicoplast seen in the TgMyoF knockdown parasites is not due to

mislocalization of TgAtg8.

DISCUSSION

The apicoplast is an essential organelle in *T. gondii*. In order to maintain a functional

apicoplast, the parasite must execute two distinct and essential cellular processes:

Division of this single copy organelle once per cell cycle to ensure each daughter

parasite inherits one apicoplast, and trafficking of nuclear encoded proteins to the

apicoplast. We investigated the role of actin and an associated myosin motor, TgMyoF,

in these processes.

Loss of MyoF and actin results in the accumulation of TgAPT1 vesicles in the

parasite cytosol. Vesicle trafficking is a multi-step process involving vesicle formation

and budding from the donor compartment, transport, docking at the target membrane

and finally vesicle fusion. While it is recognized that trafficking of nuclear encoded

proteins to the apicoplast is a vital cellular process, our understanding of this trafficking

process is incomplete and a number of trafficking pathways have been proposed [52].

Using TgAPT1-EmGFP expressing parasites, our data confirms previously published

data [14] demonstrating that vesicle formation is coordinated with the cell cycle, with

the number of vesicles increasing 7-fold during the elongation phase of apicoplast

division. We were not able to definitively determine the origin of these vesicles and we

considered two possible hypotheses: First, APT1 vesicles originate at the apicoplast

and represent a mechanism for trafficking apicoplast-synthesized lipids to other

subcellular locations, or second, these vesicles contain newly synthesized nuclear

encoded apicoplast proteins that originate at the ER, and represent a mechanism for

trafficking these proteins to the apicoplast as previously proposed [14,48]. A number of

lines of evidence led us to favor the latter hypothesis: (1) vesicle accumulation occurs

in "apicoplast-less" parasites indicating that vesicle formation is not dependent on the

apicoplast itself, as previously observed (Fig. 2) [48]. (2) APT1 protein synthesis and

vesicle formation both occur in a short window in the cell cycle when the apicoplast is

undergoing division (Fig. 1), (3) recent work has shown that the SNARE proteins

TgVAMP4-2 and TgSNAP29 localize to the apicoplast and are essential for the

trafficking of nuclear encoded proteins to the apicoplast [53,54], which highlight a role

for vesicles in protein trafficking to the apicoplast, and (4) while we cannot exclude the

possibility that these vesicles are derived from the apicoplast itself, vesicle formation at

and budding from the apicoplast has never been described in the literature.

Additionally, chloroplast to ER lipid trafficking in plants occurs by Acyl-coA's and does

not involve vesicle formation [55]. We had hoped our Pulse-Chase assay could be

used to determine the origin of the APT1 vesicles and while we did not observe

apicoplast derived vesicles (labeled with JF646), we also did not observe Halo-TMR

labeled vesicles. We speculate that the pervasive Halo-TMR signal from the ER

hampered our ability to visualize APT1-TMR vesicles in the cytosol.

A 12-hour treatment with CD to induce actin depolymerization or a 15-hour

treatment with IAA to deplete TgMyoF caused to a significant increase in the number of

vesicles in the cytosol at all phases of the cell division cycle (Fig. 2). It is possible that

these vesicles are ER-derived but fail to fuse with the apicoplast and thus accumulate

in the cytosol during subsequent cell division cycles. Alternatively, loss of these

proteins may result in vesiculation of the apicoplast itself.

MyoF and actin only play a minor role in APT1 vesicle motion. Given the role of

MyoF and actin in dense granule transport, we sought to determine if this cytoskeletal

system was also required for APT1 vesicle transport. By tracking the motion of

individual vesicles in live parasites we demonstrate that APT1 vesicle motion appears

predominately diffusive-like. With ~80% of vesicles moving in a random manner, i.e.,

those with a straightness value less than 0.5 (Fig. 3). "Directed" movements were

defined as those with a straightness value greater than 0.5 and a displacement length

greater than 500nm. Only ~10% of vesicles in control parasites met these criteria.

Actin depolymerization or TgMyoF depletion reduced this percentage by ~50% (Fig. 3).

Therefore, we conclude that only a small proportion of vesicles are transported in an

actomyosin dependent manner. Given that 90% of vesicles exhibit diffusive motion it is

unlikely that the small change in the percentage of vesicles exhibiting directed motor

driven movement can account for the large accumulation in vesicles seen after CD and

IAA treatment.

MyoF and Actin control apicoplast dynamics which mediate apicoplast-

centrosome association. The sequence of events that occur during the apicoplast

division cycle had previously been established via fixed cell immunofluorescence

assays or live-cell imaging with low temporal resolution (time between frames on the

order of minutes) [27,32] (Fig. 1A). This study reveals new insight into how this

sequence of events proceeds (Fig. 9). The first step in the division cycle is the

transition from circular to elongated morphology. Elongated apicoplasts with a length

<1.5µm were on average ~1µm away from the centrosomes. Thus, the transition from

circular to elongated occurs before, and is not dependent on centrosome association.

This transition also occurred in the absence of both actin and TgMyoF, indicating that

an as yet unidentified protein(s) must control apicoplast membrane deformation.

Since apicoplast elongation is not dependent on centrosome association a

mechanism must be in place to drive the interaction of these two organelles. The high-

temporal resolution live cell imaging presented here provided insight into how this is

achieved. Apicoplast branching, extension and retraction are needed to move the

apicoplast in close proximity to the nuclear envelope-embedded centrosomes to

facilitate an interaction with either the sides or tips of the apicoplast. Centrosome

sliding along the apicoplast side and branch formation and retraction are likely

important for positioning of the centrosome at the apicoplast tips. Consistent with this,

the localization of TgAtg8 which plays a role in mediating these interactions, is not

restricted to the apicoplast tips but found on the entire apicoplast surface (Fig. 8).

Loss of TgMyoF results in decreased apicoplast dynamics (Fig. 5) and centrosome

association (Fig. 7), which ultimately leads to apicoplast inheritance defects (Fig. 8).

Despite the defects in centrosome-apicoplast association in the absence of TgMyoF,

TgAtg8 remains localized at the apicoplast. Collectively, this data indicates that

actomyosin-driven apicoplast dynamics are important to facilitate the association

between these two organelles and for positioning of the centrosomes at the apicoplast

tips.

There are still a number of important questions that need to be addressed in future

studies. Previous work [35] has demonstrated that the actin nucleator formin 2,

relocalizes from the Golgi to the centrosomes during apicoplast division, however it is

unclear how F-actin interacts with the apicoplast in order to drive apicoplast dynamics.

Since MyoF and actin are not required for apicoplast elongation, another as yet

unidentified protein or proteins must drive membrane deformation and the dramatic

transition from a circular to elongated morphology that occurs at the start of apicoplast

division.

Overall, our results have determined that MyoF and actin control apicoplast

dynamics which is needed for apicoplast-centrosome association, a vital step of the

apicoplast division cycle.

ACKNOWLEDGEMENTS

We thank the members of the Heaslip Lab for helpful discussion during the course of

these experiments. We thank our colleagues for sharing reagents: Dr. Gary Ward

(University of Vermont) for the IMC-1 antibody; Dr. Boris Striepen (University of

Pennsylvania) for the anti-Cpn60 antibody. Dr. Ke Hu (Arizona State University) for

pTub-mcherry-TubA1 and pTub-eGFP-TubA1, pTub-FNR-RFP plasmids. Dr. Sebastian

Besteiro (Univ. Montpellier, France) for the pTub-tdTomato-TgAtg8 expression plasmid.

Schematics in figures 1 and 9 were made with Biorender.

FUNDING INFORMATION:

This work was funded by the National Institute of General Medical Sciences

1R35GM138316-01 awarded to ATH. The funders had no role in study design, data

collection and interpretation, the decision to submit the work for publication or

manuscript preparation. The authors declare that no competing interests exist.

REFERENCES

1. Torgerson PR, Mastroiacovo P. The global burden of congenital toxoplasmosis: a systematic review. Bull World Heal Organ. 2013 Jul;91(7):501–8.

2. Renaud EA, Pamukcu S, Cerutti A, Berry L, Lemaire-Vieille C, Yamaryo-Botté Y, et al. Disrupting the plastidic iron-sulfur cluster biogenesis pathway in Toxoplasma gondii has pleiotropic effects irreversibly impacting parasite viability. J Biol Chem [Internet]. 2022 Aug 1 [cited 2022 Nov 2];298(8). Available from: http://www.jbc.org/article/S0021925822006858/fulltext

3. Waller RF, Keeling PJ, Donald RGK, Striepen B, Handman E, Lang-Unnasch N, et al. Nuclear-encoded proteins target to the plastid in Toxoplasma gondii and Plasmodium falciparum. Proc Natl Acad Sci U S A. 1998;95(21):12352–7.

4. Yeh E, DeRisi JL. Chemical Rescue of Malaria Parasites Lacking an Apicoplast Defines Organelle Function in Blood-Stage Plasmodium falciparum. PLOS Biol [Internet]. 2011 Aug [cited 2022 Nov 2];9(8):e1001138. Available from: https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.1001138

5. Lim L, McFadden GI. The evolution, metabolism and functions of the apicoplast. Philos Trans R Soc B Biol Sci [Internet]. 2010 Mar 12 [cited 2022 Nov 2];365(1541):749–63. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2009.0273

6. Bergmann A, Floyd K, Key M, Dameron C, Rees KC, Brock Thornton L, et al.

Toxoplasma gondii requires its plant-like heme biosynthesis pathway for infection.

PLOS Pathog [Internet]. 2020 May 1 [cited 2022 Nov 2];16(5):e1008499.

Available from:

https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1008499

7. Sheiner L, Vaidya AB, McFadden GI. The metabolic roles of the endosymbiotic organelles of Toxoplasma and Plasmodium spp. Curr Opin Microbiol. 2013 Aug 1;16(4):452–8.

8. Brooks CF, Johnsen H, van Dooren GG, Muthalagi M, Lin SS, Bohne W, et al. The Toxoplasma Apicoplast Phosphate Translocator Links Cytosolic and Apicoplast Metabolism and Is Essential for Parasite Survival. Cell Host Microbe [Internet]. 2010 Jan 21 [cited 2022 Nov 2];7(1):62–73. Available from: http://www.cell.com/article/S1931312809004120/fulltext

9. Fichera ME, Roos DS. A plastid organelle as a drug target in apicomplexan parasites. Nature. 1997;390(6658):407–9.

10. He CY, Shaw MK, Pletcher CH, Striepen B, Tilney LG, Roos DS. A plastid segregation defect in the protozoan parasite Toxoplasma gondii. EMBO J. 2001;20(3):330–9.

11. Pfefferkorn ER, Nothnagel RF, Borotz SE. Parasiticidal effect of clindamycin on Toxoplasma gondii grown in cultured cells and selection of a drug-resistant mutant. Antimicrob Agents Chemother [Internet]. 1992 [cited 2022 Nov

731 2];36(5):1091–6. Available from:

732 https://journals.asm.org/doi/10.1128/AAC.36.5.1091

733 12. Harb OS, Chatterjee B, Fraunholz MJ, Crawford MJ, Nishi M, Roos DS. Multiple

734 functionally redundant signals mediate targeting to the apicoplast in the

735 apicomplexan parasite Toxoplasma gondii. Eukaryot Cell [Internet]. 2004 Jun

736 [cited 2022 Nov 2];3(3):663–74. Available from:

737 https://journals.asm.org/doi/10.1128/EC.3.3.663-674.2004

738 13. DeRocher A, Hagen CB, Froehlich JE, Feagin JE, Parsons M. Analysis of

739 targeting sequences demonstrates that trafficking to the Toxoplasma gondii

740 plastid branches off the secretory system. J Cell Sci [Internet]. 2000 Nov 15 [cited

741 2022 Nov 2];113(22):3969–77. Available from:

742 https://journals.biologists.com/jcs/article/113/22/3969/26332/Analysis-of-targeting-

743 sequences-demonstrates-that

744 14. Karnataki A, DeRocher A, Coppens I, Nash C, Feagin JE, Parsons M. Cell cycle-

745 regulated vesicular trafficking of Toxoplasma APT1, a protein localized to multiple

746 apicoplast membranes. Mol Microbiol [Internet]. 2007 Mar 1 [cited 2023 Jun

747 15];63(6):1653–68. Available from:

748 https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2958.2007.05619.x

749 15. Parsons M, Karnataki A, Feagin JE, DeRocher A. Protein trafficking to the

750 apicoplast: deciphering the apicomplexan solution to secondary endosymbiosis.

751 Eukaryot Cell [Internet]. 2007 Jul [cited 2022 Dec 14];6(7):1081–8. Available from:

https://pubmed.ncbi.nlm.nih.gov/17513565/

16. Derocher AE, Karnataki A, Vaney P, Parsons M. Apicoplast Targeting of a

Toxoplasma gondii Transmembrane Protein Requires a Cytosolic Tyrosine-Based

Motif. Traffic. 2012;13(5):694–704.

17. DeRocher AE, Coppens I, Karnataki A, Gilbert LA, Rome ME, Feagin JE, et al. A

thioredoxin family protein of the apicoplast periphery identifies abundant

candidate transport vesicles in Toxoplasma gondii. Eukaryot Cell [Internet].

2008;7(9):1518–29. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/18586952

18. Karnataki A, Derocher AE, Coppens I, Feagin JE, Parsons M. A Membrane

Protease is Targeted to the Relict Plastid of Toxoplasma via an Internal Signal

Sequence. Traffic [Internet]. 2007 Nov 1 [cited 2022 Nov 4];8(11):1543–53.

Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/j.1600-

0854.2007.00637.x

19. Karnataki A, DeRocher AE, Feagin JE, Parsons M. Sequential processing of the

Toxoplasma apicoplast membrane protein FtsH1 in topologically distinct domains

during intracellular trafficking. Mol Biochem Parasitol. 2009 Aug 1;166(2):126–33.

20. Prasad A, Mastud P, Patankar S. Dually localised proteins found in both the

apicoplast and mitochondrion utilize the Golgi-dependent pathway for apicoplast

targeting in Toxoplasma gondii. Biol Cell [Internet]. 2021 Jan 1 [cited 2023 Jun

15];113(1):58–78. Available from:

https://onlinelibrary.wiley.com/doi/full/10.1111/boc.202000050

21. McFadden GI, Yeh E. The apicoplast: now you see it, now you don't. Int J

Parasitol [Internet]. 2017 Feb 1 [cited 2022 Dec 14];47(2–3):137–44. Available

from: https://pubmed.ncbi.nlm.nih.gov/27773518/

22. Van Dooren GG, Tomova C, Agrawal S, Humbel BM, Striepen B. Toxoplasma

gondii Tic20 is essential for apicoplast protein import. Proc Natl Acad Sci U S A

[Internet]. 2008 Sep 9 [cited 2022 Dec 14];105(36):13574–9. Available from:

https://pubmed.ncbi.nlm.nih.gov/18757752/

23. Sheiner L, Fellows JD, Ovciarikova J, Brooks CF, Agrawal S, Holmes ZC, et al.

Toxoplasma gondii Toc75 Functions in Import of Stromal but not Peripheral

Apicoplast Proteins. Traffic [Internet]. 2015 Dec 1 [cited 2022 Dec

14];16(12):1254–69. Available from: https://pubmed.ncbi.nlm.nih.gov/26381927/

24. Felsner G, Sommer MS, Gruenheit N, Hempel F, Moog D, Zauner S, et al. ERAD

components in organisms with complex red plastids suggest recruitment of a

preexisting protein transport pathway for the periplastid membrane. Genome Biol

Evol [Internet]. 2011 [cited 2022 Dec 14];3(1):140–50. Available from:

https://pubmed.ncbi.nlm.nih.gov/21081314/

25. Sommer MS, Gould SB, Lehmann P, Gruber A, Przyborski JM, Maier UG. Der1-

mediated preprotein import into the periplastid compartment of chromalveolates?

Mol Biol Evol [Internet]. 2007 Apr [cited 2022 Dec 14];24(4):918–28. Available

from: https://pubmed.ncbi.nlm.nih.gov/17244602/

26. Mallo N, Fellows J, Johnson C, Sheiner L. Protein Import into the Endosymbiotic

Organelles of Apicomplexan Parasites. Genes (Basel) [Internet]. 2018 Aug 14

[cited 2022 Dec 14];9(8). Available from:

https://pubmed.ncbi.nlm.nih.gov/30110980/

27. Striepen B, Crawford MJ, Shaw MK, Tilney LG, Seeber F, Roos DS. The plastid

of Toxoplasma gondii is divided by association with the centrosomes. J Cell Biol.

2000;151(7):1423–34.

28. Lévêque MF, Berry L, Cipriano MJ, Nguyen HM, Striepen B, Besteiro S.

Autophagy-related protein ATG8 has a noncanonical function for apicoplast

inheritance in toxoplasma gondii. MBio [Internet]. 2015 Oct 27 [cited 2022 Nov

4];6(6). Available from: https://journals.asm.org/doi/10.1128/mBio.01446-15

29. Cheng L, Tian Y, Wang Y, Wang T, Yao Y, Yu H, et al. Toxoplasma TgAtg8-

TgAtg3 Interaction Primarily Contributes to Apicoplast Inheritance and Parasite

Growth in Tachyzoite . Microbiol Spectr [Internet]. 2022 Feb 23 [cited 2022 Nov

4];10(1). Available from: https://journals.asm.org/doi/10.1128/spectrum.01495-21

30. Mizushima N, Sahani MH. ATG8 localization in apicomplexan parasites.

http://dx.doi.org/104161/auto32183 [Internet]. 2014 Sep 1 [cited 2022 Nov

4];10(9):1487–94. Available from:

https://www.tandfonline.com/doi/abs/10.4161/auto.32183

31.   Walczak M, Ganesan SM, Niles JC, Yeh E. ATG8 Is Essential Specifically for an

Autophagy-Independent Function in Apicoplast Biogenesis in Blood-Stage Malaria

Parasites. MBio [Internet]. 2018 Jan 1 [cited 2022 Nov 4];9(1). Available from:

https://pubmed.ncbi.nlm.nih.gov/29295911/

32.   Nishi M, Hu K, Murray JM, Roos DS. Organellar dynamics during the cell cycle of

Toxoplasma gondii. J Cell Sci [Internet]. 2008;121(Pt 9):1559–68. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/18411248

33.   Francia ME, Striepen B. Cell division in apicomplexan parasites. Nat Rev

Microbiol [Internet]. 2014;12(2):125–36. Available from:

http://dx.doi.org/10.1038/nrmicro3184

34.   van Dooren GG, Reiff SB, Tomova C, Meissner M, Humbel BM, Striepen B. A

novel dynamin-related protein has been recruited for apicoplast fission in

Toxoplasma gondii. Curr Biol [Internet]. 2009;19(4):267–76. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/19217294

35.   Tosetti N, Pacheco NDS, Favre DS, Jacot D. Three f-actin assembly centers

regulate organelle inheritance, cell-cell communication and motility in toxoplasma

gondii. Elife. 2019 Feb 1;8.

36.   Stortz JF, Del Rosario M, Singer M, Wilkes JM, Meissner M, Das S. Formin-2

drives polymerisation of actin filaments enabling segregation of apicoplasts and

cytokinesis in Plasmodium falciparum. Elife. 2019;

37.   Jacot D, Daher W, Soldati-Favre D. Toxoplasma gondii myosin F, an essential

motor for centrosomes positioning and apicoplast inheritance. EMBO J [Internet].

2013/05/23. 2013; Available from: http://www.ncbi.nlm.nih.gov/pubmed/23695356

38.   Heaslip AT, Nelson SR, Warshaw DM. Dense granule trafficking in Toxoplasma

gondii requires a unique class 27 myosin and actin filaments. Mol Biol Cell

[Internet]. 2016;27(13):2080–9. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/27146112

39.   Carmeille R, Lomoriello PS, Devarakonda PM, Kellermeier JA, Heaslip AT. Actin

and an unconventional myosin motor, TgMyoF, control the organization and

dynamics of the endomembrane network in Toxoplasma gondii [Internet]. Vol. 17,

PLoS Pathogens. 2021. 1–32 p. Available from:

http://dx.doi.org/10.1371/journal.ppat.1008787

40.   Brown K, Long S, Sibley L. Conditional Knockdown of Proteins Using Auxin-

inducible Degron (AID) Fusions in Toxoplasma gondii. BIO-PROTOCOL. 2018;

41.   Brown KM, Long S, Sibley LD. Plasma membrane association by N-acylation

governs PKG function in Toxoplasma gondii. MBio. 2017;

42.   Roos DS, Donald RG, Morrissette NS, Moulton AL. Molecular tools for genetic

dissection of the protozoan parasite Toxoplasma gondii. Methods Cell Biol

[Internet]. 1994;45:27–63. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/7707991

43.   Fox BA, Ristuccia JG, Gigley JP, Bzik DJ. Efficient gene replacements in

Toxoplasma gondii strains deficient for nonhomologous end joining. Eukaryot Cell

[Internet]. 2009 Apr [cited 2022 Nov 4];8(4):520–9. Available from:

https://journals.asm.org/doi/10.1128/EC.00357-08

44.   Huynh MH, Carruthers VB. Tagging of endogenous genes in a Toxoplasma gondii

strain lacking Ku80. Eukaryot Cell [Internet]. 2009 Apr [cited 2022 Nov

4];8(4):530–9. Available from: https://journals.asm.org/doi/10.1128/EC.00358-08

45.   Shen B, Brown KM, Lee TD, David Sibley L. Efficient gene disruption in diverse

strains of Toxoplasma gondii using CRISPR/CAS9. MBio [Internet]. 2014 May 13

[cited 2022 Dec 14];5(3). Available from:

https://pubmed.ncbi.nlm.nih.gov/24825012/

46.   Donald RGK, Roos DS. Insertional mutagenesis and marker rescue in a

protozoan parasite: cloning of the uracil phosphoribosyltransferase locus from

Toxoplasma gondii. Proc Natl Acad Sci U S A [Internet]. 1995 Jun 6 [cited 2022

Dec 14];92(12):5749–53. Available from:

https://pubmed.ncbi.nlm.nih.gov/7777580/

47.   Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. SuperPlots: Communicating

reproducibility and variability in cell biology. J Cell Biol. 2020;

48.   Bouchut A, Geiger JA, Derocher AE, Parsons M. Vesicles bearing Toxoplasma

apicoplast membrane proteins persist following loss of the relict plastid or Golgi

body disruption. PLoS One [Internet]. 2014 Nov 1 [cited 2022 Dec 14];9(11).

Available from: https://pubmed.ncbi.nlm.nih.gov/25369183/

49.   Hartmann J, Hu K, He CY, Pelletier L, Roos DS, Warren G. Golgi and centrosome

cycles in Toxoplasma gondii. Mol Biochem Parasitol. 2006 Jan;145(1):125–7.

50.   Suvorova ES, Francia M, Striepen B, White MW. A Novel Bipartite Centrosome

Coordinates the Apicomplexan Cell Cycle. PLOS Biol [Internet]. 2015 Mar 3 [cited

2022 Nov 4];13(3):e1002093. Available from:

https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.1002093

51.   Hu K. Organizational changes of the daughter basal complex during the parasite

replication of Toxoplasma gondii. PLoS Pathog. 2008;4(1):0108–21.

52.   Vaishnava S, Striepen B. The cell biology of secondary endosymbiosis – how

parasites build, divide and segregate the apicoplast. Mol Microbiol [Internet]. 2006

Sep 1 [cited 2023 Jun 17];61(6):1380–7. Available from:

https://onlinelibrary.wiley.com/doi/full/10.1111/j.1365-2958.2006.05343.x

53.   Cao S, Yang J, Fu J, Chen H, Jia H. The dissection of SNAREs reveals key

factors for vesicular trafficking to the endosome-like compartment and apicoplast

via the secretory system in toxoplasma gondii. MBio [Internet]. 2021 Aug 1 [cited

2022 Nov 4];12(4). Available from:

https://journals.asm.org/doi/10.1128/mBio.01380-21

54.   Fu J, Zhao L, Pang Y, Chen H, Yamamoto H, Chen Y, et al. Apicoplast

biogenesis mediated by ATG8 requires the ATG12–ATG5-ATG16L and SNAP29

complexes in Toxoplasma gondii. https://doi.org/101080/1554862720222123639

[Internet]. 2022 [cited 2023 Jun 17]; Available from:

https://www.tandfonline.com/doi/abs/10.1080/15548627.2022.2123639

55.   Schnurr JA, Shockey JM, De Boer GJ, Browse JA. Fatty Acid Export from the

Chloroplast. Molecular Characterization of a Major Plastidial Acyl-Coenzyme A

Synthetase from Arabidopsis. Plant Physiol [Internet]. 2002 Aug 1 [cited 2023 Jun

17];129(4):1700–9. Available from: https://dx.doi.org/10.1104/pp.003251

 Devarakonda et al – Figure Legends

Figure 1. TgAPT1 vesicle trafficking occurs in the elongation phase of apicoplast division

(A) Schematic of the apicoplast division cycle in *T. gondii.* During interphase the apicoplast is

circular and located at the parasite's apical end. At the start of the cell division, apicoplast

undergoes elongation and associates with a pair of duplicated centrosomes. Centrosome

movement into daughter parasites, results in a U-shaped apicoplast. Membrane fission is driven

by TgDrpA and results in a single apicoplast being inherited by each daughter cell. Inset:

Depiction of apicoplast morphology. The apicoplast contains 4 membranes: outermost,

periplastid (green) and two innermost membrane (black). TgAPT1 localizes to the apicoplast

membranes. Cpn60 localizes to the apicoplast lumen.

915
B) Maximum intensity projection of deconvolved epifluorescent images of the parasites in

interphase, elongation and post-fission stages of the apicoplast division cycle. Parasites are

expressing TgAPT1-EmGFP (green) and mCherry-TubulinA1 (a marker for mother and

daughter parasite pellicle shown in magenta). Parasites were fixed and stained with an anti-

Cpn60 antibody (a marker for the apicoplast lumen shown in yellow) and DAPI (cyan). Scale bar

represents 5 μm. Inset: Brightness and contrast of insets was adjusted to facilitate visualization

of dimmer TgAPT1 vesicles (white arrows). Inset scale bar represents 1 μm.

(C) Quantification of the number of TgAPT1-EmGFP vesicles per parasite at interphase,

elongation, and post-fission from three independent experiments (N=60 parasites). The mean

from each experiment is indicated with large circles. These values were used to calculate the

mean, standard error of the mean and p-value. Raw data is shown with smaller colored circles.

Experiment 1 in orange, experiment 2 in grey, experiment 3 in blue. Statistical significance was

determined by unpaired Student's *t* test.

(D) Pulse-chase assay to track APT1 protein synthesis. *Upper:* Assay outline. Parasites

expressing APT1-Halo were labeled with JF646 for 15 minutes (pulse). After a 6 hour out-

growth the parasites were labeled with Halo-TMR for 15 minutes (chase). *Lower:* Maximum

intensity projections of deconvolved images of Halo-APT1 expressing parasites after Halo

pulse-chase assay. No TMR labeling was observed in parasites in interphase, indicated by

circular apicoplasts, demonstrating that APT1 is not expressed in interphase parasites. During

apicoplast elongation, TMR-Halo staining was observed in the ER and nuclear envelope (NE)

and in the apicoplast. Green arrow heads indicate accumulations of APT1-Halo/TMR in the ER

at the apical and basal sides of the nucleus. Images are maximum intensity projections of

deconvolved images. Scale bar represents 5 μm.

Figure 2. Actin depolymerization and TgMyoF knockdown results in an accumulation of

APT1 vesicles

(A) TgAPT1-EmGFP parasites treated with DMSO (control) or 0.2 μM cytochalasin D for 12

hours were categorized as either interphase, elongation or post-fission based on the stage of

the apicoplast division cycle and the size and presence of daughter parasites. Magenta arrow

indicates parasite that does not contain an apicoplast. White box indicates area used to make

inset in panel B. Images are maximum intensity projections of deconvolved images. Scale bar

represents 5 μm.

(B) Inset panels from (A). Brightness and contrast of insets was adjusted to facilitate

visualization of dimmer TgAPT1 vesicles. Inset scale bar represents 1 μm.

(C) Quantification of the number of TgAPT1-EmGFP vesicles per parasite during interphase,

elongation and post-fission stages of the apicoplast division cycle. Parasites with no apicoplast

were categorized separately. DMSO: n= 42/apicoplast division stage, n= 126 total; CD: n= 30-

42/apicoplast division stage, n = 156 total. TgAPT1 vesicle number per parasite are as follows

(Mean±SEM): DMSO - interphase: 1.05±0.1; elongation: 8.26±0.73; and post-fission: 4.19±0.39.

CD interphase: 3.59±0.15; elongation: 13.30±0.46; post-fission: 6.83±0.35; no apicoplast:

19±0.46). The mean from each experiment is indicated with large circles. These values were

used to calculate the mean, standard error of the mean and p-value. Raw data is shown with

smaller colored circles. Experiment 1 in orange, experiment 2 in grey, experiment 3 in blue.

Statistical significance was determined by unpaired Student's *t* test.

(D) TgMyoF-mAID:TgAPT1-EmGFP parasites treated with ethanol (control) or 500 µM IAA for

15 hours were categorized as either interphase, elongation or post-fission based on the stage of

the apicoplast division cycle and the size and presence of daughter parasites. Magenta arrow

indicates parasite that does not contain an apicoplast. White box indicates area used to make

inset in panel E. Images are maximum intensity projections of deconvolved images. Scale bar

represents 5 µm.

(E) Inset panels from (D). Brightness and contrast of insets was adjusted to facilitate

visualization of dimmer TgAPT1 vesicles. Inset scale bar represents 1 µm.

(F) Quantification of the number of TgAPT1-EmGFP vesicles per parasite during interphase,

elongation and post-fission in control and TgMyoF knockdown parasites. Parasites with no

apicoplast were categorized separately. (EtOH: n= 60/apicoplast division stage, n= 180 total;

IAA: n= 60/apicoplast division stage, n = 240 total). TgAPT1 vesicle number per parasite are as

follows (Mean±SEM): Ethanol - interphase: 0.8±0.14; elongation: 7.57±0.27; and post-fission:

1.83±0.16. IAA interphase: 9.9±0.52; elongation: 10.7±0.43; post-fission: 6.6±0.24; no

apicoplast = 17.1±0.58). The mean from each experiment is indicated with large circles. These

values were used to calculate the mean, standard error of the mean and p-value. Raw data is

shown with smaller colored circles. Experiment 1 in orange, experiment 2 in grey, experiment 3

in blue. Statistical significance was determined by unpaired Student's *t* test.

Figure 3. APT1 vesicles exhibit predominately diffusive motion

(A) Localization TgAPT1-EmGFP parasites in live parasites. TgAPT1 is localized to both the

apicoplast and vesicles. White boxes indicate areas used to make insets 1 and 2. Inset 1

depicts TgAPT1+ vesicle moving in a directed manner. Inset 2 depicts a TgAPT1+ vesicle

moving with diffusive-like movement. Scale bar = 5µM. Inset scale bar = 1 µM.

(B) The Z-stack movie of TgAPT1-EmGFP expressing parasites were imported into Imaris. Raw

data from frame 1 of the movie is shown. Isosurfaces of the apicoplast were reconstructed to

prevent aberrant tracking inside the apicoplast itself. Vesicle tracks observed in the first 20

seconds of imaging are color-coded by straightness value.

(C) Frequency distribution of trajectory straightness values of TgAPT1 vesicles. Straightness

value close to zero indicates random motion. Straightness value close to 1 indicates

directed/linear movement. Mean straightness value is 0.18±0.15.

(D) Frequency distribution of displacement lengths of TgAPT1+ vesicles. Vesicles with a

straightness value less than 0.5 (black) had a mean displacement length of 0.3±0.27µm

(Mean±SD), while vesicles with a straightness value greater than 0.5 (orange) had a mean

displacement length of 0.59±0.3µm (Mean±SD).

(E) Localization TgAPT1-EmGFP parasites in live parasites treated with DMSO or CD for 30

minutes prior to imaging. Raw data from frame 1 of the movie is shown. Z-stack movies were

imported into Imaris, isosurfaces of the apicoplast and vesicles were reconstructed. Vesicle

tracks observed in the first 20 seconds of imaging are color-coded by straightness value.

(F) Localization TgAPT1-EmGFP parasites in live parasites treated with ethanol or IAA for 15

hours prior to imaging. Raw data from frame 1 of the movie is shown. Z-stack movies were

imported into Imaris, isosurfaces of the apicoplast and vesicles were reconstructed. Vesicle

tracks observed in the first 20 seconds of imaging are color-coded by straightness value.

(G) Frequency distribution of trajectory straightness values of TgAPT1 vesicles from DMSO

treated (black) and cytochalasin D (CD) treated (blue) parasites. Mean straightness values are

for DMSO and CD treated are 0.19±0.18 and 0.14±0.13 (Mean±SD) respectively. Straightness

value close to zero indicates random motion. Straightness value close to 1 indicates directed

movement.

(H) Frequency distribution of trajectory straightness values of APT1 vesicles from ethanol

(black) and IAA (green) treated parasites. Mean straightness values are for ethanol and IAA

treated are 0.2±0.18 and 0.17±0.17 (Mean±SD) respectively. Straightness value close to zero

indicates random motion. Straightness value close to 1 indicates directed movement.

(I) Bar graph indicating the percentage of vesicles exhibiting directed motion, defined as a

straightness value greater than 0.5 and a track displacement length of greater than 500nm.

Actin depolymerization with CD or TgMyoF knockdown with IAA results in a significant (~50%)

decrease in the percentage of directed trajectories.

Figure 4. TgMyoF knockdown results in reduced apicoplast length

(A) TgMyoF-mAID parasites expressing eGFP-tubulin (magenta) were grown for 12 hours before treatment with ethanol (control) or IAA (TgMyoF KD) for 6 hours. Parasites were stained with an anti-Cpn60 antibody (yellow) and DAPI (cyan). Scale bar represents 5 μm.

(B) Quantification of the apicoplast length in control and TgMyoF KD parasites from 3 independent experiments (20 parasites per experiment). Mean apicoplast length was 2.12±0.05 and 1.55±0.05 μm for ethanol and IAA treated parasites respectively. The mean from each experiment is indicated with large circles. These values were used to calculate the mean, SEM and p-value. Raw data is shown with smaller colored circles. Experiment 1 in orange, experiment 2 in grey, experiment 3 in blue. Statistical significance was determined by unpaired Student's *t* test.

Figure 5. Apicoplast dynamics require actin and TgMyoF

(A) Time lapse images of TgAPT1-EmGFP parasites during the elongation phase of apicoplast division. *Upper:* The apicoplast is dynamic and exhibits branch formation and movement over the ~2-minute imaging period. *Lower:* An example of a static elongated apicoplast. Apicoplast does not change shape over the imaging period. White box indicates area used to make time-lapse inset. Inset indicates change in apicoplast morphology over time. Time in seconds is indicated in the top right corner of each image. Scale bar represents 5 μm. Inset scale bar = 1 μm. Lower:

(B) Quantification of apicoplast dynamics. Apicoplast was categorized as dynamic if one of the following was observed: dramatic changes in apicoplast shape such as bending, branch formation, branch movement or retraction. Apicoplast was categorized as static of there were subtle or no changes in apicoplast shape or position. N=70 from four independent experiments. 63±6.2% of parasites contained dynamic apicoplasts. 37±6.2 contained static apicoplasts. Data

is represented as Mean ± SEM. Statistical significance was determined by unpaired Student's *t*

test.

(C) Time lapse images of TgAPT1-EmGFP parasites during the elongation phase of apicoplast

division imaged after treatment with DMSO or cytochalasin D. White box indicates area used to

make time-lapse inset. Inset indicates change in apicoplast morphology over time. Time in

seconds is indicated in the top right corner of each image. Scale bar represents 5 µm. Inset

scale bar = 1 µm.

(D) Quantification of apicoplast dynamics after DMSO or CD treatment. 84±2.9% of DMSO

treated parasites (grey) contained dynamic apicoplasts. 9±6.7% of CD treated parasites (blue)

contained dynamic apicoplasts. Data is represented as Mean ± SEM. N= 32 and 41 for DMSO

and CD respectively from three independent experiments. Statistical significance was

determined by unpaired Student's *t* test.

(E) Time lapse images of TgMyoF-mAID:TgAPT1-EmGFP parasites during the elongation

phase of apicoplast division imaged after treatment with ethanol or IAA. White box indicates

area used to make time-lapse inset. Inset indicates change in apicoplast morphology over time.

Time in seconds is indicated in the top right corner of each image. Scale bar represents 5 µm.

Inset scale bar = 1 µm.

(F) Quantification of apicoplast dynamics after ethanol or IAA treatment. 86±4% of ethanol

treated parasites (grey) contained dynamic apicoplasts. 22±13% of IAA treated parasites

(green) contained dynamic apicoplasts. Data is represented as Mean ± SEM. N = 38 and 40 for

ethanol and IAA respectively. Statistical significance was determined by unpaired Student's *t*

test.

Figure 6: Centrosome-apicoplast interaction is maintained during apicoplast dynamics

(A) Maximum intensity projection of a deconvolved images of parasites expressing eGFP-centrin1 (centrosome marker; magenta) and FNR-RFP (apicoplast marker; green) during the elongation phase of apicoplast division. Parasites were imaged on 5 focal planes for a total of 10 minutes. Z-slice step size was 0.4 μm. Dashed boxes indicate areas used to make time lapse insets. Inset 1: The apicoplast is attached to a single centrosome and slides in relation to the static centrosome, associating with both the tips and sides of the apicoplast. Inset 2: The same apicoplast at a later time point undergoes dramatic morphological change, extending to ~3 μm in length towards the second centrosome before collapsing backward. Inset 3: Two centrosomes associated with the sides of the apicoplast. Apicoplast "branching" (magenta arrow) is observed at the site of centrosome association. Imaging time points in minutes are indicated in the top right-hand corner of each image. Scale bar represents 5 μm, inset scale bar represents 2 μm.

(B) Interphase (upper) and post-fission (lower) apicoplasts did not exhibit morphological changes throughout the course of imaging. Imaging time points in minutes are indicated in the top right-hand corner of each image. Scale bar represents 2 μm.

Figure 7. TgMyoF knockdown disrupts the apicoplast and centrosome interaction

(A) TgMyoF-mAID parasites expressing eGFP-Centrin1 (Green) were grown for 18h before treatment with either ethanol or IAA for 6h. Parasites were fixed and stained with an anti-Cpn60 (magenta) antibody and DAPI (grey). Images are maximum intensity projections of deconvolved images. Based on the length of the apicoplast control parasites were categorized as in either early or late elongation. Since the apicoplast elongation length was reduced (Fig. 4) in TgMyoF-KD parasites, these parasites could not be categorized into early or late stages of elongation.

White box in indicates area used to make inset. Inset: White dashed line and white number

indicate apicoplast length. Green dashed line and green number indicate the distance between

the apicoplast and the centrosome. Scale bar represents 5 µm. Inset scale bar represents 2

µm.

(B) Distance between the centrosome and apicoplast were plotted as a function of apicoplast

length in control and MyoF-KD parasites. Experiment 1 in orange, experiment 2 in grey,

experiment 3 in blue. In control parasites, the apicoplast-centrosome distance is short when

apicoplast length is above 1.5µm.

(C) Parasites were categorized has having an apicoplast greater or less than 1.5µm and the

mean apicoplast-centrosome distance was calculated for each category in control and TgMyoF-

KD parasites. In control parasites, the apicoplast centrosome distance is significantly shorter in

parasites containing a long apicoplast (>1.5µm) compared with control parasites containing a

short apicoplast (<1.5µm). In TgMyoF-KD parasites the mean apicoplast-centrosome distance

was ~1µm, regardless of apicoplast length.

Figure 8: Apicoplast inheritance defects are not due to changes in Atg8 localization.

1098

(A) TgMyoF-AID parasites expressing eGFP-tubulin (magenta) were treated with ethanol or IAA

for 6 hours then fixed and stained with an anti-Cpn60 antibody (yellow). Control parasites show

that the elongated apicoplasts are in proximity to the respective daughter parasites (white

arrowheads). Later in division as indicated by larger daughter parasites, the tips of the

elongated apicoplast are localized within the growing daughters (magenta arrowheads). Upon

TgMyoF knockdown, there is an asymmetric localization of the elongated apicoplast to one of

the daughter cells (white arrow) and absent from second daughter parasite (asterisks). Scale

bar represents 5 μm.

(B) Quantification of the percentage of daughter parasites in close proximity to the elongated

apicoplast in control parasites and TgMyoF KD parasites. Parasites were categorized as

follows: 2D – elongated apicoplast near or within both daughter parasites. 1D - elongated

apicoplast in close proximity to one of the daughter parasites, and "none" - elongated apicoplast

not in close proximity to either of the daughter parasites. In control parasites, 100% of

apicoplast were in close proximity to both daughter parasites compared to 13.33±6% of

TgMyoF-KD parasites. In TgMyoF-KD parasites, 70±2.9% and 16.7±4.4% of apicoplasts were

associated with one daughter parasite or neither daughter parasite respectively (Mean±SEM).

Bar graph shows quantification of combined data from 3 independent experiments (20 parasites

per experiment). Statistical significance was determined by unpaired Student's *t* test.

(C) TgMyoF-mAID parasites ectopically expressing 3xTy-Atg8 under the Atg8 promoter were

grown for 12 hours before treatment with ethanol (control) or IAA (TgMyoF KD) for 6h. Cells

were stained with anti-Ty (magenta) and anti-Cpn60 antibodies (yellow). Maximum intensity

projections of deconvolved images of parasites at interphase, elongation and post-fission show

that loss of TgMyoF does not affect Atg8 localization. Scale bar represents 5 μm

Figure 9. Illustration of the role of actin and TgMyoF in apicoplast division

In interphase parasites, *T. gondii* contains a single apicoplast and centrosome. At the onset of

apicoplast division, the apicoplast elongates in a manner independent of centrosome

association and there is an increase in the number TgAPT1+ vesicles in the cytosol, which

could supply the apicoplast with additional proteins and lipids prior to duplication. In the

elongation phase, the apicoplast is highly dynamic exhibiting bending and branch formation. At

this stage, the centrosome is dynamically associated with both the sides and the tips of the

apicoplast. Association of the apicoplast tips with the centrosome facilitates inheritance of a

single apicoplast by each daughter cell upon apicoplast fission. In the absence of TgMyoF and

actin, TgAPT1+ vesicles accumulate in the cytosol. These vesicles could represent either

vesiculation/fragmentation of the apicoplast or ER-derived vesicles that have failed to fuse with

the apicoplast. Also during elongation, the apicoplast is static and the apicoplast frequently fails

to associate with one or both centrosomes. This results in the asymmetric division of the

apicoplast into daughter parasites.

Figure 1

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Figure 9.