Estimation of the expectation of species richness difference of two-subcommunities from site-suitability difference in a species-independent stochastic model

Ryosuke Iritani [*][†][‡][§]

January 1, 2023

Abstract

Understanding how community assembly in initial faunization phases determines the difference in the number of species present (richness difference) is a fundamental question in ecology. However, little prediction exists for when, and the degree to which, species richness emerges. Here, assuming that species may have different and independent presence probabilities in a two-sites system, we investigate the expectations of richness difference in a pair of subcommunities, and explore for predictive indices for richness difference. We found that (i) species differences (average among two sites of species presence abilities divided by the total presence probability) tend to have nonconsistent effects on richness difference; (ii) spatial heterogeneity (average taken across species of the differences in species presence probabilities in two sites), if strong, may have a good predictive power for the emergence of richness difference; and (iii) the absolute difference in average presence probabilities (site-suitability difference) has a robust predictive power for richness differences. The power becomes so pronounced that it provides even an accurate approximation, but this power tends to drop as site-suitability index per se becomes smaller. The finding suggests that although it is tempting and challenging to incorporate heterogeneous factors into islands biogeography theory, the average suitability of site occupancy for any species suffices to predict patterns of richness differences in two-sites. This work provides a theoretical framework for accurate prediction of species richness differences.

Key words: Heterogeneity; Nestedness; Null models; Species richness; Stochasticity.

[*]RIKEN Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS), RIKEN, Wako, Saitama, Japan;
[†]Department of Biology, Faculty of Sciences, The University of Tokyo
[‡]lambtani@gmail.com
[§]ORCID iD:0000-0002-2396-1109

1 Introduction

Understanding the differences in species richness ('richness difference') is one of the fundamental problems in ecology [1, 2]. For presence-absence data, richness difference is defined as the absolute difference between two local species richnesses. The richness difference may be caused by various assemblage processes such as distance, area, species differences in colonization ability, and spatial heterogeneity [3–6]. Because richness difference provides essential information for the spatial biodiversity, it is becoming more and more important to predict factors shaping richness difference in the course of deterministic and stochastic community assemblage processes.

A challenging problem associated with predicting richness differences stems from complex, heterogeneous nature of the real world [7–10]. In the presence of heterogeneity, there are at least three factors that may contribute to richness differences. For example, spatial structure mediated by heterogeneity in spatial locations, distance, degree of fragmentation, and colonization ability can readily result in richness differences and subsequently form richness gradient [11, 12]. Surprisingly little theory exists as to how richness differences are influenced by such factors and how one can make accurate predictions for richness differences, both limiting our understanding of richness variations in nature. Iritani *et al.* [13], by regarding species incidence (presence or absence) as a probabilistic variable [14], studied how species difference and spatial heterogeneity affect compositional dissimilarity between two subcommunities (beta-diversity), but did not consider any properties of richness differences. Similarly, based on the assumption that species incidence is independent, Calabrese *et al.* [15] and Carmona & Pärtel [16] took the probabilistic approach to estimate the species richness and its stochastic variations. However, neither did their studies consider richness difference. Taken together, the current methodological tools have not yet succeeded in providing theoretical predictions of how richness difference can be shaped by stochastic processes when there are species difference and spatial heterogeneity.

In this article, we derive analytical, exact formula for richness difference of species richness between two subcommunities, assuming that: (i) species incidence is independent across space and species, and (ii) differential values are taken among the presence probabilities of species i in subcommunity j (for $i = 1, ..., S$ and $j = 1, 2$). Then we examine three factors for their potential for good prediction of richness differences. The present theoretical work provides useful guide to predict how richness differences can occur in natural and experimental environments, especially by nullifying correlations among species incidence.

2 Method and result

2.1 General analysis

We write $p_{i,j}$ for the probability that species i is present in subcommunity $j(= 1, 2)$, and x for the binary state of species i's incidence in subcommunity j: if species i is present in subcommunity j, we write $x_{i,j} = 1$, otherwise absent $x_{i,j} = 0$. We assume that species-pool is of limited capacity, such that the number S of species in total can be present in at least one of the sites. The species richness, or alpha-diversity, in j can be written as:

$$\alpha_j(\mathbf{X}) := \sum_{i=1}^{S} x_{i,j}, \tag{1}$$

which is a nonnegative integer value, $\alpha_j = 0, 1, ..., S$. In statistics, α_j is said to follow Poisson-Binomial distribution with parameters $\mathbf{p}_j := (p_{1,j}, p_{2,j}, ..., p_{S,j})$. The expected value of alpha-diversity over all possible $\mathbf{X}$s is defined by:

$$\mathbf{E}[\alpha_j] = \sum_{i=1}^{S} p_{i,j}. \tag{2}$$

The minimum of the species richness is defined by:

$$\mu_-(\mathbf{X}) := \min\{\alpha_1(\mathbf{X}), \alpha_2(\mathbf{X})\}. \tag{3}$$

We can write down its expected value, $\mathbf{E}[\mu_-]$, in a complicated form (Appendix).

The expectation of the minimum of species richness for two subcommunities is immediately translated down to the expectation of the difference in species richness between two subcommunities. Let $D(\mathbf{X}) := |\alpha_1(\mathbf{X}) - \alpha_2(\mathbf{X})|$ be the difference of species richness between two subcommunities. Using the identity $|\alpha_1 - \alpha_2| \equiv \alpha_1 + \alpha_2 - 2\min\{\alpha_1, \alpha_2\}$ yields:

$$\mathbf{E}[D] = \mathbf{E}[\,|\alpha_1 - \alpha_2|\,] = \mathbf{E}[\alpha_1 + \alpha_2] - 2\mathbf{E}[\mu_-]. \tag{4}$$

2.2 Predictors

We use three parameters to measure heterogeneity in the presence probabilities, $p_{i,j}$. First is species difference denoted w, which is defined as the proportion of the average taken for two sites of absolute deviation of the presence probabilities from within-site average presence probability to the sum of presence probabilities for all species for two sites [13]. Second is spatial heterogeneity denoted h, which is defined as the average, taken across species, of the difference of presence probabilities for two sites. These are given by:

$$w := \frac{1}{\mu_1 + \mu_2} \sum_{i=1}^{S} \frac{1}{2} \sum_{j=1}^{2} |p_{i,j} - \mu_j|, \tag{5}$$

$$h := \frac{1}{S} \sum_{i=1}^{S} |p_{i,1} - p_{i,2}| \tag{6}$$

[13]. Finally, we propose the following quantity:

$$\delta := \frac{1}{S} \left| \sum_{i=1}^{S} (p_{i,1} - p_{i,2}) \right| = \frac{1}{S} |\mathbf{E}[\alpha_1] - \mathbf{E}[\alpha_2]| = |\mu_1 - \mu_2|, \tag{7}$$

which represents the absolute difference of alpha-diversity and describes which site is on average more suitable. Note that $\delta \leq h$ by the triangle inequality, in which the equality ($h = \delta$) occurs when one site is predominantly more suitable than the other: $p_{i,1} > p_{i,2}$ for all $i = 1, ..., S$ or either $p_{i,1} < p_{i,2}$ for all $i = 1, ..., S$.

To illustrate the dependence of richness difference against these three quantities, we carried out stochastic simulations (Monte Carlo simulations), based on the following scheme. (1) Setting $S = 16$, choose two probability distributions to generate $\mathbf{p}_j = (p_{1,j}, ..., p_{S,j})$; we refer to such distributions as 'incidence-generators' in this article. The number of simulations is 300. (2) We compute the species difference, spatial heterogeneity, site-suitability difference, and expected richness difference, and then plot scatter plots.

We first of all find that whereas neither species difference and spatial heterogeneity on their own may provide clear predictions for richness difference, the site suitability difference best explains richness differences. This prediction seems to strongly depend on the probabilistic distributions to generate the values of $p_{i,j}$. To illustrate this, we choose three sets of the uniform distributions for incidence-generators (and we write "LHS ~ Beta[range]" to mean that LHS follows the Beta distribution with parameters indicated by RHS, $[\cdot, \cdot]$).

(1) "Random model": $p_{i,1}$ ~ Beta[1.10, 1.09] and $p_{i,2}$ ~ Beta[1.09, 1.10];

(2) "Overlap model": $p_{i,1}$ ~ Beta[1.28, 0.96] and $p_{i,2}$ ~ Beta[0.96, 1.28]

(3) "Dominance model": $p_{i,1}$ ~ Beta[1.6, 0.4] and $p_{i,2}$ ~ Beta[0.4, 1.6]

For the random model, neither species difference nor spatial heterogeneity

For the overlap model, richness difference has no clear dependence on species difference. The predictive abilities of both spatial heterogeneity and site-suitability for richness difference are improved over the random pattern, but the site-suitability's predictive ability decreases with decreased site-suitability and richness difference.

Finally, the dominance model predicts that, whereas species difference tends to have a low predictive ability, spatial heterogeneity and site-suitability difference both almost perfectly (and linearly) predict richness difference; it should be however noted that this coincidence is due to the equality $\delta = h$ for the dominance model.

In sum, we examined how accurately species difference index, spatial heterogeneity index, and site-suitability

difference index can predict the expectation of richness difference. The predictive abilities of these indices strongly depend on the nature of probability distributions for species incidence. When species' presence probabilities in two sites are very similar as in the random model, it is difficult to predict the richness difference, but there seems to be a governing relation of site-suitability difference with richness difference. As the probability distributions for the two sites become dissimilar (from random, overlap, to dominance model), the predictive abilities of the site-suitability and spatial heterogeneity tend to be improved.

Figure 1: The performance of species difference (left), spatial heterogeneity (middle), and site-suitability difference (right), for predicting the expectation of richness difference, in three models producing different probability distributions of $\mathbf{p}_1$ and $\mathbf{p}_2$. In the right panels, the dotted segment is a line with slope S. Inset panels are the incidence-generators of $\mathbf{p}_1$ (orange) and $\mathbf{p}_2$ (blue). (A) When $\mathbf{p}_1$ and $\mathbf{p}_2$ are drawn from nearly indentical distributions (incidence-generator), richness difference is systematically small because the alpha diversity tends to be similar between two sites. Neither can three predictors perform well, although site-suitability difference seems to have a formal relationship with richness difference. (B) When the two incidence-generators are partially (but not fully) overlapped, the predictive ability of site-suitability difference for species richness. (C) When one site is dominantly more likely to be inhabited by species, the predictive ability of site-suitability difference for the species becomes almost perfect. Parameters: $S = 16$; the number of simulations in each of panels (A–C): 300.

3 Discussion

Understanding and predicting differences in species richness have been of great interest in community ecology. Using probabilistic approach, we investigated how the expectation of richness difference can be predicted by three indices. These indices can be estimated from long time-series data of species incidence. We found that the expectation of richness difference has a complicated, and almost intractable, analytical form. While species differences and site-heterogeneity both tend to have limited predictive ability of the expected richness difference, a site-suitability index, defined as the average of the absolute difference of alpha diversity between two sites, provides a good predictor.

However, the predictive ability of the index decreases as the site-suitability index becomes small. The dependence of richness difference on site-suitability difference seems to be positive and nonlinear (though asymptotically linear).

The degree of spatial heterogeneity can have a good predictive power for richness difference, but the predictiveness is sensitive to the probability distributions with which species are present in each site. Specifically, if one site is a priori known to be likely inhabited by predominantly more species over the other site, then spatial heterogeneity estimated from long-time series data of species presence-absence can be used as a measure of richness difference. On the other hand, without such a priori knowledge, i.e., if the average probabilities of species to be present in one site and in the other are similar, it would be difficult to deduce richness difference from site-heterogeneity measure per se. This is because site-heterogeneity measure is invariant with the permutation of sites for each species: exchanging $p_{i,1}$ and $p_{i,2}$ for each species i does not change the value of h, and thus is not designed to detect richness difference.

In addition, our key finding predicts that richness difference may be adequately predicted by the site-suitability index, all things being equal. This mathematically suggests that it does not much matter exchanging the order of taking expectation and taking absolute value. Biologically, despite the presence of heterogeneity, the absolute difference in alpha-diversity provides a good estimate for the expected absolute difference in species richness. However, the fit becomes less accurate when the site-suitability and richness difference are both low, suggesting that it is difficult to estimate the richness difference from empirical data that typically contains a limited number of unique species. This result can be reasoned as follows. When site-suitability difference is large, it is immediately clear that one site is predominantly more likely to be inhabited by various species than the other, so that richness difference becomes more pronounced. In contrast, a small value of site-suitability means that two sites are equally likely to be inhabited by species. In such a case, richness difference may emerge also merely by small-sample effect.

Our model can be used as a framework of environmentally constrained null-model analyses of richness difference [7]. Despite the intuition that incorporating species differences and spatial heterogeneity matters, the present work showed that the effects of such heterogeneity are overall small unless the absolute difference of the expectations of alpha-diversity in two sites is low. Specifically, by averaging long time-series data of incidence, e.g., [17], one may estimate presence probabilities, assuming that presence probabilities are at dynamical equilibrium [14]. In this vein, deviations of observed richness difference from the expectation of the present work provide a signal of the effect of certain nullified factors. Since we here assumed that species incidence are uncorrelated and also that two sites are uncorrelated, such deviation can be explained by these factors.

Our work has further implications for predicting beta-diversity, because richness difference is a subcomponent beta-diversity (spatial variations in species richness; [5, 18]). Specifically, Baselga [5] proposed to partition incidence-based beta-diversity measures, e.g., Jaccard or Sørensen's [19–21], into additive components of richness difference and turnoever. Iritani *et al.* [13] derived the exact formula for the expectation of Jaccard dissimilarity in the same setting as the current work's, but further methodology has yet to be developed for the additive partitioning scheme proposed by Baselga [5, 18]. Future studies may apply the methodology to the additive partitioning frameworks and thereby advance the theory of beta-diversity for the heterogeneous world.

The present study has several limitations. First, the methodology is limited to the two-sites case, and therefore a multi-site scenario should be throughly considered. Second, the species-independence assumption, by which we consider that presence probabilities of each species are uncorrelated, may be less realistic, although it provides the null-model analytical framework. Third, related to the second limitation, more realistic spatial structure should be considered. Finally, generalizing the method to abundance-based nestedness components would be a challenging problem. That beings said, these limitations rather suggest that the current methodology has a number of research avenue to make testable predictions for, and improve our understanding of, the repeated patterns of spatial biodiversity [22].

In conclusion, we derived the analytical formula for the expected richness-difference based on the usual assumption that species incidences are uncorrelated across space. We performed stochastic simulations and found that richness difference is very well explained by site-suitability, but the explanatory power is reduced when site-suitability index is small, i.e., when the variation in spatial conditions for species presence versus absence is small. Richness difference is a fundamental quantity and has the potential for future studies to produce interesting hypotheses of beta-diversity patterns.

Acknowledgement

We thank JSPS KAKENHI (grant no. 19K22457, 19K23768 and 20K15882).

References

[1] Darlington Junior, P. J. *Zoogeography: the geographical distribution of animals*. Wiley, 1957, pp. 675–675.

[2] Ulrich, W. & Almeida-Neto, M. "On the meanings of nestedness: back to the basics". *Ecography* **35**.10 (2012), pp. 865–871. DOI: `10.1111/j.1600-0587.2012.07671.x`.

[3] Carvalho, J. C., Cardoso, P., & Gomes, P. "Determining the relative roles of species replacement and species richness differences in generating beta-diversity patterns". *Global Ecology and Biogeography* **21**.7 (2011), pp. 760–771. DOI: `10.1111/j.1466-8238.2011.00694.x`.

[4] Podani, J. & Schmera, D. "A new conceptual and methodological framework for exploring and explaining pattern in presence - absence data". *Oikos* **120**.11 (2011), pp. 1625–1638. DOI: `10.1111/j.1600-0706.2011.19451.x`.

[5] Baselga, A. "Partitioning the turnover and nestedness components of beta diversity". *Global Ecology and Biogeography* **19**.1 (2010), pp. 134–143. DOI: `10.1111/j.1466-8238.2009.00490.x`.

[6] Qian, H., Ricklefs, R. E., & White, P. S. "Beta diversity of angiosperms in temperate floras of eastern Asia and eastern North America". *Ecology Letters* **8**.1 (2004), pp. 15–22. DOI: `10.1111/j.1461-0248.2004.00682.x`.

[7] Peres-Neto, P. R., Olden, J. D., & Jackson, D. A. "Environmentally constrained null models: site suitability as occupancy criterion". *Oikos* **93**.1 (2001), pp. 110–120. DOI: `10.1034/j.1600-0706.2001.930112.x`.

[8] Willig, M., Kaufman, D., & Stevens, R. "Latitudinal gradients of biodiversity: pattern, process, scale, and synthesis". *Annual Review of Ecology, Evolution, and Systematics* **34**.1 (2003), pp. 273–309. DOI: `10.1146/annurev.ecolsys.34.012103.144032`.

[9] Connolly, S. R. "Macroecological theory and the analysis of species richness gradients". *Marine macroecology*. University of Chicago Press, 2009, pp. 279–309. DOI: `10.7208/chicago/9780226904146.003.0011`.

[10] Gotelli, N. J., Anderson, M. J., Arita, H. T., Chao, A., Colwell, R. K., Connolly, S. R., Currie, D. J., Dunn, R. R., Graves, G. R., Green, J. L., Grytnes, J.-A., Jiang, Y.-H., Jetz, W., Lyons, S. K., McCain, C. M., Magurran, A. E., Rahbek, C., Rangel, T. F., Soberón, J., Webb, C. O., & Willig, M. R. "Patterns and causes of species richness: a general simulation model for macroecology". *Ecology Letters* **12**.9 (2009), pp. 873–886. DOI: `10.1111/j.1461-0248.2009.01353.x`.

[11] Gaston, K. J. "Global patterns in biodiversity". *Nature* **405**.6783 (2000), pp. 220–227. DOI: `10.1038/35012228`.

[12] Hillebrand, H. "On the Generality of the Latitudinal Diversity Gradient". *The American Naturalist* **163**.2 (2004), pp. 192–211. DOI: `10.1086/381004`.

[13] Iritani, R., Ontiveros, V. L. J., Alonso, D., Capitan, J. A., Godsoe, W., & Tatsumi, S. "Jaccard dissimilarity in stochastic community models based on the species-independence assumption" (2022). DOI: `10.1101/2022.12.13.520233`.

[14] MacArthur, R. H. & Wilson, E. O. "An equilibrium theory of insular zoogeography". *Evolution* **17**.4 (1963), p. 373. DOI: `10.2307/2407089`.

[15] Calabrese, J. M., Certain, G., Kraan, C., & Dormann, C. F. "Stacking species distribution models and adjusting bias by linking them to macroecological models". *Global Ecology and Biogeography* **23**.1 (2013), pp. 99–112. DOI: `10.1111/geb.12102`.

[16] Carmona, C. P. & Pärtel, M. "Estimating probabilistic site-specific species pools and dark diversity from co-occurrence data". *Global Ecology and Biogeography* **30**.1 (2020), pp. 316–326. DOI: `10.1111/geb.13203`.

[17] Dornelas, M., Magurran, A. E., Buckland, S. T., Chao, A., Chazdon, R. L., Colwell, R. K., Curtis, T., Gaston, K. J., Gotelli, N. J., Kosnik, M. A., McGill, B., McCune, J. L., Morlon, H., Mumby, P. J., Øvreås, L., Studeny, A., & Vellend, M. "Quantifying temporal change in biodiversity: challenges and opportunities". *Proceedings of the Royal Society B: Biological Sciences* **280**.1750 (2013), p. 20121931. DOI: `10.1098/rspb.2012.1931`.

[18] Baselga, A. "The relationship between species replacement, dissimilarity derived from nestedness, and nestedness". *Global Ecology and Biogeography* **21**.12 (2012), pp. 1223–1232. DOI: `10.1111/j.1466-8238.2011.00756.x`.

[19] Jaccard, P. "Nouvelles recherches sur la distribution florale". *Bull. Soc. Vaud. Sci. Nat.* **44** (1908), pp. 223–270.

[20] Jaccard, P. "The Distribution of the flora in the alpine zone". *New Phytologist* **11**.2 (1912), pp. 37–50. DOI: `10.1111/j.1469-8137.1912.tb05611.x`.

[21] Sørensen, T. A. "A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons". *Biol. Skar.* **5** (1948), pp. 1–34.

[22] MacArthur, R. H. *Geographical ecology: patterns in the distribution of species*. Princeton University Press, 1984.

Appendices

A Notation

· i: label for species

· S: the maximum number of species that can be present in at least one of the sites

· j: label for site

· $x_{i,j}$: incidence of species i in site j

· $\mathbf{X}$: matrix (table) with elements $x_{i,j}$

· $p_{i,j}$: the probability of species i present in site j

· $\mathbf{E}[]$: expectation over all possible $\mathbf{X}$s

· $\alpha_j(\mathbf{X})$: the species richness in site j

B Expectation of the minimum value of local species richness

In general, for any nonnegative integer-valued stochastic variable z, the following identity holds true:

$$\mathbf{E}[z] \equiv \sum_{k \geq 1} \mathbf{P}[z \geq k]. \tag{SI 1}$$

If we take $z = \min\{z_1, z_2\}$, that is, if we wish to calculate the minimum of two independent random variables, then:

$$
\begin{aligned}
\mathbf{E}\big[\min\{z_1, z_2\}\big] &= \sum_{k \geq 1} \mathbf{P}\big[\min\{z_1, z_2\} \geq k\big] \\
&= \sum_{k \geq 1} \prod_{j=1}^{2} \mathbf{P}\big[z_j \geq k\big] \\
&= \sum_{k \geq 1} \prod_{j=1}^{2} \Big(1 - \mathbf{P}\big[0 \leq z_j \leq k - 1\big]\Big).
\end{aligned}
\tag{SI 2}
$$

By taking $z_j = \alpha_j$, the formula leads to:

$$\mathbf{E}\big[\min\{\alpha_1, \alpha_2\}\big] = \sum_{k \geq 1} \prod_{j=1}^{2} \left(1 - \sum_{n=0}^{k-1} \mathbf{P}\big[\alpha_j = n\big]\right). \tag{SI 3}$$

Since α_j follows the Poisson-Binomial distribution (PBD) with parameters $\mathbf{p}_j\big(:= (p_{1,j}, ..., p_{S,j})\big)$,

$$\mathbf{P}\big[\alpha_j = n\big] = \frac{1}{S+1}\left(\prod_{i=1}^{S} p_{i,j}\right) \sum_{\ell=0}^{S} e^{-\frac{2\pi\imath n\ell}{S+1}} \prod_{m=1}^{S} \left(e^{\frac{2\pi\imath\ell}{S+1}} - \rho_{m,j}\right) \tag{SI 4}$$

[23], where $\imath$ represents the imaginary unit, and $\rho_{m,j}$ equals $-(1 - p_{m,j}) / p_{m,j}$. Therefore,

$$\mathbf{P}\big[0 \leq \alpha_j \leq k-1\big] = \sum_{n=0}^{k-1} \mathbf{P}\big[\alpha_j = n\big] = \frac{1}{S+1}\left(\prod_{i=1}^{S} p_{i,j}\right) \sum_{\ell=0}^{S} \frac{1 - e^{-\frac{2\pi\imath k\ell}{S+1}}}{1 - e^{-\frac{2\pi\imath\ell}{S+1}}} \prod_{m=1}^{S} \left(e^{\frac{2\pi\imath\ell}{S+1}} - \rho_{m,j}\right). \tag{SI 5}$$

Note that the factor with $\ell = 0$ in the summand is a removal singularity for the fraction in the summand. Inserting this into Equation (SI 2) supplies the expectation of the minimum richness. Using the identity $\alpha_1 + \alpha_2 - 2 \min\{\alpha_1, \alpha_2\} \equiv |\alpha_1 - \alpha_2|$ and taking the expectation in both sides gives the expectation of richness difference.

References

[23]   Fernandez, M. & Williams, S. "Closed-form expression for the Poisson-binomial probability density function". *IEEE Transactions on Aerospace and Electronic Systems* **46**.2 (2010), pp. 803–817. DOI: `10.1109/taes.2010.5461658`.