Functional Organization of Auditory and Reward Systems in Healthy Aging

Alexander Belden[1], Milena Aiello Quinci[1], Maiya Geddes[2], Nancy Donovan[3], Suzanne B. Hanser[4], Psyche Loui[1*]

[1] Northeastern University

[2] McGill University

[3] Brigham and Women's Hospital and Harvard Medical School

[4] Berklee College of Music

* corresponding author: p.loui@northeastern.edu

Abstract

The intrinsic organization of functional brain networks is known to change with age, and is affected by perceptual input and task conditions. Here, we compare functional activity and connectivity during music listening and rest between young (N=24) and older (N=24) adults, using whole brain regression, seed-based connectivity, and ROI-ROI connectivity analyses. We found that although auditory and reward regions follow expected patterns of activation in response to music listening across both groups, younger adults show higher within-network connectivity of auditory and reward regions as compared to older adults during both rest and music listening. We also show that this age-related difference was reduced during music listening, especially in individuals showing high self-reported musical reward. Furthermore, young adults showed higher connectivity between auditory network and medial prefrontal cortex (mPFC) that was specific to music listening and varied as a function of liking ratings. Meanwhile, older adults showed a more globally diffuse pattern of connectivity, including higher connectivity to bilateral lingual and inferior frontal gyri. Finally, connectivity between auditory and reward regions was higher during well-liked musical stimuli across all participants, and especially higher when listening to music selected by the participant. These results highlight the roles of aging and reward sensitivity on auditory and reward networks. Results may inform the design of music-based interventions in aging populations, and improve our understanding of functional network dynamics as a whole.

Keywords: music, listening, aging, fMRI, older adults, young adults, reward, auditory

Introduction

Recent work has shown that the aging process has a significant impact on functional network dynamics of the brain, primarily characterized by a loss of network connectivity (Sala-Llonch, Bartrés-Faz & Junqué, 2015; Grady et al, 2016). The default mode network (DMN), for example, has been shown to be altered in both cognitively unimpaired and cognitively impaired older adults with neuroimaging evidence of early Alzheimer's disease pathophysiologic change (Sheline et al, 2010; Buckner et al., 2005; Hafkemeijer et al, 2012). Other networks too have shown decreases in function cognitively unimpaired older adults: the reward system shows decreased sensitivity to both reward and punishment cues when compared to young adults (Geddes et al, 2018), and the auditory network shows age-related decreases in connectivity to itself and to other networks such as DMN and salience network (Onoda et al, 2012).

Given these findings, it would be of great interest to learn how age-related changes in functional network dynamics interact with the reward response to music. Music has a unique power to elicit moments of intense emotional and physiological responses (Harrison & Loui, 2014), experiences that are intrinsically rewarding for most people (Zatorre, 2015). While music listening habits are known to be linked to psychological well-being well into old age (Laukka, 2007), the relationship between aging and the reward responses to music is not well understood.

For older adults, music-based interventions are hypothesized to be more enjoyable than other, non-musical cognitive training programs (Sutcliffe et al, 2020). On the other hand, older adults show less sensitivity to reward (Eppinger, Nystrom, & Cohen, 2012), as subserved by differences in the reward system during gambling tasks (Geddes et al, 2018). Behavioral studies have also shown that sensitivity to musical reward, as assessed by self-report through the Barcelona Music Reward Questionnaire (BMRQ; Mas-Herrero et al., 2013), is lower in older

adults than in young adults (Belfi et al, 2021; Cardona et al, 2022). Addressing these gaps in our understanding of age-related differences in reward and functional network dynamics will have implications for the design of music-based interventions for healthy aging.

Functional neuroimaging of the underlying auditory and reward circuitry in different age groups may help shed light on the intrinsic and task-related differences in music reward sensitivity across the lifespan. During music listening, regions involved with auditory processing, including superior temporal gyrus (STG) and Heschl's gyrus (HG) (Koelsch et al, 2005; Ellis et al, 2013), have been shown to coactivate with multiple areas, including lateral prefrontal networks important for auditory-motor and linguistic processes as well as reward associated brain structures including the medial prefrontal cortex (mPFC) and the dorsal and ventral striatum (Ferreri et al, 2019; Gold et al, 2019; Salimpoor et al, 2011; 2013). The STG receives input from primary auditory regions (Yeterian & Pandya, 1998; Kaas & Hackett, 2000). It is structurally and functionally connected to the IFG and anterior insula (Loui et al, 2009; Frey et al, 2008), the latter of which also plays a role in reward system processing (Sachs et al, 2016; Ai et al, 2022). The STG also shows structural and functional connectivity to the nucleus accumbens (Loui et al, 2017; Martinez-Molina et al, 2016). The latter connects to regions including the mPFC, caudate, and substantia nigra, together constituting the reward system (Karlsgodt et al, 2015). Auditory connections to the reward system are hypothesized to encode hedonic responses to music (Zatorre, 2015; Belfi & Loui, 2020).

Despite our understanding of the reward system, much is still not understood about the driving factor behind state-level as well as trait-level differences in musical reward processing. The case of musical anhedonia represents an extreme lack of sensitivity to musical reward: neuroimaging studies have reported individual differences in reward sensitivity to music listening being related to differences in white matter connectivity (Sachs et al, 2016; Loui et al, 2017; Martinez-Molina et al, 2019) and engagement of dopamine systems, specifically in the striatum

(Salimpoor et al., 2011; Zatorre, 2015). Music listening habits also vary by personality traits, with different musical preferences mapping onto specific personality traits (Rentfrow et al, 2011). Even within the same individual, differences in musical reward processing can arise based on the listener's familiarity with the piece, autobiographical memories related to the piece, and one's overall appreciation for the intentionality of the performer or composer (Alluri et al, 2017; Aydogan et al, 2018; Janata, 2009). Agency in music listening also impacts the processing of musical information, with self-selection of musical stimuli leading to more active engagement for listeners (Quinci et al, 2022), possibly due to predictive processes that can be more strongly engaged when listening to self-selected music (Vuust et al, 2022).

Here, we compare the functional connectivity of auditory and reward networks in a sample of older adults (OA; N = 24; aged 54-89) and young adults (YA; N = 24; aged 18-23) in intrinsic functional connectivity, as captured using resting state fMRI, and during music listening as captured by fMRI during a music listening task. Our hypotheses pertaining to these data are threefold. First, we expect well-liked music to elicit greater activity from auditory processing and reward associated brain regions. We expect this to be true of both young and older adults, and for differences in activity to arise between the two age groups. Second, we expect music listening to increase functional connectivity among and between auditory processing and reward associated brain regions as compared to rest. We expect this to be true of both young adults and older adults, and that these connections will be particularly strong while listening to well-liked musical stimuli. Third and finally, we expect that connectivity among and between auditory and reward associated regions will vary with age. While we expect this effect to be present during both rest and music listening, we expect that music listening will change connectivity of auditory and reward-associated regions in a way that will change with age, as shown by an age x task interaction effect.

We test these hypotheses through a combination of whole-brain linear regression, seed-based connectivity, and region-of-interest-based (ROI-to-ROI) connectivity approaches. Whole-

brain linear regression is used to test our first hypothesis, providing a metric for general metabolic activity during music listening. Seed-based and ROI-to-ROI connectivity measures are used to test our second and third hypotheses. While seed-based connectivity is intended as a more global method for determining functional connectivity, determining how auditory and reward networks (taken as a whole) may connect to the rest of the brain, ROI-to-ROI connectivity is intended as a more hypothesis-driven approach to determining the connectivity among and between known hubs of the auditory and reward networks.

Materials and Methods

Participants

We recruited equal samples of older and younger adults guided by median sample sizes of neuroimaging studies published in top neuroimaging journals in 2017 (23 participants) and 2018 (24 participants) (Szucs and Ioannidis, 2020). Twenty-four young adults (20 females, 3 males, 1 non-binary) between the ages of 18 and 23 (M=18.58, SD=1.21) were recruited through the Northeastern University student population. They had to meet the following inclusion criteria: 1) were 18 years of age or older, 2) had normal hearing, and 3) and pass an MRI screening. Twenty-eight older adults were recruited as part of a longitudinal study (Quinci et al., 2022) and met inclusion criteria to participate. Four were unable to complete the MRI portion of the study, resulting in a final sample of twenty-four older adults (13 males and 11 females) between the ages of 54 and 89 (*M*=66.67, *SD*=7.79). Recruitment for the older adults took place through community outreach as well as online recruitment engines including craigslist.org and BuildClinical.com. Participants were included if they 1) were at least 50 years old, 2) had normal hearing, and 3) passed MRI screening. Participants were excluded if they 1) changed medications within 6 weeks of screening; 2) had a history of psychotic or schizophrenic episodes, major neurological diagnosis, or other medical condition that might impair cognition;

3) had a history of chemotherapy within the past 10 years; or 4) experienced serious physical trauma or were diagnosed with a serious chronic health condition requiring medical treatment and monitoring within 3 months of screening. Participants were compensated for their time through either payment or course credit. This study was approved by the Northeastern University Institutional Review Board. This study is preregistered at https://osf.io/zxd42.

Procedure

All participants completed a pre-screening phone call before being enrolled to ensure they met the inclusion/exclusion criteria described above. All participants were also asked to provide the researcher with the names of six of their favorite songs.

Young adults were recruited from Northeastern University for a single-session study in return for course credit. Older adults were recruited as part of a longitudinal music-based intervention study that included an MRI, blood draw, battery of neuropsychological tests, and series of surveys (as described separately in Quinci et al., 2022). For the present analyses, we only used pre-intervention fMRI data from the older adults. All participants filled out an online version of the Barcelona Music Reward Questionnaire (BMRQ, Mas-Herrero et al, 2013) to assess individual differences in reward sensitivity. The BMRQ is a twenty-item questionnaire in which participants rate on a five-point scale the extent to which they agree with statements indicating their degree of musical reward across five dimensions: musical seeking, emotion evocation, mood regulation, sensory-motor, and social reward.

Special Considerations due to COVID-19

For the young adults, data collection occurred exclusively during the COVID-19 pandemic, between October 2021 and March 2022. Data collection for older adults began in

July 2019 and finished in November 2021, with nine participants completing their visits during the pandemic.

For the collection of human subject data during the COVID-19 pandemic, a Resumption of Research Plan (RRP) was developed and approved in consultation with the Institutional Review Board (IRB) of Northeastern University. Participants were called one day before their scheduled session and screened for COVID-19 symptoms. If they were symptom free, they were invited to come into the lab the next day. When arriving in the lab, they were asked the same COVID-19 screening questions and their temperatures were taken. If participants answered no to all of the questions and had a temperature reading of less than 100.4° F, they were allowed to stay to complete their in-person session. Participants and researchers wore personal protective equipment (PPE: surgical masks, face shields, lab coats, and surgical gloves) in accordance with Northeastern regulations, which changed over the course of data collection. Social distancing was enforced as much as possible during in-person sessions. Lab surfaces were disinfected before and after every study visit. In accordance with our RRP, neuropsychological testing or consenting was occasionally moved remotely using a HIPAA-compliant version of Zoom to minimize the amount of time participants spent in the lab for in-person testing.

fMRI Task

The fMRI task consisted of 24 trials altogether. In each trial, participants were first presented with a musical stimulus (lasting 20 seconds), then they were given the task of rating how familiar they found the music to be (familiarity rating lasted 2 seconds), and how much they liked the music (liking rating also lasted 2 seconds). Musical stimuli for the MRI task consisted of 24 different audio excerpts, each of which was 20 seconds in duration. Each audio stimulus was from one of the following three categories: participant self-selected music (6/24 stimuli),

researcher-selected music including well-known excerpts spanning multiple musical genres (10/24 stimuli) and novel music spanning multiple genres (8/24 stimuli). Stimuli were presented in a randomized order, and participants made ratings of familiarity and liking on the scales of 1 to 4: for familiarity: 1=very unfamiliar, 2=unfamiliar, 3=familiar, 4=very familiar; for liking: 1=hate, 2=neutral, 3=like, 4=love. Participants made these ratings by pressing a corresponding button on a button-box (Cambridge Research Systems) inside the scanner. Participants wore MR-compatible over-the-ear headphones (Cambridge Research Systems) over musician-grade silicone ear plugs during MRI data acquisition. The spatial mapping between buttons and the numerical categories of ratings were counterbalanced between subjects to reduce any systematic association between familiarity or liking and the motor activity resulting from making responses.

Behavioral Data Analysis

The proportion of each liking rating (hate, neutral, like, love) was averaged across participants for each group (young adults, older adults). A chi-square test was run on the 2 x 4 matrix relating the frequency of each liking rating (hate, neutral, like, love) for each group (young adults, older adults).

fMRI Data Acquisition

Images were acquired using a Siemens Magnetom 3T MR scanner with a 64-channel head coil at Northeastern University. For task fMRI data, continuous acquisition was used for 1440 volumes with a fast TR of 475 ms, for a total acquisition time of 11.4 minutes. Forty-eight axial slices (slice thickness = 3 mm, anterior to posterior, z volume = 14.4 mm) were acquired as echo-planar imaging (EPI) functional volumes covering the whole brain (TR = 475 ms, TE = 30 ms, flip angle = 60°, FOV = 240mm, voxel size = 3 x 3 x 3 mm^3). The resting state scan followed the same parameters and included 947 continuous scans, for a total scan length of

approximately 7 and a half minutes. T1 images were also acquired using a MPRAGE sequence, with one T1 image acquired every 2400 ms, for a total task time of approximately 7 minutes. Sagittal slices (0.8 mm thick, anterior to posterior) were acquired covering the whole brain (TR = 2400 ms, TE = 2.55 ms, flip angle = 8°, FOV= 256, voxel size = 0.8 x 0.8 x 0.8 mm^3) (as described in Quinci et al., 2022).

fMRI Data Analysis

Pre-processing

Task and resting state fMRI data were preprocessed using the Statistical Parametric Mapping 12 (SPM12) software (Penny et al, 2011) with the CONN Toolbox (Whitfield-Gabrieli & Nieto-Castanon, 2012). Preprocessing steps included functional realignment and unwarp, functional centering, functional slice time correction, functional outlier detection using the artifact detection tool, functional direct segmentation and normalization to MNI template, structural centering, structural segmentation and normalization to MNI template, and functional smoothing to an 8mm gaussian kernel (Friston et al., 1995). Denoising steps for seed-based connectivity analyses included white matter and cerebrospinal fluid confound correction (Behzadi et al, 2007), motion correction, global signal regression, and bandpass filtering to 0.008-0.09 Hz.

Univariate Whole-Brain Analyses

Behavioral responses from the fMRI task were imported into R Studio, and onset values were extracted for each trial based on the liking rating participants awarded each stimulus (hate, neutral, like, love). For each participant, data were converted from 4D to 3D images, resulting in 1440 scans. The model was specified using the following criteria: interscan interval = 0.475 seconds, microtime resolution = 16, microtime onset = 8, duration = 42 as to only include data from when the participant was listening to music. First-level main effects of each liking rating

were obtained using the SPM12 toolbox. Second-level main effects were analyzed for each group (young and older adults) for each liking rating (hate, neutral, like, love) using a one-sample t-test. Between-group contrasts (young vs. older adults) were also run for each liking rating. Results were corrected at the voxel and cluster threshold of $p < 0.05$ (FDR-corrected). Second-level results were visualized using the CONN Toolbox (Whitfield-Gabrieli, & Nieto-Castanon, 2012).

ROI-ROI Analysis

ROI-ROI analyses were performed using 18 auditory network ROIs and 18 reward network ROIs derived from the CONN default atlas. The auditory network seed consisted of all subdivisions of bilateral superior temporal gyrus (STG), middle temporal gyrus (MTG), inferior temporal gyrus (ITG), and Heschl's gyrus, and the reward network seed consisted of anterior cingulate, posterior cingulate, and bilateral insular cortex, frontal orbital cortex, caudate, putamen, pallidum, hippocampus, amygdala, and nucleus accumbens, as defined by Wang et al., 2020. Preprocessed ROI timeseries data were extracted from participants' first-level SPM.mat files using the Marsbar Toolbox (Brett et al., 2002). Values associated with each liking rating (hate, neutral, like, love) were separated for each participant and averaged across all stimuli with that particular rating. For each liking rating, ROI time-series data were correlated across all auditory regions (auditory-auditory), across all reward regions (reward-reward), and between auditory and reward regions (auditory-reward).

Follow-up analyses of ROI-ROI connectivity included a repeated-measures MANCOVA for participants who used all four liking ratings during the music listening task (N = 20 OA, 18 YA), comparing groups in auditory-auditory, auditory-reward, reward-reward, auditory-mPFC, auditory-PCC, reward-mPFC, and reward-PCC connectivity across our four levels of musical liking. Covariates included gender, self-reported musical reward, and within-group age.

Seed-Based Connectivity Analysis

Seed-based connectivity analysis was performed using the CONN Toolbox (Whitfield-Gabrieli, & Nieto-Castanon, 2012). Above defined auditory and reward network ROIs were combined, resulting in one seed for each network. For both seeds, contrasts were drawn between task (Task>Rest), age groups (YA>OA), and interactions between age and task. Further contrasts determined linear and quadratic effects of music liking (as determined by in-scanner ratings), as well as liking x age interactions. Finally, to rule out any confounding effects of within-group age differences, gender, and musical reward as measured by the BMRQ, direct subject effect contrasts of these metrics were performed for each condition and seed effect. Group, task, and liking effect contrasts were also repeated using within-group age and BMRQ as covariates of no interest, and relevant differences to the main contrasts will be presented when applicable.

Results

Behavioral Results:Barcelona Music Reward Questionnaire

The Barcelona Music Reward Questionnarie (BMRQ) was higher in young adults (M = 83.2, SD = 7.28) than in older adults (M = 72.3, SD = 9.97), as confirmed by an independent samples t-test (t(46) = 4.34, $p < 0.001$), indicating that young adults experience greater self-reported music reward, replicating previous reports (Mas-Herrero et al, 2013; Belfi et al, 2020; Cardona et al, 2022).

Frequency of Liking Ratings per Group

While both groups used all liking ratings, older adults used more extreme ratings: a Chi-squared test showed a significant association between group (young adults vs older

adults) and proportion of liking ratings for $X^2(3) = 12.7$, *p* = 0.005. Table 1 shows the proportion of each liking rating given by young and older adults respectively.

<insert Table 1 here>

Univariate Whole-Brain Results

For both young and older adults, music rated as hated, neutral, liked, and loved all activated auditory regions including the Heschl's gyrus, planum temporale, planum polare, STS, STG, medial temporal gyrus (MTG), and the hippocampus. For young adults, music rated as hated and liked also activated the parahippocampal gyrus, and music rated as liked additionally activated the brainstem. When listening to loved music, several additional regions became active in both groups: the medial prefrontal cortex (mPFC), paracingulate gyrus, posterior cingulate cortex, precuneus, orbitofrontal cortex and lateral occipital cortex. Also during loved music, young adults showed activity in the superior frontal gyrus, parahippocampal gyrus, brainstem, and supplementary motor area (SMA), whereas older adults showed activity in the ventral striatum/ nucleus accumbens, brainstem, and cerebellum (see Table 2 and Figure 1). Whole-brain comparisons for young vs. older adult contrasts showed no significant clusters at the p < .05 FDR-corrected level.

<Insert Figure 1 here>
<insert Table 2 here>

ROI-ROI Analysis

Results of our ROI-ROI analysis are summarized in Figure 2. Three repeated-measures ANOVAs were run to evaluate the main effects of liking ratings and age and the rating x age interaction for connectivity among auditory and reward regions and between auditory and

reward regions. We also ran linear and quadratic contrasts to further investigate the relationship between ratings and connectivity for the conditions listed above.

For connectivity among auditory regions, there was a significant main effect of liking $(F(3, 108) = 3.06$, p = 0.032, $\eta_p^2 = 0.078$). Bonferroni corrected pairwise comparisons revealed that beta-weights for stimuli rated as loved were significantly greater than those rated as neutral. There was no significant main effect of age $(F(1, 36) = 0.804$, p = 0.376, $\eta_p^2 = 0.022)$ and no significant rating x age interaction $(F(3, 108) = 0.807$, p = 0.492, $\eta_p^2 = 0.022)$. While the linear contrast was only marginally significant $(F(1, 36) = 3.42$, p = 0.073, $\eta_p^2 = 0.087)$, there was a significant quadratic contrast $(F(1, 36) = 6.95$, p = 0.012, $\eta_p^2 = 0.162)$, suggesting that hated and loved music had greater functional connectivity compared to neutral and liked music.

For connectivity among reward regions, there was a significant main effect of liking $(F(3,108) = 3.19$, p = 0.027, $\eta_p^2 = 0.081)$. Bonferroni-corrected pairwise comparisons revealed that beta-weights for stimuli rated as loved were significantly greater than those rated as neutral and those rated as liked. There was a significant main effect of age $(F(1,36) = 5.13$, p = 0.030, $\eta_p^2 = 0.125)$, suggesting that young adults have greater connectivity between reward regions compared to older adults. There was no significant rating x age interaction $(F(1,108) = 1.43$, p = 0.237, $\eta_p^2 = 0.038)$. There was no significant linear contrast $(F(1, 36) = 2.28$, p = 0.139, $\eta_p^2 = 0.060)$, but there was a significant quadratic contrast $(F(1, 36) = 8.26$, p = 0.007, $\eta_p^2 = 0.187)$. Thus, hated and loved music had greater connectivity compared to neutral and liked music.

For connectivity between auditory and reward regions, there was a significant main effect of liking $(F(3, 108) = 3.60$, p = 0.016, $\eta_p^2 = 0.091)$. Bonferroni-corrected pairwise comparisons revealed that beta-weights for stimuli rated as loved were significantly greater than those rated as neutral. There was no significant main effect of age $(F(3, 36) = 0.002$, p = 0.962,

η_p^2 <0.001) and no significant rating x age interaction (F(3, 108) = 1.49, p = 0.221, η_p^2 = 0.040).

There was both a significant linear contrast (F(1, 36) = 4.20, p = 0.048, η_p^2 = 0.104) and a

significant quadratic contrast (F(1, 36) = 7.28, p = 0.011, η_p^2 = 0.168); in other words, auditory-

reward connectivity scaled both with liking and with valence.

Follow-up MANCOVA analysis (see Supplementary Figure 4) indicated a significant

main effect of group (F(1, 36) = 4.21, p = 0.003, η_p^2 = 0.522). Also observed were quadratic, but

not linear, interactions between liking and group in auditory-mPFC (F(1, 36 = 4.31, p = 0.046, η_p^2

= 0.116) and auditory-PCC (F(1, 36) = 5.57, p = 0.024, η_p^2 = 0.144) connectivity, as well

between liking and BMRQ in auditory-auditory (F(1, 36) = 6.03, p = 0.010, η_p^2 = 0.155), auditory-

mPFC (F(1, 36) = 7.357, p = 0.011, η_p^2 = 0.182), and auditory-PCC (F(1, 36) = 4.27, p = 0.047,

η_p^2 = 0.115). Linear interactions between liking and within-group age were also observed in

auditory-auditory (F(1, 36) = 5.02, p = 0.032, η_p^2 = 0.132) and auditory-reward connectivity (F(1,

36) = 4.53, p = 0.041, η_p^2 = 0.121).

<insert Figure 2 here>

Seed-Based Connectivity

We seeded the auditory and reward networks separately and investigated the effects of

age, task, musical liking ratings, and BMRQ on functional connectivity from these seed

networks. All seed-based analyses presented were performed at the p < 0.05, FDR-corrected

level for both voxel height and cluster size. Significant clusters across all analyses are

summarized in Table 3. Statistical maps of all significant results can be found at

https://neurovault.org/collections/QSMRNLRW/

<insert Table 3 here>

Effects of Age

Contrasts between older adult and younger adults (YA>OA) revealed significant effects of auditory and reward network-seeded connectivity during both music listening and rest. Young adults showed significantly higher auditory network-seeded connectivity to other regions in the auditory network, including bilateral STG and MTG, during rest (Figure 3A-B). During music listening, these effects were limited to the right hemisphere, and young adults additionally showed higher connectivity between auditory network and mPFC, hippocampus, and anterior cingulate compared to older adults (see Figure 3C-D). Older adults showed higher connectivity to somatosensory, reward, and posterior default-mode processing regions compared to younger adults, for which effects were similar during music listening as during rest. When including BMRQ as a covariate of no interest, significant effects were more constrained (see Supplementary Figure 1). During rest, regions of higher auditory network connectivity for the YA group were limited to the right hemisphere, and for the OA group, higher connectivity in the somatosensory and frontal clusters no longer reached significance. During music listening, only higher connectivity in the precuneus for the OA group remained significant, suggesting that the other effects are explained by individual differences in BMRQ.

<insert Figure 3 here>

At rest, young adults showed significantly higher reward-network seeded connectivity to anterior cingulate (AC) while older adults showed significantly higher connectivity to posterior DMN and somatosensory regions (Figure 4A-B). During music listening, young adults once again showed connectivity to reward-associated regions, now including both AC and bilateral pallidum, while older adults now showed significant clusters of increased connectivity in bilateral IFG, in addition to effects observed at rest (Figure 4C-D). When including BMRQ as a covariate

of no interest, only the AC cluster showing higher connectivity in the YA group at rest and the IFG cluster showing higher connectivity in the OA group during music listening remained significant (see Supplementary Figure 2).

<Insert Figure 4 here>

Considering differences in age range across our two samples, all seed-based connectivity analyses were repeated using age, demeaned based on mean age for each sample, as a covariate of no interest. In all cases, there were no notable differences in the observed main effects of group or task. There were also no significant main effects of within-group age across all conditions.

Effects of Task

Auditory network-seeded connectivity showed significant differences between task and rest within both of our age groups. In young adults, connectivity was higher during music listening than rest in auditory cortex, reward, and default-mode regions, and at rest, connectivity was most notably higher than music listening in somatosensory regions and portions of the ITG (Figure 5A-B). In older adults, many of the observed effects were similar to young adults, although positive effects of music listening were generally more limited, and there was no increased connectivity observed within the mPFC (see Figure 5C-D). Reward network-seeded main effects of task were only observed in the YA group, including an increase in connectivity to bilateral lateral occipital cortex (LOC) and angular gyrus (AG) during music listening (see Figure 5E). Inclusion of BMRQ as a covariate of no interest resulted in no significant clusters for this contrast. No age by task interactions were observed.

<Insert Figure 5 here>

Musical Stimulus Effects

Effects contrasting different musical stimuli were also observed. There was significantly higher connectivity across all subjects between reward network and bilateral auditory cortex in the participant-selected music condition as compared to researcher-selected familiar western music (see Figure 6A). There was also a significant linear effect of musical liking ratings across all subjects in auditory network seed. This took the form of increased connectivity to auditory regions, including bilateral STG and MTG (see Figure 6B). Inclusion of BMRQ as a covariate of no interest resulted in additional significant clusters for this effect in the amygdala and brainstem (Supplementary Figure 3). Reward network-seeded connectivity to auditory cortex also showed a significant linear effect of liking; however this effect was limited to right lateral STG (see Figure 6C). There were no significant quadratic effects of liking observed, and no significant stimulus x group interaction effects.

Effect of BMRQ and Gender

There were no significant main effects of BMRQ or gender for either seed network in rest and music listening, and no significant BMRQ x task or gender x task interaction effects.

<Insert Figure 6>

Discussion

Here we characterize for the first time age-related differences in the neural substrates of musical reward. These data provide insight into the mechanisms underlying observed differences in musical reward in young and older adults (Belfi et al., 2020; Cardona et al, 2022; Mas-Herrero et al, 2013), characterize the contributions of auditory and reward networks during

music listening at varying levels of musical preference, and provide specific targets for the development of music-based interventions in older adult populations.

Univariate analyses showed that both young and older adults activated multiple regions within the auditory network during music listening, as well as the mPFC during music listening that was rated as loved. The engagement of mPFC when listening to preferred music is consistent with its observed role in musical reward processing (Salimpoor et al, 2013). Both groups showed specific activation of the hippocampus during preferred music listening. The hippocampus is known to be involved in both memory formation and emotion processing (Bannerman et al, 2004), and has been shown to be involved in the processing of complex emotions in voice and music (Frühholz et al, 2014).

The majority of main effects of music listening were similar between groups, suggesting that the overall activity patterns associated with music listening is preserved during aging. Nevertheless, univariate results, especially when listening to loved music, showed some clusters that were only significant in one group or another. Although a between-groups contrast was not statistically significant, main effects of music listening in each group pointed to differences when listening to loved music. Older adults showed significant precuneus activity whereas young adults showed significant SMA activity. The precuneus is a highly metabolically active area in the brain and is tied to episodic memory retrieval as well as the experience of agency (Cavanna & Trimble, 2006; Darby, Burke & Fox, 2018). The finding of larger clusters of activity in the precuneus in older adults may reflect more episodic memory processing and/or a greater sense of agency during music listening.

Only young adults showed SMA activity when listening to loved music. Since SMA is often activated in auditory imagery (Lima et al; 2016), especially during beat perception in rhythmic music (Nombela et al, 2013), this result may reflect that the experience of rhythm during music listening is more salient for young adults. Consistent with this view, neurophysiological responses comparing young and older adults during music listening have

shown that both groups are broadly sensitive to the beat in neural phase locking measures, but subtle differences persist in topographical distributions of phase locking at different levels of the beat (Tichko et al, 2022).

In ROI-ROI analyses, well-liked music resulted in higher connectivity among and between auditory and reward regions, as compared to music rated neutrally. The difference between young and older adults in functional connectivity was especially accentuated when listening to preferred music. We see that connectivity of auditory and reward regions during music listening is not a linear function in preference for the music, but rather determined by a u-shaped relationship with liking ratings. This stands in contrast to the effect of liking observed in our seed-based connectivity analysis, in which auditory network seed showed a significant linear effect, but not a quadratic effect, in connectivity to specific auditory regions. We interpret these different effects as resulting from differences in the methodologies of SBC and ROI-ROI analyses. SBC started with the auditory and reward networks taken as a whole, and thus we observed that the strongest connections arising within these networks lie within regions that show a linear effect of liking. Meanwhile, our ROI-ROI approach granted equal weight to each region identified as contributing to the auditory and reward network. In this regard, vertices showing strong linear effects of liking were accompanied by vertices that showed stronger connectivity during hated music, resulting in the observed U-shaped curve.

We also observed higher overall connectivity among reward regions in young adults as compared to older adults. This is in keeping with observed decreases in overall hedonic response with age (Eppinger, 2012), and are consistent with previously observed losses of functional network connectivity in older adults, both generally (Sala-Llonch, Bartrés-Faz & Junqué, 2015; Sheline et al, 2010) and specific to reward sensitive systems (Geddes et al, 2018).

Seed-based connectivity results further support the finding that young adults have stronger within-network seed-based connectivity than older adults. This was seen in auditory-

auditory and reward-reward connections during both rest and music listening. However, music listening and individual differences in musical reward did mediate these effects. Auditory-auditory group effects were limited to the right hemisphere during the music listening task, as well as when BMRQ was included as a covariate of no interest during rest. Research into hemispheric differences in auditory processing has long supported a right-hemispheric dominance in music processing as compared to speech (Zatorre, 1992). Musical training can also lead to a reduction in the rightward asymmetry of musical feature processing (Ono et al, 2011). As such, one possible explanation is that primary musical processing regions in the right hemisphere show a more robust effect of age, while left hemispheric differences can be accounted for by differences in task and individual differences in musical reward.

Alternatively, left auditory cortex is known to be more sensitive to rapid temporal information, whereas the right auditory cortex for processing spectral information (Zatorre & Belin, 2001; Boemio et al, 2005). These differences are also reflected in the way people listen to music, with a focus on spectral features leading to increased activity of right HG, and those focused on more holistic features showing increased left-lateralization (Schneider & Wengenroth, 2009). Therefore, another potential interpretation is that age-related differences in spectral processing systems are more robust than holistic auditory processing systems, the latter being explained by differences in musical reward and recovered by active music listening. Furthermore, inclusion of BMRQ during the music listening task removed all auditory-auditory and reward-reward effects favoring younger adults. These results suggest that individual differences in musical reward account for some of the observed differences between age groups, and that music listening transiently restores age-related deficits in auditory and reward network coherence.

Turning now to specific connectivity effects between our seed networks and other, out-of-network regions, one major finding is that music listening led to greater connectivity of mPFC from auditory network in young adults as compared to older adults. This can be observed in

both the task > rest contrast, in which the effect was seen in young adults but not older adults, and YA > OA contrast, in which the effect was seen during music listening, but not rest. This, taken together with findings from whole brain regression, suggests that while mPFC is activated during preferred music listening across age groups, the inclusion of mPFC into auditory system during music listening in general is more readily observable in young adults specifically. This is also supported by the results of follow-up MANCOVA analysis comparing the average pairwise correlations between auditory regions and mPFC, which resulted in a significant group by liking interaction. This is consistent with previous work demonstrating that functional connectivity of the default-mode network, and especially mPFC, decreases with age (Sambataro et al, 2010; Staffaroni et al, 2018), and continues to decline with age-related illnesses such as Alzheimer's disease (Hafkemeijer et al, 2015; Schouten et al, 2016; Staffaroni et al, 2018).

Another region that showed a significant interaction between music listening and age was the lateral occipital cortex (LOC). Auditory-seeded connectivity to LOC was higher in music listening than rest in both groups; however only the YA group showed increased connectivity between reward network and LOC. This result was not expected, considering LOC's primarily observed role in object perception (Malach et al, 2002; Nagy et al, 2012). However, some work has shown preferential connectivity between precuneus and LOC in trained musicians (Tanaka & Kirino, 2016) and increased activity and structural connectivity of LOC during music reading as compared to word reading (Bouhali et al, 2020), suggesting a potential role of LOC in music processing that is not yet fully understood.

Meanwhile, older adults showed greater connectivity between auditory-seeded and reward-seeded networks and out-of-network regions, including MFG, precentral gyrus, and postcentral gyrus. Older adults also showed significantly higher connectivity to precuneus, lending further support to the effect observed within the global linear regression model. For the most part, effects favoring older adults took the form of moderately positive effect sizes in the OA group compared to moderately negative or near-zero effect sizes in the YA group. This

suggests that while older adults show a decrease of network coherence, there is simultaneously a gain of more distributed connectivity between functionally distinct brain regions.

However, some of the specific effects favoring older adults do warrant further discussion. For one, older adults showed higher connectivity between reward network and bilateral IFG as compared to younger adults that was specific to the music listening task, and the right-lateral component of this effect was the only group effect within that contrast to survive correction for differences in BMRQ. IFG is known to play a role in musical syntax processing (Koelsch et al, 2005; Kunert et al, 2015), and right IFG in particular is dominant over left IFG for the perception of pitch in song (Merrill et al, 2012) and for the integration of harmonic information into auditory and motor systems (Bianco et al, 2016). The fact that this result is only observed during music listening may suggest that the IFG is specifically responding to musical stimuli, and the effect favoring older adults may reflect compensatory changes in aging, and suggest that older adults may be more efficiently using IFG to incorporate musical information into multiple systems, including the reward network.

Another region that showed increased connectivity to auditory and reward networks in older adults is the lingual gyrus (LG). LG plays a critical role in the encoding and retrieval of visual memories (Bogousslavsky et al, 1987; Machielsen et al, 2000). It has also shown increased structural and functional connectivity related to creativity (Belden et al, 2020; Takeuchi et al, 2010), likely drawing on its role in memory retrieval for the generative and evaluative processes involved in creative behavior. Older adults have also been shown to have higher relative activation than young adults in LG when comparing positive relative to negative visual stimuli (Kehoe et al, 2013). Here, older adults may be showing increased connectivity to lingual gyrus in part due to increased life experience, resulting in increased memory retrieval. In addition, the more pronounced connectivity effect observed during music listening may be due to a combination of memory retrieval and the aforementioned positivity effect, as the music used

is generally considered to be of positive valence and has the potential to evoke autobiographical memories in listeners.

Finally, there was a pronounced effect across all subjects, in which listening to their own self-selected music led to increased connectivity between reward network and auditory regions as compared to other recognizable music within the larger western canon. Agency in music listening is known to be an important factor in health outcomes of music-based interventions (Ruud, 1997; Cassidy & Mcdonald, 2009; Howlin et al, 2022). However, here for the first time, we present direct evidence that agency in song selection leads to increased connectivity of auditory and reward networks. This supports the use of these networks as neural targets for agency-centered music-based interventions, and helps to explain these previously observed behavioral effects.

These results represent a novel set of findings characterizing age-related differences in the musical reward response. These findings suggest that auditory and reward systems are strong targets for music-based interventions in older adults, and provide neurological support for the use of participant-selected music in such interventions. However, some limitations in the present study should be addressed. Gender disparities do exist between our two samples: the young adult sample had more female participants than the older adults sample, which was relatively gender-balanced. To account for these differences, we ran follow-up analyses to observe any main effects of gender. No significant effects were observed, suggesting that the gender differences were not confounding the present patterns of results. The age span was also not consistent between our two samples. Our older adult sample spanned a total of 35 years of age, while our young adult sample only spanned 5. Follow-up analyses revealed no main effects of within-sample age, and including within-sample age as a covariate did not have a significant impact on the observed effects, thus increasing the confidence with which we claim that effects of age primarily arose from differences between the young and older adult samples.

Our data are also limited in the extent to which participants were asked to report on their feelings about a particular piece. We did not ask about the particular emotions evoked by each piece, nor did we ask participants about the autobiographical context behind their self-selected pieces. In future studies, a more granular approach to emotional classification of pieces may be warranted to better understand the relevant contributions to auditory-reward connectivity. Furthermore, the young adult sample falls into an age range in which the brain is still developing (Giedd et al, 1999; Sowell et al, 2001), and thus for future analyses we would recommend expanding the sample to include a broader range of the lifespan, covering all age groups. Nevertheless, this novel set of findings demonstrates that different age groups do in fact vary in their auditory reward processing, further explaining observed differences in musical reward experience (Belfi et al, 2020), and adding to the depth of knowledge required for the understanding of musical reward processing.

Data availability statement

Preregistration can be found at https://osf.io/zxd42. Statistical maps of all significant results can be found at https://neurovault.org/collections/QSMRNLRW/

Author contribution

Formal Analysis, Data Curation, Methodology, Software, Validation, Investigation, Visualization, Writing - Original Draft Preparation, Writing - Review & Editing: A.B. Investigation, Project Administration, Visualization, Writing - Original Draft Preparation, Writing - Review & Editing, M.A.Q. Writing - Review & Editing: M.G., N.D., S.B.H. Conceptualization, Data Curation, Funding Acquisition, Investigation, Resources, Supervision, Visualization, Writing - Original Draft Preparation, Writing - Review & Editing: PL.

Acknowledgements and Funding Information

We acknowledge funding support from NIH R01AG078376, NIH R21AG075232, and NSF-CAREER 1945436 to PL. We acknowledge MIND Lab members Nicholas Kathios, Valerie Goutama, Dayang Gong, Jacob Ostapenko, Grace Neale, Catherine Zhou, Ritu Amarnani, Amira Toivoinen, Itamar Zik, Nicole Page, Jasper Olson, and Parker Tichko for assistance with fMRI data acquisition and analyses. We thank all our participants, and staff of Northeastern University Biomedical Imaging Center (Fred Bidmead, Valur Olafsson, Susan Whitfield-Gabrieli) for help with MRI data acquisition and analysis.

References

Ai, M., Loui, P., Morris, T. P., Chaddock-Heyman, L., Hillman, C. H., McAuley, E., & Kramer, A. F. (2022). Musical Experience Relates to Insula-Based Functional Connectivity in Older Adults. *Brain Sciences, 12(11)*, 1577.

Alluri, V., Toiviainen, P., Burunat, I., Kliuchko, M., Vuust, P., & Brattico, E. (2017). Connectivity patterns during music listening: Evidence for action-based processing in musicians. *Hum Brain Mapp*.

Aydogan, G., Flaig, N., Ravi, S. N., Large, E. W., McClure, S. M., & Margulis, E. H. (2018). Overcoming Bias: Cognitive Control Reduces Susceptibility to Framing Effects in Evaluating Musical Performance. *Scientific Reports, 8*(1), 6229.

Bannerman, D. M., Rawlins, J. N. P., McHugh, S. B., Deacon, R. M. J., Yee, B. K., Bast, T., ... & Feldon, J. (2004). Regional dissociations within the hippocampus—memory and anxiety. *Neuroscience & biobehavioral reviews, 28(3)*, 273-283.

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage, 37*(1), 90-101.

Belden, A., Zeng, T., Przysinda, E., Anteraper, S. A., Whitfield-Gabrieli, S., & Loui, P. (2020). Improvising at rest: Differentiating jazz and classical music training with resting state functional connectivity. *NeuroImage, 207*, 116384.

Belfi, A. M., & Loui, P. (2020). Musical anhedonia and rewards of music listening: current advances and a proposed model. *Annals of the New York Academy of Sciences, 1464*, 99-114.

Belfi, A. M., Moreno, G. L., Gugliano, M., & Neill, C. (2021). Musical reward across the lifespan. *Aging & Mental Health*, 1-8.

Bianco, R., Novembre, G., Keller, P. E., Kim, S. G., Scharf, F., Friederici, A. D., ... & Sammler, D. (2016). Neural networks for harmonic structure in music perception and action. *Neuroimage, 142,* 454-464.

Boemio, A., Fromm, S., Braun, A., & Poeppel, D. (2005). Hierarchical and asymmetric temporal sensitivity in human auditory cortices. *Nature neuroscience, 8(3)*, 389-395.

Bogousslavsky, J., Miklossy, J., Deruaz, J. P., Assal, G., & Regli, F. (1987). Lingual and fusiform gyri in visual processing: a clinico-pathologic study of superior altitudinal hemianopia. *Journal of Neurology, Neurosurgery & Psychiatry, 50(5)*, 607-614.

Bouhali, F., Mongelli, V., de Schotten, M. T., & Cohen, L. (2020). Reading music and words: The anatomical connectivity of musicians' visual cortex. *NeuroImage, 212*, 116666.

Brett, M., Anton, J.-L., Valabregue, R., & Poline, J.-B. (2002). *Region of interest analysis using an SPM toolbox*. Paper presented at the 8th International Conference on Functional Mapping of the Human Brain, Sendai, Japan.

Buckner, R. L., Snyder, A. Z., Shannon, B. J., LaRossa, G., Sachs, R., Fotenos, A. F., ... & Mintun, M. A. (2005). Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *Journal of neuroscience, 25(34)*, 7709-7717.

Cassidy, G., & MacDonald, R. (2009). The effects of music choice on task performance: A study of the impact of self-selected and experimenter-selected music on driving game performance and experience. *Musicae Scientiae, 13(2),* 357-386.

Cardona, G., Ferreri, L., Lorenzo-Seva, U., Russo, F. A., & Rodriguez-Fornells, A. (2022). The Forgotten Role Of Absorption In Music Reward *Annals of the New York Academy of Sciences*, In press.

Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: a review of its functional anatomy and behavioural correlates. *Brain, 129(3)*, 564-583.

Darby, R. R., Joutsa, J., Burke, M. J., & Fox, M. D. (2018). Lesion network localization of free will. *Proceedings of the National Academy of Sciences*, *115*(42), 10792-10797.

Ellis, R. J., Bruijn, B., Norton, A. C., Winner, E., & Schlaug, G. (2013). Training-mediated leftward asymmetries during music processing: a cross-sectional and longitudinal fMRI analysis. *NeuroImage, 75*, 97-107.

Eppinger, B., Nystrom, L. E., & Cohen, J. D. (2012). Reduced sensitivity to immediate reward during decision-making in older than younger adults. *PloS one, 7*(5), e36953-e36953.

Ferreri, L., Mas-Herrero, E., Zatorre, R. J., Ripollés, P., Gomez-Andres, A., Alicart, H., ... & Rodriguez-Fornells, A. (2019). Dopamine modulates the reward experiences elicited by music. *Proceedings of the National Academy of Sciences, 116(9)*, 3793-3798.

Frey, S., Campbell, J. S., Pike, G. B., & Petrides, M. (2008). Dissociating the human language pathways with high angular resolution diffusion fiber tractography. *J Neurosci, 28*(45), 11435-11444.

Friston, K. J., Ashburner, J., Frith, C. D., Poline, J. B., Heather, J. D., & Frackowiak, R. S. (1995). Spatial registration and normalization of images. *Human brain mapping, 3*(3), 165-189.

Frühholz, S., Trost, W., & Grandjean, D. (2014). The role of the medial temporal limbic system in processing emotions in voice and music. *Progress in neurobiology, 123,* 1-17.

Geddes, M. R., Mattfeld, A. T., de los Angeles, C., Keshavan, A., & Gabrieli, J. D. (2018). Human aging reduces the neurobehavioral influence of motivation on episodic memory. *NeuroImage, 171*, 296-310.

Giedd, J. N., Blumenthal, J., Jeffries, N. O., Castellanos, F. X., Liu, H., Zijdenbos, A., . . . Rapoport, J. L. (1999). Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci, 2*(10), 861-863.

Gold, B. P., Mas-Herrero, E., Zeighami, Y., Benovoy, M., Dagher, A., & Zatorre, R. J. (2019). Musical reward prediction errors engage the nucleus accumbens and motivate learning. *Proceedings of the National Academy of Sciences, 116(8),* 3310-3315.

Grady, C., Sarraf, S., Saverino, C., & Campbell, K. (2016). Age differences in the functional interactions among the default, frontoparietal control, and dorsal attention networks. *Neurobiology of aging*, *41*, 159-172.

Hafkemeijer, A., van der Grond, J., & Rombouts, S. A. (2012). Imaging the default mode network in aging and dementia. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease, 1822(3),* 431-441.

Hafkemeijer, A., Möller, C., Dopper, E. G., Jiskoot, L. C., Schouten, T. M., Van Swieten, J. C., . . . Barkhof, F. (2015). Resting state functional connectivity differences between behavioral variant frontotemporal dementia and Alzheimer's disease. *Frontiers in human neuroscience, 9*, 474.

Harrison, L. D., & Loui, P. (2014). Thrills, Chills, Frissons, and Skin Orgasms: Toward an Integrative Model of Transcendent Psychophysiological Moments in Music. *Frontiers in Psychology, 5*.

Howlin, C., Stapleton, A., & Rooney, B. (2022). Tune out pain: Agency and active engagement predict decreases in pain intensity after music listening. PloS one, 17(8), e0271329.

Janata, P. (2009). The Neural Architecture of Music-Evoked Autobiographical Memories. *Cereb. Cortex*, bhp008.

Kaas, J. H., & Hackett, T. A. (2000). Subdivisions of auditory cortex and processing streams in primates. *Proceedings of the National Academy of Sciences, 97*(22), 11793-11799.

Kehoe, E. G., Toomey, J. M., Balsters, J. H., & Bokde, A. L. (2013). Healthy aging is associated with increased neural processing of positive valence but attenuated processing of emotional arousal: an fMRI study. *Neurobiology of aging, 34(3)*, 809-821.

Karlsgodt, K. H., John, M., Ikuta, T., Rigoard, P., Peters, B. D., Derosse, P., . . . Szeszko, P. R. (2015). The accumbofrontal tract: Diffusion tensor imaging characterization and developmental change from childhood to adulthood. *Hum Brain Mapp, 36*(12), 4954-4963.

Koelsch, S. (2005). Neural substrates of processing syntax and semantics in music. *Current Opinion in Neurobiology, 15*(2), 207-212.

Kunert, R., Willems, R. M., Casasanto, D., Patel, A. D., & Hagoort, P. (2015). Music and language syntax interact in Broca's area: an fMRI study. PloS one, 10(11), e0141069.

Laukka, P. (2006). Uses of music and psychological well-being among the elderly. *Journal of Happiness Studies, 8*(2), 215.

Lima, C. F., Krishnan, S., & Scott, S. K. (2016). Roles of supplementary motor areas in auditory processing and auditory imagery. *Trends in neurosciences, 39(8),* 527-542.

Loui, P., Alsop, D., & Schlaug, G. (2009). Tone deafness: a new disconnection syndrome? *J Neurosci, 29*(33), 10215-10220.

Loui, P., Patterson, S., Sachs, M. E., Leung, Y., Zeng, T., & Przysinda, E. (2017). White matter correlates of musical anhedonia: implications for evolution of music. *Frontiers in Psychology, 1664.*

Loui, P., & Schlaug, G. (2009). Investigating Musical Disorders with Diffusion Tensor Imaging: a Comparison of Imaging Parameters. *Annals of the New York Annual Academy of Sciences, 1169*(The Neurosciences and Music III: Disorders and Plasticity), 121-125.

Machielsen, W. C., Rombouts, S. A., Barkhof, F., Scheltens, P., & Witter, M. P. (2000). FMRI of visual encoding: reproducibility of activation. *Human brain mapping, 9(3)*, 156-164.

Malach, R., Levy, I., & Hasson, U. (2002). The topography of high-order human object areas. *Trends in cognitive sciences, 6*(4), 176-184.

Martínez-Molina, N., Mas-Herrero, E., Rodríguez-Fornells, A., Zatorre, R. J., & Marco-Pallarés, J. (2016). Neural correlates of specific musical anhedonia. *Proceedings of the National Academy of Sciences, 113*(46), E7337–E7345.

Mas-Herrero, E., Marco-Pallares, J., Lorenzo-Seva, U., Zatorre, R. J., & Rodriguez-Fornells, A. (2013). Individual differences in Music Reward experiences. *Music Perception: An Interdisciplinary Journal, 31*(2), 118-138.

Mas-Herrero, E., Zatorre, R. J., Rodriguez-Fornells, A., & Marco-Pallares, J. (2014). Dissociation between Musical and Monetary Reward Responses in Specific Musical Anhedonia. *Curr Biol, 24*(6), 699-704.

Merrill, J., Sammler, D., Bangert, M., Goldhahn, D., Lohmann, G., Turner, R., & Friederici, A. D. (2012). Perception of words and pitch patterns in song and speech. *Frontiers in psychology, 3*, 76.

Nagy, K., Greenlee, M. W., & Kovács, G. (2012). The lateral occipital cortex in the face perception network: an effective connectivity study. *Frontiers in psychology, 3*, 141.

Nombela, C., Hughes, L. E., Owen, A. M., & Grahn, J. A. (2013). Into the groove: can rhythm influence Parkinson's disease?. *Neuroscience & Biobehavioral Reviews, 37(10)*, 2564-2570.

Ono, K., Nakamura, A., Yoshiyama, K., Kinkori, T., Bundo, M., Kato, T., & Ito, K. (2011). The effect of musical experience on hemispheric lateralization in musical feature processing. *Neuroscience letters, 496(2),* 141-145.

Onoda, K., Ishihara, M., & Yamaguchi, S. (2012). Decreased functional connectivity by aging is associated with cognitive decline. *Journal of cognitive neuroscience, 24(11)*, 2186-2198.

Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J., & Nichols, T. E. (2011). *Statistical parametric mapping: the analysis of functional brain images*: Elsevier.

Quinci, M. A., Belden, A., Goutama, V., Gong, D., Hanser, S., Donovan, N. J., . . . Loui, P. (2022). Longitudinal changes in auditory and reward systems following receptive music-based intervention in older adults. *Sci Rep, 12*(1), 11517.

Rentfrow, P. J., Goldberg, L. R., & Levitin, D. J. (2011). The structure of musical preferences: A five-factor model (Vol. 100, pp. 1139-1157). *US: American Psychological Association.*

Ruud, E. (1997). Music and the quality of life. *Nordic journal of music therapy, 6(2)*, 86-97.

Sachs, M. E., Ellis, R. J., Schlaug, G., & Loui, P. (2016). Brain Connectivity Reflects Human Aesthetic Responses to Music. *Social, Cognitive, and Affective Neuroscience, 11*(6), 884-891.

Sala-Llonch, R., Bartrés-Faz, D., & Junqué, C. (2015). Reorganization of brain networks in aging: a review of functional connectivity studies. *Frontiers in psychology, 6,* 663.

Salimpoor, V. N., Benovoy, M., Larcher, K., Dagher, A., & Zatorre, R. J. (2011). Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nat Neurosci, 14*(2), 257-262.

Salimpoor, V. N., van den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R. J. (2013). Interactions between the nucleus accumbens and auditory cortices predict music reward value. *Science, 340*(6129), 216-219.

Sambataro, F., Murty, V. P., Callicott, J. H., Tan, H.-Y., Das, S., Weinberger, D. R., & Mattay, V. S. (2010). Age-related alterations in default mode network: impact on working memory performance. *Neurobiology of aging, 31*(5), 839-852.

Schneider, P., & Wengenroth, M. (2009). The neural basis of individual holistic and spectral sound perception. *Contemporary music review, 28(3)*, 315-328.

Schouten, T. M., Koini, M., de Vos, F., Seiler, S., van der Grond, J., Lechner, A., . . . Rombouts, S. A. R. B. (2016). Combining anatomical, diffusion, and resting state functional magnetic resonance imaging for individual classification of mild and moderate Alzheimer's disease. *NeuroImage: Clinical, 11*, 46-51.

Sheline, Y. I., Raichle, M. E., Snyder, A. Z., Morris, J. C., Head, D., Wang, S., & Mintun, M. A. (2010). Amyloid plaques disrupt resting state default mode network connectivity in cognitively normal elderly. *Biological psychiatry, 67*(6), 584-587.

Sowell, E. R., Thompson, P. M., Tessner, K. D., & Toga, A. W. (2001). Mapping continued brain growth and gray matter density reduction in dorsal frontal cortex: Inverse relationships during postadolescent brain maturation. *Journal of Neuroscience, 21*(22), 8819-8829.

Staffaroni, A. M., Brown, J. A., Casaletto, K. B., Elahi, F. M., Deng, J., Neuhaus, J., . . . Saloner, R. (2018). The longitudinal trajectory of default mode network connectivity in healthy older adults varies as a function of age and is associated with changes in episodic memory and processing speed. *Journal of Neuroscience, 38*(11), 2809-2817.

Sutcliffe, R., Du, K., & Ruffman, T. (2020). Music Making and Neuropsychological Aging: A Review. *Neurosci Biobehav Rev*.

Szucs, D., & Ioannidis, J. P. (2020). Sample size evolution in neuroimaging research: An evaluation of highly-cited studies (1990-2012) and of latest practices (2017-2018) in high-impact journals. *NeuroImage, 221*, 117164. https://doi.org/10.1016/j.neuroimage.2020.117164

Tanaka, S., & Kirino, E. (2016). Functional connectivity of the precuneus in female university students with long-term musical training. *Frontiers in human neuroscience, 10*, 328.

Tanaka, S., & Kirino, E. (2019). Increased functional connectivity of the angular gyrus during imagined music performance. *Frontiers in human neuroscience, 13*, 92.

Takeuchi, H., Taki, Y., Sassa, Y., Hashizume, H., Sekiguchi, A., Fukushima, A., & Kawashima, R. (2010). White matter structures associated with creativity: evidence from diffusion tensor imaging. *Neuroimage, 51(1)*, 11-18.

Tichko, P., Page, N., Kim, J. C., Large, E. W., & Loui, P. (2022). Neural Entrainment to Musical Pulse in Naturalistic Music Is Preserved in Aging: Implications for Music-Based Interventions. *Brain Sciences, 12(12)*, 1676.

Trost, W., Ethofer, T., Zentner, M., & Vuilleumier, P. (2011). Mapping Aesthetic Musical Emotions in the Brain. *Cereb Cortex*.

Utevsky, A. V., Smith, D. V., & Huettel, S. A. (2014). Precuneus is a functional core of the default-mode network. *Journal of Neuroscience, 34*(3), 932-940.

Vuust, P., Heggli, O. A., Friston, K. J., & Kringelbach, M. L. (2022). Music in the brain. *Nature Reviews Neuroscience*.

Wang, D., Belden, A., Hanser, S., Geddes, M., & Loui, P. (2020). Resting-State Connectivity of Auditory and Reward Systems in Alzheimer's Disease and Mild Cognitive Impairment. *Front. Hum. Neurosci., 14*, 280.

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connect, 2*(3), 125-141.

Yeterian, E. H., & Pandya, D. N. (1998). Corticostriatal connections of the superior temporal region in rhesus monkeys. *J Comp Neurol, 399*(3), 384-402.

Zatorre, R. J., Evans, A. C., Meyer, E., & Gjedde, A. (1992). Lateralization of phonetic and pitch discrimination in speech processing. *Science, 256(5058)*, 846-849.

Zatorre, R. J., & Belin, P. (2001). Spectral and temporal processing in human auditory cortex. *Cerebral cortex, 11(10),* 946-953.

Zatorre, R. J. (2015). Musical pleasure and reward: mechanisms and dysfunction. *Annals of the New York Academy of Sciences, 1337*(1), 202-211.

Tables

Table 1 | Liking Rating Frequencies by Group

	Rating			
	Hate	Neutral	Like	Love
Young Adults				
% within Group	13.80%	32.50%	23.20%	30.50%
% within Rating	43.60%	57.90%	46.30%	48.90%
% of Total	6.90%	16.20%	11.60%	15.30%
Older Adults				
% within Group	17.80%	23.50%	26.80%	31.90%
% within Rating	56.40%	42.10%	53.70%	51.10%
% of Total	8.90%	11.80%	13.40%	16.00%
Total				
% of Total	15.80%	28.00%	25.00%	31.20%

Table 2 | Univariate Whole-Brain Results

Region(s)/Clusters	Side	X	Y	Z	Cluster Size
Young Adults					
Hate					
STG/STS/MTG/Heshl's Gyrus	Right	54	-6	-3	455
Planum Temporale/Polare	Left	-51	-18	3	302
Parahippocampal Gyrus	Left	-12	-9	-27	20
Hippocampus	Right	21	-15	-24	12
Brainstem	Left	-3	-33	-66	11
		-18	-78	-45	10
Neutral					
STG/STS/MTG/Heshl's Gyrus	Right	48	-18	6	506
Planum Temporale/Polare	Left	-48	-15	0	378
Parahippocampal Gyrus	Right	15	-15	-24	12
Like					
STG/STS/MTG/Heshl's Gyrus	Right	63	-21	6	493
Planum Temporale/Polare	Left	-48	-18	0	366
Brainstem	Left	-18	-81	-48	14
Parahippocampal Gyrus Hippocampus	Right	18	-12	-24	11
Love					
STG/STS/MTG/Heshl's Gyrus	Left	-48	-15	0	1064
Planum Temporale/Polare	Right	63	-24	6	918
Paracingulate Gyrus Medial Prefrontal Cortex	Left	-6	57	-9	333
Hippocampus	Right	12	-30	-12	147
Superior Frontal Gyrus	Left	-21	24	39	52
Lateral Occipital Cortex	Left	-36	-78	33	34
Precuneus Posterior Cingulate Cortex	Left	-9	-54	12	34
Brainstem	Right	9	-48	-42	32
Parahippocampal Gyrus	Left	-18	-36	-21	19
Orbitofrontal Cortex	Left	-30	33	-12	14
Supplementary Motor Area	Right	6	6	63	12
Older Adults					
Hate					
STG/STS/MTG/Heshl's Gyrus	Right	63	-24	0	320
Planum Temporale/Polare	Left	-54	-18	3	254
Neutral					
STG/STS/MTG/Heshl's Gyrus	Right	51	-6	-3	316
Planum Temporale/Polare	Left	-51	-18	3	284
Like					
STG/STS/MTG/Heshl's Gyrus	Right	48	-12	0	557
Planum Temporale/Polare	Left	-51	-18	3	490
Hippocampus	Right	18	-15	-21	31
	Left	-18	-12	-21	20
Love					
STG/STS/MTG/Heshl's Gyrus	Right	51	-6	-3	644
Planum Temporale/Polare	Left	-45	-27	6	632
Posterior Cingulate Cortex Precuneus	Left	-6	-54	15	194
Medial Prefrontal Cortex Paracingulate Gyrus	Left	-3	51	-15	139
Lateral Occipital Cortex	Left	-42	-72	33	103
Hippocampus	Right	21	-18	-21	47
	Left	-18	-21	-21	26
Brainstem	Right	3	-36	-12	38
Ventral Striatum/ Nucleus Accumbens	Left	-3	9	-12	33
Orbitofrontal Cortex	Left	-36	30	-15	19
Cerebellum	Right	42	-69	-42	16

Voxel Threshold: $p < 0.05$ p-FDR corrected; Cluster Threshold: $k > 10$ cluster-size

Table 3 | Seed-Based Connectivity Results

Region	Hemisphere	X	Y	Z	k	Direction
YA(1) OA(-1); Rest; Auditory Network Seed						
Precuneous	Bilateral	24	-72	32	20652	OA>YA
MFG	Right	34	42	38	713	OA>YA
MTG/STG	Left	-68	-18	-4	554	YA>OA
MTG/STG	Right	72	-28	0	479	YA>OA
MFG	Left	-36	50	26	427	OA>YA
LOC	Left	-38	-72	44	383	YA>OA
PostCG	Left	-48	-24	32	319	OA>YA
Caudate	Right	28	6	12	262	OA>YA
Putamen	Left	-20	6	14	221	OA>YA
PostCG	Right	58	-18	38	217	OA>YA
YA(1) OA(-1); Music; Auditory Network Seed						
Precuneous	Bilateral	-12	-78	2	22094	OA>YA
MFG	Left	-36	44	18	1321	OA>YA
MFG	Right	36	38	32	1285	OA>YA
PreCG	Left	-56	2	6	935	OA>YA
Insular Cortex	Right	30	14	6	742	OA>YA
Thalamus	Right	30	-30	-12	489	OA>YA
STG/MTG	Right	68	-42	14	359	YA>OA
White Matter	Left	-36	-46	12	313	YA>OA
Cerebellum	Right	30	-40	-42	283	OA>YA
Hippocampus	Right	12	-6	-24	221	YA>OA
AC	Bilateral	0	32	-4	214	YA>OA
mPFC	Bilateral	-2	20	-16	194	YA>OA
YA(1) OA(-1); Rest; Reward Network Seed						
Precuneous	Bilateral	22	-76	30	14053	OA>YA
AC	Bilateral	6	18	18	1531	YA>OA
HG	Right	58	-28	8	445	OA>YA
PreCG	Right	42	-10	42	415	OA>YA
SFG	Right	22	32	42	298	OA>YA
MFG	Right	40	14	42	244	OA>YA
HG	Left	-44	-34	8	189	OA>YA
PreCG	Left	-56	2	20	171	OA>YA
YA(1) OA(-1); Music; Reward Network Seed						
Lingual Gyrus	Bilateral	-2	-70	0	1282	OA>YA
PreCG	Bilateral	-6	-24	74	760	OA>YA
IFG	Left	-38	44	-4	739	OA>YA
IFG	Right	42	42	-12	629	OA>YA
AC	Bilateral	-6	-4	30	437	YA>OA
PostCG	Right	36	-36	62	292	OA>YA
Precuneous	Bilateral	4	-72	38	245	OA>YA
PreCG	Left	-42	-12	30	208	OA>YA
PreCG	Right	46	-12	36	186	OA>YA
Pallidum	Left	-20	-6	-4	143	YA>OA
Pallidum	Right	18	2	-4	129	YA>OA
YA; Music(1) Rest(-1); Auditory Network Seed						
PostCG	Left	-42	-36	56	14668	Rest>Music
PostCG	Right	34	-54	60	6882	Rest>Music
Lingual Gyrus	Left	-18	-52	8	5748	Music>Rest
STG/MTG	Left	-56	-16	2	3770	Music>Rest
STG/MTG	Right	52	6	-10	2841	Music>Rest
Cuneal Cortex	Bilateral	-12	-88	36	2252	Music>Rest
mPFC	Bilateral	-2	50	-12	1883	Music>Rest
ITG	Right	54	-24	-22	1653	Rest>Music
OFusG	Right	24	-88	-12	739	Rest>Music
Brain Stem	Bilateral	10	-18	-42	526	Rest>Music
Forb	Left	-48	42	-10	411	Rest>Music
FOrb	Right	24	32	-12	374	Rest>Music
Insular Cortex	Right	36	-6	12	362	Rest>Music
ITG	Right	54	-10	-42	316	Rest>Music
LOC	Left	-42	-84	12	284	Music>Rest
PreCG	Right	64	12	20	248	Rest>Music
AG	Right	66	-50	26	233	Rest>Music
FOrb	Left	-24	30	-18	227	Music>Rest
OFusG	Left	-20	-88	-16	226	Rest>Music
OA; Music(1) Rest(-1); Auditory Network Seed						
PostCG	Left	-44	-18	44	7422	Rest>Music
PreCG	Right	28	-54	60	5459	Rest>Music
Precuneous	Bilateral	-6	-48	0	1182	Music>Rest
STG/MTG	Left	-66	-28	0	889	Music>Rest
STG/MTG	Right	52	-12	-10	726	Music>Rest
MTG	Left	-56	-52	2	458	Rest>Music
Thalamus	Bilateral	0	-4	0	211	Music>Rest
LOC	Left	-42	-72	36	187	Music>Rest
Cerebellum	Left	-8	-54	-30	158	Music>Rest
PreCG	Left	-54	2	24	138	Rest>Music
Insular Cortex	Left	-36	-10	12	103	Rest>Music
Lingual Gyrus	Left	-2	-66	-10	96	Music>Rest
ITG	Left	-44	2	-46	94	Rest>Music
YA; Music(1) Rest(-1); Reward Network Seed						
AG	Right	54	-58	18	119	Music>Rest
LOC	Left	-44	-72	14	95	Music>Rest
All Subjects; Self-Selected(1) Familiar Western(-1); Reward Network Seed						
STG/MTG	Left	-50	-30	0	1958	SS>FW
STG/MTG	Right	48	-18	-4	1929	SS>FW
Forb	Left	-42	18	-10	108	SS>FW
All Subjects; Hate(-3) Neutral(-1) Like(1) Love(3); Auditory Network Seed						
STG/MTG	Left	-62	-24	6	1142	
STG/MTG	Right	52	-18	-4	820	
All Subjects; Hate(-3) Neutral(-1) Like(1) Love(3); Reward Network Seed						
STG/MTG	Right	46	-22	-4	42	

Voxel height threshold p < 0.05, FDR-corrected; Cluster size threshold p < 0.05, FDR-Corrected

Figure Captions

Figure 1: Univariate 2nd level results showing activity for (A) Young Adults and (B) Older Adults for different liking ratings. (C) Subcortical structures involved with listening to loved music for both Young (red), Older (blue), both (purple).

Figure 2: Age Related Differences in ROI-ROI Connectivity for Various Levels of Liking. (A) Correlation matrices showing the relationship between auditory and reward regions for different liking ratings for young and older adults. (B) Bar graphs depicting overall connectivity between auditory-auditory, reward-reward, and auditory-reward regions for young (orange) and older (blue) adults for hated, neutral, liked, and loved musical stimuli.

Figure 3: Age Differences in Auditory Network Seed-Based Connectivity: Significant clusters for YA>OA contrast in auditory network seed, significant at the $p < 0.05$, FDR-corrected level for both voxel height and cluster size. (A) Visualization of significant clusters in right hemisphere during resting state scan, including clusters favoring young adults (warm tones) and clusters favoring older adults (cool tones). (B) Bar graphs depicting effect sizes for all significant clusters in resting state contrast. Clusters with higher connectivity in young adults (red) are depicted in the top graph, and clusters with higher connectivity in older adults (blue) are depicted in the bottom graph. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster. (C) Visualization of significant clusters in right hemisphere during music listening task, including clusters favoring young adults (warm tones) and clusters favoring older adults (cool tones). (D) Bar graphs depicting effect sizes for all significant clusters in music listening contrast. Clusters with higher connectivity in young adults (red) are depicted in the top graph, and clusters with higher connectivity in older adults (blue) are depicted in the bottom graph. (E) Line graph

depicting ROI-ROI connectivity between auditory network regions and mPFC comparing across groups and liking ratings. Error bars depict +/- two standard errors, and marginal means are adjusted for covariates of gender, BMRQ, and within group age.

Figure 4: Age Differences in Reward Network Seed-Based Connectivity: Significant clusters for YA>OA contrast in reward network seed, significant at the $p < 0.05$, FDR-corrected level for both voxel height and cluster size. (A) Visualization of significant clusters in right hemisphere during resting state scan, including clusters favoring young adults (warm tones) and clusters favoring older adults (cool tones). (B) Bar graphs depicting effect sizes for all significant clusters in resting state contrast. Clusters with higher connectivity in young adults (red) are depicted in the top graph, and clusters with higher connectivity in older adults (blue) are depicted in the bottom graph. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster. (C) Visualization of significant clusters in right hemisphere during music listening task, including clusters favoring young adults (warm tones) and clusters favoring older adults (cool tones). (D) Bar graphs depicting effect sizes for all significant clusters in music listening contrast. Clusters with higher connectivity in young adults (red) are depicted in the top graph, and clusters with higher connectivity in older adults (blue) are depicted in the bottom graph.

Figure 5: Task Differences in Seed-Based Connectivity: Significant clusters for music > rest contrast, significant at the $p < 0.05$, FDR-corrected level for both voxel height and cluster size. (A) Visualization of significant clusters from auditory network seed in right hemisphere of young adults, including clusters favoring music listening (warm tones) and clusters favoring rest (cool tones). (B) Bar graphs depicting effect sizes for all significant clusters in young adult contrast. Clusters with higher connectivity during music listening (red) are depicted in the top graph, and clusters with higher connectivity during rest (blue) are depicted in the bottom graph.

Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster. (C) Visualization of significant clusters from auditory network seed in right hemisphere of older adults, including clusters favoring music listening (warm tones) and clusters favoring rest (cool tones). (D) Bar graphs depicting effect sizes for all significant clusters in older adult contrast. Clusters with higher connectivity during music listening (red) are depicted in the top graph, and clusters with higher connectivity during rest (blue) are depicted in the bottom graph. (E) Visualization of significant clusters from Reward network seed in young adults, and bar graphs depicting effect sizes for all significant clusters. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster.

Figure 6: Differences of Musical Preference in Seed-Based Connectivity: Significant clusters for effects of musical selection (self-selected > familiar western) and linear effects of musical liking in auditory and reward network seed-based connectivity ([-3,-1,1,3], two-tailed T-test; $p < 0.05$, FDR-corrected for voxel height and cluster size). (A) Visualization of significant clusters from reward network seed showing effects of selection, and bar graphs depicting effect sizes. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster. (B) Visualization of significant clusters from auditory network seed showing linear effects of liking, and bar graphs depicting effect sizes. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster. (C) Visualization of significant clusters from reward network seed showing linear effects of liking, and bar graphs depicting effect sizes. Cluster labels are MNI coordinates for center of mass of each significant cluster, as well as the most prevalent anatomical region included within each cluster.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figures

Figure S1

Supplementary Figure 1: Age Differences in Auditory Network Seed-Based Connectivity Corrected for Differences in BMRQ: Significant clusters for YA>OA contrast in auditory-network seed when including BMRQ as a covariate of no interest, significant at the p < 0.05, FDR-corrected level for both voxel height and cluster size. (A) Visualization of significant clusters during resting state scan, including clusters favoring young adults (warm tones) and clusters favoring older adults (cool tones). (B) Visualization of significant clusters favoring older adults during music listening task.

Figure S2

Supplementary Figure 2: Age Differences in Reward Network Seed-Based Connectivity

Corrected for Differences in BMRQ: Significant clusters for YA>OA contrast in reward-network

seed when including BMRQ as a covariate of no interest, significant at the p < 0.05, FDR-

corrected level for both voxel height and cluster size. (A) Visualization of significant clusters

favoring young adults during resting state scan. (B) Visualization of significant clusters favoring

older adults during music listening task.

Figure S3

Supplementary Figure 3: Auditory Network Seed-Based Connectivity of Musical Preference

Corrected for Differences in BMRQ: Significant clusters for linear liking contrast (hate, neutral,

like, love [-3,-1,1,3]) in auditory-network seed across all subjects when including BMRQ as a

covariate of no interest. Clusters are significant at the p < 0.05, FDR-corrected level for both

voxel height and cluster size.

Figure S4

Supplementary Figure 4: ROI-ROI MANCOVA Results. Bar graph comparing older adults (blue) and young adults (red) across liking ratings in measures of auditory-auditory (A), reward-reward (B), auditory-reward (C), auditory-mPFC (D), Reward-mPFC (E), Auditory-PCC (F) and reward-PCC (G) connectivity. Error bars represent +/- two standard errors, and covariates appearing in the model are evaluated at the following values: Gender = 0.3947, BMRQ = -0.3344, within-group age = -0.1491.