Single-cell transcriptional analysis of lamina propria lymphocytes in the jejunum reveals ILC-like cells in pigs

Junhong Wang [1,2,3,4]，Mingyang Cheng [1,2,3,4], Ming Gao [1,2,3,4], Yu Sun [1,2,3,4], Yiyuan Lu [1,2,3,4], Xiaoxu Li [1,2,3,4], Chunwei Shi [1,2,3,4], Jianzhong Wang [1,2,3,4], Nan Wang [1,2,3,4], Wentao Yang [1,2,3,4], Yanlong Jiang [1,2,3,4], Haibin Huang [1,2,3,4], Guilian Yang [1,2,3,4], Yan Zeng [1,2,3,4]*, Chunfeng Wang [1,2,3,4]*, Xin Cao [1,2,3,4]*

[1] College of Veterinary Medicine, Jilin Agricultural University, Changchun, China.

[2] Jilin Provincial Key Laboratory of Animal Microecology and Healthy Breeding, Jilin Agricultural University, Changchun, China.

[3] Jilin Provincial Engineering Research Center of Animal Probiotics, Jilin Agricultural University, Changchun, China.

[4] Key Laboratory of Animal Production and Product Quality Safety of Ministry of Education, Jilin Agricultural University, Changchun, China.

*Correspondence: zengyan@jlau.edu.cn; wangchunfeng@jlau.edu.cn; xinc@jlau.edu.cn

Abstract

Pigs are the most suitable model to study various therapeutic strategies and drugs for human beings, while knowledge about tissue- and cell type-specific transcriptomes and heterogeneity is poorly available. Here, we focused on the intestinal immunity of pigs. Through single-cell sequencing (scRNA-seq) and flow cytometry analysis of the types of immune cells in the jejunum of pigs, we found that innate lymphoid cells (ILCs) existed in the lamina propria lymphocytes (LPLs) of the jejunum. Then, through flow sorting of Live/Dead (L/D)⁻Lineage(LIN)⁻CD45⁺ cells and scRNA-seq, we found that ILCs in the porcine jejunum were mainly ILC3s, with a small number of ILC1s, ILC2s and NK cells. Through a gene expression map, we found that ILCs coexpressed IL-7Rα, ID2 and other genes and differentially expressed RORC, GATA3 and other genes but did not express the CD3 gene. According to their gene expression profiles, ILC3s can be divided into four subgroups, and genes such as CXCL8, CXCL2, IL-22, IL-17 and NCR2 are differentially expressed. To further detect and identify ILC3s, we prepared RORC monoclonal antibodies and verified the classification of ILCs in the porcine jejunum subgroup and the expression of related hallmark genes at the protein level by flow cytometry. For systematically characterizing of ILCs in the porcin intestines, we combined our pigs ILCs dataset with publicly available humans and mice ILCs data and identified that the humans and pigs ILCs shared more common features than that of mice in gene signatures and cell

states. Our results for the first time showed in detail the gene expression of porcine jejunal ILCs, the subtype classification of ILCs and the markers of various ILCs, which provides a basis for in-depth exploration of porcine intestinal mucosal immunity.

Key words: innate lymphoid cells; swine; jejunum; scRNA-seq; mucosal immunity

Introduction

As an ideal animal model that is expected to replace human beings as study subjects, pigs can be used to simulate complex human diseases, for example, pigs can be used as a computational model for predicting human pharmacokinetics (PK) (Pedersen et al., 2022), as an animal model for tissue expansion experiments (Han et al., 2022), and as an animal model for burn research (Abdullahi et al., 2014). In addition, because they are highly similar to human beings in organ size, structure, anatomy, genetics and physiological functions, they can be used as ideal donor sources for organ transplantation (Wolf et al., 2019; Sykes and Sachs, 2019). Regarding the immune system, the immune cell subsets expressed by pigs are similar to those expressed by humans in phenotype, and the two species have high nucleotide and protein sequence homology in immune molecules (Käser, 2021; Starbæk et al., 2018). Therefore, pigs have great research value and potential, but basic immunology research related to pigs is still in the primary stage, and there are few reports on the intestinal immunity of pigs, especially on innate lymphoid cells (ILCs).

In recent years, it has been found that innate lymphoid cells (ILCs) play important functions in immunity and tissue homeostasis (Harly et al., 2019). The mutual interaction between the microbiome and immune system coordinates the influence of many microorganisms on the host, and ILCs are the frontline that continuously interact with symbiotic flora in the intestine (Mortha et al., 2014; Yu et al., 2022; Zhou et al., 2020a; Ronan et al., 2021). ILCs are a new family of innate lymphocytes that come from bone marrow lymphoprogenitor cells and lack classical rearranged antigen receptors (Vivier et al., 2016). They migrate to various tissue sites, mucous membranes, etc., where they participate in antipathogen reactions, inflammation, tissue homeostasis and immune tolerance (Diefenbach et al., 2020; Spits and Cupedo, 2012). According to their functions, four main subgroups of ILCs are defined: ILC1s, ILC2s, ILC3s and NK cells (Vivier et al., 2018; Warner and Ohashi, 2019; Fachi et al., 2020; Xu et al., 2019; Seillet et al., 2020). In addition to ILCs with effector functions, some ILCs with regulatory functions (ILCregs) have been defined (Wang et al., 2017; Seehus et al., 2017). ILC3s have signature gene expression such as KIT (CD117), IL-7Rα (CD127) (Pascual-Reguant et al., 2021), and IL-23R (Schleussner et al., 2018). ILC3s can work with commensal microbes to promote intestinal homeostasis and regulate inflammation and mucosal immunity (Liu et al., 2020). ILC1s have the closest relationship with Cytotoxic CD8[+] T cells, expressing higher levels of genes encoding cytotoxic molecules than other ILC/T types (Hidalgo et al., 2008). A previous study found higher levels of Th2/ILC2-type cells in the lung mucosa (Tibbitt et al., 2019; Zeis et al., 2020). ILC2s have functional similarities to Th2 cells in terms of the expression of the master transcription factor GATA3 and the production of functional cytokines and can play important roles in the expulsion of parasites and induction of allergy through secretion of IL-5 and IL-13, including myeloid cell recruitment, eosinophil maturation, smooth muscle contraction, and mucin production by epithelial cells (von Moltke et al., 2016; Gurram and Zhu, 2019). To date, NK cells are the only identified porcine non-B/non-T-lymphocyte subset and are an important cell population of the

innate immune system. With the ability to attack natural pathogens with infected and malignant body cells as well as to produce immunomodulatory cytokines, NK cells are commonly identified by flow cytometry as CD3ε⁻CD8α⁺ (Gerner et al., 2009).

Single-cell RNA sequencing (scRNA-seq) has been used to describe the gene expression of immune cells in many tissues and organs of pigs, including peripheral blood (Herrera-Uribe et al., 2021), thymus (Gu et al., 2022), lung and embryonic tissue (Zhang et al., 2021; Kong et al., 2020; Liu et al., 2021). The heterogeneity of endothelial cells, microglia regulation and transcriptome changes under single-cell resolution of virus-infected porcine macrophages were also revealed through the cell landscape at the pig single-cell level (Zheng et al., 2022). Recently, researchers have found the presence of suspected ILCs in the intestinal immune cells of pigs (Zhang et al., 2019; Wiarda et al., 2022). However, there is no report on the gene expression and cell typing of ILCs in the porcine intestine. We extracted total lamina propria lymphocytes (LPLs) from the porcine jejunum, obtained Live/Dead (L/D)⁻Lineage(LIN)⁻CD45⁺ cells (amount: 200,000) by fluorescence-activated cell sorting (FACS), and then performed scRNA-seq. We recorded and presented the ILC population and heterogeneity level in the porcine jejunum, which has not been previously described. Collectively, these data are the highest resolution transcriptome maps of ILCs in the intestinal immune landscape of pigs and can be used to further decode the phenotype and function of ILCs in the intestinal tract. By performing scRNA-seq on immune cells from the jejunum of piglets, we recorded and showed the specific typing of porcine ILCs and defined the ILC3 subgroup in detail. It was found that there was a certain transdifferentiation process of ILCs in the piglet jejunum under complex intestinal conditions. Unbiased cluster analysis of ILCs of humans, pigs and mice showed that the gene expression similarity of ILCs of humans and porcine intestines was high, but the similarity between pigs and mice was not high.

Our data provide a resource to better understand porcine ILCs of LPLs in the jejunum. We recorded and showed the specific typing of porcine ILCs and the characterizing of the ILC3 subgroup in detail. It was found that there was a certain transdifferentiation process of ILCs in the piglet jejunum under complex intestinal conditions. Cluster analysis of ILCs in humans, pigs and mice showed that the gene expression similarity of human and pig intestinal ILCs was high, but the similarity between pigs and mice was not high, which suggested that swine could be a better substitute for humans as model animals to investigate intestinal mucosal system homeostasis after various challenges.

Results
1. Characterization of immune cells in LPLs of the jejunum in pigs
The immune cells in the porcine jejunum lamina propria are abundant, and the cell types are obvious. We identified a large number of myeloid cells present in the pig lamina jejuna, including dendritic cells (CD172⁺SLAII⁺CD163⁻), macrophages (CD172⁺SLAII⁺CD163⁺), polymorphonuclear neutrophil (CD172⁺SLAII⁻CD163⁻), and monocytes (CD172⁺SLAII⁺/⁻CD163⁺/⁻). T cells in the lamina propria include two cell subsets, namely, CD3⁺γδT⁻ and CD3⁺γδT⁺, and we found more CD4⁺ cells in the CD3⁺γδT⁻ subset and more CD8⁺ cells in the CD3⁺γδT⁺ cell subset (Gerner et al., 2009). We found certain proportion of B cells expressing SLAII and CD8⁺ NK cells expressing CD11b and CD11c were present in the pig lamina propria (Jin et al., 2021; Piriou-Guzylack and Salmon, 2008) . To obtain a large number of ILCs, the cells obtained by flow sorting were subjected to subsequent single-cell transcriptome

sequencing (Figure S1A).

The analysis of single-cell transcriptome data of the pig jejunum shows that there are a large number of nonimmune cells and immune cells in the jejunum, among which immune cells (CD45) are mainly divided into myeloid cells, T cells (CD3), B cells (CD21) and ILCs. However, at this resolution, ILCs are mainly ILC3s (IL-23R), and the classification of ILCs into other subgroups is not clear. These immune cells together constitute the immune system of the porcine jejunum (Figure S1B, C) (Table S1).

To better explore the specific grouping information and expression profiles of inherent lymphoid cells in the jejunum of pigs, we removed the cells with abnormally low and high gene counts and high mitochondrial gene expression and then used Seurat to cluster the remaining cells (Stuart et al., 2019). A total of 15,763 immune cells were obtained. According to the different transcription levels of genes of different cells, they can be preliminarily divided into four types of cells (Figure 1A), among which there are 11,208 ILCs/T cells, whose surface receptors are CD127 and nuclear transcription factors such as GATA3 and RORC. There are 3,500 myeloid cells that express THBS1, TIF1, MS4A2 and other landmark genes of myeloid cells, and the gene expression of myeloid cells is heterogeneous, which is related to the functional polarization of myeloid cells. There were 717 B cells, which express B-cell markers such as MZB1, JCHAIN and TNFS17R, and 338 unknown immune cells (Figure 1B, 1C). According to the gene expression clustering UMAP analysis, the cells were divided into 13 cell subsets (Figure 1D) (Table S2), in which the symbolic marker expressed in each subgroup is shown in Figure 1E. With CD172 antibodies in Lin, the single-cell results showed that only some myeloid cells were detected, such as the cell subsets of macrophages, mast cells, granulocytes, dendritic cells, and monocytes. Although we removed most of the non-ILCs from the immune cells, some unknown cells were still present, which also suggests that the immune cell subsets in the pig intestinal lamina propria are very enriched, possibly due to the influence of the complex intestinal environment.

2. ILC3s are the most abundant cell type among Lin$^-$CD45$^+$ cells in LPLs of the jejunum in pigs

Early sorting has removed a large number of T cells and is therefore not abundant for typing T-cell subpopulations. The regrouping results showed that ILC/T cells can be divided into seven subsets according to their hallmarks, namely, ILC3s, CD8$^+$ T cells, CD4$^+$ T cells, CD3$^+$ T cells, ILC1s, ILC2s, and NK cells (Figure 2A). Studies have shown that typical cytokines and transcription factors produced by Th1, Th2 and Th17 cells are also secreted by ILC1s, ILC2s and ILC3s (Spits et al., 2013), respectively. The results of the surface molecular responses of T cells and ILCs were found to be quite different. T cells expressed CD3E, which could be divided into CD4 T cells and CD8 T cells, while ILCs expressed surface markers such as IL-23R (ILC3), PTGDR2 (ILC2) and NCR1 (NK) (Figure 2B). Both T cells and ILCs express the chemokines CXCL8 (ILC3, CD8 T, and CD3 T cells) and CCL5 (CD8 T, CD3 T, ILC1, and NK cells), and CD4 T cells express more IL-10 (Figure 2C). For transcription factors, T cells and ILCs are similar, as both express ID2 and RORC (ILC3, CD3 T); IKZF3 is mainly expressed by T cells, ILC1s and NK cells; TBX21 and EOMES are mainly expressed by NK cells; and GATA3, as an important regulator of T-cell development, is expressed in ILC/T cells (Figure 2D).

In the barrier tissues of mammals, ILCs account for approximately 0.5-5% of lymphocytes, so they are a rare cell group (Simoni et al., 2018; Halim and Takei, 2014). Our scRNA-seq results showed that there were ILCs in the intestinal lamina propria of pigs, which accounted for a certain

proportion (Figure S2A). Cluster analysis of ILCs showed that the correlation between ILC3s and ILC2s was higher, while that between NK cells and ILC1s was higher (Figure S2B). For the GSVA analysis of ILCs (Figure S2C) (Table S3), subsets of ILCs were highly enriched in pathways such as LINOLEIC_ACID_METABOLISM, RENIN_ANGIOTENSIN_SYSTEM, and MATURITY_ONSET_DIABETES_OF_THE_YOUNG, suggesting that ILCs play a crucial role in development and are involved in the regulation of morphogenesis in various tissues and organs. However, the enrichment degree in PENTOSE_PHOSPHATE_PATHWAY, ACID_METABOLISM and other pathways was relatively low.

We performed a detailed statistical analysis on the surface receptors, effector factors, transcription factors and other highly expressed genes of ILCs (Table S4). We found that except for ILC3s, other ILCs accounted for a relatively low proportion of the total ILCs.

ILC3s

We found that porcine ILC3s expressed many immune characteristic genes of this cell type. For the surface receptors, ILC3s expressed iconic surface receptors such as CD2, KIT, IL-7Rα, IL-23R, NCR and AHR (Figure 2E). IL-7Rα plays an important role in the development and differentiation of ILC3s (Bernink et al., 2015; Robinette et al., 2017), and NCR2+ ILC3s have a unique function in maintaining intestinal homeostasis (Song et al., 2015). ILC3s highly expressed the powerful chemokines CXCL2 and CXCL8 of neutrophils (Figure 2F), which revealed their regulatory effect on granulocyte-affecting migration and survival (Furue et al., 2020; Croxatto et al., 2016). Similar to TH17 cells, ILC3s express the TNF, IL-22, IL-17A and IL-17F genes, which is the main reason why ILC3s play an important role in the innate immune response, especially in the intestinal mucosa (Riding et al., 2022; Serafini et al., 2022). ILC3s represent a series of transcription factors (Figure 2G), among which ID2 is an important transcription factor that regulates many cellular processes (Zhong et al., 2020); RORC (RORγt) has been used as a symbolic transcription factor of ILC3s (Croft et al., 2022); and Runx3 is necessary for the normal development of ILC1s and ILC3s. ETS1 is involved in stem cell development, cell senescence and death (Lamber et al., 2008), and FOS is a regulator of cell proliferation (Güller et al., 2008), differentiation and transdifferentiation. We also found that ILC3s in the porcine intestine highly expressed the gonadotropic development-related gene TAC3 (Figure 2H) (Bianco and Kaiser, 2009), and the TMIGD2 gene was involved in the positive regulation of T-cell activation (Li et al., 2022b). The transcribed genes of ILC3s were enriched in pathways such as HEMATOPOIETIC_CELL_LINEAGE, NITROGEN_METABOLISM, CYTOKINE_CYTOKINE_RECEPTOR_INTERACTION, and ARACHIDONIC_ACID_METABOLISM. ILC3s are involved in innate and adaptive inflammatory host defense, cell growth, differentiation, cell death, angiogenesis, and developmental and repair processes aimed at restoring homeostasis and regulating nitrogen metabolism, arachidonic acid metabolism and other processes (Figure S2D).

ILC1s

ILC1s still express other genes related to the CD3 complex, such as CD3G and CD247 (encoding CD3z) (Figure 2E). The antigen presentation-related genes CD96 and SLA-DRA (Georgiev et al., 2018) and the gene ITGAE (integrin subunit αE), which mediates the adhesion of T lymphocytes to epithelial cells, were also expressed. The expression of genes encoding cytotoxic molecules, including GZMA, GZMB, GZMH and GNLY (Hidalgo et al., 2008), was significantly increased (Figure 2F). Additionally, the gene PRF1 encoding the perforin protein was

found (Horne et al., 2008). The main regulator gene of activation-induced alternative splicing in T cells is HNRNPLL (Yaman et al., 2009). The transcription factors identified for ILC1s include SOX4 and the transcription inhibitor MXI1 (Figure 2G). It was also found that ILC1s were highly responsive to the transcription factor GRHL1 and the transcription inhibitor BATF3. The EVL gene had high expression in T cells and ILC1s (Figure 2H). It has been reported that the PRDX2 gene may contribute to the antiviral activity of CD8+ T cells (Besgen et al., 2010). The ILC1-enriched pathways are all related to autoimmune diseases, inducing cytotoxic T lymphocyte (CTL) and natural killer (NK) cell responses (Figure S2D).

ILC2s

Our results showed that ILC2s expressed TH2/ILC2 surface marker genes, including CD44, CD81, and CD164 (Figure 2E), which regulate cell development, activation, growth, movement, adhesion and migration. ILC2s also expressed PTGDR2; the chemokine receptor genes CCR9; interleukin-6 receptor (IL-6R) and interleukin-9 receptor (IL-9R); the receptor gene Fcgr3a encoding the fc part of immunoglobulin g; and the IgE receptor encoding gene FCER1A. ILC2s express TH2 cytokines such as IL-4, IL-5 and IL-13 and express CSF2 (Figure 2F), which controls the production, differentiation and function of granulocytes and macrophages. ILC2s also express the cytokine signaling inhibitory gene SOCS3 and the chemokine gene CXCL14, which are involved in the homeostasis of macrophages. ILC2s significantly expressed GATA3, NFKB1, RORA, ZBTB16, YBX3, CREB3L1, BATF, KLF4 and other transcription factors (Figure 2G). ILC2s expressed the negative regulatory gene TP53I11, which is involved in cell population proliferation (Figure 2H) (Xiao et al., 2019); the TESC gene, which regulates molecular metabolic processes and bone marrow cell differentiation (Zhou et al., 2020b); the IFN-induced antiviral protein family genes IFITM1 and IFITM3; and NFKBIZ, which encodes the nuclear I kappa B protein. The gene sets enriched in ILC2s are related to intestinal immunity. A significant feature of intestinal immunity is that it can produce a large number of noninflammatory immunoglobulin A (IgA) antibodies, which can serve as the first line of defense against microorganisms (Figure S2D).

NK cells

NK cells of pigs expressed CD8a, CD11b (Gerner et al., 2009) and NCR1 (Mair et al., 2012) (Figure 2E). These cells express several genes (killer cell lectin-like receptor) of superfamily C, KLRB1, KLRFI and KLRK1; the IL-12 receptor gene IL-12RB2; and ADGRE5, which mediates cell–cell interactions. In terms of cytokines, we found that NK cells are similar to CD8 T cells/ILC1s (Figure 2F), both of which express NKG7 and the chemokine genes CCL4 and CCL5, which are involved in immune regulation and inflammatory processes (Malarkannan, 2020). In terms of transcription factors, NK cells expressed the transcription factors EOMES (Wagner et al., 2020; Zhang et al., 2018) and TBX21 (coexpressed by ILC1s and NK cells) (Figure 2G) (Fang et al., 2022; Spits et al., 2016); the transcription factors IKZF3 and IRF8 (Romero et al., 1999; Yamashita et al., 2021; Salem et al., 2014), which regulate lymphocyte development; KLF2, which is associated with early animal development (Basu et al., 2005); a gene that controls cell differentiation, namely, BHLHE40 (Sethuraman et al., 2018); and the apoptosis-related gene DDIT3 (Yamaguchi and Wang, 2004; Ohoka et al., 2005). On the other hand, NK cells also express PLAC8, PLEK, LYST, ISG15, GIMAP4, S100A1, CEP104 and other genes (Figure 2H), among which it has been reported that LYST can regulate the lysis of granules in cytotoxic T cells and NK cells (Gil-Krzewska et al., 2016). ISG15 is an important ubiquitin-like protein gene in

innate immunity (Swaim et al., 2017). The survival and development of GIMAP4T lymphocytes play an important regulatory role (Nitta et al., 2006). NK cells were shown to be enriched in pathways such as KEGG_SYSTEMIC_LUPUS_ERYTHEMATOSUS and KEGG_GLYCOSPHINGOLIPID_BIOSYNTHESIS_LACTO_AND_NEOLACTO_SERIES (Figure S2D).

According to the statistical results of interacting genes among ILC subsets, we found that there were 32 ligand binding relationships among ILC subsets (Figure S3A, B), the most significant of which was the obvious interaction between NK cell subsets and other ILCs, which also proved the important relationship between NK cells and antitumor, antiviral infection and immunomodulation (Figure S3C) (Poznanski and Ashkar, 2019; Bi and Wang, 2020). The obvious ligand CD74-MIF plays an important role in cell-mediated immunity, immune regulation and inflammation and can produce anti-inflammatory effects by inhibiting glucocorticoids and regulating macrophage function in host defense (Fan et al., 2013). EPHB6-EFNB1 may play a role in cell adhesion and the development or maintenance of the nervous system (Davis et al., 1994). The binding of TNF- and its receptor is involved in the response of cells to stimuli such as cytokines and stress and plays a key role in regulating the immune response to infection (Nie et al., 2013). C5AR1-RPS19 participates in the biochemical pathway of the innate and adaptive immune response-complement system (Robertson et al., 2018). CD44-HBEGF is involved in a variety of cellular functions, including activation, recycling and homing of T lymphocytes; hematopoiesis; inflammation; and response to bacterial infection (Vikesaa et al., 2006; Bourguignon et al., 2004).

3. Cluster analysis of ILC3s

To classify and analyze the subsets of ILC3s in more detail, we found that the ILC3 group can be divided into four subgroups (Figure 3A) according to their gene expression differences. The correlation analysis results show that ILC3b and ILC3c have a high correlation, while ILC3d has a low correlation with other ILC subgroups (Figure 3B). Gene expression profiles of the ILC3 gene list show the four subsets (Figure 3C) (Table S5), and significant differences in the transcription levels among the ILC3 subgroups were found (Figure 3D). The high expression of the heat shock protein genes HSPA6 and DNAJB1 (Figure 3E) in the ILC3a subset suggests that ILC3a cells may play an important role in maintaining the normal function of intestinal cells, thus regulating the balance between the survival and death of cells. The expression of inflammatory cytokine genes such as CXCL2, CXCL8 and IL-22 suggests that this cell subset plays an important role in the inflammatory response. Further GO enrichment analysis of ILC3a cells showed that they mainly play a regulatory role in BP (biological process) enrichment (Figure S2E) for entries such as positive regulation of multicellular organismal processes, positive regulation of biological processes, positive regulation of developmental processes, and regulation of molecular functions. There was no significant difference between ILC3b cells and high-response genes. ILC3c cells overexpress the receptor gene FCGR3A encoding the Fc of immunoglobulin G, which has the function of regulating NK cells (Lee et al., 2015; Srpan et al., 2018); IL-17F is an effector cytokine of the innate and adaptive immune system and participates in antibacterial host defense and the maintenance of tissue integrity (Puel et al., 2011; Goepfert et al., 2017). The protein encoded by the CD74 gene is related to class II major histocompatibility complex (MHC) and is an important chaperone for regulating antigen presentation of the immune response. ZBTB16 is an important transcriptional repressor gene involved in cell cycle processes and organism development (Figure 3F) (Constantinides et al., 2014). ILC3d cells highly expressed the tumor

suppressor genes SPINT2 and the involved cellular defense gene NCR2. Moreover, the important chemotactic receptor gene GPR183 was expressed in all ILC3 subsets (Figure 3G) (Benned-Jensen et al., 2012; Zhang et al., 2012). The results of GO enrichment analysis of the ILC3d subset show that it mainly plays a role in regulating the biological process and development of cells in BP, with enrichment (Figure S2F) of terms such as singular organic process, regulation of multicellular organic development, and negative regulation of biological process.

We have successfully described the typing as well as the localization of different cells in the pig gut from a transcriptomic landscape through the different expression of genes among different cells. However, in terms of protein expression, we still need to perform certain experiments to confirm the porcine intestinal innate immunity typing and to detect protein targets. Thus, a monoclonal antibody against porcine RORC was generated and validated by flow cytometry. Then, we detected the subsets of ILCs by flow cytometry (Figure S4A), which was consistent with the expression of scRNA-seq. ILC3s were most abundant among all ILCs in the lamina propria of the porcine jejunum, accounting for approximately 80%, while the proportions of ILC2s, ILC1s and NK cells were relatively low, and the expression of GATA3 in ILC3s was low. At the same time, through flow analysis of ILCs in the ileum, colon, lung and blood, we found that ILCs existed in various tissues of pigs, and the composition of ILCs in different parts was quite different. In the small intestine, including the jejunum and ileum, ILC3s were dominant, while ILC1s/NK cells were dominant in the lung and blood (Figure S4B).

4. Pseudotemporal analysis of ILCs

ILC subsets develop from specific progenitor cells (Harly et al., 2018); however, the fate of ILC subsets is not fixed. The environment regulates the transdifferentiation of ILCs through epigenetics (Shao et al., 2021), contributes to the plasticity of ILC subgroups, and changes their functional polarization to adapt their response to different tissues and different pathogenic stimuli (Bal et al., 2020). Our results show that there is transdifferentiation among the intestinal ILC subgroups of pigs under certain conditions, and there are different enrichment positions in the whole pseudotime-series trajectory, which suggests that there may be a transdifferentiation relationship among our ILC subgroups, among which most ILC3 subgroup cells are enriched in the early stage of the pseudotime-series trajectory, especially ILC3b subgroup cells, which are obviously enriched in the early stage of development. ILC1s, ILC2s and NK cells were enriched at the late stage of the pseudotime trajectory (Figure 4A), which indicated that ILC3s in our gut had high plasticity and could be transformed into ILC1s, ILC2s and NK cells. Because cells may differentiate into different types, there are many branches, and the differential genes between different branches may be the key to determining the fate of cell differentiation. Through the analysis of the branch nodes in the whole pseudotime sequence process, we found that there are three differentiation trends at the two differentiation nodes: the first trend is differentiation into ILC1 and NK cell subsets, the second trend is differentiation into the ILC2 subgroup, and the third trend is differentiation into the ILC3 subgroup. We express the differences in the differentiation of nodes in different directions in Figure 4B.

We found that the changes in some important landmark genes in each ILC subgroup during the pseudotime process, such as the highly expressed genes CCL5 and EOMES and other genes of NK cells and ILC1s, increased with the pseudotime response. The symbolic gene RORC of ILC3s changes with the pseudochronological process (Figure 4C). At the same time, we found some

important expression levels (Figure 4D) that regulate the transdifferentiation genes of ILCs. It has been reported that Aiolos (IKZF3) (a TGF-β imprinted molecule) and TBX21 jointly promote the transdifferentiation of ILC3s to ILC1s (Cella et al., 2019; Sciumé et al., 2012; Mazzurana et al., 2019). We found that IKZF3 and T-bet are mainly expressed by ILC1 and NK cells in ILCs. We found that the responses of IKZF3 and IKZF1 promoted the differentiation of ILC3s into ILC1s and NK cells and promoted the transcriptional response of related genes (Mazzurana et al., 2019). Vonarbourg et al. found that IL-12 and IL-15 accelerated the conversion of ILC3s into ILC1s (Vonarbourg et al., 2010). The C-Maf (CMIP) gene regulates ILC3s, and it can restrict the transdifferentiation of ILC3s to ILC1s (Parker et al., 2020). In our results, some ILC3 subgroups (ILC3a and ILC3c), ILC1s and ILC2s were mainly expressed CMIP. The IL-4 gene secreted by ILC2s can also promote the transdifferentiation of ILC3s into ILC2s and maintain the identity of ILC2s (Golebski et al., 2019). T-bet and Notch signals from NK cells and ILC1s can drive NCR⁻ ILC3s to NCR⁺ ILC3s for ILC3 subgroup cells (Chea et al., 2016; Rankin et al., 2013; Klose et al., 2013). TGFβ is expressed in ILCs, studies have shown that this gene can transform NCR⁺ ILC3s into NCR- ILC3s and can also induce ILC3s to differentiate into ILC regs and c-Kit⁻ ILC2s into c-Kit⁺ ILC2s (Viant et al., 2016; Verrier et al., 2016; Bernink et al., 2019). It has also been found that RORγt-dependent ILC1s may be transdifferentiated from ILC3s in the presence of IL-12 (Eisenring et al., 2010). Although our results show that there is transdifferentiation of ILCs in the porcine intestine, much research data may be needed to prove the specific transdifferentiation direction among ILC subgroups in the porcine intestine and the related molecules that mediate this process.

5. Porcine and human ILC clusters are transcriptionally conserved but distinct from ILC subsets in mice

Through the analysis of ILCs in the human intestinal lamina propria and those in the mouse intestinal lamina propria, we found that ILCs in the human intestinal tract were mainly ILC3s, but a small number of ILC1s were found, whereas no ILC2s or NK cells were found. Certain proportions of ILC2s and ILC3s were found in mice, but ILC1s and NK cells (Figure S5A) were not found. Cluster analysis of ILC subsets in the intestinal tract shows that ILC3s in the human intestinal tract can be divided into four subgroups, namely, ILC3a, ILC3b, ILC3c and ILC3d, and we show the genes with significant differential expression in each ILC3 subgroup and the ILC1 group (Figure S5B). ILCs in the porcine intestine are mainly ILC3s (Figure S5C). We found that the proportion of ILC3s in the intestinal tract of mice was still high, but there was also a high proportion of ILC2s (compared with humans and pigs). We showed the genes with significant differential expression in each ILC3 subgroup and the ILC2 group (Figure S5D).

To compare the transcriptional landscape of ILCs in the intestinal lamina propria among different species, we integrated our ILC dataset with published ILC scRNA-seq data from healthy and mice (Qi et al., 2021; Zhou et al., 2022). We found that the similarity of gene expression in ILCs of human-pig-mouse was high, with 9017 common genes. There were 1425 unique coexpressed genes between humans and pigs, 478 between humans and mice, and only 281 between pigs and mice. At the same time, we found that there were 842, 524 and 290 genes expressed by humans, pigs and mice, respectively (Figure 5A). Although we found that there was a high similarity among human-pig-mouse, the abundance of gene expression of ILCs within the three species was quite different (Figure 5B) (Table S6).

By performing unbiased cluster analysis between human and pig ILCs, we found that clusters

8 and 10 were NK and ILC1 subgroups, cluster 11 was the ILC2 subgroup, and the others were ILC3 subgroups (clusters 0, 1, 2, 3, 4, 5, 6, 7, and 9) (Figure 5C). Some ILCs clustered together with the pig intestinal tract, including ILC3 subgroup clusters 5 and 6, while human ILC1s clustered together with pig NK cells. We found that there were some similarities in the gene expression of ILCs between humans and pigs, such as the expression of the ILC signature genes IL-2RA, CCR6, KIT, RORC, and GATA3 (Figure 5D). At the same time, we found the differential expression of some genes, such as the expression of NCR1 in human ILC3s, which was greater than that of NCR2 in pig ILC3s (Figure 5E). And CD109 had high expression in pigs but hardly any expression in humans. The expression level of the IL-22 gene is similar to that of human ILC3s, but the expression level of IL-17 in porcine ILC3s is higher than that of humans. The expression level of the TAC3 gene in porcine ILC3s was very high, but there was no expression in human ILC3s.

For cluster 5, cluster 6 and cluster 8, which cluster humans and pigs together, we found that humans and pigs highly coexpressed genes in cluster 5 including HSPA6, BAG3, DNAJB4, ZFAND2A, CKS2 and others (Figure S6A) and had a higher enrichment in BP and CC (Figure S6B). Cohighly expressed genes in cluster 6 included ZFP36L1, ST8SIA4, F2R, SLC8A1, and TGFBR2 (Figure S6C), and there was a higher enrichment in BP and CC (Figure S6D). Cohighly expressed genes in cluster 8 included SAMD3, NKG7, IRF8, SYTL3, and ADGRE5 (Figure SE), and there was a higher enrichment in CC and MF (Figure S6F).

The intestinal ILCs of mice were detected by flow cytometry. After CD45$^+$LIN$^-$CD127$^+$ cells were gated, we detected the expression levels of RORγt and GATA3 and found that the ILC3:ILC2 ratio was approximately 2:1 (Figure 6A), which was consistent with the scRNA-seq data. The results of cluster analysis showed that the gene expression of ILCs between mice and pigs was quite different compared to humans versus pigs, and subgroups were divided into 12 clusters, but almost no common cluster subgroup was found. There were an NK subgroup (cluster 12), ILC1 subgroup (cluster 0), ILC2 subgroup (clusters 7, 8, and 11) and ILC3 subgroup (clusters 0, 1, 2, 3, 4, 5, 6, and 9) (Figure 6B). We found that the expression levels of the hallmark genes of ILCs between pigs and mice were similar, such as IL-7Rα, IL-23R, GATA3, RORC, and ID2. Among them, IL-23R and RORC were both expressed by ILC3s, while ILC2s expressed higher levels of GATA3 (Figure 6C). At the same time, we found that the cytokines IL-4, IL-5 and IL-13 in the ILC2 subgroup were more strongly expressed in mouse ILC2s, while CXCL14 was more strongly expressed in porcine ILC2s (Figure 6D). The transcription of the cytokines CXCL2, CXCL8, IL-22, TIMP1, TNF, ZBTB16 and other genes in the ILC3 subgroup was more intensively expressed in porcine ILC3s (Figure 6E). For example, CD90 (THY1), a member of the superfamily encoding cell surface glycoprotein and immunoglobulin, is only expressed in mouse ILCs. The expression level of AHR is the highest in mouse intestinal ILC2s. The mouse ILC3s highly expressed the chemokine gene CX3CL1, cytokine signaling inhibitor gene SOCS3 and chemokine receptor gene CCR7 (Figure 6F).

According to our human-pig-mouse coclustering analysis, the gene expression species similarity of intestinal ILCs in humans, pigs and mice was high. However, the gene expression abundance of ILCs varies greatly between species. For the clustering of ILCs between humans and pigs, we found common-clustered ILC3 subsets, and human ILC1s were also clustered together with porcine NK cells. However, the intestinal ILCs between pigs and mice were quite different and did not cluster obviously.

Discussion

We sorted 200,000 L/D⁻LIN⁻CD45⁺ cells from porcine jejunum LPLs by flow cytometry for scRNA-seq and finally obtained 15,763 translational landscapes after filtering. We found helper-like ILCs (including ILC1s, ILC2s and ILC3s), with results similar to Wiarda et al (Wiarda et al., 2022). Interestingly, we performed a more detailed analysis of ILCs in jejunal LPLs. We found that porcine intestinal ILCs mainly included four groups, namely, ILC3s, ILC1s, ILC2s and NK cells (sorted according to the proportion of cell number). Among them, ILC3s were the main ILC in the pig jejunal LPLs, and a further subset analysis of ILC3s based on the gene expression profiles revealed four subsets of ILC3s and identified transdifferentiation among subpopulations of ILCs. However, we did not find LTi cells in the intestinal ILCs of pigs, there were no clustered subgroups, and no gene list of LTA (the marker gene of human LTi cells) was found.

In the porcine jejunum, CD2, as a broad-spectrum and relatively stable whole T-cell antigen, is expressed not only in mature T cells and thymocytes but also in ILCs (Li et al., 2022a). The jejunal ILCs are mainly ILC3s and have signature transcribed genes similar to human genes such as IL-23R, hydrocarbon receptor (AHR) , IL-7Rα, NCR2, RORC and GATA3 (Figure 2E, G). Expression of the highly expressed aryl AHR of ILC3s can regulate the ILC balance and control chromatin accessibility at the AHR locus through positive feedback (Li et al., 2018). The host can also regulate the secretion of IL-22 by ILC3s by participating in the AHR pathway, thereby generating appropriate immunity against various pathogens (Li et al., 2019). AHR expression is also important for maintaining NKp46⁺ ILC3s (Xia et al., 2017). Studies have found that ILCs expressing NKp44 mainly develop into ILC3s (Nagasawa et al., 2019). IL-7-dependent maintenance of ILC3s is required for normal entry of lymphocytes into lymph nodes (Yang et al., 2018). Runx3 is required for the normal development of both ILC1s and ILC3s and controls the survival of ILC1s. Runx3 is also required for the expression of the transcription factor RORγt and its downstream target transcription factor AHR in ILC3s (Ebihara et al., 2015). At the same time, we found that ILC3s have rich subtypes, such as ILC3a with high expression of CXCL2 and IL-22, ILC3c with high expression of IL-17F, and ILC3d with high expression of CD74 (Figure 3C). We also found that ILC3s have the potential to differentiate into ILC2s, ILC1s, and NK cells. Although ILC3b cells have no obvious highly expressed genes, they have more obvious plasticity in the transdifferentiation among ILC subgroups. At the same time, we found some genes related to this process, such as IKZF1 and TGFB1, that were coexpressed by ILCs. IKZF3, TBX21 and Notch are expressed by ILC1s and NK cells, and CMIP is expressed by ILC3s, ILC2s and ILC1s. These genes play an important role in regulating the transdifferentiation of ILCs (Croft et al., 2022).

The difference between NK cells and ILC1s is usually the expression of the transcription factor exoderm protein (Eomes) (Artis and Spits, 2015; Almeida and Belz, 2016; Daussy et al., 2014). ILC1s and NK cells have overlapping phenotypes and functions, but they depend on different developmental pathways. NKG7, CCL5 and LITAF showed a higher expression in both ILC1 and NK (Figure 2F). The transcriptional repressor of ILC1s, MXI1, can negatively regulate MYC function and is therefore a potential tumor suppressor (Figure 2G). For the marker genes of NK such as LYST，ISG15 and GIMAP4T (Figure 2H), some studies have reported that LYST functions in regulating lysis granules in cytotoxic T cells and NK cells (Gil-Krzewska et al., 2016). ISG15 is a ubiquitin-like protein gene that is important in innate immunity (Swaim et al., 2017). GIMAP4T plays an important role in regulating the survival and development of lymphocytes

(Nitta et al., 2006). The hallmark receptor gene PTGDR2 for ILC2s can mediate the proinflammatory chemotaxis of eosinophils, basophils, and lymphocytes produced during allergic inflammation (Wojno et al., 2015; Pérez-Novo et al., 2010). IL-4, IL-5, IL-13, and CSF2, with high expression in ILC2s (Figure 2F), belong to the same gene cluster and are located on the chromosome 5q31 region, which is related to the differentiation of Th2 cells/ILC2s (Meyer et al., 2020). Among the transcription factors of ILC2s, GATA3 is an important regulator of the cell development of ILCs, and it is also very important for the development of ILC3s (Zhong et al., 2020; Kasal and Bendelac, 2021; Ren et al., 2022). RORA has been proven to contribute to the transcriptional regulation of some genes involved in circadian rhythm and is important for the development of ILC2s and ILC3s and the regulation of macrophages and Treg cells (Lee et al., 2021). ZBTB16 encodes a zinc finger transcription factor and participates in the progression of the cell cycle (Usui et al., 2021; Xu et al., 2019), the single-cell map of lung ILC2s shows that the tissue-resident ILC progenitor cells (ILCPs) and effector cells heterogeneously express ZBTB16 (Zeis et al., 2020). The YBX3 gene is involved in the negative regulation of the cell response to tumor necrosis factor and programmed cell death (Qin et al., 2020; Cooke et al., 2019). In steady-state tissues, ILC2s are more recognized as a major source of type 2 cytokines than Th2 cells and are a key driver of allergic reactions in the early stages of the disease (Kabata et al., 2018; Seehus et al., 2017).

Compared with pig and mouse ILCs, human and pig intestinal ILCs are transcriptionally conserved. ILCs were mainly ILC3-predominant in humans and porcine intestines, with an almost complete absence of ILC2s, while ILC2s were present in high proportions in mice. We also found some ILC marker genes that are common highly expressed in humans, pigs and mice. The transcription factor GATA3 is essential for the development of all ILCs, and studies have reported that GATA3 can not only control IL-7Rα but can also regulate the transcription factors TBX21 and RORC (Zhong et al., 2016). Some transcription factors, such as RORC, TBX21 and Notch (Figure 2G), are also essential for the differentiation of ILC subgroups. In the differentiation of human ILCs, some studies have found that the Notch→RORC→IL-23R pathway plays an important role, and these genes are expressed similarly in pig and human intestinal ILCs (Croft et al., 2022; Schroeder et al., 2022). At the same time, there is some specific expression of genes in porcine jejunum ILCs (Figure 5E). For example, in porcine ILCs, the NCR1 gene is only expressed in NK cells but not in other ILC subgroups, and porcine ILC3s showed a higher transcription level of the IL-17 gene. We also found the SLA1 gene, which specifically responds to porcine ILCs and regulates the signal transduction of the T-cell receptor (TCR). In summary, the results of cross-species analysis and comparison of ILCs have laid a foundation for further exploration of pigs as a human medical model (Hryhorowicz et al., 2020; Käser, 2021; Pabst, 2020; Sinkora and Butler, 2016). We found a high similarity of intestinal ILCs between pigs and humans, and this finding confirms the feasibility of pigs as a human surrogate model for studies related to intestinal mucosal immunity. Especially in research concerning the cell target of ILC-related and intestinal mucosal immune diseases such as enteritis and inflammatory bowel disease (IBD), pigs have a natural advantage as model animals.

Studies have reported the existence of regulatory relationships such as the gut-brain axis (Li et al., 2022c), gut-lung axis (Wypych et al., 2019) and gut-kidney axis (Xie et al., 2022). In our results, porcine ILC3s was found to express TAC3 gene. The TAC3 gene encodes neurokinin B (NKB), whose receptor is NK3R (Young et al., 2010), and the combination of NKB and NK3R

can promote gonadotrophin-releasing hormone (GnRH) release (Billings et al., 2010). GnRH binds to specific receptors on the pituitary gonadotropin to promote the release of sex hormones. Therefore, TAC3 play an important role in the gonadotrope axis, We proposed the gut-gonadal axis, where porcine intestinal ILC3s regulate gonadal development.

Our research on ILCs in porcine intestines can promote the intestinal health of pigs, and at the same time, it is of great significance to biomedical research, animal models and global food health. However, due to the limited research on the intestinal tract of pigs, especially ILCs, a new lymphocyte family, there is not enough biological information at present. To the best of our knowledge, our discovery is the first single-cell description of the ILC transcriptome landscape in the pig jejunum, focusing on research on ILCs in the jejunum. In addition, other lymphocyte types in the intestine also play an important role in intestinal health, but they are outside the scope of this study. Our work is based on the intestinal health of the jejunum, identifying the transcription information of ILCs in the jejunum by scRNA-seq and then inferring the functional subgroup. This work also provides more biological knowledge related to this species of pig.

Although our data can be used as favorable evidence for the existence of ILCs in porcine intestines and can help us to explore which genes the ILCs express to play an immunomodulatory role and how they transdifferentiate under complex environmental conditions in the intestines, this understanding is only applicable to the intestines at present, while the roles of ILCs in other tissues of pigs are still unclear. Therefore, our data provide an exciting starting point for the study of porcine ILCs, and it is necessary to further analyze ILCs in other parts of pigs in the future to analyze the tissue heterogeneity and developmental characteristics of porcine ILCs.

Limitations of the study

Although this study provided a comprehensive transcription analysis of ILCs in the lamina propria of the porcine intestine, we only analyzed single age/sex pigs. In this study, to obtain more ILCs, we enriched lymphocytes in the lamina propria with Lin⁻, which led to the loss of a large number of T/B lymphocyte groups and nonlymphocyte groups. For example, the CD8 and CD4 antibodies we used could lead to the loss of NK and LTi cells. In addition, although we found that there is a great transcription difference between the porcine intestinal ILC subgroup and mouse intestinal ILC subgroup, there is a high similarity with human intestinal ILCs, and we cannot rule out that the differences among human, porcine and mouse species are due to biological and technical effects, such as differences in physiological age, tissue preparation methods and sequencing saturation. In the future, if more reagents or better methods are available, we should use a variety of methods to verify our findings.

Materials and Methods
Animals and sample collection

All Specific pathogen Free (SPF) experimental pigs were purchased from Harbin Veterinary Research Institute, and all piglets were weaned at 3 weeks old. When piglets were 4 weeks old, we used flow cytometry to analyze the immune cells of the jejunum lamina propria. Then, we collected samples from the jejunum of 4 SPF female pigs, collected cell samples by flow sorting, performed scRNA-seq, and made sections of tissue samples for immunofluorescence experiments. All animal experiments met the requirements of the Animal Management and Ethics Committee of Jilin Agricultural University.

Cell separation

For cell isolation of the jejunum, immediately after euthanasia of piglets, approximately 5 cm of the jejunum was collected and stored in phosphate buffered saline (PBS) at 4°C. The outer muscle layer was peeled off, and the intestine was cut longitudinally to expose the lumen. The tissue was rinsed gently with PBS to remove the intestinal contents and cut into 1-cm large and small intestine segments with scissors.

The tissue was transferred to the separation solution (15 mL of RPMI-1640, 1% double antibody, 1% HEPES, 5 mM EDTA, 2 mM DTT, and 2% heat-inactivated fetal bovine serum (FCS)) and incubated in a shaking incubator at 37°C and 200 rpm for 28 minutes. After shaking violently for 2 minutes, the remaining intestinal tissue was retained, the washing liquid was removed, and PBS at 4°C was used for washing.

The remaining tissues were transferred to the digestive juice solution (8 mL RPMI-1640 medium, 1% double antibody, 1% HEPES, 50 mg collagenase IV, 1 mg DNase I, and 2% FCS) and incubated in a shaking incubator at 37°C and 250 rpm for 20 minutes. Digestion was continued for 20 minutes, after which the samples were shaken vigorously for 2 minutes, and then the digestive juice was collected through a 70-µm cell membrane.

Culture solution (10 ml RPMI-1640, 5% FCS) was added to the collected digestive juice, centrifuged at 4°C and 450xg for 8 minutes, resuspended in 40% Percoll solution, then slowly added to 80% Percoll solution, and centrifuged at 24°C and 500xg.

Antibodies staining

Monoclonal antibody against porcine RORC for flow cytometry was produced by Jieyu Company (RORC Gene ID: 100622477) (clone 11F4). Hybridoma cells were preserved by the China Center for Type Culture Collection (CCTCC) (No. C2022219), and related products have been patented (No. 202210980598. X).

Antibodies for pigs were as follows: Invitrogen: CD2 monoclonal antibody (14-0029-82), CD3e monoclonal antibody (MA5-28774), CD3e monoclonal antibody (Biotin) (MA5-28771), CD11b monoclonal antibody (MA5-16604), CD11c monoclonal antibody (APC) (17-0116-42), CD21 monoclonal antibody (A1-19243), CD21 monoclonal antibody (PE) (MA1-19754), CD45 monoclonal antibody (FITC) (MA5-28383), CD117 (c-Kit) monoclonal antibody (APC) (17-1171-82), CD163 monoclonal antibody (PE) (MA5-16476), CD172a monoclonal antibody (MA5-28299), CD335 (NKp46/NCR1) monoclonal antibody (PE) (MA5-28352), SLA Class II DR monoclonal antibody (MA5-28503), GATA3 monoclonal antibody (PE-Cyanine7) (25-9966-42), T-bet monoclonal antibody (PE) (12-5825-82), and goat anti-mouse IgG H&L (PE-Cyanine5) (M35018).

BD Pharmingen: Fixable Viability Stain 780 (565388), purified rat anti-mouse CD16/CD32 (Mouse BD Fc Block) (553142), mouse anti-pig CD3ε (PE) (561485), mouse anti-pig CD4a (PE) (559586), mouse anti-pig CD8a (FITC) (551543), mouse anti-pig CD8a (Alexa Fluor 647) (561475), rat anti-pig γδ T Lymphocytes (PE) (551543), rat anti-pig γδ T lymphocytes (561486), and mouse anti-GATA3 (APC) (558686).

Abcam: goat anti-mouse IgG H&L (FITC) (ab6785), goat F(ab')2 anti-mouse IgG H&L (PE) (ab7002), goat F(ab')2 anti-rat IgG H&L (PE-Cyanine5) (ab130803), and rat anti-mouse IgG1 H&L (PE) (ab99605).

BioLegend: CD11b monoclonal antibody (APC) (101212), CD11c monoclonal antibody (PE) (301606).

R&D: porcine CD34 antibody (AF3890).

Antibodies for mouse were as follows: BD: Fixable Viability Stain 780 (565388), CD11b (Biotin) (557395), γδ T (Biotin) (553176), CD19 (Biotin) (553784), TCRβ (Biotin) (553168), Ly6G/C (Biotin) (553124), TER-119 (Biotin) (553672), streptavidin protein (APC-cy7) (554063), CD45 (FITC) (551874), CD127 (PE-cy7) (560733), RORγt (PE) (562607), and GATA3 (Alexa Fluor 647) (560068).

scRNA-seq

Enriched ILCs were obtained by flow sorting. We stained the total LPL extracted from the lamina propria with antibodies against LIFE, CD45, lin1 (CD3, CD21, γδT, CD11c, CD172a) and lin2 (CD4, CD8, CD163, CD11b) and then selected LIFE-CD45+Lin1-Lin2- cells by flow sorting. The concentration was 900 cells/μL, the cell viability was > 75, the total number of cells was adjusted to approximately $5x10^6$, and then SCRNA-SEQ was performed.

The cell suspension was loaded into Chromium microfluidic chips with 30 chemistry and barcoded with a 10× Chromium Controller (10× Genomics). RNA from the barcoded cells was subsequently reverse-transcribed, and sequencing libraries were constructed with reagents from a Chromium Single Cell 30 reagent kit (10× Genomics) according to the manufacturer's instructions. Sequencing was performed with Illumina (Novaseq PE150) according to the manufacturer's instructions (Illumina).

Data analysis

We used FastQC to perform basic statistics on the quality of the raw reads. Generally, celltanger count supports FASTQ files from raw base call (BCL) files generated by Illumina sequencers as input files. 10× Genomics® does not recommend additional processing of the sequence.

Raw reads were demultiplexed and mapped to the reference genome by the 10× Genomics Cell Ranger pipeline (https://support.10xgenomics.com/single-cell-geneexpression/software/pipelines/latest/what-is-cell-ranger) using default parameters. All downstream single-cell analyses were performed using Cell Ranger and Seurat unless mentioned specifically. In brief, for each gene and each cell barcode (filtered by CellRanger), unique molecule identifiers were counted to construct digital expression matrices. For secondary filtration by Seurat, a gene with expression in more than 3 cells was considered expressed, and each cell was required to have at least 200 expressed genes. Then, some of the foreign cells were filtered out.

Secondary analysis of gene expression

Cellranger reanalyze takes feature-barcode matrices produced by cellranger count or cellranger aggr and reruns the dimensionality reduction, clustering, and gene expression algorithms using cellranger default parameter settings.

The Seurat package was used to perform data normalization, dimensionality reduction, clustering, and differential expression. We used the Seurat alignment method canonical correlation analysis (CCA) for integrated analysis of datasets. For clustering, highly variable genes were selected, and the principal components based on those genes were used to build a graph, which was segmented with a resolution of 0.6 (Stuart et al., 2019).

To perform global analysis between samples, based on the filtered gene expression matrix by Seurat, between-samples differential expression analysis was carried out using the edgeR package to obtain zone-specific marker genes (McCarthy et al., 2012).

For enrichment analysis of marker genes, Reactome pathway-based analysis of marker genes was implemented by the ReactomePA R package. REACTOME is an open-source, open access, manually curated and peer-reviewed pathway database (https://reactome.org/).

The GSVA analysis was performed using the msigdb database (https://www.gsea-msigdb.org/gsea/msigdb/index.jsp)c2.cp.kegg.v7.4.symbols.gmt.

The results from CellDB were presented for the cell type interaction network using iTALK software.

For trajectory analysis, Monocle2 was used to calculate the correlation of cells to obtain the minimum spanning tree, find the minimum path, project all other data points to the minimum path, and finally obtain the cell differentiation trajectory diagram (Qiu et al., 2017).

Dataset integration

The porcine jejunal scRNA-seq data was derived from GSE175411 (Fan et al., 2022). We compared our data with published human intestinal ILC (GSA-Human: HRA000919) (Qi et al., 2021) and mouse intestinal ILC (GSE166266) (Zhou et al., 2022) datasets. The Ensembl genome browser (Ensembl Genes 105) was used to convert human (GRCh38) and mouse (GRCm39) gene names to the corresponding pig gene names prior to integration (https://www.ensembl.org/biomart/martview/). The indiscriminate clustering algorithm was analyzed later (Gu et al., 2022).

DATA AVAILABILITY STATEMENT

Data and scripts will be available upon publication.

AUTHOR CONTRIBUTIONS

Cell isolation, J.H.W. and M.Y.C.; data analysis, M.G, Y.S. and Y.Y.L ; manuscript preparation and writing, J.H.W., Y.Z., X.X.L., and C.W.S.; supervision and project administration, C.F.W. and X.C. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (32273043, 32202890, U21A20261)，the Science and Technology Development Program of Changchun City (21ZY42) the Science and Technology Development Program of Jilin Province (20200402041NC), and China Agriculature Research System of MOF and MARA (CARS-35). We thank the Novogene Co., LTD for providing technical services.

ETHICS

The animal management procedures and all laboratory procedures abided by the regulations of the Animal Care and Ethics Committees of Jilin Agriculture University, China.

REFERENCES

Abdullahi, A., Amini-Nik, S., Jeschke, M., 2014. Animal models in burn research. Cellular and molecular life sciences : CMLS 71(17), 3241-3255.

Almeida, F., Belz, G., 2016. Innate lymphoid cells: models of plasticity for immune homeostasis and rapid responsiveness in protection. Mucosal immunology 9(5), 1103-1112.

Artis, D., Spits, H., 2015. The biology of innate lymphoid cells. Nature 517(7534), 293-301.

Bal, S., Golebski, K., Spits, H., 2020. Plasticity of innate lymphoid cell subsets. Nature reviews. Immunology 20(9), 552-565.

Basu, P., Morris, P., Haar, J., Wani, M., Lingrel, J., Gaensler, K., Lloyd, J., 2005. KLF2 is essential for primitive erythropoiesis and regulates the human and murine embryonic beta-like globin genes in vivo. Blood 106(7), 2566-2571.

Benned-Jensen, T., Norn, C., Laurent, S., Madsen, C., Larsen, H., Arfelt, K., Wolf, R., Frimurer, T., Sailer, A., Rosenkilde, M., 2012. Molecular characterization of oxysterol binding to the Epstein-Barr virus-induced gene 2 (GPR183). The Journal of biological chemistry 287(42), 35470-35483.

Bernink, J., Krabbendam, L., Germar, K., de Jong, E., Gronke, K., Kofoed-Nielsen, M., Munneke, J., Hazenberg, M., Villaudy, J., Buskens, C., Bemelman, W., Diefenbach, A., Blom, B., Spits, H., 2015. Interleukin-12 and -23 Control Plasticity of CD127(+) Group 1 and Group 3 Innate Lymphoid Cells in the Intestinal Lamina Propria. Immunity 43(1), 146-160.

Bernink, J., Ohne, Y., Teunissen, M., Wang, J., Wu, J., Krabbendam, L., Guntermann, C., Volckmann, R., Koster, J., van Tol, S., Ramirez, I., Shrestha, Y., de Rie, M., Spits, H., Romero Ros, X., Humbles, A., 2019. c-Kit-positive ILC2s exhibit an ILC3-like signature that may contribute to IL-17-mediated pathologies. Nature immunology 20(8), 992-1003.

Besgen, P., Trommler, P., Vollmer, S., Prinz, J., 2010. Ezrin, maspin, peroxiredoxin 2, and heat shock protein 27: potential targets of a streptococcal-induced autoimmune response in psoriasis. Journal of immunology (Baltimore, Md. : 1950) 184(9), 5392-5402.

Bi, J., Wang, X., 2020. Molecular Regulation of NK Cell Maturation. Frontiers in immunology 11, 1945.

Bianco, S., Kaiser, U., 2009. The genetic and molecular basis of idiopathic hypogonadotropic hypogonadism. Nature reviews. Endocrinology 5(10), 569-576.

Billings, H.J., Connors, J.M., Altman, S.N., Hileman, S.M., Holaskova, I., Lehman, M.N., McManus, C.J., Nestor, C.C., Jacobs, B.H., Goodman, R.L., 2010. Neurokinin B Acts via the Neurokinin-3 Receptor in the Retrochiasmatic Area to Stimulate Luteinizing Hormone Secretion in Sheep. Endocrinology 151(8), 3836-3846.

Bourguignon, L., Singleton, P., Diedrich, F., 2004. Hyaluronan-CD44 interaction with Rac1-dependent protein kinase N-gamma promotes phospholipase Cgamma1 activation, Ca(2+) signaling, and cortactin-cytoskeleton function leading to keratinocyte adhesion and differentiation. The Journal of biological chemistry 279(28), 29654-29669.

Cella, M., Gamini, R., Sécca, C., Collins, P., Zhao, S., Peng, V., Robinette, M., Schettini, J., Zaitsev, K., Gordon, W., Bando, J., Yomogida, K., Cortez, V., Fronick, C., Fulton, R., Lin, L., Gilfillan, S., Flavell, R., Shan, L., Artyomov, M., Bowman, M., Oltz, E., Jelinsky, S., Colonna, M., 2019. Subsets of ILC3-ILC1-like cells generate a diversity spectrum of innate lymphoid cells in human mucosal tissues. Nature immunology 20(8), 980-991.

Chea, S., Perchet, T., Petit, M., Verrier, T., Guy-Grand, D., Banchi, E., Vosshenrich, C., Di Santo, J., Cumano, A., Golub, R., 2016. Notch signaling in group 3 innate lymphoid cells modulates their plasticity. Science signaling 9(426), ra45.

Constantinides, M., McDonald, B., Verhoef, P., Bendelac, A., 2014. A committed precursor to innate lymphoid cells. Nature 508(7496), 397-401.

Cooke, A., Schwarzl, T., Huppertz, I., Kramer, G., Mantas, P., Alleaume, A., Huber, W., Krijgsveld, J., Hentze, M., 2019. The RNA-Binding Protein YBX3 Controls Amino Acid Levels by Regulating SLC mRNA Abundance. Cell reports 27(11), 3097-3106.e3095.

Croft, C., Thaller, A., Marie, S., Doisne, J., Surace, L., Yang, R., Puel, A., Bustamante, J., Casanova, J., Di Santo, J., 2022. Notch, RORC and IL-23 signals cooperate to promote multi-lineage human innate lymphoid cell differentiation. Nature communications 13(1), 4344.

Croxatto, D., Micheletti, A., Montaldo, E., Orecchia, P., Loiacono, F., Canegallo, F., Calzetti, F., Fulcheri, E., Munari, E., Zamò, A., Venturini, P., Moretta, L., Cassatella, M., Mingari, M., Vacca, P., 2016. Group 3 innate lymphoid cells regulate neutrophil migration and function in human decidua. Mucosal immunology 9(6), 1372-1383.

Daussy, C., Faure, F., Mayol, K., Viel, S., Gasteiger, G., Charrier, E., Bienvenu, J., Henry, T., Debien, E., Hasan, U., Marvel, J., Yoh, K., Takahashi, S., Prinz, I., de Bernard, S., Buffat, L., Walzer, T., 2014. T-bet and Eomes instruct the development of two distinct natural killer cell lineages in the liver and in the bone marrow. The Journal of experimental medicine 211(3), 563-577.

Davis, S., Gale, N., Aldrich, T., Maisonpierre, P., Lhotak, V., Pawson, T., Goldfarb, M., Yancopoulos, G., 1994. Ligands for EPH-related receptor tyrosine kinases that require membrane attachment or clustering for activity. Science (New York, N.Y.) 266(5186), 816-819.

Diefenbach, A., Gnafakis, S., Shomrat, O., 2020. Innate Lymphoid Cell-Epithelial Cell Modules Sustain Intestinal Homeostasis. Immunity 52(3), 452-463.

Ebihara, T., Song, C., Ryu, S., Plougastel-Douglas, B., Yang, L., Levanon, D., Groner, Y., Bern, M., Stappenbeck, T., Colonna, M., Egawa, T., Yokoyama, W., 2015. Runx3 specifies lineage commitment of innate lymphoid cells. Nature immunology 16(11), 1124-1133.

Eisenring, M., vom Berg, J., Kristiansen, G., Saller, E., Becher, B., 2010. IL-12 initiates tumor rejection via lymphoid tissue-inducer cells bearing the natural cytotoxicity receptor NKp46. Nature immunology 11(11), 1030-1038.

Fachi, J., Sécca, C., Rodrigues, P., Mato, F., Di Luccia, B., Felipe, J., Pral, L., Rungue, M., Rocha, V., Sato, F., Sampaio, U., Clerici, M., Rodrigues, H., Câmara, N., Consonni, S., Vieira, A., Oliveira, S., Mackay, C., Layden, B., Bortoluci, K., Colonna, M., Vinolo, M., 2020. Acetate coordinates neutrophil and ILC3 responses against C. difficile through FFAR2. The Journal of experimental medicine 217(3).

Fan, B., Zhou, J., Zhao, Y., Zhu, X., Zhu, M., Peng, Q., Li, J., Chang, X., Shi, D., Yin, J., Guo, R., Li, Y., He, K., Fan, H., Li, B., 2022. Identification of Cell Types and Transcriptome Landscapes of Porcine Epidemic Diarrhea Virus-Infected Porcine Small Intestine Using Single-Cell RNA Sequencing. Journal of immunology (Baltimore, Md. : 1950).

Fan, C., Rajasekaran, D., Syed, M., Leng, L., Loria, J., Bhandari, V., Bucala, R., Lolis, E., 2013. MIF intersubunit disulfide mutant antagonist supports activation of CD74 by endogenous MIF trimer at physiologic concentrations. Proceedings of the National Academy of Sciences of the United States of America 110(27), 10994-10999.

Fang, D., Cui, K., Cao, Y., Zheng, M., Kawabe, T., Hu, G., Khillan, J., Li, D., Zhong, C., Jankovic, D., Sher, A., Zhao, K., Zhu, J., 2022. Differential regulation of transcription factor T-bet induction during NK cell development and T helper-1 cell differentiation. Immunity 55(4), 639-655.e637.

Furue, M., Furue, K., Tsuji, G., Nakahara, T., 2020. Interleukin-17A and Keratinocytes in Psoriasis. International journal of molecular sciences 21(4).

Georgiev, H., Ravens, I., Papadogianni, G., Bernhardt, G., 2018. Coming of Age: CD96 Emerges as Modulator of Immune Responses. Frontiers in immunology 9, 1072.

Gerner, W., Käser, T., Saalmüller, A., 2009. Porcine T lymphocytes and NK cells--an update. Developmental and comparative immunology 33(3), 310-320.

Gil-Krzewska, A., Wood, S., Murakami, Y., Nguyen, V., Chiang, S., Cullinane, A., Peruzzi, G., Gahl, W., Coligan, J., Introne, W., Bryceson, Y., Krzewski, K., 2016. Chediak-Higashi syndrome: Lysosomal trafficking regulator domains regulate exocytosis of lytic granules but not cytokine secretion by natural killer cells. The Journal of allergy and clinical immunology 137(4), 1165-1177.

Goepfert, A., Lehmann, S., Wirth, E., Rondeau, J., 2017. The human IL-17A/F heterodimer: a two-faced cytokine with unique receptor recognition properties. Scientific reports 7(1), 8906.

Golebski, K., Ros, X., Nagasawa, M., van Tol, S., Heesters, B., Aglmous, H., Kradolfer, C., Shikhagaie, M., Seys, S., Hellings, P., van Drunen, C., Fokkens, W., Spits, H., Bal, S., 2019. IL-1β, IL-23, and TGF-β drive plasticity of human ILC2s towards IL-17-producing ILCs in nasal inflammation. Nature communications 10(1), 2162.

Gu, W., Madrid, D., Joyce, S., Driver, J., 2022. A single-cell analysis of thymopoiesis and thymic iNKT cell development in pigs. Cell reports 40(1), 111050.

Güller, M., Toualbi-Abed, K., Legrand, A., Michel, L., Mauviel, A., Bernuau, D., Daniel, F., 2008. c-Fos overexpression increases the proliferation of human hepatocytes by stabilizing nuclear Cyclin D1. World journal of gastroenterology 14(41), 6339-6346.

Gurram, R., Zhu, J., 2019. Orchestration between ILC2s and Th2 cells in shaping type 2 immune responses. Cellular & molecular immunology 16(3), 225-235.

Halim, T., Takei, F., 2014. Isolation and characterization of mouse innate lymphoid cells. Current protocols in immunology 106, 3.25.21-23.25.13.

Han, T., Lee, T., Ledwon, J., Vaca, E., Turin, S., Kearney, A., Gosain, A., Tepole, A., 2022. Bayesian calibration of a computational model of tissue expansion based on a porcine animal model. Acta biomaterialia 137, 136-146.

Harly, C., Cam, M., Kaye, J., Bhandoola, A., 2018. Development and differentiation of early innate lymphoid progenitors. The Journal of experimental medicine 215(1), 249-262.

Harly, C., Kenney, D., Ren, G., Lai, B., Raabe, T., Yang, Q., Cam, M., Xue, H., Zhao, K., Bhandoola, A., 2019. The transcription factor TCF-1 enforces commitment to the innate lymphoid cell lineage. Nature immunology 20(9), 1150-1160.

Herrera-Uribe, J., Wiarda, J., Sivasankaran, S., Daharsh, L., Liu, H., Byrne, K., Smith, T., Lunney, J., Loving, C., Tuggle, C., 2021. Reference Transcriptomes of Porcine Peripheral Immune Cells Created Through Bulk and Single-Cell RNA Sequencing. Frontiers in genetics 12, 689406.

Hidalgo, L., Einecke, G., Allanach, K., Halloran, P., 2008. The transcriptome of human cytotoxic T cells: similarities and disparities among allostimulated CD4(+) CTL, CD8(+) CTL and NK cells. American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons 8(3), 627-636.

Horne, A., Ramme, K., Rudd, E., Zheng, C., Wali, Y., al-Lamki, Z., Gürgey, A., Yalman, N., Nordenskjöld, M., Henter, J., 2008. Characterization of PRF1, STX11 and UNC13D genotype-phenotype correlations in familial hemophagocytic lymphohistiocytosis. British journal of haematology 143(1), 75-83.

Hryhorowicz, M., Lipiński, D., Hryhorowicz, S., Nowak-Terpiłowska, A., Ryczek, N., Zeyland, J., 2020. Application of Genetically Engineered Pigs in Biomedical Research. Genes 11(6).

Jin, Y., Cao, X., Shi, C., Feng, B., Huang, H., Jiang, Y., Wang, J., Yang, G., Yang, W., Wang, C., 2021. Lactobacillus rhamnosus GG Promotes Early B Lineage Development and IgA Production in the Lamina Propria in Piglets. Journal of immunology (Baltimore, Md. : 1950) 207(8), 2179-2191.

Kabata, H., Moro, K., Koyasu, S., 2018. The group 2 innate lymphoid cell (ILC2) regulatory network and its underlying mechanisms. Immunological reviews 286(1), 37-52.

Kasal, D., Bendelac, A., 2021. Multi-transcription factor reporter mice delineate early precursors to the ILC and LTi lineages. The Journal of experimental medicine 218(2).

Käser, T., 2021. Swine as biomedical animal model for T-cell research-Success and potential for transmittable and non-transmittable human diseases. Molecular immunology 135, 95-115.

Klose, C., Kiss, E., Schwierzeck, V., Ebert, K., Hoyler, T., d'Hargues, Y., Göppert, N., Croxford, A., Waisman, A., Tanriver, Y., Diefenbach, A., 2013. A T-bet gradient controls the fate and function of CCR6-RORγt+ innate lymphoid cells. Nature 494(7436), 261-265.

Kong, Q., Yang, X., Zhang, H., Liu, S., Zhao, J., Zhang, J., Weng, X., Jin, J., Liu, Z., 2020. Lineage specification and pluripotency revealed by transcriptome analysis from oocyte to blastocyst in pig. FASEB journal : official publication of the Federation of American Societies for Experimental Biology 34(1), 691-705.

Lamber, E., Vanhille, L., Textor, L., Kachalova, G., Sieweke, M., Wilmanns, M., 2008. Regulation of the transcription factor Ets-1 by DNA-mediated homo-dimerization. The EMBO journal 27(14), 2006-2017.

Lee, J., Kim, H., Baek, S., 2021. Unraveling the physiological roles of retinoic acid receptor-related orphan receptor α. Experimental & molecular medicine 53(9), 1278-1286.

Lee, J., Zhang, T., Hwang, I., Kim, A., Nitschke, L., Kim, M., Scott, J., Kamimura, Y., Lanier, L., Kim, S., 2015. Epigenetic modification and antibody-dependent expansion of memory-like NK cells in human cytomegalovirus-infected individuals. Immunity 42(3), 431-442.

Li, B., Lu, Y., Zhong, M., Qian, J., Li, R., Davidson, D., Tang, Z., Zhu, K., Argenty, J., de Peredo, A., Malissen, B., Roncagalli, R., Veillette, A., 2022a. Cis interactions between CD2 and its ligands on T cells are required for T cell activation. Science immunology 7(74), eabn6373.

Li, S., Bostick, J., Ye, J., Qiu, J., Zhang, B., Urban, J., Avram, D., Zhou, L., 2018. Aryl Hydrocarbon Receptor Signaling Cell Intrinsically Inhibits Intestinal Group 2 Innate Lymphoid Cell Function. Immunity 49(5), 915-928.e915.

Li, Y., Lv, C., Yu, Y., Wu, B., Zhang, Y., Lang, Q., Liang, Z., Zhong, C., Shi, Y., Han, S., Xu, F., Tian, Y., 2022b. KIR3DL3-HHLA2 and TMIGD2-HHLA2 pathways: The dual role of HHLA2 in immune responses and its potential therapeutic approach for cancer immunotherapy. Journal of advanced research.

Li, Y., Xie, H., Zhang, W., Wei, Y., Sha, R., Xu, L., Zhang, J., Jiang, Y., Guo, T., Zhao, B., 2019. Type 3 innate lymphoid cells are altered in colons of C57BL/6 mice with dioxin exposure. The Science of the total environment 662, 639-645.

Li, Z., Lai, J., Zhang, P., Ding, J., Jiang, J., Liu, C., Huang, H., Zhen, H., Xi, C., Sun, Y., Wu, L., Wang, L., Gao, X., Li, Y., Fu, Y., Jie, Z., Li, S., Zhang, D., Chen, Y., Zhu, Y., Lu, S., Lu, J., Wang, D., Zhou, H., Yuan, X., Li, X., Pang, L., Huang, M., Yang, H., Zhang, W., Brix, S., Kristiansen, K., Song, X., Nie, C., Hu,

S., 2022c. Multi-omics analyses of serum metabolome, gut microbiome and brain function reveal dysregulated microbiota-gut-brain axis in bipolar depression. Molecular psychiatry.

Liu, B., Ye, B., Zhu, X., Yang, L., Li, H., Liu, N., Zhu, P., Lu, T., He, L., Tian, Y., Fan, Z., 2020. An inducible circular RNA circKcnt2 inhibits ILC3 activation to facilitate colitis resolution. Nature communications 11(1), 4076.

Liu, T., Li, J., Yu, L., Sun, H., Li, J., Dong, G., Hu, Y., Li, Y., Shen, Y., Wu, J., Gu, Y., 2021. Cross-species single-cell transcriptomic analysis reveals pre-gastrulation developmental differences among pigs, monkeys, and humans. Cell discovery 7(1), 8.

Mair, K., Essler, S., Patzl, M., Storset, A., Saalmüller, A., Gerner, W., 2012. NKp46 expression discriminates porcine NK cells with different functional properties. European journal of immunology 42(5), 1261-1271.

Malarkannan, S., 2020. NKG7 makes a better killer. Nature immunology 21(10), 1139-1140.

Mazzurana, L., Forkel, M., Rao, A., Van Acker, A., Kokkinou, E., Ichiya, T., Almer, S., Höög, C., Friberg, D., Mjösberg, J., 2019. Suppression of Aiolos and Ikaros expression by lenalidomide reduces human ILC3-ILC1/NK cell transdifferentiation. European journal of immunology 49(9), 1344-1355.

McCarthy, D., Chen, Y., Smyth, G., 2012. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic acids research 40(10), 4288-4297.

Meyer, A., Engevik, A., Madorsky, T., Belmont, E., Stier, M., Norlander, A., Pilkinton, M., McDonnell, W., Weis, J., Jang, B., Mallal, S., Peebles, R., Goldenring, J., 2020. Group 2 Innate Lymphoid Cells Coordinate Damage Response in the Stomach. Gastroenterology 159(6), 2077-2091.e2078.

Mortha, A., Chudnovskiy, A., Hashimoto, D., Bogunovic, M., Spencer, S., Belkaid, Y., Merad, M., 2014. Microbiota-dependent crosstalk between macrophages and ILC3 promotes intestinal homeostasis. Science (New York, N.Y.) 343(6178), 1249288.

Nagasawa, M., Heesters, B., Kradolfer, C., Krabbendam, L., Martinez-Gonzalez, I., de Bruijn, M., Golebski, K., Hendriks, R., Stadhouders, R., Spits, H., Bal, S., 2019. KLRG1 and NKp46 discriminate subpopulations of human CD117CRTH2 ILCs biased toward ILC2 or ILC3. The Journal of experimental medicine 216(8), 1762-1776.

Nie, H., Zheng, Y., Li, R., Guo, T., He, D., Fang, L., Liu, X., Xiao, L., Chen, X., Wan, B., Chin, Y., Zhang, J., 2013. Phosphorylation of FOXP3 controls regulatory T cell function and is inhibited by TNF-α in rheumatoid arthritis. Nature medicine 19(3), 322-328.

Nitta, T., Nasreen, M., Seike, T., Goji, A., Ohigashi, I., Miyazaki, T., Ohta, T., Kanno, M., Takahama, Y., 2006. IAN family critically regulates survival and development of T lymphocytes. PLoS biology 4(4), e103.

Ohoka, N., Yoshii, S., Hattori, T., Onozaki, K., Hayashi, H., 2005. TRB3, a novel ER stress-inducible gene, is induced via ATF4-CHOP pathway and is involved in cell death. The EMBO journal 24(6), 1243-1255.

Pabst, R., 2020. The pig as a model for immunology research. Cell and tissue research 380(2), 287-304.

Parker, M., Barrera, A., Wheaton, J., Zuberbuehler, M., Allan, D., Carlyle, J., Reddy, T., Ciofani, M., 2020. c-Maf regulates the plasticity of group 3 innate lymphoid cells by restraining the type 1 program. The Journal of experimental medicine 217(1).

Pascual-Reguant, A., Köhler, R., Mothes, R., Bauherr, S., Hernández, D., Uecker, R., Holzwarth, K., Kotsch, K., Seidl, M., Philipsen, L., Müller, W., Romagnani, C., Niesner, R., Hauser, A., 2021.

Multiplexed histology analyses for the phenotypic and spatial characterization of human innate lymphoid cells. Nature communications 12(1), 1737.

Pedersen, K., Gradel, A., Ludvigsen, T., Christoffersen, B., Fuglsang-Damgaard, C., Bendtsen, K., Madsen, S., Manfé, V., Refsgaard, H., 2022. Optimization of pig models for translation of subcutaneous pharmacokinetics of therapeutic proteins: Liraglutide, insulin aspart and insulin detemir. Translational research : the journal of laboratory and clinical medicine 239, 71-84.

Pérez-Novo, C., Holtappels, G., Vinall, S., Xue, L., Zhang, N., Bachert, C., Pettipher, R., 2010. CRTH2 mediates the activation of human Th2 cells in response to PGD(2) released from IgE/anti-IgE treated nasal polyp tissue. Allergy 65(3), 304-310.

Piriou-Guzylack, L., Salmon, H., 2008. Membrane markers of the immune cells in swine: an update. Veterinary research 39(6), 54.

Poznanski, S., Ashkar, A., 2019. What Defines NK Cell Functional Fate: Phenotype or Metabolism? Frontiers in immunology 10, 1414.

Puel, A., Cypowyj, S., Bustamante, J., Wright, J., Liu, L., Lim, H., Migaud, M., Israel, L., Chrabieh, M., Audry, M., Gumbleton, M., Toulon, A., Bodemer, C., El-Baghdadi, J., Whitters, M., Paradis, T., Brooks, J., Collins, M., Wolfman, N., Al-Muhsen, S., Galicchio, M., Abel, L., Picard, C., Casanova, J., 2011. Chronic mucocutaneous candidiasis in humans with inborn errors of interleukin-17 immunity. Science (New York, N.Y.) 332(6025), 65-68.

Qi, J., Crinier, A., Escalière, B., Ye, Y., Wang, Z., Zhang, T., Batista, L., Liu, H., Hong, L., Wu, N., Zhang, M., Chen, L., Liu, Y., Shen, L., Narni-Mancinelli, E., Vivier, E., Su, B., 2021. Single-cell transcriptomic landscape reveals tumor specific innate lymphoid cells associated with colorectal cancer progression. Cell reports. Medicine 2(8), 100353.

Qin, Z., Qu, X., Lei, L., Xu, L., Pan, Z., 2020. Y-Box-Binding Protein 3 (YBX3) Restricts Influenza A Virus by Interacting with Viral Ribonucleoprotein Complex and Imparing its Function. The Journal of general virology 101(4), 385-398.

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H., Trapnell, C., 2017. Reversed graph embedding resolves complex single-cell trajectories. Nature methods 14(10), 979-982.

Rankin, L., Groom, J., Chopin, M., Herold, M., Walker, J., Mielke, L., McKenzie, A., Carotta, S., Nutt, S., Belz, G., 2013. The transcription factor T-bet is essential for the development of NKp46+ innate lymphocytes via the Notch pathway. Nature immunology 14(4), 389-395.

Ren, G., Lai, B., Harly, C., Baek, S., Ding, Y., Zheng, M., Cao, Y., Cui, K., Yang, Y., Zhu, J., Hager, G., Bhandoola, A., Zhao, K., 2022. Transcription factors TCF-1 and GATA3 are key factors for the epigenetic priming of early innate lymphoid progenitors toward distinct cell fates. Immunity 55(8), 1402-1413.e1404.

Riding, A., Loudon, K., Guo, A., Ferdinand, J., Lok, L., Richoz, N., Stewart, A., Castro-Dopico, T., Tuong, Z., Fiancette, R., Bowyer, G., Fleming, A., Gillman, E., Suchanek, O., Mahbubani, K., Saeb-Parsy, K., Withers, D., Dougan, G., Clare, S., Clatworthy, M., 2022. Group 3 innate lymphocytes make a distinct contribution to type 17 immunity in bladder defence. iScience 25(7), 104660.

Robertson, N., Rappas, M., Doré, A., Brown, J., Bottegoni, G., Koglin, M., Cansfield, J., Jazayeri, A., Cooke, R., Marshall, F., 2018. Structure of the complement C5a receptor bound to the extra-helical antagonist NDT9513727. Nature 553(7686), 111-114.

Robinette, M., Bando, J., Song, W., Ulland, T., Gilfillan, S., Colonna, M., 2017. IL-15 sustains IL-7R-independent ILC2 and ILC3 development. Nature communications 8, 14601.

Romero, F., Martínez-A, C., Camonis, J., Rebollo, A., 1999. Aiolos transcription factor controls cell death in T cells by regulating Bcl-2 expression and its cellular localization. The EMBO journal 18(12), 3419-3430.

Ronan, V., Yeasin, R., Claud, E., 2021. Childhood Development and the Microbiome-The Intestinal Microbiota in Maintenance of Health and Development of Disease During Childhood Development. Gastroenterology 160(2), 495-506.

Salem, S., Langlais, D., Lefebvre, F., Bourque, G., Bigley, V., Haniffa, M., Casanova, J., Burk, D., Berghuis, A., Butler, K., Leahy, T., Hambleton, S., Gros, P., 2014. Functional characterization of the human dendritic cell immunodeficiency associated with the IRF8(K108E) mutation. Blood 124(12), 1894-1904.

Schleussner, N., Merkel, O., Costanza, M., Liang, H., Hummel, F., Romagnani, C., Durek, P., Anagnostopoulos, I., Hummel, M., Jöhrens, K., Niedobitek, A., Griffin, P., Piva, R., Sczakiel, H., Woessmann, W., Damm-Welk, C., Hinze, C., Stoiber, D., Gillissen, B., Turner, S., Kaergel, E., von Hoff, L., Grau, M., Lenz, G., Dörken, B., Scheidereit, C., Kenner, L., Janz, M., Mathas, S., 2018. The AP-1-BATF and -BATF3 module is essential for growth, survival and TH17/ILC3 skewing of anaplastic large cell lymphoma. Leukemia 32(9), 1994-2007.

Schroeder, J., Howard, J., Lord, G., 2022. Transcription factor-driven regulation of ILC1 and ILC3. Trends in immunology 43(7), 564-579.

Sciumé, G., Hirahara, K., Takahashi, H., Laurence, A., Villarino, A., Singleton, K., Spencer, S., Wilhelm, C., Poholek, A., Vahedi, G., Kanno, Y., Belkaid, Y., O'Shea, J., 2012. Distinct requirements for T-bet in gut innate lymphoid cells. The Journal of experimental medicine 209(13), 2331-2338.

Seehus, C., Kadavallore, A., Torre, B., Yeckes, A., Wang, Y., Tang, J., Kaye, J., 2017. Alternative activation generates IL-10 producing type 2 innate lymphoid cells. Nature communications 8(1), 1900.

Seillet, C., Luong, K., Tellier, J., Jacquelot, N., Shen, R., Hickey, P., Wimmer, V., Whitehead, L., Rogers, K., Smyth, G., Garnham, A., Ritchie, M., Belz, G., 2020. The neuropeptide VIP confers anticipatory mucosal immunity by regulating ILC3 activity. Nature immunology 21(2), 168-177.

Serafini, N., Jarade, A., Surace, L., Goncalves, P., Sismeiro, O., Varet, H., Legendre, R., Coppee, J., Disson, O., Durum, S., Frankel, G., Di Santo, J., 2022. Trained ILC3 responses promote intestinal defense. Science (New York, N.Y.) 375(6583), 859-863.

Sethuraman, A., Brown, M., Krutilina, R., Wu, Z., Seagroves, T., Pfeffer, L., Fan, M., 2018. BHLHE40 confers a pro-survival and pro-metastatic phenotype to breast cancer cells by modulating HBEGF secretion. Breast cancer research : BCR 20(1), 117.

Shao, F., Yu, D., Xia, P., Wang, S., 2021. Dynamic regulation of innate lymphoid cells in the mucosal immune system. Cellular & molecular immunology 18(6), 1387-1394.

Simoni, Y., Fehlings, M., Kløverpris, H., McGovern, N., Koo, S., Loh, C., Lim, S., Kurioka, A., Fergusson, J., Tang, C., Kam, M., Dennis, K., Lim, T., Fui, A., Hoong, C., Chan, J., Curotto de Lafaille, M., Narayanan, S., Baig, S., Shabeer, M., Toh, S., Tan, H., Anicete, R., Tan, E., Takano, A., Klenerman, P., Leslie, A., Tan, D., Tan, I., Ginhoux, F., Newell, E., 2018. Human Innate Lymphoid Cell Subsets Possess Tissue-Type Based Heterogeneity in Phenotype and Frequency. Immunity 48(5), 1060.

Sinkora, M., Butler, J., 2016. Progress in the use of swine in developmental immunology of B and T lymphocytes. Developmental and comparative immunology 58, 1-17.

Song, C., Lee, J., Gilfillan, S., Robinette, M., Newberry, R., Stappenbeck, T., Mack, M., Cella, M., Colonna, M., 2015. Unique and redundant functions of NKp46+ ILC3s in models of intestinal inflammation. The Journal of experimental medicine 212(11), 1869-1882.

Spits, H., Artis, D., Colonna, M., Diefenbach, A., Di Santo, J., Eberl, G., Koyasu, S., Locksley, R., McKenzie, A., Mebius, R., Powrie, F., Vivier, E., 2013. Innate lymphoid cells--a proposal for uniform nomenclature. Nature reviews. Immunology 13(2), 145-149.

Spits, H., Bernink, J., Lanier, L., 2016. NK cells and type 1 innate lymphoid cells: partners in host defense. Nature immunology 17(7), 758-764.

Spits, H., Cupedo, T., 2012. Innate lymphoid cells: emerging insights in development, lineage relationships, and function. Annual review of immunology 30, 647-675.

Srpan, K., Ambrose, A., Karampatzakis, A., Saeed, M., Cartwright, A., Guldevall, K., De Matos, G., Önfelt, B., Davis, D., 2018. Shedding of CD16 disassembles the NK cell immune synapse and boosts serial engagement of target cells. The Journal of cell biology 217(9), 3267-3283.

Starbæk, S., Brogaard, L., Dawson, H., Smith, A., Heegaard, P., Larsen, L., Jungersen, G., Skovgaard, K., 2018. Animal Models for Influenza A Virus Infection Incorporating the Involvement of Innate Host Defenses: Enhanced Translational Value of the Porcine Model. ILAR journal 59(3), 323-337.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W., Hao, Y., Stoeckius, M., Smibert, P., Satija, R., 2019. Comprehensive Integration of Single-Cell Data. Cell 177(7), 1888-1902.e1821.

Swaim, C., Scott, A., Canadeo, L., Huibregtse, J., 2017. Extracellular ISG15 Signals Cytokine Secretion through the LFA-1 Integrin Receptor. Molecular cell 68(3), 581-590.e585.

Sykes, M., Sachs, D.H., 2019. Transplanting organs from pigs to humans. Sci Immunol 4(41).

Tibbitt, C., Stark, J., Martens, L., Ma, J., Mold, J., Deswarte, K., Oliynyk, G., Feng, X., Lambrecht, B., De Bleser, P., Nylén, S., Hammad, H., Arsenian Henriksson, M., Saeys, Y., Coquet, J., 2019. Single-Cell RNA Sequencing of the T Helper Cell Response to House Dust Mites Defines a Distinct Gene Expression Signature in Airway Th2 Cells. Immunity 51(1), 169-184.e165.

Usui, N., Berto, S., Konishi, A., Kondo, M., Konopka, G., Matsuzaki, H., Shimada, S., 2021. Zbtb16 regulates social cognitive behaviors and neocortical development. Translational psychiatry 11(1), 242.

Verrier, T., Satoh-Takayama, N., Serafini, N., Marie, S., Di Santo, J., Vosshenrich, C., 2016. Phenotypic and Functional Plasticity of Murine Intestinal NKp46+ Group 3 Innate Lymphoid Cells. Journal of immunology (Baltimore, Md. : 1950) 196(11), 4731-4738.

Viant, C., Rankin, L., Girard-Madoux, M., Seillet, C., Shi, W., Smyth, M., Bartholin, L., Walzer, T., Huntington, N., Vivier, E., Belz, G., 2016. Transforming growth factor-β and Notch ligands act as opposing environmental cues in regulating the plasticity of type 3 innate lymphoid cells. Science signaling 9(426), ra46.

Vikesaa, J., Hansen, T., Jønson, L., Borup, R., Wewer, U., Christiansen, J., Nielsen, F., 2006. RNA-binding IMPs promote cell adhesion and invadopodia formation. The EMBO journal 25(7), 1456-1468.

Vivier, E., Artis, D., Colonna, M., Diefenbach, A., Di Santo, J., Eberl, G., Koyasu, S., Locksley, R., McKenzie, A., Mebius, R., Powrie, F., Spits, H., 2018. Innate Lymphoid Cells: 10 Years On. Cell 174(5), 1054-1066.

Vivier, E., van de Pavert, S., Cooper, M., Belz, G., 2016. The evolution of innate lymphoid cells. Nature immunology 17(7), 790-794.

von Moltke, J., Ji, M., Liang, H., Locksley, R., 2016. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. Nature 529(7585), 221-225.

Vonarbourg, C., Mortha, A., Bui, V., Hernandez, P., Kiss, E., Hoyler, T., Flach, M., Bengsch, B., Thimme, R., Hölscher, C., Hönig, M., Pannicke, U., Schwarz, K., Ware, C., Finke, D., Diefenbach, A., 2010. Regulated expression of nuclear receptor RORγt confers distinct functional fates to NK cell receptor-expressing RORγt(+) innate lymphocytes. Immunity 33(5), 736-751.

Wagner, J., Wong, P., Schappe, T., Berrien-Elliott, M., Cubitt, C., Jaeger, N., Lee, M., Keppel, C., Marin, N., Foltz, J., Marsala, L., Neal, C., Sullivan, R., Schneider, S., Keppel, M., Saucier, N., Cooper, M., Fehniger, T., 2020. Stage-Specific Requirement for Eomes in Mature NK Cell Homeostasis and Cytotoxicity. Cell reports 31(9), 107720.

Wang, S., Xia, P., Chen, Y., Qu, Y., Xiong, Z., Ye, B., Du, Y., Tian, Y., Yin, Z., Xu, Z., Fan, Z., 2017. Regulatory Innate Lymphoid Cells Control Innate Intestinal Inflammation. Cell 171(1), 201-216.e218.

Warner, K., Ohashi, P., 2019. ILC regulation of T cell responses in inflammatory diseases and cancer. Seminars in immunology 41, 101284.

Wiarda, J., Trachsel, J., Sivasankaran, S., Tuggle, C., Loving, C., 2022. Intestinal single-cell atlas reveals novel lymphocytes in pigs with similarities to human cells. Life science alliance 5(10).

Wojno, E., Monticelli, L., Tran, S., Alenghat, T., Osborne, L., Thome, J., Willis, C., Budelsky, A., Farber, D., Artis, D., 2015. The prostaglandin D_2 receptor CRTH2 regulates accumulation of group 2 innate lymphoid cells in the inflamed lung. Mucosal immunology 8(6), 1313-1323.

Wolf, E., Kemter, E., Klymiuk, N., Reichart, B., 2019. Genetically modified pigs as donors of cells, tissues, and organs for xenotransplantation. Animal frontiers : the review magazine of animal agriculture 9(3), 13-20.

Wypych, T., Wickramasinghe, L., Marsland, B., 2019. The influence of the microbiome on respiratory health. Nature immunology 20(10), 1279-1290.

Xia, P., Liu, J., Wang, S., Ye, B., Du, Y., Xiong, Z., Han, Z., Tong, L., Fan, Z., 2017. WASH maintains NKp46 ILC3 cells by promoting AHR expression. Nature communications 8, 15685.

Xiao, T., Xu, Z., Zhang, H., Geng, J., Qiao, Y., Liang, Y., Yu, Y., Dong, Q., Suo, G., 2019. TP53I11 suppresses epithelial-mesenchymal transition and metastasis of breast cancer cells. BMB reports 52(6), 379-384.

Xie, Y., Hu, X., Li, S., Qiu, Y., Cao, R., Xu, C., Lu, C., Wang, Z., Yang, J., 2022. Pharmacological targeting macrophage phenotype via gut-kidney axis ameliorates renal fibrosis in mice. Pharmacological research 178, 106161.

Xu, W., Cherrier, D., Chea, S., Vosshenrich, C., Serafini, N., Petit, M., Liu, P., Golub, R., Di Santo, J., 2019. An Id2-Reporter Mouse Redefines Innate Lymphoid Cell Precursor Potentials. Immunity 50(4), 1054-1068.e1053.

Yamaguchi, H., Wang, H., 2004. CHOP is involved in endoplasmic reticulum stress-induced apoptosis by enhancing DR5 expression in human carcinoma cells. The Journal of biological chemistry 279(44), 45495-45502.

Yaman, E., Gasper, R., Koerner, C., Wittinghofer, A., Tazebay, U., 2009. RasGEF1A and RasGEF1B are guanine nucleotide exchange factors that discriminate between Rap GTP-binding proteins and mediate Rap2-specific nucleotide exchange. The FEBS journal 276(16), 4607-4616.

Yamashita, M., Kuehn, H., Okuyama, K., Okada, S., Inoue, Y., Mitsuiki, N., Imai, K., Takagi, M., Kanegane, H., Takeuchi, M., Shimojo, N., Tsumura, M., Padhi, A., Zhang, K., Boisson, B., Casanova, J.,

Ohara, O., Rosenzweig, S., Taniuchi, I., Morio, T., 2021. A variant in human AIOLOS impairs adaptive immunity by interfering with IKAROS. Nature immunology 22(7), 893-903.

Yang, J., Cornelissen, F., Papazian, N., Reijmers, R., Llorian, M., Cupedo, T., Coles, M., Seddon, B., 2018. IL-7-dependent maintenance of ILC3s is required for normal entry of lymphocytes into lymph nodes. The Journal of experimental medicine 215(4), 1069-1077.

Young, J., Bouligand, J., Francou, B., Raffin-Sanson, M., Gaillez, S., Jeanpierre, M., Grynberg, M., Kamenicky, P., Chanson, P., Brailly-Tabard, S., Guiochon-Mantel, A., 2010. TAC3 and TACR3 defects cause hypothalamic congenital hypogonadotropic hypogonadism in humans. The Journal of clinical endocrinology and metabolism 95(5), 2287-2295.

Yu, L., Agirman, G., Hsiao, E., 2022. The Gut Microbiome as a Regulator of the Neuroimmune Landscape. Annual review of immunology 40, 143-167.

Zeis, P., Lian, M., Fan, X., Herman, J., Hernandez, D., Gentek, R., Elias, S., Symowski, C., Knöpper, K., Peltokangas, N., Friedrich, C., Doucet-Ladeveze, R., Kabat, A., Locksley, R., Voehringer, D., Bajenoff, M., Rudensky, A., Romagnani, C., Grün, D., Gasteiger, G., 2020. In Situ Maturation and Tissue Adaptation of Type 2 Innate Lymphoid Cell Progenitors. Immunity 53(4), 775-792.e779.

Zhang, J., Marotel, M., Fauteux-Daniel, S., Mathieu, A., Viel, S., Marçais, A., Walzer, T., 2018. T-bet and Eomes govern differentiation and function of mouse and human NK cells and ILC1. European journal of immunology 48(5), 738-750.

Zhang, L., Shih, A., Yang, X., Kuei, C., Wu, J., Deng, X., Mani, N., Mirzadegan, T., Sun, S., Lovenberg, T., Liu, C., 2012. Identification of structural motifs critical for epstein-barr virus-induced molecule 2 function and homology modeling of the ligand docking site. Molecular pharmacology 82(6), 1094-1103.

Zhang, L., Zhu, J., Wang, H., Xia, J., Liu, P., Chen, F., Jiang, H., Miao, Q., Wu, W., Zhang, L., Luo, L., Jiang, X., Bai, Y., Sun, C., Chen, D., Zhang, X., 2021. A high-resolution cell atlas of the domestic pig lung and an online platform for exploring lung single-cell data. Journal of genetics and genomics = Yi chuan xue bao 48(5), 411-425.

Zhang, W., Wu, Q., Zhu, Y., Yang, G., Yu, J., Wang, J., Ji, H., 2019. Lactobacillus rhamnosusProbiotic GG Induces Alterations in Ileal Microbiota With Associated CD3CD19T-betIFNγ Cell Subset Homeostasis in Pigs Challenged With Serovar 4,[5],12:i. Frontiers in microbiology 10, 977.

Zheng, Y., Li, S., Li, S., Yu, S., Wang, Q., Zhang, K., Qu, L., Sun, Y., Bi, Y., Tang, F., Qiu, H., Gao, G., 2022. Transcriptome profiling in swine macrophages infected with African swine fever virus at single-cell resolution. Proceedings of the National Academy of Sciences of the United States of America 119(19), e2201288119.

Zhong, C., Cui, K., Wilhelm, C., Hu, G., Mao, K., Belkaid, Y., Zhao, K., Zhu, J., 2016. Group 3 innate lymphoid cells continuously require the transcription factor GATA-3 after commitment. Nature immunology 17(2), 169-178.

Zhong, C., Zheng, M., Cui, K., Martins, A., Hu, G., Li, D., Tessarollo, L., Kozlov, S., Keller, J., Tsang, J., Zhao, K., Zhu, J., 2020. Differential Expression of the Transcription Factor GATA3 Specifies Lineage and Functions of Innate Lymphoid Cells. Immunity 52(1), 83-95.e84.

Zhou, B., Yuan, Y., Zhang, S., Guo, C., Li, X., Li, G., Xiong, W., Zeng, Z., 2020a. Intestinal Flora and Disease Mutually Shape the Regional Immune System in the Intestinal Tract. Frontiers in immunology 11, 575.

Zhou, L., Zhou, W., Joseph, A., Chu, C., Putzel, G., Fang, B., Teng, F., Lyu, M., Yano, H., Andreasson, K., Mekada, E., Eberl, G., Sonnenberg, G., 2022. Group 3 innate lymphoid cells produce the growth factor HB-EGF to protect the intestine from TNF-mediated inflammation. Nature immunology 23(2), 251-261.

Zhou, Z., Chen, J., Zhang, R., Ye, L., Wang, J., Pan, Y., Wang, X., Li, W., Zhang, Y., Xu, L., Chen, M., 2020b. Tescalcin is an unfavorable prognosis factor that regulats cell proliferation and survival in hepatocellular carcinoma patients. Cancer communications (London, England) 40(8), 355-369.

Figure legend

Fig. 1 | Characterization of immune cells in the pig jejunum after flow cytometry sorting.

A. UMAP plots showing the immune landscape of the 4 subpopulations of intestinal cells (selection by flow cytometry). Cells are color-coded according to the defined subset (ILCs/T cells: red; myeloid cells: blue; B cells: green; undifferentiated cells: yellow). **B**. Pie charts showing the quantitative differences for these four classes of cell clusters, ILCs/T cells: 11,208 cells (71.1%); myeloid cells: 3500 cells (22.2%); B cells: 717 cells (4.6%); undefined: 338 cells (2.1%). **C**. Violin plot representation of the differentially expressed marker genes for the four clusters. Three signature genes are displayed in each cluster, except for the undefined cells. **D**. UMAP plots showing the immune landscape of the 13 clusters of intestinal cells. **E**. Heatmap showing the marker genes with the five highest differentially expressed genes in each cluster. The abscissa shows different clusters, and the ordinate shows differentially expressed gene names.

Fig. 2 | Detailed typing research of ILCs in pig jejunum.

A. UMAP plots showing the immune landscape of the 7 clusters of intestinal cells (ILC/T-cell regrouping). Violin statistics of the marker genes for ILC and T-cell differential expression. **B**. receptor genes. **C**. cytokine genes. **D**. transcription factor genes. Detailed statistics of the expressed genes in the four subsets of ILCs. Dot plot showing Z scored mean expression of selected marker genes in the ILC clusters. The size of the dot represents the percentage of the gene in a subset of cells. The colors of the dots represent the average expression of the gene in a subset of cells. **E**. receptor genes. **F**. cytokine genes. **G**. transcription factor genes. **H**. other genes.

Fig. 3 | Detailed typing research of ILC3s in pig jejunum

A. UMAP plots showing the immune landscape of the ILC3 clusters. Cells are color-coded according to the defined subset (ILC3a: blue; ILC3b: orange; ILC3c: green; ILC3d: red). **B**. Spearman correlation analysis between subsets of ILC3s. Colors represent the degree of correlation between the ILC3 clusters, and tree branches represent correlation cluster grading. **C**. Heatmap showing the differential gene average expression of ILC3 clusters. The top 10 differential genes were taken from the ILC3a, ILC3b, ILC3c and ILC3d subgroups. Among these, only one differentially expressed gene in ILC3b cells was upregulated, while the remaining differentially expressed genes were downregulated. **D**. Top ten expressed genes from the receptor protein and top ten expressed genes encoding secreted protein, transcription factor, and other protein-encoding significantly differentiating among the groups of ILCs. Gene symbols and annotations were retrieved from public databases. Receptor protein genes are color-coded in blue, secreted protein genes in orange, transcription factor genes in red, and other protein-encoding genes in green. Genes are ranked by pvalue. **E**. Violin plots of the ILC3a highly expressed genes. **F**. Violin plots of the ILC3c highly expressed genes. **G**. Violin plots of the ILC3d highly expressed genes.

Fig. 4 | Pseudotemporal analysis of ILCs

A. Pseudotime traces created by Monocle2 using the 7 clusters of ILCs. Arrows indicate the direction of differentiation and show the enrichment position of each cluster in the pseudotime (different colors represent the different clusters). Black circles mark positions that are possible nodes of pseudotime. **B**. Gene heatmap of cell differences between different branches. The top is

the node branch in pseudotemporal order, each row represents a gene, each column is the pseudotime point, the color represents the average expression value of the gene at the current time point, and the color is gradually reduced from red to blue. Genes with similar expression trends will cluster together to form different clusters. **C**. Statistical map of the changes in marker genes in ILC clusters with pseudotime. **D**. Violin plots of some genes that play regulatory transdifferentiation roles in subsets of ILCs.

Fig. 5 | The pig and human clusters of intestinal ILCs are transcriptionally conserved.
A. Venn statistics of genes expressed in human-pig-mouse intestinal ILCs (Statistical mapping after normalization according to the number of expressed gene names). **B**. Statistical heatmap of the top 20 highly expressed genes in human, pig and mouse intestinal ILCs. **C**. UMAP plots showing the immune landscape of human and pig intestinal ILC enterocyte cluster subsets (divided into 12 clusters). UMAP with color coding according to donor origin. **D**. Six marker genes coexpressed in human and pig ILCs are shown. **E**. Five marker genes are displayed as differentially expressed in human and pig ILCs.

Fig. 6 | The pig and mouse clusters of intestinal ILCs with differentially expressed genes
A. Mouse intestinal ILCs were examined by flow cytometry. **B**. UMAP plots showing the immune landscape of human and mouse intestinal ILC clusters. **C**. Six marker genes coexpressed in pig and mouse ILCs are shown. **D**. Differential expression of 4 cytokine marker genes in pig and mouse ILC2s is shown. **E**. Six cytokine marker genes are displayed as differentially expressed in pig and mouse ILC3s. **F**. Five marker genes are differentially expressed in pig and mouse ILCs.

Graphical abstract

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6