Genomic relationships among diploid and polyploid species of the genus *Ludwigia* L.

section *Jussiaea* using a combination of molecular cytogenetic, morphological, and

crossing investigations

D. Barloy[1*], L. Portillo - Lemus[1], S. A. Krueger-Hadfield[3], V. Huteau[2], O. Coriton[2]

[1] DECOD (Ecosystem Dynamics and Sustainability), Institut Agro, INRAE, IFREMER

35042 Rennes, France

[2] Molecular Cytogenetics Platform, IGEPP (Institute for Genetics, Environment and Plant

Protection), INRAE, Institut Agro, Univ Rennes, 35653, Le Rheu, France

[3] Department of Biology, The University of Alabama at Birmingham, 1300 University Blvd,

Birmingham, AL 35294, USA

[*] Corresponding author: dominique.barloy@agrocampus-ouest.fr

ABSTRACT

The genus *Ludwigia* L. section *Jussiaea* is composed of a polyploid species complex with 2x,

4x, 6x and 10x ploidy levels, suggesting possible hybrid origins. The aim of the present study

is to understand the genomic relationships among diploid and polyploid species in the section

Jussiaea. Morphological and cytogenetic observations, controlled crosses, genomic *in situ*

hybridization (GISH), and flow cytometry were used to characterize species, ploidy levels,

ploidy patterns, and genomic composition across taxa. Genome sizes obtained were in

agreement with the diploid, tetraploid, hexaploid, and decaploid ploidy levels. Results of

GISH showed that progenitors of *Ludwigia stolonifera* (4x) were *Ludwigia peploides* subsp.

montevidensis (2x) and *Ludwigia helminthorrhiza* (2x), which also participated for one part

(2x) to the *Ludwigia ascendens* genome (4x). *Ludwigia grandiflora* subsp. *hexapetala* (10x)

resulted from the hybridization between *L. stolonifera* (4x) and *Ludwigia grandiflora* subsp.

grandiflora (6x). One progenitor of *L. grandiflora* subsp. *grandiflora* was identified as *L.*

peploides (2x). Our results suggest the existence of several processes of hybridization, leading

to polyploidy, and possibly allopolyploidy, in the section Jussiaea due to the diversity of

ploidy levels. The success of GISH opens up the potential for future studies to identify other

missing progenitors in *Ludwigia* L. as well as other taxa.

Keywords: GISH, invasive plant, *Ludwigia* L., Onagraceae, polyploidy, phylogenetics

INTRODUCTION
36
Polyploidization is widespread in plants and is considered as a major driving force in

plant speciation and evolution (Husband et al., 2013; Alix et al., 2017; Otto and Whitton,

2000). Autopolyploid plants arise from the duplication of one genome within one species and

allopolyploid plants result from the association of two or more divergent genomes through

interspecific hybridization and subsequent genome duplication (Alix et al., 2017; Soltis et al.,

2015). Furthermore, some polyploids can arise from both auto- and allopolyploidy events

because of their evolutionary histories and are called auto-allo-polyploid. Genomic analyses

have revealed that all angiosperms have been subjected to at least one round of polyploidy in

their evolutionary history and are thus considered paleopolyploids (Garsmeur et al., 2014).

Thus, understanding the origins of polyploid taxa is integral to understanding angiosperm

evolution.

Polyploid plants are often thought to be more resilient to extreme environments than

diploids because of their increased genetic variation (Husband et al., 2013). Their duplicated

genes act as a buffer and can include gene conversion events, activation of transposable

elements, chromatin remodelling, and DNA methylation changes (Hollister, 2015).

Polyploidy might confer an advantage with both abiotic and biotic stress by increasing

tolerance to salt or drought stress or by improving resistance to bioagressors (Van de Peer et

al., 2021). Thus, polyploids are able to occupy new ecological niches (Stebbins, 1985; Blaine

Marchant et al., 2016) and often show greater adaptability than their progenitors (McIntyre,

2012; Allario et al., 2013; Baniaga et al., 2020; Akiyama et al., 2021; Van de Peer et al.,

2021). Van de Peer et al. (2021) suggested that as in a constant environment, polyploidization

may play an important role in response to habitat disturbance, nutritional stress, physical

stress, and climate change (Wei et al., 2019). For example, Baniaga et al. (2020) showed that

ecological niches of polyploid plants differentiated often faster than found in their diploid

relatives. A polyploid advantage has also been reported in invasive plants and their success in

non-native habitats (Te Beest et al., 2012). However, Lobato-de Magalhães et al. (2021)

observed little difference in the incidence of each ploidy state within a set of 49 of the world's

most invasive aquatic weeds and concluded there is no consistent evidence of polyploid

advantage in invasiveness. Nevertheless, *Spartina anglica*, an invasive neoallopolyploid weed

species that appeared around 1890, has increased fitness with its prolific seed production,

fertility, and extensive clonal growth as compared to its progenitors (Baumel et al., 2002). A

recent study including 50 alien non-invasive aquatic plant species and 68 alien invasive

species across various aquatic habitats in the Kashmir Himalayas found that invasive species

are largely polyploids whereas non-invasive species tend to diploids (Wani et al., 2018).

Ludwigia L., a worldwide wetland genus of 83 species, forms a strongly monophyletic

lineage sister to the rest of the Onagraceae. It is currently classified as members of 23 sections

(Levin et al., 2003, 2004). Sections were clustered into three main groups by Raven (1963).

The first group concerned the Myrtocarpus complex, comprising 14 sections (Raven, 1963;

Eyde, 1977; Ramamoorthy, 1979; Zardini and Raven, 1992). The second group included

species in the section *Eujussiaea* Munz (Munz, 1942), also referred to as a sect.

Oligospermum (Raven, 1963) but now correctly called sect. *Jussiaea* (Hoch et al., 1993). The

third group combined species in sect. *Isnardia*, sect. *Ludwigia*, sect. *Microcarpium*, and sect.

Miquelia P.H. Raven (Raven, 1963; Wagner et al., 2007). Liu et al. (2017) provided the first

comprehensive molecular phylogeny of *Ludwigia* genus using both nuclear and chloroplast

DNA regions. Sixty of 83 species in the *Ludwigia* genus were distributed in the two clades A

and B, with the sub-clade B1 which consisted of only sect. *Jussiaea*. This section included

seven species: three diploid species (2n=2x=16) (*Ludwigia torulosa* (Arn.) H. Hara, *Ludwigia*

helminthorrhiza (Mart.) H. Hara, *Ludwigia peploides* (Kunth) P.H. Raven); two tetraploid

species (2n=4x=32) (*Ludwigia adscendens* (L.) H. Hara, *Ludwigia stolonifera* (Guill. &Perr.)

P.H. Raven); one hexaploid species (2n=6x=48) (*Ludwigia grandiflora* subsp. *grandiflora*);

and one decaploid species (2n=10x=80) (*Ludwigia grandiflora* subsp. *hexapetala*). While

most species are native to the New World, particularly South America, two species are

restricted to the Old World, *Ludwigia stolonifera* and *Ludwigia adscendens*, in Africa and

tropical Asia, respectively (Wagner et al., 2007) (Table S1). It is not easy to distinguish

between the hexaploid and decaploid species morphologically and both have previously been

treated as a single species (*Ludwigia uruguayensis* (Cambess.) H. Hara; Zardini et al., 1991).

Octoploid hybrids between *L. grandiflora* subsp. *hexapetala (Lgh)* and *L. grandiflora* subsp.

grandiflora (*Lgg*) were found in southern Brazil which for both species is their native area

(Zardini et al, 1991). Studies of Liu et al (2017) confirmed close relationship between *Lgg*

and *Lgh*. So, Nesom and Kartesz (2000) suggested that as *Lgg* and *Lgh* shared genomic

portions and possible hybridization between them, both species were recognized as subspecies

within the single species *L. grandiflora*. However, several authors, including Okada et al.

(2009) and Grewell et al (2016), continue to recognize two distinct species. In this paper, species

were named as described by Nesom and Kartesz (2000) and Armitage et al (2013), i.e., considered

as two subspecies of *L. grandiflora* (*Lgg* and *Lgh*). So, phylogenetic studies (Lui et al 2017)

revealed that the *L. peploides* (2x) or a relative and the *L. adscendens* (4x) are probably

progenitors to *L. stolonifera* and *L. × taiwanensis* (3x), respectively. Furthermore, based on

morphological observations, Zardini et al. (1991) suggested that *Lgh* may be result of

interspecific hybridization between *Lgg* and *L. hookeri*. So, in view of the diversity of ploidy

levels present in the *ludwigia* sect. *Jussiaea*, results of morphological and molecular analysis,

polyploid species could be probably the result of hybridization between diploid species or

combinations of diploid and polyploid species. In this study, we focused on species belonging

to the second group, sect. *Jussiaea*. Most species of the section grow in warm temperate to

subtropical moist or wet habitats worldwide. Some of these species, such as *Ludwigia*

peploides subsp. *montevidensis* (Kunth) P.H. Raven, *Ludwigia grandiflora* (syn. *L.*

grandiflora subsp. *grandiflora*), *Ludwigia hexapetala* (Hook. & Arn.) Zardini, H.Y. Gu &

P.H. Raven (syn. *L. grandiflora* subsp. *hexapetala*) (Hook. & Arn.) Zardini, H. Y. Gu & P. H.

Raven, can be invasive weeds in wetlands and other wet areas in the USA (Grewell et al.,

2016), Europe (Portillo-Lemus et al., 2021), Japan (Hieda et al., 2020), and Korea (Kim et al,

2019). Recently, Méndez Santos and González-Sivilla (2020) revealed that *L. helminthorrhiza*

(Mart.) H. Hara must be treated and managed as an invasive alien species in Cuba.

Reproductive systems in *Ludwigia* L. are both clonal with production of asexual fragments

and sexual with seeds production. Okada et al. (2009) showed that clonal spread through

asexual reproduction is the primary regeneration mode of *L. grandiflora* subsp. *grandiflora*

and *L. grandiflora* subsp. *hexapetala* in California. Furthermore, Dandelot (2004) reports that

all the populations of *L. grandiflora* subsp. *hexapetala* in the French Mediterranean area

could have originated from a single clone. Similarly, Reddy et al. (2021) observed low

genotypic diversity in both *L. grandiflora* subsp. *grandiflora* and *L. grandiflora* subsp.

hexapetala in the United State with as example an analysis of multiple invasive populations of

L. grandiflora subsp. *hexapetala* in Alabama, California, Oregon, Washington, and Florida

identified a single genotype.

The aim of this study is to explore the complicated evolutionary history of genus

Ludwigia L. section *Jussiaea* using a combination of cytogenetic, morphological, and

crossing investigations. This is a difficult puzzle to elucidate, with taxa ranging from diploid

to decaploid and with both allo- and autopolyploidy involved in the history of these taxa. The

occurrence of different ploidy levels of *Ludwigia* species belonging to the same clade might

indicate that a diploid species in this clade could be the progenitor of the polyploids analysed.

However, while many authors have highlighted the possibility of interspecific hybridization

between the species presents in the Jussieae section, there is a lack of data enabling the

polyploid origin of these species to be identified, i.e., the auto or allopolyploid origin as well

as that of the progenitor species. First, we observed some morphological traits as a simple

verification step to prove that the species collected were those expected. Second, we

characterized the different species by analysis of their genome size using flow cytometry and

their ploidy level by cytogenetic observations. We identified the genomic relationships by

Genomic *in situ* Hybridization (GISH) and evaluated the ability of inter-species hybridization

after controlled pollination. The genomic relationships between diploid and polyploid species

are reported for the first time in sect. *Jussiaea*.

MATERIAL AND METHODS

Plant material

Two diploid, two tetraploid, one hexaploid, and one decaploid *Ludwigia* species were

analysed. Fifteen plants of *Ludwigia peploides* subsp. *montevidensis* (2x) (hereafter, *Lpm*) and

of *L. grandiflora* subsp. *hexapetala* (hereafter, *Lgh*) (10x) were collected in France at the

marshes of la Musse (47°14'27.5"N, 1°47'21.3"W) and Mazerolles (47°23'16.3"N,

1°28'09.7"W), respectively. Ten plants of the diploid species *L. helminthorrhiza* (hereafter,

Lh) was purchased in aquarium store (provider Ruinemans Aquarium B.V. Netherland). Five

plants of *Ludwigia adscendens* (L.) H. HARA (4x) (hereafter, *La*), and of *L. stolonifera* (4x)

(hereafter, *Ls*) and ten of *L. grandiflora* subsp. *grandiflora* (6x) (hereafter, *Lgg*) were

collected in Flores island, Indonesia (Pulau Flores; 8°49'40.8"S, 120°48'39.0"E), Lebanon

(Hekr al Dahri; 34°37'54.5"N, 36°01'28.9"E), and the USA (Co. Rd 73, outside Greensboro,

AL; 32°61'51.41"N, 87°68'65.4"W), respectively. As all *Ludwigia* species reproduce

preferentially by clonal reproduction ; each plant was used as mother plant giving new plants

from the development of buds present on its stem which are then used for all experiments

(Okada et al., 2009; Glover et al., 2015). The plants were easily maintained in the greenhouse

at Institut Agro Rennes - Angers before analysis (Portillo-Lemus et al, 2021).

Morphology

To confirm that the collected *Ludwigia* species corresponded to the expected species,

we carried out qualitative observations using simple visual morphological traits such as the

color of the flowers and roots and the pneumatophore form as reported in Table S1.

Morphological observations for each species were made on at least 30 plants in the

greenhouse and confirmed in natura on 15 plants in 15 and 36 populations of *Lpm* and *Lgh* in

France, respectively.

Chromosome counting

At least 40 root tips of 0.5 - 1.5 cm in length were taken for each *Ludwigia* sp. as follows

from 15 *Lpm*; ten *Lh*; five *La*; five *Ls*; ten *Lgg* and 15 *Lgh* different plants and were incubated

in 0.04% 8-hydroxiquinoline for 2 hours at room temperature in the dark, followed by 2h at

4°C to accumulate metaphases. Chromosome preparations were performed according to

procedures detailed in Ksiazczyk et al. (2011). At least four roots per species were observed.

The 4',6-diamidino-2-phenylindole (DAPI) staining chromosome counts per species were

estimated on a total of 20 cells at the mitotic metaphase stage using the visualization software

Zen 2 PRO (Carl Zeiss, Germany).

Genome size estimation by flow cytometry

To explore the genome size among the different *Ludwigia* spp., we used flow cytometry.

Approximately 4 mg of fresh roots or leaves from five plants of *Ludwigia* spp. and of fresh

leaves from five plants of *Trifolium rupens* (2C DNA = 2.23 pg) *or Zea mays* (2C DNA =

5.55 pg) (Zonneved et al, 2019) (used as an internal reference standard for *Lpm*, *Lh* and *Lgh*

species and *Ls*, *La*, *Lgg* and *Lgh* species, respectively) were harvested and transferred to a

Petri dish. Estimation of genome size for each species was obtained as described by Boutte et

al, 2020. For the different *Ludwigia* spp., two or three measures of genome size were made,

excepted for *Ls* (only one measure). From each species, the mean ratio of DNA content was

calculated (mean + CI (Confidence Interval), p-value= 0.05)). Genome sizes were converted

from picograms (pg) to Megabases (Mb) using $1\,pg\,=\,978$ Mbp (Dolezel et al., 2003).

Genomic in situ hybridization (GISH)

GISH is used to distinguish chromosomes from different genomes in interspecific/intergeneric

hybrids or allopolyploids. Total genomic DNA of a genitor involved in the formation of a

hybrid is used at the same time as an unlabeled DNA from another genitor, at a higher

concentration, which serves as a blocking DNA, hybridizing with the sequences in common

with both genomes. This method is based on repetitive sequences which are more often in

plant species-specific. Thus, we compared the level of relatedness between the genomes of the

studied species and hypothetical parental species.

DNA was extracted from 30 mg of freeze-dried buds taken from 15 *Lpm*, ten *Lh*, five

Ls, five *La*, ten *Lgg,* and 15 *Lgh* plants, using the Macherey-Nagel extraction kit

NucleoSpin® Food to which we have made following modifications to obtain a

polysaccharide free DNA: (1) after lysis step with Buffer CF, we mixed freeze-dried buds

with an equivalent volume of PCIA 25:24:1 (parts of phenol, chloroform, isoamyl alcohol) for

5 minutes; (2) then we transferred the whole in a tube containing phase-look gel and

centrifuged at 800rpm for 5 minutes (Quantabio, Massachusetts, USA); (3) then the DNA was

precipitated using absolute ethanol at -18°C instead of QW and C5 buffers. Finally, the DNA

was resuspended after an incubation of 5 min in 100 ml elution buffer with 5 mM TRIS at pH

8.5 at 65°C. 500 ng of total genomic DNA were labelled by random priming with biotin-14-

dCTP (Invitrogen by Thermo Fisher Scientific) used as probes.

Total genomic DNA used as a blocking DNA was autoclaved to yield fragments of 100-300

bp. The ratio DNA probe / blocking DNA was 1:50. The hybridized probes correspond to the

chromosomes present on the slide (i.e., same species) and genomic DNA (blocking DNA)

from different species were used as competitors in to block the common sequences at both

species. Genomic In Situ Hybridization (GISH) was carried out as described in Coriton et al,

2019, using a 5 µg of blocking DNA (~50-fold excess). Biotinylated probes were

immunodetected by Texas Red avidin DCS (Vector Laboratories, Burlingame, CA, USA) and

the signal was amplified with biotinylated anti-avidin D (Vector Laboratories). The

chromosomes were mounted and counterstained in Vectashield (Vector Laboratories)

containing 2.5µg/mL 4',6-diamidino-2-phenylindole (DAPI). Fluorescence images were

captured using an ORCA-Flash4 (Hamamatsu, Japan) on an Axioplan 2 microscope (Zeiss,

Oberkochen, Germany) and analysed using Zen 2 PRO software (Zeiss, Oberkochen,

Germany). For each *Ludwigia* species, at least three independent slides were made with a total

of 20 cells observed per species. The images were processed using Photoshop v.8.0.1 (Adobe

Systems Inc., San Jose, CA, USA).

Controlled interspecific crosses

Controlled interspecific pollinations were carried out in the greenhouse between

Ludwigia species which putatively shared the same parental genome. Thus, interspecific

hybridizations were made between *L. peploides* subsp. *montevidensis*, *L. stolonifera* and/or *L.*

grandiflora subsp. *hexapetala* used as male or as female. Ten plants of each species were

used for crosses. *Ludwigia* spp. produced flowers on a shoot until July to October, with at one

time only one flower per shoot at the good stage of mature for pollination. To carry out

interspecific pollinations, flowers were enclosed in cellophane bags to protect them from

external pollen before and after pollination. Flowers used as 'female' were emasculated

before anthesis. A mix of pollen from flowers of five different plants for each of other species

was used to pollinate emasculated flowers. Between two to 25 interspecific crosses were

made according to the availability of flowers. To control efficiency of pollination in

greenhouse, we also conducted at the same time 45, 75 and 50 intraspecific crosses for *Lpm,*

Lgh and *Ls*, respectively.

Pollination success for interspecific crosses was estimated by the number of fruits,

fruit size and weight, the number of seeds, viable plantlets, and the number of plants

ultimately produced. For intraspecific crosses, the number of fruits obtained were noted.

RESULTS

Morphological traits of Ludwigia species

The qualitative traits observed in the species collected, namely the color of flowers

and roots and the pneumatophore form, were consistent with those reported for these species

in literature (see Table S1 for traits and authors).

For the diploid species, red roots, yellow flowers, and rare cylindric pneumatophores were

observed in *Lpm*. In contrast, in *Lh*, we observed red roots, creamy white petals with narrow

yellow base, and abundant, clustered conical pneumatophores (Figure 1). For the tetraploid

species, *La* had pink roots, white petals with yellow base, and had few conical

pneumatophores. *Ls* had white roots, petal color light yellow and similar form of

pneumatophores as those of *La*. For the hexaploid species *Lgg*, only roots were observed and

were pink. The decaploid species *Lgh* had white roots, flowers with yellow petals, and few,

long cylindrical pneumatophores per node. Color of roots and pneumatophore number and

form were confirmed in natura for the different populations of *Lpm* and *Lgh* observed (Figure

1).

Genome size and ploidy level

The chromosome numbers were as expected: for both diploids, *Lpm* and *Lh,* 2n = 16;

for both tetraploids *Ls* and *La,* 2n =32; for hexaploid *Lgg*, 2n=48 and for decaploid *Lgh,*

2n=80 (Table 1, Appendix S2). *Ludwigia* spp. exhibited an ~0.77-fold range of C-values. The

lowest value, 0.53 pg/2C, was found in *Lpm* and the highest, 2.9pg/2C, in *Lgh* (Table 1,

Appendix S3). The tetraploid species *Ls* (1.07pg/2C) and *La* (1.06pg/2C) have C-values that

were twice that the value for the diploid *Lpm* (0.53pg/2C) and *Lh* (0.55pg/2C). The hexaploid

species *Lgg* had C-value 1.77pg/2C. Thus, the genome size by ploidy level revealed that the

monoploid genome sizes (1Cx-value, 0.133-0.147 pg) of the tetraploid, hexaploid, and

decaploid species are the same (0.34-0.49 pg/1Cx). The difference is accounted for by the

higher ploidy levels.

Ludwigia genome sizes of diploid and tetraploid species were similar between species

with the same ploidy level and varied proportionally with ploidy levels (i.e., 2x≈260 Mb, 4x≈

500 Mb; Table 1, Appendix S3). The genome size of hexaploid and decaploid species were

closer than those expected with regard to ploidy level (i.e., ratio (6x/2x) = 1.07; ratio (10x/2x)

= 1.06; Table 1) with 864 Mb and 1419 Mb, respectively.

Genomic relationships using the GISH technique

For the diploid species, when we hybridized slides of *Lpm* with a *Lpm* probe (red) and *Lh*

blocking DNA (grey), 16 chromosomes were tagged in red signals and zero chromosome

showed a grey signal (Figure 2A). Thus, the *Lh* blocking DNA did not block any sequence

present in the *Lpm* probe, meaning that no *Lh* genome was shared with *Lpm*. But, when slides

of *Lh* were hybridized with a *Lh* probe and *Lpm* blocking DNA, ten chromosomes of *Lh*

showed grey signal corresponding to *Lpm* chromosomes (Figure 2B). This observation

suggests genome homology with the *Lpm* genome but four chromosomes were stained in red,

meaning that there are nevertheless differences in *Lpm* and *Lh* genomes. Due to the absence

of chromosomes marked by *Lh* blocking DNA in *Lpm*, we can suggest that even some

homology exist, *Lpm* and *Lh* most likely correspond to different genomes, arbitrarily noted A

for *Lpm* and B for *Lh*.

For the tetraploid species *Ls* and *La*, we hybridized *Ls* slides with a *Ls* probe and three

different blocking DNA combinations from species having different ploidy levels – *Lpm* (2x),

Lh (2x) and *La* (4x) – and for *La* slides, with a *La* probe and *Lh* blocking DNA (Table 2,

Figure 3). When *Lpm* DNA was hybridized over *Ls*, the blocking DNA *Lpm* blocked 16

chromosomes (grey) and the other 16 chromosomes tagged in red by the *Ls* probe (Figure

3A). A similar result was obtained with the blocking DNA of *Lh*, with 16 chromosomes

showing red signals and 16 grey (Figure 3B). Thus, the tetraploid *Ls* would be the result of an

interspecific hybridization between the two diploid species *Lpm* and *Lh*. Based on the genome

naming proposed here, the genomic composition of *L. stolonifera* could be AABB.

300   After use of *La* blocking DNA over *Ls* chromosomes, we observed 16 chromosomes

tagged in red and 16 chromosomes tagged in grey (Figure 3C). The hybridization performed

with *Lh* blocking DNA on the second tetraploid, *La*, identified 16 red chromosomes and 16

grey chromosomes (Figure 3D). Both results suggested that the two tetraploid species *La* and

Ls shared a same genome coming from *Lh* (BB component). Thus, *Lh* would also be one of

the components of the tetraploid *La*, with a XXBB putative genome composition, where the

XX genome corresponds to an unknown *Ludwigia* diploid species.

307   For the hexaploid species *Lgg*, slides of *Lgg* were hybridized with a *Lgg* probe and

four blocking DNA of different ploidy levels – *Lpm* (2x), *Lh* (2x), *Ls* (4x), *La* (4x), and *Lgh*

(10x) Table 2). The *Lpm* competitor DNA blocked 16 chromosomes (tagged in grey) and 32

chromosomes showing red signals were hybridized with the *Lgg* probe DNA (Figure 4 A). A

similar hybridization was obtained with the *Ls* blocking DNA in which slides of *Lgg* had 16

grey chromosomes and 32 chromosomes with red signals (Figure 4 B). Thus, the hexaploid

species *Lgg* contains an identical genomic component found in *Ls* (4x) and in *Lpm* (2x; i.e.,

AA genomic part).

315   Hybridizations performed on slides of *Lgg* with *Lh* (2x) and *La* (4x) blocking DNA

316   exhibited hybridization profiles that were more challenging to interpret with 48 red

317   chromosomes, but with different hybridization intensities (with 16 more intense signals with

318   *La* blocking DNA and 8 less intense signals with *Lh* blocking DNA (Table 2, Figures 4C,

319   4D). The 16 more intense signals could correspond to a 2x component (16 chromosomes)

320   specific to *Lgg*.

321   For the decaploid species, *Lgh*, slides were hybridized with a *Lgh* probe and five

blocking DNA of different ploidy levels, including *Lpm*, *Lh*, *Ls*, *La* and *Lgg*, respectively

(Table 2). The *Lpm* DNA competitor blocked 32 chromosomes with grey signals whereby 48

chromosomes showing red signals (Figure 5A). An identical hybridization result was obtained

with the *Ls* blocking DNA with 48 chromosomes with red signals and 32 grey chromosomes

(Figure 5C). Thus, the 2x component, *Lp*, also present in *Ls* (4x), is found in a double dose

(32 chromosomes) in *Lgh* (10x). The results obtained with the *Lh* and *La* DNA blocking

showed 80 red chromosomes but 16 with lower intensity (Table 2, Figures 5B, 5D). After

GISH hybridization of *Lgg* (6x) DNA on *Lgh* (10x) chromosomes, 32 of 80 *Lgh*

chromosomes showed a red signal (Figure 5E). This result revealed that *Lgg* was probably

one of progenitors of *Lgh*.

Interspecific hybridization

Interspecific hybridization between species sharing the AA genome were carried out

and reproductive success was observed by fruit production when the species used as female

possessed the lower ploidy level (Figure 6, Table S2). No fruits were obtained after crosses

between *Ls* (4x) used as female and *Lp* (2x) used as male or between *Lgh* (10x) used as

female and *Lpm* (2x) or *Ls* (4x) used as male. Thus, all interspecific crosses with the diploid

species *Lpm* (2x) used as female and *Ls* (2x) or *Lgh* (10x) used as male gave fruits showing

similar weight and length (Figure 6, Table S2). The fruits obtained from the *Lpm* (2x) x *Lgh*

(10x) crosses had very large seeds whose development led to the bursting of the fruit walls

(Figure S5). However, only 53.4% and 3.9% of seeds from *Lpm* (2x) x *Ls* (4x) and *Lpm* (2x) x

Lgh (10x) crosses germinated. If all germinated seeds gave plantlets for *Lpm* (2x) x *Ls* (2x)

crosses, only three plants developed for *Lpm* (2x) x *Lgh* (10x). Finally, no plants survived 90

days after seedling, as all plants showed chlorotic signs and at the end of the observation

period, they were not able to survive (Figure 6, Table S2, Figure S3). Similarly, fruits were

produced after *Ls* (4x) x *Lgh* (10x) crosses with a mean number of seeds per fruit of 23.5

(Figure 6, Table S2) but no seed has germinated. Unfortunately, chlorotic plants from *Lpm*

(2x) x *Ls* (4x) and *Lpm* (2x) x *Lgh* (10x) crosses did not develop enough roots for

chromosome observations. For control intraspecific crosses *Lpm* x *Lpm*, *Lgh* x *Lgh* and *Ls* x

Ls, all crosses produced fruits revealing effectiveness of the greenhouse pollination

conditions.

DISCUSSION

To better understand the evolutionary history of genus Luwigia, we have evaluated the

genomic relationships between diploid and polyploid species using the molecular cytogenetic

and crossing investigations.

Validation of Ludwigia species sect. Jussieae studied and identification of new
discriminating traits.

Wagner et al. (2007) summarized the complex history of the Onagraceae. The genus

Ludwigia forms a lineage separate from the rest of the Onagraceae family (Eyde, 1981, 1982)

The long-standing taxonomic confusion surrounding aquatic *Ludwigia* species required a

approach combining morphometric and cytogenetic evaluations to differentiate the species

and improve taxonomic identification (Grewell et al., 2016). Furthermore, distinguishing

Ludwigia species represents a real challenge.

In this study, qualitative morphological traits were observed for the six Lg ssp. grown

in a common garden, allowing compare these species growing under the same conditions. Our

results confirmed that all plants collected corresponded to the expected species. However, our

cross observations of the different species in a common garden revealed additional differences

between these species. For example, the red roots of *Lpm* were never described before, but are

visible on the seedlings as soon as the seeds germinate until the plant reaches maturity in

natura (Appendix S5). *Lh* plants studied had these same characteristics as those described

(Rocha and Melo, 2020), but the petals were more creamy-white than white and were sharply

narrow at the petiole. We found that the pneumatophore form, petal and root coloration could

also differentiate these species (Figure 1). For the tetraploid species, flowers of *La* are

described as creamy white petals with yellow at the base (Wagner et al., 2007) but we

observed white petals similar to *Lh* (Appendix S6). As *Ls* had light yellow petals, the flower

color may be a good characteristic with which to distinguish these two *tetraploid* species in

natura (Appendix S4). For the hexaploid species *Lgg*, we only saw pink roots and more

morphological investigations are required. Finally, the decaploid species *Lgh* had white roots

and bright yellow petals (Figure 1).

Grewell et al., (2016) reported that distinguish in field *Lgg* and *Lgh* was complicated.

Nesom and Kartesz (2000) suggested that few morphological distinctions between *Lgh* and

Lgg exist and broadly overlapping: plants with larger leaves and flowers and less dense

vestiture characterize *Lgh*, whereas smaller leaves and flowers and denser vestiture would

describe *Lgg*. However, comparing flower morphology in sterile and fertile French *Lgh*

populations, two flower sizes were observed which may call into question the criterion for

distinguishing flower size between *Lgh* and *Lgg* (Appendix S5, Portillo-Lemus et al., 2021).

As for the distinction between *Lpm* and *Lgh*, the differences in stipule shape are often cited,

reniform for *Lpm* and oblong and acuminate for *Lgh* (Thouvenot et al., 2013), but this

character is also not easily used. For all these reasons, we propose new criteria to help field

managers based on the root color. *Lpm* has red roots, whereas *Lgh* has white roots.

Importantly, this character can be observed at different stages of plant development

(Appendix S5). *Lgg* seems to have pink roots at a young plant stage. Whether this

characteristic is also true at all stages of *Lgg* development, it could also be a promising way to

distinguish *Lgg* and *Lgh*.

Genomic relationships and origins of polyploids in section Jussieae

We propose the first hypotheses regarding diploid-polyploid relationships of *Ludwigia* diploid

to decaploid species belonging to the section *Jussiaea* (Figure 6). The diploid species studied

here were composed of two different genomes, we have called AA and BB for *Lpm* and *Lh*,

respectively. Both diploid *Lpm* and *Lh* were the progenitors of *Ls,* with the latter composed of

AABB (Figure 3). We also found that *Lh* was a progenitor of *La* (BB), sharing same genome

with *Ls* even though the *La*, native to Asian-Pacific, and *Ls*, native to African, do not

currently co-occur (Table S1). Our results are in agreement with the phylogenetic analysis of

Liu et al. (2017) who suggested through analysis of nuclear DNA sequences that *Lp* or a close

relative contributed to the origin of *Ls* and shared a same genome (here designated as genome

AA). Similarly, Liu et al (2017) reported that *L. adscendens* (4x) is close to *L.*

helminthorrhiza (2x) (genome BB). GIS analysis revealed that *Lh* and *Ls* shared at least one

genome, which was not shown by Liu et al (2017) phylogenetic analysis.

411   Furthermore, considering the genome sizes of both diploid species *Lpm* and *Lh* and

assuming additivity, our genome size data fit perfectly with our scenarios of tetraploid *Ls* and

La origin. On the other hand, we showed that *Lpm* also participated for one part (2x) to the

origin of the hexaploid *Lgg* genome. The decaploid species *Lgh* seems to have emerged from

interspecific hybridization and allopolyploidization events between the tetraploid species *Ls*

(4x) and the hexaploid species *Lgg*. Liu et al. (2017) also demonstrated a close relationship

between *Lgg* and *Lgh* using nuclear and chloroplast DNA regions as molecular markers. In

addition, *Lgh* shares the same pneumatophore form as *Lpm* and the same root color as *Ls*,

which may provide further evidence that both species are progenitors of *Lgh*.

420   All chromosomes of *Lgg* and *Lgh* were tagged by *Lh* blocking DNA, but had strong or

light hybridization intensities for 16 chromosomes respectively. The intensity of fluorescence

could be explained by repetitive sequences shared among closely related species or specific

for given species. Thus, Liu et al. 2008 could distinguish the subgenomes of Triticeae

allopolyploids due to differences in element abundance and the resulting probe signal

intensity. In addition, in a Silene hybrid, Markova et al, 2007 showed that the intensity of

fluorescence varied quantitatively based on the relatedness of the species. These results may

suggest genome divergence between *Lgg* or *Lgh* and *Lh*. The intensity level of the signal over

the majority of the chromosomes likely indicates a mixing of genomic sequences between

parental genomes, in particular for the *Lh* genome (BB), in the hexaploid and decaploid

formation. The effectiveness of GISH is much reduced, with clear evidence of considerable

mixing of genomic sequence between parental DNA. Lim et al. (2007) have shown that within

1 million years of allopolyploid *Nicotiana* divergence, there is considerable exchange of

repeats between parental chromosome sets. After *c.*5 million years of divergence GISH fails.

Repetitive sequences, including dispersed repeats, such as transposable elements (Tes), or

tandem repeats such as satellite DNAs, represent an important fraction of plant genomes that

impact evolutionary dynamics (Vicient and Casacuberta, 2017; Giraud et al., 2021). Yet, no

exhaustive investigations have been undertaken to evaluate the nature and dynamics of

repetitive sequences between different species of *Ludwigia* that probably diversified since

hexaploid and decaploid events when the *Ludwigia* family originated at least 50 m.y. ago

(Raven and Tai, 1979).

***Success of interspecific hybridization and contribution to origin of* Ludwigia**

species, sect. Jussieae

In addition to these results, interspecific crosses between *Ludwigia* species sharing the

A genome produced fruits only when the female parent possessed lower ploidy level

suggesting that efficiency of pollination was possible through the presence of the same

genome in both species. In interspecific crosses differences also exist according to the ploidy

level of the female parent. For example in *Brassica ssp.*, more hybrids formed when

allotetraploid species, *Brassica napus* is used as female in crosses with diploid species used as

male (Kerlan et al., 1992). In contrast, several crosses between *Triticum aestivum* L. and

diploid wild relatives were successful when the female parent had the lower chromosome

number (Sharma, 1995). Liu et al (2017) observed through the cp tree analysis that *La* and *Ls*

are grouped suggesting that both decaploid species shared at least one maternally inherited

genome, probably the BB genome from *Lh*. Unfortunately, *Lh* was not include in cp tree

analysis by Liu et al (2017). The combined data from the interspecific crosses carried out in

this study and the phylogenetic analysis carried out by Liu et al (2017) allows us to

hypothesize that in *Ludwigia* sp. sect. Jussieae, interspecific hybrids can be obtained when the

species used as a female has the lowest ploidy level.

Natural hybrids within section *Jussieae* have been reported between *La* (2n = 4x= 32)

and *L. peploides* subsp. *stipulacea* (2n= 2x =16), with production of a triploid sterile hybrid

(2n = 3x= 24) named *L. x taiwanensis* (Peng, 1990). Between *Lgg* (2n = 6x = 48) and *Lgh* (2n

= 10x = 80), an octoploid hybrid was produced (2n = 8x = 64) and between *Lgg* (2n = 6x =

48) and *L. hookeri* (2n = 2x = 32), a pentaploid hybrid was produced (2n = 5x = 40) (Zardini

et al., 1991; Zardini and Raven, 1992). For our *Lpm* x *Lgh* crosses, we obtained fruit

production after each pollination. Despite the production of a significant seed number, very

low germination was found, with no viable plants. Dandelot (2004) reported that in France,

hybrids between *Lpm* and *Lgh* have never been recorded in nature, whereas hybrids have been

created under experimental conditions. But if Dandelot (2004) obtained fruit from *Lpm* x *Lgh*

crosses, the ability of seeds to germinate and viability of plantlets were not analyzed. As

found by Dandelot (2004), we found zero fruit production when *Lgh* was used as female.

All interspecific crosses using the lower ploidy of *Ludwigia* ssp. as female were

functional and fruits were produced. But depending on the type of interspecific crosses, no

viable seeds or necrotic plants were obtained. Crosses between related species or parents with

different ploidy are often impossible due to post-zygotic reproductive barriers in which the

hybrid progeny fails to develop or becomes sterile. Thus, in crosses between *B. napus* and a

more distant species such as *Sinapis alba*, the interspecific hybridization efficiency is also

extremely low and embryos need to be rescued using fertilized ovary culture (Chèvre et al.,

1994). This indicated an early abortion of seeds after fertilization and the parental genome

dosage in the endosperm plays an important role for seed collapse.

Interspecific hybrids between *Ludwigia* spp. in section Jussieae seem possible only if

interspecific crosses occur between a female plant with lower ploidy level than male plant,

and probably at a very low success rate in natura. However, observing fruit production is not

enough, thus, we recommend observing seed germination, plantlet viability, plant survival,

and chromosome counts.

CONCLUSION

Thus, in this study we demonstrated the interest of a truly novel combination of data to

identify genomic relationships and origins of polyploids in a poorly understood *Ludwigia*

complex. One way to investigate phylogenetic relationship in a polyploid complex is to use

flow cytometric analyses complemented with chromosome counts, as recently described for

the analysis of the polyploid complex *Linum suffruticosum s.l.* (Linaceae) (Afonso et al.,

2021). Another way involves (i) the use of organellar DNA (chloroplast or nuclear regions) as

molecular markers as it was described for phylogenetic analysis of the genus *Isoëtes* (Pereira

et al., 2019) or the diploid and autohexaploid cytotypes of *Aster amellus* (Mairal et al., 2018);

or (ii) OMICS-data tools as RAD-Seq (restriction site-associated DNA sequencing) as

described in the evolutionary processes of apomictic polyploid complexes on the model

system *Ranunculus* (Karbstein et al., 2022). Thus, the various approaches used in this study,

combining morphological and cytogenetic analyses, in situ hybridization and interspecific

crosses, could constitute a first step towards phylogenetic studies of species belonging to

poorly understood complexes for which there are few genomic resources.

Our results suggest that allopolyploidy played an important role in the evolutionary

history of the *Ludwigia* L., section *Jussieae*, giving rise to complex relationships among

species. However, some species are missing in our analyses as well as in Liu et al. (2017).

The missing species of section *Jussiaea* are the four diploids, *Ludwigia peploides* (Kunth)

P.H.Raven subsp. *glabrescens* (O. Kuntze) P.H.Raven, *Ludwigia peploides* subsp. *peploides*,

Ludwigia peploides subsp. *stipulacea* (Ohwi) P.H.Raven, *Ludwigia torulosa* (Arn.) H.Hara.

and the two tetraploid species, *Ludwigia hookeri* (Micheli) H.Hara, *Ludwigia peduncularis*

(C.Wright ex Griseb.) M.Gómez (Hoch et al., 2015). As part of the phylogenetic relationships

remains unresolved, new GISH experiments should be done with these species, especially to

identify the progenitor of the unknown 2x and 4x genome of *Lgg* and *Lgh*, respectively.

Furthermore, as based on morphological observations, Zardini et al. (1991) suggested that

Lgh may be result of interspecific hybridization between *Lgg* and *L. hookeri*, the tetraploid

species *L. hookeri* could be one of progenitor of missing genomes of *Lgg* and *Lgh* species.

514
Conflict of interest

The authors declare they have no conflict of interest relating to the content of this article.

Acknowledgments

This research was supported by FEDER funds from Région Centre-Val de Loire and by

Agence de l'eau Loire-Bretagne (grant Nature 2045, programme 9025 (AP 2015 9025).

FEDER also financed the doctoral grant of L. Portillo – Lemus. Collections in Alabama were

supported by the Department of Biology at the University of Alabama at Birmingham with

field help from S. Heiser and S. Shainker-Connelly. The authors thank also Biogenouest (the

western French network of technology core facilities in life sciences and the environment,

supported by the Conseil Regional Bretagne). The authors address many thanks to PCI

recommender Malika Ainouche and the two reviewers, Alex Baumel and Karol Marhold for

their constructive and high-quality comments which greatly brush up the article and once

again to Malika Ainouche for accepting to be recommender and organizing an efficient and

constructive peer-review.

Author Contributions

DB, LMP and OC contributed to conception and design of experiments. LPM and DB provide

roots and DNA of all Lg species for *in situ* hybridization and ploidy level analysis. SAKH

collected Lgg samples from USA. VH and OC acquired GISH and cytological data. BD, LMP

and OC carried out analysis and interpretation of data. DB write the draft of this manuscript

and DB, LPM, SAKH and OC revised the manuscript. All authors gave a final approval of the

version to be published.

Supporting Information

Appendix S1: Original images of In situ genomic hybridization analyses of somatic

metaphase chromosomes, a) of *L. peploides* subsp. *montevidensis* 2(n= 2x =16).

corresponding to figure 2 , b) ; of the tetraploid species, *L. stolonifera* and *L. adscendens*

(2n=4x=32) corresponding to figure 3 ; of *L. grandiflora subsp. grandiflora* (2n=6x=48)

corresponding to figure 4 ; from *L. grandiflora* subsp. *hexapetala* (2n=10X=80)

corresponding to figure 5.

Appendix S2: Polyploidy levels of different species of *ludwigia* sp section Jussieae. (**A**)

Ludwigia peploides subsp. *montevidensis* chromosomes (2n=2x=16), (**B**) *Ludwigia*

helminthorrhiza chromosomes (2n=2x=16); (**C**) *Ludwigia stolonifera* chromosomes

(2n=4x=32); (**D**) *Ludwigia adscendens* chromosomes (2n=4x=32); (**E**) *Ludwigia grandiflora*

subsp. *grandiflora* (2n=6x=48); *Ludwigia grandiflora* subsp. *hexapetala* (2n=10x=80).

Chromosome number correspond to ploidy level: 16 chromosomes for diploid species (**A**) and

(**B**); 32 chromosomes for tetraploid species (**C**) and (**D**); 48 chromosomes for hexaploid

species (**E**) and 80 chromosomes for decaploid species (F)

Appendix S3: Flow Cytometry results (**A**) and examples of peak profiles (logarithmic) in the

flow cytometer of nuclei stained from roots with propidium iodide (PI) (**B**). The 'trifolium

repens' peak (1C=1,12 pg) or "Zea mays" peak (1C=2,77 pg) is used as internal standard to

determinate the DNA contents of the sample nuclei (*). (1) *Ludwigia peploides* subsp.

montevidensis; (2) *L. helminthorrhiza;* (3) *L. adscendens*; (4) *L. grandiflora* subsp.

grandiflora and (5) *L. grandiflora* sp. *Hexapetala.* [1] : 1 pg DNA = 978 Mbp (from Doležel et

al. 2003) ; [2] : Zonneveld et al, 2019

Appendix S4: Fruit production and seedling from interspecific hybridization between

ludwigia species possessing A genome; *Lpm = Ludwigia* peploides subsp. montevidensis

(2n=16, AA); *Ls = Ludwigia stolonifera* (2n=32, AABB); *Lgh = Ludwigia grandiflora* subsp.

hexapetala (2n=80, AAAABBXXXX/XXYY). (a) the seeds produced from *Lpm* x *Lgh*

interspecific cross are large, which has led to the fruit bursting. (b) 30 days after seedling,

green plantlets from *Lpm* x *Ls* interspecific cross were obtained. But, 60 days later, plants

showed chlorotic development, stopped growing and died. Das: Number of day after seedling

Appendix S5: Morphological traits to distinguish *Ludwigia peploides* subsp. *montevidensis*

and *Ludwigia grandiflora* subsp. *hexapetala,* (**a**) roots at seedling stage ; (**b**) adult roots in

natura ; (**c**) pneumatophores in natura ; (**d**) flowers.

Appendix S6: Size and color of *Ludwigia* sp. flowers. **a**: Flower of *L. grandiflora* subsp.

hexapetala in sterile population (10x), **b**: Flower of *L. grandiflora* subsp. *hexapetala* in fertile

population (10x), **c**: Flower of *L. peploides* subsp. *montevidensis* (2x), **d**: Flower of *L.*

adscendens (4x) and **e**: Flower of *L. stolonifera* (4x)

References

Afonso, A., J. Loureiro, J. Arroyo, E. Olmedo-Vicente, and S. Castro. 2021. Cytogenetic
diversity in the polyploid complex Linum suffruticosum sl. (Linaceae). *Botanical*
Journal of the Linnean Society 195: 216-232.

Akiyama, R., J. Sun, M. Hatakeyama, H. E. Lischer, R. V. Briskine, A. Hay, X. Gan, et al.
2021. Fine☐scale empirical data on niche divergence and homeolog expression patterns in
an allopolyploid and its diploid progenitor species. *New Phytologist* 229: 3587–3601.

Alix, K., P. R. Gérard, T. Schwarzacher, and J. S. Heslop-Harrison. 2017. Polyploidy and
interspecific hybridization: partners for adaptation, speciation and evolution in plants.
Annals of botany 120: 183–194.

Allario, T., J. Brumos, J. M. COLMENERO☐FLORES, D. J. Iglesias, J. A. Pina, L. Navarro,
M. Talon, et al. 2013. Tetraploid Rangpur lime rootstock increases drought tolerance via
enhanced constitutive root abscisic acid production. *Plant, cell & environment* 36: 856–
868.

Armitage, J. D., K. Könyves, J. P. Bailey, J. C. David, and A. Culham. 2013. A molecular,
morphological and cytological investigation of the identity of non-native Ludwigia
(Onagraceae) populations in Britain. *New Journal of Botany* 3: 88–95.

Baniaga, A. E., H. E. Marx, N. Arrigo, and M. S. Barker. 2020. Polyploid plants have faster
rates of multivariate niche differentiation than their diploid relatives. *Ecology Letters* 23:
68–78.

Baumel, A., M. L. Ainouche, R. J. Bayer, A. K. Ainouche, and M. T. Misset. 2002. Molecular
Phylogeny of Hybridizing Species from the Genus Spartina Schreb. (Poaceae). *Molecular*
Phylogenetics and Evolution 22: 303–314.

Blaine Marchant, D., D. E. Soltis, and P. S. Soltis. 2016. Patterns of abiotic niche shifts in
allopolyploids relative to their progenitors. *New Phytologist* 212: 708–718.

Boutte, J., L. Maillet, T. Chaussepied, S. Letort, J.M. Aury, C. Belser, F. Boideau, A. Brunet,
O. Coriton, G. Deniot, C. Falentin, V. Huteau, M. Lodé-Taburel, J. Morice, G. Trotoux,
A.M. Chèvre, M. Rousseau-Gueutin and J. Ferreira de Carvalho. 2020. Genome Size
Variation and Comparative Genomics Reveal Intraspecific Diversity in *Brassica rapa*.
Front Plant Sci 11:577536. doi: 10.3389/fpls.2020.577536.

Chèvre, A. M., F. Eber, E. Margale, M. C. Kerlan, C. Primard, F. Vedel, M. Delseny, and G.
Pelletier. 1994. Comparison of somatic and sexual *Brassica napus – Sinapis alba* hybrids
and their progeny by cytogenetic studies and molecular characterization. *Genome* 37: 367–
374.

Dandelot, S. 2004. Les Ludwigia spp. Invasives du Sud de la France: Historique,
Biosystématique, Biologie et Ecologie. Aix-Marseille 3.

Dolezel, J., J. Bartos, H. Voglmayr, and J. Greilhuber. 2003. Nuclear DNA content and
genome size of trout and human. *Cytometry Part A* 51: 127–8; author reply 129.

Eyde, R. H. 1982. Evolution and Systematics of the Onagraceae: Floral Anatomy. *Annals of*
the Missouri Botanical Garden 69: 735–747.

Eyde, R. H. 1981. Reproductive Structures and Evolution in Ludwigia (Onagraceae). III.
Vasculature, Nectaries, Conclusions. *Annals of the Missouri Botanical Garden* 68: 379–
412.

Eyde, R. H. 1977. Reproductive structures and evolution in Ludwigia (Onagraceae). I.
Androecium, placentation, merism. *Annals of the Missouri Botanical Garden*: 644–655.

Garsmeur, O., J. C. Schnable, A. Almeida, C. Jourda, A. D'Hont, and M. Freeling. 2014. Two
evolutionarily distinct classes of paleopolyploidy. *Molecular biology and evolution* 31:
448–454.

Giraud, D., O. Lima, V. Huteau, O. Coriton, J. Boutte, A. Kovarik, A. R. Leitch, et al. 2021.
Evolutionary dynamics of transposable elements and satellite DNAs in polyploid Spartina
species. *Plant Science* 302: 110671.

Glover, R., R. E. Drenovsky, C. J. Futrell, and B. J. Grewell. 2015. Clonal integration in
Ludwigia hexapetala under different light regimes. *Aquatic Botany* 122: 40–46.

Grewell, B. J., M. D. Netherland, and M. J. Skaer Thomason. 2016. Establishing research and
management priorities for invasive water primroses (Ludwigia spp.). *U.S. Army Corps of*
Engineers, Engineer Research and Development Center/Environmental Laboratory;
Vicksburg, MS, USA.

Hieda, S., Y. Kaneko, M. Nakagawa, and N. Noma. 2020. Ludwigia grandiflora (Michx.)
Greuter & Burdet subsp. hexapetala (Hook. & Arn.) GL Nesom & Kartesz, an invasive
aquatic plant in Lake Biwa, the largest lake in Japan. *Acta Phytotaxonomica et*
Geobotanica 71: 65–71.

Hoch, P. C., W. L. Wagner, and P. H. Raven. 2015. The correct name for a section of
Ludwigia L. (Onagraceae). *PhytoKeys* 50: 31.

Hoch, P. C., J. V. Crisci, H. Tobe, and P. E. Berry. 1993. A Cladistic Analysis of the Plant
Family Onagraceae. *Systematic Botany* 18: 31–47.

Hollister, J.D. 2015. Polyploidy: adaptation to the genomic environment. *New Phytol* 205:
1034–1039.

Husband, B. C., S. J. Baldwin, and J. Suda. 2013. The incidence of polyploidy in natural plant
populations: major patterns and evolutionary processes. *Plant genome diversity* 2: 255–
276. Springer.

Karbstein, K., S.Tomasello, L. Hodač, N.Wagner, P. Marinček, B. H., Barke, C. Paetzold, et
al. 2022. Untying Gordian knots: Unraveling reticulate polyploid plant evolution by
genomic data using the large Ranunculus auricomus species complex. *New Phytologist*
235: 2081-2098.

Kerlan, M. C., A. M. Chèvre, F. Eber, A. Baranger, and M. Renard. 1992. Risk assessment of
outcrossing of transgenic rapessed to related species: I. Interspecific hybrid production

under optimal conditions with emphasis on pollination and fertilization. *Euphytica* 62:
145–153.

Kim, H. W., D. C. Son, S. H. Park, C. Jang, E. Sun, H. Jo, S. M. Yun et al. 2019. Unrecorded
alien plant on South Korea: Ludwigia peploides subsp.montevidensis (Spreng). P.H.
Raven. *Korean journal of Plant Research* 32: 201-206.
663

Levin, R. A., W. L. Wagner, P. C. Hoch, W. J. Hahn, A. Rodriguez, D. A. Baum, L. Katinas,
et al. 2004. Paraphyly in Tribe Onagreae: Insights into Phylogenetic Relationships of
Onagraceae Based on Nuclear and Chloroplast Sequence Data. *Systematic Botany* 29: 147–
164.

Levin, R. A., W. L. Wagner, P. C. Hoch, M. Nepokroeff, J. C. Pires, E. A. Zimmer, and K. J.
Sytsma. 2003. Family-level relationships of Onagraceae based on chloroplast rbcL and
ndhF data. *American Journal of Botany* 90: 107–115.

Lim, K. Y., R. Matyasek, A. Kovarik, and A. Leitch. 2007. Parental Origin and Genome
Evolution in the Allopolyploid Iris versicolor. *Annals of Botany* 100: 219–224.

Liu, S.-H., P. C. Hoch, M. Diazgranados, P. H. Raven, and J. C. Barber. 2017. Multi-locus
phylogeny of Ludwigia (Onagraceae): Insights on infra- generic relationships and the
current classification of the genus. *TAXON* 66: 1112–1127.

Liu, Z., W. Yue, D. Li, R.R. Wang, X. Kong, K. Lu, G. Wang, Y. Dong, W. Jin, and X.
Zhang. 2008. Structure and dynamics of retrotransposons at wheat centromeres and
pericentromeres. Chromosoma 117:445-56. doi: 10.1007/s00412-008-0161-9.
679

Lobato-de Magalhães, T., K. Murphy, A. Efremov, V. Chepinoga, T. A. Davidson, and E.
Molina-Navarro. 2021. Ploidy state of aquatic macrophytes: Global distribution and
drivers. *Aquatic Botany* 173: 103417.

Mairal, M., M. Šurinová, S. Castro, and Z. Münzbergová. 2018. Unmasking cryptic
biodiversity in polyploids: origin and diversification of Aster amellus aggregate. *Annals of*
Botany 122:1047-1059

Markova M, E. Michu, B. Vyskot, B. Janousek, and J. Zluvova. 2007. An interspecific hybrid
as a tool to study phylogenetic relationships in plants using the GISH technique.
Chromosome Res. 15:1051-9. doi: 10.1007/s10577-007-1180-8.

McIntyre, P. J. 2012. Polyploidy associated with altered and broader ecological niches in the
Claytonia perfoliata (Portulacaceae) species complex. *American Journal of Botany* 99:
655–662.

Méndez Santos, I. E. M., and R. González-Sivilla. 2020. Expansión de Ludwigia
helminthorrhiza (Onagraceae) en Cuba. *Anales del Jardín Botánico de Madrid* 77: 7.

Munz, P. A. 1942. Studies in Onagraceae XII: A Revision of the New World Species of
Jussiaea. *Darwiniana* 4: 179–284.

Nesom, G. L., and J. T Kartesz. 2000. Observations on the Ludwigia uruguayensis Complex
(Onagraceae) in the United States. *Castanea* 65: 123–125.
698

Okada, M., B. J. Grewell, and M. Jasieniuk. 2009. Clonal spread of invasive Ludwigia
hexapetala and L. grandiflora in freshwater wetlands of California. *Aquatic Botany* 91:
123–129.

Otto, S. P., and J. Whitton. 2000. Polyploid incidence and evolution. *Annual review of*
genetics 34: 401–437.

Peng, C. I. 1990. Ludwigia taiwanensis (Onagraceae), a new species from Taiwan, and its
origin. *Botanical Bulletin of Academia Sinica* 31: 343–349.*genetics* 6: 836–846.

Pereira, J. B., P. H. Labiak, T. Stützel, and C. Schulz. 2019. Nuclear multi-locus phylogenetic
inferences of polyploid Isoëtes species (Isoëtaceae) suggest several unknown diploid
progenitors and a new polyploid species from South America. *Botanical Journal of the*
Linnean Society 189: 6-22.
710
Portillo-Lemus, L. O., M. Bozec, M. Harang, J. Coudreuse, J. Haury, S. Stoeckel, and D.
Barloy. 2021. Self-incompatibility limits sexual reproduction rather than environmental
conditions in an invasive water primrose. *Plant-Environment Interactions* 2: 74–86.

Ramamoorthy, T. P. 1979. A Sectional Revision of Ludwigia Sect. Myrtocarpus S. Lat.
(Onagraceae). *Annals of the Missouri Botanical Garden* 66: 893–896.

Raven, P. H. 1963. The old world species of Ludwigia (including hissiaea), with a synopsis of
the genus (Onagraceae). *REINWARDTIA* 6: 327–427.

Raven, P. H., and W. Tai. 1979. Observations of Chromosomes in Ludwigia (Onagraceae).
Annals of the Missouri Botanical Garden 66: 862–879.

Reddy, A. M., P. D. Pratt, B. J. Grewell, N. E. Harms, G. Cabrera Walsh, M. C. Hérnandez,
A. Faltlhauser, and X. Cibils-Stewart. 2021. Biological control of invasive water
primroses, Ludwigia spp., in the United States: A feasibility assessment. *J. Aquat. Plant.*
Manag.

Rocha, A. M., and J. I. M. de Melo. 2020. Diversity and distribution of Ludwigia
(Onagraceae) in Paraíba State, Northeastern Brazil. *European Journal of Taxonomy*.

Sharma H. C. 1995. How wide can a wide cross be? Euphytica 82: 43-64.

Soliman, A. T., R. S. Hamdy, and A. B. Hamed. 2018. Ludwigia stolonifera (Guill. & Perr.)
P.H. Raven, Insight into its Phenotypic Plasticity, Habitat Diversity and Associated
Species. *Egyptian Journal of Botany* 58: 605–626.

Soltis, P. S., D. B. Marchant, Y. Van de Peer, and D. E. Soltis. 2015. Polyploidy and genome
evolution in plants. *Current opinion in genetics & development* 35: 119–125.

Stebbins, G. L. 1985. Polyploidy, hybridization, and the invasion of new habitats. *Annals of*
the Missouri Botanical Garden: 824–832.

Te Beest, M., J. J. Le Roux, D. M. Richardson, A. K. Brysting, J. Suda, M. Kubešová, and P.
Pyšek. 2012. The more the better? The role of polyploidy in facilitating plant invasions.
Annals of botany 109: 19–45.

Thouvenot, L., J. Haury, and G. Thiebaut. 2013. A success story: Water primroses, aquatic
plant pests. *Aquatic Conservation: Marine and Freshwater Ecosystems* 23.

Van de Peer, Y., T.-L. Ashman, P. S. Soltis, and D. E. Soltis. 2021. Polyploidy: an
evolutionary and ecological force in stressful times. *The Plant Cell* 33: 11–26.

Vicient, C. M., and J. M. Casacuberta. 2017. Impact of transposable elements on polyploid
plant genomes. *Annals of Botany* 120: 195–207.

Wagner, W. L., P. C. Hoch, and P. H. Raven. 2007. Revised classification of the Onagraceae.
Systematic Botany Monographs.

Wani, G. A., M. A. Shah, Z. A. Reshi, and M. A. Dar. 2018. Polyploidy determines the stage
of invasion: clues from Kashmir Himalayan aquatic flora. *Acta Physiologiae Plantarum*
40: 58.

Wei, N., R. Cronn, A. Liston, and T.-L. Ashman. 2019. Functional trait divergence and trait
plasticity confer polyploid advantage in heterogeneous environments. *New Phytologist*
221: 2286–2297.

Zardini, E. M., H. Gu, and P. H. Raven. 1991. On the Separation of Two Species within the
Ludwigia uruguayensis Complex (Onagraceae). *Systematic Botany* 16: 242–244.

Zardini, E., and P. H. Raven. 1992. A New Section of Ludwigia (Onagraceae) with a Key to
the Sections of the Genus. *Systematic Botany* 17: 481–485.

Zonneveld, B.J. 2019. The DNA weights per nucleus (genome size) of more than 2350
species of the Flora of The Netherlands, of which 1370 are new to science, including the
pattern of their DNA peaks. *Forum Geobotanicum* **8**: 24–78.

Tables

Table 1: Ploidy levels, chromosome numbers and genome sizes estimated by flow cytometry

in *Ludwigia* L. spp. sect. *Jussiaea*.

Species names are mentioned according to the revised nomenclature by Hoch et al. (2015).

Genome sizes were converted from picograms (pg) to Megabases (Mb) using 1 pg = 978

Mbp.

Species name	Ploidy and chromosome numbers	DNA nucleao content (1C in pg)	Genome size (Mb)
Ludwigia peploides subsp. *montevidensis (Lpm)*	2n= 2x = 16	0.265	262
Ludwigia helminthorrhiza (Lh)	2n= 2x = 16	0.275	268
Ludwigia adscendens (La)	2n= 4x = 32	0.53	520
Ludwigia stolonifera (Ls)	2n= 4x = 32	0.535	522
Ludwigia grandiflora subsp. *grandiflora (Lgg)*	2n= 6x = 48	0.885	864
Ludwigia grandiflora subsp. *hexapetala (Lgh)*	2n= 10x = 80	1.045	1419

Table 2: Results of GISH with different *Ludwigia* L. probes (red) combined with blocking DNA (grey) on *L. peploides* subsp. *montevidensis* (Lpm), *L. helminthorrhiza* (Lh), *L. adscendens* (La), *L. grandiflora* subsp. *grandiflora* (Lgg) and *L. grandiflora* subsp. *hexapetala* (Lgh) chromosomes.

Chromosomes of one species tagged in red correspond to DNA of this species and chromosomes tagged in grey are blocked by DNA of others species.

Chromosomes / Blocking DNA	Lpm (2n = 16)	Lh (2n = 16)	Ls (2n = 32)	La (2n= 32)	Lgg (2n= 48)	Lgh (2n= 80)
Lpm (2n = 16)		4 red signals + 10 grey signals	16 red signals + 16 grey signals		32 red signals + 16 grey signals	48 red signals + 32 grey signals
Lh (2n = 16)	16 red signals		16 red signals + 16 grey signals	16 red signals + 16 grey signals	48 red signals (8 less intense)	80 red signals (16 less intense)
Ls (2n = 32)					32 red signals + 16 grey signals	48 red signals + 32 grey signals
La (2n= 32)			16 red signals + 16 grey signals		48 red signals (16 more intense)	80 red signals (16 less intense)
Lgg (2n = 48)						32 red signals + 48 grey signals

Table 3: Reproductive success after controlled interspecific crosses between different

Ludwigia L. spp. belonging to the section *Jussiaea*.

Interspecific hybridization (female x male) between the three species, *Ludwigia peploides*

subsp. *montevidensis* (Lpm), *Ludwigia stolonifera* (Ls) and/or *Ludwigia grandiflora* subsp.

hexapetala (Lgh, AAAA BB XXXX/XXYY) used as female or male. All species possess

same genome A: Lpm (2x, AA); Ls (4x, AABB); Lgh (10x, AAAA BB XXXX or XXYY).

Number of plantlets and plants were counted three (21 days) and 8 weeks (56 days) after seed

germination, respectively.　NA: data not available. (+/-= confidence interval, α=0.05). For

control interspecific crosses *Lgh* x *Lgh* and *Lpm* x *Lpm*, a set of randomly selected plantlets

were followed until 56 days after seed germination.

Controlled interspecific crosses	*Lpm* x *Ls*	*Lpm* x *Lgh*	*Ls* x *Lpm*	*Ls* x *Lgh*	*Lgh* x *Lpm*	*Lgh* x *Ls*	*Lgh* x *Lgh*	*Lpm* x *Lpm*
Number of cross pollination	8	25	10	2	10	10	75	45
Number of fruits	8	25	0	2	0	0	75	45
Mean length of fruit (mm)	15.08 (+/- 0.78)	16.64 (+/- 0.82)	/	NA	/	/	7	NA
Mean fruit weight (g)	62.04 (+/- 6.46)	64.64 (+/- 6.02)	/	NA	/	/	NA	NA
Number of total seed	221	1101	/	47	/	/	3750	1980
Number of germinated seeds	118	34	/	0	/	/	3375	1881
Number of plantlets 21 days	118	3	/	0	/	/	3750	1881
Number of plants 56	0	0	/	0	/	/	100 from a set of	50 from a set of 50

days							100	

799
800

 Legends of figures:

 Figure 1: Morphological traits of *Ludwigia* L. species in section *Jussiaea*.

 Ludwigia L. species are classified in a phylogenic tree as proposed by Liu et al (2017). Three

 morphological traits were observed (color of roots, pneumatophore form, color of flower).

 Figure 2: Genomic in situ hybridization (GISH) on mitotic metaphase chromosomes from

 Ludwigia peploides subsp. *montevidensis* (2n= 2x =16) using *Ludwigia peploides* subsp.

 montevidensis probe (2x) (red) and *Ludwigia helminthorrhiza* (2x) (10µg) as blocking DNA

 (A) and from *L. helminthorrhiza* (2n= 2x =16) using *L. helminthorrhiza* probe (2x) and *L.*

 peploides subsp. *montevidensis* (2x) (10µg) as blocking DNA (B).

 Thus, GISH reveals specifically 16 red signals (white stars) and 0 *L. peploides* subsp.

 montevidensis chromosomes (grey) (A) and 4 red signals (white stars) and 10 *L.*

 helminthorrhiza chromosomes (grey) (B). Chromosomes were counterstained with DAPI

 (grey). Bar represents 5 µm.

 Figure 3: Genomic in situ hybridization (GISH) on mitotic metaphase chromosomes from the

 tetraploid species, *Ludwigia stolonifera* and *Ludwigia adscendens* (2n= 4x =32).

 GISH was carried out for *L. stolonifera* using *L. stolonifera* probe (4x) (red) and *Ludwigia*

 peploides subsp. *montevidensis* (2x) (10µg) as DNA blocking (A), *Ludwigia helminthorrhiza*

 (2x) as block (B) and *L. adscendens* (4x) as block (C) and for *L. adscendens* (4x) using *L.*

 adscendens probe (4x) (red) and *L. helminthorrhiza* (2X) (10µg) as block (D). Thus, GISH

 revealed for *L. stolonifera* specifically 16 red signals (white stars) and 16 *L. peploides* subsp.

 montevidensis chromosomes (grey) (A), 16 red signals (white stars) and 16 *L. helminthorrhiza*

 chromosomes (grey) (B), 16 red signals (white stars) and 16 *L. adscendens* chromosomes

 (grey) (C) and for *L. adscendens* 16 red signals (white stars) and 16 *L. helminthorrhiza*

chromosomes (grey) (D). Chromosomes were counterstained with DAPI (grey). Bar

represents 5 µm.

Figure 4: Genomic in situ hybridization (GISH) on mitotic metaphase chromosomes from *L.*

grandiflora subsp. *grandiflora* (2n= 6x =48) using *Ludwigia grandiflora* subsp. *grandiflora*

probe (6x) (red) and *Ludwigia peploides* subsp. *montevidensis* (2x) (A), *Ludwigia*

helminthorriza (2x) (10µg) as block (B), *Ludwigia stolonifera* (4x) (10µg) as block (10µg) as

block (C), *Ludwigia adscendens* (4x) (10µg) as block (D), *Ludwigia grandiflora* subsp.

hexapetala (10x) as block (E).

Thus, GISH reveals specifically 32 red signals (white star) and 16 *L. peploides* chromosomes

(grey) (A), 48 red signals with 8 present less intensity (white star) (B), 32 red signals (white

star) and 16 *L. stolonifera* chromosomes (grey) (C) and 48 red signals with 16 present more

intensity (white star) (D). Chromosomes were counterstained with DAPI (grey). Bar

represents 5 µm.

Figure 5: Genomic in situ hybridization (GISH) on mitotic metaphase chromosomes from

from *Ludwigia grandiflora* subsp. *hexapetala* (2n= 10X =80) using *L. grandiflora* subsp.

hexapetala probe (10x) (red) and *Ludwigia peploides* subsp. *montevidensis* (2x) (10µg) as

block (A), *Ludwigia helminthorrhiza* (2x) as block (B), *Ludwigia stolonifera* (4x) (10µg) as

block (C), *Ludwigia adscendens* (4x) as block (D) and *L. grandiflora* subsp. *grandiflora* (6x)

as block (E).

Thus, GISH reveals specifically 48 red signals and 32 *L. peploides* chromosomes (grey) (A),

80 red signals and 16 present less intensity (white stars) (B), 48 red signals and 32 *L.*

stolonifera chromosomes (grey) (C), 80 red signals and 16 present less intensity (white stars)

(D) and 32 red signals and 48 *L. grandiflora* subsp. *grandiflora* (grey) (E). Chromosomes

were counterstained with DAPI (grey). Bar represents 5 µm.

Figure 6: Hypothetical phylogenetic history of *Ludwigia* L. species of section Jussieae

861 Figure 1: Morphological traits of *Ludwigia* L. species in section *Jussiaea*.

862   *Ludwigia* L. species are classified in a phylogenic tree as proposed by Liu et al (2017). Three

863 morphological traits were observed (color of roots, pneumatophore form, color of flower).

866
867

Figure 2: In situ genomic hybridation analyses of somatic metaphase chromosomes from *Ludwigia peploides* subsp. *montevidensis* (2n=2x=16) using *L. peploides* subsp. *montevidensis* probe (2x) (red) and *Ludwigia helminthorrhiza* (2x) (10µg) as blocking DNA (A) and from *L. helminthorrhiza* (2n=2x=16) using *L. helminthorrhiza* probe (2x) and *L. peploides* subsp. *montevidensis* (2x) (10µg) as blocking DNA (B). Thus, GISH reveals specifically 16 red signals (white stars) and 0 *L. peploides* subsp. *montevidensis* chromosomes (grey) (A) and 4 red signals (white stars) and 10 *L. helminthorrhiza* chromosomes (grey) (B). Chromosomes were counterstained with DAPI (grey). Bar represents 5 µm.

877
878

879
Figure 3: In situ genomic hybridation analyses of somatic metaphase chromosomes of the

tetraploid species, *Ludwigia stolonifera* and *Ludwigia adscendens* (2n=4x=32).

GISH was carried out for *L. stolonifera* using *L. stolonifera* probe (4x) (red) and *Ludwigia*

peploides subsp. *montevidensis* (2x) (10µg) as DNA blocking (A), *Ludwigia helminthorrhiza*

(2x) as block (B) and *L. adscendens* (4x) as block (C) and for *L. adscendens* (4x) using *L.*

adscendens probe (4x) (red) and *L. helminthorrhiza* (2X) (10µg) as block (D) . Thus, GISH

revealed for *L. stolonifera* specifically 16 red signals (white stars) and 16 *L. peploides* subsp.

montevidensis chromosomes (grey) (A), 16 red signals (white stars) and 16 *L. helminthorrhiza*

chromosomes (grey) (B), 16 red signals (white stars) and 16 *L. adscendens* chromosomes

(grey) (C) and for *L. adscendens* 16 red signals (white stars) and 16 *L. helminthorrhiza*

chromosomes (grey) (D). Chromosomes were counterstained with DAPI (grey). Bar

represents 5 µm.

894
895
Figure 4: In situ genomic hybridation analyses of somatic metaphase chromosomes from

Ludwigia grandiflora subsp. grandiflora (2n=6x=48) using *L. grandiflora subsp. grandiflora*

probe (6x) (red) and *Ludwigia peploides* subsp. *montevidensis* (2x) (**A**), *Ludwigia*

helminthorriza (2x) (10µg) as block (**B**), *Ludwigia stolonifera* (4x) (10µg) as block (10µg) as

block (**C**), *Ludwigia adscendens* (4x) (10µg) as block (**D**), *Ludwigia grandiflora* subsp.

hexapetala (10x) as block (**E**).

Thus, GISH reveals specifically 32 red signals (white star) and 16 *L. peploides* chromosomes

(grey) (**A**), 48 red signals with 8 present less intensity (white star) (**B**), 32 red signals (white

star) and 16 *L. stolonifera* chromosomes (grey) (**C**) and 48 red signals with 16 present more

intensity (white star) (**D**). Chromosomes were counterstained with DAPI (grey). Bar

represents 5 µm.

907
908

L. grandiflora subsp. *hexapetala*

L. grandiflora subsp. *hexapetala*

L. grandiflora subsp. *hexapetala*

L. grandiflora subsp. *hexapetala*

L. grandiflora subsp. *hexapetala*

A

B

C

D

E

909
910

Figure 5: In situ genomic hybridation analyses of somatic metaphase chromosomes from *Ludwigia grandiflora* subsp. *hexapetala* (2n=10X=80) using *L. grandiflora* subsp. *hexapetala* probe (10x) (red) and *Ludwigia peploides* subsp. *montevidensis* (2x) (10μg) as block (**A**), *Ludwigia helminthorrhiza* (2x) as block (**B**), *Ludwigia stolonifera* (4x) (10μg) as block (**C**), *L. adscendens* (4x) as block (**D**) and *Ludwigia grandiflora* subsp. *grandiflora* (6x) as block (**E**).

Thus, GISH reveals specifically 48 red signals and 32 *L. peploides* chromosomes (grey) (**A**), 80 red signals and 16 present less intensity (white stars) (**B**), 48 red signals and 32 *L.*

919   *stolonifera* chromosomes (grey) (**C**)*,* 80 red signals and 16 present less intensity (white stars)

920   (**D**) and 32 red signals and 48 *L. grandiflora* subsp. *grandiflora* (grey) (E)*.* Chromosomes

921   were counterstained with DAPI (grey). Bar represents 5 µm.

922
923

924
925
926   Figure 6: Hypothetical phylogenetic history of *ludwigia* species of section Jussieae
927
928
929
930