Indoline CD4-mimetic Compounds Mediate Potent and Broad HIV-1 Inhibition and Sensitization to Antibody-dependent Cellular Cytotoxicity

Christopher Fritschi,[†,1] Saumya Anang,[‡,1] Zhen Gong,[⊥,1] Mohammadjavad Mohammadi,[∇] Jonathan Richard, [¶,§] Catherine Bourassa,[¶] Kenny T. Severino,[#] Hannah Richter,[#] Derek Yang,[†] Hung-Ching Chen,[†] Ta-Jung Chiu,[†] Michael Seaman,[#] Navid Madani,[‡] Cameron Abrams,[∇] Andrés Finzi,[¶,§] Wayne A. Hendrickson,[⊥, +,2] Joseph Sodroski,[‡, §, ‖,2] and Amos B. Smith, III[†,2]

[†] Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States

[‡] Department of Cancer Immunology and Virology, Dana-Farber Cancer Institute and [§]Department of Microbiology, Harvard Medical School, Boston, Massachusetts 02115, United States

[⊥] Department of Biochemistry and Molecular Biophysics and [+]Department of Physiology and Cellular Biophysics, Columbia University, New York, New York 10032, United States

[∇] Department of Chemical and Biological Engineering, Drexel University, Philadelphia, Pennsylvania 19104, United States

[¶] Centre de Recherche du CHUM, Montreal, Quebec, Canada

[§] Departement de Microbiologie, Infectiologie et Immunologie, Universite de Montreal, Montreal, Quebec, Canada

[#] Center for Virology and Vaccine Research, Beth Israel Deaconess Medical Center, Boston, Massachusetts 02215

[‖] Department of Immunology and Infectious Diseases, Harvard School of Public Health, Boston, Massachusetts 02115, United States

[1]These authors contributed equally to this work.

[2]To whom correspondence may be addressed. Email: wah2@cumc.columbia.edu, smithab@sas.upenn.edu, or joseph_sodroski@dfci.harvard.edu.

Author Contributions: C.F., S.A., Z.G., W.A.H., J.S., and A.B.S. designed research; C.F., S.A., Z.G., M.M., J.R., C.B., K.T.S., H.R., D.Y., H.C.C. and T.J.C. performed research; C.F., S.A., Z.G., M.M., J.R., C.B., K.T.S., H.R., D.Y., H.C.C., T.J.C., M.S., N.M., C.A., A.F., W.A.H., J.S. and A.B.S. analyzed data; and C.F., S.A., Z.G., W.A.H., J.S., and A.B.S. wrote the paper..

Competing Interest Statement: The authors declare no competing financial interest.

Keywords: retrovirus, gp120, entry inhibitor, structure-based drug design, small molecule, antiretroviral therapy, antibody-dependent cellular cytotoxicity

This PDF file includes:

Abstract

Binding to the host cell receptors, CD4 and CCR5/CXCR4, triggers large-scale conformational

changes in the human immunodeficiency virus (HIV-1) envelope glycoprotein (Env) trimer

[(gp120/gp41)$_3$] that promote virus entry into the cell. CD4-mimetic compounds (CD4mcs)

comprise small organic molecules that bind in the highly conserved CD4-binding site of gp120

and prematurely induce inactivating Env conformational changes, including shedding of gp120

from the Env trimer. By inducing more "open," antibody-susceptible Env conformations, CD4mcs

also sensitize HIV-1 virions to neutralization by antibodies and infected cells to antibody-

dependent cellular cytotoxicity (ADCC). Here, we report the design, synthesis and evaluation of

novel CD4mcs based on an indoline scaffold. Compared with our current lead indane scaffold

CD4mc, BNM-III-170, several indoline CD4mcs exhibit increased potency and breadth against

HIV-1 variants from different geographic clades. Viruses that were selected for resistance to the

lead indane CD4mc, BNM-III-170, are susceptible to inhibition by the indoline CD4mcs. The

indoline CD4mcs also potently sensitize HIV-1-infected cells to ADCC mediated by plasma from

HIV-1-infected individuals. Crystal structures indicate that the indoline CD4mcs gain potency

compared to the indane CD4mcs through more favorable π-π overlap from the indoline pose and

by making favorable contacts with the vestibule of the CD4-binding pocket on gp120. The rational

design of indoline CD4mcs thus holds promise for further improvements in antiviral activity,

potentially contributing to efforts to treat and prevent HIV-1 infection.

Main Text

Introduction

The human immunodeficiency virus (HIV-1) establishes persistent infections that, if untreated,

lead to the life-threatening acquired immunodeficiency syndrome (AIDS). The HIV-1 pandemic

represents a significant challenge to global health, with 38 million people currently infected and

1.5 million new infections occurring annually (1). Antiretroviral treatments have extended the lives

of infected individuals, but treatment needs to be continued indefinitely to prevent viral rebound

in most cases (2). Furthermore, development of drug-resistant HIV-1 strains and drug side effects

can limit effective treatment options (3). To date, practical measures to prevent HIV-1

transmission by antiretroviral drugs or vaccines remain elusive (2, 3). Additional approaches to

limit HIV-1 replication would complement ongoing efforts to address these challenges.

Entry of HIV-1 into host cells is mediated by the envelope glycoprotein (Env) trimer, which consists

of three gp120 exterior subunits noncovalently associated with three gp41 transmembrane

subunits (4-6). As the only virus-specific protein on the surface of virions and infected cells, Env

serves as a target for host antibodies that neutralize viruses and kill infected cells through

antibody-dependent cellular cytotoxicity (ADCC) (7-11). The unliganded Env largely resides in a

"closed" pretriggered conformation that resists the binding of potentially neutralizing or ADCC-

mediating antibodies elicited during natural infection (12-16). CD4 binding drives Env into more

"open" conformations that engage the CCR5/CXCR4 co-receptor, promoting additional Env

transitions into a gp41 six-helix bundle that fuses the viral and cell membranes (17-22).

The binding site for CD4 is a conserved Env gp120 structure that is conformationally altered by

CD4 binding (23, 24). In turn, CD4 binding creates an internal cavity (the Phe 43 cavity) in Env

bounded by well conserved residues from gp120 and a single phenylalanine residue (Phe 43)

from CD4 (*Figure 1B*). CD4-mimetic compounds (CD4mcs) comprise organic small-molecule

HIV-1 entry inhibitors that bind in the Phe 43 cavity, as well as occupy the gp120 "vestibule"

leading into the cavity (25). Highly conserved gp120 residues in the vestibule make critical

contacts with CD4; therefore, by occupying the gp120 vestibule, the CD4mcs directly compete

with CD4 (26, 27). In addition, CD4mcs prematurely trigger conformational changes in Env similar

to those induced by CD4 (26, 28, 29). In the absence of coreceptor-expressing target cells, these

prematurely activated Envs rapidly and irreversibly become non-functional, in some cases

shedding gp120 (28, 29).   At concentrations that do not completely inhibit HIV-1 infection,

CD4mcs induce "open" conformations of Env, thereby sensitizing HIV-1 virions to neutralization

or HIV-1-infected cells to ADCC by otherwise ineffectual antibodies (30-34). CD4mcs have been

shown to synergize with ADCC-mediating antibodies to decrease the HIV-1-infected cell reservoir

in humanized mice (35). CD4mcs directly protect humanized mice from vaginal acquisition of HIV-

1 and, in combination with vaccine-elicited antibodies, CD4mcs protect monkeys from a stringent

heterologous simian-human immunodeficiency virus (SHIV) mucosal challenge (36, 37).

Early CD4mcs discovered by Debnath and co-workers (27) using a gp120-CD4 screen exhibited

weak antiviral potency against a limited range of HIV-1 isolates. Replacement of the tetramethyl-

piperidine scaffold of these prototypic CD4mcs with an indane scaffold represented a key advance

(24, 37). Rationally designed indane CD4mcs (*Figure 1A and C*) were able to achieve more

contacts with the gp120 vestibule, resulting in increased antiviral potency and breadth (38).

Nonetheless, some primary HIV-1 strains remain resistant even to the most potent indane

CD4mc, BNM-III-170 (**1**). HIV-1 passaged in the presence of BNM-III-170 develops resistance to

this CD4mc as a result of Env changes in the Phe 43 cavity and in a gp120 element that

allosterically modulates CD4 binding (39).

The practical utility of CD4mcs as antiviral agents would be enhanced by the availability of more

potent analogs with better coverage of HIV-1 strains. Guided by crystal structures of BNM-III-170

and congeners thereof, we found that 3-substituents on the indane ring of BNM-III-170 can

contact a vestibule pocket situated at the interface of the gp120 inner and outer domains, between

the α1 and α5 helices (*Figure 1B*). As a result, some 3-substituted indane analogs, such as CJF-

III-049-S (**2**, *Figure 1C*), exhibited modest (3-fold) improvements in antiviral activity compared

with BNM-III-170; however, only the viruses with His 105 in the gp120 α1 helix showed such an

increase (40).

Figure 1. Targeting the Phe 43 cavity and surrounding vestibule of HIV-1 gp120. (**A**) General core of an indane or

indoline CD4mc with the core's atom positions numbered. (**B**) HIV-1$_{BG505}$ gp120 (PDB: 5THR) is displayed with the

crucial structural features for CD4mc design highlighted. The α5 (gold) and α1 (magenta) helices, as well as the β20/21

loop (cyan), harbor residues that contribute to native CD4 binding. Chain E was used for visualization with a 70%

transparent surface placed on gp120. The association of BNM-III-170, from an HIV-1$_{C1086}$ complex (PDB: 5F4P), with

the Phe 43 cavity is shown in the enlarged insert. (**C**) Evolution of more potent CD4mcs from BNM-III-170 (**1**) (38).

Based on the above observations, we hypothesized that replacing the C3 carbon with a nitrogen

(*Figure 1A and C*) to create an indoline scaffold might have several benefits. First, the N3 nitrogen

of the indoline scaffold would permit facile incorporation of diverse functional groups (*vide infra*).

Second, the nitrogen at N3 would be expected to limit conformational freedom of the 5-membered

ring, thereby decreasing the number of conformational states that the oxalamide (at C1) and

methyl guanidinium (at C2) can adopt. Only one state has been observed by X-ray for the indane

CD4mc complexes with gp120 (24, 37, 39), but an entropic penalty would be expected in selecting

this pose upon complexation. Lastly, the rigidified 5-membered ring of the indolines may lead to

different geometries of the substituents at the N3 position, compared to the reported indane

congeners (40). This would permit different branching angles of the substituents, thereby allowing

exploration of broader chemical space, given the highly modular nature of the indoline scaffold.

Results

A. Synthetic Chemistry

Based on the rationale above, we set out to design a synthetic route to obtain the *indoline* CD4mc

framework. As illustrated in *Figure 2A*, anthranilic ester **4** was first protected as a benzyl

carbamate. This was followed by one-pot alkylation with methyl bromoacetate (**5**), and then an

intramolecular Claisen condensation employing cesium carbonate to furnish the oxindole **6**.

Judicious choice of the aniline nitrogen protecting group proved important for late-stage

functionalization, as we desired to react selectively at this N3 nitrogen in the presence of both the

scaffold's guanidinium (at C2) and methylamino methyl (at C5) substituents. Taking inspiration

from our process synthesis developed for BNM-III-170 (41), a dynamic kinetic resolution was

performed on the oxindole **6** to give the β-hydroxy ester **7** (42). From **7**, it was necessary to install

an amine at the benzylic position of the indoline core. To accomplish this, we first reduced the

ester with $LiBH_4$, followed by protection of the primary alcohol.

Figure 2. Synthesis of indoline core CD4mcs. (**A**) Synthetic route towards a common intermediate for analog synthesis starting from commercially available methyl 2-amino-4-bromobenzoate **4**. This route affords the indoline scaffold-containing intermediate **14** in 15-steps and 1.1% overall yield. (**B**) Functionalization strategy of the common intermediate **14** for analog synthesis.

Stereoinversion was then accomplished via displacement of the secondary alcohol using diphenylphosphorylazide (DPPA) to furnish an azide, that was next reduced with tin (II) chloride to give the desired primary amine **8**. Installation of our previously optimized oxalamide (43) utitlizing 9 on the derived primary amine gave intermediate **10**, which was subjected to tetrabutylammonium fluoride (TBAF) to effect removal of the silyl protecting group, followed by

Mitsunobu union of the resulting primary alcohol with tri-N-bocguanidine to yield intermediate **11**.

With these two crucial structural features installed, we turned towards installation of the

methylamino methyl side-chain. Suzuki coupling (44) of aryl bromide **11** with potassium

vinyltrifluoroborate (**12**), oxidative cleavage, followed by reductive amination of the resulting

aldehyde with methylamine, and protection of the free amine as the *tert*-butyl carbamate provided

intermediate **13**.

Initially, removal of the benzyl carbamate proved difficult, as employing standard hydrogenation

conditions resulted in dechlorination of the *m*-F-*p*-Cl aromatic ring. Instead, transfer

hydrogenation utilizing Pd(OAc)$_2$ and Et$_3$SiH as the hydride source permitted selective removal

of the benzyl carbamate protecting group to give **14**. It should be noted that the indoline scaffold

is very sensitive to acidic conditions. Significant transfer of the *tert*-butyloxycarbonyl from the

adjacent protected guanidine to the indoline nitrogen was observed even when subjected to weak

acids, such as pyridinium hydrochloride. With release of the indoline nitrogen having been

achieved, acylation with triphosgene and 2,6-lutidine as base yielded carbamoyl chloride **15**,

which proved to be stable, and as such could be used directly without purification in conjunction

with various alcohols to produce carbamates such as **16** in excellent yield. Finally, removal of the

protecting groups resulted in the carbamates as TFA salts (**17-30**, *Figure 2B* and *Figure 3*).

Synthetic modifications at the indoline N3 position proceeded with prioritization based on a

combination of chemical feasibility, structural considerations, and feedback from viral inhibition

assays. Molecular modeling of the CD4mc-gp120 complexes was informed by the disposition of

the indane C3 substituent in CJF-III-049-S (**2**), similarity between the indoline and indane

scaffolds, and the topography of the gp120 vestibule surface (*Figure S5 and S6)*. Crystal

structures of gp120 complexes with selected compounds tested and refined hypotheses in this

structure-based development course.

B. *Inhibition of HIV-1 Infection by Indoline CD4mcs*

Initially, several carbamate analogs were synthesized and tested in an assay measuring the

inhibition of single-round HIV-1$_{JR-FL}$ infection. We were curious to know if the simplest carbamate

(**17**, X = NCO$_2$Me) would retain the ability to inhibit Env function. Gratifyingly, this analog displayed

a modest 2.5-fold increase in potency (*Figure 3A*) compared to the previous lead CD4mc, BNM-

III-

Figure 3. Indoline CD4mcs inhibit HIV-1 entry more effectively than the previous lead indane CD4mc BNM-III-170. (**A**)

Inhibition of infectivity of recombinant HIV-1 pseudotypes by CD4mc analogs. The bar graph shows the inhibitory

activity of indoline CD4mcs against infection of recombinant virus pseudotyped with the Env of HIV-1$_{JR-FL}$, normalized

to that of the IC$_{50}$ of BNM-III-170. The table beneath reports the IC$_{50}$ values of indoline CD4mcs against viruses

pseudotyped with the Envs of the AD8, JR-FL and BG505 HIV-1 strains. Only three of the compounds (DY-III-075,

CJF-IV-055 and CJF-IV-056) inhibited the control virus pseudotyped by the amphotropic murine leukemia virus (A-

MLV) Env at concentrations of 300 μM or less. The table also reports the free energy of binding for the indoline CD4mc

analogs predicted by an HIV-1$_{BG505}$ gp120 docking model. (**B**) Comparison of the inhibition of infectivity by BNM-III-170

(**38**) and CJF-III-288 against a panel of recombinant viruses pseudotyped with the Envs of HIV-1 strains from multiple

phylogenetic clades. Compared with BNM-III-170, CJF-III-288 displays an approximately 10-fold increase in potency

for all HIV-1 pseudotypes examined, with the exception of clade A/E recombinants. Neither BNM-III-170 nor CJF-III-

288 inhibited the control virus pseudotyped by the amphotropic murine leukemia virus (A-MLV) Env at concentrations

of 100 μM or less. (**C**) The cell proliferation assay was conducted at varying concentrations of BNM-III-170 and indoline

CD4mcs prepared from stock solutions in 100% DMSO. Compounds were incubated with TZM-bl cells for 72 h, after

which the relative viability of the cells was evaluated with respect to the cells without treatment. The volume of DMSO

used in the DMSO control corresponds to the volume of the CD4mc used at each indicated concentration.

Measurements were performed in triplicate. Error bars indicate mean ± SEM. The levels of TZM-bl cell toxicity for the

indoline CD4mcs are indicated as minimal (green), moderate (orange) or severe (red).

170 (**1**). We also found that indoline **17** bound to gp120 in a similar manner as for indanes **1** and

2. These results suggested that a wide series of indoline CD4mc warranted further investigation.

Carbamates in the alkyl series of ethyl (**18**), propyl (**19**), butyl (**20**), pentyl (**21**), and hexyl (**22**)

carbon chain replacements for the methyl group were readily synthesized and tested for inhibition

of JR-FL virus pseudotypes (*Figure 3A*). As the carbon chain length grew, virus inhibition initially

increased and then decreased, with peak antiviral activity 24-fold higher than for BNM-III-170 for

the three-carbon chain [CJF-III-288 (**19**), *Figure 3A*].

We also explored the consequences of unsaturated bonds in the carbamate system to see if any

effect of addition of an electron pi-cloud could be observed. Allyl carbamate **23** and benzyl

carbamate **26** were therefore constructed and tested. The allyl analog **23** proved to be less potent

than its propyl counterpart **19**, thus providing no indication of significant pi interactions in the

targeted area (*Figure 3A*). However, the benzyl carbamate **26** demonstrated an 11-fold increase

in potency relative to BNM-III-170 (*Figure 3A*), indicating that the proposed targeted area on Env

is sufficient to accommodate an aromatic ring without issue. In general, hydrophobic substituents

on the N3-carbamate appear to interact efficiently with the targeted Env region, leading to more

potent viral inhibition.

To probe further the structural features of the CD4mc that could increase antiviral potency, we

examined the effect of deletion of the one-carbon spacer between the phenyl ring and the

carbamate oxygen in CJF-III-192 (**26**). To this end, the phenyl carbamate derivative **25** was

constructed and tested. Interestingly, the antiviral potency of this CD4mc was decreased

compared with that of the benzyl carbamate, CJF-III-192 (**26**) (*Figure 3A*). The crystal structure

of CJF-III-192 (**26**) complexed to gp120 shows that the one-carbon spacer is essential for

providing the rotatable bond needed for the phenyl ring to adopt a favorable binding conformation.

Introduction of a *para*-bromine on the phenyl ring of **26** to yield **27** decreased the antiviral activity

(*Figure 3A*); this decrease is not readily explained by the crystal structure of **26**, since the para

position projects away from gp120 into the solvent.

Allyl carbamate **23** proved useful for analog synthesis as well. A 1,2-diol functionality was readily

introduced from the alkene via dihydroxylation. The resulting diol **24** is structurally similar to the

indane CD4mc CJF-III-049-S (*Figure 1C*), but the butanediol substituent is placed further out from

the scaffold. A 2.5-fold improvement in viral inhibition by **24** (*Figure 3A*) compared to BNM-III-

170 was observed. These results indicate that **24** is comparable to CJF-III-049-S (**2**) (*Figure 1C*)

in antiviral activity but less active than **23**; therefore, the addition of a diol at this location did not

apparently enhance positive interactions with Env.

Selective fluorine incorporation has proved useful in pharmaceutical development (45); we

therefore explored such derivatization here for CD4mc indoline compounds. In addition, given the

positive effect of 3-substituent hydrophobicity observed in the alkyl and aryl series, we were

intrigued by the potential contribution of fluorine incorporation to this effect. Specifically,

pentafluorobenzyl carbamate **30**, 1,1,1-trifluoroethyl carbamate **28** and 1,1,1-trifluoropropyl

carbamate **29**, analogs of indolines **26**, **18** and **19**, respectively, were synthesized and tested for

the ability to inhibit HIV-1$_{JR-FL}$ infection. The fluorinated analogs **30** and **28** showed 2.1- to 2.7-fold

enhanced antiviral activity compared to their respective fully hydrogenated counterparts **26** and

18 (Figure 3). Pleasingly, the pentafluorobenzyl carbamate in **30** (CJF-IV-046) resulted in a 23-

fold improvement in viral inhibition compared to BNM-III-170. The antiviral activity of **29** was

similar to that of **19**, indicating that the addition of fluorine did not enhance potency against HIV-

1 in this instance. Thus, for some indoline CD4mcs, fluorination can be beneficial to antiviral

potency (*Figure 3A*).

To examine the breadth of the observed antiviral activity, we tested selected indoline CD4mcs

against HIV-1$_{AD8}$, a different clade B virus, and against HIV-1$_{BG505}$, a clade A virus that has proven

to be more difficult to inhibit with current CD4mcs. The tested indoline CD4mcs (**19**, **23**, **25**-**26, 28**

and **30**) all inhibited HIV-1$_{AD8}$ with IC$_{50}$ values 5-10-fold better than those for HIV-1$_{JR-FL}$ (*Figure*

3A). The clade A HIV-1$_{BG505}$ was not susceptible to inhibition by our previous lead CD4mc, BNM-

III-170 (**1**, IC$_{50}$>300 µM) or other tested indane CD4mcs. By contrast, most of the tested *indoline*

CD4mcs inhibited HIV-1$_{BG505}$ infection with IC$_{50}$ values in the micromolar range (*Figure 3A*). Of

these CD4mcs, CJF-IV-046 (**30**), CJF-III-288 (**19**) and DY-III-065 (**29**) stood out as the most

potent CD4mcs, with BG505 IC$_{50}$ values of 10.1 ± 6.8 µM, 27.0 ± 8.8 µM, and 35.0 ± 24.0 µM

respectively. To further investigate the underlying binding mechanism of these CD4mcs to HIV-

1$_{BG505}$, we then used molecular docking and the free energy of binding calculation using molecular

mechanics/generalized born surface area (MM-GBSA) to evaluate binding mode of the CD4mcs

and their correlation with experimental IC$_{50}$ values (*Figure S5 and S6*). These results indicate that

the carbamate substituents at the indoline N3 nitrogen can confer not only improved potency but

also greater breadth against HIV-1 strains.

We next examined the ability of two potent indoline CD4mc analogs, CJF-III-288 (**19**) and CJF-

IV-046 (**30**), to inhibit infection of TZM-bl target cells by a panel of global HIV-1 strains from

phylogenetic clades A, B, C, D and G, as well as circulating recombinant forms AE, AG, BC and

CD. With the exception of the clade AE HIV-1, which have a His 375 residue that fills the Phe 43

cavity, all of the viruses were inhibited by CJF-III-288 (**19**) and CJF-IV-046 (**30**); 92% and 98% of

these viruses were inhibited by CJF-III-288 and CJF-IV-046, respectively, with IC_{50} values less

than 15 µM (*Figure 3B, Table S1*). By contrast, 25% of the viruses were either not inhibited or

required more than 15 µM to be inhibited by BNM-III-170 (**1**). The IC_{50} values of BNM-III-170 (**1**),

CJF-III-288 (**19**) and CJF-IV-046 (**30**) for this panel of viruses were strongly correlated, with the

slope of the regression line indicating an ~10-fold increase in potency for the two indoline CD4mcs

over BNM-III-170 (*Figure 3B, Table S3, Figure S4*). These results indicate that the properties of

the HIV-1 Env that determine sensitivity/resistance apply to both indane and indoline CD4mcs. A

weaker correlation was observed between the IC_{50} values ofCJF-III-288 (**19**) and sCD4-Ig *(Figure*

S4).

C. *Cytotoxicity of the indoline CD4mcs*

With the exception of DY-III-075 (**20**), CJF-IV-055 (**21**) and CJF-IV-056 (**22**), none of the indoline

CD4mc analogs tested exhibited non-specific or toxic activity at concentrations up to 300 µM in

the antiviral assays using Cf2Th-CD4/CCR5 target cells. However, at concentrations higher than

20 µM, CJF-IV-046 (**30**) exerted non-specific or toxic effects on the TZM-bl target cells used to

evaluate the inhibition of infection by the panel of global HIV-1 strains (*Table S1*). To investigate

potential effects on cell proliferation and viability, the potent indoline CD4mcs **19**, **20**, **28**, **26**, **29**,

30 and BNM-III-170 were incubated with TZM-bl cells at varying concentrations (*Figure 3C*). We

observed that CD4mcs bearing aromatic carbamate functionalities (**26**, **30**) inhibited TZM-bl cell

proliferation beginning at a concentration of 50 μM. Most of the CD4mcs with alkyl carbamate

substituents (**19**, **28**, **29**) did not affect cell viability; however, at high (100 μM) concentrations of

20, toxicity for TZM-bl cells was observed. These results indicate that non-specific toxicity for

TZM-bl cells is associated with indoline analogs with aromatic carbamates, a consideration for

future indoline CD4mc design. Fortunately, most alkyl carbamate substituents appear to be non-

toxic replacements for the aromatic carbamates, giving rise to CD4mcs that are equally effective

in terms of antiviral potency. Moreover, the inclusion of fluorination in the ethyl or propyl

carbamates (**28**, **29**) did not adversely affect cell viability. It is worth noting that DMSO also

decreases the TZM-bl cell viability at higher concentrations.

D. Crystal Structures of Indoline CD4mcs

In parallel with indoline CD4mc synthesis and evaluation of antiviral activity, we also analyzed

crystal structures of selected compounds in complexes with the HIV-1$_{C1086}$ gp120 core$_e$ protein,

which preferentially assumes a CD4-bound conformation (46). Pre-formed crystals were soaked

with individual indoline CD4mcs [CJF-III-214 (**17**), CJF-III-289 (**18**), CJF-III-288 (**19**), CJF-III-192

(**26**), CJF-IV-047 (**28**), DY-III-065 (**29**), or CJF-IV-046 (**30**)], and the atomic structures were refined

against X-ray diffraction measurements for the compounds at resolutions from 2.5 to 1.9 Å (Table

S1). For comparisons, structures were also re-determined in this P2$_1$2$_1$2$_1$ lattice at improved

resolution for the previously reported complexes with the indane CD4mcs BNM-III-170 (38) and

CJF-III-049-S (40). Electron density distributions were well defined, as shown in Figure 4A for **29**

and for each ligand in Figures S1 A-I. The densities for each of the four copies in the asymmetric

unit are highly similar (e.g., Figure S2 A-D for **29**); accordingly, we constrained the fitted gp120-

CD4mc atomic structures to be identical. The gp120 binding modes of all seven indoline CD4mcs

are the same (Figure S3A).

Figure 4. Crystal structures of CD4mcs in complex with HIV-1$_{C1086}$ gp120 core$_e$. (**A**) Crystal structure of indoline CD4mc DY-III-065 (**29**) as complexed with gp120 displayed in stick representation. Hydrogen bonds are shown as black dashed lines. The carbon atoms of DY-III-065, carbon atoms of gp120, and the oxygen, nitrogen, chlorine and fluorine atoms are shown in magenta, beige, red, blue, dark green and light green, respectively. The C$_\alpha$ atoms of terminal residues in gp120 segments are represented in balls. The density is from an 2Fo-Fc synthesis contoured at about 1.5σ and selected for coverage within 2 Å of ligand atoms. (**B**) Comparison between indoline CD4mc DY-III-065 (**29**) and indane CD4mc CJF-III-049-S (**1**) after superimposition of the gp120 cores. The carbon atoms of DY-III-065 and CJF-III-049-S are in magenta and cyan, respectively. The insert image compares DY-III-065 and CJF-III-049-S after superimposition of their aromatic six-membered rings. (**C**) π-π interactions between the aromatic six-membered ring of DY-III-065 (**29**) and the G473-D474 peptide unit of gp120. The black dashed line connects the centroid of the aromatic six-membered ring of DY-III-065 and center of the G473-D474 peptide bond. (**D-F**) Hydrogen bonds and non-bonded contacts that DY-III-065 (**D**), CJF-III-049-S (**E**) or CJF-IV-046 (**F**) form with gp120 and surrounding water molecules. Hydrogen bonds and non-bonded contacts are shown as black and green dashed lines, respectively. (**G**) Fluorine atoms of DY-III-065 (**29**) form favored non-bonded contacts with the C$_\alpha$ of E429 in gp120 and with its own carbamate carbon. (**H**) A water network at the protein-ligand interface in the structure of the DY-III-065 (**29**) complex associates the strictly conserved gp120 D368 carboxylate with the guanidinium group from C2 of the CD4mc.

The substituents at indane positions 1, 2 and 5 of BNM-III-170 (**1**) are preserved identically in CJF-III-049-S (**2**) and in all of the indoline CD4mcs of this study. The methylamino methyl group

at C5 makes no gp120 contacts, but the other two are stereo-specifically fixed by hydrogen

bonding to the gp120 backbone. Thus, despite varied substitutions at N3 or C3, the binding poses

of the indoline CD4mcs are similar to those of the indane CD4mcs (Figures 4B and S3B);

however, because of geometric differences between the indoline and indane CD4mcs in the

dispositions of the fixed C1 and C2 substituents on the five-membered rings, their very similar

ring skeletons are oriented differently when bound to gp120 (Figure 4B). As a consequence, the

aromatic six-membered ring is stacked against the G473-D474 peptide unit (Figure 4C) more

closely for the indoline CD4mcs, which should provide greater π-π overlap and strengthened

binding. Moreover, this redisposition in the indoline CD4mcs (Figure 4D) relative to the indane

CD4mcs (Figure 4E) also moves several non-bonded contacts into highly favorable van der Waals

interactions.

The N3 carbamate ester substituents of the most efficacious indoline CD4mcs studied here (**19**,

29 and **30**) all occupy spaces in the Phe 43 vestibule of gp120 that are similar to, but distinctive

from that for the C3 butanediol substituent of indane CD4mc **2** (Figures 4E and S3B). The

carbamate ester cores of these indolines are chemically and structurally the same through the

first carbon of the carbamate substituent, yet the antiviral activities of these compounds increase

from the methyl to propyl, benzyl, trifluoropropyl and pentafluorobenzyl by factors of 4.3, 2.0, ~8

and 4.2, respectively. The paucity of gp120 contacts from the carbamate substituents confounds

the analysis, especially as compared with those from the C3 substituent in indane **2** (Table S2),

which shows many more contacts despite having activities >17-fold lower than the best indoline

CD4mcs (**19**, **29** and **30**). The benzyl carbamate (**26**) and its pentafluorobenzyl counterpart (**30**)

adopt identical conformations as bound to gp120 (Figure S3C); and the propyl (**19**) and

trifluoropropyl (**29**) carbamate are also very similar (Figure S3D). Thus, the fully hydrogenated

and partially fluorinated pairs make very similar gp120 contacts, altogether hydrophobic for the

propyl carbamate; nevertheless, for **30** (Figures 4F) and **29** (Figure 4G), the fluorine contacts

appear to potentiate the interactions with the gp120 core$_e$ protein, in keeping with their enhanced

antiviral potencies.

The structure of the gp120 core$_e$ complex with DY-III-065 (**29**) reveals water structure at protein-

ligand interfaces. One water site, shown in Figure 4D, is hydrogen-bonded to the carbonyl oxygen

of G473, the amino group of M475, and the carbonyl oxygen of W427 (not shown) and just outside

hydrogen-bonding distance from the carbamate-linking oxygen atom. A second set of waters

associates the C2 guanidinium group of **29** with the carboxyl group of the stringently conserved

D368 of gp120, which is critical for CD4 binding (23). The C2 guanidinium nitrogens of **29** H-

bond to two water molecules, one of which H-bonds to D368; water-mediated H-bonding also

bridges D368 to a C1 oxalamide oxygen of **29** (Figure 4H). These water sites are also present in

the indane CD4mc complexes, but they only become clearly evident here with the highest

resolution yet achieved for this system. Synthetic efforts to create new indoline CD4mc analogs

that mimic these water-based interactions with D368 are in progress.

It is noteworthy that interactions between DY-III-065 (**29**) and gp120 primarily involve main chain

atoms or the side chain of the conserved CD4-binding residue D368 (23, 24)). Water molecules

that mediate the contacts between **29** and the side chains of polymorphic HIV-1 gp120 residues

(e.g., His/Gln 105) can potentially form hydrogen bonds despite this variation. These

characteristics should militate against the evolution of HIV-1 escape mutants.

E. Antiviral Activity of Indoline CD4mcs Against Resistant HIV-1 Variants

Previously, CD4mc-resistant variants of HIV-1$_{AD8}$ were selected by incubating infected cultures

with increasing concentrations of BNM-III-170 (39). After selection, the derived mutant HIV-1$_{AD8}$

130-C was completely resistant to entry inhibition by concentrations of BNM-III-170 up to 300 μM.

In this case, two changes in the Phe 43 cavity (I424T and S375N) and a change (E64G) in Layer

of the gp120 inner domain individually contributed to BNM-III-170 resistance, and acted additively to increase resistance to BNM-III-170 (39).

Given the improved breadth of infection inhibition exhibited by the indoline CD4mcs, we were curious if this class of CD4mcs could overcome these resistance-associated adaptations and inhibit the infection of these HIV-1 mutants. The evolved BNM-III-170-resistant HIV-1$_{AD8}$ mutant 130-C and the HIV-1$_{AD8}$ mutant containing the three resistance-associated gp120 changes (E64G+S375N+I424T) were inhibited by high concentrations of CJF-III-192 (**26**) (*Table 1*); the CD4mcs CJF-III-288 (**19**) and CJF-IV-046 (**30**) were less effective against these mutants (*Table 1*). The indoline CD4mcs were also evaluated for their ability to inhibit the HIV-1$_{AD8}$ mutants containing single or pairwise combinations of the resistance-associated gp120 changes. All of the tested indoline CD4mcs (**19, 26** and **30**) inhibited the single-residue HIV-1$_{AD8}$ mutants with IC$_{50}$ values less than 10 µM. The inhibition of infection was diminished for the pairwise combinations of the gp120 changes; however, variants containing all three permutations of the pairwise mutations were still inhibited by the indoline CD4mcs **19**, **26** and **30** with IC$_{50}$ values less than 75 µM. In all cases, the indoline CD4mcs were able to inhibit these resistance-selected HIV-1$_{AD8}$ mutants more effectively than BNM-III-170 (39). Thus, although the gp120 changes that determine resistance to our current lead indane CD4mc, BNM-III-170, also result in relative resistance to the indoline CD4mcs, the increased potency of the indoline CD4mcs allows measurable inhibition of HIV-1 variants that are resistant to BNM-III-170.

Table 1. Sensitivity of BNM-III-170-resistant HIV-1 variants to inhibition by indoline CD4mcs

HIV-1(AD8) Env variant	Fold resistance to:				IC$_{50}$ (µM)		
	BNM-III-170	CJF-III-288	CJF-IV-046	CJF-III-192	CJF-III-288	CJF-IV-046	CJF-III-192
WT	1	1	1	1	0.20 ± 0.03	0.20 ± 0.01	0.20 ± 0.01
E64G	109	25	20	32.5	5.0 ± 1.2	4.0 ± 2.0	6.5 ± 1.01
S375N	5.2	7.5	8.5	9	1.5 ± 0.17	1.7 ± 0.02	1.8 ± 0.91
I424T	5.2	11	8.5	9	2.2 ± 0.08	1.7 ± 0.02	1.8 ± 0.84
E64G+S375N	>1500	300	200	200	60.0 ± 7.2	40.0 ± 10.0	40.0 ± 5.0
E64G+I424T	>1500	375	200	200	75.0 ± 6.1	40.0 ± 12.2	40.0 ± 6.3
S375N+I424T	209	51	41	45	10.1 ± 2.2	8.2 ± 2.08	9.0 ± 3.0
E64G+S375N+I424T	>1500	1450	700	650	290.2 ± 10	140.0 ± 15.0	130 ± 35.0
130-C	>1500	1305	700	400	260.9 ± 30	140.0 ± 20.7	80 ± 20.0
A-MLV	>1500	>1500	>1500	>1500	>300	>300	>300

The sensitivity of recombinant luciferase-expressing viruses pseudotyped with the indicated HIV-1$_{AD8}$ Env variants to the indoline CD4mcs was determined using Cf2Th-CD4/CCR5 target cells, as described in Materials and Methods. The IC$_{50}$ values (in µM) are shown in the right three columns. The level of resistance relative to that of the wild-type (WT) AD8 Env is shown in the left columns; the values for the indane CD4mc, BNM-III-170, were derived from reference 39.

F. ADCC and Antibody Recognition

CD4mcs have the potential to sensitize HIV-1-infected cells to killing by ADCC (30). We therefore examined the ability of the indoline CD4mc inhibitors: (1) to enhance recognition of Env on the surface of infected cells by antibodies (Ab) present in plasma from HIV-1-infected individuals (HIV+ plasma); (2) to mediate recognition of antibody-bound infected cells by FcγRIIIa receptors

in the presence of HIV+ plasma; and (3) to sensitize infected cells to ADCC mediated by HIV+

plasma. Potent indoline CD4mcs were evaluated first for their ability to enhance HIV+ plasma

binding to primary CD4+ T cells infected by HIV-1$_{JR-FL}$. As demonstrated in *Figure 5A and D,* the

indoline CD4mcs CJF-III-288 (**19**), CJF-III-192 (**26**), and CJF-IV-046 (**30**) enhanced the binding

of plasma from HIV-1-infected individuals to HIV-1$_{JR-FL}$-infected cells more efficiently than BNM-

III-170. We also evaluated whether the CD4mcs were able to enhance HIV+ plasma binding to

HIV-1$_{JR-FL}$-infected cells and promote subsequent FcγRIIIa receptor engagement (*Figure 5B and*

E). All of the indoline CD4mcs examined [CJF-III-288 (**19**), CJF-III-192 (**26**), and CJF-IV-046 (**30**)]

improved recombinant soluble dimeric FcγRIIIa engagement by HIV+ plasma to a greater degree

than BNM-III-170. In addition, the indoline CD4mcs CJF-III-288 (**19**), CJF-III-192 (**26**) and CJF-

IV-046 (**30**) all enhanced ADCC killing of HIV-1$_{JR-FL}$-infected CD4+ T cells to a greater extent than

BNM-III-170 (*Figure 5C and F*).

In addition to enabling the recognition of HIV-1$_{JR-FL}$-infected CD4+ T-cells by HIV+ plasma,

indoline CD4mc CJF-III-288 (**19**) enhanced HIV+ plasma binding to primary CD4+ T cells infected

with a panel of HIV-1 strains ~2-fold more effectively than BNM-III-170 (*Figure 5G*).

Figure 5. Indoline CD4mcs enhance recognition of HIV-1-infected cells by HIV+ plasma, enabling ADCC. (**A-F**) Primary CD4+ T cells were infected with HIV-1$_{JR-FL}$ and used as targets for HIV+ plasma and FcγRIIIa binding and for their susceptibility to HIV+ plasma mediated-ADCC. (**A**) HIV+ plasma binding (MFI) to CD4+ T cells infected with HIV-1$_{JR-FL}$ in the presence of DMSO or the indicated CD4mc. (**B**) FcγRIIIa binding (MFI) in the presence of HIV+ plasma and DMSO or a CD4mc. (**C**) HIV+ plasma-mediated ADCC (%) in the presence of DMSO or the indicated CD4mc. (**D**) HIV+ plasma binding (MFI) in the presence of DMSO or a CD4mc. (**E**) FcγRIIIa binding (MFI) in the presence of HIV+ plasma and DMSO or a CD4mc. (**F**) HIV+ plasma-mediated ADCC (%) in the presence of DMSO or a CD4mc. (**G**) Comparison of BNM-III-170 and CJF-III-288 stimulation of HIV+ plasma binding to primary CD4+ T cells infected with the indicated HIV-1 strain. Statistical significance was tested using (**A-F**) paired *t* test or Wilcoxon test based on statistical normality (**P<0.01; ***P<0.001 ****P<0.0001) and (**G**) Spearman rank correlation test. Concentrations of CD4mc used in each assay were: BNM-III-170 (50 μM), CJF-III-192 (50 μM), CJF-IV-046 (25 μM), or CJF-III-288 (50 μM).

Discussion

CD4mcs are unique small-organic-molecule HIV-1 entry inhibitors that competitively block CD4 binding, trigger premature conformational changes in the Env leading to functional inactivation, and "open" Env, sensitizing virions to antibody neutralization and infected cells to ADCC. Here, we report that certain indoline CD4mc analogs exhibit 10-20-fold increases in specific anti-HIV-1 potency, compared with the earlier lead indane CD4mc, BNM-III-170. Accompanying this increase in potency, we observed greater breadth of activity against a range of primary HIV-1 strains. Natural HIV-1 variants exhibit approximately 1000-fold differences in sensitivity to indane CD4mcs, with a substantial fraction of variants requiring greater than 100 µM concentrations of BNM-III-170 to achieve 50% inhibition of infection. Remarkably, regardless of their phylogenetic classification into clade A, B, C, D, AG, BC or CD, diverse HIV-1 strains exhibited increased sensitivity to the more potent indoline CD4mc analogs. Only clade AE recombinant HIV-1, which has the imidazole side chain of His 375 occupying the Phe 43 cavity, were resistant to potent indoline CD4mcs such as CJF-III-288.

Likewise, an HIV-1$_{AD8}$ variant (130-C) selected for resistance to BNM-III-170, as well as recombinant HIV-1 containing key resistance-associated changes in Env, were found to be inhibited more efficiently by the indoline CD4mcs than by BNM-III-170. Apparently, Env changes near or within the Phe 43 cavity (such as S375N and I424T in the 130-C virus) that are unfavorable for CD4mc binding can be compensated by contacts made by the indoline CD4mcs with the gp120 vestibule. These novel contacts of the indoline CD4mcs also appear to compensate for the more distant E64G change, which may decrease the propensity of Env to stabilize the CD4-bound conformation favored by the CD4mc (39). Thus, although the indoline CD4mcs are still affected by the resistance-associated gp120 changes, the overall increase in the relative antiviral potency of the indoline CD4mcs allows inhibition of the resistant viruses at achievable concentrations.

Although further study is needed, the increased antiviral potency of the indoline CD4mcs,

compared with BNM-III-170, likely derives from novel contacts with the gp120 surface made by

the substituents on the N3 nitrogen. Crystal structures suggest that these contacts involve a

conserved hydrophobic pocket at the junction of the gp120 inner and outer domains (*Figure 1B*).

This hydrophobic pocket is flanked by the polymorphic residues His/Gln 105 in the α1 helix and

Arg/Lys 476 in the α5 helix. These natural polymorphisms apparently exert only minimal effects

on HIV-1 sensitivity to the more potent indoline CD4mcs, because several susceptible natural

variants exhibit polymorphic changes in these residues. These observations suggest that indoline

CD4mc potency may benefit from interactions of the N3 substituent that are tolerant of some

natural HIV-1 diversity. A consequence of the additional gp120 contacts made by the N3

substituent may thus be better stabilization of the binding pose shared by the indoline and indane

cores. This pose promotes hydrogen bonding with multiple mainchain atoms (Asn 425, Met 426,

Trp 427, Gly 473) that contribute to the ability of CD4mcs to recognize gp120 glycoproteins from

multiple strains.

Strong correlations were also observed between the sensitivities of HIV-1 variants to the indane

CD4mc BNM-III-170 and to potent indoline CD4mcs, CJF-III-288 and CJF-IV-046. As the

pseudotyped recombinant viruses used in the assays differ only in Env, the observed correlations

indicate that Env-specific variables determine sensitivity to CD4mcs in general. The nature of the

Env variables accounting for the wide range of sensitivities to CD4mcs is clearly worthy of further

investigation. We note that polymorphisms in residues 105 and 476, which can be contacted by

some indoline analogs, but not by BNM-III-170, do not result in deviations from the linear

correlation in *Figure 3B*, supporting the assertion that these natural changes exert minimal impact

on the efficacy of the more potent indoline CD4mcs.

Finally, three of the indoline CD4mcs, CJF-III-192, CJF-III-288 and CJF-IV-046, were more

effective than BNM-III-170 in exposing vulnerable Env epitopes on the surface of HIV-1-infected

cells relevant to ADCC. These results justify ongoing studies in animal models to evaluate the

ability of indoline CD4mc to sensitize HIV-1-infected cells to ADCC.

Indoline CD4mc with improved potency should expedite exploration of the therapeutic and

prophylactic potential of these interesting small molecules.

Materials and Methods

Ethics statement. Written informed consent was obtained from all study participants [the

Montreal Primary HIV Infection Cohort (47, 48) and the Canadian Cohort of HIV Infected Slow

Progressors (49-51) and research adhered to the ethical guidelines of CRCHUM and was

reviewed and approved by the CRCHUM Institutional Review Board (Ethics Committee, approval

number CE 16.164 - CA). Research adhered to the standards indicated by the Declaration of

Helsinki. All participants were adult and provided informed written consent prior to enrollment in

accordance with Institutional Review Board approval.

Cell lines and primary cells. 293T cells and TZM-bl cells were grown in Dulbecco's modified

Eagle's medium (DMEM) (Life Technologies, Wisent Inc.) supplemented with 10% fetal bovine

serum (FBS) (Life Technologies, VWR) and 100 μg/ml of penicillin-streptomycin (Life

Technologies, Wisent Inc.). Cf2Th cells stably expressing the human CD4 and CCR5 coreceptors

for HIV-1 were grown in the same medium supplemented with 0.4 mg/ml of G418 and 0.2 mg/ml

of hygromycin.

Human peripheral blood mononuclear cells (PBMCs) from healthy HIV-1-negative individuals

were obtained by leukapheresis and Ficoll-Paque density gradient isolation and were

cryopreserved in liquid nitrogen until use. CD4+ T lymphocytes were purified from resting PBMCs

by negative selection using immunomagnetic beads according to the manufacturer's instructions

(StemCell Technologies). The cells were activated with phytohemagglutinin-L (10 µg/mL) for 48

h and then maintained in RPMI 1640 complete medium supplemented with recombinant

interleukin-2 (rIL-2) (100 U/mL).

Viral production and infection

Vesicular stomatitis virus G (VSVG)-pseudotyped viruses were produced and titrated as

previously described (52). Viruses were used to infect activated primary CD4+ T cells from healthy

HIV-1-negative donors by spin infection at 800 x g for 1 h in 96-well plates at 25°C.

CD4-mimetic compounds (CD4mcs). The CD4mc analogs were synthesized as described in

detail in the Supporting Information. The compounds were dissolved in DMSO at a stock

concentration of 10 mM and stored at -20°C until use. The CD4mcs were diluted to 1 mM in

DMEM for pseudovirus inhibition assays, 50 or 25 µM in PBS for cell-surface staining and 50 or

25 µM in RPMI 1640 complete medium for ADCC assays.

Antibodies and plasma. The sCD4-Ig used in the virus inhibition assays is described in

reference 39. The anti-coreceptor binding site 17b monoclonal antibody (mAb) (NIH HIV Reagent

Program) and plasma from HIV-1-infected individuals were used to assess cell-surface Env

conformation. Plasma from the Montreal Primary HIV Infection Cohort (47, 48) and the Canadian

Cohort of HIV Infected Slow Progressors (49-51) were collected, heat-inactivated and conserved

at −80 °C until use. The conformation-independent anti-gp120 outer-domain 2G12 mAb (NIH HIV

Reagent Program) was used to normalize Env expression. Goat anti-human IgG (H+L) (Thermo

Fisher Scientific) antibodies pre-coupled to Alexa Fluor 647 and streptavidin conjugated to Alexa

Fluor 647 were used as a secondary antibody in flow cytometry experiments.

Inhibition of HIV-1 infection by CD4mcs. HIV-1 inhibition assays were performed as described

previously (29,39,43). To produce recombinant luciferase-expressing HIV-1, 293T cells were

transfected with plasmids expressing HIV-1 Env variants and Rev, the pCMVΔP1Δenv HIV-1

Gag-Pol packaging construct and the firefly luciferase-expressing HIV-1 vector at a 1:1:3 µg DNA

ratio using Effectene transfection reagent (Qiagen). Recombinant luciferase-expressing viruses

capable of a single round of replication were released into the cell medium and were harvested

48 h later. The virus-containing supernatants were clarified by low-speed centrifugation (600 x g

for 10 min) and used for single-round infections. To measure inhibition of infection, the

recombinant viruses were incubated with the CD4mcs for 1 h at 37°C. The mixture was then

added to Cf2Th-CD4/CCR5 target cells expressing CD4 and CCR5. After 48 h of culture, the

target cells were lysed and the luciferase activity was measured.

The pseudotyped virus inhibition assays used to generate the data in Figure 3B and Table S1

were conducted as previously described (38). Single-round HIV-1 pseudotyped by the Envs from

a panel of multiclade HIV-1 were incubated for 1 h at 37°C with 5-fold serial dilutions of CD4mc

or sCD4-Ig in duplicate. TZM-bl cells were then added in growth medium containing DEAE-

dextran at a final concentration of 11 µg/ml. The assay plates were incubated for 48 h at 37°C,

5% CO_2, after which the cells were lysed and luciferase reporter gene expression was measured

using Bright-Glo luciferase reagent (Promega) and a Victor 3 or GloMax Navigator luminometer

(Perkin Elmer/Promega). The 50% and 80% inhibitory concentrations (IC_{50} and IC_{80}, respectively)

were calculated based on the relative luciferase unit (RLU) activity in the treated sample

compared with that in the untreated control, after subtraction of the background RLU in control

uninfected TZM-bl cells.

Cell viability assay. TZM-bl cells were seeded in 96-well assay plates containing 100 µl of culture medium. CD4mcs in stock solutions of 100% DMSO were added at the specified final concentrations in triplicates. For the DMSO control, a volume of DMSO was added that corresponded to the volume of the CD4mc added at each indicated concentration. After seventy-two h, cell viability was measured using the CellTiter 96® Non-Radioactive Cell Proliferation Assay (MTT) kit from Promega. Briefly, 15 µl of dye solution (from the kit) was added to each well followed by incubation of the plate at 37°C for 4 h in a humidified CO_2 incubator. One hundred µl of solubilization/stop solution (from the kit) was then added to each well. Absorbance at 570 nm and 630 nm was measured using a 96-well plate reader.

Flow cytometry analysis of cell-surface staining. Cell surface staining of primary infected cells was performed 48 h post-infection as previously described (30). HIV-1-infected cells were incubated for 1 h at 37 °C with HIV+ plasma (1:1000 dilution) in the presence of CD4mcs (25 or 50 µM) or with equivalent volume of vehicle (DMSO). Cells were then washed once with PBS and stained with appropriate Alexa Fluor-647-conjugated (Invitrogen) secondary Abs (2 µg/mL) for 20 minutes at room temperature. Alternatively, the binding of HIV+ plasma was detected using a biotin-tagged dimeric recombinant soluble FcγRIIIa (0.2 µg/mL) followed by the addition of Alexa Fluor 647-conjugated streptavidin (Thermo Fisher Scientific; 2 µg/mL). Infected cells were then stained intracellularly for HIV-1 p24, using the Cytofix/Cytoperm Fixation/ Permeabilization Kit (BD Biosciences, Mississauga, ON, Canada) and the fluorescent anti-p24 mAb (PE or FITC-conjugated anti-p24, clone KC57; Beckman Coulter/Immunotech). The percentage of infected cells (p24+ cells) was determined by gating the living cell population on the basis of AquaVivid viability dye staining. Samples were analyzed on an LSRII cytometer (BD Biosciences), and data analysis was performed using FlowJo vX.0.7 (Tree Star, Ashland, OR, USA).

ADCC FACS-based assay. Measurement of ADCC using the FACS-based assay was performed

at 48 h post-infection as previously described (30, 34). Briefly, HIV-1$_{JR-FL}$-**i**nfected primary CD4+

T cells were stained with viability (AquaVivid; Thermo Fisher Scientific) and cellular (cell

proliferation dye eFluor670; eBioscience) markers and used as target cells. Autologous PBMC

effectors cells, stained with another cellular marker (cell proliferation dye eFluor450; eBioscience),

were added at an effector: target ratio of 10:1 in 96-well V-bottom plates (Corning, Corning, NY).

Briefly, infected primary CD4+ T cells were incubated with HIV+ plasma (1:1000), in the presence

of CD4mcs (25 or 50 µM) or an equivalent volume of vehicle (DMSO). The plates were

subsequently centrifuged for 1 min at 300 g, and incubated at 37°C, 5% CO_2 for 4 to 6 h before

being fixed in a 2% PBS-formaldehyde solution. Cells were then stained intracellularly for HIV-1

p24 as described above. Samples were analyzed on an LSRII cytometer (BD Biosciences). Data

analysis was performed using FlowJo vX.0.7 (Tree Star). The percentage of ADCC was

calculated with the following formula: (% of p24+ cells in Targets plus Effectors) − (% of p24+

cells in Targets plus Effectors plus plasma) / (% of p24+ cells in Targets) by gating on infected

live target cells.

Statistical analysis. Statistics were analyzed using GraphPad Prism version 9.4.1 (GraphPad,

San Diego, CA, USA). Every data set was tested for statistical normality and this information was

used to apply the appropriate (parametric or nonparametric) statistical test. P values <0.05 were

considered significant; significance values are indicated as * P<0.05, ** P<0.01, *** P<0.001, ****

P<0.0001.

Protein Purification and X-ray Crystallography. The plasmid expressing the gp120 core$_e$

protein of the clade C HIV-1$_{1086}$ was donated by Lei Chen from Peter Kwong's laboratory (National

Institutes of Health) (46). An 8-histidine (8His) tag was added to the C-terminus of the gp120 core$_e$

protein to assist affinity purification. About 1 mg of the expressor plasmid was transfected into 1

liter of HEK293S GnTI- suspension cells at 2 x 10^6 cells/mL. After expression of the core$_e$ protein,

the supernatant was filtered using a 0.45 µm filter and passed through a pre-packed Ni-NTA

column. The Ni-NTA column was washed with 40 mM imidazole in 1x phosphate-buffered saline

(PBS) and the 8His-tagged gp120 core$_e$ protein was eluted with 250 mM imidazole in 1x PBS.

The eluted gp120 core$_e$ protein was deglycosylated overnight in a 37 °C water bath with

Endoglycosidase Hf (New England Biolabs). Afterwards, the deglycosylated gp120 was further

purified using Concanavalin A (Con A)-Sepharose (Cytiva) to remove any incompletely

deglycosylated proteins. The flowthrough of the Con A-Sepharose column was collected and

further purified using a pre-packed Ni-NTA column, which removed Endoglycosidase Hf in the

flowthrough. The completely deglycosylated gp120 core$_e$ protein was eluted with 250 mM

imidazole in 1x PBS and further polished with a Superdex 200 Increase column (Cytiva). The

purified deglycosylated gp120 core$_e$ protein was concentrated to 10 mg/mL for crystallization.

Crystallization of the unliganded gp120 core$_e$ protein was performed at 20 °C using the hanging

drop vapor diffusion method. The crystals grew in drops consisting of 1 µL of protein solution and

1 µL of reservoir solution to equilibrate against 600 µL of reservoir solution 14 to 16 %(w/v) PEG

1500, 0.1 M CaCl$_2$, 0.1 M imidazole pH 6.5. For each experiment, the compound of interest was

dissolved in 100% DMSO. A single crystal was picked from the mother liquor and soaked in 5 µL

of a stabilization buffer that contained 26% PEG 1500 (w/v), 0.1 M CaCl$_2$, 0.1 M imidazole pH 6.5,

2.5 mM Tris-HCl pH 7.5, 350 mM NaCl, 0.02% NaN$_3$, 5% (v/v) DMSO and 2 mM of the compound.

The gp120 core$_e$ crystals were soaked for 30 to 60 min in the stabilization buffer and then

transferred for 5 seconds in cryo-protectant, which is the stabilization buffer but with 30% ethylene

glycol. Diffraction data were collected on 24ID-C and 24ID-E beamline at Advanced Photon

Source (APS) at Argonne National Laboratory. Crystal structures were solved by the molecular

replacement module in PHENIX using the unliganded HIV-1C$_{1086}$ gp120 core$_e$ structure (PDB ID:

3TGR) and refined with phenix.refine.

Data Availability

Data deposition: Atomic coordinates and structure factors have been deposited in the Research Collaboratory for Structural Bioinformatics (RCSB) Protein Data Bank, www.rcsb.org under accession code 8FLY for BNM-III-170 (53), 8FLZ for CJF-III-049-S (54), 8FM0 for CJF-III-214 (55), 8FM2 for CJF-III-289 (56), 8FM3 for CJF-III-288 (57), 8FM4 for CJF-IV-047 (58), 8FM5 for DY-III-065 (59), 8FM7 for CJF-III-192 (60) and 8FM8 for CJF-IV-046 (61).

Acknowledgments

We thank Irwin Chaiken and all the members of the P01 Consortium Structure-Based Antagonism of HIV-1 Envelope Function in Cell Entry (NIH Grant No. AI150471). We also thank the staff at NE-CAT beamlines at APS for their support during data collection. Dr. Charles Ross, and Dr. Jun Gu (University of Pennsylvania) are also acknowledged for their assistance obtaining mass and NMR spectra, respectively. The authors also thank Dr. Mark Hogarth for kindly providing recombinant dimeric FcgRIIIa and Dr Beatrice Hahn for providing the following infectious molecular clones: HIV_{CH77TF}, $HIV_{CH185TF}$, $HIV_{CH850TF}$ and HIV_{CH167}. A.F. is the recipient of a Canada Research Chair on Retroviral Entry #RCHS0235 950-232424.

References

1.   Nations U (2021) UNAIDS Fact Sheet 2021.

2.   Eholié SP, Aoussi FE, Ouattara IS, Bissagnéné E, & Anglaret X (2012) HIV treatment and care in resource-constrained environments: challenges for the next decade. *Journal of the International AIDS Society* 15(2):17334.

3.   Deeks SG, Overbaugh J, Phillips A, & Buchbinder S (2015) HIV infection. *Nature Reviews Disease Primers* 1(1):15035.

4.   Wyatt R & Sodroski J (1998) The HIV-1 envelope glycoproteins: fusogens, antigens, and immunogens. *Science* 280(5371):1884-1888.

5.   Robey WG*, et al.* (1985) Characterization of Envelope and Core Structural Gene Products of HTLV-III with Sera from AIDS Patients. *Science* 228(4699):593-595.

708   6.   Allan JS, *et al.* (1985) Major glycoprotein antigens that induce antibodies in AIDS

709   patients are encoded by HTLV-III. *Science* 228(4703):1091-1094.

710   7.   Haynes BF, *et al.* (2016) HIV-Host Interactions: Implications for Vaccine Design. *Cell*

711   *host & microbe* 19(3):292-303.

712   8.   Fauci AS (2016) An HIV Vaccine: Mapping Uncharted Territory. *JAMA* 316(2):143-144.

713   9.   Hoxie JA (2010) Toward an antibody-based HIV-1 vaccine. *Annual review of medicine*

714   61:135-152.

715   10.   Karlsson Hedestam GB, *et al.* (2008) The challenges of eliciting neutralizing antibodies

716   to HIV-1 and to influenza virus. *Nature reviews. Microbiology* 6(2):143-155.

717   11.   Richard J, Prévost J, Alsahafi N, Ding S, & Finzi A (2018) Impact of HIV-1 Envelope

718   Conformation on ADCC Responses. *Trends in microbiology* 26(4):253-265.

719   12.   Munro JB, *et al.* (2014) Conformational dynamics of single HIV-1 envelope trimers on

720   the surface of native virions. *Science* 346(6210):759.

721   13.   Herschhorn A, *et al.* (2016) Release of gp120 Restraints Leads to an Entry-Competent

722   Intermediate State of the HIV-1 Envelope Glycoproteins. *mBio* 7(5).

723   14.   Ma X, *et al.* (2018) HIV-1 Env trimer opens through an asymmetric intermediate in which

724   individual protomers adopt distinct conformations. *eLife* 7.

725   15.   Haim H, *et al.* (2011) Contribution of intrinsic reactivity of the HIV-1 envelope

726   glycoproteins to CD4-independent infection and global inhibitor sensitivity. *PLoS*

727   *pathogens* 7(6):e1002101.

728   16.   Wei X, *et al.* (2003) Antibody neutralization and escape by HIV-1. *Nature* 422(6929):307-

729   312.

730   17.   Wilen CB, Tilton JC, & Doms RW (2012) Molecular mechanisms of HIV entry. *Advances*

731   *in experimental medicine and biology* 726:223-242.

732   18.   Melikyan GB*, et al.* (2000) Evidence that the transition of HIV-1 gp41 into a six-helix

733   bundle, not the bundle configuration, induces membrane fusion. *The Journal of cell*

734   *biology* 151(2):413-423.

735   19.   Kuhmann SE, Platt EJ, Kozak SL, & Kabat D (2000) Cooperation of multiple CCR5

736   coreceptors is required for infections by human immunodeficiency virus type 1. *J Virol*

737   74(15):7005-7015.

738   20.   Weissenhorn W, Dessen A, Harrison SC, Skehel JJ, & Wiley DC (1997) Atomic structure

739   of the ectodomain from HIV-1 gp41. *Nature* 387(6631):426-430.

740   21.   Chan DC, Fass D, Berger JM, & Kim PS (1997) Core structure of gp41 from the HIV

741   envelope glycoprotein. *Cell* 89(2):263-273.

742   22.   Lu M, Blacklow SC, & Kim PS (1995) A trimeric structural domain of the HIV-1

743   transmembrane glycoprotein. *Nature structural biology* 2(12):1075-1082.

744   23.   Kwong PD*, et al.* (1998) Structure of an HIV gp120 envelope glycoprotein in complex

745   with the CD4 receptor and a neutralizing human antibody. *Nature* 393(6686):648-659.

746   24.   Wyatt R*, et al.* (1998) The antigenic structure of the HIV gp120 envelope glycoprotein.

747   *Nature* 393(6686):705-711.

748   25.   Courter JR*, et al.* (2014) Structure-Based Design, Synthesis and Validation of CD4-

749   Mimetic Small Molecule Inhibitors of HIV-1 Entry: Conversion of a Viral Entry Agonist to

750   an Antagonist. *Accounts of Chemical Research* 47(4):1228-1237.

751   26.   Schön A*, et al.* (2006) Thermodynamics of Binding of a Low-Molecular-Weight CD4

752   Mimetic to HIV-1 gp120. *Biochemistry* 45(36):10973-10980.

753   27.   Zhao Q*, et al.* (2005) Identification of N-phenyl-N′-(2,2,6,6-tetramethyl-piperidin-4-yl)-

754   oxalamides as a new class of HIV-1 entry inhibitors that prevent gp120 binding to CD4.

755   *Virology* 339(2):213-225.

756   28.   Haim H*, et al.* (2009) Soluble CD4 and CD4-mimetic compounds inhibit HIV-1 infection

757   by induction of a short-lived activated state. *PLoS pathogens* 5(4):e1000360.

758   29.   Madani N, *et al.* (2017) Activation and Inactivation of Primary Human Immunodeficiency

759   Virus Envelope Glycoprotein Trimers by CD4-Mimetic Compounds. *J Virol* 91(3).

760   30.   Richard J, *et al.* (2015) CD4 mimetics sensitize HIV-1-infected cells to ADCC.

761   *Proceedings of the National Academy of Sciences* 112(20):E2687.

762   31.   Madani N, *et al.* (2014) CD4-mimetic small molecules sensitize human

763   immunodeficiency virus to vaccine-elicited antibodies. *J Virol* 88(12):6542-6555.

764   32.   Yoshimura K, *et al.* (2010) Enhanced exposure of human immunodeficiency virus type 1

765   primary isolate neutralization epitopes through binding of CD4 mimetic compounds. *J*

766   *Virol* 84(15):7558-7568.

767   33.   Ding S, *et al.* (2017) Short Communication: Small-Molecule CD4 Mimetics Sensitize

768   HIV-1-Infected Cells to Antibody-Dependent Cellular Cytotoxicity by Antibodies Elicited

769   by Multiple Envelope Glycoprotein Immunogens in Nonhuman Primates. *AIDS research*

770   *and human retroviruses* 33(5):428-431.

771   34.   Ding S, *et al.* (2019) A New Family of Small-Molecule CD4-Mimetic Compounds

772   Contacts Highly Conserved Aspartic Acid 368 of HIV-1 gp120 and Mediates Antibody-

773   Dependent Cellular Cytotoxicity. *Journal of Virology* 93(24):e01325-01319.

774   35.   Rajashekar JK, *et al.* (2021) Modulating HIV-1 envelope glycoprotein conformation to

775   decrease the HIV-1 reservoir. *Cell host & microbe* 29(6):904-916.e906.

776   36.   Madani N, *et al.* (2018) A CD4-mimetic compound enhances vaccine efficacy against

777   stringent immunodeficiency virus challenge. *Nature Communications* 9(1):2363.

778   37.   Princiotto AM, *et al.* (2018) A Small-Molecule CD4-Mimetic Compound Protects Bone

779   Marrow–Liver–Thymus Humanized Mice From HIV-1 Infection. *The Journal of Infectious*

780   *Diseases* 218(3):471-475.

781   38.   Melillo B, *et al.* (2016) Small-Molecule CD4-Mimics: Structure-Based Optimization of

782   HIV-1 Entry Inhibition. *ACS Medicinal Chemistry Letters* 7(3):330-334.

783   39.   Anang S*, et al.* (2022) Characterization of Human Immunodeficiency Virus (HIV-1)

784   Envelope Glycoprotein Variants Selected for Resistance to a CD4-Mimetic Compound.

785   *Journal of Virology* 96(17):e00636-00622.

786   40.   Fritschi CJ*, et al.* (2021) Identification of gp120 Residue His105 as a Novel Target for

787   HIV-1 Neutralization by Small-Molecule CD4-Mimics. *ACS Medicinal Chemistry Letters*

788   12(11):1824-1831.

789   41.   Chen J*, et al.* (2019) Development of an Effective Scalable Enantioselective Synthesis of

790   the HIV-1 Entry Inhibitor BNM-III-170 as the Bis-trifluoroacetate Salt. *Organic Process*

791   *Research & Development* 23(11):2464-2469.

792   42.   Luo Z*, et al.* (2018) Stereogenic cis-2-substituted-N-acetyl-3-hydroxy-indolines via

793   ruthenium(II)-catalyzed dynamic kinetic resolution - asymmetric transfer hydrogenation.

794   *Chemical Communications* 54(96):13503-13506.

795   43.   Madani N*, et al.* (2008) Small-molecule CD4 mimics interact with a highly conserved

796   pocket on HIV-1 gp120. *Structure* 16(11):1689-1701.

797   44.   Miyaura N, Yamada K, & Suzuki A (1979) A new stereospecific cross-coupling by the

798   palladium-catalyzed reaction of 1-alkenylboranes with 1-alkenyl or 1-alkynyl halides.

799   *Tetrahedron Letters* 20(36):3437-3440.

800   45.   Müller K, Faeh C, & Diederich F (2007) Fluorine in pharmaceuticals: looking beyond

801   intuition. *Science* 317(5846):1881-1886.

802   46.   Kwon YD*, et al.* (2012) Unliganded HIV-1 gp120 core structures assume the CD4-bound

803   conformation with regulation by quaternary interactions and variable loops. *Proc Natl*

804   *Acad Sci U S A* 109(15):5663-5668.

805   47.   Fontaine J, Chagnon-Choquet J, Valcke HS, Poudrier J, & Roger M (2011) High

806   expression levels of B lymphocyte stimulator (BLyS) by dendritic cells correlate with HIV-

807   related B-cell disease progression in humans. *Blood* 117(1):145-155.

48.   Fontaine J, *et al.* (2009) HIV infection affects blood myeloid dendritic cells after

successful therapy and despite nonprogressing clinical disease. *J Infect Dis*

199(7):1007-1018.

49.   Pereyra F, *et al.* (2010) The major genetic determinants of HIV-1 control affect HLA

class I peptide presentation. *Science* 330(6010):1551-1557.

50.   Kamya P, *et al.* (2011) Receptor-ligand requirements for increased NK cell

polyfunctional potential in slow progressors infected with HIV-1 coexpressing

KIR3DL1*h/*y and HLA-B*57. *J Virol* 85(12):5949-5960.

51.   Peretz Y, *et al.* (2007) Functional T cell subsets contribute differentially to HIV peptide-

specific responses within infected individuals: correlation of these functional T cell

subsets with markers of disease progression. *Clinical immunology (Orlando, Fla.)*

124(1):57-68.

52.   Prévost J, *et al.* (2020) The HIV-1 Env gp120 Inner Domain Shapes the Phe43 Cavity

and the CD4 Binding Site. *mBio* 11(3).

53.   Z. Gong et al., HIV-1 gp120 complex with BNM-III-170. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FLY. Deposited 22 December 2022.

54.   Z. Gong et al., HIV-1 gp120 complex with CJF-III-049-S. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FLZ. Deposited 22 December 2022.

55.   Z. Gong et al., HIV-1 gp120 complex with CJF-III-214. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM0. Deposited 22 December 2022.

56.   Z. Gong et al., HIV-1 gp120 complex with CJF-III-289. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM2. Deposited 22 December 2022.

57.   Z. Gong et al., HIV-1 gp120 complex with CJF-III-288. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM3. Deposited 22 December 2022.

58.   Z. Gong et al., HIV-1 gp120 complex with CJF-IV-047. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM4. Deposited 22 December 2022.

59.   Z. Gong et al., HIV-1 gp120 complex with DY-III-065. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM5. Deposited 22 December 2022.

60.   Z. Gong et al., HIV-1 gp120 complex with CJF-III-192. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM7. Deposited 22 December 2022.

61.   Z. Gong et al., HIV-1 gp120 complex with CJF-IV-046. Protein Data Bank.

https://www.rcsb.org/structure/unreleased/8FM8. Deposited 22 December 2022.